A Syntenic Pangenome for Gardnerella Reveals Taxonomic Boundaries and Stratification of Metabolic and Virulence Potential across Species

Heather Bouzek[*] [1,2], Martha A. Zepeda-Rivera[*] [1,2], Sujatha Srinivasan[2], Elliot M. Lee[2], Dakota S. Jones[2], Elsa F. McMahon[2], Susan M. Strenk[2], Tina L. Fiedler[2], Michael France[3,4], Jacques Ravel[3,4], David N. Fredricks[2#], Christopher D. Johnston[1,2#]

[1]Department of Genomic Medicine, University of Texas MD Anderson Cancer Center, Houston, TX, USA

[2] Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Center, Seattle, Washington, USA

[3] Center for Advanced Microbiome Research and Innovation, Institute for Genome Sciences, University of Maryland School of Medicine, Baltimore, Maryland, USA

[4]Department of Microbiology and Immunology, University of Maryland School of Medicine, Baltimore, Maryland, USA

[#]**Address correspondence to**:

David N. Fredricks, MD, dfredric@fredhutch.org

Christopher D. Johnston, PhD, cdjohnston@mdanderson.org

---

[*] These authors contributed equally to this work.

Abstract

     Bacterial vaginosis (BV) is a prevalent condition characterized by an imbalance in the vaginal microbiota, often involving the presence of *Gardnerella* species. The taxonomic complexity and inconsistent nomenclature of *Gardnerella* have impeded progress in understanding the role of this genus in health and disease. In this study, we obtained genomes from 392 *Gardnerella* isolates from asymptomatic individuals and those with BV. With stringent curation of these and additional publicly-available genomes, we conducted a comprehensive genomic analysis (final dataset: n=312). We resolve taxonomic ambiguities using current species thresholds and propose updated *Gardnerella* nomenclature to facilitate formal naming of these twenty-one identified lineages (eleven species, fifteen subspecies). Syntenic pangenome analysis revealed lineage-specific genetic factors and putative operons associated with virulence, metabolism, and genetic defense mechanisms. We also report on the first cryptic plasmids naturally present within the *Gardnerella* genus, which should serve to advance mechanistic studies within these species. Our findings provide a unified framework for *Gardnerella* taxonomy and nomenclature, and advance our understanding of species-specific functional capabilities, with implications for *Gardnerella* research, diagnostics, and targeted therapeutics in BV.

Introduction

     Bacterial vaginosis (BV) is a common and global condition characterized by an imbalance in the vaginal microbiota. This imbalance typically involves a reduction in beneficial *Lactobacillus* species and an overgrowth of obligate and facultative anaerobes[1], leading to symptoms such as unusual discharge, odor, and discomfort[2]. BV is associated with significant health risks, including

increased susceptibility to sexually transmitted infections[3] and adverse pregnancy outcomes such

as preterm birth and low birth weight[4,5].

While molecular methods have associated many bacterial species with BV[6-12], *Gardnerella*

vaginalis was the first bacterium isolated from individuals affected with BV[13] and was long

thought to be the sole causative agent of this condition[14]. This assumption was based on

Gardnerella species exhibiting a high virulence potential, including adherence to the vaginal

epithelium[15], production of cytotoxins[16], formation of biofilms[17], and its presence in 95–100% of

BV cases[18-20]. Moreover, *Gardnerella* species thrive in the vaginal environment, even at the low

pH typical of the *Lactobacillus*-dominant genital tract microbiota[21]. Although vaginal

concentrations of *Gardnerella* are greater in those with BV[18], *Gardnerella* species are also present

in asymptomatic individuals, suggesting the potential for both commensal and pathogenic lineages

or states. Our previous research revealed that while *Gardnerella vaginalis* is correlated with a

greater risk ratio for BV than other validly named *Gardnerella* species, no single *Gardnerella*

species is specific for BV based on a *cpn60* qPCR. In fact, the presence of a high *Gardnerella*

species diversity is a risk factor for the development of BV in our longitudinal study[18]. However,

the genetic capabilities of individual *Gardnerella* lineages remain unclear, partially due to

inconsistent taxonomic assignments and overly complex *Gardnerella* nomenclature[22], despite

laudable efforts from multiple research groups[23-31]. In this study, we used various comparative

genomic approaches to taxonomically identify *Gardnerella* lineages and propose a single naming

scheme for all currently known species that is reflective of phylogenetic placement and historical

context. We obtained genomes from 392 *Gardnerella* isolates from asymptomatic individuals and

those affected with BV. We stringently curated additional publicly available *Gardnerella* genomes

for a final genome set (n=312), which was analyzed via *cpn60* gene typing, average nucleotide

identity, digital DNA/DNA hybridization, and phylogenetic approaches. Our analyses identified

eleven distinguishable *Gardnerella* species, encompassing fifteen subspecies. Here, in addition to

proposing an updated nomenclature scheme for twenty-one distinct *Gardnerella* lineages, we

identify genomic features associated with each through pangenome analysis. Our work aims to

advance knowledge on the genetic characteristics and predicted functional capabilities of lineages

within the *Gardnerella* genus.

Results

***Gardnerella* and *Bifidobacterium* are distinct genera**

We obtained genomes for 392 *Gardnerella* isolates (Fredricks and Ravel bacterial isolate

collections) from asymptomatic individuals and those with BV. We curated these and additional

publicly available *Gardnerella* genomes (n=387 within the NCBI database as of May 2025) using

stringent quality metrics to remove duplicate and low-quality genomes (**Methods,**

Supplementary Table 1). This resulted in a final curated set of 312 genomes for comparative

genomic analyses (**Table 1**, **Supplementary Table 2**). Analysis of the *cpn60* gene (which encodes

a 60 kDa chaperonin) currently provides the most reliable method for *Gardnerella* species typing.

These species cannot be distinguished by traditional 16S rRNA gene sequence typing[32,33]. BLAST

analysis of the *cpn60* gene of these isolates supported their identification as *Gardnerella* species

(**Supplementary Table 2**).

Despite the classification of these genomes as *Gardnerella* against and within the NCBI

database, initial speciation against the Genome Taxonomy Database (GTDB) assigned many of

these genomes to *Bifidobacterium* (**Supplementary Table 2**). However, *Bifidobacterium* is known

to be a polyphyletic group, consisting of at least six distinct clades[34,35], with its members exhibiting

high genetic diversity. The separation of *Gardnerella* from *Bifidobacterium* has previously been

demonstrated through comparisons of genome length, G+C percentage, and overall genome

composition.[28,36] Comparisons of genome length and G+C percentage between our *Gardnerella*

genome set, *Bifidobacterium,* and closely related *Cardovia* (e.g., *Alloscardovia, Aeroscardovia,*

etc.) genomes (n=227) indicate that *Gardnerella* genomes are significantly shorter ($p<0.001$,

Wilcoxon *post hoc* test) and have lower G+C content ($p<0.001$, Wilcoxon *post hoc* test) than other

Bifidobacterium members (**Supplementary Figure 1, Supplementary Table 3**). Additionally,

average nucleotide identity (ANI) values fell below generally accepted genus thresholds of 73-

75% (**Supplementary Figure 1**). These results confirmed that our final curated genome set

included only *Gardnerella* species, further supporting previous evidence that *Gardnerella* and

Bifidobacterium represent distinct genera within the *Bifidobacteriaceae* family.

Lineage	Abbreviation	No. Genomes	Genome Length	GC %
G. aggregata	Ga	n=86	1.37-1.72 Mbp	41.64%-42.50%
G. aggregata subsp. aggregataa	Gaa	n=5	1.48-1.53 Mbp	42.11%-42.50%
G. aggregata subsp. hutchinsoni	Gah	n=11	1.52-1.66 Mbp	42.03%-42.43%
G. aggregata subsp. leopoldii	Gal	n=24	1.37-1.67 Mbp	42.07%-42.34%
G. aggregata subsp. swidsinskii	Gas	n=46	1.52-1.72 Mbp	41.64%-42.29%
G. bivia	Gb	n=6	1.55-1.64 Mbp	37.90%-38.31%
G. coleohominis	Gc	n=6	1.64-1.72 Mbp	40.53%-40.76%
G. insolita	Gi	n=3	1.59-1.66 Mbp	41.27%-43.80%
G. insolita subsp. insolita	Gii	n=2	1.61-1.66 Mbp	41.27%-43.80%
G. insolita subsp. libera	Gil	n=1	1.59 Mbp	43.60%
G. kenyensis	Gk	n=1	1.66 Mbp	43.62%
G. media	Gm	n=26	1.49-1.65 Mbp	42.90%-43.50%
G. media subsp. greenwoodii	Gmg	n=13	1.49-1.65 Mbp	42.90%-43.44%
G. media subsp. media	Gmm	n=2	1.53-1.55 Mbp	43.04%-43.05%
G. media subsp. pittsburghi	Gmp	n=11	1.50-1.64 Mbp	43.05%-43.50%
G. nova	Gn	n=2	1.62-1.70 Mbp	45.35%-45.46%
G. rara	Gr	n=6	1.47-1.58 Mbp	43.01%-43.29%
G. sialicivorans	Gs	n=60	1.50-1.68 Mbp	42.00%-42.74%
G. sialicivorans subsp. picketti	Gspk	n=40	1.51-1.68 Mbp	42.00%-42.74%
G. sialicivorans subsp. piotii	Gspt	n=17	1.50-1.59 Mbp	42.22%-42.61%
G. sialicivorans subsp. sialicivorans	Gss	n=3	1.50-1.53 Mbp	42.39%-42.48%
G. vaginalis	Gv	n=111	1.57-1.78 Mbp	40.99%-41.82%
G. vaginalis subsp. regina	Gvr	n=15	1.62-1.72 Mbp	41.11%-41.42%
G. vaginalis subsp. transversa	Gvt	n=17	1.57-1.67 Mbp	41.25%-41.62%
G. vaginalis subsp. vaginalis	Gvv	n=79	1.58-1.78 Mbp	40.99%-41.82%
G. washingtoni	Gw	n=5	1.56-1.65 Mbp	42.89%-43.11%

Table 1: *Gardnerella* lineages referenced in this study

Table 1 shows the *Gardnerella* lineages identified and referenced in this study along with their proposed names and unique abbreviations. Summary statistics, including the number of genomes in this collection, the range of genome length and the G+C content are shown for each lineage.

Identification and proposed revised nomenclature for *Gardnerella* lineages

Given the importance of *Gardnerella* species diversity in BV[18], we wanted to identify the number of lineages present in our genome collection. Although different *Gardnerella* species have been identified through various approaches[27-30,37,38], the naming of these groups has been a

consistent challenge for the *Gardnerella* field despite extensive and laudable efforts from multiple

research groups (**Figure 1**). In summary, Piot *et al.*[23] initially identified *Gardnerella* biotypes

based on sugar hydrolysis phenotypes. When this approach proved insufficient to capture the

diversity of the genus*,* it was expanded to 17 biotypes by adding three fermentation targets[24]. The

use of early genomic methods, including restriction fragment length polymorphism (RFLP)[25] and

amplified ribosomal DNA restriction analysis (ARDRA)[26], identified distinct *Gardnerella*

genotypes. Whole-genome comparisons brought about another naming scheme seeking to

encompass all strains, where "A", "B", "C", and "D"[27] were proposed designations for a newer

condensed species assignment[22,33]. Nearly simultaneously, another method was proposed[39], using

"1", "2", "3", and "4"[21] for the same clades, which led to a disjointed hybrid designation (C/1, B/2,

A/4, and D/3). The species were then renamed as reassorted ecotypes[28], using yet another

nomenclature, one that incorporated the genomic context of the distinct types. After whole-genome

sequencing became widely available, three more competing standards based on ANI emerged,

describing thirteen genomic species (Gsp)[29], nine genomospecies (GS)[30], and nine different

genotypes (GGtypes)[36]. Formal naming of a limited subset of these species, including *G.*

vaginalis[13], *G. pickettii*[40]*, G. piotii*[29], *G. leopoldii*[29], *G. swidsinskii*[29], and *G. greenwoodii*[40], has

further compounded this complex nomenclature (**Figure 1**). Despite the use of different

classification schemes, prior studies consistently identified two distinct *Gardnerella* subgenera

(Set A and Set B), each encompassing specific lineages[27].

Consistent with this, *cpn60* gene sequences and reference-free whole-genome phylogenies

from our collection clearly separated all genomes into Set A or Set B (**Figure 2**, **Supplementary**

Figure 2). To determine the number of *Gardnerella* species represented, we compared the

genome-wide ANI values and identified a 94% threshold that delineated intra-species comparisons

(**Supplementary Figure 3**), slightly below the prokaryotic species ANI threshold of 95 ± 0.5 %

reported by Jain *et al.*[41]. ANI and digital DNA-DNA hybridization (dDDH) values were co-linear

(**Supplementary Figure 3**) and as such dDDH did not provide additional information. Using a

94% ANI threshold, we identified eleven *Gardnerella* species within our dataset. Whole-genome

phylogenies combined with ANI values further highlighted subspecies groups in five of these

species, resulting in twenty-one distinct *Gardnerella* lineages (**Figure 2, Supplementary Table**

4). These species designations were congruent with the classification proposed by Potter *et al.*[30].

Given the extensive history of conflicting classification schemes (**Figure 1**), we next sought to

standardize species and subspecies names across all lineages identified (**Figure 1**, **Figure 2**, **Table**

1, **Supplementary Table 2**).

Details on the rationale for each proposed nomenclatural revision, including species-

specific functional attributes and ecological isolation sources, are provided in the **Protologues**.

Our aim was to integrate previously used species and subspecies names to maintain a consistent

framework. Historically recognized "*G. vaginalis*" corresponds to a subspecies group that we now

propose to designate as *G. vaginalis* subspecies *vaginalis*, preserving historical continuity while

reflecting its phylogenetic placement (**Figure 1**, **Table 1**, **Supplementary Table 1**). Similarly, we

propose distinct species names for lineages with well-supported genomic and phenotypic

divergence, including *G. sialicivorans*, *G. aggregata,* and *G. media*, among others. To minimize

ambiguity in referring to *Gardnerella* species in broader studies, we developed guidelines for

assigning unique shorthand notation for each lineage (**Methods**). Species names are to be

abbreviated with two letters (*Ga*, *Gb*, *Gc*, *Gi*, *Gk, Gm, Gn, Gr*, *Gs*, *Gv* and *Gw*), while individual

subspecies should be designated with three-letter abbreviations (except for *Gs* subsp. *pickettii* and

Gs subsp. *piotii,* which require four letters, *Gspk* and *Gspt,* to distinguish them from other named

subspecies) (**Figure 1**, **Table 1**, **Methods**).

Figure 1: Historical context of *Gardnerella* nomenclature

A) From top to bottom schematic shows the identification method by type, publication first author,

names given to *Gardnerella* or naming framework, date of publication, and resulting conclusions

from naming schemes. **B**) *Gardnerella* subtyping resulting from the publication in (A) where each

column represents a subset of *Gardnerella* named. Lineages between different publications are

connected by arrows. The final two columns contain the updated lineages with corresponding

species and subspecies names proposed within this study.

Figure 2: Average nucleotide identity values between *Gardnerella* genomes

Average nucleotide identity values between *Gardnerella* genomes in historical **A**) Set A and **B**) Set B. Key for each panel indicates corresponding values to each color. White represents values below 90%. Genomes are organized based on a kSNP4[42]-generated whole-genome reference-free phylogeny.

A syntenic pangenome for *Gardnerella*

After stringently identifying the *Gardnerella* species and subspecies lineages in our genome set and assigning each a standardized name, we were able to accurately investigate the genetic features associated with each. Pangenome analysis provides a comprehensive view of the genetic repertoire within a group of bacteria by identifying genetic features that are well-conserved across most (>95%) genomes ("core genome") versus those that are only present in a subset of genomes ("accessory genome"). While the core genome often contains genes involved in essential cellular processes[43,44], analysis of the accessory genomes can provide key insights about genetic diversity, adaptation, and evolutionary relationships. For example, previous pangenome efforts in *Gardnerella* have expanded taxonomic groupings, identified metabolically essential genes, and uncovered virulence and niche adaptation features[31,45,46]. However, bacterial genes often function in co-regulated units or clusters involved in specific biological processes (e.g. operons). To identify putative operons within the *Gardnerella* pangenome, we applied a syntenic approach that accounts for gene order by using the Partitioned PanGenome Graph of Linked Neighbors (PPanGGOLiN) tool[43] (**Figure 3**, **Figure 4, Supplementary Table 5**). Briefly, the coding sequences of genes are grouped into "nodes" based on amino acid similarity and visualized as circular nodes, with node size reflecting the relative presence of that gene group across the genome set (**Figure 4A-C**).

Nodes representing neighboring genes are linked into syntenic units called "modules"

(**Supplementary Figure 4**). This analysis clearly distinguished between the core ("persistent"

partition) and accessory genomes, and enabled identification of accessory genetic elements present

at intermediate frequency ("shell" partitions) or very low frequency ("cloud" partition) (**Figure 3**).

By leveraging modules and pangenome partitions, we can trace conserved arrangements of genes

across multiple genomes and simultaneously gain insight into the functional and evolutionary

dynamics driving diversity at the genus, species, and strain levels.

The core genome in our *Gardnerella* dataset reflected approximately 16% of identified

nodes, enriched in essential cellular functions including translation and ribosomal structure,

transcription, RNA processing, signal transduction, energy production, and metabolism of

nucleotides, amino acids, inorganic ions, and lipids (**Figure 3, Supplementary Figure 4**).

Periodically, these conserved nodes are interrupted by shell-associated nodes (**Figure 4A,** nodes

in distinct colors, branching from orange nodes), which mark points of evolutionary divergence.

These shell nodes often represent alternate evolutionary pathways, commonly associated with

horizontal gene transfer (HGT) events. We noted node differences between shell partitions that

were consistent with virulence, metabolism, and defense systems, warranting further investigation.

Figure 3: *Gardnerella* pangenome partitions

Heatmap displays the presence and absence of nodes in each partition where each row represents a node, and each column corresponds to a *Gardnerella* genome. Colors on the side indicate the core/persistent (orange), phage (red), shells (green), and cloud (blue) partitions. Genomes are ordered by kSNP4[42] phylogenies of Set A and Set B, with colors on dendrogram indicating lineage.

A

B

C

D

Tree Scale:10 Tree Scale:10

Type I pullanase

Sialidase (NanH1)
Sialidase (NanH2)
Sialidase (NanH3)
Sialidase (NanH4)
Vaginolysin

G. vaginalis *G. sialicivorans* *G. rara* *G. aggregata* *G. bivia* *G. coleohominis* *G. media* *G. washingtoni* *G. insolita* *G. kenyensis* *G. nova*

Gvr *Gspk* *Gaa* *Gmm* *Gii*
Gvt *Gspt* *Gah* *Gmg* *Gil*
Gvv *Gss* *Gal* *Gmp*
Gas

E Interset Comparisons

NanH1
p<0.00001
Proportion of Genomes
Set A   Set B

NanH2
p<0.00001
Proportion of Genomes
Set A   Set B

NanH3
p<0.00001
Proportion of Genomes
Set A   Set B

NanH4
p=0.00782
Proportion of Genomes
Set A   Set B

Vaginolysin
p<0.00001
Proportion of Genomes
Set A   Set B

F Lineage-specific associations

NanH2
p<0.00001
Proportion of Genomes
Gr   Other

NanH3
p<0.00001
Proportion of Genomes
Gs   Other

NanH4
p<0.00001
Proportion of Genomes
Gv   Other

Vaginolysin
p=0.0006
Proportion of Genomes
Gvv   Other

Figure 4: Syntenic maps of the *Gardnerella* pangenome

A-C) Visualizations of the PPanGGOLiN[43]-generated *Gardnerella* pangenome. Each node represents a gene group, with syntenic nodes representing neighboring genes. Size of node indicates relative presence across *Gardnerella* genomes. Colors indicate core (orange) or shell (other colors) partitions. Core and shell partitions are shown individually in **B**) and **C**), respectively. **D**) Presence-absence heatmap of canonical *Gardnerella* virulence factors and the well-conserved pullulanase across each *Gardnerella* lineage. Heatmap is organized by kSNP4[42] phylogenies of Set A and Set B. Colors indicate lineage and presence/absence as denoted in key. **E-F**) Graphs show the virulence factor presence compared by genome proportion between *Gardnerella* **E**) Set A and Set B and **E**) specific lineages and all others. Statistical analysis was carried out using a two-sample z-test, two-tailed.

Virulence factors and predicted pathogenicity across *Gardnerella* lineages

  Gardnerella colonization of the vaginal niche, as well as its pathogenesis, is dependent on mucin degradation, adherence to the vaginal epithelium, and the formation of stable biofilms[15]. Canonical *Gardnerella* virulence factors include four sialidase variants[47] (NanH1, NanH2, NanH3, NanH4) that aid in mucin degradation, and vaginolysin[16,48], which binds and lyses host cells in the vaginal mucosa. In our pangenome dataset we identified specific nodes associated with each of these virulence factors (**Supplementary Table 5**).

  The Type I pullulanase protein was well conserved (present in 100% of *Gardnerella* genomes), in agreement with previous reports (**Figure 4D**). Comparisons between *Gardnerella* Set A and Set B showed that the four sialidase proteins (NanH1-NanH4) are significantly enriched in Set A genomes, while vaginolysin is significantly enriched in Set B genomes (**Figure 4E**).

Lineage-specific associations were also observed for three of the sialidase variants as well as

vaginolysin (NanH2-*Gr*, NanH3-*Gs*, NanH4-*Gv*, vaginolysin-*Gvv*) (**Figure 4F**). Our results

support the hypothesis that variable lineage-dependent pathogenic potential exists across

Gardnerella.

Predicted metabolic capabilities across *Gardnerella* lineages

Given the observed difference in metabolism-related nodes, we bioinformatically predicted

the biosynthesis of amino acids and the catabolism of small carbon sources across our *Gardnerella*

genome dataset (**Figure 5A, Supplementary Table 6**). The biosynthesis of glutamine, glycine,

and proline appeared to be well conserved (>99% of genomes) while a high proportion of genomes

(89%) harbored the capacity for cysteine biosynthesis (**Figure 5A**). Compared to Set A, genomes

across Set B were significantly enriched for asparagine, threonine, and methionine biosynthesis

(**Figure 5B**). Within Set B, asparagine biosynthesis was significantly depleted in *Ga*, threonine

biosynthesis was significantly depleted in *Gn,* and methionine biosynthesis was significantly

depleted in *Gm* and *Gb* and significantly enriched in *Ga* (**Figure 5C**). Although asparagine and

threonine biosynthesis were significantly enriched in Set B, we noted their species-specific

enrichments within Set A. Compared to all other Set A genomes, asparagine and threonine

biosynthesis were significantly enriched in *Gr* and *Gv*, respectively (**Figure 5C**). *Gardnerella*

species comparisons indicated that asparagine biosynthesis was significantly enriched in *Gm,*

cysteine biosynthesis was significantly depleted in *Gc* and *Gr*, and proline biosynthesis was

significantly depleted in *Gn* (**Figure 5D**). Lineage comparisons within species indicated that

asparagine biosynthesis was significantly enriched in *Gal*, methionine biosynthesis was

significantly enriched in *Gmp,* asparagine biosynthesis was significantly depleted in *Gmm*, and

cysteine biosynthesis was significantly depleted in *Gss* (**Figure 5D**).

Genes for the catabolism of ethanol appear to be a well-conserved feature of *Gardnerella*

genomes (**Figure 5A**). Genomes across Set A were further significantly enriched for the catabolism

of alanine, gluconate, and serine (**Figure 5E**). Notably, lineage-specific associations within Set A

were observed for the catabolism of each of these small carbon sources. While gluconate

catabolism is significantly enriched in *Gv*, alanine catabolism is significantly depleted in *Gvr*

(**Figure 5F**). Serine catabolism was completely absent in *Gr* and within Set B genomes was

significantly depleted in *Gmp* and *Gas* compared to other lineages within their respective species

(**Figure 5G**). Interestingly, the catabolism of acetate and deoxyribose appeared to be lineage-

specific and unique to *Gb* and *Gv*, respectively (**Figure 5G**). In the case of deoxyribose, within *Gv*

its catabolism was significantly enriched in subspecies *Gvv* (**Figure 5G**). Our results indicate that

Set A and Set B genomes are distinguished by their relative enrichment of pathways for catabolism

of small carbon sources or synthesis of distinct amino acids with lineage-specific associations.

Figure 5: Predictions for metabolic capabilities across *Gardnerella* lineages

A) Presence-absence heatmap for GapMind-predicted[49-51] pathways to synthesize amino acids
(AA) or catabolize small carbon sources (SCS) where each row represents a *Gardnerella* genome.

257   Genomes are ordered by kSNP4[42] phylogenies of Set A and Set B. Presence by proportion of

258   genomes between Set A and Set B (**B** and **E**), within Set A or Set B (**C** and **F**), and between specific

259   lineages (**D** and **G**) are shown. Statistical analysis was carried out using a two-sample z-test, two-

260   tailed. Colors represent each lineage.

261
Phage and antiphage defense mechanisms across *Gardnerella* genomes

In our pangenome data set, we noted that multiple nodes were related to phage-associated

proteins that encompassed two distinct *Siphoviridae* family *Gardnerella* prophage, including the

previously described vB_Gva_AB1[52,53] and a novel phage we have tentatively designated as

"phage_gamma" (**Supplementary Table 7**). Although vB_Gva_AB1 is sometimes reported as

several different prophages, the discrete synteny within our PPanGGOLiN pangenome suggests

that these variants represent a single phage (**Supplementary Table 5**). It is notable that some

strains appear to have remnant phage genes, although these appeared to be non-syntenic and

scattered throughout the genome. An arbitrary cutoff of greater than ten phage_gamma genes and

greater than twenty-five vB_Gva_AB1 genes was considered for phage presence (**Supplementary**

Figure 5). While both vB_Gva_AB1 and phage_gamma were identified in Set A and Set B

genomes, vB_Gva_AB1 was significantly enriched in Set B genomes (**Supplementary Figure 5**).

To counteract phages, bacteria have evolved an array of defense systems[54], with the most

widespread and well-characterized being restriction-modification (RM) and CRISPR-Cas

mechanisms[55]. Across our *Gardnerella* genome dataset we identified the presence of sixty different

systems, with an average of twelve defenses per genome (**Figure 6A-B**, **Supplementary Figure**

6, Supplementary Table 8). Of the defense systems with described mechanisms, dXTPase

systems[56] were well conserved (99% genomes). Type I RM systems were prevalent (56% of

genomes) and did not differ significantly between *Gardnerella* sets. Set A genomes were

significantly enriched for some Abi proteins (AbiD[57] and AbiL[58]), CBASS defenses[59,60], and DRT

mechanisms[61,62], while Set B showed enrichment for a larger number of defenses, including those

with Abi-mechanisms (AbiE[63], Gabija[64,65], Lamassu[64,66,67]), CRISPR-Cas (CRISPR arrays and Cas

Type IE proteins), and Type II RM defenses (**Supplementary Figure 7**). Across *Gardnerella*

lineages, *Gas* showed an enrichment for DNA-modification system (DMS) proteins, while *Gvv*

showed an enrichment for AbiL and DRT defenses (**Supplementary Figure 7**).

Although defense mechanisms evolved primarily to protect against bacteriophages, their

activity extends against other horizontally transferred genetic elements, as well as plasmid vectors

used during *in vitro* transformation. RM defenses, which are dependent upon detection and

destruction of inappropriately methylated foreign DNA, are one of the most prevalent across

bacteria. Further, it is notable that approaches to overcome these RM systems during genetic

engineering have been shown to significantly improve transformation efficiency across multiple

bacterial species[68-71]. Thus, we sought to characterize RM defenses across a subset of our

Gardnerella genome dataset. We obtained and analyzed methylomes for 42 *Gardnerella* isolates

across thirteen lineages within the Fredricks isolate collection (**Supplementary Table 2**). A total

of 39 methylated motifs were detected and assigned to Type I (25 motifs), Type II (11 motifs), or

Type III (3 motifs) RM systems (**Figure 6C**, **Supplementary Table 9**).

Figure 6: Defense systems and cryptic plasmids across _Gardnerella_ genomes

A-B) Presence-absence heatmaps for PADLOC-predicted[66,72] defense mechanisms. Genomes are ordered by kSNP4[42] phylogenies of **A)** Set A and **B)** Set B. **C)** Sunburst plot shows the sequence of each methyl-modified motif detected within each *Gardnerella* lineage. The size of the section represents the prevalence within each lineage. **D)** Schematics depict plasmid annotations for pDNF1199S-01, p0756_546_1_1_BHK10-01, and pNR047-01.

Identification of a native Gardnerella cryptic plasmid

There is renewed interest in developing genetic tools for engineering *Gardnerella* species[73,74]. Because *Gardnerella* were long thought to lack mobile genetic elements, such as native plasmids, most efforts have focused on adapting non-*Gardnerella* genetic tools for use in this genus. Recent studies have been extensive and show promising results, achieving targeted chromosomal integrations and deletions using *E. coli*-based suicide vectors[73], as well as reproducible transformation with a *Bifidobacterium-E.coli* shuttle vector, albeit with low efficiency[74]. During genome sequencing of the Fredricks clinical isolate collection, we observed an independent circular contig, with high coverage, forming a second replicon within the otherwise closed genome sequence of *Gspk* strain DNF01199S (**Figure 6D, Supplementary Table 10**). Annotation of this contig showed three coding sequences for hypothetical proteins and a single putative replication initiation protein (**Supplementary Table 11**). Subsequent plasmid isolation and validation confirmed the episomal nature of this contig, marking pDNF01199S-01 as the first cryptic plasmid identified within the *Gardnerella* genus.

Motivated by this discovery, we assessed whether this or additional cryptic plasmids were present in our *Gardnerella* pangenome. We reasoned that the current limited knowledge of such plasmids reflects their rarity. As such, we focused on the cloud partition of the accessory genome,

which highlights strain-specific genetic factors and collectively comprises 47.23% of the

Gardnerella pangenome. Using structural sequence insights from pDNF01199S-01, we identified

two additional putative plasmid sequences within the cloud partition: one in *Gvv* strain

0756_546_1_1_BHK10[75] (p0756_546_1_1_BHK10-01), which was isolated from cervicovaginal

mucus in South Africa, and another in *Gk* strain NR047 (pNR047-01)[76], isolated from a vaginal

swab in Nairobi, Kenya. All three plasmids are approximately 4 kb in size, contain coding

sequences for putative hypothetical proteins, and share a highly similar replication initiation

protein (84.48%-95.26% similar), suggesting a potentially conserved replication mechanism for

cryptic plasmids within the *Gardnerella* genus (**Figure 6D**, **Supplementary Tables 10-11**).

Given that pDNF01199S-01 stably replicates, we next sought to engineer a *Gardnerella-*

E.coli shuttle vector that could be used for the transformation of other *Gardnerella* isolates. To

establish an antibiotic selectable marker, we noted that within our pangenome dataset, 22% of

genomes contained the presence of *tetM*, including *Gspk* DNF01199S. This was consistent with

previous reports on the variable tetracycline resistance phenotypes across *Gardnerella* strains[77].

Thus, we assessed the ability of both *tetM+* and *tetM-* isolates to grow in the presence of

supplemented tetracycline. As predicted, *tetM+* isolates were resistant to tetracycline while *tetM-*

isolates showed susceptibility, indicating that the *tetM* cassette confers tetracycline resistance *in*

vitro, and could serve as a selectable marker for this vector. To enable the propagation of our

Gardnerella-E.coli shuttle vector in *E. coli* strains, we combined pDNF01199S-01, *tetM* cloned

from DNF01199S, and the *E. coli* minicircle plasmid backbone (pMC) into a single parental

plasmid (herein designated pG01-pp) (**Supplementary Figure 8A-C**). As previously reported[68],

we induced and purified the pDNF01199S-tetM-MC minicircle vector (pG01-MC). We

subsequently attempted to transform this minicircle into the *tetM-* strain, *Gvv* ATCC 14018

(**Supplementary Figure 8D**). While this yielded no transformants, we predicted that overcoming

RM defenses present in these strains would increase transformation efficiency. A previously

reported RM system modifying "GGCC" sites has been described in ATCC 14018[78], and bypassing

this system was shown to improve transformation efficiency with suicide-based vectors[73]. Thus,

we used commercially available *HaeIII* methyltransferase (MTase) to methyl-modify these sites

in pG01-MC prior to transformation. In combination with further optimization of growth and

transformation conditions, transformation efficiency was substantially increased through these

approaches from a minimum of 2.30×10^0 transformants per µg of DNA to 6.85×10^2

(**Supplementary Figure 8D**). Collectively, these results show that pG01 is the first replicative

shuttle vector based upon a native *Gardnerella* cryptic plasmid. Together with recent work by the

Dillard and Hill groups[73,74], these efforts should establish a robust platform for more mechanistic

investigations of *Gardnerella* biology and pathogenesis.

Discussion

Taxonomic inconsistencies and fluctuating nomenclature have made it inherently

challenging to delineate the roles of *Gardnerella* lineages in health and disease states. Our work

seeks to resolve these ambiguities by integrating whole-genome analyses to produce a more refined

Gardnerella taxonomic classification system. We further propose an updated and unified

Gardnerella nomenclature that is reflective of these lineages. Proposing nomenclatural changes,

particularly for well-established organisms such as '*G. vaginalis'*, (updated here to *G. vaginalis*

subsp. *vaginalis*), can often be contentious and may face resistance within the scientific

community. However, updating *Gardnerella* lineages to appropriate species and subspecies

designations, using established criteria, is necessary to prevent further taxonomic confusion. We

noted that while the *cpn60* gene, often used for *Gardnerella* typing, can largely distinguish between

Gardnerella species, its resolution is insufficient for subspecies assignments. Accurate subspecies

assignments require whole-genome approaches through either phylogenetic assessment or average

nucleotide identity comparisons. When additional *Gardnerella* lineages are isolated, whole-

genome sequencing efforts should allow for continuous integration into this updated taxonomic

and nomenclature scheme. Ultimately, our naming efforts seek to increase clarity and ensure that

the functional role of specific species in the genital tract can be accurately interpreted, while

preserving previous naming efforts, where possible.

With this refined taxonomic and nomenclature framework, we probed for genetic features

and putative operons associated with each lineage by applying a syntenic pangenome approach.

We noted that the vast majority of *Gardnerella* gene content (~84%) is species and strain-variable

with many differences associated with virulence, metabolism, and genetic defense systems.

Gardnerella Set A genomes (*Gr, Gs,* and *Gv*) showed an enrichment for sialidases and an increased

potential for the catabolism of small carbon sources. In contrast, Set B genomes (*Ga, Gb, Gc, Gi,*

Gk, Gm, Gn, and *Gw*) were enriched in putative adhesion and virulence proteins, displayed a

greater capacity for amino acid biosynthesis, had a higher propensity to harbor the vG_gva_AB1

prophage, and contained a broader repertoire of genetic defense mechanisms.

Mechanistic studies to assess the roles of *Gardnerella* genetic factors, including those

identified in this study, are currently hindered by limited genetic tools available for *Gardnerella*.

In fact, it has previously been reported that *Gardnerella* species lack mobile genetic elements, such

as plasmids[27]. However, our extensive genome sequencing efforts reveal the first *Gardnerella*

cryptic plasmid naturally occurring within a single cultivated strain, along with additional

homologous sequences in publicly available genomes. Coupled with recent advances in

Gardnerella species engineering, this discovery of extrachromosomal and self-replicative genetic

elements within the *Gardnerella* genus should facilitate the development of advanced genetic

tools. Collectively, overcoming genetic intractability within *Gardnerella* species will greatly

advance our ability to investigate the role of the genus in health and disease states.

Acknowledgements

Research reported in this publication was supported in part by the National Human Genome

Research Institute as part of the Human Microbiome Project Initiative focused on technology

development under award number R01HG005816 to D.N.F., the National Institute for Allergy and

Infectious Diseases of the National Institutes of Health under award number U19AI084044 to J.R.,

the National Institute of Dental and Craniofacial Research (NIDCR) of the National Institutes of

Health Office of the Director (OD) Common Fund under award number R01DE027850 to C.D.J.

Materials and Methods

Collection, isolation, sequencing, and initial typing of Gardnerella species from the human

vaginal microbiota

Gardnerella strains subjected to whole-genome sequencing were cultivated either from the

Fredricks collection, harboring isolates from human study participants with and without BV in

Seattle, Washington, USA[18,79] or from the Ravel isolate collection, harboring strains from multiple

geographic locations, including USA and Africa (Supplementary Table 2). Bacterial isolates

consistent with *Gardnerella* morphotypes on Gram stain were subjected 16S rRNA gene amplicon

sequencing to confirm phylogenetic placement in the *Gardnerella* genus and *cpn60* gene PCR with

amplicon sequencing to further specify these bacteria[18]. Phylogenetically diverse *Gardnerella*

species were selected for whole-genome sequencing. High molecular weight DNA was extracted

from each isolate using the MasterPure DNA purification kit (Lucigen), with two

phenol/chloroform cleanups prior to DNA precipitation where needed. DNA extraction was

quantified on a TapeStation 2200 instrument run with a Genomic DNA tape (Agilent).

Genome sequencing and assembly

Sequencing on the Pacific Biosciences SeqI/II Platforms

A total of forty-two strains from the Fredricks collection, and seventeen from the Ravel collection,

were selected for single-molecule real-time (SMRT) long-read sequencing using either the PacBio

SeqI or SeqII instrument (Pacific Biosciences) at the Fred Hutchinson Cancer Center or the

Genomic Resource Center of the University Maryland School of Medicine, respectively.

Sequencing libraries were prepared using the SMRTBell Template Prep Kit 1.0 or 2.0 and were

size-selected using a BluePippin (Sage Science) with a cutoff of 3-5 kb. Sequencing reads were

processed via Microbial Assembly software within the Pacific Biosciences' SMRTAnalysis

pipeline (v.9.0.0.92188).

Sequencing on the Illumina HiSeq 4000 Platform

A total of 332 isolates were selected for Illumina sequencing on the HiSeq4000 platform. Short

read libraries were prepared using the KAPA HyperPlus Kit (Kapa Biosystems) with KAPA

Single-Indexed Adapter Kit Set B. A fixed volume (35 µL) of genomic DNA was used as input,

and libraries were prepared following the manufacturer's protocol with modifications based on

their amount of input DNA. For samples with 0.5 or 0.2 ng input DNA, the fragmentation enzyme

was diluted 1:2 or 1:5 with water. All samples were fragmented at 37°C for 5 min. Adapter

concentrations varied according to the input, and the adapter ligation was carried out overnight at

4°C for all samples. The post-ligation cleanup was performed with 0.8x Ampure XP beads

(Beckman Coulter, Indianapolis IN) and 20 µL of sample was used in library amplification. Post-

amplification cleanup was performed with 1x Ampure XP beads; libraries with remaining adapter

dimer peaks were cleaned a second time. The final elution was in 25 µL of nuclease-free water.

Libraries were run on a TapeStation instrument with a D1000 tape (Agilent) to assess quality and

concentration. Libraries were sequenced (8 libraries/lane) on an Illumina HiSeq 4000 instrument

using the 150 bp paired-end protocol. Reads were assembled using Unicycler.

Publicly available genomes from the National Center for Biotechnology Information (NCBI)

We used publicly available genomes both to provide a direct reference for previous studies and to

supplement our own sequencing efforts. We searched GenBank[80] assembly databases for the genus

Gardnerella and family *Bifidobacteraceae* (without genus assignment). Using the Type (Strain)

Genome Server (TYGS)[81] to confirm the genus of these genomes, we discovered that several of

the unassigned *Bifidobacteraceae* bacteria in NCBI were *Gardnerella* and included them in our

analysis. *Bifidobacteraceae* (without genus assignment) were included if they were collected as

part of the same NCBI BioProject which added 20 *Gardnerella* that were not typed further than

the *Bifidobacteraceae* family. Additionally, we included three sequences (350_GVAG,

476_GVAG, and 842_GVAG) that were initially assigned to *Gardnerella* but were later renamed

to *Alloscardovia* to verify this placement. The initial number of potential genomes was n = 779.

To verify the taxonomy of these strains we used the TYGS type strain server[81] to validate the genus

and removed genomes that were assigned outside of *Gardnerella*. Draft genomes with more than

250 contigs were removed due to inherent unreliability and genomes that were modified in the

laboratory were also removed.

Bifidobacterium strains were collected and curated in much the same manner as the

Gardnerella strains. A total of 98 strains from all known species of the *Bifidobacterium* genus and

the closely related *Cardovia* genera—*Alloscardovia, Aeroscardovia*, *Bombiscardovia*,

Pseudoscardovia, and *Scardovia*—were downloaded from NCBI. Whenever possible, the type

strains were collected. Sequences were assessed using CheckM, TYGS, and ANI via cutoffs

similar to those used for the *Gardnerella* strains. *Bifidobacteriaceae* groups were assigned

according to Alessandri *et al.*[82].

Genome quality assessments

The completeness and contamination of all genomes (sequenced or available on NCBI) were

assessed via CheckM[83]. The genomes were annotated via RAST[84]. Genomes with contamination

greater than 1% or completeness less than 90% by CheckM were excluded. Genome sequences

were marked as duplicates if they shared an average nucleotide identity (ANI) greater than 99.5%

with other genomes in the collection (**Supplementary Table 1**). Notes regarding criteria for which

a specific representative was chosen can be found in Supplementary Table 1.

Average nucleotide identity and digital DNA-DNA hybridization comparisons

Average nucleotide identity (ANI) metrics were obtained for Gardnerella Set A and Set B

genomes, separately, using fastANI[41] and visualized using the R packages ComplexHeatmap[85] and

"seriation"[86]. Digital DNA-DNA hybridization (dDDH) [87] values were calculated via the Genome-

to-Genome Distance Calculator[81] and visualized in R.

kSNP4 whole-genome phylogeny

kSNP4[42] was used to construct a reference-free phylogenetic tree for *Gardnerella* Set A (kmer=17,

FCK=0.167) and *Gardnerella* Set B (kmer=17, FCK=0.797) genomes, separately. *Bombiscardovia*

bombi strain DSM19703 was included to root both dendrograms. The resulting parsimony trees

were visualized iTOL v7[88], with GapMind[49-51] metabolism predictions and PADLOC [66,72] defense

mechanism predictions annotated where appropriate.

cpn60 whole-genome phylogeny

RAST-annotated nucleotide sequences for *cpn60* were MUSCLE-aligned and used to generate a

maximum-likelihood dendrogram using MEGA v11[89]. The resulting tree was visualized and

annotated using iTOL v7[88].

PPanGGOLiN Gardnerella pangenome

We used PPanGGOLiN[43] on our curated genome set to graph and cluster the genes from all strains,

find regions of genomic plasticity and synteny[90], and separate the pangenome gene clusters into

core, shell, and cloud partitions, with clustering set to 50% coverage and 50% identity. EggNOG[91],

InterPro[92] and KEGG Mapper[93] were run separately on each genome to gather and confirm the

CAZyme[94], Pfam[95], and KEGG designations of the gene clusters. COG categories[96] were assigned

on the basis of the consensus of these designations. We added the annotation "Unclassified

Metabolism" to capture gene clusters that are clearly metabolism related, but the metabolite is not

defined or could be of multiple metabolic types.

Isolation of pDNF1199S-01 plasmid

Gvr strain DNF1199S was grown in New York City III (NYCIII) liquid media in an AS-580

anaerobic chamber (Anaerobe Systems, Morgan Hill, CA) in a gas mixture of 5% CO_2, 5% H_2,

and 90% N_2 at 37°C. A QIAamp DNA Miniprep kit (Qiagen, Hilden, Germany) was used to isolate

pDNF1199S-01 from a 5mL culture of the bacteria. Isolated DNA was digested with *XhoI* (New

England Biolabs, Ipswich, MA) and visualized by gel electrophoresis to confirm plasmid presence.

***Gardnerella* tetracycline resistance testing**

Gardnerella strains were grown anaerobically in an AS-580 anaerobic chamber (Anaerobe

Systems, Morgan Hill, CA) in a gas mixture of 5% CO_2, 5% H_2, and 90% N_2 at 37°C. Bacteria

were recovered from frozen stocks on NYCIII agar plates and sub-cultured twice in 2mL of

NYCIII liquid media. Culture purity was verified by Gram stain. The final cultures were

normalized to 0.5 McFarland turbidity using fresh NYCIII broth and 25µL was spotted onto an

NYCIII agar plate containing 16µg/mL tetracycline and an antibiotic-free NYCIII agar plate. 25µL

of *L. reuteri* CF48-3A was spotted onto each tetracycline plate as a tetracycline-resistant control[97].

Tetracycline resistance was determined according to Clinical and Laboratory Standards Institute

antimicrobial susceptibility testing protocols and any growth on tetracycline NYCIII agar plates

was considered as resistant to tetracycline.

pMC-pDNF1199S-tetM construction and minicircle preparation

pDNF1199S-01 and *tetM* were both cloned from gDNA isolated from strain DNF1199S. Gene

fragments were assembled with pMC by Gibson assembly[98]. The parental construct was

transformed into *E. coli* ZYCY10P3S2T (System Biosciences, Palo Alto, CA) by heat shocking

according to the SBI Minicircle DNA Technology user manual. Transformant colonies were

selected by growing on Luria-Burtani (LB) agar plates containing 50 µg/mL kanamycin. A

successfully transformed colony was chosen and stocked, then induction of minicircles was

performed according to the SBI Minicircle DNA Technology user manual by growing the bacteria

in induction media containing arabinose. pDNF1199S-*tetM* minicircle DNA was then purified

from the induced *E. coli* using a Qiagen Plasmid Midi Kit (Qiagen, Hilden, Germany). The

sequence of purified plasmid was confirmed against the construct sequence by whole-plasmid

sequencing performed by Plasmidsaurus (Plasmidsaurus, Eugene, OR). "GGCC" sites on the

plasmid were protected by incubating 1μg of the DNA with 1μL of *HaeIII* MTase (New England

Biolabs, Ipswich, MA) at 37°C for 4hrs in the presence of 160μM S-Adenosyl methionine (New

England Biolabs, Ipswich, MA). Following enzyme treatment, DNA was purified with Monarch

PCR & DNA Cleanup Kit (New England Biolabs, Ipswich, MA).

Preparation of competent *Gardnerella* cells

Gvv type strain ATCC 14018 was grown in an AS-580 anaerobic chamber (Anaerobe Systems,

Morgan Hill, CA) in a gas mixture of 5% CO_2, 5% H_2, and 90% N_2 at 37°C. Cells were recovered

from frozen stocks on NYCIII agar plates and subcultured into NYCIII liquid media. 500μL of

culture was subcultured into 5mL NYCIII media and grown overnight. 4mL of this culture was

inoculated into 40mL of modified de Man-Rogosa-Sharpe media[99] supplemented with 5g/L

glucose, 0.4g/L L-cysteine HCl, 100mL/L filter-sterilized horse serum, and 100 mM NaCl when

stated. Once the target OD_{600} was reached, cell were harvested by centrifugation at 4°C at 4,052x*g*

for 10min. Cells were washed in one of the following three ways: 1) twice in 25mL ice-cold

maltose citrate buffer (0.2625g/L citric acid and 0.5M maltose at pH 5.8), 2) twice in 25mL ice-

cold autoclaved 10% glycerol, or 3) once in 25mL ice-cold autoclaved water from a Milli-Q

IQ7000 pure water system (Millipore Sigma, St. Louis, MO, USA) followed by a second wash in

25mL ice-cold maltose citrate buffer. Washed cells were then resuspended in 200μL of maltose

citrate buffer, mixed with 200μL 80% glycerol, and stored at -80°C in aliquots.

Gardnerella electrotransformation and verification

Competent cells were mixed with plasmid DNA on ice and transferred into pre-chilled 0.1cm gap

Gene Pulser/MicroPulser Electroporation Cuvettes (Bio-Rad, Hercules, CA). Electroporation was

then performed on a Gene Pulser Xcell (Bio-Rad, Hercules, CA) at 25µF, 200Ω, and 2,000V.

Immediately following electroporation, 950µL of pre-warmed NYCIII broth was added to the cells

and they were transferred to an AS-580 anaerobic chamber (Anaerobe Systems, Morgan Hill, CA)

in a gas mixture of 5% CO_2, 5% H_2, and 90% N_2 to recover at 37°C for 1 hour. 100µL – 200µL of

transformed cells were then plated on NYCIII agar plates with and without 16 µg/mL tetracycline.

The plates were incubated anaerobically at 37°C for 48 hours and transformed colonies on the

tetracycline plates were counted to determine transformation efficiency. After the first successful

transformation, transformant colonies were grown in 5mL NYCIII media with 16 µg/mL

tetracycline and pDNF1199S-*tetM* was recovered from them using a QIAamp DNA Miniprep kit

(Qiagen, Hilden, Germany). The sequence of the purified plasmid was confirmed by sequencing

performed by Plasmidsaurus (Plasmidsaurus, Eugene, OR). Following transformation of

Gardnerella strain DNF00354 with pDNF1199S-tetM, the transformation reaction was plated onto

NYCIII agar plates with and without 16 µg/mL tetracycline. A small amount of growth from a

transformant colony on the tetracycline plate or lawn growth from the control plate was

resuspended in sterile water and colony PCR was performed to amplify and detect the *tetM* gene.

1µL of resuspended bacteria was added to a reaction mix of 12.5µL GoTaq Green master mix

(Promega, Madison, WI), 0.5µL of each primer, and 6.5µL H_2O. PCR was carried out for 28 cycles

with an annealing temperature of 60°C and an extension time of 2.5 min.

Primer sequences were as follows (5' → 3'):

Forward: CGCGACCAAATATTGGTACATTATTACAGCTATTTTGTAATCACGTACTC

Reverse: GAGACTGTCGTATTGATGTCACGGGCTTAATAGGTCTTATGTCGG

References

1 Fredricks, D. N., Fiedler, T. L. & Marrazzo, J. M. Molecular identification of bacteria
associated with bacterial vaginosis. *N Engl J Med* **353**, 1899-1911 (2005).
https://doi.org/10.1056/NEJMoa043802
2 Amsel, R. *et al.* Nonspecific vaginitis. Diagnostic criteria and microbial and epidemiologic
associations. *Am J Med* **74**, 14-22 (1983). https://doi.org/10.1016/0002-9343(83)91112-9
3 Alcendor, D. J. Evaluation of Health Disparity in Bacterial Vaginosis and the Implications
for HIV-1 Acquisition in African American Women. *Am J Reprod Immunol* **76**, 99-107
(2016). https://doi.org/10.1111/aji.12497
4 Klebanoff, M. A. & Brotman, R. M. Treatment of bacterial vaginosis to prevent preterm
birth. *Lancet* **392**, 2141-2142 (2018). https://doi.org/10.1016/S0140-6736(18)32115-9
5 Berman, H. L., Goltsman, D. S. A., Anderson, M., Relman, D. A. & Callahan, B. J.
Gardnerella diversity and ecology in pregnancy and preterm birth. *mSystems* **9**, e0133923
(2024). https://doi.org/10.1128/msystems.01339-23
6 Fredricks, D. N., Fiedler, T. L., Thomas, K. K., Mitchell, C. M. & Marrazzo, J. M. Changes
in vaginal bacterial concentrations with intravaginal metronidazole therapy for bacterial
vaginosis as assessed by quantitative PCR. *J Clin Microbiol* **47**, 721-726 (2009).
https://doi.org/10.1128/JCM.01384-08
7 Fredricks, D. N., Fiedler, T. L., Thomas, K. K., Oakley, B. B. & Marrazzo, J. M. Targeted
PCR for detection of vaginal bacteria associated with bacterial vaginosis. *J Clin Microbiol*
45, 3270-3276 (2007). https://doi.org/10.1128/JCM.01272-07
8 Hummelen, R. *et al.* Deep sequencing of the vaginal microbiota of women with HIV. *PLoS*
One **5**, e12078 (2010). https://doi.org/10.1371/journal.pone.0012078
9 Ravel, J. *et al.* Vaginal microbiome of reproductive-age women. *Proc Natl Acad Sci U S A*
108 Suppl 1, 4680-4687 (2011). https://doi.org/10.1073/pnas.1002611107
10 Shipitsyna, E. *et al.* Composition of the vaginal microbiota in women of reproductive age-
-sensitive and specific molecular diagnosis of bacterial vaginosis is possible? *PLoS One* **8**,
e60670 (2013). https://doi.org/10.1371/journal.pone.0060670
11 Srinivasan, S. *et al.* Bacterial communities in women with bacterial vaginosis: high
resolution phylogenetic analyses reveal relationships of microbiota to clinical criteria.
PLoS One **7**, e37818 (2012). https://doi.org/10.1371/journal.pone.0037818
12 van de Wijgert, J. H. *et al.* The vaginal microbiota: what have we learned after a decade of
molecular characterization? *PLoS One* **9**, e105998 (2014).
https://doi.org/10.1371/journal.pone.0105998
13 Gardner, H. L. & Dukes, C. D. Haemophilus vaginalis vaginitis: a newly defined specific
infection previously classified non-specific vaginitis. *Am J Obstet Gynecol* **69**, 962-976
(1955).
14 Schwebke, J. R., Muzny, C. A. & Josey, W. E. Role of *Gardnerella vaginalis* in the
pathogenesis of bacterial vaginosis: a conceptual model. *J Infect Dis* **210**, 338-343 (2014).
https://doi.org/10.1093/infdis/jiu089
15 Patterson, J. L., Stull-Lane, A., Girerd, P. H. & Jefferson, K. K. Analysis of adherence,
biofilm formation and cytotoxicity suggests a greater virulence potential of *Gardnerella*
vaginalis relative to other bacterial-vaginosis-associated anaerobes. *Microbiology*
(Reading) **156**, 392-399 (2010). https://doi.org/10.1099/mic.0.034280-0

627 16 Gelber, S. E., Aguilar, J. L., Lewis, K. L. & Ratner, A. J. Functional and phylogenetic
628   characterization of Vaginolysin, the human-specific cytolysin from *Gardnerella vaginalis*.
629   *J Bacteriol* **190**, 3896-3903 (2008). https://doi.org/10.1128/JB.01965-07

630 17 Castro, J. *et al.* Using an in-vitro biofilm model to assess the virulence potential of bacterial
631   vaginosis or non-bacterial vaginosis *Gardnerella vaginalis* isolates. *Sci Rep* **5**, 11640
632   (2015). https://doi.org/10.1038/srep11640

633 18 Munch, M. M. *et al. Gardnerella* Species and Their Association With Bacterial Vaginosis.
634   *J Infect Dis* **230**, e171-e181 (2024). https://doi.org/10.1093/infdis/jiae026

635 19 Muzny, C. A. *et al.* An Updated Conceptual Model on the Pathogenesis of Bacterial
636   Vaginosis. *J Infect Dis* **220**, 1399-1405 (2019). https://doi.org/10.1093/infdis/jiz342

637 20 Janulaitiene, M. *et al.* Prevalence and distribution of Gardnerella vaginalis subgroups in
638   women with and without bacterial vaginosis. *BMC Infect Dis* **17**, 394 (2017).
639   https://doi.org/10.1186/s12879-017-2501-y

640 21 Balashov, S. V., Mordechai, E., Adelson, M. E. & Gygax, S. E. Identification,
641   quantification and subtyping of *Gardnerella vaginalis* in noncultured clinical vaginal
642   samples by quantitative PCR. *J Med Microbiol* **63**, 162-175 (2014).
643   https://doi.org/10.1099/jmm.0.066407-0

644 22 Shvartsman, E., Hill, J. E., Sandstrom, P. & MacDonald, K. S. *Gardnerella* Revisited:
645   Species Heterogeneity, Virulence Factors, Mucosal Immune Responses, and Contributions
646   to Bacterial Vaginosis. *Infect Immun* **91**, e0039022 (2023).
647   https://doi.org/10.1128/iai.00390-22

648 23 Piot, P. *et al.* Biotypes of *Gardnerella vaginalis*. *J Clin Microbiol* **20**, 677-679 (1984).
649   https://doi.org/10.1128/jcm.20.4.677-679.1984

650 24 Benito, R., Vazquez, J. A., Berron, S., Fenoll, A. & Saez-Neito, J. A. A modified scheme
651   for biotyping *Gardnerella vaginalis*. *J Med Microbiol* **21**, 357-359 (1986).
652   https://doi.org/10.1099/00222615-21-4-357

653 25 Nath, K., Devlin, D. & Beddoe, A. M. Heterogeneity in restriction patterns of *Gardnerella*
654   *vaginalis* isolates from individuals with bacterial vaginosis. *Res Microbiol* **143**, 199-209
655   (1992). https://doi.org/10.1016/0923-2508(92)90009-d

656 26 Ingianni, A., Petruzzelli, S., Morandotti, G. & Pompei, R. Genotypic differentiation of
657   *Gardnerella vaginalis* by amplified ribosomal DNA restriction analysis (ARDRA). *FEMS*
658   *Immunol Med Microbiol* **18**, 61-66 (1997). https://doi.org/10.1111/j.1574-
659   695X.1997.tb01028.x

660 27 Ahmed, A. *et al.* Comparative genomic analyses of 17 clinical isolates of *Gardnerella*
661   *vaginalis* provide evidence of multiple genetically isolated clades consistent with
662   subspeciation into genovars. *J Bacteriol* **194**, 3922-3937 (2012).
663   https://doi.org/10.1128/JB.00056-12

664 28 Cornejo, O. E., Hickey, R. J., Suzuki, H. & Forney, L. J. Focusing the diversity of
665   Gardnerella vaginalis through the lens of ecotypes. *Evol Appl* **11**, 312-324 (2018).
666   https://doi.org/10.1111/eva.12555

667 29 Vaneechoutte, M. *et al.* Emended description of *Gardnerella vaginalis* and description of
668   *Gardnerella leopoldii* sp. nov., *Gardnerella piotii* sp. nov. and *Gardnerella swidsinskii* sp.
669   nov., with delineation of 13 genomic species within the genus *Gardnerella*. *Int J Syst Evol*
670   *Microbiol* **69**, 679-687 (2019). https://doi.org/10.1099/ijsem.0.003200

Potter, R. F., Burnham, C. D. & Dantas, G. In Silico Analysis of *Gardnerella* Genomospecies Detected in the Setting of Bacterial Vaginosis. *Clin Chem* **65**, 1375-1387 (2019). https://doi.org/10.1373/clinchem.2019.305474

Innamorati, K. A. *et al.* Metronidazole response profiles of Gardnerella species are congruent with phylogenetic and comparative genomic analyses. *Genome Med* **17**, 28 (2025). https://doi.org/10.1186/s13073-025-01446-4

Links, M. G., Dumonceaux, T. J., Hemmingsen, S. M. & Hill, J. E. The chaperonin-60 universal target is a barcode for bacteria that enables de novo assembly of metagenomic sequence data. *PLoS One* **7**, e49755 (2012). https://doi.org/10.1371/journal.pone.0049755

Paramel Jayaprakash, T., Schellenberg, J. J. & Hill, J. E. Resolution and characterization of distinct cpn60-based subgroups of *Gardnerella vaginalis* in the vaginal microbiota. *PLoS One* **7**, e43009 (2012). https://doi.org/10.1371/journal.pone.0043009

Zhang, G., Gao, B., Adeolu, M., Khadka, B. & Gupta, R. S. Phylogenomic Analyses and Comparative Studies on Genomes of the Bifidobacteriales: Identification of Molecular Signatures Specific for the Order Bifidobacteriales and Its Different Subclades. *Front Microbiol* **7**, 978 (2016). https://doi.org/10.3389/fmicb.2016.00978

Lugli, G. A. *et al.* Comparative genomic and phylogenomic analyses of the Bifidobacteriaceae family. *BMC Genomics* **18**, 568 (2017). https://doi.org/10.1186/s12864-017-3955-4

Tarracchini, C. *et al.* Assessing the Genomic Variability of Gardnerella vaginalis through Comparative Genomic Analyses: Evolutionary and Ecological Implications. *Appl Environ Microbiol* **87** (2020). https://doi.org/10.1128/AEM.02188-20

Bohr, L. L., Mortimer, T. D. & Pepperell, C. S. Lateral Gene Transfer Shapes Diversity of *Gardnerella* spp. *Front Cell Infect Microbiol* **10**, 293 (2020). https://doi.org/10.3389/fcimb.2020.00293

Schellenberg, J. J. *et al. Gardnerella vaginalis* Subgroups Defined by cpn60 Sequencing and Sialidase Activity in Isolates from Canada, Belgium and Kenya. *PLoS One* **11**, e0146510 (2016). https://doi.org/10.1371/journal.pone.0146510

Hill, J. E., Albert, A. Y. K. & Group, V. R. Resolution and Cooccurrence Patterns of *Gardnerella leopoldii, G. swidsinskii, G. piotii,* and *G. vaginalis* within the Vaginal Microbiome. *Infect Immun* **87** (2019). https://doi.org/10.1128/IAI.00532-19

Sousa, M. *et al. Gardnerella pickettii* sp. nov. (formerly *Gardnerella* genomic species 3) and *Gardnerella greenwoodii* sp. nov. (formerly *Gardnerella* genomic species 8) isolated from female urinary microbiome. *Int J Syst Evol Microbiol* **73** (2023). https://doi.org/10.1099/ijsem.0.006140

Jain, C., Rodriguez, R. L., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* **9**, 5114 (2018). https://doi.org/10.1038/s41467-018-07641-9

Hall, B. G. & Nisbet, J. Building Phylogenetic Trees From Genome Sequences With kSNP4. *Mol Biol Evol* **40** (2023). https://doi.org/10.1093/molbev/msad235

Gautreau, G. *et al.* PPanGGOLiN: Depicting microbial diversity via a partitioned pangenome graph. *PLoS Comput Biol* **16**, e1007732 (2020). https://doi.org/10.1371/journal.pcbi.1007732

Flores Ramos, S. *et al.* Genomic Stability and Genetic Defense Systems in *Dolosigranulum pigrum*, a Candidate Beneficial Bacterium from the Human Microbiome. *mSystems* **6**, e0042521 (2021). https://doi.org/10.1128/mSystems.00425-21

Dillard, L. R., Glass, E. M., Lewis, A. L., Thomas-White, K. & Papin, J. A. Metabolic Network Models of the Gardnerella Pangenome Identify Key Interactions with the Vaginal Environment. *mSystems* **8**, e0068922 (2023). https://doi.org/10.1128/msystems.00689-22

Teixeira, P., Sousa, M., Gonçalves Ribeiro, T., Grosso F., Peixe L. Unraveling the *Gardnerella* Pangenome: Insights into Genetic Diversity, Virulence, and Antibiotic Resistance. *bioRxiv* **2025.03.03.641253** (2025). https://doi.org/doi: https://doi.org/10.1101/2025.03.03.641253

Chen, L., Li, J. & Xiao, B. The role of sialidases in the pathogenesis of bacterial vaginosis and their use as a promising pharmacological target in bacterial vaginosis. *Front Cell Infect Microbiol* **14**, 1367233 (2024). https://doi.org/10.3389/fcimb.2024.1367233

Rottini, G. *et al.* Identification and partial characterization of a cytolytic toxin produced by *Gardnerella vaginalis*. *Infect Immun* **58**, 3751-3758 (1990). https://doi.org/10.1128/iai.58.11.3751-3758.1990

Price, M. N., Shiver A.L., Day, L.A., Torres, M., Lesea, H.P., Owens, T.K., Deutschbauer, A.M., Arkin, A.P. . Improving the annotation of amino acid biosynthesis pathways: GapMind 2024. *Biorxiv* **2024.10.14.618325** (2024). https://doi.org/https://doi.org/10.1101/2024.10.14.618325

Price, M. N., Deutschbauer, A. M. & Arkin, A. P. GapMind: Automated Annotation of Amino Acid Biosynthesis. *mSystems* **5** (2020). https://doi.org/10.1128/mSystems.00291-20

Price, M. N., Deutschbauer, A. M. & Arkin, A. P. Filling gaps in bacterial catabolic pathways with computation and high-throughput genetics. *PLoS Genet* **18**, e1010156 (2022). https://doi.org/10.1371/journal.pgen.1010156

Bordigoni, A., Bouchard, S. & Desnues, C. Isolation and Characterization of Gardnerella Phage vB_Gva_AB1, a Bacteriophage Infecting a Clinical Strain of *Gardnerella vaginalis*. *Microbiol Resour Announc* **10** (2021). https://doi.org/10.1128/MRA.00053-21

Malki, K. *et al.* Genomes of *Gardnerella* Strains Reveal an Abundance of Prophages within the Bladder Microbiome. *PLoS One* **11**, e0166757 (2016). https://doi.org/10.1371/journal.pone.0166757

Georjon, H. & Bernheim, A. The highly diverse antiphage defence systems of bacteria. *Nat Rev Microbiol* **21**, 686-700 (2023). https://doi.org/10.1038/s41579-023-00934-x

Tesson, F. *et al.* Systematic and quantitative view of the antiviral arsenal of prokaryotes. *Nat Commun* **13**, 2561 (2022). https://doi.org/10.1038/s41467-022-30269-9

Tal, N. *et al.* Bacteria deplete deoxynucleotides to defend against bacteriophage infection. *Nat Microbiol* **7**, 1200-1209 (2022). https://doi.org/10.1038/s41564-022-01158-0

McLandsborough, L. A., Kolaetis, K. M., Requena, T. & McKay, L. L. Cloning and characterization of the abortive infection genetic determinant abiD isolated from pBF61 of Lactococcus lactis subsp. lactis KR5. *Appl Environ Microbiol* **61**, 2023-2026 (1995). https://doi.org/10.1128/aem.61.5.2023-2026.1995

Deng, Y. M., Liu, C. Q. & Dunn, N. W. Genetic organization and functional analysis of a novel phage abortive infection system, AbiL, from Lactococcus lactis. *J Biotechnol* **67**, 135-149 (1999). https://doi.org/10.1016/s0168-1656(98)00175-8

Cohen, D. *et al.* Cyclic GMP-AMP signalling protects bacteria against viral infection. *Nature* **574**, 691-695 (2019). https://doi.org/10.1038/s41586-019-1605-5

Millman, A., Melamed, S., Amitai, G. & Sorek, R. Diversity and classification of cyclic-oligonucleotide-based anti-phage signalling systems. *Nat Microbiol* **5**, 1608-1615 (2020). https://doi.org/10.1038/s41564-020-0777-y

Gao, L. *et al.* Diverse enzymatic activities mediate antiviral immunity in prokaryotes. *Science* **369**, 1077-1084 (2020). https://doi.org/10.1126/science.aba0372

Mestre, M. R. *et al.* UG/Abi: a highly diverse family of prokaryotic reverse transcriptases associated with defense functions. *Nucleic Acids Res* **50**, 6084-6101 (2022). https://doi.org/10.1093/nar/gkac467

Dy, R. L., Przybilski, R., Semeijn, K., Salmond, G. P. & Fineran, P. C. A widespread bacteriophage abortive infection system functions through a Type IV toxin-antitoxin mechanism. *Nucleic Acids Res* **42**, 4590-4605 (2014). https://doi.org/10.1093/nar/gkt1419

Doron, S. *et al.* Systematic discovery of antiphage defense systems in the microbial pangenome. *Science* **359** (2018). https://doi.org/10.1126/science.aar4120

Cheng, R. *et al.* A nucleotide-sensing endonuclease from the Gabija bacterial defense system. *Nucleic Acids Res* **49**, 5216-5229 (2021). https://doi.org/10.1093/nar/gkab277

Payne, L. J. *et al.* Identification and classification of antiviral defence systems in bacteria and archaea with PADLOC reveals new system types. *Nucleic Acids Res* **49**, 10868-10878 (2021). https://doi.org/10.1093/nar/gkab883

Millman, A. *et al.* An expanded arsenal of immune systems that protect bacteria from phages. *Cell Host Microbe* **30**, 1556-1569 e1555 (2022). https://doi.org/10.1016/j.chom.2022.09.017

Johnston, C. D. *et al.* Systematic evasion of the restriction-modification barrier in bacteria. *Proc Natl Acad Sci U S A* **116**, 11454-11459 (2019). https://doi.org/10.1073/pnas.1820256116

Finn, M. B., Ramsey, K. M., Tolliver, H. J., Dove, S. L. & Wessels, M. R. Improved transformation efficiency of group A Streptococcus by inactivation of a type I restriction modification system. *PLoS One* **16**, e0248201 (2021). https://doi.org/10.1371/journal.pone.0248201

Hu, S. *et al.* Altering under-represented DNA sequences elevates bacterial transformation efficiency. *mBio* **14**, e0210523 (2023). https://doi.org/10.1128/mbio.02105-23

Chen, C. *et al.* Development of a Pre-Modification Strategy to Overcome Restriction-Modification Barriers and Enhance Genetic Engineering in Lactococcus lactis for Nisin Biosynthesis. *Int J Mol Sci* **26** (2025). https://doi.org/10.3390/ijms26052200

Payne, L. J. *et al.* PADLOC: a web server for the identification of antiviral defence systems in microbial genomes. *Nucleic Acids Res* **50**, W541-W550 (2022). https://doi.org/10.1093/nar/gkac400

Klimowicz, A. K., Garcia, E.M., Jefferson, K.K., Dillard, J.P. Genetic transformation of *Gardnerella* species and characterization of vaginolysin and sialidase mutants. *bioRxiv* **2025.05.09.653190** (2025). https://doi.org/https://doi.org/10.1101/2025.05.09.653190

Kularatne, B. & Hill, J. E. Transformation of Gardnerella vaginalis with a Bifidobacterium-Escherichia coli shuttle vector plasmid. *Microbiol Spectr* **13**, e0048125 (2025). https://doi.org/10.1128/spectrum.00481-25

Zhu, M. *et al.* Vaginal *Lactobacillus* fatty acid response mechanisms reveal a novel strategy for bacterial vaginosis treatment. *bioRxiv* (2023). https://doi.org/10.1101/2023.12.30.573720

76   Khan, S., Vancuren, S. J. & Hill, J. E. A Generalist Lifestyle Allows Rare *Gardnerella* spp.
to Persist at Low Levels in the Vaginal Microbiome. *Microb Ecol* **82**, 1048-1060 (2021).
https://doi.org/10.1007/s00248-020-01643-1
77   Kharsany, A. B., Hoosen, A. A. & Van den Ende, J. Antimicrobial susceptibilities of
Gardnerella vaginalis. *Antimicrob Agents Chemother* **37**, 2733-2735 (1993).
https://doi.org/10.1128/AAC.37.12.2733
78   Bulavaite, A., Dalgediene, I., Michailoviene, V. & Pleckaityte, M. Type II Restriction-
Modification System from Gardnerella vaginalis ATCC 14018. *Pathogens* **9** (2020).
https://doi.org/10.3390/pathogens9090703
79   Srinivasan, S. *et al.* More Easily Cultivated Than Identified: Classical Isolation With
Molecular Identification of Vaginal Bacteria. *J Infect Dis* **214 Suppl 1**, S21-28 (2016).
https://doi.org/10.1093/infdis/jiw192
80   Clark, K., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J. & Sayers, E. W. GenBank. *Nucleic
Acids Res* **44**, D67-72 (2016). https://doi.org/10.1093/nar/gkv1276
81   Meier-Kolthoff, J. P. & Goker, M. TYGS is an automated high-throughput platform for
state-of-the-art genome-based taxonomy. *Nat Commun* **10**, 2182 (2019).
https://doi.org/10.1038/s41467-019-10210-3
82   Alessandri, G., van Sinderen, D. & Ventura, M. The genus bifidobacterium: From
genomics to functionality of an important component of the mammalian gut microbiota
running title: Bifidobacterial adaptation to and interaction with the host. *Comput Struct
Biotechnol J* **19**, 1472-1487 (2021). https://doi.org/10.1016/j.csbj.2021.03.006
83   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* **25**, 1043-1055 (2015). https://doi.org/10.1101/gr.186072.114
84   Aziz, R. K. *et al.* The RAST Server: rapid annotations using subsystems technology. *BMC
Genomics* **9**, 75 (2008). https://doi.org/10.1186/1471-2164-9-75
85   Gu, Z. Complex heatmap visualization. *Imeta* **1**, e43 (2022).
https://doi.org/10.1002/imt2.43
86   Hahsler, M., Hornik, K., Buchta, C. . Getting Things in Order: An Introduction to the R
Package Seriation. *Journal of Statistical Software* **25(3),1-34** (2008).
https://doi.org/doi.org/10.18637/jss.v025.i03
87   Auch, A. F., von Jan, M., Klenk, H. P. & Goker, M. Digital DNA-DNA hybridization for
microbial species delineation by means of genome-to-genome sequence comparison. *Stand
Genomic Sci* **2**, 117-134 (2010). https://doi.org/10.4056/sigs.531120
88   Letunic, I. & Bork, P. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic
tree display and annotation tool. *Nucleic Acids Res* **52**, W78-W82 (2024).
https://doi.org/10.1093/nar/gkae268
89   Tamura, K., Stecher, G. & Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis
Version 11. *Mol Biol Evol* **38**, 3022-3027 (2021). https://doi.org/10.1093/molbev/msab120
90   Bazin, A., Gautreau, G., Medigue, C., Vallenet, D. & Calteau, A. panRGP: a pangenome-
based method to predict genomic islands and explore their diversity. *Bioinformatics* **36**,
i651-i658 (2020). https://doi.org/10.1093/bioinformatics/btaa792
91   Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and phylogenetically
annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids
Res* **47**, D309-D314 (2019). https://doi.org/10.1093/nar/gky1085

92 Blum, M. *et al.* InterPro: the protein sequence classification resource in 2025. *Nucleic*
Acids Res **53**, D444-D456 (2025). https://doi.org/10.1093/nar/gkae1082
93 Kanehisa, M., Furumichi, M., Sato, Y., Matsuura, Y. & Ishiguro-Watanabe, M. KEGG:
biological systems database as a model of the real world. *Nucleic Acids Res* **53**, D672-
D677 (2025). https://doi.org/10.1093/nar/gkae909
94 Drula, E. *et al.* The carbohydrate-active enzyme database: functions and literature. *Nucleic*
Acids Res **50**, D571-D577 (2022). https://doi.org/10.1093/nar/gkab1045
95 Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res* **49**, D412-
D419 (2021). https://doi.org/10.1093/nar/gkaa913
96 Galperin, M. Y. *et al.* COG database update: focus on microbial diversity, model organisms,
and widespread pathogens. *Nucleic Acids Res* **49**, D274-D281 (2021).
https://doi.org/10.1093/nar/gkaa1018
97 Egervarn, M., Roos, S. & Lindmark, H. Identification and characterization of antibiotic
resistance genes in *Lactobacillus reuteri* and *Lactobacillus plantarum*. *J Appl Microbiol*
107, 1658-1668 (2009). https://doi.org/10.1111/j.1365-2672.2009.04352.x
98 Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred
kilobases. *Nat Methods* **6**, 343-345 (2009). https://doi.org/10.1038/nmeth.1318
99 Douwe van Sinderen, M. V. *Bifidobacteria: Methods and Protocols*. 1 edn, (Humana,
2021).
870