Immunological characterization of peritoneal exudate cells in liver cirrhosis

patients

Authors:

Shiori Kaji, MD[1], Izumi Sasaki, PhD[2,*], Satoko Kunimoto, MD[3], Naoko Wakaki-Nishiyama, BS[2],

Emi Tosuji, MD[4], Ayano Kurara, HSG[2], Takashi Kato, MD, PhD[2,5], Chisako Mutsukawa, BS[2],

Daisuke Okuzaki, PhD[6], Chizuyo Okamoto, BS[2], Yukiko Yamano, MD[3], Ryo Shimizu, MD[1],

Shuya Maeshima, MD[1], Yoshiyuki Ida, MD, PhD[1], Yasunobu Yamashita, MD, PhD[1], Tadamichi

Hashimoto, MD, PhD[7], Sadahiro Iwabuchi, PhD[8], Shinichi Hashimoto, PhD[8], Shin-Ichiroh Saitoh,

PhD[9], Shin-ichi Araki, MD, PhD[3], Masayuki Kitano, MD, PhD[1,*] and Tsuneyasu Kaisho, MD,

PhD[2,*]

Affiliations:

[1] Second Department of Internal Medicine, Wakayama Medical University, Wakayama, Japan.

[2] Department of Immunology, Institute of Advanced Medicine, Wakayama Medical University,

Wakayama, Japan.

[3] Department of Nephrology, Wakayama Medical University, Wakayama, Japan.

[4] Department of Dermatology, Wakayama Medical University, Wakayama, Japan.

[5] Department of Rheumatology and Clinical Immunology, Wakayama Medical University,

Wakayama, Japan.

[6] Laboratory for Human Immunology (Single Cell Genomics), WPI Immunology Frontier

Research Center, Osaka University, Suita-City, Osaka, Japan.

[7] Medical Corporation Hashimoto Hospital, Wakayama, Japan.

[8] Department of Molecular Pathophysiology, Institute of Advanced Medicine, Wakayama Medical

University, Wakayama, Japan.

[9] Department of Intractable Disorders, Institute of Advanced Medicine, Wakayama Medical

University, Wakayama, Japan.

Address correspondence:

*Corresponding author. Email: izumisas@wakayama-med.ac.jp (I.S); kitano@wakayama-

med.ac.jp (M.K) ; tkaisho@wakayama-med.ac.jp (T.Kai.)

Conflicts of interest: nothing to report.

Financial support: This work was supported in part by 2023 Wakayama Medical Award for Young

Researchers (to S.K.). This work was also supported by Grant-in-Aid for Transformative Research

Areas (JP22H05182 and JP22H05187 to T. Kai. and JP22H05187 to I.S.), for Scientific Research

(B) (JP26293106, JP17H04088, JP20H03505 and JP24K02298 to T. Kai.), for Scientific Research

(C) (JP19K07628 and JP22K07006 to I.S.), for Scientific Research on Innovative Areas

(JP17H05799 and JP19H04813 to T. Kai.), for Exploratory Research (JP17K19568, JP21K19384

and JP23K18222 to T. Kai.), for Research Activity Start-up (JP23K19487 to T.Kat) from the Japan

Society for the Promotion of Science, Takeda Science Foundation (to T. Kai. and I.S.), GSK Japan

Research Grant 2021 (to I.S.), Kowa Life Science Foundation (to I.S.).

Keywords: continuous ambulatory peritoneal dialysis, dendritic cells, GATA6-expressing

macrophages, mucosal-associated invariant T cells, Child-Pugh classification

Abbreviations: LC, liver cirrhosis; MAIT cells, mucosal-associated invariant T cells; PECs,

peritoneal exudate cells; CAPD, continuous ambulatory peritoneal dialysis; DCs, dendritic cells;

scRNA-seq, single-cell RNA sequencing; UMAP, Uniform Manifold Approximation and

Projection; KEGG, Kyoto Encyclopedia of Genes and Genomes; CP-B, Child-Pugh class B; bulk

RNA-seq, bulk RNA sequencing; CP, Child-Pugh; CP-C, Child-Pugh class C; *GATA6*[+] MPs,

GATA6-expressing macrophages; *FCGR3A*[+] MPs, *FCGR3A*-expressing macrophages.

ABSTRACT

BACKGROUND AND AIMS: Liver cirrhosis (LC) is the end stage of liver fibrosis caused by various chronic liver diseases. Patients with LC often develop ascites containing peritoneal exudate cells (PECs). However, those cells have not been fully immunologically characterized. In this study, we clarify immune cell profiles of PECs from patients with LC.

APPROACH AND RESULTS: PECs were collected from patients with LC or patients receiving continuous ambulatory peritoneal dialysis (CAPD) as a non-cirrhotic control and subjected to single-cell RNA sequencing (scRNA-seq), bulk RNA sequencing (RNA-seq) and flowcytometry analyses. Analysis of scRNA-seq revealed that dendritic cells (DCs) and macrophages were major populations in CAPD patient-derived PECs, while those cells were decreased and T cells were most abundant in LC patient-derived PECs. Notably, *FCGR3A*-expressing macrophages were dominant over DCs and *GATA6*-expressing macrophages in LC patient-derived PECs. Bulk RNA-seq analysis further clarified expression of a set of genes was up- or down-regulated along with LC severity. Especially, expression of T cell signature genes was featured by its increase at Child-Pugh class B, but decrease at Child-Pugh class C. Flowcytometry analysis showed increase of T cells, decrease of DCs and macrophages, and increased expression of CD16 and CD163 in CD1c^{int}CD14^{high} cells corresponding to *FCGR3A*-expressing macrophages in LC patient-derived PECs.

CONCLUSIONS: In LC patient-derived PECs, myeloid and T cell populations and their gene expression profiles were fluctuated with severity. Our findings should contribute to further development of diagnosis or therapeutic maneuver for LC.

INTRODUCTION

Liver cirrhosis (LC) is an end stage of liver fibrosis and causes more than one million deaths

every year all over the world.[1] As liver fibrosis progresses, patients often develop ascites, mainly

due to portal hypertension. A variety of immune cells can be found in the ascites and they change

their profiles, depending on the pathological state of the patients. For example, in patients with

LC, CD8[+] mucosal-associated invariant T (MAIT) cells and CD16[high] CD163[high] macrophages are

increased in the ascites but not in peripheral bloods.[2] In mice, peritoneal cavity macrophages

migrate into the liver upon liver injury,[3] and regulate the pathological conditions in the liver.[4,5]

These studies indicate peritoneal cavity cells should have critical roles in liver pathogenesis.

However, it is unclear whether or how various types of immune cells fluctuate in the ascites of

patients with LC.

In this study, we analyzed the peritoneal exudate cells (PECs) of patients with LC in comparison

with those of patients undergoing continuous ambulatory peritoneal dialysis (CAPD). We then

found increase in frequency of T cells and macrophage subsets and decrease in frequency of

dendritic cells (DCs) in the ascites of LC patients. Furthermore, we have also revealed some

parameters correlating with disease severity.

MATERIALS AND METHODS

Patients

Patients with LC and patients with end-stage renal disease undergoing CAPD at Wakayama

Medical University were eligible for this study (Supplementary Table S1-3). Written informed

consent was obtained from all patients, and the study protocol conforms to the ethical guidelines

of the 2013 Declaration of Helsinki and the 2018 Declaration of Istanbul. All studies were

performed in accordance with appropriate ethical regulations and were pre-approved by the

Wakayama Medical University Research Ethics Committee (#3918).

Preparation of peritoneal exudate cells

PECs from patients with LC were collected from ascites by puncture and drainage at the

diagnosis or treatment. PECs from patients undergoing CAPD were collected from 1.5-2 L of

CAPD effluent according to the standard CAPD prescription. Ascites of patients with LC or

CAPD effluent were centrifuged at 4℃ 2000 rpm 5 min. Cell pellets were washed three times

with phosphate buffered saline (PBS) and resuspended in PBS supplemented with 1% fetal calf

serum (FCS), 5 mM ethylenediaminetetraacetic acid (EDTA) and 0.01% sodium azide (NaN_3).

Turk's solution was used to analyze cell viability to ensure single-cell suspension with more

than 90% viability.

Single-cell RNA sequencing analysis

Single-cell RNA sequencing (scRNA-seq) analysis was performed using Chromium Controller

(10x Genomics, Pleasanton, CA). Libraries were sequenced on NovaSeq 6000 (Illumina, San

Diego, CA) in a 101+101 base paired-end mode. Illumina RTA v3.4.4 software was used for

base calling. Generated reads were input into the cellranger pipeline (cellranger-7.1.0, 10x

Genomics) for UMI counting for each gene-cell (barcode) combination. The cluster analysis

based on Uniform Manifold Approximation and Projection (UMAP) plot and differential

expression analysis were performed by Loupe Browser (v7.0.1, 10x Genomics).

Bulk RNA sequencing analysis

Total RNA was extracted with RNeasy micro kit (QIAGEN, Hilden, Germany). Full-length

cDNA was prepared using a SMART-Seq HT Kit (Takara Bio, Japan). According to the

SMART-Seq kit instructions, an Illumina library was then prepared using a NexteraXT DNA

Library Preparation Kit (Illumina). Sequencing was performed on NovaSeq 6000 platform in a

101-base single-end mode. Illumina RTA v3.4.4 software was used for base calling. Generated

reads were mapped to the human (hg19) reference genome using TopHat v2.1.1 in combination

with Bowtie2 ver. 2.2.8 and SAMtools ver. 0.1.18. Fragments per kilobase of exon per million

mapped fragments (FPKMs) were calculated using Cuffdiff 2.2.1 with parameter -max-bundle-

frags 50000000. The DAVID Bioinformatics database and Kyoto Encyclopedia of Genes and

Genomes (KEGG) annotations were used for enrichment analysis. The web tools ClustVis and

VolcaNoseR[6] were used to draw heatmaps and volcano plots, respectively.

Flowcytometry analysis

PECs were collected in Eppendorf tubes from patients undergoing CAPD or patients with LC,

and subjected to flowcytometry analysis. PECs were stained with the antibodies listed in

Supplementary Table S4. Dead cells detected as positive for AmCyan channels were excluded

using the LIVE/DEAD Fixable Dead Cell Aqua Stain Kit (cat; L34966, Invitrogen, Waltham,

MA). Stained cells were analyzed with FACS Verse and data were processed with FlowJo

Version 10.8.1 software (TreeStar, Ashland, OR).

Magnetic beads cell isolation

$CD1c^{high}$ $CD14^{int}$, $CD1c^{int}$ $CD14^{high}$ or $CD1c^{low-int}$ $CD14^{low}$ cells were isolated from PECs by

using magnetic-activated cell sorting (MACS). PECs were stained with biotinylated anti-CD1c

antibody for 20 min on ice. After washing, stained PECs were applied to streptavidin microbeads

(Miltenyi Biotec, Bergisch Gladbach, Germany) and then magnetically sorted with LS columns

as per the manufacturer's instructions. Streptavidin microbeads-conjugated cells were used as

$CD1c^{high}$ $CD14^{int}$ cells. The flow-through cells including $CD1c^{int}$ $CD14^{high}$ and $CD1c^{low-int}$ $CD14^{low}$

cells were stained with PE-conjugated anti-CD14 antibody for 20 min on ice. After washing, cells

were applied to PE microbeads (Miltenyvi Biotec) and then magnetically sorted with LS columns

as per the manufacturer's instructions. PE microbeads-conjugated cells were used as CD1c^{int}

CD14^{high} cells and the flow-through cells were used as CD1c$^{low-int}$ CD14^{low} cells. Purity of

CD1c^{high} CD14^{int}, CD1c^{int} CD14^{high} or CD1c$^{low-int}$ CD14^{low} cells were verified by flowcytometry

analysis.

Quantitative real-time PCR

Using PrimeScript RT Reagent Kit (TaKaRa, Kusatsu, Japan), 0.5 μg of total RNA was reverse

transcribed into complementary DNA. Relative expression of RNA transcripts was quantified

using gene-specific primers, TB Green Premix Ex Taq II (TaKaRa) and the StepOnePlus Real-

Time PCR System (Applied Biosystems, Waltham, MA). TaqMan probes (TaqMan Gene

Expression Assay, Applied Biosystems) were used for 18S rRNA (internal control). Expression

of all genes was normalized to that of 18S rRNA and is represented as the ratio to the indicated

reference samples. Gene-specific primers used in the study were obtained from FASMAC, and

all primers were validated for linear amplification. The primer sequences were designed and

synthesized by FASMAC as follows: *GATA6*, forward primer: 5'-

GCCACTACCTGTGCAACGCCT-3', reverse primer: 5'-CAATCCAAGCCGCCGTGATGAA-

3'.

Statistical analysis

Statistical significance was determined by an unpaired two-tailed Student's t test. *p* values are

indicated by *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. $p < 0.05$ was considered

statistically significant.

Data availability statement

Bulk RNA sequencing (bulk RNA-seq) and scRNA-seq data generated in this study have been

deposited at GEO under accession number GSE287363 and GSE287364, respectively.

RESULTS

Single-cell RNA-seq analysis of LC and CAPD patient-derived PECs

We first collected PECs from a patient with LC at Child-Pugh class B (CP-B) and from a

patient undergoing CAPD on the same day and performed scRNA-seq analysis (Fig. 1A).

According to several signature genes, five cell clusters including T cells, B cells, *GATA6*-

expressing macrophages, DCs and *FCGR3A*-expressing macrophages were defined (Fig. 1B,C).

In CAPD patient-derived PECs, the frequencies of T cells, B cells, *GATA6*-expressing

macrophages, DCs and *FCGR3A*-expressing macrophages were 18%, 1.4%, 9.8%, 30% and

40%, respectively. DCs and macrophages occupied around 80% of total PECs. In LC patient-

derived PECs, T cells were significantly increased to 73% and became the most abundant

population. Meanwhile, *GATA6*-expressing macrophages, DCs and *FCGR3A*-expressing

macrophages were decreased to 1.6%, 2.1% and 16%, respectively. Due to the most prominent

decrease of DCs, *FCGR3A*-expressing macrophages were dominant over *GATA6*-expressing

macrophages and DCs in LC patient-derived PECs.

We then investigated expression of DC signature genes. According to the heatmap, feature plot

and violin plot analyses, expression of DC signature genes, including CD1 family members

(*CD1A, CD1B, CD1C, CD1D, CD1E*), *FLT3*, *FCER1A* and HLA family gene (*HLA-DRA*), were

much lower in DCs of LC patient-derived PECs than in DCs of CAPD patient-derived PECs

(Fig. 1C,D).

FCGR3A-expressing macrophages are featured by expression of *CD163*, *MARCO*, *SIGLEC1*

and *LYVE1*, and this type of macrophage has been reported to be involved in tissue repair.[7]

CD163, *MARCO*, *SIGLEC1* and *LYVE1* were expressed in *FCGR3A*-expressing macrophages in

CAPD patient-derived PECs (Fig. 1C,E). Expression levels of these genes was increased in LC

patient-derived PECs.

In mice, macrophages expressing a transcription factor, GATA6, occupy around 90% of

resident peritoneal macrophages.[8-10] GATA6 is essential for generation of resident peritoneal

macrophages and its target genes include *Serpinb2*, *Nt5e*, *Itga6*, *Thbs1*, *Tgfb2*, *Ltbp1*[10] and

CD9.[9] *GATA6*-expressing macrophages also exist in humans, although they represent less than

10% and are not a major population.[11] *CD9*, *SERPINB2, NT5E, ITGA6, THBS1, TGFB2* and

LTBP1 were highly expressed in *GATA6*-expressing macrophages in CAPD patient-derived

PECs (Fig. 2A-C). Expression levels of all of these genes except *CD9* and *NT5E* were

decreased, while *CD9* and *NT5E* expression was increased in LC patient-derived PECs.

PDGFRB, *KRT5*, *ACTA2*, *COL1A1* and *UPK3B* were expressed in peritoneal mesothelial cells

and classified as peritoneal mesothelial cell signature genes.[12] All of these genes were expressed

selectively in *GATA6*-expressing macrophages in CAPD patient-derived PECs (Fig. 2D-F).

Furthermore, among them, *PDGFRB*, *KRT5*, *ACTA2*, *COL1A1* and *UPK3B* were expressed at

higher levels in LC than in CAPD patient-derived PECs.

We then analyzed T cell populations. In LC patient-derived PECs, T cell percentages among

PECs were increased and *CD8A*[+] T cells dominated over *CD4*[+] T cells, but the ratio of *CD4*[+] to

CD8A[+] T cells were not so different from that of CAPD patient-derived PECs (Fig. 3A-C). In

both LC and CAPD patient-derived PECs, *CD8A*[+] T cells included a significant fraction of cells

expressing *CD8A*, *TRAV1-2* (encoding TCR Vα7.2) and *KLRB1* (encoding CD161), which

represent MAIT cells.

Bulk RNA-seq analysis revealed down- or up- regulated genes in LC patient-derived PECs

We also performed bulk RNA-seq analysis in PECs from patients with LC and patients

undergoing CAPD (Fig. 4A). In comparison to PECs from patients undergoing CAPD,

expression of 1107 and 906 genes were more than 2-fold down- and up-regulated, respectively,

with significance in PECs from patients with LC (Fig. 4B). The top five downregulated genes in

PECs from patients with LC included DC signature genes, such as *CD1C*, *CD1E*, *CD1B* and

FCER1A (Fig. 4C).

KEGG pathway analysis of the top 100 downregulated genes in PECs from patients with LC

showed that 'Hematopoietic cell lineage' was the most enriched pathway (Fig. 4D). The

pathway contained CD1 family genes (*CD1A*, *CD1B*, *CD1C* and *CD1E*) and *FLT3*, *IL1R2* and

HLA-DQR2, all of which are dominantly expressed in DCs (Supplementary Figure 1A,B).

These CD1 family genes (*CD1A*, *CD1B*, *CD1C* and *CD1E*), *IL1R2* and *LAMC2* were also

contained in the second enriched pathway, 'Amoebiasis'. The third enriched pathway

was 'ECM-receptor interaction' pathway, which contained *TNC*, *NPNT*, *COL6A3* and *THBS1*.

Among the four genes, *TNC*, *NPNT* and *COL6A3* were dominantly expressed in *GATA6*-

expressing macrophages and *THBS1* was expressed in DCs and *GATA6*-expressing

macrophages. According to the scRNA-seq analysis, we categorized the top 100 downregulated

genes in PECs from patients with LC. 82 genes were expressed in DCs or *GATA6*-expressing

macrophages. Among them, 37 genes were expressed in both DCs and *GATA6*-expressing

macrophages. 45 genes were dominantly expressed in *GATA6*-expressing macrophages, while

no genes were expressed dominantly in DCs (shown in light blue in Fig. 4C). Thus, the gene

expression analysis indicates that DCs and *GATA6*-expressing macrophages mainly decreased in

PECs from patients with LC.

KEGG pathway analysis of 100 upregulated genes in PECs from patients with LC

showed 'Graft-versus-host disease' was the most enriched pathway (Fig.4E, F). This pathway

contained *GZMB*, *KIR3DL1*, *FASLG*, *KIR3DL2* and *KIR2DL3*, which are highly expressed in T

cells (Supplementary Figure 1C,D). Among 100 upregulated genes, 24 genes were dominantly

expressed in *FCGR3A*-expressing macrophages (shown in green in Fig. 4E). Those genes

include complement genes such as *C1QB*, *C1QC* and *C2*, which belong to the 'Alcoholic liver

disease' pathway. They also include *ABCA1*, *APOE* and *PLTP*, which belong to the 'Cholesterol

metabolism' pathway. *C1QB*, *C1QC*, *C2*, *ABCA1*, *APOE* and *PLTP* were highly expressed in

FCGR3A-expressing macrophages in PECs from patients with LC (Supplementary Figure

1C,D). These results indicate that T cells and *FCGR3A*-expressing macrophages were increased

in PECs from patients with LC.

Comprehensive gene expression analysis in correlation with LC severity

We then performed comprehensive gene expression analysis in correlation with LC severity

(Fig. 5A, left). Severity of LC was assessed by Child-Pugh (CP) classification, which are widely

used.[13] Among eight patients with LC, four patients, including a patient subjected to scRNA-seq

analysis, were CP-B and the other 4 patients were Child-Pugh class C (CP-C).

We first searched the genes, expression of which decreased with disease severity. Expression

of 956 genes were more than 2-fold downregulated in PECs from patients with LC CP-B,

compared with PECs from patients undergoing CAPD (Fig.5A, middle). Expression of 443

genes were more than 2-fold downregulated in PECs from LC CP-C patients, compared with

PECs from patients with LC CP-B (Fig.5A, right). Comparison of these genes revealed that

expression of six genes (*CTLA4, C15orf48, PMAIP1, FAM86HP, TNNT1,* and *EPC1*) was

decreased as LC gets worse (Fig. 5B,C). Among these genes, *CTLA4, C15orf48, PMAIP1* and

EPC1 were highly expressed in DCs and *GATA6*-expressing macrophages, while *TNNT1* was

highly expressed in DCs and *FCGR3A*-expressing macrophages (Supplementary Figure 2A,B).

We then searched the genes, the expression of which increased with disease severity.

Expression of 1045 genes were more than 2-fold upregulated in PECs from patients with LC

CP-B, compared with PECs from patients undergoing CAPD (Fig.5A, middle). Expression of

295 genes were more than 2-fold upregulated in PECs from patients with LC CP-C, compared

with PECs from patients with LC CP-B (Fig.5A, right). Comparison of these genes clarified that

expression of four genes (*IDH1, CMKLR1, SLC7A8* and *SLC46A1*) was increased as LC

becomes more severe (Fig.5D, E). All four of these genes were highly expressed in *GATA6-* and

FCGR3A-expressing macrophages (Supplementary Figure 2C,D).

T cell signature genes are downregulated in PECs from patients with LC CP-C

We have also searched genes, expression of which is upregulated at CP-B and then

downregulated at CP-C. Expression of 126 genes were more than 2-fold upregulated in PECs

from patients with LC CP-B, compared with PECs from patients with LC CP-C and PECs from

patients undergoing CAPD (Fig.6 A,B). KEGG pathway analysis of these 126 genes

showed 'Natural killer cell mediated cytotoxicity' pathway was enriched with significance

(Fig.6B, C). Expression of *GZMB*, *IFNG*, *SH2D1A* and *LCK* in the pathway was elevated in

PECs from patients with LC CP-B. According to scRNA-seq analysis, these four genes were

highly expressed in T cells (Supplementary Figure 3A,B). These results suggest that T cells

increased in PECs from patients with LC CP-B, and then decreased in PECs from patients with

LC CP-C.

Decrease of CD1c^high CD14^int DCs and increase of CD8α⁺ MAIT cells in LC patient-derived

PECs

We analyzed surface expression of several lineage markers in PECs. CD45$^+$ cells of PECs are

shown as the CD1c versus CD14 profile (Fig.7A). This profile showed three major clusters,

CD1c^{high} CD14^{int} cells, CD1c^{int} CD14^{high} cells, and CD1c$^{low-int}$ CD14^{low} cells. According to high

expression of CD1c, CD1c^{high} CD14^{int} cells can be regarded as DCs. CD1c^{int} CD14^{high} cells can

be regarded as *FCGR3A*-expressing macrophages, because they express CD16 but not CD1c

(Fig. 7D). Meanwhile, CD1c$^{low-int}$ CD14^{low} cells are heterogenous and T cells were a major

population of them. Notably, quantitative real-time PCR analysis among fractions of CD45$^+$

PECs revealed that expression of *GATA6* was found only in CD1c$^{low-int}$ CD14^{low} cells (Fig. 7B),

indicating that *GATA6*-expressing macrophages are contained in CD1c$^{low-int}$ CD14^{low} cells.

According to the scRNA-seq analysis, it can be assumed that CD9 expression can be used to

identify *GATA6*-expressing macrophages. Then, we have examined surface CD9 expression in

CD1c$^{low-int}$ CD14^{low} cells in CAPD or LC patient-derived PECs. The frequency of CD1c$^{low-int}$

CD14^{low} CD9$^+$ cells was decreased in LC patient-derived PECs (Fig.7C).

Then we have examined the fluctuation of immune cell populations in LC patient-derived

PECs. CD1c^{high} CD14^{int} cells, that is, DCs were present in CAPD patient-derived PECs, while

they were decreased or absent in LC patient-derived PECs (Fig.7D). CD1c^{int} CD14^{high} cells were

decreased in LC patient-derived PECs compared with in CAPD patient-derived PECs. We have

then analyzed expression of CD16 and CD163 in CD1c^{int} CD14^{high} cells (Fig. 7E). Expression of

CD16 as well as CD163 was higher in LC patient-derived PECs than in CAPD patient-derived

PECs. Meanwhile, expression of CD16 and CD163 on DCs and CD1c$^{low-int}$ CD14^{low} cells

remained low in LC patient-derived PECs. Thus, expression of CD16 and CD163 was elevated

only in CD1c^{int} CD14^{high} cells among LC patient-derived PECs. In patients with LC CP-B, T

cells including CD161[+] TCR Vα7.2[+] MAIT cells were increased (Fig.7F). These results were

consistent with both scRNA-seq analysis and bulk RNA-seq analysis.

DISCUSSION

The current study is the first report on comparative analysis to characterize immune cell

profiles of PECs between patients with LC and patients undergoing CAPD. Five main cell

clusters, including T cells, B cells, DCs and *GATA6-* and *FCGR3A*-expressing macrophages,

were found in PECs. In CAPD patient-derived PECs, DCs and *GATA6-* and *FCGR3A-*

expressing macrophages were dominant cell populations. This immune cell profile is consistent

with previous reports.[11,12] Meanwhile, in LC patient-derived PECs, DCs and two clusters of

macrophages decreased and T cells became dominant. Among DCs and macrophages, *FCGR3A-*

expressing macrophages were dominant in LC patient-derived PECs. Bulk RNA-seq analysis

further clarified up- or down-regulated gene lists between PECs from patients with LC and

those from patients undergoing CAPD. The results were consistent with scRNA-seq analysis of

PECs.

FCGR3A-expressing macrophages express *CD163*, *MARCO* and *LYVE1* and represent M2

type and anti-inflammatory phenotype. Meanwhile, DCs, which express *CD1C* and *CD1E*,

show M1 type and proinflammatory phenotype. Therefore, LC PECs tend to be skewed to M2.

In the liver, MAIT cells can promote liver fibrosis through the recruitment of profibrotic

monocytes.[14] In this context, increase of MAIT cells in the peritoneal cavity might cause

recruitment of *FCGR3A*-expressing macrophages. It should be intriguing to clarify the roles of

such recruited *FCGR3A*-expressing macrophages in progression of liver fibrosis.

GATA6-expressing macrophages are resident in the peritoneal cavity and also profibrotic, as

they express collagen gene such as *COL1A1* and repair-related genes (*THBS1*, *TGFB2*) (Fig.2A-

F). *GATA6*-expressing macrophages represent a minor population in LC patient-derived PECs.

However, it is still possible that such macrophages migrate to the liver and provoke fibrosis,

because *GATA6*-expressing macrophages in the peritoneal cavity are required for liver

protection in murine liver injury model.[3]

We have further characterized gene expressions with LC severity. Compared with PECs from

patients undergoing CAPD, expression of six genes was decreased in PECs from patients with

LC CP-B and their expression levels were further reduced in PECs from patients with LC CP-C.

Those genes were expressed mainly in DCs and *GATA6*-expressing macrophages. Among those

genes, *CTLA4* encodes CTLA4, which can inhibit T cell proliferation.[15] *C15orf48* encodes

modulator of cytochrome C oxidase during inflammation, which is a mitochondrial protein

critically involved in autophagy induction and *C15orf48*-deficient mice showed development of

autoimmunity, indicating that *C15orf48* regulates immune functions.[16] Decreased expression of

these immunosuppressive molecules might perturb peritoneal immune homeostasis in patients

with LC.

Meanwhile, four genes (*IDH1*, *CMKLR1*, *SLC7A8* and *SLC46A*) were found to increase in

their expression as the LC stage progresses. Those genes were highly expressed in *GATA6*- and

FCGR3A-expressing macrophages. *IDH1* encodes IDH, which generates α-ketoglutarate from

isocitrate, thereby promoting differentiation of macrophages towards M2.[17,18] Expression of

CMKLR1 is high in macrophages from broncho alveolar lavages of patients with cystic fibrosis,

indicating involvement of *CMKLR1* in fibrosis.[19] Expression of *SLC7A8* is also correlated with

that of CD163 in M2 macrophages.[20] Thus, elevated expression of these genes might be related

to upregulation of fibrotic or M2 type macrophage functions. It should be interesting whether or

how those gene products contribute to progression of liver fibrosis.

Expression of some genes was elevated in PECs from patients with LC CP-B, but decreased in

PECs from patients with LC CP-C. KEGG pathway analysis of such gene enrichment of the

'Natural killer cell mediated cytotoxicity' pathway was enriched (Fig. 6C). This enrichment can

be assumed to be due to increase of T cells (Supplementary Figure 3A,B). Especially cytotoxic

T cell signature genes, *GZMB* and *IFNG,* were prominently elevated. MAIT cells, which are

also cytotoxic, are decreased in peripheral bloods, but increased in PECs from patients with

decompensated LC and spontaneous bacterial peritonitis (SBP).[2,21] In that study, patients with

LC were not categorized by CP classification. Meanwhile, we studied patients with LC at CP-B

and CP-C in the absence of SBP (Supplementary Table S1). It is interesting to elucidate how T

cells including MAIT cells are increased at CP-B and then decreased at CP-C. CD163^{high} M2

type macrophages can inhibit T cell proliferation.[22] Therefore, the T cell decrease can be

ascribed to dominancy of M2 macrophages at CP-C. It is also possible that activated T cells

leave the peritoneal cavity and migrate to the liver.

FACS analysis showed that, in LC patient-derived PECs, CD1c^{high} CD14^{int} DCs were

decreased, while CD16^{high} CD163^{high} cells were increased (Fig. 7D,E). We further found that

CD1$^{low-int}$ CD14^{low} cells include *GATA6*-expressing macrophages (Fig. 7B) and that these cells

were decreased in LC patient-derived PECs. Furthermore, T cells including MAIT cells were

increased and existed as a major population in LC patient-derived PECs. These flowcytometry

analysis results were consistent with the results of scRNA-seq and bulk RNA-seq analyses.

We thus characterized immune cell profiles of PECs from patients with LC. Whereas DCs and

macrophages were dominant in PECs from patients undergoing CAPD, T cells were the main

population in PECs from patients with LC. Although DCs and macrophages were decreased in

PECs from patients with LC, *FCGR3A*-expressing macrophages, which are profibrotic, were

relatively dominant over DCs and *GATA6*-expressing macrophages. We also compared gene

expression profiles between CP-B and CP-C and clarified that T cells were increased at CP-B

but decreased at CP-C. We further identified a set of genes expression of which was increased or

decreased with LC severity. Our present findings should have a potential to lead to further

development of diagnosis or therapeutic maneuver for LC.

ACKNOWLEDGEMENTS

We thank Ms. Sumiko Okura and Ms. Aoi Matsumura-Tawaki for secretarial assistance. We

acknowledge proofreading and editing by Benjamin Phillis at the Clinical Study Support Center,

Wakayama Medical University.

REFERENCES

1.   GBD 2017 Cirrhosis Collaborators. The global, regional, and national burden of

cirrhosis by cause in 195 countries and territories, 1990-2017: a systematic analysis for the Global

Burden of Disease Study 2017. *Lancet Gastroenterol Hepatol*. Mar 2020;5(3):245-266.

doi:10.1016/s2468-1253(19)30349-8

2.   Niehaus CE, Strunz B, Cornillet M, et al. MAIT Cells Are Enriched and Highly

Functional in Ascites of Patients With Decompensated Liver Cirrhosis. *Hepatology*. Oct

2020;72(4):1378-1393. doi:10.1002/hep.31153

3.   Wang J, Kubes P. A Reservoir of Mature Cavity Macrophages that Can Rapidly Invade

Visceral Organs to Affect Tissue Repair. *Cell*. Apr 21 2016;165(3):668-78.

doi:10.1016/j.cell.2016.03.009

4.   Hossain M, Shim R, Lee WY, Sharpe AH, Kubes P. Gata6(+) resident peritoneal

macrophages promote the growth of liver metastasis. *Nat Commun*. Jul 29 2022;13(1):4406.

doi:10.1038/s41467-022-32080-y

5.   Chang H, Ni Y, Shen C, et al. Peritoneal GATA6(+) macrophage drives hepatic

immunopathogenesis and maintains the T(reg) cell niche in the liver. *Immunology*. Sep

2022;167(1):77-93. doi:10.1111/imm.13519

6.   Goedhart J, Luijsterburg MS. VolcaNoseR is a web app for creating, exploring, labeling

and sharing volcano plots. *Sci Rep*. Nov 25 2020;10(1):20560. doi:10.1038/s41598-020-76603-3

7.   Chakarov S, Lim HY, Tan L, et al. Two distinct interstitial macrophage populations

coexist across tissues in specific subtissular niches. *Science*. Mar 15 2019;363(6432).

doi:10.1126/science.aau0964

8.   Ghosn EE, Cassado AA, Govoni GR, et al. Two physically, functionally, and

developmentally distinct peritoneal macrophage subsets. *Proc Natl Acad Sci U S A*. Feb 9

2010;107(6):2568-2573. doi:10.1073/pnas.0915000107

9.   Rosas M, Davies LC, Giles PJ, et al. The transcription factor Gata6 links tissue

macrophage phenotype and proliferative renewal. *Science*. May 9 2014;344(6184):645-648.

doi:10.1126/science.1251414

10.   Okabe Y, Medzhitov R. Tissue-specific signals control reversible program of

localization and functional polarization of macrophages. *Cell*. May 8 2014;157(4):832-44.

doi:10.1016/j.cell.2014.04.016

11.   Han J, Gallerand A, Erlich EC, et al. Human serous cavity macrophages and dendritic

cells possess counterparts in the mouse with a distinct distribution between species. *Nat Immunol*.

Jan 2024;25(1):155-165. doi:10.1038/s41590-023-01688-7

12.   Hu W, Li G, Dong W, et al. Single-cell sequencing reveals peritoneal environment and

insights into fibrosis in CAPD patients. *iScience*. Apr 21 2023;26(4):106336.

doi:10.1016/j.isci.2023.106336

13.   Durand F, Valla D. Assessment of the prognosis of cirrhosis: Child-Pugh versus MELD.

J Hepatol. 2005;42 Suppl(1):S100-7. doi:10.1016/j.jhep.2004.11.015

14.   Mabire M, Hegde P, Hammoutene A, et al. MAIT cell inhibition promotes liver fibrosis

regression via macrophage phenotype reprogramming. *Nat Commun*. Apr 1 2023;14(1):1830.

doi:10.1038/s41467-023-37453-5

15.   Laurent S, Carrega P, Saverino D, et al. CTLA-4 is expressed by human monocyte-

derived dendritic cells and regulates their functions. *Hum Immunol*. Oct 2010;71(10):934-41.

doi:10.1016/j.humimm.2010.07.007

16.   Takakura Y, Machida M, Terada N, et al. Mitochondrial protein C15ORF48 is a stress-

independent inducer of autophagy that regulates oxidative stress and autoimmunity. *Nat Commun*.

Feb 1 2024;15(1):953. doi:10.1038/s41467-024-45206-1

17.   Liu PS, Wang H, Li X, et al. α-ketoglutarate orchestrates macrophage activation through

metabolic and epigenetic reprogramming. *Nat Immunol*. Sep 2017;18(9):985-994.

doi:10.1038/ni.3796

18.   Mouton AJ, Aitken NM, Moak SP, et al. Temporal changes in glucose metabolism reflect

polarization in resident and monocyte-derived macrophages after myocardial infarction. *Front*

Cardiovasc Med. 2023;10:1136252. doi:10.3389/fcvm.2023.1136252

19.   Mannes PZ, Barnes CE, Biermann J, et al. Molecular imaging of chemokine-like

receptor 1 (CMKLR1) in experimental acute lung injury. *Proc Natl Acad Sci U S A*. Jan 17

2023;120(3):e2216458120. doi:10.1073/pnas.2216458120

20.   Zhao X, Jin L, Liu Y, Liu Z, Liu Q. Bioinformatic analysis of the role of solute carrier-

glutamine transporters in breast cancer. *Ann Transl Med*. Jul 2022;10(14):777. doi:10.21037/atm-

22-2620

21.   Ibidapo-Obe O, Stengel S, Köse-Vogel N, et al. Mucosal-Associated Invariant T Cells

Redistribute to the Peritoneal Cavity During Spontaneous Bacterial Peritonitis and Contribute to

Peritoneal   Inflammation.   *Cell*   *Mol*   *Gastroenterol*   *Hepatol*.   2020;9(4):661-677.

doi:10.1016/j.jcmgh.2020.01.003

22.   Han Q, Shi H, Liu F. CD163(+) M2-type tumor-associated macrophage support the

suppression of tumor-infiltrating T cells in osteosarcoma. *Int Immunopharmacol*. May

2016;34:101-106. doi:10.1016/j.intimp.2016.01.023

FIG.1. Single-cell RNA sequencing analysis revealed major cell clusters and gene expression

profiles of CAPD and LC patient-derived PECs

(A) Schematic workflow of scRNA-seq analysis of human PECs.

(B) UMAP shows five distinct clusters of CAPD PECs (CAPD patient-derived PECs) and LC

PECs (LC patient-derived PECs). Clusters include T cells, B cells, *GATA6*-expressing

macrophages (*GATA6*+ MPs), DCs and *FCGR3A*-expressing macrophages (*FCGR3A*+ MPs).

(C) Heatmap shows five clusters of CAPD and LC PECs and the expression of immune cell

signature genes.

(D) Feature plots and violin plots of DC signature gene expression in CAPD and LC PECs.

(E) Feature plots or violin plots of expression of *CD14*, *FCGR3A*, *CD163*, *MARCO*, *SIGLEC1*

and *LYVE1* in CAPD and LC PECs.

FIG.2. Single-cell RNA sequencing analysis revealed gene expression profiles of *GATA6-*

expressing macrophages in CAPD and LC patient-derived PECs

(A-C) Gene expression of *GATA6*, *CD9*, *SERPINB2*, *NT5E*, *ITGA6*, *THBS1*, *TGFB2* and *LTBP1*

is shown in heatmap (A), violin plots (B) and feature plots (C).

(D-F) Gene expression of *PDGFRB*, *KRT5*, *ACTA2*, *COL1A1* and *UPK3B* is shown in heatmap

(D), violin plots (E) and feature plots (F).

FIG.3. Single-cell RNA sequencing analysis revealed gene expression profiles of T cells in

CAPD and LC patient-derived PECs

(A-C) Gene expression of *CD3E*, *CD4*, *CD8A*, *TRAV1-2* and *KLRB1* is shown in heatmap (A),

violin plots (B) and feature plots (C).

FIG.4. Bulk RNA sequencing analysis revealed that DCs and *GATA6*-expressing

macrophages mainly decreased in LC patient-derived PECs

(A) Schematic workflow of bulk RNA-seq analysis of human PECs.

(B) Volcano plots of gene expression in PECs show a fold change (FC) and p value for the

comparison of LC PECs versus CAPD PECs. Up- or down- regulated genes (FC $\geq$ 2, $p <$

0.05) are depicted in red or blue, respectively.

(C) Heatmap of the top 100 downregulated genes in LC PECs.

(D) KEGG pathway analysis of downregulated genes in (C).

(E) Heatmap of the top 100 upregulated genes in LC PECs.

(F) KEGG pathway analysis of upregulated genes in (E).

FIG.5. Bulk RNA sequencing analysis revealed up- or down- regulated genes in correlation

with LC severity

(A) Volcano plots of gene expression show a FC and *p* value for the comparisons of LC CP-B

PECs versus CAPD PECs (middle) or LC CP-C PECs versus LC CP-B PECs (right). Two-

fold up- or down- regulated genes (FC ≥ 2, *p* < 0.05) with significance are depicted in red or

blue, respectively.

(B) Venn diagram depicting the numbers of 2-fold downregulated genes with significance in LC

CP-B PECs (A, middle) or LC CP-C PECs (A, right), then expression of the overlapped six

genes showed in heatmap.

(C) Expression of the six genes (B). Statistical significance was determined by an unpaired two-

tailed Student's t test. p values are indicated by $*p < 0.05$, $**p < 0.01$. $p < 0.05$ was considered

statistically significant.

(D) Venn diagram depicting the numbers of 2-fold upregulated genes with significance in LC CP-

B PECs (A, middle) or LC CP-C PECs (A, right), then expression of the overlapped four

genes showed in heatmap.

(E) Expression of the four genes (D). Statistical significance was determined by an unpaired two-

tailed Student's t test. p values are indicated by $*p < 0.05$, $**p < 0.01$. $p < 0.05$ was considered

statistically significant.

A

Up-regulated in
LC CP-B
1045 genes

(related to Fig.5)

Down-regulated in
LC CP-C
443 genes

(related to Fig.5)

919 | 126 | 317

B

C

-Log₁₀ (*p* value)

Natural killer cell mediated cytotoxicity
PD-L1 expression and PD-1 checkpoint pathway in cancer
Chagas disease
T cell receptor signaling pathway

1.3 (*p* = 0.05)

FIG.6. Bulk RNA sequencing analysis revealed that T cell signature genes were upregulated

in LC CP-B but not LC CP-C

(A) Venn diagram depicting the numbers of upregulated genes in LC CP-B (Fig.5A, middle) or

those of downregulated genes in LC CP-C (Fig.5A, right).

(B) Heatmap of overlapped genes in (A).

(C) KEGG pathway analysis of overlapped genes in (A).

FIG.7. Flowcytometry analysis revealed decrease of DCs and increase of CD16^{high} CD163^{high}

macrophages and CD8α$^+$ MAIT cells in LC PECs

(A) Gating strategy to identify CD1c^{high} CD14^{int}, CD1c^{int} CD14^{high} and CD1c$^{low-int}$ CD14^{low} cells

in CAPD PECs. Numbers indicate percentages of gated cells.

(B) CD1c^{high} CD14^{int}, CD1c^{int} CD14^{high} and CD1c$^{low-int}$ CD14^{low} cells were purified from CAPD

PECs by MACS system. Expression of *GATA6* among these three subsets is shown (N=3).

(C) Frequency of CD1c$^{low-int}$ CD14^{low} CD9$^+$ cells in CAPD PECs and LC PECs. Numbers

indicate percentages of gated cells.

(D) Surface expression of CD1c, CD14, CD16 and CD163 in CD1c^{high} CD14^{int}, CD1c^{int}

CD14^{high} and CD1c$^{low-int}$ CD14^{low} cells in CAPD PECs or LC PECs. Red dots indicate

CD1c^{high} CD14^{int} cells, blue dots indicate CD1c^{int} CD14^{high} cells and orange dots indicate

CD1c$^{low-int}$ CD14^{low} cells, respectively. Numbers indicate percentages of gated cells.

(E) Surface expression of CD16 and CD163 in CD1c$^{low-int}$ CD14^{high} cells in CAPD PECs (N=4)

or LC PECs (N=3). MFI indicates mean fluorescence intensity. Statistical significance was

determined by an unpaired two-tailed Student's t test. *p* values are indicated by *p < 0.05,

***p < 0.001. *p* < 0.05 was considered statistically significant.

(F) Surface expression of T cell markers in CAPD PECs or LC PECs. Numbers indicate

percentage of gated cells.

Supplementary Figure 1

Gene expression profiles of up- or down- regulated genes in LC patient-derived PECs by

scRNA-seq analysis

(A,B) Expression of *HLA-DQA2*, *IL1R2*, *LAMC2*, *TNC*, *COL6A3*, *NPNT* and *THBS1* is shown in

feature plots (A) and violin plots (B).

(C,D) Expression of *GZMB*, *KIR3DL1*, *FASLG*, *KIR3DL2*, *KIR2DL3*, *C1QB*, *C1QC*, *C2*, *ABCA1*,

APOE and *PLTP* is shown in feature plots (C) and violin plots (D).

Supplementary Figure 2

Gene expression profiles of up- or down- regulated genes with LC severity by scRNA-seq

analysis

(A,B) Expression of *CTLA4*, *C15orf48*, *PMAIP1*, *FAM86HP*, *TNNT1* and *EPC1* is shown in

feature plots (A) and violin plots (B).

(C,D) Expression of *IDH1*, *CMKLR1*, *SLC7A8* and *SLC46A1* is shown in feature plots (C) and

violin plots (D).

580 **Supplementary Figure 3**

581 **Gene expression profiles of upregulated genes in LC CP-B patient-derived PECs by scRNA-**

582 **seq analysis**

583 (A,B) Expression of *GZMB*, *IFNG*, *SH2D1A* and *LCK* is shown in feature plots (A) and violin

584 plots (B).

 Supplementary Table S1. Patients undergoing CAPD or with LC enrolled in this study

Patient	Group	Etiology of LC	Experiment performed	SBP
#1	CAPD	-	sample 1 ; Bulk RNA-seq sample 2 ; scRNA-seq sample 3 ; Flowcytometry	not tested
#2	CAPD	-	sample 1 ; Bulk RNA-seq sample 2 ; Flowcytometry	not tested
#3	CAPD	-	sample 1 ; Bulk RNA-seq sample 2, 3 ; Flowcytometry	not tested
#4	CAPD	-	Bulk RNA-seq	not tested
#5	CAPD	-	Bulk RNA-seq	not tested
#6	CAPD	-	Bulk RNA-seq	not tested
#7	LC CP-B	MASLD	Bulk RNA-seq, Flowcytometry	Negative
#8	LC CP-B	Hepatitis C	Bulk RNA-seq, Flowcytometry	Negative
#9	LC CP-B	PBC	Flowcytometry	Negative
#10	LC CP-B	ALD	Bulk RNA-seq, Flowcytometry	Negative
#11	LC CP-B	MASLD	sample 1 ; scRNA-seq, Bulk RNA-seq sample 2 ; Flowcytometry	Negative

#12	LC CP-C	PSC+AIH	**Bulk RNA-seq, Flowcytometry**	**Negative**
#13	LC CP-C	postoperative cholangitis	**Bulk RNA-seq, Flowcytometry**	**Negative**
#14	LC CP-C	PSC	**Bulk RNA-seq, Flowcytometry**	**Negative**
#15	LC CP-C	ALD	**Bulk RNA-seq**	**Negative**
#16	CAPD	-	**Flowcytometry**	**not tested**
#17	LC CP-B	ALD	**Flowcytometry**	**Negative**
#18	LC CP-B	unknown	**Flowcytometry**	**Negative**

Summary of each assay is shown.

SBP, spontaneous bacterial peritonitis; MASLD, metabolic dysfunction associated steatotic liver

disease; PBC, primary biliary cholangitis; ALD, alcoholic liver disease; PSC, primary sclerosing

cholangitis; AIH, autoimmune hepatitis

Supplementary Table S2. Clinical characteristics of the study individuals for scRNA-seq

analysis

Characteristic	Analysis of scRNA-seq	
	CAPD (N = 1)	LC CP-B (N = 1)
Age, y	69	76
Male sex	Male	Male
Serum leukocytes (×10^9/l)	5.86	3.09
Plt (×10^9 /l)	248	52
Serum PT (%)	-	60.9
Serum Albumin (g/dl)	3.0	2.9
AST (U/l)	13	15
ALT (U/l)	13	11
T-Bil (mg/dl)	0.5	0.9

602 **Supplementary Table S3.** Clinical characteristics of the study individuals for bulk RNA-seq

603 analysis

	Analysis of Bulk RNA-seq		
Characteristic	**CAPD (N = 6)**	**LC CP-B (N = 4)**	**LC CP-C (N = 4)**
Age, y	63.5 (45-77)	74 (65-81)	59.5 (45-69)
Sex (male / female)	4 / 2	2 / 2	3 / 1
Body Mass Index	25.1 (21.3-32.5)	22.5 (17.9-33.0)	20.6 (19.0-20.9)
Serum leukocytes ($\times 10^9$/l)	5.445 (3.580-11.46)	4.280 (3.090-6.190)	7.980 (5.480-10.08)
Plt ($\times 10^9$/l)	203.5 (170-292)	54.5 (52-245)	111 (82-128)
Serum PT (%)	-	77.25 (60.9-87.9)	52.7 (49-66.1)
Serum Albumin (g/dl)	3.15 (2.9-3.6)	2.8 (2.6-2.9)	2.05 (2.0-2.8)
AST (U/l)	15.5 (11-22)	41 (15-65)	50.5 (30-454)
ALT (U/l)	14 (6-45)	13 (9-50)	35.5 (13-120)
T-Bil (mg/dl)	-	1.3 (0.5-1.7)	8.95 (0.7-23.4)
Creatinine	11.71 (9.33-12.04)	0.975 (0.82-1.17)	1.085 (0.52-4.16)

608 **Supplementary Table S4.** Antibodies used for flowcytometry analysis

Antibody name	Color	Clone	Cat	Antibody source
anti-CD45	APC	HI30	304037	BioLegend
anti-CD1c	Biotin	L161	331504	BioLegend
anti-CD14	PE	61D3	12-0149-42	Invitrogen
anti-CD9	PerCPCy5.5	HI9A	312110	BioLegend
anti-CD3	APCCy7	UCTH1	300426	BioLegend
anti-CD16 (encoded by *FCGR3A*)	Pacific Blue	3G8	980106	BioLegend
anti-CD163	FITC	GHI/61	333617	BioLegend
anti- CD161	PECy7	HP-3G10	339918	BioLegend
anti-TCR Vα7.2	Brilliant violet 421	3C10	351716	BioLegend
anti-CD4	PerCPCy5.5	RPA-T4	300530	BioLegend
anti-CD8α	FITC	RPA-T8	301060	BioLegend