# Somatic repetitive element insertions Define New Biomarkers in Pan-cancer genome

Lin Xia[1#], Kailing Tu[1#], Huan Wang[1#], Xuyan Liu[2], Yahui Liu[2], Qilin Zhang[2], Ziting Feng[2], Chunyan Yu[1], Maolin Li[1], Yan Huang[1], Jing Wang[1], Ranlei Wei[1], Tianfu Zeng[2], Xuenan Pi[1], Chen Ling[3], Qianwen Zhang[3], Yixin Qiao[4], Shengzhong Hou[5], Zhuoyuan Zhang[6,7], Ailin Wei[8], Haotian Liao[9], Wenhao Guo[10], Jifeng Liu[4,11], Zhong Wu[5], Zhoufeng Wang[12,13], Ang Li[5], Weiming Li[12,13], Dan Xie[1,2*]

1. Laboratory of Omics Technology and Bioinformatics, Frontiers Science Center for Disease-related Molecular Network, West China Hospital, Sichuan University, Chengdu, Sichuan, China

2. State Key Laboratory of Biotherapy, West China Hospital, Sichuan University, Chengdu, Sichuan, China

3. Department of Gynecology and Obstetrics, Key Laboratory of Birth Defects and Related Diseases of Women and Children, Ministry of Education, West China Second Hospital, Sichuan University, Chengdu, China

4. Department of Otolaryngology Head and Neck Surgery, West China Hospital, Sichuan University, Chengdu, China.

5. Division of Pancreatic Surgery, Department of General Surgery, West China Hospital, Sichuan University

6. Department of Head and Neck Cancer Surgery, West China School of Stomatology, Sichuan University, Chengdu, Sichuan, China

7. State Key Laboratory of Oral Diseases, West China School of Stomatology, Sichuan University, Chengdu, Sichuan, China

8. Guangan People's Hosital, Guangan, Sichuan, China

9. Division of Liver Surgery, Department of General Surgery and Laboratory of Liver Surgery, and State Key Laboratory of Biotherapy, West China Hospital, Sichuan University, Chengdu, Sichuan, China.

10. Department of Abdominal Oncology, Cancer Center and State Key Laboratory of

Biotherapy, West China Hospital, West China Medical School, Sichuan

University,Chengdu, China.

11. Deep Underground Space Medical Center, West China Hospital, Sichuan University,

Chengdu, China.

12. Institute of Respiratory Health, Frontiers Science Center for Disease-related

Molecular Network, West China Hospital, Sichuan University, Chengdu, Sichuan,

China

13. Precision Medicine Key Laboratory of Sichuan Province, West China Hospital,

Sichuan University, Chengdu, Sichuan, China

* Corresponding authors. Email: danxie@scu.edu.cn

# These authors contributed equally

# Abstract

Repetitive elements insertions (REIs), including tandem repeat insertions (TRIs) and

transposable element insertions (TEIs), constitute >90% of large-scale (>50bp) somatic

insertions and are linked to diseases and cancer biomarkers. However, the genomic

landscape of large-scale REIs in pan-cancer has not been well-characterized due to

short-read sequencing limitations. Here, we detected somatic REIs in 325 cancer

genomes across 12 cancer types using long-read whole-genome sequencing and

customized algorithms. The diverse sequence patterns of somatic TRIs and the insertion

mechanisms of somatic TEIs were characterized across different cancer types. We

identified 152 high-frequency somatic TRIs, with 10 showing pan-cancer characteristics,

including TRIs in *SHROOM2, PALMD*, and the enhancer of *PTPRZ1*. The MHC class II

gene cluster exhibited the highest somatic TEI abundance, affecting ~40% tumors.

Adenocarcinomas and squamous cell carcinomas exhibited differing TEI distributions

within this region. Our study highlights REIs as important but under-explored markers for

personalized cancer diagnosis and immunotherapy.

# Introduction

Cancer is formidable in its vast diversity, although all cancers share common hallmarks[1,2]. Understanding both the commonalities and variations among cancer types is crucial for uncovering the fundamental mechanisms of oncogenesis and advancing precision medicine. Pan-cancer studies, which aim at identifying shared genomic patterns and alterations driving cancer progression across cancers [3-7], not only enhances our understanding of cancer biology, but also holds considerable promise for advancing early diagnosis and developing personalized treatment strategies. These efforts have significantly expanded the knowledge of cancer driver genes, with *TP53*, *PIK3CA*, *KRAS*, *PTEN*, and *ARID1A* among the most frequently mutated genes across various cancers [8]. Furthermore, pan-cancer studies have uncovered new non-coding driver genes [9,10], mutational signatures [11], and evolutionary patterns [12-14], as well as their downstream effects on gene expression [15] and cancer-specific characteristics [16,17]. Despite these advances, many findings have yet to translate into clinical benefits, highlighting the pressing need for the identification of other critical genetic variants in cancer progression.

Previous pan-cancer research has predominantly focused on small-scale mutations, such as single nucleotide variations (SNVs) and small insertions/deletions (INDELs, < 50bp). Structural variations (SVs), which impact a broader range of nucleotide change [18], have increasingly garnered attention due to their significant role in cancer progression [19]. With short-read sequencing (SRS) data, researchers have identified various genomic SVs, including whole-genome duplications [20], chromothripsis [21], chromosomal instability [17], and complex structural variants [22], all associated with poorer prognoses. Nevertheless, large-scale insertions (>50 bp) in tumors remain less explored, particularly for repetitive element insertions (REIs), including tandem repeat insertions (TRIs) and transposable element insertions (TEIs) [23]. Previous population studies and tumor analyses indicate that over 80% of large-scale insertions are repetitive element insertions (REIs) [24-26]. Both TRIs [27,28] and TEIs [29-32] are drawing increasing attention as potential diagnostic and therapeutic target for various types of cancer. Despite extensive mapping efforts to characterize TRI

and TEI events in cancer genomes [27,28,33-35], only a limited number of recurrent somatic

REIs have been identified. The short read lengths of SRS technologies have hindered the

in-depth exploration of REI variation patterns in cancer genomes [23]. These challenges

underscore the urgent need for advancements in sequencing technologies and

bioinformatics approaches to enhance the detection and analysis of somatic REIs in

cancer.

The continuous advancement of third-generation sequencing (TGS) technology,

particularly Oxford Nanopore Technologies (ONT) and Pacific Biosciences (PacBio), has

significantly enhanced long-read sequencing (LRS). Technology of ONT produces reads

exceeding 10 kb, with some extending to millions of bases, enabling the complete capture

of large-scale SVs and their flanking regions in a single read. This improvement enhanced

the sensitivity and accuracy of SV detection [26,36-38]. LRS has enabled researchers to

accurately map TR alternations across populations, revealing complex TR alternations

that were previously undetectable by SRS [39,40]. LRS were also used in population studies

to uncover the widespread presence of short interspersed nuclear element (SINE)

insertions, which are more difficult to capture than long interspersed nuclear element

(LINE) insertions typically identified by SRS [36,37]. However, most population studies have

focused on germline genomic alterations, leaving the study of somatic REIs in tumor

genome largely unexplored [41-43]. While some studies have applied long-read sequencing

to examine structural variations in tumors, the lack of large tumor sample cohorts and

suitable bioinformatics tools has limited the comprehensive analyses.

In this study, we conducted long-read nanopore whole genome sequencing (LRS WGS)

on 325 paired tumor and blood samples from 12 cancer types. With specialized algorithms

developed, we identified 39,661 somatic REIs. The diversity of somatic TRIs sequence

patterns and variable TEI insertion mechanisms across cancer types were elucidated. In

total, we identified 152 high-frequency somatic TRIs, where 10/152 are shared in

diverse cancer types, including somatic TRI events were located within the intron of

*SHROOM2* and *PALMD*, and in the distal enhancer region upstream of *PTPRZ1*.

Regarding TEIs, approximately 40% of tumors harbored somatic TEIs in the MHC class II

gene cluster, with adenocarcinomas showing greater TEI enrichment than squamous cell

carcinomas. Notably, adenocarcinoma cells with a higher TEI burden within this region

showed significantly enhanced T cell activation compared to those with a lower burden,

indicating that TEIs may contribute to the regulation of tumor immune responses. Overall,

this study provides a valuable resource to the scientific and clinical community and a

paradigm for character the somatic genomic alternations in pan-cancer genomes using

LRS.

# Results

## Section 1: Specimens, Tumor Types and Dataset

This pan-cancer study encompassed 325 tumor samples and 300 paired blood samples from 12 distinct cancer types (Figure 1a). Solid tumor types were from gynecologic (ovarian [OV], uterine corpus endometrial carcinoma [UCEC], cervical squamous cell carcinoma [CCA]), core gastrointestinal (stomach adenocarcinoma [GC], colon adenocarcinoma [IC]), developmental gastrointestinal (liver hepatocellular carcinoma [HCC], pancreatic adenocarcinoma [PAAD]), head and neck (Oral squamous cell carcinoma [OSCC], Laryngocarcinom [LC]), thoracic (lung adenocarcinoma [LUAD], lung squamous cell carcinoma [LUSC]), and central nervous system cancers (glioblastoma multiforme [GBM]). 4, 2, 27 and 2 multiple synchronous tumors were obtained from 9 patients with LUAD (2), GBM (1), HCC (5) and PAAD (1), separately (Table S1). All patients provided written informed consent for genomic studies approved by local institutional review boards.

Whole-genome sequencing (WGS) was performed on all tumor and paired blood samples using PromethION nanopore sequencer, yielding long sequencing reads (Table S2). A total of 45.73 terabases (Tb) of long clean reads were generated, averaging 93.03 gigabases (Gb) per tumor sample and 51.63 Gb per paired blood sample. The mean read depth reached 30 X (13 X~46 X) for tumor samples and 17 X (3 X~43 X) for paired blood samples (Figure S1a), with mean read N50 length across all samples was 24.9 kbp (4.5 kbp~43.1 kbp) for clean long reads (Figure S1b). Due to an update in ONT technology during data acquisition, two versions of sequencing chips were employed in this study. Specifically, 83.06% (510/614) of the LRS WGS data were generated using the R9 chips, while the remaining data were obtained using the R10 chips (Figure S1c). Notably, the data produced with the R10 chips exhibited significantly higher quality compared to that generated with the R9 chips (Wilcoxon test, p < 2.2e-16) (Figure S1d). Additionally, RNA-seq datasets were generated for 89 NSCLC samples on an NGS platform [25] (Table S3).

## Section 2: Accurate identification of somatic REIs in pan-cancer genomes

In our previous research, nearly 90% somatic insertions in NSCLC genomes were identified as somatic TRIs and TEIs, where somatic TRIs were predominantly located within annotated simple repeat regions (77.8%) [25]. The high repetitiveness of these sequences complicates alignment accuracy, often resulting in a high false-positive rate. To address this issue, we designed a more precise algorithm for detecting somatic TRIs across tumor genomes (STAR Methods) (Figure 1b and S2a). First, cleaned LRS reads were mapped to the hg38 reference genome using Minimap2 software (version 2.17) [44]], achieving an average mapping rate of 99.91% (93.07%-100%). Next, widely used SV detection softwares were applied to identify loci harboring somatic TRIs. Specifically, Sniffles2 (version 2.0.6) [45] was employed to detect INSs within tandem repeat regions annotated by RepeatMasker in the hg38 reference genome. Repeat expansions (REs) in tumor genomes were then identified using Straglr (version 1.4.1) [46]. Regions identified by either tool were combined to form a final list of candidate somatic tandem repeat insertion windows for further analysis. For each paired tumor-blood samples, somatic TRIs within these candidate windows were detected based on the clustering of reads according to editing distance and the TDScope algorithm [47] (Figure S2a, STAR Methods).

Somatic transposable element insertions (TEIs) were identified by optimizing an in-house pipeline for somatic structural variant (SV) detection [25] (Figure 1c and S2b). First, tumor sample alignments were merged with those of the paired blood sample. Next, large-scale INSs were detected using Sniffles2 (version 2.0.6) [45]. A somatic INS was considered valid if more than 3 supporting reads from the tumor sample, but no read from the paired blood sample (STAR Methods). Next, Iris software (version 1.0.5)[48] was used to correct the sequences and breakpoints of the candidate somatic INSs. To false positives due to reference-individual genome differences, a custom pipeline was developed. The reference genome at each INS loci was polished using Racon (version 1.4.16)[49], based on alignments from the paired blood samples. Tumor read were then realigned to the

polished reference, and those with more than 3supporting reads were retained as

candidate somatic INSs. Finally, consensus sequences of somatic INSs were categorized

using RepeatMasker (version 4.1.2)[50], Tandem Repeat Finder (version 4.10.0) [51], and

Sdust (version 0.1)[52] (STAR Methods). Somatic INSs with consensus sequences

containing TE segments were classified as somatic TEIs (STAR Methods).

A total of 16,382 somatic TRIs (Table S4) were identified across 325 tumor genomes, with

counts ranging from 1 to 849, and an average of 50. Similarly, 23,279 somatic TEIs (Table

S5) were identified across 325 tumor genomes, with counts ranging from 0 to 1,776, and

an average of 72 (Figure 1d and S2c). No correlation was identified between tumor purity

(mean 46.33%, Table S1) and the count of somatic TRIs or somatic TEIs (Figure S3a-S3d,

STAR Methods). Squamous carcinomas (SCC) exhibited a significantly higher burden of

somatic TEIs compared to other cancer types (Welch two sample t-test, p value=0.0014

with Other, p value=1.6e-05 with Adenocarcinoma, p value=2.9e-07 with high burden)

(Figure S3e and S3f). Notably, 75.88% (3351/4416) of the somatic TEIs were specific to

SCC. However, only 43of the 3351 of the SCC-specific somatic TEIs were shared

between two SCC samples, suggesting that while TEs are more active in SCC genomes,

but SCC-specific TEIs were mostly low-frequency mutations. The distribution of somatic

TRI and TEI counts across various cancer types did not show significant differences

(Figure 1c, S3g).

Interestingly, a subset of 11 non-SCC tumor samples exhibited markedly higher numbers

of both somatic TRIs (ranging from 592 to 849) and somatic TEIs (ranging from 621 to

1776) compared to the remaining tumors (TRIs: ranging from 1 to 139, TEIs: ranging from

0 to 279) (Figure 1d and S2c). This group, termed high-burden samples, included 2 HCC

samples, 1 GBM sample, 2 GC samples, and 6 LUAD samples. Using the Genomic

Regions Enrichment of Annotations Tool (GREAT)[53], the high-burden-specific somatic

TRIs (Table S6) were linked to distal cis-regulatory effects on genes involved in cell cycle

regulation, telomere length, and chromosome localization (Figure S3h). Additionally,

transcriptomic analysis of available NSCLC NGS RNA-seq data revealed that high-burden

LUAD samples exhibited significantly lower expression of DNA repair-related genes,

including *ERCC3*, *RAD51*, and *XRCC4* (Wilcoxon rank test, p < 0.05,

|log2(FoldChange)| > 1), compared to other LUAD samples (Figure S3i). Reduced

expression of these DNA repair genes may lead to the accumulation of unrepaired DNA

damage, contributing to genomic instability [54-56].

Somatic TRIs were significantly enriched in subtelomeric regions (5.308-fold compared to

the rest region of chromosome arm in average, Student t-test, p-value = 9.81e-14),

whereas no enrichment of somatic TEIs was detected in subtelomeric regions (Student

t-test, p-value = 0.6275) (Figure S4a, Table S7). Similar to the previous LRS study [57], the

long arms of human chromosomes were more likely exhibited a broader zone of somatic

TRIs enrichment (5–10 Mbp) compared to the short arms (Figure S4b). Similar to a large

healthy Chinese cohort genomic study [58] and a recent TE recombination study [59], somatic

TEIs were frequently enriched in centromeric and pericentromeric regions, particularly on

chromosomes 20, 4, and 9 (Figure S4c). Notably, solo LINE insertions, solo SINE

insertions, and complex TEIs containing LTR segments exhibited the most pronounced

enrichment in these regions. (Figure S4c). In contrast, a comparable analysis of TEs in

the reference genome and all somatic TEIs revealed no such patterns for solo LINE

insertions.

To identify hotspot regions of somatic TRIs and TEIs, the genome was segmented into

non-overlapping 1 Mb windows, and their distribution was investigated (Figure 1e).

Hotspot regions containing genes that overlapped or were adjacent to somatic TRIs and

TEIs were identified. These included *MUC16*, associated with various cancers [60-62] and

widely used as a biomarker for ovarian cancer (CA125) [63] *STK24*, a prognostic marker for

HCC, LUAD, PAAD, and KIRC (p < 0.001, log-rank test), involved in apoptosis regulation

and cell migration[64,65]; *NBPF1*, part of the neuroblastoma breakpoint family, linked to

genomic instability in neuroblastoma and gliomas[66,67]; and *CELF2*, a prognostic marker

for HNSC and KIRC (p < 0.001, log-rank test), implicated in apoptosis and cancer

progression [68]. Genes such as *ZNF716* and *ZNF717*, which encoding zinc finger proteins

involved in transcriptional regulation and signaling [69]; *PTPN7* and *PTPRZ1*, which were

members of the protein tyrosine phosphatase family [70,71], were also identified as highly

recurrent but whose role in cancer remain to be fully characterized. In addition to genes

with established roles in tumor progression, we also identified a number of genes with

high-frequency somatic REI events across pan-cancer samples, including those involved

in cytoskeleton regulation (e.g., *SHROOM2*, *PALMD*, and *CROCC*), suggesting their

potential as biomarkers.

We next examined the enrichment of somatic TRIs and somatic TEIs in gene and

regulatory regions by comparing the real dataset ('O', observed) to datasets representing

the genomic coordinates of gene and regulatory regions ('E', expected) (Figure S4d).

52.5% somatic TRIs were located within gene regions, while 47.9% somatic TEIs were

detected in these regions. Notably, somatic TRIs were significantly enriched in candidate

cis-regulatory elements predicted by ENCODE (cCRE)) (log2(O/E) ranging from 0.27 to

2.07, binomial test, FDR < 0.05), suggesting their potential roles in chromosomal structure

and genome regulation. In contrast, somatic TEIs were enriched in regions proximal to

genebody (±2.5 kb, log2(O/E) ranging from 0.1 to 0.2, binomial test, FDR < 0.05),

implicating them in noncanonical splicing events, as previously reported [31].

## Section 3: The Sequence Features and Mechanisms of

## Somatic REIs

To explore the mutational patterns of somatic TRI, we applied pairwise alignment

algorithms to compare the consensus sequences of germline and somatic events within

the somatic TRI windows across various samples. Sequences identified exclusively in

somatic TRI events, relative to their paired germline events, were classified as somatic

TRI mutational sequences. These sequences were classified into three categories based

on the repeat motifs involved. The first category, 'Expansion', referred to somatic TRI

events where a pre-existing germline repeat motif is expanded in the somatic sample

(Figure 2a). For example, in the somatic TRI window (chr18:80,195,548-80,195,600) of

sample UCEC12, the AT motif was notably expanded (Figure S5a), reflecting an increase

in repeat length. The second category, 'Motif-Change', involved somatic TRI events where

a new repeat motif emerges, absent in the germline sequence (Figure 2a). In sample

Lung136, the somatic sequence showed the appearance of the AGATATATAT motif, which

was not present in the germline TR window (chrX:145,918,007-145,918,062) (Figure S5a).

The third category includes somatic TRI events where the inserted sequence did not

match any recognizable repeat motif, classified as 'Undefined' (Figure 2a and S5a).

As expected, 95% of somatic TRI events were primarily characterized by the expansion of

repeat motifs (Figure 2b, 2c and S5b). Among these, 67.8% involved the expansion of

pre-existing germline motifs, with the proportion of mutational sequences ranging from

40.28% to 100% (average 87.89%, median 91.48%). 16.7% were driven by the expansion

of newly emerged motifs, with the proportion of mutational sequences ranging from

50.15% to 100% (average 81.90%, median 82.12%). Furthermore, 10.5% of somatic TRI

events exhibited a combination of both germline motif expansion (average 47.85%,

median 49.03%) and the expansion of new motifs (average 45.96%, median 44.98%). In

addition to these, 2.8% of somatic TRI events were classified as 'Undefined', where the

repeat motifs in the somatic INSs could not be identified. These events had an

undetermined insertion frequency ranging from 63.25% to 99.84% (average 86.64%,

median 86.84%). Furthermore, 2.2% of somatic TRI events displayed a combination of all

three mutational types, preventing classification into a predominant pattern. These events

were categorized as "Ambiguous". Given the challenges associated with nanopore

sequencing, particularly with long tandem repeat sequences [72], we propose that the

"Undefined" and "Ambiguous" categories may be affected by sequencing errors.

In terms of the sequence pattern of TEIs, we identified two types of somatic TEIs:

'complex TEIs', whose inserted sequences comprise multiple TE fragments (Figure 2d),

and 'solo TEIs', which consist of a single TE fragment. Among the somatic TEIs, 56%

were classified as solo TEIs, while 44% were complex TEIs (Figure 2e). Complex TEIs

were significantly longer than solo TEIs (average length: 1,823.8 bp vs. 570.5 bp,

Student's t-test, p-value < 2.2e-16) (Figure S5c). Approximately 80% of complex TEIs

exceeded 300 bp, beyond the typical read length of short-read sequencing (SRS), making

them difficult to resolve with SRS. Notably, 3.4% of complex TEIs matched the known

somatic recombination pattern of Alu and L1 [33,59]. The remaining complex TEIs exhibited

more intricate compositions of TE families and inverted TE segments (Table S5) —

observations not systematically identified in SRS studies.

Somatic TEIs containing LINE elements were most prevalent (57.8%), followed by those

containing SINE elements (44.7%) (Figure 2c, S5d and S5e). The distribution of somatic

TEIs varied across different tumor types. In SCC, gastrointestinal cancers (including IC

and GC), and UCEC, somatic TEIs predominantly contained LINE segments

(45.4%-76.0%), consistent with previous SRS studies showing a higher frequency of

somatic solo L1 insertions in SCC [73,74]. These tumors exhibited a typical ~300 bp

somatic TEI profile with no distinct peaks (Figure S5f). In contrast, other tumors a higher

proportion of somatic SINE-containing somatic TEIs (54.6%-70.4%). Peaks were

identified around 300 bp and 6 kb (Figure S5f), corresponding to SINE- and

LINE-containing somatic TEIs, respectively. High-burden samples exhibited a noticeable

peak at approximately 300 bp, with 48.2% (5422/11238) being solo SINEs. These findings

suggested differential TE activity across cancer types. Notably, younger TE subfamilies,

such as AluY and L1HS, significantly contributed to the tumor genome (Figure S5g),

highlighting their activity in tumor development and their key role in shaping the somatic

TEI composition of the cancer genome.

To explore the insertion mechanisms, we analyzed the sequences features of somatic

TEIs (Supplementary Note). By aligning somatic TEI sequences with the flanking regions

of insertion sites (STAR Methods), we identified 82.1% of somatic TEIs contained

homologous sequences. Based on the positions and lengths of homologous sequences,

we classified somatic TEIs into two mechanisms: retrotransposition and double-strand

break (DSB) repair via homologous recombination[59]. 58.2% of somatic TEIs were

associated with target site duplications (TSDs), a hallmark of retrotransposition (Figure 2g

and Figure S6a). The median TSD length was 14.6 bp (Figure S6h), consistent with

previous findings [74]. 31.3% of somatic TEIs exhibited homologous sequences (average:

760.8 bp) on the one side of the breakpoint, suggesting the homologous recombination.

Among these, 12.3% had homologous sequences longer than 100 bp (average: 262 bp,

ranging from 101 bp to 19,659 bp), similar to the non-allelic homologous recombination

(NAHR) mechanism described previously ([59,75] (Figure 2g and Figure S6b). 12.9% of

somatic TEIs carried shorter homologous sequences (<100 bp, average: 7 bp), indicating

microhomology-mediated recombination (Figure 2g and Figure S6c). The remaining

somatic TEIs were classified as 'other' indicating unknown mechanisms (Figure 2g and

Figure S6d).

We further uncovered the mechanisms underlying the formation of solo and complex TEIs.

Somatic solo TEIs were predominantly generated via retrotransposition (69.2%), with only

17.4% arising from recombination (Figure 2e). In contrast, complex TEIs exhibited a more

balanced distribution, with 38.9% arising from recombination and 40.3% by

retrotransposition (Figure 2e). Notably, the mechanisms of somatic complex TEIs varied

across cancer types. GC, GBM, and HCC had a larger proportion of somatic complex

TEIs generated by NAHR recombination (Chi-square test, p-value < 2.2e-16, Fold

Change = 2.56). Digestive system cancers (GC, IC, PAAD, and HCC) displayed a larger

proportion of somatic complex TEIs with unidentified mechanisms (Chi-square test,

p-value < 2.2e-16, Fold Change = 1.96). These findings suggest that the formation of

somatic complex TEIs is characterized by significant stochastic variability rather than a

strictly ordered process.

# Section 4: High-Frequency Somatic TRI Events among samples are potential Pan-Cancer Biomarkers

To identify non-random somatic TRI events, we compared the observed frequency

distribution of somatic TRI events with the expected distribution under random conditions.

This analysis revealed that 152 somatic TRI events, each detected in ≥10 patients, are

exceptionally rare under the random distribution model (permutation test, p-value ≤ 0.002;

STAR Methods, Figure S7a). Notably, three of these events have also been reported as

recurrent repeat expansion events (rREs) with subtype-specific characteristics in

previous pan-cancer studies of somatic simple repeat expansions[34] (Table S4). We

therefore defined these somatic TRI events as high-frequency somatic TRI events. Similar

to the genome-wide distribution pattern of somatic TRI events, we identified that

high-frequency TRI events are significantly underrepresented in coding sequence (CDS)

regions compared to the genome-wide background (binomial test, FDR = 0.0425; Figure

S7b). This finding suggested that the high-frequency somatic TRI events are more likely to

exert their downstream effects through alterations in non-coding regions, potentially

impacting regulatory processes. Previous studies have demonstrated that tandem repeat

(TR) sequence variations contribute to genomic regulation primarily by modulating

transcription factor (TF) binding sites [76] or by introducing CpG dinucleotide sequences

that influence methylation patterns, thereby affecting downstream gene transcription [77].

High-frequency somatic TRI events are significantly enriched in 10 out of 490 ENCODE

TF binding regions (Table S9) compared to low-frequency TRI events (Chi-square test,

p-value ≤ 0.05, absolute ratio difference > 5%; Figure S7c). Among these, the most

notable enrichment was observed in CTCF binding sites, where high-frequency somatic

TRIs displayed an odds ratio of 2.032 (p-value = 3.3e-4). This suggested that

high-frequency somatic TRIs may be involved in 3D chromatin structure alterations and

distal regulatory processes [78]. Additionally, when comparing somatic sequences of

high-frequency TRIs to their germline counterparts, we identified significant differences in

CpG dinucleotide counts in 62 out of 152 high-frequency somatic TRIs (Wilcoxon rank test,

FDR < 0.05, |log2(Fold Change)|>1, Figure S7d, Table S10). Leveraging the unique ability

of nanopore sequencing to simultaneously detect sequence variations and methylation

states [79], we found that 18 out of these 62 high-frequency somatic TRIs introduced

methylation marks at somatic extended CpG dinucleotides (Wilcoxon rank test, p-value ≤

0.05, Figure S7e, Table S11). For instance, in the sample Lung120, we investigated that a

somatic TRI region    (chr6:64,295,338–64,295,575) within an intron of the *EYS* gene

introduced 55 methylated CpG sites (Figures S7f-S7i, Supplementary Note). As

mentioned in the previous study [77], these somatic TRIs may act as potential regulatory

elements.

To investigate the association between high-frequency somatic TRIs and cancer-related

biological processes, we performed functional enrichment analysis on genes neighboring

high-frequency somatic TRIs. Compared to low-frequency somatic TRIs, genes

associated with high-frequency somatic TRIs were significantly enriched in functional

categories related to focal adhesion, apical junctions, and cytokine-cytokine receptor

interactions (Figure 3a, Chi-square test, p-value < 2e-4). Among these, the somatic TRI

event located at chrX:9,791,969–9,792,160, within the intron of SHROOM2, which is

associated with apical junction functions, exhibited the highest shared frequency across

patients (57 out of 299, Figure 3b, Table S12). Using targeted PCR and Sanger

sequencing, we validated this somatic TRI event in 25 paired tumor and blood samples

(Figure 3c, Figure S8a, Table S13 and Table S14). Additionally, this SHROOM2-related

TRI was detected in liver cancer and intestinal cancer cell lines (Figure S8b). Overall, this

somatic TRI event was present in nearly 20% of pan-cancer samples (57 out of 299) and

occurred across 11 of the 12 cancer types analyzed. Notably, this somatic TRI exhibited

particularly high recurrence rates in HCC (58.82%; 10 out of 17) and IC (40%; 6 out of 15).

Intriguingly, all multi-site biopsies from three HCC patients with this somatic TRI

consistently contained it, while biopsies from two HCC patients without this somatic TRI

consistently lacked it. This result indicated a high degree of intra-tumor consistency for

this somatic TRI. Based on length distribution, somatic TRI sequences within this window

were categorized into two groups: long-fragment expansions and short-fragment

expansions (Figure 3d). The long-fragment expansions ranged from 3,896 to 8,284 bp,

whereas the short-fragment expansions ranged from 639 to 3,259 bp (Figure 3e).

Interestingly, the two cancer types with high recurrence rates (HCC and IC) predominantly

carried somatic TRIs with long-fragment expansions (HCC: 7 out of 10; IC: 5 out of 6;

Figure 3f, Fisher's exact test, p-value = 3.29e-6). These long-fragment expansions were

primarily characterized by the amplification of the AGAAT repeat motif (Figure S9, Table

S15).

Subsequently, using our NSCLC NGS RNA-seq dataset, we performed differential gene

expression analysis within a 300 kb window surrounding all high-frequency somatic TRI

events detected in NSCLC. This analysis identified 51 differentially expressed genes

associated with 33 high-frequency somatic TRI events (Figure S10a, Table S16). Among

these, we identified the somatic TRI event located at chr7:121,603,262–121,603,506,

which was previously reported in our study to be present in nearly 40% of LUSC samples

(Figure S10b and S9c). This somatic TRI was shown to regulate the expression of its

downstream *PTPRZ1* gene by altering the 3D genomic structure [25]. Additionally, we

identified two other somatic TRI events associated with the expression of genes in the

PTP family: the somatic TRI at chr1:202,204,213–202,204,518, which was correlated with

the upregulation of *PTPN7* expression (Figure S10b and S9d), and the somatic TRI at

chr7:158,595,097–158,595,342, which was associated with the downregulation of

*PTPRN2* expression (Figure S10b and S9e). Notably, while the somatic TRI at

chr7:121,603,262–121,603,506 was significantly enriched in LUSC and LC samples, the

other two PTP-related somatic TRI events did not exhibit enrichment in specific cancer

types (Figure S10f). Furthermore, we noticed the presence of these three TRI events

across the pan-cancer cohort, encompassing 66 samples.

By performing clustering analysis based on the distribution spectrum of somatic TRI

events, we identified that samples from HCC, GBM, GC, and LUAD formed four distinct

clusters (Figure 4a). Notably, GC, GBM, and LUAD samples did not harbor somatic TRI

events with an intra-cancer frequency exceeding 30% (Figure 4b), indicating a high

degree of inter-patient heterogeneity for TRI events within these cancer types. In total, we

identified 12 somatic TRI events that were detected within single cancer types with high

frequency (Figure 4c, STAR Methods). HCC harbored the highest number of intra-cancer

high-frequency somatic TRI events (4 out of 12; Table S16, Figures 4d). These included a

previously described somatic TRI event located within the SHROOM2 intron, as well as

three additional events: one within the PALMD intron, predominantly characterized by

expansions of the AAAG motif (chr1:99,683,035–99,683,173, Figure S11a), which is

consistent with previous reports in other pan-cancer somatic tandem repeat study [34]

（Table S4）; another within the *LIPG* intron, mainly involving expansions of the GGGAGA

motif (chr18:49,579,001–49,579,348, Figure S11b); and an intergenic event featuring

expansions of the GGTTT motif (chr1:34,802,227–34,802,447, Figure S11c). Notably,

these HCC-specific high-frequency somatic TRI events were also recurrently detected in

IC, UCEC, OV, OSCC, and LC. Genes such as *SHROOM2* and *PALMD* are implicated in

the regulation of the cytoskeleton [80-83], while *LIPG* is involved in lipid metabolism [84].

High-frequency somatic TRIs shared across different cancer types may suggest common

malignant biological behaviors, making them potential diagnostic and therapeutic

biomarkers for these cancers.

Additionally, digestive system adenocarcinomas (IC and PAAD), gynecological tumors

(CCA, OV, and UCEC), and three SCCs (OSCC, LC, and LUSC) formed three distinct

clusters based on the frequency of somatic TRI events (Figure 4a). Each cluster exhibited

specific somatic TRI events. Within the IC-PAAD group, we identified three somatic TRI

events with significantly higher frequencies compared to other cancer types (Student's

t-test, p-value < 0.05): a somatic TRI event located in the *LRBA* intron

(chr4:150,658,505–150,658,775; Figure S11d), another in the *SLC22A2* intron

(chr6:160,218,690–160,218,725; Figure S11e), and the third upstream of the *SPINK1*

gene (chr5:147,837,649–147,837,716, 6073 bp upstream; Figure S12a). These genes are

associated with tumor cell growth (*LRBA*) [85], chemotherapeutic resistance (*SLC22A2*) [86],

and tumor dedifferentiation (*SPINK1*) [87], respectively. In contrast to the IC-PAAD group,

gynecological tumors (OV, CCA, and UCEC) formed a distinct cluster in the somatic TRI

frequency spectrum, but exhibited low recurrence rates for their specific somatic TRI

events. Two cluster-specific somatic TRI events (Student's t-test, p-value < 0.05) included

one upstream of the ENSG00000257323 gene (chr12:74,615,532–74,615,609; Figure

S12b) and another within the *MED16* intron (chr19:878,845–879,150; Figure S12c).

These events appeared at frequencies below 30% within any individual gynecological

tumor type. The SCC cluster, consisting of OSCC, LC, and LUSC, exhibited three

cluster-specific high-frequency somatic TRI events (Figure 4f). These included the

previously described *PTPRZ1*-related somatic TRI event (Figure S10c), as well as two

somatic TRI events located in the promoter of the OBI1 gene

(chr13:78,659,295–78,659,339; Figure 4g) and the *SLC10A6* intron

(chr4:86,842,786–86,842,877; Figure S12d). In the available samples, we validated the

cluster-specific high-frequency TRI events through PCR experiments and Sanger

sequencing, highlighting the existence of similar somatic TRI distribution patterns among

cancer types within the same cluster (Figure S12, Table S14).

## Section 5: somatic TEIs were enriched in the MHC II gene cluster

To investigate the somatic TEI distribution across cancer types, we clustered samples

based on somatic TEI breakpoints, identifying five distinct clusters (Table S1) with unique

histological compositions (Figure 5a). Cluster 1 predominantly comprised SCC samples

(67%) (Figure 5b and 5c), including 87.5% of LUSC and 68.8% of OSCC cases (Figure 5c

and S14a). Notably, approximately 20% of ADC samples were also assigned to Cluster1,

with IC (73.3%) and OV (50%) being the most represented (Figure 5c and S14a). This

suggested that IC and OV share TEI characteristics with SCC samples in Cluster 1.

Moreover, ADC samples in Cluster 1 exhibited lower somatic TEIs burden (mean: 36.7,

median: 19) compared to those in other clusters (mean: 156, median: 23) (Welch

two-sample test，p = 0.005), indicating the distinct somatic TEI profiles of ADC samples

within Cluster 1. 57.7% of Cluster 1 SCC samples further subdivided into two subclusters.

In Subcluster 1, 75% of the samples were SCC, while in Subcluster 2, nearly 60% of the

samples were SCC (Figure 5d and S14b). Subcluster 1 exhibited a significantly higher

somatic TEI burden than Subcluster 2 (Figure S14c), highlighting the substantial

molecular heterogeneity within SCC samples.

Cluster 2 encompassed tumor samples with the highest levels of TEI burden, including all

high-burden samples and two multi-site biopsies from a LUAD patient, which ranked just below the high-burden samples. Although multi-site biopsies were collected from 9 patients, with samples from the same patient clustering together, only the complete set from patient HCC10 was assigned to Cluster 3, suggesting a potential association between he multi-lesional nature of the patient and their distinct somatic TEI profile.

Clusters 4 and 5 were predominantly composed of ADC samples (68% and 66%, respectively) but differed in somatic TEI burden and distribution. Cluster 4 exhibited a higher somatic TEI burden and a greater degree of overlap between samples (Figure 5e), indicating a more homogeneous TEI profile. This cluster also included poorly differentiated, non-metastatic CCA samples and early-stage (I/II) HCC biopsies, which clustered with multi-site samples from a stage I LUAD patient. In contrast, cancers commonly associated with high heterogeneity, such as PAAD, UCEC, and GBM, were largely detected in Cluster 5, with PAAD samples showing duodenal or peripancreatic invasion (Figure S14a, Table S1).

We analyzed high- and low-frequency somatic TEI regions across the clusters, covering 1,464 1Mb genomic windows. In Cluster 1 and Cluster 5, we identified 15 and 6 regions, respectively, with significantly low-frequency somatic TEIs (Fisher's exact test, FDR < 0.01, Fold change < 0.5, and >10% of non-cluster samples showing somatic TEIs in the same window). In contrast, Cluster 2, Cluster 3, and Cluster 4 contained 1384, 45, and 14 regions, respectively, with significantly high-frequency somatic TEIs (Fisher's exact test, FDR < 0.01, Fold change > 2, and >10% of cluster samples exhibiting somatic TEIs in the same window) (Figure S14d, Table S17).

To identify non-randomly enriched TEI windows, we applied a permutation test (STAR Methods). In windows with ≥14 patients, somatic TEI frequency was significantly higher than in randomly selected background regions. This suggested that these windows are non-randomly enriched with somatic TEIs (permutation test, p-value=0.0002, Figure 6a). Given the high sequence variability and repetitive nature of centromeric and

pericentromeric regions, as well as the limited functional data and reference genome

quality in these areas, we excluded them from further analysis. After exclusion, the

chr6:32,000,000-33,000,000 window, containing the MHC II gene cluster, exhibited the

highest somatic TEI count and the greatest proportion of TEI-positive samples (40.6%)

(Figure 6b). Particularly in Cluster 1 (67% SCC samples), the proportion of samples

harboring TEI (27.8%) is significantly lower than that in clusters predominantly composed

of non-SCC samples (49.7%) (Fisher's exact test, FDR=0.0007).

Somatic TEIs in this region were primarily located in the intergenic and coding regions of

the *HLA-DRB1*, *HLA-DRB5*, and *HLA-DRB6* genes (Figure 6c). The distribution of

somatic TEIs varied across cancer types. High-burden samples showed widespread

somatic TEIs around the MHC class II gene cluster, while LUAD samples showed higher

frequencies of somatic TEIs than LUSC (Figure 6c). Notably, in the *HLA-DRB1* genebody

and its 5 kb flanking regions, somatic TEIs were absent in LUSC, but frequent in LUAD

(Fisher's exact test, FDR = 0.00066). Most somatic TEIs in this region were solo SINEs

(Figure S15a), consistent with previous findings of LINE enrichment in SCC samples.

Next, we examined the impact of somatic TEIs in the chr6:32,000,000-33,000,000 region

on gene expression using transcriptomic data from LUAD tumor samples. Comparison of

32 LUAD samples with somatic TEIs in this region to 39 samples without revealed 1,798

differentially expressed genes (p value < 0.05, |log2(FC)| > 1, Table S18, Figure S15b).

These genes were enriched in cytokine signaling and immune response pathways (Figure

S16a and S16b). 6 immunoglobulin-related and 4 T-cell receptor-related genes were

significantly downregulated in LUAD samples with somatic TEIs (Figure S15c). These

results suggested that TEIs in this region may contribute to immune evasion and tumor

progression.

Next, we further validated our bioinformatics findings through experiments. First, in a

subset of samples that underwent LRS, the authenticity of the TEI event was gauged by

Polymerase Chain Reaction (PCR). With blood DNA serving as a reference, when a TEI

event was present in the tumor tissue sample, the target band obtained from the tumor

tissue sample exceeded that of the blood tissue under identical primer amplification

conditions (Figure S15 d and e, Table S19). Moreover, through the utilization of specific

fluorescence in situ hybridization (FISH) probes (Table S20), TEI events were also

discerned in certain sample tissue sections (Figure 6e).   It is widely recognized that

tumor tissues encompass a rich and diverse spectrum of cell types, inclusive of infiltrating

immune cells. Considering that the TEI insertion site is located in the MHC II gene cluster

region, we gathered additional fresh clinical samples (Table S21).   By sorting CD45

positive cells, we further amplified the TEI events identified in this region within the clinical

sample LRS sequencing outcomes. The findings confirmed that these insertions occurred

in both CD45 positive and negative cells (considered as tumor cells) (Figures S17).

Subsequently, we detected TEI events within the MHC II gene cluster in 33 tumor cell

lines stored in the laboratory. In alignment with the sequencing results in clinical samples,

non-squamous carcinomas harbored a greater number of TEI events in contrast to

squamous carcinomas (a total of 63 TEIs were detected, with an average detection of

11.07 in squamous carcinomas and 19.35 in non-squamous carcinomas) (Figures S18,

S19, and S20a).

MHC class II molecules play a crucial role in cellular immunity. In antigen-presenting cells

(such as macrophages, dendritic cells, and B cells), MHC class II molecules are involved

in the processing and presentation of ingested exogenous antigens into peptide

fragments, and presenting them to CD4+ T cells, thereby initiating the adaptive immune

response[88]. Here, we compared the in vitro phenotypes of CD40L activation by TEI-enrich

and TEI-impoverished tumor cells through in vitro co-culture of tumor cells and mouse

CD4-positive t cells. The results showed that in squamous cell carcinoma cell lines, the

surface expression of CD40L on TEI-rich tumor cells was twice that of TEI-impoverished

cells (Figure 6e and Figure S20b); while the activation situation is reversed in

adenocarcinoma cells (Figure 6e and Figure S20b), this all suggested that the frequency

of TEI events and the sequence type of TEI event insertions may be important factors in

regulating the success of tumor adaptive immune initiation.

# Discussion

This study was designed to elucidate high-frequency somatic large-scale REIs (>50bp) in

the pan-cancer genome. Previous research has been constrained by the limitations of

SRS technology, which hindered the precise identification of these somatic SVs and their

distribution across tumor population. LRS technology offers a promising solution to these

challenges [89]. In this work, we applied nanopore LRS to sequence 325 tumor samples and

their matched blood samples from 12 cancer types, generating nearly 46 Tb of data. This

extensive LRS dataset, one of the largest to date, provides a robust resource for exploring

somatic large-scale REI events across the pan-cancer landscape.

Current LRS-based somatic SV detection algorithms are hindered by alignment biases,

particularly in repetitive genomic regions. These biases, stemming from the

seed-and-chain alignment strategy, undermine the accuracy of insertion detection [89,90].

Although assembly-based approaches hold promise, they depend on high-quality

individual genome assemblies, which are challenging to produce in the context of

heterogeneous tumor genomes [91,92]. To overcome these limitations in detecting

somatic REIs, we developed specialized bioinformatics methods tailored for somatic TRI

and TEI detection, respectively. By harnessing the technical strengths of long reads that

span somatic insertions, and combining unsupervised learning with partial order

alignment graphs and statistical analysis of local sequence features, we minimized errors

introduced by reference genomes and alignment algorithms, thus enabling more accurate

detection of somatic REIs. Recent advancements in long-read sequencing (LRS)

technology, alongside continuous improvements to the human genome

reference—particularly the completion of the Telomere-to-Telomere (T2T) project [93]—have

significantly enhanced the study of large-scale repetitive sequence variations, such as

centromeric and segmental duplications [94]. This study also benefits from these

developments, uncovering numerous related somatic REIs. However, the structural

complexity and dynamic nature of these regions, marked by variation in repeat copy

number and sequence divergence, present substantial challenges to accurate detection.

Consequently, refined bioinformatics methods and ultra-long sequencing technologies are

expected to provide critical solutions to these issues [95].

Using our developed methodology, we identified a significant number of high-frequency

somatic REIs across pan-cancer genomes. In addition to the somatic TRI events located

within the introns of the *SHROOM2* and *PALMD* genes, which exhibited remarkably high

recurrence in HCC, IC, CCA, UCEC, OV, LC, OSCC, and LUSC. Interestingly, cancers

from anatomically neighboring tissues tend to exhibit similar somatic TRI profiles. For

example, the gynecological cancers CCA, UCEC, and OV shared a similar somatic TRI

spectrum, with a somatic TRI event in the intron of *MED16* being specific to these tumor

types. The *MED16* gene is associated with estrogen receptor sensitivity in the previous

research [96], the gynecological cancers specific somatic TRI event may play a role as a

regulatory element of this gene. Similarly, squamous cell carcinomas, including LUSC, LC,

and OSCC, displayed similar somatic TRI profiles. Notably, a somatic TRI event identified

in approximately 40% of LUSC samples was found to occur at an even higher frequency

in LC samples (nearly 70%). This event has been shown to influence the

three-dimensional structure of the genome, thereby modulating the expression of the

downstream gene *PTPRZ1* through enhancer activity [25]. In addition to *PTPRZ1*, further

analysis of paired RNA-seq data from non-small cell lung cancer (NSCLC) revealed two

additional somatic TRI events (chr1:202,204,213-202,204,518 and

chr7:158,595,097-158,595,342, Figure S10a and S10b), which were significantly

associated with the expression of their downstream genes, *PTPN7* and *PTPRN2*. Notably,

20.31% of pan-cancer patients carried at least one somatic TRI event potentially

associated with the PTP gene family. As an important receptor kinase family, the PTP

genes play a crucial role in immune evasion mechanisms [71]. We propose that somatic TRI

events may function as key distal regulatory elements for the expression of this gene

family. Further investigation into the transcriptional regulatory mechanisms underlying

these somatic TRI events will be essential to understanding the role of the PTP gene

family in immune regulation. Understanding the impact of the high-frequency somatic TRI

events we identified on the tumor genome network is a key focus for future research.

The success of immune checkpoint inhibitors in certain cancers has demonstrated that

the immune system can target and reject tumors through T cell responses. Tumor cell

recognition relies on MHC-mediated presentation of tumor-associated antigens. Our

research reveals that somatic TEIs are enriched in MHC II gene regions, with over 40% of

pan-cancer samples harboring somatic TEI events in these regions. Notably, LUAD

exhibited a significantly higher somatic TEI burden in MHC II regions compared to LUSC

(Figure 6c). Although the immunological effects of these somatic transposable element

insertions (TEIs) have not been fully elucidated, co-culture experiments indicated that

TEI-enriched adenocarcinoma cell lines are more potent in suppressing CD4+ T cell

activation compared to TEI-impoverished counterparts. Furthermore, LUSC patients have

been shown to derive greater clinical benefit from PD-1 inhibitors than LUAD patients [97].

Given that TEI-enriched LUAD cell lines suppress CD40L expression on CD4+ T cells, we

hypothesize that immune activation within these tumors may be constrained. Since PD-1

inhibitors function to counteract PD-L1-mediated immune suppression of CD8+ T cells,

their efficacy in TEI-enriched LUAD may be diminished due to the absence of CD4+ T cell

activation. Nevertheless, this hypothesis warrants further in vivo validation. Moreover, we

propose that TEI enrichment in MHC II regions could serve as a potential target for tumor

immune evasion. Yet, the precise mechanisms by which TEIs modulate immune

responses remain poorly understood, highlighting a critical area for future investigation in

tumor immunology.

Here, we leverage LRS technology to investigate somatic REIs and their potential as

diagnostic and therapeutic biomarkers in cancer. By integrating optimized bioinformatics

algorithms and advancing long - read sequencing technologies, future studies will be able

to systematically characterize diverse somatic genomic alterations, such as translocations

and inversions, from the dataset we provided. Additionally, integrating multimodal

nanopore sequencing data will enable single-cell resolution analysis of large genomic

regions and their associated epigenetic modifications, as demonstrated in recent studies

[98]. These efforts will provide critical insights into the regulatory mechanisms underlying

large-scale somatic structural variations (SVs) in cancer genomes, shedding light on their

functional impact. While our current dataset offers valuable insights into pan-cancer

diagnostics, expanding the cohort to include diverse cancer types and enhancing clinical

follow-up data—such as treatment responses and patient survival—will enable a

comprehensive exploration of the relationship between somatic SVs and cancer

outcomes. To achieve this, we are actively expanding our research cohort to encompass a

broader spectrum of tumor types and patient demographics, ensuring robust

characterization in future studies.

In conclusion, our study demonstrates the transformative potential of long-read

sequencing in cancer genomics. By expanding the understanding of the genetic variants

of cancer, this work sets the stage for future studies that will explore the role of SVs in

human disease. As LRS technology and analysis methods evolve, we anticipate these

advancements will enable more accurate SVs detection and a deeper insight into their

functional consequences, ultimately guiding the development of targeted therapies and

personalized treatment approaches in oncology.

## STAR Methods

### Patient Cohort and Ethics

The matched tumor-blood samples across 8 cancer types (HCC, GBM, PAAD, LUAD, IC,

GC, LUSC, and LC) were collected from patients who undergone surgery at our hospital.

The matched tumor-blood samples across 3 gynecological cancer types (CCA, OV and

UCEC) were collected from patients who undergone surgery at West China Second

University Hospital, Sichuan University. And the matched tumor-blood samples of OSCC

were collected from patients who undergone surgery at West China Hospital of

Stomatology, Sichuan University. None of the patients were received preoperative

chemotherapy or radiotherapy. All diagnoses confirmed through histological review by a

board-certified pathologist. Cancer clinical stage was determined based on the 8th edition

of the American Joint Committee on Cancer (AJCC) TNM staging system. This study

protocol was approved by the Institutional Review Board of our hospital (Ethics: Project

identification code: 20242416). Written informed consent was obtained from all patients.
Demographic and clinical data were collected for analysis(Table S1).
722

## PromethION whole genome Sequencing

High-molecular-weight (HMW) genomic DNA (gDNA) was extracted from tissue or blood
samples using the MagAttract HMW DNA Kit (QIAGEN). DNA quality was assessed by
Novogene Co. LTRI. using three standard methods: (1) DNA purity was determined by
measuring OD260/280 and OD260/230 ratios with a NanoDrop spectrophotometer, (2)
agarose gel electrophoresis was used to evaluate DNA degradation and detect RNA
contamination, and (3) DNA concentration was quantified with the Qubit® DNA Assay Kit
and Qubit® 3.0 Fluorometer (Life Technologies, CA, USA).
731

For library preparation, 273 samples (84%) were processed using the SQK-LSK109 kit,
while 52 samples (16%) were prepared with the SQK-LSK114 kit (Oxford Nanopore
Technologies, UK), following the manufacturer's protocols (Table S1). For each sample,
48 µL of gDNA (2 µg) and nuclease-free water (NFW) were combined with 3.5 µL of
NEBNext FFPE DNA Repair Buffer, 2 µL of NEBNext FFPE DNA Repair Mix, 3.5 µL of
NEBNext Ultra II End Prep Reaction Buffer, and 3 µL of NEBNext Ultra II End Prep
Enzyme Mix. The mixture was gently combined in a PCR tube, spun down, and incubated
sequentially at 20 °C for 5 minutes and 65 °C for 5 minutes. The prepared sample was
transferred to a 1.5-mL tube, and 60 µL of AMPure XP beads was added. After gentle
mixing and two washes with 70% ethanol, the purified DNA was combined with 25 µL of
Ligation Buffer (LNB), 10 µL of NEBNext Quick T4 DNA Ligase, and 5 µL of Adapter Mix
(AMX). This mixture was incubated at 25 °C for 20 minutes. Subsequently, 40 µL of
AMPure XP beads was added, and the pellet was resuspended in 25 µL of Elution Buffer
(EB).
746

For sequencing preparation, 30 µL of flush tether (FLT) was added to PromethION Flush
Buffer (PFB), and 500 µL of the Priming Mix was loaded onto a PromethION flow cell and
equilibrated for 5 minutes at room temperature. The library preparation involved mixing 75
µL of sequencing buffer (SQB), 51 µL of loading buffer (LB), and 24 µL of DNA library.
This was followed by loading 500 µL of the Priming Mix and 150 µL of the library onto the
flow cells. DNA sequencing was performed on flow cells as per the manufacturer's
instructions. Samples processed with SQK-LSK109 were sequenced on R9 flow cells
(84%), while those prepared with SQK-LSK114 were sequenced on R10 flow cells (16%).
Base calling was performed using Guppy 3.2.8 with default parameters in real time during
sequencing.
757

## Short-read RNA-seq Data Analysis

For some NSCLC samples, paired-end reads were aligned to the hg38 reference genome using HISAT2 [99] (version 2.1.0). SAMtools [100] was used to sort and index the bam files. StringTie [101](version 1.3.6) was used to generate the GTF files containing the read count information and calculate the TPM matrix of gene expression for all genes across all samples. The raw count matrix of gene expression for all genes across all samples is calculated by htseq-count [102] (version 2.0.2) based on the gencode annotation file version 42.

## Detection of differential expression gene

In order to defined potential gene associated to somatic REIs, we classified NSCLC samples into two groups based on whether harboring target somatic REI event. Then the gene expression normalized and compared between two groups through DESeq2 [103]. Genes with p-value <= 0.05 and |log2fold-change| > 1 were defined as differential expression genes.

## Oxford Nanopore Sequencing Data Processing

Initially, raw nanopore sequencing data were processed using NanoFilt (v2.8.0, https://github.com/wdecoster/nanofilt.git, parameters: -q 7 -l 1000) to filter the fastq files and generate clean reads. The quality of these filtered reads was then assessed using NanoPlot [104](v1.39.0, parameter: --maxlength 40000). Next, the ONT long reads were aligned to the hg38 reference genome using Minimap2 [44] (v2.2.17) with the following parameters: --MD, -ax map-ont, -L, and -t 30.

## Tumor purity calculation

Somatic variants (SNVs) and copy number alterations (CNVs) were identified in tumor samples using multiple approaches. For LRS WGS sequencing data, SNVs were called using ClairS [105] (version 0.1.2) with parameters set to --platform ont_r10 --min_coverage 3. CNVs were detected using Control-FREEC [106] (version 11.6) with default settings. For the NGS WGS dataset, SNVs were identified in both tumor and matched blood samples using Mutect2 from GATK [107] (version 4.1.2.0). The resulting variants were filtered using GATK's FilterMutectCalls tool. CNVs in the WGS dataset were also detected using Control-FREEC with default parameters. Tumor purity was then estimated using both SNVs and CNVs with ABSOLUTE [108] (version 1.0.6), applying the following parameters: min.ploidy = 0.5, max.ploidy = 8, max.sigma.h = 0.2, max.neg.genome=0.005, sigma.p=0, max.as.seg.count=1500, and max.non.clonal=100.

## Somatic TRI Identification

Considering the high false-positive rate associated with detecting somatic SVs of the
tandem repeat type using current algorithms, we have developed a more accurate method
for identifying somatic tandem repeat insertions across pan-cancer samples (Extended
Figure 2a). Our preliminary screening for potential somatic tandem repeat insertion sites
involved the integration of two algorithms: Sniffles2 and Straglr, to pinpoint candidate
regions. We initially employed Sniffles2 (version 2.0.7) with parameters --minsvlen 50
--mapq 5 --minsupport 3 to analyze tumor LRS WGS data, detecting INS SV intervals.
These intervals were then intersected with tandem repeat regions annotated in the hg38
genome by RepeatMasker, utilizing the bedtools intersect command, with
RepeatMasker-annotated tandem repeats as the initial candidate windows. Subsequently,
we performed a secondary screen with straglr (version 1.4.1) using parameters
--min_str_len 2 --max_str_len 100 --min_ins_size 100 --genotype_in_size --min_support 2
--max_num_clusters 9 on the tumor LRS WGS data, uncovering additional potential
tandem repeat regions not identified by the sniffles2 workflow. These regions were added
to the pool of candidate somatic tandem repeat intervals using the bedtools 'intersect'
command. All candidate windows from each sample were further refined using the
TDScope-simple command to efficiently filter for windows likely to contain somatic tandem
repeat variations. We then consolidated all candidate somatic tandem repeat windows
across samples using bedtools merge -d 200, yielding a comprehensive set of candidate
windows for the pan-cancer cohort. Utilizing TDScope (version 1.0) with default
parameters, we assessed these windows in both cancer samples and their corresponding
normal blood samples. Candidate somatic tandem repeat intervals were identified by
filtering for intervals with ≥3 supporting reads in normal blood samples, ≥3 supporting
reads in tumor samples, an SV length (SVLEN) of ≥50, and a somatic genotype number =
1.
821

## Classification of Somatic TRI by Mutation Type

To annotate the consensus sequences of tumor and blood in TRI windows, we employed
RepeatMasker, pyTRF (https://github.com/lmdu/pytrf), and mTR[109]. Due to the superior
performance of pyTRF in annotating simple repeat sequences, we prioritized its
annotation results. The remaining unannotated sequences were subsequently annotated
using RepeatMasker and mTR. To eliminate inconsistencies between the different
software tools, we normalized motifs of simple repeats, accounting for strand orientation
and repeat unit size. This allowed us to obtain a consistent annotation of tumor and blood
consensus sequences.
831

Pairwise alignment of the tumor and blood consensus sequences was performed to
identify regions in blood characterized by gaps corresponding to insertions in tumor.

Annotation of the tumor insertion sequence was compared with all annotations of the
blood sequence in the corresponding TRI window. If the tumor insertion motif matched the
blood motif, the mutation type was classified as "Expansion." If the motifs differed, it was
categorized as "Motif-Change." If no motif was annotated for the tumor insertion, the
mutation type was classified as "Other." Using this approach, we determined the length of
each mutation type in the tumor insertion sequence.
840

**Detection of Differential Somatic TRI Methylation Levels**
To identify somatic TRI windows with significantly altered CpG patterns, we compared the
CpG counts in all high-frequency TRI window pairs between somatic and germline
sequences. A paired Wilcoxon rank-sum test was employed to assess differences in CG
counts between somatic and germline samples. P-values were adjusted for multiple
comparisons using the Benjamini-Hochberg method to calculate the false discovery rate
(FDR) for each window. The relative change in CpG counts between somatic and
germline sequences for each window was expressed as the log2 fold change
(log2(FoldChange)), calculated as: $log2FolgChange = log2(mean(somaticCG)+1\ /$
$mean(germlineCG+1))$. High-frequency TRI windows with FDR ≤ 0.05 and
|log2(FoldChange)| ≥ 1 were defined as significant CpG alteration windows, and these
were used to compare methylation level changes in somatic sequences relative to
germline sequences.
854

Specifically, reads from tumor samples classified by the TRIscope algorithm as 'somatic
support' or 'unsupport' reads were aligned to reference genomes where somatic and
germline sequences had been replaced, using Minimap2 (version 2.2.17) with the
following parameters: --MD -ax map-ont -L. The methylation levels of somatic and
germline TRI events were then calculated using the f5c pipeline (version 1.3, parameters:
call-methylation -t 20 --iop 40).
861

862   To identify CpG insertion sites in somatic TRIs relative to germline TRIs, we used a
863   pairwise alignment comparison algorithm (Python package: pairwise2 from the Biopython
864   library). Based on these comparisons, we calculated methylation level changes for the
865   inserted CpG sequences in somatic TRI events.
866

**Refinement Of High-confidence Somatic TEI Candidates**
Samtools (version 2.8.0) was employed to tag BAM format files with the following
command: samtools view -H "$bam_sort"; samtools view "$bam_sort" | awk
'BEGIN{OFS="\t"}{$1="tag"$1; print $0}') | samtools view -S -b - -o "$bam_sort_tag".
Following tagging, Samtools was again used to merge the tagged BAM files from tumor

and paired blood samples (samtools merge -h "$blood_bam" "$merge_bam" "$blood_bam"

"$tumor_bam"). Subsequently, SVs were identified using Sniffles2 (v2.0.6) based on the

merged BAM files, which reported all supporting reads of each SV in the variant call

format (VCF) file. Reads with a mapping quality below 10 were excluded from SV

detection. To improve the accuracy of SV detection, a tandem repeat file

(https://github.com/fritzsedlazeck/Sniffles/blob/master/annotations/human_GRCh38_no_

alt_analysis_set.trf.bed) was provided using the --tandem-repeats parameter, enabling

Sniffles to better resolve SVs involving repetitive sequences. SVs supported only by reads

shorter than 1,000 base pairs (bp) were also discarded. Large-scale insertions were

considered up to a maximum length of 10,000 bp. All other parameters were set to default

values. Then, an initial set of candidate somatic SVs was filtered using a custom Python

script with the following parameters: tumor reads supporting SV ≥ 3, blood reads

supporting SV = 0, and SV length ≥ 50 bp.

To improve accuracy, an additional screening of candidate somatic SVs was performed.

Specifically, SV positions were restricted to a standard deviation of less than 150 bp. For

insertions (INS) and duplications (DUP), comparisons were made between the

alignment-supported lengths and those reported by Sniffles, with outliers excluded based

on:

A.mean(Lengtha)-2*STRI(Lengtha)<=Lengths <=mean(Lengtha)+2*STRI(Lengtha)

B.Q1-1.5IQR<=Length:<=Q3+1.5IQR.

Additionally, SVs located in repeat regions were assessed for significant length

differences between tumor and blood alignments, retaining only those exhibiting clear

distinctions.

To further ensure high-confidence somatic SV identification, an error-correction process

was conducted. Candidate regions were first identified as those within a ±50 bp range

centered around insertion sites reported by Sniffles2. Alignments in these regions were

extracted, and the reference genomes were polished using Racon (version 1.4.16, default

parameters) over six consecutive iterations, utilizing paired blood reads from each region.

Tumor and blood reads were then realigned to these polished reference genomes, and

SVs were detected using Sniffles2. SVs supported by more than three tumor reads, with a

length exceeding 50 bp, were retained as high-confidence somatic SVs.

**Somatic TEI Annotation**

Iris is a software developed to improve the accuracy of insertion calling. We used Iris
software (version 1.0.4, parameters: genome_buffer=1000 --keep_long_variants) to
correct the positions and sequences of somatic SVs identified as INSs and DUPs. To
improve the accuracy of Iris, we pre-processed the data by supplementing missing
sequences for variants that Sniffles did not report. Specifically, we extracted the
highest-quality reads from those supporting the Sniffles call that also spanned the
insertion site. The insertion sequence from these reads was then identified using pysam
and used as the input for Iris correction.

After error correction, we used RepeatMasker (version 4.1.2) to annotate the corrected
insertion sequences with the following parameters: -species human -engine RMBlast -q,
along with 50 bp of reference genome sequences upstream and downstream of the
insertion site. If RepeatMasker failed to identify transposon regions, we performed
re-annotation using Tandem Repeat Finder (TRF, version 4.10.0, parameters: 2 5 7 80 10
50 2000 -l 10 -ngs -h) and Sdust (version 0.1). Insertion sequences marked as "Unmask"
by RepeatMasker were also re-annotated with TRF and Sdust if the annotated length
exceeded 5 bp.

## Classification of Somatic TEI

If the 50 bp flanking sequences and the insertion sequence are not annotated as the
same type of repeat and the insertion carries a transposable element (TE) sequence, the
somatic INS is classified as a somatic TEI. Insertions containing TE sequences are further
classified based on their structure and recombination mechanism. A single, complete
insertion in the same orientation is classified as Solo, while insertions with multiple or
fragmented TEs, or those with different orientations, are classified as Complex.

## Classification of SINE and LINE Truncations in Somatic TEIs

Each SINE and LINE insertion was annotated using RepeatMasker to compare the
insertion sequences against template sequences in the RepeatMasker database. Using
the positive strand as an example, we classified the insertion sequences based on their
alignment characteristics. If the somatic insertion aligned fully with the template, with no
gaps at either end, it was classified as Full Length. If the sequence matched entirely to the
3' end (i.e., left = 0), it was categorized as a 5' Truncation. Conversely, if it matched fully to
the 5' end (i.e., begin = 1), it was classified as a 3' Truncation. If the sequence partially
aligned with both the 3' and 5' ends but was not fully aligned at either end, it was classified
as a TE Insertion. Additionally, the alignment coordinates of each SINE and LINE insertion
against the template sequence were recorded for subsequent statistical analysis.

**Detection of SINE and LINE PolyA tails in Somatic TEI**

For an insertion sequence on the negative strand, if there were 10 or more consecutive 'A' nucleotides within the first 30 bp of the sequence, it was classified as containing a polyA tail. Conversely, for an insertion sequence on the positive strand, the presence of 10 or more consecutive 'T' nucleotides within the last 30 bp was used as the criterion for classifying the sequence as containing a polyA tail.

**Detection of Somatic TEI Homologous Sequences**

To identify homologous sequences between the ends of an insertion and the flanking regions of the reference genome, we performed BLAST analysis (version 2.14.1, parameters: -outfmt 6 -word_size 4), This analysis compared each insertion sequence with reference genomic sequences of varying lengths on both sides of the insertion site.

For gradient length determination, the maximum reference length was set equal to the length of the insertion, and the minimum length was defined by the allowable breakpoint deviation. For sequences shorter than 100 bp, we used length increments of 5 bp, while for sequences longer than 100 bp, increments were set to one-twentieth of the insertion length to optimize computational efficiency. Homologous sequences were defined as the longest segments within the breakpoint deviation that closely aligned with the ends of the insertion sequence.

Based on the relative position and length of these homologous sequences, insertions were classified into four categories:

1. Non-allelic homologous recombination (NAHR): Homologous sequences greater than 100 bp in length.

2. Tandem duplications (TSD): Homologous sequences less than 100 bp in length with a distal relative position.

3. Recombination (Else): Homologous sequences that did not meet the criteria for NAHR or TSD.

4. No homologous sequence detected: Insertions were classified as "N" when no homologous sequence was identified.

**Somatic TEIs Motif Clustering**

To examine the distribution patterns of somatic TEIs across the genome, we divided the genome into 1 Mb fixed-size windows to create a pan-cancer atlas of somatic TEI distribution. The genome was segmented into consecutive 1 Mb windows, and the presence of somatic TEIs within each window was recorded for each tumor sample. This

resulted in a binary matrix representing the presence or absence of somatic TEIs across
the genome for each sample.

Windows containing TEIs in only a single tumor sample were excluded from further
analysis. For the remaining data, we calculated the Jaccard similarity coefficient for each
pair of tumor samples to assess their genomic TEI similarity. Hierarchical clustering was
then performed using the pheatmap package in R (ward.D method) to group tumor
samples based on their somatic TEI distribution.

**Analysis of the Association Between L1 Insertion Rate and Genomic**
**Features**
The L1 insertion rate is determined by calculating the total number of somatic L1
insertions (10,140) identified in 317 out of 322 samples within each 1Mb window. To
investigate the association between L1 insertion rates and multiple predictor variables, we
employed a statistical framework based on negative binomial regression[1]. Specifically,
we categorized the genome into four bins (bin0 - bin3) based on the degree of match
between the classic L1 endonuclease motif (defined as TTTT|R, where R is A or G or
Y|AAAA, where Y is C or T) and the DNA sequence. Bin0 contains DNA motifs that have
four or more mismatches (out of five possible), which are highly dissimilar. Bin1, bin2, and
bin3 include genomic segments of the hg38 genome assembly with three, two, and at
most one mismatch, respectively. The most tightly matching DNA strands at each site are
considered.

1006

Regional analysis of genomic features utilized histone mark data (H3K9me3, H3K4me3,
H3K36me3, H3K27ac ChIP-Seq data) and DNase hypersensitivity (DHS) data from the
Roadmap Epigenomics Consortium. These data were divided into four bins based on the
average enrichment scores across eight cell types (E017, E114, E117, E118, E119, E122,
E125, E127). Bin0 includes regions with enrichment scores below baseline (Roadmap
enrichment scores relative to input <1), while bins 1 to 3 correspond to regions with
higher-than-average enrichment scores across the genome. Specifically, the DHS bins
1-3 span 132.1-136.1 Mb, the H3K9me3 bins 1-3 span 192.8-198.7 Mb, the H3K36me3
bins 1-3 span 142.3-146.5 Mb, the H3K4me3 bins 1-3 span 44.3-45.9 Mb, and the
H3K27ac bins 1-3 span 77.8-80.1 Mb.

1017

1018     Replication timing (RT) data from ENCODE were processed by averaging smoothed
1019     waveform signals from eight cell types (IMR-90, HeLa S3, HUVEC, BJ, HEP G2, NHEK,
1020     SK-N-SH, and MCF-7). The data were divided into four equal-sized bins (quartiles), where
1021     bin0 represents the latest replication and bin3 represents the earliest replication.

1022

RNA-Seq data, also from the Roadmap, were processed similarly. By averaging data from

eight cell types (E071, E096, E114, E117, E118, E119, E122, E127), the data were

divided into four genomic bins: bin0 contains non-expressed genes (logRPKM=0) and

non-expressed intergenic DNA (totaling 1173.9 Mb), while bin1 (maximum

logRPKM=0.02), bin2 (maximum logRPKM=0.15), and bin3 (logRPKM > 0.15) span 353.7

Mb, 302.2 Mb, and 382.9 Mb, respectively.

1029

1030 In all figures (Supplementary Fig. 3b), the enrichment scores compare differences in

1031 specific features (replication timing, histone marks, gene expression, and L1 motifs)

1032 between bins 1-3 and bin0. The logarithmic enrichment for bin0 is defined as 0, and thus it

1033 is not displayed in the enrichment plots. Regional analysis is limited to genomic regions

1034 with perfect alignment scores, based on the hg38 reference genome.

1035

**Fresh tissue preparation and dissociation into single cells.**

All the freshly resected surgical specimens were immediately washed with

phosphate-buffered saline (PBS) and processed to generate single cell suspensions.

Tissue digestion was performed in 15 ml tube containing 10 ml pre-warmed RPMI 1640

(ThermoFisher Scientific), 2 mg/ml dispase (Roche), 1 mg/ml type IV collagenase (Sigma)

and 10 U/µl DNase I (Roche) for 60 minutes at 37°C. The reaction was deactivated by

10% FBS. Cell suspensions were filtered using a 70 µm filter and then centrifuged at 500

rpm for 6 min at 4°C to pellet dead cells and red blood cells. The cells were washed twice

and suspended in PBS with 0.5% bovine serum albumin (BSA, Sigma), and, used CD45

MicroBeads (Miltenyibiotec ,130-045-801) to distinguish CD45+ cells from the rest of the

cells.

1047

**Validation of the Somatic TEI&TRI in Tissues and Cell Lines using PCR**

**and Sanger sequencing**

DNA extraction and PCR: DNA was extracted from tumor tissues, blood, LUSC cell lines

(H226, H520, HCC95), LUAD cell lines (PC-9, A549, H1299 and H1975), IC cell lines

(LOVO, SW620, RKO, HCT15, HCT116,T84), Breast cancer cell line (HCC1395), OSCC

cell lines (TU212, TU686, SNU46, SNU899), CCA cell lines (H8, C33A, CASK, SiHA),

PAAD cell lines (HPAC, PANC-1,BXPC-1,ASPC-1),HCC cell lines (sun387, Huh7, hcclm9,

snu499,sk-hep1), the H1581 cell line, Normal cell line BEAS-2B, HCC1395BL using the

DNeasy Blood & Tissue Kit (QIAGEN) according to the manufacturer's instructions. The

forward and reverse primers were designed according to the 100 bp upstream and

downstream sequences of the insertion position and synthesized by Sangon Biotech

(Table TEI,Table TRI). DNA was extracted from clinical samples or cell lines and

subjected to PCR using primers. Each 50-µL reaction consisted of 2× PremSTAR Max Premix 25 µL, 0.8 µL 10 µM anchor primer, 0.8 µL 10 µM test primer, and 50-150 ng DNA. PCR was performed as follows: an initial step of 2 minutes at 98 °C, 30 cycles of 10 seconds at 98 °C, 15 seconds at 60 °C and 60-300 seconds at 72 °C (size of PCR products between 200 bp and 6 kb). Finally, the samples were incubated at 72 °C for 5 minutes. The PCR products were sent to Sangon for Sanger sequencing.

## Cell Culture

Human cell lines provided by the American Type Culture Collection (Manassas, VA, USA) were used in the present study. The cells were incubated in Dulbecco's modified Eagle's medium (DMEM) or RPMI 1640 medium (Gibco; Thermo Fisher Scientific, Inc.). The medium was supplemented with 10% fetal bovine serum (FBS) (Gibco; Thermo Fisher Scientific, Inc.), and the cells were all maintained at 37 °C in 5% $CO_2$. Upon reaching 70-80% confluence, the cells were washed with PBS and detached with 0.25% trypsin/0.2% ethylenediaminetetraacetic acid (EDTA). Cell morphology was viewed under a light microscope, and the cells were suspended to a concentration of 1 x 106 cells/ml.

## Immunofluorescence and FISH

Briefly, 10 µm thick frozen sections were treated with 0.2% Triton X-100 in PBS for 15 minutes and blocked with 10% serum in PBST for 40 minutes at room temperature. They were then incubated with primary antibodies, Anti CK7/serpinB3/NES/GFAP (1:500, Huabio, ET1609-62, ET7111-21, ET1610-96, ET1601-23) overnight at 4°C and washed with PBS for 15 min, Alexa-conjugated secondary antibodies (1:1000, iFluor 488 Goat anti-rabbit IgG antibody, Huabio, HA720120F; iFluor 594 Conjugated Goat anti-rabbit IgG polyclonal Antibody, Huabio, HA1122) were incubated for 1 hour at room temperature. For FISH, probes specific to the target sequences (Table S20) were hybridized to the cells overnight at 37°C. After washing, the cells were counterstained with DAPI and analyzed under a fluorescence microscope.

## Cell co-culture

Splenic cells were collected from 6-8 weeks old C57/BL6 mice and resuspended in RoboSep™ Buffer (STEMCELL Technologies, 20104). Total splenocytes were enriched for CD4+ T cells with EasySep™ Mouse CD4+ T Cell Isolation Kit (STEMCELL Technologies, 19852). The selected CD4+ T cells and tumor cells were directly co-cultured with RPMI 1640 medium (containing 10% FBS) at a ratio of 10:1 for 24 hours, and suspended T cells were collected at 24 hours for subsequent flow experiments.

## Flow cytometry

For cell surface staining, cells were stained with the corresponding antibodies, APC
anti-mouse CD45 (1:100, Biolegend, 103112); FITC anti-mouse CD4 (1:100, Biolegend,
100405); PE anti-mouse CD154 (CD40L) (1:50, Biolegend, 157003); eBioscience™
Fixable Viability Dye eFluor™ 506 (1:100, Thermo Fisher Scientific, 65-0866-18), in the
presence of Fc receptor blocker on ice for 30 min, and then fixed with 4%
paraformaldehyde for 10 minutes, washed with 1x PBS solution, and resuspended after
centrifugation. Samples were assessed by a spectral flow cytometer (Cytoflex, BECKMAN
Biosciences,) equipped with SpectroFlo software (Cytek Biosciences). Acquired flow
cytometry results were analyzed by FlowJo software.

## Data availability

1109 All sequencing data are available at the Genome Sequence Archive for Human:

1110 https://ngdc.cncb.ac.cn/gsa-human/s/QfWwWORT,

1111 https://ngdc.cncb.ac.cn/gsa-human/s/d97GuJ2L and

1112 https://ngdc.cncb.ac.cn/gsa-human/s/MnWe8KbN.

## Code availability

1115 All code used is open source and available at github:

1116 https://github.com/LinXialab/TGS_somatic_TEI/ and

1117 https://github.com/Goatofmountain/TDScope.

# Reference

1. Swanton, C., Bernard, E., Abbosh, C., Andre, F., Auwerx, J., Balmain, A., Bar-Sagi, D., Bernards, R., Bullman, S., DeGregori, J., et al. (2024). Embracing cancer complexity: Hallmarks of systemic disease. Cell *187*, 1589-1616. 10.1016/j.cell.2024.02.009.

2. Hanahan, D. (2022). Hallmarks of Cancer: New Dimensions. Cancer Discov *12*, 31-46. 10.1158/2159-8290.CD-21-1059.

3. International Cancer Genome, C., Hudson, T.J., Anderson, W., Artez, A., Barker, A.D., Bell, C., Bernabe, R.R., Bhan, M.K., Calvo, F., Eerola, I., et al. (2010). International network of cancer genome projects. Nature *464*, 993-998. 10.1038/nature08987.

4. Cancer Genome Atlas Research, N., Weinstein, J.N., Collisson, E.A., Mills, G.B., Shaw, K.R., Ozenberger, B.A., Ellrott, K., Shmulevich, I., Sander, C., and Stuart, J.M. (2013). The Cancer Genome Atlas Pan-Cancer analysis project. Nat Genet *45*, 1113-1120. 10.1038/ng.2764.

5. Kandoth, C., McLellan, M.D., Vandin, F., Ye, K., Niu, B., Lu, C., Xie, M., Zhang, Q., McMichael, J.F., Wyczalkowski, M.A., et al. (2013). Mutational landscape and significance across 12 major cancer types. Nature *502*, 333-339. 10.1038/nature12634.

6. Lawrence, M.S., Stojanov, P., Mermel, C.H., Robinson, J.T., Garraway, L.A., Golub, T.R.,

1135        Meyerson, M., Gabriel, S.B., Lander, E.S., and Getz, G. (2014). Discovery and saturation
1136        analysis of cancer genes across 21 tumour types. Nature *505*, 495-501. 10.1038/nature12912.
1137

7.    Hutter, C., and Zenklusen, J.C. (2018). The Cancer Genome Atlas: Creating Lasting Value
beyond Its Data. Cell *173*, 283-285. 10.1016/j.cell.2018.03.042.

8.    Bailey, M.H., Tokheim, C., Porta-Pardo, E., Sengupta, S., Bertrand, D., Weerasinghe, A.,
Colaprico, A., Wendl, M.C., Kim, J., Reardon, B., et al. (2018). Comprehensive Characterization
of Cancer Driver Genes and Mutations. Cell *173*, 371-385 e318. 10.1016/j.cell.2018.02.060.

9.    Hornshoj, H., Nielsen, M.M., Sinnott-Armstrong, N.A., Switnicki, M.P., Juul, M., Madsen, T.,
Sallari, R., Kellis, M., Orntoft, T., Hobolth, A., and Pedersen, J.S. (2018). Pan-cancer screen for
mutations in non-coding elements with conservation and cancer specificity reveals
correlations with expression and survival. NPJ Genom Med *3*, 1. 10.1038/s41525-017-0040-5.

10.   Rheinbay, E., Nielsen, M.M., Abascal, F., Wala, J.A., Shapira, O., Tiao, G., Hornshoj, H., Hess,
J.M., Juul, R.I., Lin, Z., et al. (2020). Analyses of non-coding somatic drivers in 2,658 cancer
whole genomes. Nature *578*, 102-111. 10.1038/s41586-020-1965-x.

11.   Hakobyan, A., Meyenberg, M., Vardazaryan, N., Hancock, J., Vulliard, L., Loizou, J.I., and
Menche, J. (2024). Pan-cancer analysis of the interplay between mutational signatures and
cellular signaling. iScience *27*, 109873. 10.1016/j.isci.2024.109873.

12.   Hu, Z., Li, Z., Ma, Z., and Curtis, C. (2020). Multi-cancer analysis of clonality and the timing of
systemic spread in paired primary tumors and metastases. Nat Genet *52*, 701-708.
10.1038/s41588-020-0628-z.

13.   Fontana, D., Crespiatico, I., Crippa, V., Malighetti, F., Villa, M., Angaroni, F., De Sano, L., Aroldi,
A., Antoniotti, M., Caravagna, G., et al. (2023). Evolutionary signatures of human cancers
revealed via genomic analysis of over 35,000 patients. Nat Commun *14*, 5982.
10.1038/s41467-023-41670-3.

14.   Shah, Y., Verma, A., Marderstein, A.R., White, J., Bhinder, B., Garcia Medina, J.S., and
Elemento, O. (2021). Pan-cancer analysis reveals molecular patterns associated with age. Cell
Rep *37*, 110100. 10.1016/j.celrep.2021.110100.

15.   Chen, H., Li, C., Peng, X., Zhou, Z., Weinstein, J.N., Cancer Genome Atlas Research, N., and
Liang, H. (2018). A Pan-Cancer Analysis of Enhancer Expression in Nearly 9000 Patient
Samples. Cell *173*, 386-399 e312. 10.1016/j.cell.2018.03.027.

16.   Hoadley, K.A., Yau, C., Hinoue, T., Wolf, D.M., Lazar, A.J., Drill, E., Shen, R., Taylor, A.M.,
Cherniack, A.D., Thorsson, V., et al. (2018). Cell-of-Origin Patterns Dominate the Molecular
Classification of 10,000 Tumors from 33 Types of Cancer. Cell *173*, 291-304 e296.
10.1016/j.cell.2018.03.022.

17.   Drews, R.M., Hernando, B., Tarabichi, M., Haase, K., Lesluyes, T., Smith, P.S., Morrill Gavarro,
L., Couturier, D.L., Liu, L., Schneider, M., et al. (2022). A pan-cancer compendium of
chromosomal instability. Nature *606*, 976-983. 10.1038/s41586-022-04789-9.

18.   Sudmant, P.H., Rausch, T., Gardner, E.J., Handsaker, R.E., Abyzov, A., Huddleston, J., Zhang, Y.,
Ye, K., Jun, G., Fritz, M.H., et al. (2015). An integrated map of structural variation in 2,504
human genomes. Nature *526*, 75-81. 10.1038/nature15394.

19.   Zhang, Y., Yang, L., Kucherlapati, M., Chen, F., Hadjipanayis, A., Pantazi, A., Bristow, C.A., Lee,
E.A., Mahadeshwar, H.S., Tang, J., et al. (2018). A Pan-Cancer Compendium of Genes
Deregulated by Somatic Genomic Rearrangement across More Than 1,400 Cases. Cell Rep *24*,

515-527. 10.1016/j.celrep.2018.06.025.

20.    Bielski, C.M., Zehir, A., Penson, A.V., Donoghue, M.T.A., Chatila, W., Armenia, J., Chang, M.T.,
1181           Schram, A.M., Jonsson, P., Bandlamudi, C., et al. (2018). Genome doubling shapes the
1182           evolution and prognosis of advanced cancers. Nat Genet *50*, 1189-1195.
1183           10.1038/s41588-018-0165-1.

1184    21.    Consortium, I.T.P.-C.A.o.W.G. (2020). Pan-cancer analysis of whole genomes. Nature *578*,
1185           82-93. 10.1038/s41586-020-1969-6.

1186    22.    Hadi, K., Yao, X., Behr, J.M., Deshpande, A., Xanthopoulakis, C., Tian, H., Kudman, S., Rosiene,
1187           J., Darmofal, M., DeRose, J., et al. (2020). Distinct Classes of Complex Structural Variation
1188           Uncovered across Thousands of Cancer Genome Graphs. Cell *183*, 197-210 e132.
1189           10.1016/j.cell.2020.08.006.

1190    23.    Ho, S.S., Urban, A.E., and Mills, R.E. (2020). Structural variation in the sequencing era. Nat
1191           Rev Genet *21*, 171-189. 10.1038/s41576-019-0180-9.

1192    24.    Xu, L., Wang, X., Lu, X., Liang, F., Liu, Z., Zhang, H., Li, X., Tian, S., Wang, L., and Wang, Z.
1193           (2023). Long-read sequencing identifies novel structural variations in colorectal cancer. PLoS
1194           Genet *19*, e1010514. 10.1371/journal.pgen.1010514.

1195    25.    Dan Xie, L.X., Zhoufeng Wang,Tianfu Zeng,Xuenan Pi,Huan Wang,Guonian Zhu,Xinyue
1196           Wu,Yangqian Li,Yan Deng,Yawen Qi,Xuyan liu,Fengmei Zhang,Weimin Li (2023). Landscape of
1197           Large-Scale Somatic Genomic Insertions in Non-Small Cell Lung Carcinoma Revealed by
1198           Nanopore Sequencing. bioRxiv. https://doi.org/10.1101/2023.10.07.561149.

1199    26.    Reis, A.L.M., Rapadas, M., Hammond, J.M., Gamaarachchi, H., Stevanovski, I., Ayuputeri
1200           Kumaheri, M., Chintalaphani, S.R., Dissanayake, D.S.B., Siggs, O.M., Hewitt, A.W., et al. (2023).
1201           The landscape of genomic structural variation in Indigenous Australians. Nature *624*, 602-610.
1202           10.1038/s41586-023-06842-7.

1203    27.    Cortes-Ciriano, I., Lee, S., Park, W.Y., Kim, T.M., and Park, P.J. (2017). A molecular portrait of
1204           microsatellite instability across multiple cancers. Nat Commun *8*, 15180.
1205           10.1038/ncomms15180.

1206    28.    Hause, R.J., Pritchard, C.C., Shendure, J., and Salipante, S.J. (2016). Classification and
1207           characterization of microsatellite instability across 18 cancer types. Nat Med *22*, 1342-1350.
1208           10.1038/nm.4191.

1209    29.    Shah, N.M., Jang, H.J., Liang, Y., Maeng, J.H., Tzeng, S.C., Wu, A., Basri, N.L., Qu, X., Fan, C., Li,
1210           A., et al. (2023). Pan-cancer analysis identifies tumor-specific antigens derived from
1211           transposable elements. Nat Genet *55*, 631-639. 10.1038/s41588-023-01349-3.

1212    30.    Mehdipour, P., Marhon, S.A., Ettayebi, I., Chakravarthy, A., Hosseini, A., Wang, Y., de Castro,
1213           F.A., Loo Yau, H., Ishak, C., Abelson, S., et al. (2020). Epigenetic therapy induces transcription
1214           of inverted SINEs and ADAR1 dependency. Nature *588*, 169-173.
1215           10.1038/s41586-020-2844-1.

1216    31.    Jang, H.S., Shah, N.M., Du, A.Y., Dailey, Z.Z., Pehrsson, E.C., Godoy, P.M., Zhang, D., Li, D., Xing,
1217           X., Kim, S., et al. (2019). Transposable elements drive widespread expression of oncogenes in
1218           human cancers. Nat Genet *51*, 611-617. 10.1038/s41588-019-0373-3.

1219    32.    Sun, Z., Zhang, R., Zhang, X., Sun, Y., Liu, P., Francoeur, N., Han, L., Lam, W.Y., Yi, Z., Sebra, R.,
1220           et al. (2022). LINE-1 promotes tumorigenicity and exacerbates tumor progression via
1221           stimulating metabolism reprogramming in non-small cell lung cancer. Mol Cancer *21*, 147.
1222           10.1186/s12943-022-01618-5.

33.    Nam, C.H., Youk, J., Kim, J.Y., Lim, J., Park, J.W., Oh, S.A., Lee, H.J., Park, J.W., Won, H., Lee, Y.,
et al. (2023). Widespread somatic L1 retrotransposition in normal colorectal epithelium.
Nature *617*, 540-547. 10.1038/s41586-023-06046-z.

34.    Erwin, G.S., Gursoy, G., Al-Abri, R., Suriyaprakash, A., Dolzhenko, E., Zhu, K., Hoerner, C.R.,
White, S.M., Ramirez, L., Vadlakonda, A., et al. (2023). Recurrent repeat expansions in human
cancer genomes. Nature *613*, 96-102. 10.1038/s41586-022-05515-1.

35.    Grunewald, T.G., Bernard, V., Gilardi-Hebenstreit, P., Raynal, V., Surdez, D., Aynaud, M.M.,
Mirabeau, O., Cidre-Aranaz, F., Tirode, F., Zaidi, S., et al. (2015). Chimeric EWSR1-FLI1
regulates the Ewing sarcoma susceptibility gene EGR2 via a GGAA microsatellite. Nat Genet
*47*, 1073-1078. 10.1038/ng.3363.

36.    Beyter, D., Ingimundardottir, H., Oddsson, A., Eggertsson, H.P., Bjornsson, E., Jonsson, H.,
Atlason, B.A., Kristmundsdottir, S., Mehringer, S., Hardarson, M.T., et al. (2021). Long-read
sequencing of 3,622 Icelanders provides insight into the role of structural variants in human
diseases and other traits. Nat Genet *53*, 779-786. 10.1038/s41588-021-00865-4.

37.    Wu, Z., Jiang, Z., Li, T., Xie, C., Zhao, L., Yang, J., Ouyang, S., Liu, Y., Li, T., and Xie, Z. (2021).
Structural variants in the Chinese population and their impact on phenotypes, diseases and
population adaptation. Nat Commun *12*, 6501. 10.1038/s41467-021-26856-x.

38.    Gustafson, J.A., Gibson, S.B., Damaraju, N., Zalusky, M.P.G., Hoekzema, K., Twesigomwe, D.,
Yang, L., Snead, A.A., Richmond, P.A., De Coster, W., et al. (2024). High-coverage nanopore
sequencing of samples from the 1000 Genomes Project to build a comprehensive catalog of
human genetic variation. Genome Res *34*, 2061-2073. 10.1101/gr.279273.124.

39.    Ichikawa, K., Kawahara, R., Asano, T., and Morishita, S. (2023). A landscape of complex
tandem repeats within individual human genomes. Nat Commun *14*, 5530.
10.1038/s41467-023-41262-1.

40.    Dolzhenko, E., English, A., Dashnow, H., De Sena Brandine, G., Mokveld, T., Rowell, W.J.,
Karniski, C., Kronenberg, Z., Danzi, M.C., Cheung, W.A., et al. (2024). Characterization and
visualization of tandem repeats at genome scale. Nat Biotechnol *42*, 1606-1614.
10.1038/s41587-023-02057-3.

41.    Liu, L., Zhang, J., Wood, S., Newell, F., Leonard, C., Koufariotis, L.T., Nones, K., Dalley, A.J.,
Chittoory, H., Bashirzadeh, F., et al. (2024). Performance of somatic structural variant calling
in lung cancer using Oxford Nanopore sequencing technology. BMC Genomics *25*, 898.
10.1186/s12864-024-10792-3.

42.    Sakamoto, Y., Xu, L., Seki, M., Yokoyama, T.T., Kasahara, M., Kashima, Y., Ohashi, A., Shimada,
Y., Motoi, N., Tsuchihara, K., et al. (2020). Long-read sequencing for non-small-cell lung
cancer genomes. Genome Res *30*, 1243-1257. 10.1101/gr.261941.120.

43.    Fujimoto, A., Wong, J.H., Yoshii, Y., Akiyama, S., Tanaka, A., Yagi, H., Shigemizu, D., Nakagawa,
H., Mizokami, M., and Shimada, M. (2021). Whole-genome sequencing with long reads
reveals complex structure and origin of structural variation in human genetic variations and
somatic mutations in cancer. Genome Med *13*, 65. 10.1186/s13073-021-00883-1.

44.    Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics *34*,
3094-3100. 10.1093/bioinformatics/bty191.

45.    Smolka, M., Paulin, L.F., Grochowski, C.M., Horner, D.W., Mahmoud, M., Behera, S., Kalef-Ezra,
E., Gandhi, M., Hong, K., Pehlivan, D., et al. (2024). Detection of mosaic and population-level
structural variants with Sniffles2. Nat Biotechnol *42*, 1571-1580.

10.1038/s41587-023-02024-y.

46.  Chiu, R., Rajan-Babu, I.S., Friedman, J.M., and Birol, I. (2021). Straglr: discovering and
genotyping tandem repeat expansions using whole genome long-read sequences. Genome
Biol *22*, 224. 10.1186/s13059-021-02447-3.

47.  Kailing Tu, Q.Z., Yang Li,Yucong Li,Lanfang Yuan,Jie Tang,Lin Xia,Jing Wang,Wei Huang,Dan Xie
(2025). Accurate Somatic SV detection via sequence graph model-based local pan-genome
optimization. bioRxiv. https://doi.org/10.1101/2025.02.11.636543.

48.  Kirsche, M., Prabhu, G., Sherman, R., Ni, B., Battle, A., Aganezov, S., and Schatz, M.C. (2023).
Jasmine and Iris: population-scale structural variant comparison and analysis. Nat Methods
*20*, 408-417. 10.1038/s41592-022-01753-3.

49.  Vaser, R., Sovic, I., Nagarajan, N., and Sikic, M. (2017). Fast and accurate de novo genome
assembly from long uncorrected reads. Genome Res *27*, 737-746. 10.1101/gr.214270.116.

50.  Tarailo-Graovac, M., and Chen, N. (2009). Using RepeatMasker to identify repetitive elements
in genomic sequences. Curr Protoc Bioinformatics *Chapter 4*, Unit 4 10.
10.1002/0471250953.bi0410s25.

51.  Benson, G. (1999). Tandem repeats finder: a program to analyze DNA sequences. Nucleic
Acids Res *27*, 573-580. 10.1093/nar/27.2.573.

52.  Morgulis, A., Gertz, E.M., Schaffer, A.A., and Agarwala, R. (2006). A fast and symmetric DUST
implementation to mask low-complexity DNA sequences. J Comput Biol *13*, 1028-1040.
10.1089/cmb.2006.13.1028.

53.  McLean, C.Y., Bristor, D., Hiller, M., Clarke, S.L., Schaar, B.T., Lowe, C.B., Wenger, A.M., and
Bejerano, G. (2010). GREAT improves functional interpretation of cis-regulatory regions. Nat
Biotechnol *28*, 495-501. 10.1038/nbt.1630.

54.  Gao, Y., Ferguson, D.O., Xie, W., Manis, J.P., Sekiguchi, J., Frank, K.M., Chaudhuri, J., Horner, J.,
DePinho, R.A., and Alt, F.W. (2000). Interplay of p53 and DNA-repair protein XRCC4 in
tumorigenesis, genomic stability and development. Nature *404*, 897-900. 10.1038/35009138.

55.  Bhattacharya, S., Srinivasan, K., Abdisalaam, S., Su, F., Raj, P., Dozmorov, I., Mishra, R.,
Wakeland, E.K., Ghose, S., Mukherjee, S., and Asaithamby, A. (2017). RAD51 interconnects
between DNA replication, DNA repair and immunity. Nucleic Acids Res *45*, 4590-4605.
10.1093/nar/gkx126.

56.  Coin, F., Auriol, J., Tapias, A., Clivio, P., Vermeulen, W., and Egly, J.M. (2004). Phosphorylation
of XPB helicase regulates TFIIH nucleotide excision repair activity. EMBO J *23*, 4835-4846.
10.1038/sj.emboj.7600480.

57.  Audano, P.A., Sulovari, A., Graves-Lindsay, T.A., Cantsilieris, S., Sorensen, M., Welch, A.E.,
Dougherty, M.L., Nelson, B.J., Shah, A., Dutcher, S.K., et al. (2019). Characterizing the Major
Structural Variant Alleles of the Human Genome. Cell *176*, 663-675 e619.
10.1016/j.cell.2018.12.019.

58.  Niu, Y., Teng, X., Zhou, H., Shi, Y., Li, Y., Tang, Y., Zhang, P., Luo, H., Kang, Q., Xu, T., and He, S.
(2022). Characterizing mobile element insertions in 5675 genomes. Nucleic Acids Res *50*,
2493-2508. 10.1093/nar/gkac128.

59.  Pascarella, G., Hon, C.C., Hashimoto, K., Busch, A., Luginbuhl, J., Parr, C., Hin Yip, W., Abe, K.,
Kratz, A., Bonetti, A., et al. (2022). Recombination of repeat elements generates somatic
complexity in human genomes. Cell *185*, 3025-3040 e3026. 10.1016/j.cell.2022.06.032.

60.  Liu, T., Wu, L., Liu, J., Chen, H., Zhu, B., Qiao, D., Zhu, Y., Liu, T., Chen, Q., and Hu, A. (2023).

Comprehensive characterization of MUC16 mutations in lung adenocarcinoma for
immunotherapies and prognosis: An observational study. Medicine (Baltimore) *102*, e35481.
10.1097/MD.0000000000035481.

61.  Shah, A., Chaudhary, S., Lakshmanan, I., Aithal, A., Kisling, S.G., Sorrell, C., Marimuthu, S.,
Gautam, S.K., Rauth, S., Kshirsagar, P., et al. (2023). Chimeric antibody targeting unique
epitope on onco-mucin16 reduces tumor burden in pancreatic and lung malignancies. NPJ
Precis Oncol *7*, 74. 10.1038/s41698-023-00423-7.

62.  Marimuthu, S., Lakshmanan, I., Muniyan, S., Gautam, S.K., Nimmakayala, R.K., Rauth, S., Atri,
P., Shah, A., Bhyravbhatla, N., Mallya, K., et al. (2022). MUC16 Promotes Liver Metastasis of
Pancreatic Ductal Adenocarcinoma by Upregulating NRP2-Associated Cell Adhesion. Mol
Cancer Res *20*, 1208-1221. 10.1158/1541-7786.MCR-21-0888.

63.  Skates, S.J., Greene, M.H., Buys, S.S., Mai, P.L., Brown, P., Piedmonte, M., Rodriguez, G.,
Schorge, J.O., Sherman, M., Daly, M.B., et al. (2017). Early Detection of Ovarian Cancer using
the Risk of Ovarian Cancer Algorithm with Frequent CA125 Testing in Women at Increased
Familial Risk - Combined Results from Two Screening Trials. Clin Cancer Res *23*, 3628-3637.
10.1158/1078-0432.CCR-15-2750.

64.  Caputo, M., Xia, Y., Anand, S.K., Cansby, E., Andersson, E., Marschall, H.U., Konigsrainer, A.,
Peter, A., and Mahlapuu, M. (2023). STE20-type kinases MST3 and MST4 promote the
progression of hepatocellular carcinoma: Evidence from human cell culture and expression
profiling of liver biopsies. FASEB J *37*, e23105. 10.1096/fj.202300397RR.

65.  Huang, N., Lin, W., Shi, X., and Tao, T. (2018). STK24 expression is modulated by DNA copy
number/methylation in lung adenocarcinoma and predicts poor survival. Future Oncol *14*,
2253-2263. 10.2217/fon-2018-0126.

66.  Li, L., Chen, S., Tang, Y., Wu, J., He, Y., and Qiu, L. (2022). Oncogene or tumor suppressor gene:
An integrated pan-cancer analysis of NBPF1. Front Endocrinol (Lausanne) *13*, 950326.
10.3389/fendo.2022.950326.

67.  Zhang, J., Zhao, Y., Wang, J., Sneh, T., Yu, Q., Zhou, X., and Gong, C. (2020). NBPF1
independently determine the risk stratification and prognosis of patients with neuroblastoma.
Genomics *112*, 3951-3957. 10.1016/j.ygeno.2020.06.038.

68.  Li, J., Xian, L., Zhu, Z., Wang, Y., Zhang, W., Zheng, R., Xue, W., and Li, J. (2023). Role of CELF2
in ferroptosis: Potential targets for cancer therapy (Review). Int J Mol Med *52*.
10.3892/ijmm.2023.5291.

69.  Jen, J., and Wang, Y.C. (2016). Zinc finger proteins in cancer progression. J Biomed Sci *23*, 53.
10.1186/s12929-016-0269-9.

70.  Frankson, R., Yu, Z.H., Bai, Y., Li, Q., Zhang, R.Y., and Zhang, Z.Y. (2017). Therapeutic Targeting
of Oncogenic Tyrosine Phosphatases. Cancer Res *77*, 5701-5705.
10.1158/0008-5472.CAN-17-1510.

71.  Stanford, S.M., and Bottini, N. (2023). Targeting protein phosphatases in cancer
immunotherapy and autoimmune disorders. Nat Rev Drug Discov *22*, 273-294.
10.1038/s41573-022-00618-w.

72.  Fang, L., Liu, Q., Monteys, A.M., Gonzalez-Alegre, P., Davidson, B.L., and Wang, K. (2022).
DeepRepeat: direct quantification of short tandem repeats on signal data from nanopore
sequencing. Genome Biol *23*, 108. 10.1186/s13059-022-02670-6.

73.  Rodriguez-Martin, B., Alvarez, E.G., Baez-Ortega, A., Zamora, J., Supek, F., Demeulemeester, J.,

Santamarina, M., Ju, Y.S., Temes, J., Garcia-Souto, D., et al. (2020). Pan-cancer analysis of
whole genomes identifies driver rearrangements promoted by LINE-1 retrotransposition. Nat
Genet *52*, 306-319. 10.1038/s41588-019-0562-0.

74.   Helman, E., Lawrence, M.S., Stewart, C., Sougnez, C., Getz, G., and Meyerson, M. (2014).
Somatic retrotransposition in human cancer revealed by whole-genome and exome
sequencing. Genome Res *24*, 1053-1063. 10.1101/gr.163659.113.

75.   Kidd, J.M., Graves, T., Newman, T.L., Fulton, R., Hayden, H.S., Malig, M., Kallicki, J., Kaul, R.,
Wilson, R.K., and Eichler, E.E. (2010). A human genome structural variation sequencing
resource reveals insights into mutational mechanisms. Cell *143*, 837-847.
10.1016/j.cell.2010.10.027.

76.   Horton, C.A., Alexandari, A.M., Hayes, M.G.B., Marklund, E., Schaepe, J.M., Aditham, A.K.,
Shah, N., Suzuki, P.H., Shrikumar, A., Afek, A., et al. (2023). Short tandem repeats bind
transcription factors to tune eukaryotic gene expression. Science *381*, eadd1250.
10.1126/science.add1250.

77.   Depienne, C., and Mandel, J.L. (2021). 30 years of repeat expansion disorders: What have we
learned and what are the remaining challenges? Am J Hum Genet *108*, 764-785.
10.1016/j.ajhg.2021.03.011.

78.   Dubois, F., Sidiropoulos, N., Weischenfeldt, J., and Beroukhim, R. (2022). Structural variations
in cancer and the 3D genome. Nat Rev Cancer *22*, 533-546. 10.1038/s41568-022-00488-9.

79.   Simpson, J.T., Workman, R.E., Zuzarte, P.C., David, M., Dursi, L.J., and Timp, W. (2017).
Detecting DNA cytosine methylation using nanopore sequencing. Nat Methods *14*, 407-410.
10.1038/nmeth.4184.

80.   Guo, C.T., Jardin, B.D., Lin, J.S., Ambroise, R.L., Wang, Z., Yang, L.Z., Mazumdar, N., Lu, F.J., Ma,
Q., Cao, Y.P., et al. (2024). In vivo proximity proteomics uncovers palmdelphin (PALMD) as a
Z-disc-associated mitigator of isoproterenol-induced cardiac injury. Acta Pharmacol Sin *45*,
2540-2552. 10.1038/s41401-024-01348-y.

81.   Sainz-Jaspeado, M., Smith, R.O., Plunde, O., Pawelzik, S.C., Jin, Y., Nordling, S., Ding, Y.,
Aspenstrom, P., Hedlund, M., Bastianello, G., et al. (2021). Palmdelphin Regulates Nuclear
Resilience to Mechanical Stress in the Endothelium. Circulation *144*, 1629-1645.
10.1161/CIRCULATIONAHA.121.054182.

82.   Yuan, J., Chen, L., Xiao, J., Qi, X.K., Zhang, J., Li, X., Wang, Z., Lian, Y.F., Xiang, T., Zhang, Y., et al.
(2019). SHROOM2 inhibits tumor metastasis through RhoA-ROCK pathway-dependent and
-independent mechanisms in nasopharyngeal carcinoma. Cell Death Dis *10*, 58.
10.1038/s41419-019-1325-7.

83.   Farber, M.J., Rizaldy, R., and Hildebrand, J.D. (2011). Shroom2 regulates contractility to
control endothelial morphogenesis. Mol Biol Cell *22*, 795-805. 10.1091/mbc.E10-06-0505.

84.   Yu, J.E., Han, S.Y., Wolfson, B., and Zhou, Q. (2018). The role of endothelial lipase in lipid
metabolism, inflammation, and cancer. Histol Histopathol *33*, 1-10. 10.14670/HH-11-905.

85.   Wang, J.W., Gamsby, J.J., Highfill, S.L., Mora, L.B., Bloom, G.C., Yeatman, T.J., Pan, T.C., Ramne,
A.L., Chodosh, L.A., Cress, W.D., et al. (2004). Deregulated expression of LRBA facilitates
cancer cell growth. Oncogene *23*, 4089-4097. 10.1038/sj.onc.1207567.

86.   Liu, Y., Zheng, X., Yu, Q., Wang, H., Tan, F., Zhu, Q., Yuan, L., Jiang, H., Yu, L., and Zeng, S.
(2016). Epigenetic activation of the drug transporter OCT2 sensitizes renal cell carcinoma to
oxaliplatin. Sci Transl Med *8*, 348ra397. 10.1126/scitranslmed.aaf3124.

87.    Man, K.F., Zhou, L., Yu, H., Lam, K.H., Cheng, W., Yu, J., Lee, T.K., Yun, J.P., Guan, X.Y., Liu, M., and Ma, S. (2023). SPINK1-induced tumor plasticity provides a therapeutic window for chemotherapy in hepatocellular carcinoma. Nat Commun *14*, 7863. 10.1038/s41467-023-43670-9.

88.    Reith, W., LeibundGut-Landmann, S., and Waldburger, J.M. (2005). Regulation of MHC class II gene expression by the class II transactivator. Nat Rev Immunol *5*, 793-806. 10.1038/nri1708.

89.    Lin, J., Jia, P., Wang, S., Kosters, W., and Ye, K. (2023). Comparison and benchmark of structural variants detected from long read and long-read assembly. Brief Bioinform *24*. 10.1093/bib/bbad188.

90.    Sahlin, K., Baudeau, T., Cazaux, B., and Marchet, C. (2023). A survey of mapping algorithms in the long-reads era. Genome Biol *24*, 133. 10.1186/s13059-023-02972-3.

91.    Hickey, G., Monlong, J., Ebler, J., Novak, A.M., Eizenga, J.M., Gao, Y., Human Pangenome Reference, C., Marschall, T., Li, H., and Paten, B. (2024). Pangenome graph construction from genome alignments with Minigraph-Cactus. Nat Biotechnol *42*, 663-673. 10.1038/s41587-023-01793-w.

92.    Li, H., Feng, X., and Chu, C. (2020). The design and construction of reference pangenome graphs with minigraph. Genome Biol *21*, 265. 10.1186/s13059-020-02168-z.

93.    Nurk, S., Koren, S., Rhie, A., Rautiainen, M., Bzikadze, A.V., Mikheenko, A., Vollger, M.R., Altemose, N., Uralsky, L., Gershman, A., et al. (2022). The complete sequence of a human genome. Science *376*, 44-53. 10.1126/science.abj6987.

94.    Vollger, M.R., Guitart, X., Dishuck, P.C., Mercuri, L., Harvey, W.T., Gershman, A., Diekhans, M., Sulovari, A., Munson, K.M., Lewis, A.P., et al. (2022). Segmental duplications and their variation in a complete human genome. Science *376*, eabj6965. 10.1126/science.abj6965.

95.    Jain, M., Koren, S., Miga, K.H., Quick, J., Rand, A.C., Sasani, T.A., Tyson, J.R., Beggs, A.D., Dilthey, A.T., Fiddes, I.T., et al. (2018). Nanopore sequencing and assembly of a human genome with ultra-long reads. Nat Biotechnol *36*, 338-345. 10.1038/nbt.4060.

96.    Li, H., Li, K., Shu, D., Shen, M., Tan, Z., Zhang, W., Pu, D., Tan, W., Tang, Z., Jin, A., and Liu, S. (2022). MED16 Promotes Tumour Progression and Tamoxifen Sensitivity by Modulating Autophagy through the mTOR Signalling Pathway in ER-Positive Breast Cancer. Life (Basel) *12*. 10.3390/life12101461.

97.    Brahmer, J., Reckamp, K.L., Baas, P., Crino, L., Eberhardt, W.E., Poddubskaya, E., Antonia, S., Pluzanski, A., Vokes, E.E., Holgado, E., et al. (2015). Nivolumab versus Docetaxel in Advanced Squamous-Cell Non-Small-Cell Lung Cancer. N Engl J Med *373*, 123-135. 10.1056/NEJMoa1504627.

98.    Battaglia, S., Dong, K., Wu, J., Chen, Z., Najm, F.J., Zhang, Y., Moore, M.M., Hecht, V., Shoresh, N., and Bernstein, B.E. (2022). Long-range phasing of dynamic, tissue-specific and allele-specific regulatory elements. Nat Genet *54*, 1504-1513. 10.1038/s41588-022-01188-8.

99.    Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol *37*, 907-915. 10.1038/s41587-019-0201-4.

100.    Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., and Genome Project Data Processing, S. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078-2079. 10.1093/bioinformatics/btp352.

101.    Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.C., Mendell, J.T., and Salzberg, S.L. (2015).

StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat

Biotechnol *33*, 290-295. 10.1038/nbt.3122.

102.    Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with

high-throughput sequencing data. Bioinformatics *31*, 166-169.

10.1093/bioinformatics/btu638.

103.    Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550. 10.1186/s13059-014-0550-8.

104.    De Coster, W., D'Hert, S., Schultz, D.T., Cruts, M., and Van Broeckhoven, C. (2018). NanoPack:

visualizing and processing long-read sequencing data. Bioinformatics *34*, 2666-2669.

10.1093/bioinformatics/bty149.

105.    Su, J., Zheng, Z., Ahmed, S.S., Lam, T.W., and Luo, R. (2022). Clair3-trio: high-performance

Nanopore long-read variant calling in family trios with trio-to-trio deep neural networks. Brief

Bioinform *23*. 10.1093/bib/bbac301.

106.    Boeva, V., Popova, T., Bleakley, K., Chiche, P., Cappo, J., Schleiermacher, G., Janoueix-Lerosey,

I., Delattre, O., and Barillot, E. (2012). Control-FREEC: a tool for assessing copy number and

allelic content using next-generation sequencing data. Bioinformatics *28*, 423-425.

10.1093/bioinformatics/btr670.

107.    do Valle, I.F., Giampieri, E., Simonetti, G., Padella, A., Manfrini, M., Ferrari, A., Papayannidis,

C., Zironi, I., Garonzi, M., Bernardi, S., et al. (2016). Optimized pipeline of MuTect and GATK

tools to improve the detection of somatic single nucleotide polymorphisms in whole-exome

sequencing data. BMC Bioinformatics *17*, 341. 10.1186/s12859-016-1190-7.

108.    Carter, S.L., Cibulskis, K., Helman, E., McKenna, A., Shen, H., Zack, T., Laird, P.W., Onofrio, R.C.,

Winckler, W., Weir, B.A., et al. (2012). Absolute quantification of somatic DNA alterations in

human cancer. Nat Biotechnol *30*, 413-421. 10.1038/nbt.2203.

109.    Morishita, S., Ichikawa, K., and Myers, E.W. (2021). Finding long tandem repeats in long noisy

reads. Bioinformatics *37*, 612-621. 10.1093/bioinformatics/btaa865.

1469

# Acknowledgments

We thank members of the Dan Xie lab for useful discussions. This work was supported by the National Natural Science Foundation of China 82173383 and 82241238 (to D.X.), 32200508 (to L.X.), 32300522 (to K.L.T), 32300003 (to H.W.), 82371883 (to J.F.L.), 82472365 (to H.T.L.), the Sichuan Province Science and Technology Program 2025ZNSFSC1098 (to K.L.T.), and the 1·3·5 project for disciplines of excellence, West China Hospital, Sichuan University ZYYC23024 (to D.X.).

1477

# Author contributions:

Conceptualization: D.X.; Data Curation: W.L., Z.F.W., Z.W., C.L., J.F.L., Y.X.Q., Q.W.Z., H.T.L., S.Z.H., Z.Y.Z., A.L., A.L.W.; Software, Formal Analysis and Methodology: L.X.,

1481 K.L.T., X.Y.L., Q.L.Z., R.L.W., Z.T.F., X.N.P.; Validation: H.W., M.L.L., Y.H., J.W.;

1482 Supervision: D.X.; Writing – Original Draft: D.X., L.X., K.L.T., H.W., X.Y.L., Y.H.L., C.Y.Y.,

1483 Z.T.F.; Writing – Review & Editing: all authors.

1484

1485 **Declaration of interests**

1486 The authors declare no competing interests.

1487

# Figure Legends

**Figure 1. Genome-wide detection of somatic insertions in pan-cancer genomes.**

a.  Overview of a pan-cancer cohort comprising 325 paired tumor and blood samples from 299 patients across 12 cancer types: 1) glioblastoma (GBM); 2) oral squamous cell carcinoma (OSCC); 3) laryngeal cancer (LC); 4) lung adenocarcinoma (LUAD); 5) lung squamous cell carcinoma (LUSC); 6) hepatocellular carcinoma (HCC); 7) gastric carcinoma (GC); 8) pancreatic adenocarcinoma (PAAD); 9) intestinal carcinoma (IC);

10) ovarian cancer (OV); 11) uterine corpus endometrial carcinoma (UCEC); 12)

cervical squamous cell carcinoma (CCA).

b.    Scheme of the bioinformatics pipeline used to identify somatic TRIs. The top track

('reference') shows the hg38 genome. the second track ('somatic') shows reads

alignments supporting somatic SVs mapped to the reference genome, and the third

track ('germline') shows reads alignments supporting germline SVs mapped to the

reference genome. Circles represent repeat motifs, with different colors indicating

different repeat motifs.

c.    Scheme of the bioinformatics pipeline for identifying somatic TEI. The first blue track

represents alignments from the tumor sample mapped to the reference genome,

while the second orange track shows alignments from the paired blood sample. After

initial detection (Method), inserted sequences are corrected. Following annotation

(Method), somatic TEIs are classified into two categories: complex TEIs (upper track)

or solo TEIs (lower track).

d.    Somatic insertion frequency (ln) across cancer types. Somatic insertion frequencies

were calculated based on the total number of somatic TRIs and TEIs per sample.

e.    Ideogram depicting the distribution of somatic TEI (blue, left) and somatic TRI (red,

right). The ideogram shows the genomic distribution of somatic TEIs (blue) and

somatic TRIs (red), with darker colors indicating higher frequencies of TEIs or TRIs

within each 1Mb window. The top 30 high-frequency 1Mb windows are highlighted

according to the following features: regions with clustered TEs near characteristic

genes are marked with a gray background; high-frequency windows located in

telomeric, sub-telomeric, centromeric regions are enclosed in a box.

1521

1522

1523

**Figure 2. Sequence features and mechanisms of somatic TRIs and somatic TEIs.**

a. Schematic representation of three somatic TRI motif mutation patterns, namely expansion, motif change and other, arranged from top to bottom. In each subpanel, the first row illustrates the somatic repeat patterns, whereas the second row presents the corresponding germline repeat patterns. Different repeat motifs are color-coded for clarity.

b. Ternary phase diagram illustrating the relative proportions of expansion, motif-change, and other events in somatic TRIs. Categories include: Expansion (orange), Expansion & Motif-Change (yellow), Motif-Change (blue), Undefined (dark blue), and Ambiguous (gray).

c.   Proportion of the five categories of somatic TRI: Expansion, Expansion &

Motif-Change, Motif-Change, Undefined, and Ambiguous.

d.   Schematic representation of a somatic complex TEI sequence. The color coding of

the bases indicates their origin, with yellow-highlighted sequences representing

target site duplication (TSD) sequences. For LINE/L1 sequences, 5755 base pairs

were omitted, and for SINE/Alu sequences, 141 base pairs were omitted.

e.   Pie charts shows the proportion of somatic complex (blue) and solo (orange) TEI.

f.   Proportion of different TE families within all somatic TEIs. 'SINE' indicates TEIs

containing SINE elements but lacking LINE elements; 'LINE' indicates TEIs

containing LINE elements but lacking SINE elements; 'SINE+LINE' denotes TEIs

containing both SINE and LINE elements.

g.   Schematic illustration of somatic TEI insertion mechanisms.

h.   Summary of somatic TEI insertion mechanisms. The bubble chart shows the relative

proportions of insertion mechanisms for complex and solo TEIs across various

cancer types, with bubble size and color intensity corresponding to the magnitude of

these proportions. A pie chart summarizes the overall distribution of insertion

mechanisms for complex and solo somatic TEIs. TSD: target site duplication; NAHR:

non-allelic homologous recombination; microhomology: microhomology

recombination.

1553

1554

1555

**Figure 3. High-frequency somatic TRIs located in the intron of *SHROOM2*.**

a. Bar plot showing the number of patients harbored somatic TRIs in each high-frequency TRI window (>= 10 patients), along with the pathways enrichment analysis of genes located within or near these high-frequency somatic TRI windows.

b. Genome browser tracks displaying the ENCODE-annotated genes, and ENCODE-predicted cis-regulatory elements (cCREs) surrounding somatic TRIs in the SHROOM2 gene locus. Lower panel: Epigenomic signals around somatic TRIs in SHROOM2, as visualized by chromatin marks.

c. Detecting somatic TRI events harbored by *SHROOM2* through PCR in clinical samples, blood DNA as a control.

d. The detailed sequence overview of somatic TRIs harbored by *SHROOM2* in different

1567      clusters. The ATCG bases are colored differently, with gaps represented as short

1568      dashes.

1569    e. Distribution of somatic TRI lengths in *SHROOM2* across different clusters.

1570    f. Bar chart showing the distribution of cancer types in patients harboring somatic TRIs

1571      within SHROOM2 across the different clusters.

1572

1573

1574

**Figure 4. The characteristic of somatic TRI clusters**

a. Clustering heatmap of somatic TRI frequencies across cancer types.

b. Proportion of tumor samples with somatic TRIs. The number represents the proportion

of mutations within the cancer type. Different colors represent various percentage

categories.

c.  Frequency distribution of somatic TRIs across cancer types. The bars at the bottom

represent the mutation frequency of the mutation within individual cancer types, with

colors indicating different cancer types.

d.  Bar plots showing the frequency of HCC associated high-frequency somatic TRI

across cancer types. The high frequency cancer types (over 30%) are filled in red.

e-f. Bar plots showing the frequency distribution of high-frequency somatic TRI in

IC&PAAD (e), as well as OSCC, LC and LUSC (f) clusters compared to other cancer types.

Bars of significant cancer types are filled in red.

g.   Browser plot displaying the epigenomic signal, ENCODE-annotated gene transcript,

and cCREs (Candidate Cis-Regulatory Elements) predicted by ENCODE around the

upstream somatic TRIs of the OBI1 gene. The detailed sequence overview of different

clusters surrounding the upstream somatic TRIs of OBI1 is presented in the lower

enlarged panel, with the length distribution of these 2 clusters color-coded.

1593

**Figure 5. Clusters of somatic TEIs.**

a. Heatmap showing the clustering results of somatic TEIs. Each column and each row represent Jaccard similarities of a sample compared to others. Hierarchical clustering was performed according to Jaccard similarity with Ward.D linkage. Each cluster has a pie chart to indicate the proportions of adenocarcinoma, squamous cell carcinoma, and others. Two SCC-subclusters of cluster 1 are distinguished.

b. Pies showing the proportional of 5 clusters within the three major histology categories:

1603   SCC (squamous cell carcinoma), ADC (adenocarcinoma), and Others.

1604 c. A stacked bar chart displays the distribution percentages of different cancer types

1605   across five clusters.

1606 d. Proportional distribution of ADC (adenocarcinoma), SCC (squamous cell carcinoma),

1607   and others within subclusters of Cluster 1.

1608 e. A boxplot depicting the number of somatic TEIs (log10) within 5 clusters, with the

1609   p-value calculated using t-test. ns $p>0.05$, $*p < 0.05$, $**p < 0.005$, $***p < 0.0005$

1610

1611

1612

**Figure 6. The MHC II gene family harbored high frequency somatic TRIs.**

a.  Distribution of the number of shared TEI samples. Gray: random distribution (Methods); Orange: observed distribution. The x-axis represents the number of samples sharing a somatic TEI, and the y-axis represents the count of somatic TEIs.

b.  Scatter plot illustrating the number of samples sharing somatic TEIs and the count of somatic TEIs. Each dot represents a 1Mb genomic window, with the color indicating the source of its chromosomal coordinates. The genomic window containing the highest number of TEIs, chr6:32,000,000-33,000,000, is highlighted.

c.   Bar plots showing the number of somatic transposable element insertions (TEIs)

within 500 bp windows in LUAD (first track), LUSC (second track), and high-burden

samples (third track). The fourth track displays the cumulative number of somatic

TEIs within 500 bp windows, with bars color-coded according to the type of TEI. Each

bar represents a 500 bp window. The browser plot includes epigenomic signals,

ENCODE-annotated gene transcripts, and candidate cis-regulatory elements (cCREs)

predicted by ENCODE around the somatic TEIs within the MHC II gene family.

d.   Detecting somatic TEIs events through PCR in some clinical samples, blood DNA as

a control (A: ALL; H: HCC; L: LC; D: DRB5).

e.   Detecting somatic TEIs events through FISH in in some sample tissue slices,

CK7/serpinB3/NES/GAFP is the marker for GC/CCA/LUNG/GBM respectively. WT

refers to the sequence without somatic TEIs events, TE contains the sequence of

somatic TEIs events. Flow cytometric analyses of CD40L mobilization in CD4+ T cells

after 24 hours of in vitro co-culture with tumor cells.

1635

1636

1637

**Figure S1. Summary of long-read sequencing data.**

a. Sequencing depth of long-read data from 325 paired tumor (below) and paired blood (above) samples across 12 cancer types and high-burden samples.

b. Scatter plots showing the N50 and mean quality of LRS reads derived from paired tumor (gray) and blood (white) samples, spanning 12 cancer types and high-burden samples. Each dot represents a sample.

1644     c.    Box plots comparing sequencing quality between R9 (blue) and R10 (yellow) chips,

1645           with statistical significance assessed by the Wilcoxon rank test. Each dot represents

1646           a sample.

1647     d.    Pie chart illustrating the distribution of sequencing chip types used across various

1648           cancer types and high-burden samples. Each color represents a different chip

1649           version.

1650

1651

**Figure S2. Overview of Methods for Identifying Somatic Genomic Alterations and The Frequency of Somatic TRIs/TEIs.**

a.  Schematic representation of the workflow for identifying somatic TRI. i. Candidate somatic TRI regions are identified using Sniffles2 and Straglr; ii. Reads spanned these regions are extracted, and edit distances are calculated between sequences spanned potential somatic TRI region, followed by clustering to initial screen the presence of somatic TRIs; iii. Somatic TRIs are confirmed using TDscope; iv. Repeat

motif of somatic TRIs are annotated (see Methods for details).

b.  Schematic representation of the workflow for identifying somatic TEI. i. Reads from

tumor and blood samples are aligned to the reference genome; ii. Candidate somatic

SVs are identified using Sniffles2.; iii. After polishing the reference genome, reads are

realigned to the refined reference to validate the presence of candidate SVs; iv.

Somatic SVs are further refined using IRIS; v. Somatic SVs are annotated (see

Methods for details).

c.  The upper panel shows the frequency (log10 scale) of somatic TRIs, while the lower

panel displays the frequency of somatic TEIs across various cancer types. Each dot

represents an individual sample, with distinct colors indicating different cancer types.

Samples with high burden of TRIs or TEIs are labeled accordingly.

1670

1671

1672

**Figure S3. Tumor Purity and Characteristics of Somatic REIs.**

a.  Box plot of tumor purity across different cancer types. Each box represents a specific

cancer type, with the y-axis indicating tumor purity percentage.

b-d. Scatter plots showing the correlation between tumor purity and the count of somatic

TRIs (e), somatic TEIs (f), and the combined count of somatic TRIs and TEIs (d). Data points represent individual samples, colored by cancer type. Pearson correlation coefficients (R²) and p-values are displayed in the upper-right corner of each plot. Boxplot showing the log-transformed number of somatic TRIs (blue) and somatic TEIs (yellow) across different cancer types: "Other" (including HCC, GBM), Squamous (LUSC, CCA, OSCC, LC), and Adenocarcinoma (LUAD, PAAD, GC, IC, OV, UCEC). High-burden samples are indicated separately. Statistical significance is assessed using a two-tailed t-test.

e.    Boxplot illustrating the log-transformed number of somatic TRIs (blue) and somatic TEIs (yellow) in Adenocarcinoma, Squamous, Other and high-burden samples. p-values were calculated using the Wilcoxon rank-sum test.

f.    Boxplot illustrating the log-transformed number of somatic TRIs and somatic TEIs in LUSC (yellow) and LUAD (orange) samples. p-values were calculated using the Wilcoxon rank-sum test.

g.    A bar chart depicting the proportion of somatic TEIs (yellow) and somatic TRIs (blue) across cancer types and high-burden samples. Each bar represents a distinct sample. A pie chart above summarizes the overall percentage of somatic TRIs and TEIs within each cancer type.

h.    Bar plot showing the GO Molecular Function enrichment analysis of genes associated with high-burden specific TRIs, as predicted by the GREAT. The x-axis represents the -log10 transformed FDR from the binomial test.

i.    Boxplots comparing the expression levels of *ERCC3*, *RAD51*, and *XRCC4* in high-burden LUAD samples (red) and low-burden LUAD samples (blue). Statistical significance was assessed using a two-tailed t-test.

1702

1703

**Figure S4. The genome distribution of somatic TRIs and TEIs.**

a.  Distribution of the distance from somatic TRIs (yellow, above) and somatic TEIs (blue,

below) to telomeres. The x-axis represents the distance from somatic insertions to

the telomeric ends, while the y-axis indicates the number of samples. Two red dashed

lines mark the positions of 0 bp (telomere ends) and 5 Mbp from the telomere ends.

b. Cumulative distribution of somatic TRI (left) and somatic TEI (right). The plot reveals

a more rapid saturation of somatic TRIs at the ends of chromosome arms. A sliding

window of 500,000bp, starting from the telomeric regions and extending toward the

centromere, was used to cumulatively count somatic TRIs and TEIs. The x-axis is

truncated at 20 Mbp.

c. Chromosome-level distribution of various somatic TEI types ("observed") compared

to TEs in the hg38 reference genome ("expected"). The fold enrichment of observed

TEIs relative to random expectations is calculated refer to the previous study [58]. Each

density plot corresponds to a distinct type of TEI, represented by different colors.

d. Enrichment of somatic TRIs (left) and somatic TEIs (right) on genomic elements. The

ratio of observed insertions to random expectation (O/E) is calculated using a

hypergeometric test. *p < 0.05. The y-axis is presented on a log scale for O/E values.

Somatic TRIs and TEIs are shown in different colors to distinguish between the two

insertion types.

e. Oncoplot shows high frequency (Sample frequency > 5% ) COSMIC cancer genes

with somatic REI events located within gene body and 2kbp flanking regions. The bar

plot on the top and right show the number of somatic REI events located within

samples and gene flank regions with color coded in the bottom,

**Figure S5. Sequence charateristic of somatic TRIs and somatic TEIs.**

a. Schematic representation of somatic TRI patterns for Expansion, Motif Change, and Other. The top row represents somatic sequences, the middle row indicates aligned/unaligned, and the bottom row shows germline sequences.

b. Proportions of each somatic TRI category in cancer types: Expansion (orange), Expansion & Motif-Change (yellow), Motif-Change (blue), Ambiguous (dark blue) and

Unknown (gray).

c.   Violin plot displaying the log-transformed length distribution for somatic complex (blue)

and solo (yellow) TEI.

d.   Proportions of somatic complex and solo TEI across various cancer types. 'complex

SINE' indicates complex TEIs containing SINE elements but lacking LINE elements;

'complex LINE' indicates complex TEIs containing LINE elements but lacking SINE

elements; 'complex SINE+LINE' denotes TEIs containing both SINE and LINE

elements.

e.   Frequency (log10) of SINE (orange), LINE (blue), LTR (yellow) and Retroposon

(purple) subfamilies in adenocarcinoma, squamous cell carcinoma and other cancer

type. Each column represents a subfamily. 'ADC': adenocarcinoma, 'SCC':

squamous cell carcinoma, 'other': HCC and GBM.

f.   Length density distribution (log10 scale) of somatic TEIs in squamous cell carcinoma,

other cancer types, adenocarcinoma, and high-burden samples. Different colors

represent different cancer types, similar to panel h.

g.   Log-transformed counts of different subfamilies of somatic TEIs across cancer types.

"ADC" represents adenocarcinoma, "SCC" stands for squamous cell carcinoma, and

"other" includes HCC and GBM.

h.   Density plot of log-transformed length distribution of TSD across different cancer

types.

1755

**Figure S6. Examples of somatic TEI insertion mechanisms**

a. The schematic illustrates example of Target Site Duplication. In IGV, the reference genome comprises assembled sequences containing TEI-supporting reads, with all blood reads and tumor TEI-supporting reads aligned to this assembly. Complex TEI sequences are highlighted in light green, and homologous sequences are marked in

yellow-green. LINE elements are colored in blue, while SINE elements are colored in

orange-red. Target site duplications are marked in bright yellow.

b. The schematic illustrates example of Recombination (NAHR). In IGV, the reference

genome comprises assembled sequences containing TEI-supporting reads, with all

blood reads and tumor TEI-supporting reads aligned to this assembly. Complex TEI

sequences are highlighted in light green, and homologous sequences are marked in

yellow-green. LINE elements are colored in blue, while SINE elements are colored in

orange-red. Homologous sequences are marked in bright yellow.

c. The schematic illustrates example of Recombination (Microhomology). In IGV, the

reference genome comprises assembled sequences containing TEI-supporting reads,

with all blood reads and tumor TEI-supporting reads aligned to this assembly.

Complex TEI sequences are highlighted in light green, and homologous sequences

are marked in yellow-green. LINE elements are colored in blue, while SINE elements

are colored in orange-red. Homologous sequences are marked in bright yellow.

d. The schematic illustrates example of Other. In IGV, the reference genome comprises

assembled sequences containing TEI-supporting reads, with all blood reads and

tumor TEI-supporting reads aligned to this assembly. Complex TEI sequences are

highlighted in light green, and homologous sequences are marked in yellow-green.

LINE elements are colored in blue, while SINE elements are colored in orange-red.

1782

1783

**Figure S7. Overview of TRI distribution**

a.  Bar plot depicting the number of patients sharing the same somatic TRI events. Gray

bars indicate the random distribution (Method), while orange bars represent the

observed distribution. The x-axis represents the number of patients sharing somatic

TRIs, and the y-axis represents the count of TRIs.

b.  Bar plot illustrating the percentage distribution of low-frequency, high-frequency, and

background TRIs located within intergenic, intronic, and coding sequence (CDS)

regions.

c. Volcano plot depicting the differential enrichment between high and low frequency

somatic TRIs on known 490 TF binding regions from ENCODE project. Elements with

p-value < 0.05 (Chi-square test) and |log2(Odds Ratio)| > 1 are highlighted in red.

d. Volcano plot illustrating high-frequency TRIs with significant changes in CpG counts

between somatic and matched germline TRI sequences. Orange: TRIs with FDR <

0.05 and log2(FoldChange) > 1 (Wilcoxon test); Blue:FDR < 0.05 and

log2(FoldChange) < 1 (Wilcoxon test).

e. Box plot displaying the proportions of methylated and unmethylated samples at 18

high-frequency somatic TRIs introduced methylation marks at generated CpG

dinucleotides (Wilcoxon test, p < 0.05). Each dot representing the proportion of

methylated (blue) and unmethylated (pink) states for each sample within each TRI

window.

f. Browser plot displaying the epigenomic signal, ENCODE-annotated gene transcript,

and cCREs (Candidate Cis-Regulatory Elements) predicted by ENCODE around the

somatic TRI located in the intron of the EYS gene.

g. An enlarged panel providing a detailed sequence overview of different clusters

surrounding the somatic TRIs in the EYS intron.

h. Density plot depicting the length distribution of two clusters, color-coded for clarity.

i. Methylation analysis of the TRI sequence from a representative sample: the top panel

displays overall sequences of somatic and corresponding germline with pairwise

alignment details; the middle panel highlights the specific somatic CpG insertion site;

the bottom panel shows all tumor reads supporting the TRI event, along with

methylation status at the somatic CpG insertion site. Methylated CpG sites are

colored in red and unmethylated CpG sites are colored in blue.

1816

1817

1818 **Figure S8. Sanger sequencing of Shroom2-related SRE in clinical samples and**

1819 **detection in cell lines**

1820 a. Visual display of sanger sequencing of Shroom2-related SRE in clinical samples.

1821 b. Detection of Shroom2-related SRE in liver cancer and intestinal cancer cell lines.

1822

1823

1824

**Figure S9. The browser plot of somatic TRIs located in the intron of *SHROOM2*,**

**along with the somatic TRIs sequence displays for HCC and IC.**

The browser plot illustrates epigenomic signals from ENCODE project, including cell line

H3K27ac chip-seq and DNase signal, gene transcript Candidate Cis-Regulatory Elements

(cCREs), and H3K27ac marks identified by ENCODE.The somatic TRI (chrX:

9791969-9792160) is highlighted in blue. The bottom panel illustrates detailed somatic

sequences of HCC and IC samples harbored the somatic TRI event

with the motif AATAG abbreviated and highlighted in red.

1833

1834

**Figure S10. Three somatic TRIs that associated with the expression of PTP family genes in NSCLC.**

a.  Volcano plot showing differentially expressed genes between NSCLC samples with and without high-frequency somatic TRIs. Genes from the PTP family are highlighted with text. Dotted lines parallel to the X-axis indicates where p = 0.05, while two vertical dashed lines represent log2(Fold change) values of -1 and 1, respectively.

b.   Box plot of the expression levels of differentially expressed PTP family genes with and

without somatic TRI. Orange: with TRI; Gray: without TRI. P values were calculated by

t-test.

c.   The browser plot of somatic TRIs (chr7:121603262-121603506) illustrates the

following epigenomic signals: (1) H3K27ac in the A673 cell line, (2) H3K27ac in the

SK-N-MC cell line, (3) DNase-seq in the A673 cell line, (4) DNase-seq in the SK-N-MC

cell line. Additionally,it displays the ENCODE-annotated gene transcript (5), Candidate

Cis-Regulatory Elements (cCREs) predicted by ENCODE (6), and H3K27ac marks

identified by ENCODE (7). The plot concludes with the CTTC motif of the repeat. An

enlarged panel below outlines a detailed sequence overview of various clusters, with

vertical labels indicating cluster and cancer type information.

d.   The browser plot of somatic TRIs (chr1:202204213-202204518) illustrates the

following epigenomic signals: (8) DNase-seq in the H7 cell line, (9) DNase-seq in the

A549 cell line, (10) H3K27ac in the RWPE1 cell line, (11) H3K27ac in the A549 cell

line. Additionally,it displays the ENCODE-annotated gene transcript (5), Candidate

Cis-Regulatory Elements (cCREs) predicted by ENCODE (6), and H3K27ac marks

identified by ENCODE (7). The plot concludes with the CA motif of the repeat. An

enlarged panel below outlines a detailed sequence overview of various clusters, with

vertical labels indicating cluster and cancer type information.

e.   The browser plot of somatic TRIs (chr1:202204213-202204518) illustrates the

following epigenomic signals: (12) H3K27ac in the RWPE1 cell line, (13) H3K27ac in

the H9 cell line, (14) DNase-seq in the RWPE1 cell line, (15) DNase-seq in the H9 cell

line. Additionally,it displays the ENCODE-annotated gene transcript (5), Candidate

Cis-Regulatory Elements (cCREs) predicted by ENCODE (6), and H3K27ac marks

identified by ENCODE (7). The plot concludes with the TGGTGA motif of the repeat.

An enlarged panel below outlines a detailed sequence overview of various clusters,

with vertical labels indicating cluster and cancer type information.

f.   Frequency of somatic TRI occurrence around PTP genes across cancer types.

Orange: with TRI; Gray: without TRI.

g.   Oncoplot of samples associated with the PTP Family somatic TRI windows. The bar

charts on the top and right show the counts, while the percentages on the left indicate

the proportion among all samples.

1873

1874

1875

**Figure S11. Display of representative somatic TRI events**

a. The browser plot of somatic TRIs (chr1:99683035-99683173) with significantly high frequency in HCC, OSCC and LC illustrates the following epigenomic signals: (1) H3K27ac in the HeLa-S3 cell line, (2) H3K27ac in the SJSA1 cell line, (3) H3K27ac in the ACC112 cell line, (4) DNase-seq in the HeLa-S3 cell line, and (5) DNase-seq in

the SJSA1 cell line. Additionally, it displays the ENCODE-annotated gene transcript

(6), Candidate Cis-Regulatory Elements (cCREs) predicted by ENCODE (7), and

H3K27ac marks identified by ENCODE (8). The plot concludes with the AAAG motif of

the repeat. An enlarged panel below outlines a detailed sequence overview of various

clusters, with vertical labels indicating cluster and cancer type information.

b.   The browser plot of somatic TRIs (chr18:49579001-49579348) with significantly high

frequency in HCC and IC illustrates the following epigenomic signals: (9) H3K27ac in

the Caco-2 cell line, (10) H3K27ac in the Karpas-422 cell line, (11) DNase-seq in the

Caco-2 cell line, (12) DNase-seq in the Karpas-422 cell line, it displays the

ENCODE-annotated gene transcript (6), Candidate Cis-Regulatory Elements (cCREs)

predicted by ENCODE (7), and H3K27ac marks identified by ENCODE (8). The plot

concludes with the GGGAGA motif of the repeat. An enlarged panel below outlines a

detailed sequence overview of various clusters, with vertical labels indicating cluster

and cancer type information.

c.   The browser plot of somatic TRIs (chr1:34802227-34802447) with significantly high

frequency in HCC and OV illustrates the following epigenomic signals: (13) H3K27ac

in the HFFc6 cell line, (14) DNase-seq in the K562 cell line, (15) DNase-seq in the

HFFc6 cell line, and H3K27ac marks identified by ENCODE (8). The plot concludes

with the GGTTT motif of the repeat. An enlarged panel below outlines a detailed

sequence overview of various clusters, with vertical labels indicating cluster and

cancer type information.

d.   The browser plot of somatic TRIs (chr4:150658505-150658775) with significantly high

frequency in IC and PAAD illustrates the following epigenomic signals: (16) H3K27ac

in the H9 cell line, (17) DNase-seq in the H9 cell line, (18) DNase-seq in the KBM-7

cell line, it displays the ENCODE-annotated gene transcript (6), Candidate

Cis-Regulatory Elements (cCREs) predicted by ENCODE (7), and H3K27ac marks

identified by ENCODE (8). The plot concludes with the GGTTT motif of the repeat. An

enlarged panel below outlines a detailed sequence overview of various clusters, with

vertical labels indicating cluster and cancer type information.

e.   The browser plot of somatic TRIs (chr6:160218690-160218725) with significantly high

frequency in IC and PAAD illustrates the following epigenomic signals: (19) H3K27ac

in the WTC11 cell line, (20) DNase-seq in the WTC11 cell line, (21) DNase-seq in the

RCC cell line. Additionally, it displays the ENCODE-annotated gene transcript (6),

Candidate Cis-Regulatory Elements (cCREs) predicted by ENCODE (7), and

H3K27ac marks identified by ENCODE (8). The plot concludes with the

CAGCAGCAACAG motif of the repeat. An enlarged panel below outlines a detailed

sequence overview of various clusters, with vertical labels indicating cluster and

cancer type information.

1919

**Figure S12. Display of representative somatic TRI events**

a. The browser plot of somatic TRIs (chr5:147837649-147837716) with significantly high frequency in IC and PAAD illustrates the following epigenomic signals: (1) H3K27ac in the 22Rv1 cell line, (2) H3K27ac in the A549 cell line, (3) DNase-seq in the Caco-2 cell line, (4) DNase-seq in the A549 cell line. Additionally, it displays the ENCODE-annotated gene transcript (5), Candidate Cis-Regulatory Elements (cCREs) predicted by ENCODE (6), and H3K27ac marks identified by ENCODE (7). The plot concludes with the TTTC motif of the repeat. An enlarged panel below outlines a detailed sequence overview of various clusters, with vertical labels indicating cluster and cancer type information.

b. The browser plot of somatic TRIs (chr12:74615532-74615609) illustrates the following

1933 epigenomic signals: (8) H3K27ac in the MCF-7 cell line, (9) DNase-seq in the MCF-7

1934 cell line, (10) DNase-seq in the RCC 7860 cell line. Additionally, it displays the

1935 ENCODE-annotated gene transcript (5), Candidate Cis-Regulatory Elements (cCREs)

1936 predicted by ENCODE (6), and H3K27ac marks identified by ENCODE (7). The plot

1937 concludes with the TACATATTTA motif of the repeat. An enlarged panel below outlines

1938 a detailed sequence overview of various clusters, with vertical labels indicating cluster

1939 and cancer type information.

1940 c. The browser plot of somatic TRIs (chr19:878845-879150) illustrates the following

1941 epigenomic signals: (11) H3K27ac in the Caco-2 cell line, (12) DNase-seq in the

1942 Caco-2 cell line, (13) DNase-seq in the L1-S8R cell line, Additionally, it displays the

1943 ENCODE-annotated gene transcript (5), Candidate Cis-Regulatory Elements (cCREs)

1944 predicted by ENCODE (6). The plot concludes with the CCCAG motif of the repeat. An

1945 enlarged panel below outlines a detailed sequence overview of various clusters, with

1946 vertical labels indicating cluster and cancer type information.

1947 d. The browser plot of somatic TRIs (chr4:86842786-86842877) illustrates the following

1948 epigenomic signals: (14) H3K27ac in the RWPE1 cell line, (15) H3K27ac in the

1949 RWPE2 cell line, (16) H3K27ac in the MCF 10A cell line, (17) DNase-seq in the

1950 RWPE2 cell line, and (18) DNase-seq in the MCF 10A cell line. Additionally, it displays

1951 the ENCODE-annotated gene transcript (5), Candidate Cis-Regulatory Elements

1952 (cCREs) predicted by ENCODE (6), and H3K27ac marks identified by ENCODE (7).

1953 The plot concludes with the TTTC motif of the repeat. An enlarged panel below

1954 outlines a detailed sequence overview of various clusters, with vertical labels

1955 indicating cluster and cancer type information.

1956

1957

1958 **Figure S13. Validation of clinical samples for TRI events.**

1959 a. Detecting somatic TRI events harbored by *PALMD* through PCR in clinical samples,

1960 blood DNA as a control.

1961 b. Detecting somatic TRI events harbored by *SPINK1* through PCR in clinical samples,

1962 blood DNA as a control.

1963 c. Detecting somatic TRI events harbored by *SLC10A6* through PCR in clinical samples,

1964 blood DNA as a control.

1965 d. Detecting somatic TRI events harbored by *OBI1* through PCR in clinical samples,

1966 blood DNA as a control.

1967

1968

1969

**Figure S14. The characteristic of somatic TEI clusters.**

a. Distribution of the 5 clusters of somatic TEI patterns in 12 cancer types.

b. Distribution of the 12 cancer types in subclusters of Cluster 1. 'other' presents samples out of subcluster1 and subcluster2.

c. Density plot showing the distribution of somatic transposable element insertions (TEIs) in Cluster 1, separated into subcluster1 (yellow), subcluster2 (blue), and the remaining Cluster 1 samples (gray). The x-axis represents the number of somatic TEIs.

d. Volcano plot of significant enriched/depleted somatic TEI windows (1Mbp) between samples in a cluster and samples not in the cluster (Fisher's exact test, p value < 0.05).

e. Oncoplot of top 10 enriched/depleted somatic TEI windows of each cluster (FDR <

1981   0.05). The color of the left heatmap represents the significance level (Fisher's exact

1982   test, log10(FDR)) of enrichment or depletion of a specific cluster in a specific 1Mb

1983   window. The color of the middle heatmap represents the log2(Fold Change) of

1984   enrichment or depletion of a specific cluster in a specific 1Mb window. The color of the

1985   middle heatmap represents the log2(Fold Change) of enrichment or depletion of a

1986   specific cluster in a specific 1Mb window. The color of the right heatmap represents

1987   the fraction of samples harbored somatic TEIs in the specific 1Mb window.

1988

1989

1990

**Figure S15. The proportion of different types of TEIs inserted in MHC II gene family,**

**and differentially expressed genes related to these somatic TEIs.**

a. Pie chart of somatic TEI type distribution. 'solo SINE': TEIs contained solo SINE
   segments. 'complex SINE': TEIs contained multi SINE segments and not contained
   other segments. 'solo LINE': TEIs contained solo LINE segments. 'complex LINE':
   TEIs contained multi LINE segments and not contained other segments. 'SINE+LINE':
   TEIs both contained SINE segments and LINE segments. 'solo LTR': TEIs consist of
   solo LTR segments. 'other': the rest type of TEI.

1999    b. Volcano plot showing the differential gene expression between LUAD with somatic

2000        TEIs and LUAD without somatic TEIs.

2001    c. Box plot of gene expression (log2) in samples with (blue) and without (gray) somatic

2002        TEIs.

2003    d. Detection of somatic TEI events via PCR: PCR analysis was performed in IC12, with

2004        blood DNA used as a control to distinguish somatic TEI events.

2005    e. Sanger sequencing visualization of TEI: The reference genome sequence

2006        incorporates both the flanking region (highlighted in red) and the TEI sequence

2007        (highlighted in blue). The TEI annotation is displayed above the reference genome for

2008        precise localization and identification.

2009

2010

2011

**Figure S16. Bubble plot of pathway and GO enrichment.**

a. Sankey plots of KEGG pathway and corresponding genes are shown on the left, and bubble plots of enrichment results are shown on the right. The x-axis represents the -log10(p-value), y-axis lists the pathways, with the color gradient from purple to yellow

2016       representing the gene ratio. The size of the dots corresponds to the term ratio, with

2017       larger dots representing a greater magnitude of change. The connecting lines

2018       represent relationships or pathways between the genes.

2019   b.  Sankey plots of GO biological process and corresponding genes are shown on the left,

2020       and bubble plots of enrichment results are shown on the right. The x-axis represents

2021       the -log10(p-value), y-axis lists the pathways, with the color gradient from purple to

2022       yellow representing the gene ratio. The size of the dots corresponds to the term ratio,

2023       with larger dots representing a greater magnitude of change. The connecting lines

2024       represent relationships or pathways between the genes.

2025

**Figure S17. DetectingTEI events through PCR in additional fresh clinical samples after CD45+ cell sorting.**

The green dots mark the TEI events that were only amplified in the tumor.

**Figure S18. Detecting TEI events through PCR in tumor cell lines.**

The green dots mark the potential TEI events.

2035

2036

**Figure S19. Detecting TEI events through PCR in tumor cell lines.**

The green dots mark the potential TEI events.

**Figure S20. Statistics of TEI events in tumor cell lines and Flow sytometric analysis.**

a. Statistics of TEI events in tumor cell lines. Each column represents a TEI site, and

each row represents a cell line. The first three tracks correspond to cell lines from

adenocarcinoma, other types, and squamous, respectively. The fourth track

summarizes TEI events from 33 cell lines in the region chr6:32,000,000-33,000,000,

with different colors indicating the categories of these TEI events. The lower enlarged

panel displays a browser plot showing ENCODE-annotated gene transcripts, cCREs

predicted by ENCODE, and epigenomic signals around the somatic TEIs within the

MHC II gene family.

b.  Violin plot showing the expression level of PD-1 and PD-L1 in LUAD samples

harboring somatic TEIs within the HLA-DRB1 genebody and its ±5kb flanking regions

('with'), compare to LUAD samples not harboring somatic TEIs within these regions

('without').

c.  Flow cytometric analyses of CD40L mobilization in CD4+ T cells after 24 hours of in

vitro co-culture with tumor cells.

2056

2057