Possible exposure to unidentified coronaviruses in roe deer (*Capreolus capreolus*) populations suggested by SARS-CoV-2 serological investigation in France

Grégoire Perez*[,a,1,2,3,4] & Lucas D. Lalande*[,a,1,5], Vincent Legros[6], Angéli Kodjo[7], Hélène Verheyden[8,9], Vincent Bourret[8,9], Nicolas Cèbe[8,9], Yannick Chaval[8,9], Paul Revelli[1], Maryline Pellerin[10], Jean-François Lemaître[1], Carole Peroz[1,11], Benjamin Rey[1], François Débias[1], Rebecca Garcia[1], Gilles Bourgoin[1,11], Emmanuelle Gilot-Fromont[1,11]

* Corresponding authors :
Grégoire Perez (gregoire.perez@cirad.fr), Lucas D. Lalande (lalande.luke@gmail.com)

[a] Contributed equally

[1] Université de Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie Evolutive UMR 5558, F-69622 Villeurbanne, France

[2] CIRAD, UMR ASTRE, Conakry, Guinée

[3] ASTRE, CIRAD, INRAE, Université de Montpellier, 34398 Montpellier, France.

[4] CERFIG, Université Gamal Abdel Nasser de Conakry, Conakry, Guinée

[5] Université de Toulouse, Institut National Universitaire Jean-Francois Champollion, BTSB-UR 7417, Place de Verdun, 81000 Albi, France

[6] Université de Lyon, Université Lyon 1, CNRS, Centre International de Recherche en Infectiologie, Inserm U1111 UMR5308, ENS de Lyon, Lyon, France

[7] Université de Lyon, VetAgro Sup, Laboratoire des Leptosires et d'Analyses Vétérinaires, 69280 Marcy L'Etoile, France

[8] Université de Toulouse, INRAE, CEFS, Castanet Tolosan, 31326, France

[9] LTSER ZA PYrénées GARonne, Auzeville-Tolosane, 31320, France

[10] Office Français de la Biodiversité, Direction de la Recherche et de l'Appui Scientifique, Service Conservation et Gestion Durable des Espèces Exploités, 52210 Châteauvillain, France

[11] Université de Lyon, VetAgro Sup, 69280 Marcy l'Etoile, France

Abstract

The risk of viral transmissions from domestic and wild animals to humans is of high concern for human health. Humans can also transmit viral infections back to domestic and wild animals, which can then act as reservoir for the maintenance of viruses, with the risk of epidemic diseases re-emergence. The Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), causing the COVID-19, likely originated from wildlife and has already been evidenced to be transmitted from humans to captive, domestic and wild animals. In particular, white-tailed deer (*Odocoileus virginianus*) show high-prevalence of SARS-CoV-2 following human contamination, and recent data suggest that white-tailed deer could act as an emerging virus reservoir. Here, we investigated whether SARS-CoV-2 could also have emerged in several longitudinally-monitored populations of European roe deer (*Capreolus capreolus*) in direct contact with humans in France. We performed indirect tests (serological ELISAs and seroneutralization) on serum collected pre- and post-emergence of the virus in human populations. We also performed direct tests (nasal swabs followed by RT-PCR) to detect the presence of SARS-CoV-2 RNA in those populations in 2022. We also investigated the exposure and prevalence of the virus in three other cervid species in France. ELISA tests were positive for 2.20 % of the sera tested, both pre- and post-emergence of the virus, but all positive samples were negative with seroneutralization. Direct PCR testing showed no positive results in 2022. Thus, although seroprevalence increased after the COVID-19 emergence (2022) in one out of three roe deer population, our results suggest that SARS-CoV-2 has not emerged in those populations, and that ELISA cross-reaction with another circulating virus, possibly an unidentified bovine coronavirus, is possible.

Keywords: COVID-19, wild ungulates, cervids, spill-over, spill-back, zoonoses

1. Introduction

The potential transmission of viral infections inducing an epidemic from livestock or wildlife to humans is of high concern for human health worldwide, as vertebrates constitute the main source of human emerging diseases (Jones et al., 2008). Conversely, human pathogens can be transmitted to domestic or wild animal species, potentially inducing new epidemics in these species (*e.g.* human coronavirus OC43 outbreak in wild chimpanzees; Patrono et al. 2018). Thus, there is a need to identify pathogens features (*e.g.* mutation rate, transmission route) and ecological contexts (*e.g.* species interactions, habitat use) that promote such events.

Coronaviruses are hosted by many mammal species and can be transmitted from one species to another, including humans (Alluwaimi et al., 2020; Monchatre-Leroy et al., 2017; Patrono et al., 2018). The COVID-19 pandemic, due to the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), most likely originated from wildlife (*e.g.* Lytras et al. 2021, Domingo 2022). Since the beginning of the COVID-19 pandemic, transmission events have been documented from humans back to domestic animals, for example farmed minks (*Neovison vison*; Oreshkova et al. 2020, Oude Munnink et al. 2021), domestic cats (*Felis catus*) and dogs (*Canis lupus familiaris*; Hobbs and Reid 2021, Krafft et al. 2021) or wild felids kept in zoos (McAloose et al., 2020). These first observed transmission events occurred in domestic, captive or intensive farming contexts and could be explained by elevated rates of direct contacts between humans and animals in such specific contexts. However, multiple events of transmission of SARS-CoV-2 from human to free-living white-tailed deer (*Odocoileus virginianus*) have also been documented in North America. These events were followed by transmissions within local white-tailed deer population, resulting in populations displaying high prevalence levels which were up to 40% by seroprevalence, and 82.5% and 70% in retropharyngeal lymph node and nasal swabs tested by RT-PCR,

respectively (Chandler et al., 2021; Feng et al., 2023; Hale et al., 2022; Kuchipudi et al.,

2022). Eventually, three cases of transmission from the newly infected white-tailed deer

populations to humans were detected, suggesting that the white-tailed deer may become an

emerging virus reservoir in North America (Feng et al., 2023). Similar contamination

pathways can be expected in Europe, as several closely-related species of cervids are highly

abundant and can be exposed to viral transmission. For instance, first cases of seropositivity

have been recently observed in fallow deer (*Dama dama*) and red deer (*Cervus elaphus*) in

free-living suburban populations in Spain, suggesting that the host tropism of SARS-CoV-2

may change as new variants emerge (Encinas et al., 2023).

Among European cervid species, the roe deer (*Capreolus capreolus*) is particularly

interesting. Indeed, the species occupies a large variety of habitats, including agricultural

areas with proximity to domestic species and human activities (A. J. Hewison et al., 2001),

making roe deer potentially exposed to human-borne viruses. The species may also display a

gregarious behaviour in winter in open landscapes (Gerard et al., 1992), favouring disease

transmission. Furthermore, the hunting regulations and the absence of large predator during

the last 50 years resulted in high abundance of roe deer all over Europe. Between 1984 and

2005, the number of roe deer in Europe was estimated to increase from 6.2 to 9.5 million and

hunting grew from 1.7 to 2.7 million of hunted individuals yearly (Burbaite & Csanyi, 2009).

In France, the roe deer is widely hunted (*c.* 600,000 individuals officially hunted in 2022, *i.e.*

ten times the number hunted 50 years ago; OFB, Réseau Ongulés sauvages 2022). The rate of

potential contact with humans, one of the potential routes of transmission of the virus from

humans to cervids, and particularly roe deer, is therefore particularly high in France

(Fernández-Bastit et al., 2023).

Benefiting from longitudinal studies of four free-living (including three enclosed) and

one captive populations of roe deer in France, we investigated whether SARS-CoV-2 had

recently emerged in these populations subject to hunting or direct handling as part of capture-mark-recapture programs. Because of annual captures, roe deer from these populations are particularly exposed to human viruses' transmission. Moreover, the continuous follow-up of the same individuals from 2010 to 2022 was an opportunity to detect virus arrival in populations and to explore associated individual factors, namely age and sex. Indeed, older individuals and males are expected to have higher seroprevalence as observed in white-tailed deer, likely driven by sex- and age-specific behaviours during reproductive periods (Caserta et al., 2023; Hale et al., 2022). The virus was searched for by means of either direct PCR testing (on individuals captured in 2022) or indirect serological testing to assess exposure before and after SARS-CoV-2 emergence in human in France in 2020. We further investigated exposure and presence of the virus in other cervid samples collected in 2022 during capture or hunting operations: red deer, fallow deer and sika deer (*Cervus nippon*).

2. Material and methods

2.1. Sampling

Sera sampling is summarized in Figure 1 and Table 1.

2.1.1. Roe deer populations

Roe deer sera were sampled from five populations: Trois-Fontaines (TF), Chizé (CH), Aurignac (AU), Gardouch (GA) and Domain of Praillebard (DP). Among these populations, three are hunted wild populations: TF, CH and AU (Fig. 1, red squares). TF is a 1,360 ha fenced forest in north-eastern France (48°43'N, 4°55'E) hosting an enclosed wild roe deer population; CH is a 2,614 ha fenced Integral Biological Reserve (unmanaged forest) since 2006 located in western France (46°05'N, 0°25'W) hosting also a closed wild roe deer population; AU is an open study area covering about 7,500 ha in south-western France (43°16'N, 0°53'E) including forests and agricultural landscapes with free-ranging roe deer.

Captive roe deer were sampled for their sera in GA (Fig. 1, orange triangle), an

experimental station in southern France (43°22', 1°40'E) hosting captive roe deer in small 0.5

ha enclosures or semi-captive roe deer in a large enclosure composed of 9 ha of forest and 5

ha of meadows.

Roe deer sera were also sampled at the DP (Fig. 1, yellow diamond), a private fenced

area of 148 ha in eastern France (45°57'N, 4°55'E) hosting a small roe deer population

introduced from TF and CH in 2019-2020.

2.1.2. Sera collection

Sera were collected in these populations during 2,134 capture events, involving 1,289

different individuals. Each individual was sampled one to nine times during the study period

(2010-2022), with 874 individuals sampled only once. All samples were collected each year

in winter between December (year *t-1)* and March (year *t*), except at GA where sampling

occurred until May. Sera were immediately stored at -80°C after collection on the field and

during transportation to the laboratory, where they were stored at -20°C until analyses.

2.1.3. Nasal swabs collection

For direct virus detection, nasal swabs were collected in 2022 during the roe deer

capture-mark-recapture programs from TF (n = 49), CH (n = 95) and AU (n = 61). Nasal

swabs were immediately stored at -80°C after collection and during transportation to the

laboratory where they were immediately tested upon arrival.

2.1.4. Additional sampling on other cervid species in other locations

Additionally, we sampled cervids in other locations: the Vercors Vie Sauvage (VVS)

private reserve located in eastern France (44°57'N, 5°14'E; Fig. 1, green star) and three

departments of central France (Allier, Cher and Loire; Fig. 1, blue areas).

We sampled 30 sera from other cervids captured in 2022 in the VVS, including 16

European fallow deer, five sika deer and nine red deer (Table 1). In the three central France

departments, nasal swabs were also collected on hunted roe deer (Allier: n = 3, Cher: n = 29,

and Loire: n = 4), and on red deer in one department (Cher: n = 3) (Fig. 1, blue areas).

We defined two periods, either before COVID-19 emergence in France (pre-COVID-

19 emergence: 2010-2020, n = 1,752, 'pre-Cov' hereafter) or after COVID-19 emergence

(post-COVID-19 emergence: 2021-2022, n = 382, 'post-Cov' hereafter).

Legend

■ Wild populations sampled from 2010 to 2022 included in the serological study
  and tested by RT-PCR in 2022

▲ Captive population sampled from 2019 to 2022 included in the serological study
  and tested by RT-PCR in 2022

◆ Wild population sampled in 2022 included in the serological study only

★ Wild red deer, sika deer, and fallow deer populations sampled in 2022
  included in the serological study only

▢ Departments with roe deer and/or red deer samples tested by RT-PCR in 2022

173
174
175   **Figure 1**. Locations of deer (Cervidae) populations sampled for the sera used in the SARS-
176   CoV-2 serological study and the nasal swabs used in the SARS-CoV-2 RNA RT-PCR screening.

177 **Table 1.** Summary of the sampling scheme and the SARS-CoV-2 serological results for roe deer (*Capreolus capreolus*) and other cervids for pre-
178 (*i.e.* 2010-2020) and post- (*i.e.* 2021-2022) COVID-19 emergence, set in March 2020. Number of positive results and seroprevalence were
179 determined from ELISA assays.

Location (sampling period)	Species	Number of distinct individuals	Number of samples (pre-covid/2021/2022)	Number positive results (pre-covid/2021/2022)	Seroprevalence % (pre-covid/2021/2022)
Trois-Fontaines (2010-2022)	Roe deer	539	942 (832/29/81)	17 (16/1/0)	1.8 (1.9/3.5/0.00)
Chizé (2010-2022)	Roe deer	557	965 (817/29/119)	23 (15/0/8)	2.4 (1.8/0/6.7)
Aurignac (2019-2022)	Roe deer	180	205 (97/49/59)	7 (4/2/1)	3.4 (4.1/2.0/1.7)
Gardouch (2019-2022)	Roe deer	9	18 (6/4/8)	0 (0/0/0)	0.0 (0.0/0.0/0.0)
Domaine of Praillebard (2022)	Roe deer	4	4 (0/0/4)	0 (-/-/0)	0.0 (-/-/0.0)
Vercors Vie Sauvage (2022)	Fallow deer	16	16 (0/0/16)	1 (-/-/1)	6.3 (-/-/6.3)
	Red deer	9	9 (0/0/9)	0 (-/-/0)	0.0 (-/-/0.0)
	Sika deer	5	5 (0/0/5)	1 (-/-/1)	20.0 (-/-/20.0)
Total	**Roe deer**	**1,289**	**2,134 (1,752/111/271)**	**47 (35/3/9)**	**2.2 (2.0/2.7/3.3)** 95%CI: [1.6; 2.9] ([1.4; 2.8]/[0.6; 7.7]/[1.5; 6.2])
	Fallow deer	**16**	**16 (0/0/16)**	**1 (-/-/1)**	**6.3 (-/-/6.3)** 95%CI: [0.2; 30.2] (-/-/[0.2; 30.2])
	Red deer	**9**	**9 (0/0/9)**	**0 (-/-/0)**	**0.0 (-/-/0.0)** 95%CI: [0.0; 33.6] (-/-/[0.0; 33.6])
	Sika deer	**5**	**5 (0/0/5)**	**1 (-/-/1)**	**20.0 (-/-/20.0)** 95%CI: [0.5; 71.6] (-/-/[0.5; 71.6])
	All species	**1,319**	**2,164 (1,752/111/301)**	**49 (35/3/11)**	**2.3 (2.0/2.7/3.7)** 95%CI: [1.7; 3.0] ([1.4; 2.8]/[0.6; 7.7]/[1.8; 6.4])

2.2. Initial ELISA serological screening

Serum samples were first screened using the ID Screen® SARS-CoV-2 Double Antigen Multi-species ELISA kit (ID Vet, France). This kit is designed for the detection of antibodies (IgG, IgM) against the nucleocapsid (N-protein) of SARS-CoV-2, and thus does not allow to discriminate between past and recent infection. The optical density (OD) was measured spectrophotometrically at 450 nm (Safas MP96, Monaco) following the manufacturer's instructions. The seropositivity threshold was S/P% $\geq$ 60, with $S/P = \frac{OD_{sample} - OD_{negative\ control}}{OD_{positive\ control} - OD_{negative\ control}} \times 100$.

2.3. Confirmatory seroneutralization assay

To confirm or refute positive ELISA results, the corresponding samples were further analysed using seroneutralization as described in Pozzetto et al. (2021). Briefly, Murine leukemia virus-based pseudoparticles carrying a green-fluorescent protein (GFP) reporter pseudotyped with SARS-CoV-2 spike protein (SARS-CoV-2pp) were used to measure the neutralizing antibody activity in sera. A sample of *ca.* 10^3 pseudoparticles was incubated with a 1/100 dilution of serum for 1 h at 37 °C before infection of the Vero-E6R cells (ATCC CRL-1586). The percentage of GFP-positive cells was compared to cells infected with SARS-CoV-2pp incubated without serum to set the percentage of neutralization. The positive control used was a rabbit monoclonal neutralizing antibody targeting the receptor-binding domain of the SARS-CoV-2 (2019-nCoV) spike protein (Sino Biological, 40592-R001), with a 100-fold dilution, as used in Fritz et al., 2021 and Pozzetto et al., 2021. As negative control, we used a subset of 30 samples that were negative to ELISA tests, corresponding to the pre-pandemic period (Fritz et al., 2021).

2.4. Direct detection

Nasal swabs (n = 244) were analysed using reverse transcriptase polymerase chain reaction (RT-PCR) technique. In an experimental study on white-tailed deer, RT-PCR on

nasal swabs allowed to detect SARS-CoV-2 viral RNA at least 22 days post-infection, while

infectious virions could only be detected for up to six days post-infection (Martins et al.,

2022). Briefly, automated extraction instruments (KingFisher Flex 96 instrument by Thermo

Fisher Scientific, Waltham, MA, USA, or the Auto Pure 96 instrument by Allsheng

Instruments CO, Hangzhou, China) were used to extract viral RNA from swabs using the

BioExtract Premium Mag (Biosellal, Dardilly, France). Subsequently, SARS-CoV-2 RNA

detection was performed by qRT-PCR with the Bio-T kit TriStar COVID-19 (Biosellal,

France), which targets the envelope gene (E) of Sarbecoviruses (among which SARS-CoV-1,

SARS-CoV-2), and RNA-dependent RNA polymerase (RdRp) in the Orf1ab gene, according

to the manufacturer's instructions. The AriaMx Real-Time PCR Thermal Cycler System

(Agilent Technologies, Santa Clara, CA, USA) was used for amplification. The condition

consisted of 1 cycle of 20 min at 50 °C, 1 min at 95 °C, followed by 40 cycles of 10 s at 95

°C, and 45 s at 60 °C. The results were analysed using the Agilent Aria 1.6 software, in which

a cycle threshold value (Ct-value) < 40 for all target genes was defined as a positive result.

2.5. Analysis of seropositivity in roe deer

We analysed the distribution of ELISA results to better understand the origin of

seropositive signals. First, under the hypothesis of SARS-CoV-2 emergence in roe deer

populations following introduction from humans, a higher seroprevalence was expected in the

post-COVID-19 human emergence period compared to the pre-COVID human emergence

period. To test for this hypothesis, we compared the seroprevalences observed during two

periods: either before COVID-19 emergence in France (pre-COVID-19 emergence: 2010-

2020, n = 1,752, 'pre-Cov' hereafter) or after COVID-19 emergence (post-COVID-19

emergence: 2021-2022, n = 382, 'post-Cov' hereafter), for all populations together, with Chi-

square tests (alpha = 0.05). We used samples of the first period as controls to test for potential

cross-reaction with other coronaviruses circulating in these populations. We included the year

2020 in the pre-emergence period because sampling occurred between December 10[th] 2019

and March 12[th] 2020, before the virus spread across France and the lockdown was enforced

on March 17[th] (Bernard-Stoecklin et al., 2020).

However, because animal manipulation was reduced in 2021 as a consequence of the

implementation of social restrictions during the pandemic, the virus could have emerged in

roe deer populations later than 2021 (*i.e.* in 2022). Thus, considering a possible delayed

exposition of the roe deer population, we ran a second test comparing the 2010-2021 *v.* 2022

periods. We thus defined two different periods: either before a delayed COVID-19 emergence

in France (pre-delayed COVID-19 emergence: 2010-2021, n = 1,863, 'pre-dCov' hereafter) or

after a delayed COVID-19 emergence with relaxed social restrictions (post-delayed COVID-

19 emergence: 2022, n = 271, 'post-dCov' hereafter).

Serological results were further analysed to identify individual and population factors

related to ELISA seropositivity. In this aim, we used 902 samples from TF, 866 from CH, and

202 from AU, corresponding to individuals of known age and sex. A binomial Generalised

Linear Model (GLM) was used to model individual serological status. First, seropositivity

could be cumulative with age, leading to autocorrelation with time in seropositive status.

Thus, the probability that an individual with a positive result is also tested positive a

following year was compared to the probability of an individual being positive a single year

using a binomial test ($P(positive\,twice|positive)$ *v.* $P(positive|tested)$, alpha = 0.05). Then,

several potential explanatory variables were included in the GLM to explain individual

serological result. To test for a contrast in seroprevalence after the COVID-19 emergence,

pre-Cov and post-Cov periods as well as pre-dCov and post-dCov periods were included as

distinct explanatory factors in the GLM but we excluded the possibility to retain both

variables simultaneously. Seroprevalence may also have changed linearly over time for other

reasons (*e.g.* environmental condition variation, forest management), so the temporal trend

was also tested by using the sampling year as a linear integer (2010-2022). Seroprevalence

was also expected to increase with age in the case of a cumulative effect due to antibodies

persistence. This effect was tested adding age as a continuous variable (linear integer, 1-16

years of age). In TF and CH, age is precisely determined by observation of the dentition

eruption of yearlings (A. Hewison et al., 1999) so that only individuals caught during their

first winter are known-aged individuals. For AU, the exact age was not known for all

individuals, instead, age was estimated using five age classes through animal dentition

observations (< 1, 1, 2-3, 4-5 and ≥ 6 years of age). In this case, the youngest possible age

was considered (*e.g.* for an individual of the 4-5 years age class, it was considered as a 4-year-

old individual) so that age was treated as a continuous variable as in TF and CH. As sex is

also known to potentially influence susceptibility to infection (*e.g.* Channappanavar et al.

2017, Hewitt et al. 2024) this factor was also included in the models. One can also expect a

higher increase in seroprevalence in enclosed populations with tight contacts with humans

because of the follow-up research program. Population was thus included in the models in

interaction with the COVID-19 emergence periods. As the effect of age can differ between

sexes as observed in white-tailed deer (Caserta et al., 2023), models included the age:sex

interaction. Because population-specific characteristics can act differently on individual

condition with advancing age, as it is for example the case for immunosenescence patterns

that differ between TF and CH (Cheynel et al., 2017; Lalande et al., 2024), models included

an age:population interaction. Finally, because temporal trends can differ among populations,

the population:year interaction were also included in the models.

Model selection was then performed to select the best model based on the Akaike

Information Criteria corrected for small sample sizes (AICc): models with ΔAICc < 2 were

kept and the significance of the variables included in those retained models was evaluated

using type III ANOVAs. To evaluate the significance of the difference between groups for

significant categorical variables, *post hoc* between groups paired comparisons were conducted using binomial tests. All statistics were done with R version 4.2.2 (R Core Team, 2023) in Rstudio 2023.03.0 (Posit Team, 2023), and with the *car* 3.1-2 (Fox & Weisberg, 2019), *MuMIn* 1.47.5 (Bartoń, 2023), *readr* 2.1.4 (Wickham et al., 2023) and *visreg* 2.7.0 (Breheny & Burchett, 2020) packages.

3. Results

3.1. Serological results

A total of 47 out of 2,134 ELISA tests (2.2% [95%CI: 1.6; 2.9]) performed on roe deer sera had positive results for anti-SARS-CoV-2 antibodies against the N-protein (Table 1). It included samples collected during both the pre-Cov (35 positives: 2.0%) and post-Cov (3 positives: 2.7% and 9 positives: 3.3%, in 2021 and 2022, respectively) periods (Table 1). Among the 47 positive results, 17 out of 942 (1.8%) were found in TF (pre-Cov: 16 out of 832 (1.9%); 2021: 1 out 29 (3.5%); 2022: 0 out of 81), 23 out of 965 (2.4%) in CH (pre-Cov: 15 out of 817 (1.8%); 2021: 0 out of 29, 2022: 8 out of 119 (6.7%)) and 7 out of 205 (3.4%) in AU (pre-Cov: 4 out of 97 (4.1%); 2021: 2 out of 49 (2.0%); 2022: 2 out of 59 (1.7%)).

The S/P% values of positive results ranged from 60.7 to 824.4% and no evident bimodal distribution was observed, as expected when a proportion of the population have been exposed to a pathogen and displays a high antibody concentration. Although seroprevalence was higher in the post-COVID-19 emergence period (delayed or not) than before SARS-CoV-2 emergence, this difference, when considering all roe deer populations together, was not statistically significant (pre-Cov *v.* post-Cov Chi² test: $\chi^2 = 1.41$, df = 1, $p = 0.24$; pre-dCov *v.* post-dCov Chi² test: $\chi^2 = 1.26$, df = 1, $p = 0.26$). Additionally, one roe deer was tested twice during the same year (with a three-week interval). Both samples from this individual were positive and observed S/P% declined from 328% to 120% between the two samples. Only the first observation was kept for the subsequent analyses.

Among the sera from other cervids inhabiting the VVS reserve, ELISA tests

performed on one fallow deer out of 16 (6.3% [0.2; 30.2]) and one sika deer out of five

(20.0% [0.5; 71.6]) yielded seropositive results (Table 1).

SARS-CoV-2 seroneutralization assays were performed on nine positive and 15

negative roe deer samples of the post-COVID-19 emergence period, and on four positive and

15 negative roe deer samples of the pre-COVID-19 emergence period. All seroneutralization

assays, including the 13 carried out on ELISA-positive samples, yielded negative results.

3.2. Direct virus detection

RT-PCR carried out on nasal swabs collected during winter 2022 on 205 captured roe

deer (of which 10 were positive with the ELISA assay) and 36 hunted roe deer and three

hunted red deer, did not reveal any positive result.

3.3. Analysis of seropositivity in roe deer

The 47 positive sera originated from 46 individuals from CH, TF, and AU (see Table

1), including 28 individuals sampled and tested only once, seven tested twice, six tested

thrice, four tested four times and one tested five times. During the follow up of the 19 positive

individuals tested multiple times, both appearance and disappearance of antibodies were

evidenced, as 13 seronegative individuals were found seropositive in the following years (six

on the next year, seven two years or more later), while 11 negative results were observed on

samples that were previously positive (four on the next year, seven two years or more later).

Only one individual was tested positive twice during two different years (2016 and 2018) but

no sample was available in-between (*i.e.* 2017). 20 positive results were from individuals in

their first year. The probability that an individual was positive two different years was not

different from expected by chance based on the observed prevalence (binomial test: $p = 1$).

So, seropositivity events (47 sera in 46 individuals) were considered as independent in the

following analyses.

Individual seropositivity was analysed in 1,970 samples obtained on 1,141 individuals

for which population, sampling year, age and sex were known in AU, CH and TF populations.

The models retained according to the Akaike Information criterion (AICc)-based selection

procedure (ΔAICc $\leq$ 2) are presented in Table 2. The most parsimonious model of

seroprevalence included the positive linear effect of age ($\beta \pm SE = 0.31 \pm 0.24$), the delayed

COVID-19 emergence variable (dCov: 0.80 ± 1.11) and the effect of population (AU

[intercept]: -5.11 ± 1.39, CH: 3.33 ± 1.46, TF: -13.48 ± 748.18). We also found a significant

interaction between the delayed emergence variable and population (AU [intercept]: $-5.11 \pm$

1.39, CH: -2.33 ± 1.21, TF: 13.84 ± 748.18), showing a higher seropositivity in CH in 2022

compared to the 2010-2021 period (Figure 2). We evidenced a significant interaction between

age and population (AU [intercept]: -5.11 ± 1.39, CH: -0.60 ± 0.27, TF: -0.30 ± 0.26),

showing a decrease in seropositivity with age in CH only (Figure 3). Note that the variable

comparing pre-Cov and post-Cov periods was never retained in the higher-ranked models

(Table 2). In contrast, those models all included the pre-dCov *v.* post-dCov periods variable.

In these models, the effect of the population and its interaction with the delayed emergence

variable and with age were also always retained.

Finally, we conducted *post hoc* between groups paired comparisons on the retained

model interactions. We found that the seroprevalence was higher during the post-dCov (*i.e.*

2022) in CH (pre-dCov: 1.60% [95%CI: 0.83-2.79%], post-dCov: 6.78%, [2.97-12.92%], *p <*

0.01; Fig. 2), and decreased with age at CH, but not in TF and AU. Seroprevalence decreased

from 4.50% [95%CI: 2.48-7.44%] during the first year of life to 0.68% [95%CI: 0.08-2.43%]

in individuals aged 5 years old and more in average for both periods (*p < 0.01*; Fig. 3) but was

more marked in post-dCov.

Table 2. Summary of seroprevalence models retained by the AICc-based selection procedure. Only models within 2 ΔAICc are presented.

Int	Age	Cov	dCov	Pop	Sex	Year	Age:Pop	Age:Sex	Cov:Pop	dCov:Pop	R²	df	logLik	AICc	ΔAICc	weight
-5.11	**0.31**		**+**	**+**			**+**			**+**	**0.01**	**9**	**-191.62**	**401.33**	**0.00**	**0.47**
-132.89	0.31		+	+		0.06	+			+	0.01	10	-191.11	402.32	0.99	0.29
-5.28	0.32		+	+	+		+			+	0.01	10	-191.29	402.69	1.36	0.24

355 Int: intercept; Cov: COVID-19 emergence period (2010-2020 *v.* 2021-2022); dCov: delayed COVID-19 emergence period (2010-2021 *v.* 2022);
356 Pop: population (Aurignac, Chizé, Trois-Fontaines); df: degrees of freedom; logLik: log-likelihood; AICc: Akaike Information Criterion corrected;
357 ΔAICc: difference of AICc from the best model; weight: relative model weight. See text for further details.

358
Figure 2. SARS-CoV-2 seroprevalence of roe deer (*Capreolus capreolus*) population and the
delayed COVID-19 emergence periods. The plot displays fitted seroprevalence with 95%
confidence interval for Aurignac (AU), Chizé (CH) and Trois-Fontaines (TF) populations for the
pre- (2010-2021; "Pre"; upper panel) and post-delayed COVID-19 (2022; "Post"; lower panel)
emergence periods; "a" indicates a significant difference ($p < 0.05$) in *post hoc* between
groups paired comparisons.

Figure 3. SARS-CoV-2 seroprevalence of roe deer (*Capreolus capreolus*) in the population of
Chizé (CH) as a function of age and delayed COVID-19 emergence periods. The plot displays
fitted seroprevalence in CH during the pre- (2010-2021, 'pre', blue dots) and post-delayed
COVID-19 (2022, 'post', orange triangles) emergence periods (see text for more details) with
95% confidence intervals.

4. Discussion

This study is the first long-term individual monitoring of serological status of wild-living cervids for SARS-CoV-2. Our survey intended to detect, at the individual or population levels, an increase in seropositivity or seroprevalence following the emergence of SARS-CoV-2 in France.

Interpreting serological results in terms of viral infection is not straightforward. SARS-CoV-2 ELISA tests indicated an overall seroprevalence of 2.20% in roe deer. We also observed positive results in other cervids from VVS despite a low sample size (one out of 16 fallow deer and one out of five sika deer). However, all sera submitted to the seroneutralization assay, and all swabs tested for direct detection of the virus by RT-PCR in 2022 gave negative results. This indicates that sarbecoviruses, among which SARS-CoV-2, are unlikely to circulate in the studied populations of roe deer.

The absence of confirmation of SARS-CoV-2 antibodies by seroneutralization assay is consistent with other studies conducted in Germany, Austria and the UK on roe deer, red deer, fallow deer, sika deer, Reeves's muntjac (*Muntiacus reevesi*), and Chinese water deer (*Hydropotes inermis*) (Encinas et al., 2023; Holding et al., 2022; Moreira-Soto et al., 2022; Wernike, Fischer, et al., 2022). Negative seroneutralization assays despite positive ELISA tests have also been reported in wild boars (*Sus scrofa*), red foxes (*Vulpes vulpes*) and jackals (*Canis aureus moreoticus*) (Jemeršić et al., 2021; Wernike, Fischer, et al., 2022). It can be explained by an inadequate specificity of the ELISA test allowing cross-reaction with antibodies directed to other closely-related viruses (Krüttgen et al., 2021). Also, while ELISA used in this study detect antibodies targeting SARS-CoV-2 nucleoprotein, the seroneutralization assay detects neutralizing antibodies targeting spike glycoprotein. Previous studies showed a poor correlation between antibodies targeting nucleoproteins and neutralizing antibodies in SARS-CoV-2-infected patients (Legros et al., 2021). This could

also explain why a serum can score positive in a nucleoprotein-based ELISA assay, but

negative in a seroneutralization assay. Such results underline the importance of not relying on

ELISA tests alone before drawing any conclusions.

Nonetheless, these positive results suggest that the ELISA test used might cross-react

with one or several other coronaviruses circulating in the studied roe deer populations.

Viruses close to bovine coronaviruses (BCoVs) have already been found in red deer and sika

deer (Amer, 2018), and are potential candidates that could have been detected by our ELISA

analyses. Positive serological results were observed in all populations of roe deer, except DP

and GA for which only four and 18 sera were tested, respectively. If a virus of bovine origin

is involved, one would expect a lower seroprevalence in the two isolated roe deer populations

of CH and TF compared to the free-roaming roe deer population of AU exposed to cattle. Our

results do not show such pattern and in principle would not support the BCoVs hypothesis.

The circulation of phylogenetically close coronaviruses transmitted by humans (*e.g.*

HCoV-OC43; Vijgen et al. 2006) is also possible. In case of a human transmitted coronavirus,

a higher seroprevalence in captive roe deer in GA would be expected, but no positive

individual was observed there (acknowledging, however, a small sample size). Thus, our

results do not support the human coronavirus hypothesis either.

Other possibilities would include: (*i*) long-lasting BCoVs infection maintenance in the

isolated populations since the enclosure of the areas, (*ii*) repeated introduction of BCoVs by

infected animals entering the enclosure by some gaps in the fences, which is possible for roe

deer or another susceptible species like wild boar, or (*iii*) exposure of cervid populations to

coronaviruses from other wildlife species present in the studied areas. Indeed, we cannot

exclude the eventuality that coronaviruses are transmitted to roe deer by bats, rodents, rabbits

or hedgehogs (*e.g.* alpha- and beta-coronaviruses; Monchatre-Leroy et al. 2017, Arnaout et al.

2023), or that another, still unknown, coronavirus is circulating in cervid populations, as

suggested elsewhere (Vijgen et al., 2006; Wernike, Fischer, et al., 2022). The seropositivity of

a fallow deer and a sika deer, which were not in such close contact with human or cattle than

studied roe deer, supports the idea of an exposure of wild cervid populations to at least one

cross-reacting coronavirus.

However, as the virus(es) responsible of the observed seroprevalence is unknown, one

cannot exclude the possibility that the different populations are exposed to distinct viruses

(*e.g.* BCoVs in AU, other coronaviruses in CH and TF). More investigations are required

before drawing any firm conclusion, for instance by using more specific tests to confirm or

infirm the presence of BCoVs antibodies in ELISA positive roe deer sera (Ulrich et al., 2020),

or conduct High Throughput Sequencing analyses on cervid samples to identify candidate

viruses (Fitzpatrick et al., 2021).

When studying more closely the spatio-temporal and individual factors of

seropositivity in the three roe deer populations with a sufficient sample size (AU, CH and

TF), we detected a significant statistical interaction between the pre- and post-COVID-19

delayed emergence periods and populations. Although these results must be interpreted

cautiously because of the low explained variance by the models, seroprevalence increased in

CH from about 1.60% to 6.78% in 2022 while it fell to 0.00% in TF and did not change

substantially in AU. We found no evidence that seroprevalence increased with age, and even

found that seroprevalence decreased with age in CH. Furthermore, among 11 positive

individuals that were tested the following years, only one was found seropositive twice. Other

individuals showing seropositive status were tested negative during the following years,

including four individuals tested only one year later. Together, these results are consistent

with a short persistence of the antibodies (less than one year), but for at least several weeks

(>3). Although a rapid decline of humoral response has been described, antibodies after

natural SARS-CoV-2 infection in humans can persist for more than a year (Ghasemi et al.,

2022; Sarjomaa et al., 2022). In white-tailed deer, it should last at least 13 months (Hamer et

al., 2022), but persistence in other species is not well known. Thus, according to available

data in white-tailed deer, as persistence time is expected to be longer and juveniles are not

expected to have a higher seroprevalence, our observations are not consistent with a SARS-

CoV-2 emergence in the studied roe deer populations (Hamer et al., 2022; Hewitt et al.,

2024). Data from coming years may clarify these temporal trends in the different populations.

Nonetheless, without formally identifying the responsible virus(es), observed patterns are

difficult to explain.

Transmission of the SARS-CoV-2 virus from human to roe deer is thus unlikely to

have happened to date in the studied populations, contrary to what has been observed among

white-tailed deer populations in North America (Chandler et al., 2021; Hale et al., 2022;

Kuchipudi et al., 2022). Our results confirm the results of previous studies evidencing an

absence of contact between the most common European cervids and the SARS-CoV-2 virus

(Holding et al., 2022; Moreira-Soto et al., 2022; Wernike, Fischer, et al., 2022), except in

densely populated areas (Encinas et al., 2023). Indeed, roe deer is not living in peri-urban

areas as frequently as white-tailed deer, and thus contact rate between roe deer and human is

expected to be lower than for white-tailed deer. Roe deer is also not as close to white-tailed

deer relative to their ACE2 (which interacts with the SARS-CoV-2 spike protein) and likely

less susceptible to SARS-CoV-2 infection (Damas et al., 2020). Despite the low probability of

roe deer, or other European cervids, to serve as a reservoir host for SARS-CoV-2, the

circulation of one or several other coronaviruses, maybe BCoVs, in roe deer populations

might raise the question of a possible virus cross-recombination leading to a new virus

variant. Further studies should be carried out to identify coronaviruses circulating in wild

cervids and domestic ruminants, and to continue testing roe deer populations to enable track

temporal trends. Maintaining serological surveillance and implementing efficient virological

surveillance to detect viruses in wildlife is necessary to evaluate the possibility of

transmission between human and other domestic or wild animals (Wernike, Böttcher, et al.,

2022).

Conflict of interest disclosure

478 The authors declare that they have no conflict of interests relating to the content of this article.
479

Data and code availability

481 All data and codes used for the present analyses have been deposited on the CIRAD dataverse
482 and are publicly available: https://doi.org/10.18167/DVN1/AR6UDN (Perez & Lalande,
483 2025).

Fundings

486 This project was funded by the *Université de Lyon* (Project RESPOND and Doctoral grant),
487 *VetAgro Sup* and the *Office Français de la Biodiversité* (OFB, project CNV-REC-2019-08).
488 Funders were not involved in the study design, the collection, analysis and interpretation of
489 data, the writing of the report and the decision to submit the article for publication.

Ethics and authorisations

The research presented in this article was done according to all institutional and/or national
guidelines. The protocol of capture and blood sampling of roe deer is under the authority of
the *Office Français de la Biodiversité* (OFB) and was approved by the Director of Food,
Agriculture and Forest (Prefectoral order 2009-14 from Paris). All procedures were approved
by the Ethical Committee of Lyon 1 University (project DR2014-09, June 5th, 2014). The
Trois-Fontaines population is part of the long-term Studies in Ecology and Evolution (SEE-
Life) program of the CNRS. The data from Aurignac population came from a long-term study
approved by ethical committee n°115 and by French Ministry of Higher Education and
Research (Apafis#39320).
501

Authors contributions

Study conceptualisation: GB and EGF
Methodological design: GP, GB and EGF
Field sampling: GP, LL, HV, VB, NC, YC, PR, MP, JFL, CP, BR, FD, RG, GB and EGF
Immunological assays: GP, LL and BR
Seroneutralization assays: VL
RT-PCR screening: AK

Study coordination and supervision: GB and EGF

Further supervision: MP and JFL

Funding: JFL, GB and EGF

Statistical analyses: GP

Writing – original draft: GP and LL

Writing – first editing: GB and EGF

Writing – review and editing: GP, LL, VL, AK, HV, VB, JFL, CP, BR, GB and EGF

References

Alluwaimi, A. M., Alshubaith, I. H., Al-Ali, A. M., & Abohelaika, S. (2020). The coronaviruses of animals
and birds⬚: Their zoonosis, vaccines, and models for SARS-CoV and SARS-CoV-2. *Frontiers in*
Veterinary Science, 7, 582287. https://doi.org/10.3389/fvets.2020.582287

Amer, H. M. (2018). Bovine-like coronaviruses in domestic and wild ruminants. *Animal Health*
Research Reviews, 19(2), 113⬚124. https://doi.org/10.1017/S1466252318000117

Arnaout, Y., Picard-Meyer, E., Robardet, E., Cappelle, J., Cliquet, F., Touzalin, F., Jimenez, G., &
Djelouadji, Z. (2023). Assessment of virus and *Leptospira* carriage in bats in France. *PLOS*
ONE, 18(10), e0292840. https://doi.org/10.1371/journal.pone.0292840

Bartoń, K. (2023). *MuMIn⬚: Multi-Model Inference* (Version 1.47.5) [Logiciel].
https://cran.rstudio.com/web/packages/MuMIn/index.html

Bernard-Stoecklin, S., Rolland, P., Silue, Y., Mailles, A., Campese, C., Simondon, A., Mechain, M.,
Meurice, L., Nguyen, M., Bassi, C., Yamani, E., Behillil, S., Ismael, S., Nguyen, D., Malvy, D.,
Lescure, F. X., Georges, S., Lazarus, C., Tabaï, A., … Investigation team. (2020). First cases of
coronavirus disease 2019 (COVID-19) in France⬚: Surveillance, investigations and control
measures, January 2020. *Euro Surveillance: Bulletin Europeen Sur Les Maladies Transmissibles*
(European Communicable Disease Bulletin), 25(6), 2000094. https://doi.org/10.2807/1560-
7917.ES.2020.25.6.2000094

Breheny, P., & Burchett, W. (2020). *visreg⬚: Visualization of regression models* (Version 2.7.0)
[Logiciel]. https://CRAN.R-project.org/package=visreg

Burbaite, L., & Csanyi, S. (2009). Roe deer population and harvest changes in Europe. *Estonian*
Journal of Ecology, 58(3), 169⬚181.

Caserta, L. C., Martins, M., Butt, S. L., Hollingshead, N. A., Covaleda, L. M., Ahmed, S., Everts, M. R. R.,
Schuler, K. L., & Diel, D. G. (2023). White-tailed deer (*Odocoileus virginianus*) may serve as a
wildlife reservoir for nearly extinct SARS-CoV-2 variants of concern. *Proceedings of the*
National Academy of Sciences, 120(6), e2215067120.
https://doi.org/10.1073/pnas.2215067120

Chandler, J. C., Bevins, S. N., Ellis, J. W., Linder, T. J., Tell, R. M., Jenkins-Moore, M., Root, J. J., Lenoch,
J. B., Robbe-Austerman, S., DeLiberto, T. J., Gidlewski, T., Kim Torchetti, M., & Shriner, S. A.
(2021). SARS-CoV-2 exposure in wild white-tailed deer (*Odocoileus virginianus*). *Proceedings*
of the National Academy of Sciences, 118(47), e2114828118.
https://doi.org/10.1073/pnas.2114828118

Channappanavar, R., Fett, C., Mack, M., Ten Eyck, P. P., Meyerholz, D. K., & Perlman, S. (2017). Sex-
based differences in susceptibility to SARS-CoV infection. *Journal of Immunology*, *198*(10),
4046⬛4053. https://doi.org/10.4049/jimmunol.1601896
Cheynel, L., Lemaître, J.-F., Gaillard, J.-M., Rey, B., Bourgoin, G., Ferté, H., Jégo, M., Débias, F.,
Pellerin, M., Jacob, L., & Gilot-Fromont, E. (2017). Immunosenescence patterns differ
between populations but not between sexes in a long-lived mammal. *Scientific Reports*, *7*(1),
Article 1. https://doi.org/10.1038/s41598-017-13686-5
Damas, J., Hughes, G. M., Keough, K. C., Painter, C. A., Persky, N. S., Corbo, M., Hiller, M., Koepfli, K.-
P., Pfenning, A. R., Zhao, H., Genereux, D. P., Swofford, R., Pollard, K. S., Ryder, O. A., Nweeia,
M. T., Lindblad-Toh, K., Teeling, E. C., Karlsson, E. K., & Lewin, H. A. (2020). Broad host range
of SARS-CoV-2 predicted by comparative and structural analysis of ACE2 in vertebrates.
Proceedings of the National Academy of Sciences, *117*(36), 22311⬛22322.
https://doi.org/10.1073/pnas.2010146117
Delaune, T., Ouattara, M. S., Ballot, R., Sausse, C., Felix, I., Maupas, F., Chen, M., Morison, M.,
Makowski, D., & Barbu, C. (2021). Landscape drivers of pests and pathogens abundance in
arable crops. *Ecography*, *44*(10), 1429⬛1442. https://doi.org/10.1111/ecog.05433
Domingo, J. L. (2022). An updated review of the scientific literature on the origin of SARS-CoV-2.
Environmental Research, *215*, 114131. https://doi.org/10.1016/j.envres.2022.114131
Encinas, P., Escalera, A., Aydillo, T., Iglesias, I., Nelson, M. I., García-Sastre, A., & del Real, G. (2023).
SARS-CoV-2 neutralizing antibodies in free-ranging fallow deer (*Dama dama*) and red deer
(*Cervus elaphus*) in suburban and rural areas in Spain. *Transboundary and Emerging Diseases*,
2023, e3324790. https://doi.org/10.1155/2023/3324790
Feng, A., Bevins, S., Chandler, J., DeLiberto, T. J., Ghai, R., Lantz, K., Lenoch, J., Retchless, A., Shriner,
S., Tang, C. Y., Tong, S. S., Torchetti, M., Uehara, A., & Wan, X.-F. (2023). Transmission of
SARS-CoV-2 in free-ranging white-tailed deer in the United States. *Nature Communications*,
14(1), Article 1. https://doi.org/10.1038/s41467-023-39782-x
Fernández-Bastit, L., Vergara-Alert, J., & Segalés, J. (2023). Transmission of severe acute respiratory
syndrome coronavirus 2 from humans to animals⬛: Is there a risk of novel reservoirs? *Current*
Opinion in Virology, *63*, 101365. https://doi.org/10.1016/j.coviro.2023.101365
Fischer, S., Weiland, E., & Frölich, K. (1998). Characterization of a bovine viral diarrhea virus isolated
from roe deer in Germany. *Journal of Wildlife Diseases*, *34*(1), 47⬛55.
https://doi.org/10.7589/0090-3558-34.1.47
Fitzpatrick, A. H., Rupnik, A., O'Shea, H., Crispie, F., Keaveney, S., & Cotter, P. (2021). High
Throughput Sequencing for the detection and characterization of RNA viruses. *Frontiers in*
Microbiology, *12*. https://doi.org/10.3389/fmicb.2021.621719
Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression* (3rd éd.). Sage.
https://socialsciences.mcmaster.ca/jfox/Books/Companion/
Fritz, M., Rosolen, B., Krafft, E., Becquart, P., Elguero, E., Vratskikh, O., Denolly, S., Boson, B.,
Vanhomwegen, J., Gouilh, M. A., Kodjo, A., Chirouze, C., Rosolen, S. G., Legros, V., & Leroy, E.
M. (2021). High prevalence of SARS-CoV-2 antibodies in pets from COVID-19+ households.
One Health (Amsterdam, Netherlands), *11*, 100192.
https://doi.org/10.1016/j.onehlt.2020.100192
Frölich, K. (1995). Bovine virus diarrhea and mucosal disease in free-ranging and captive deer
(Cervidae) in Germany. *Journal of Wildlife Diseases*, *31*(2), 247⬛250.
https://doi.org/10.7589/0090-3558-31.2.247

Gerard, J.-F., Poulle, M.-L., Bideau, E., Maublanc, M.-L., & Vincent, J.-P. (1992). Influence of season
and habitat openness on European roe deer (*Capreolus capreolus* L.) group size. *Ungulates*,
91, 279–281.

Ghasemi, D., Araeynejad, F., Maghsoud, O., Gerami, N., Keihan, A. H., Rezaie, E., Mehdizadeh, S.,
Hosseinzadeh, R., Mohammadi, R., Bahardoust, M., & Heiat, M. (2022). The Trend of IgG and
IgM antibodies during 6-month period after the disease episode in COVID-19 patients.
Iranian Journal of Science and Technology. Transaction A, Science, *46*(6), 1555–1562.
https://doi.org/10.1007/s40995-022-01382-7

Hale, V. L., Dennis, P. M., McBride, D. S., Nolting, J. M., Madden, C., Huey, D., Ehrlich, M., Grieser, J.,
Winston, J., Lombardi, D., Gibson, S., Saif, L., Killian, M. L., Lantz, K., Tell, R. M., Torchetti, M.,
Robbe-Austerman, S., Nelson, M. I., Faith, S. A., & Bowman, A. S. (2022). SARS-CoV-2
infection in free-ranging white-tailed deer. *Nature*, *602*(7897), Article 7897.
https://doi.org/10.1038/s41586-021-04353-x

Hamer, S. A., Nunez, C., Roundy, C. M., Tang, W., Thomas, L., Richison, J., Benn, J. S., Auckland, L. D.,
Hensley, T., Cook, W. E., Pauvolid-Corrêa, A., & Hamer, G. L. (2022). Persistence of SARS-CoV-
2 neutralizing antibodies longer than 13 months in naturally infected, captive white-tailed
deer (*Odocoileus virginianus*), Texas. *Emerging Microbes & Infections*, *11*(1), 2112–2115.
https://doi.org/10.1080/22221751.2022.2112913

Hewison, A. J., Vincent, J. P., Joachim, J., Angibault, J. M., Cargnelutti, B., & Cibien, C. (2001). The
effects of woodland fragmentation and human activity on roe deer distribution in agricultural
landscapes. *Canadian Journal of Zoology*, *79*(4), 679–689. https://doi.org/10.1139/z01-032

Hewison, A., Vincent, J. P., Angibault, J. M., Delorme, D., Laere, G. V., & Gaillard, J. M. (1999). Tests of
estimation of age from tooth wear on roe deer of known age : Variation within and among
populations. *Canadian Journal of Zoology*, *77*(1), 58–67. https://doi.org/10.1139/z98-183

Hewitt, J., Wilson-Henjum, G., Collins, D. T., Linder, T. J., Lenoch, J. B., Heale, J. D., Quintanal, C. A.,
Pleszewski, R., McBride, D. S., Bowman, A. S., Chandler, J. C., Shriner, S. A., Bevins, S. N.,
Kohler, D. J., Chipman, R. B., Gosser, A. L., Bergman, D. L., DeLiberto, T. J., & Pepin, K. M.
(2024). Landscape-scale epidemiological dynamics of SARS-CoV-2 in white-tailed deer.
Transboundary and Emerging Diseases, *2024*(1), 7589509.
https://doi.org/10.1155/2024/7589509

Hobbs, E. C., & Reid, T. J. (2021). Animals and SARS-CoV-2 : Species susceptibility and viral
transmission in experimental and natural conditions, and the potential implications for
community transmission. *Transboundary and Emerging Diseases*, *68*(4), 1850–1867.
https://doi.org/10.1111/tbed.13885

Holding, M., Otter, A. D., Dowall, S., Takumi, K., Hicks, B., Coleman, T., Hemingway, G., Royds, M.,
Findlay-Wilson, S., Curran-French, M., Vipond, R., Sprong, H., & Hewson, R. (2022). Screening
of wild deer populations for exposure to SARS-CoV-2 in the United Kingdom, 2020–2021.
Transboundary and Emerging Diseases, *69*(5), e3244–e3249.
https://doi.org/10.1111/tbed.14534

Jemeršić, L., Lojkić, I., Krešić, N., Keros, T., Zelenika, T. A., Jurinović, L., Skok, D., Bata, I., Boras, J.,
Habrun, B., & Brnić, D. (2021). Investigating the presence of SARS-CoV-2 in free-living and
captive animals. *Pathogens*, *10*(6), Article 6. https://doi.org/10.3390/pathogens10060635

Jones, K. E., Patel, N. G., Levy, M. A., Storeygard, A., Balk, D., Gittleman, J. L., & Daszak, P. (2008).
Global trends in emerging infectious diseases. *Nature*, *451*(7181), 990–993.
https://doi.org/10.1038/nature06536

Krafft, E., Denolly, S., Boson, B., Angelloz-Pessey, S., Levaltier, S., Nesi, N., Corbet, S., Leterrier, B.,
Fritz, M., Leroy, E. M., Gouilh, M. A., Cosset, F.-L., Kodjo, A., & Legros, V. (2021). Report of
one-year prospective surveillance of SARS-CoV-2 in dogs and cats in France with various
exposure risks : Confirmation of a low prevalence of shedding, detection and complete
sequencing of an alpha variant in a cat. *Viruses*, *13*(9), 1759.
https://doi.org/10.3390/v13091759

Krüttgen, A., Cornelissen, C. G., Dreher, M., Hornef, M. W., Imöhl, M., & Kleines, M. (2021).
Determination of SARS-CoV-2 antibodies with assays from Diasorin, Roche and IDvet. *Journal*
of Virological Methods, *287*, 113978. https://doi.org/10.1016/j.jviromet.2020.113978

Kuchipudi, S. V., Surendran-Nair, M., Ruden, R. M., Yon, M., Nissly, R. H., Vandegrift, K. J., Nelli, R. K.,
Li, L., Jayarao, B. M., Maranas, C. D., Levine, N., Willgert, K., Conlan, A. J. K., Olsen, R. J., Davis,
J. J., Musser, J. M., Hudson, P. J., & Kapur, V. (2022). Multiple spillovers from humans and
onward transmission of SARS-CoV-2 in white-tailed deer. *Proceedings of the National*
Academy of Sciences, *119*(6), e2121644119. https://doi.org/10.1073/pnas.2121644119

Lalande, L. D., Bourgoin, G., Carbillet, J., Cheynel, L., Debias, F., Ferté, H., Gaillard, J.-M., Garcia, R.,
Lemaître, J.-F., Palme, R., Pellerin, M., Peroz, C., Rey, B., Vuarin, P., & Gilot-Fromont, E.
(2024). Early-life glucocorticoids accelerate lymphocyte count senescence in roe deer.
General and Comparative Endocrinology, *357*, 114595.
https://doi.org/10.1016/j.ygcen.2024.114595

Legros, V., Denolly, S., Vogrig, M., Boson, B., Siret, E., Rigaill, J., Pillet, S., Grattard, F., Gonzalo, S.,
Verhoeven, P., Allatif, O., Berthelot, P., Pélissier, C., Thiery, G., Botelho-Nevers, E., Millet, G.,
Morel, J., Paul, S., Walzer, T., … Pozzetto, B. (2021). A longitudinal study of SARS-CoV-2-
infected patients reveals a high correlation between neutralizing antibodies and COVID-19
severity. *Cellular & Molecular Immunology*, *18*(2), 318 327. https://doi.org/10.1038/s41423-
020-00588-2

Lytras, S., Xia, W., Hughes, J., Jiang, X., & Robertson, D. L. (2021). The animal origin of SARS-CoV-2.
Science, *373*(6558), 968 970. https://doi.org/10.1126/science.abh0117

Martins, M., Boggiatto, P. M., Buckley, A., Cassmann, E. D., Falkenberg, S., Caserta, L. C., Fernandes,
M. H. V., Kanipe, C., Lager, K., Palmer, M. V., & Diel, D. G. (2022). From Deer-to-Deer : SARS-
CoV-2 is efficiently transmitted and presents broad tissue tropism and replication sites in
white-tailed deer. *PLOS Pathogens*, *18*(3), e1010197.
https://doi.org/10.1371/journal.ppat.1010197

McAloose, D., Laverack, M., Wang, L., Killian, M. L., Caserta, L. C., Yuan, F., Mitchell, P. K., Queen, K.,
Mauldin, M. R., Cronk, B. D., Bartlett, S. L., Sykes, J. M., Zec, S., Stokol, T., Ingerman, K.,
Delaney, M. A., Fredrickson, R., Ivančić, M., Jenkins-Moore, M., … Diel, D. G. (2020). From
people to *Panthera* : Natural SARS-CoV-2 infection in tigers and lions at the Bronx Zoo. *mBio*,
11(5), 10.1128/mbio.02220-20. https://doi.org/10.1128/mbio.02220-20

Monchatre-Leroy, E., Boué, F., Boucher, J.-M., Renault, C., Moutou, F., Ar Gouilh, M., & Umhang, G.
(2017). Identification of alpha and beta coronavirus in wildlife species in France : Bats,
rodents, rabbits, and hedgehogs. *Viruses*, *9*(12), Article 12.
https://doi.org/10.3390/v9120364

Moreira-Soto, A., Walzer, C., Czirják, G. Á., Richter, M. H., Marino, S. F., Posautz, A., De Yebra Rodo,
P., McEwen, G. K., Drexler, J. F., & Greenwood, A. D. (2022). Serological evidence that SARS-
CoV-2 has not emerged in deer in Germany or Austria during the COVID-19 pandemic.
Microorganisms, *10*(4), 748. https://doi.org/10.3390/microorganisms10040748

OFB, Réseau Ongulés sauvages. (2022). *Prélèvements ongulés sauvages. Saison 2022-2023*. Office
Français de la Biodiversité.
https://www.ofb.gouv.fr/sites/default/files/Fichiers/Lettre%20information/ficheTCGG_2022
_2023.pdf
Oreshkova, N., Molenaar, R. J., Vreman, S., Harders, F., Munnink, B. B. O., Honing, R. W. H. der,
Gerhards, N., Tolsma, P., Bouwstra, R., Sikkema, R. S., Tacken, M. G., Rooij, M. M. de,
Weesendorp, E., Engelsma, M. Y., Bruschke, C. J., Smit, L. A., Koopmans, M., Poel, W. H. van
der, & Stegeman, A. (2020). SARS-CoV-2 infection in farmed minks, the Netherlands, April
and May 2020. *Eurosurveillance*, *25*(23), 2001005. https://doi.org/10.2807/1560-
7917.ES.2020.25.23.2001005
Oude Munnink, B. B., Sikkema, R. S., Nieuwenhuijse, D. F., Molenaar, R. J., Munger, E., Molenkamp,
R., van der Spek, A., Tolsma, P., Rietveld, A., Brouwer, M., Bouwmeester-Vincken, N.,
Harders, F., Hakze-van der Honing, R., Wegdam-Blans, M. C. A., Bouwstra, R. J.,
GeurtsvanKessel, C., van der Eijk, A. A., Velkers, F. C., Smit, L. A. M., … Koopmans, M. P. G.
(2021). Transmission of SARS-CoV-2 on mink farms between humans and mink and back to
humans. *Science (New York, N.Y.)*, *371*(6525), 172‒177.
https://doi.org/10.1126/science.abe5901
Patrono, L. V., Samuni, L., Corman, V. M., Nourifar, L., Röthemeier, C., Wittig, R. M., Drosten, C.,
Calvignac-Spencer, S., & Leendertz, F. H. (2018). Human coronavirus OC43 outbreak in wild
chimpanzees, Côte d'Ivoire, 2016. *Emerging Microbes & Infections*, *7*(1), 1‒4.
https://doi.org/10.1038/s41426-018-0121-2
Perez, G., & Lalande, L. D. (2025). *SARS-CoV-2 serological results of five roe deer (Capreolus
capreolus) populations in France sampled between 2010 and 2022* [Dataset]. CIRAD
Dataverse. https://doi.org/10.18167/DVN1/AR6UDN
Posit Team. (2023). *RStudio: Integrated Development Environment for R* (Version 2023.03.0)
[Logiciel]. Posit Software, PBC. http://www.posit.co/
Pozzetto, B., Legros, V., Djebali, S., Barateau, V., Guibert, N., Villard, M., Peyrot, L., Allatif, O., Fassier,
J.-B., Massardier-Pilonchéry, A., Brengel-Pesce, K., Yaugel-Novoa, M., Denolly, S., Boson, B.,
Bourlet, T., Bal, A., Valette, M., Andrieu, T., Lina, B., … Trouillet-Assant, S. (2021).
Immunogenicity and efficacy of heterologous ChAdOx1–BNT162b2 vaccination. *Nature*,
600(7890), 701‒706. https://doi.org/10.1038/s41586-021-04120-y
R Core Team. (2023). *R: A language and environment for statistical computing* (Version 4.2.2)
[Logiciel]. R Foundation for Statistical Computing. https://www.R-project.org
Sarjomaa, M., Diep, L. M., Zhang, C., Tveten, Y., Reiso, H., Thilesen, C., Nordbø, S. A., Berg, K. K.,
Aaberge, I., Pearce, N., Kersten, H., Vandenbroucke, J. P., Eikeland, R., & Fell, A. K. M. (2022).
SARS-CoV-2 antibody persistence after five and twelve months: A cohort study from South-
Eastern Norway. *PLoS ONE*, *17*(8), e0264667. https://doi.org/10.1371/journal.pone.0264667
Ulrich, L., Wernike, K., Hoffmann, D., Mettenleiter, T. C., & Beer, M. (2020). Experimental Infection of
cattle with SARS-CoV-2. *Emerging Infectious Diseases*, *26*(12), 2979‒2981.
https://doi.org/10.3201/eid2612.203799
Vijgen, L., Keyaerts, E., Lemey, P., Maes, P., Van Reeth, K., Nauwynck, H., Pensaert, M., & Van Ranst,
M. (2006). Evolutionary history of the closely related group 2 coronaviruses: Porcine
hemagglutinating encephalomyelitis virus, bovine coronavirus, and human coronavirus OC43.
Journal of Virology, *80*(14), 7270‒7274. https://doi.org/10.1128/JVI.02675-05

Wernike, K., Böttcher, J., Amelung, S., Albrecht, K., Gärtner, T., Donat, K., & Beer, M. (2022).
Antibodies against SARS-CoV-2 suggestive of single events of spillover to cattle, Germany.
Emerging Infectious Diseases, *28*(9), 1916⊡1918. https://doi.org/10.3201/eid2809.220125
Wernike, K., Fischer, L., Holsteg, M., Aebischer, A., Petrov, A., Marquart, K., Schotte, U., Schön, J.,
Hoffmann, D., Hechinger, S., Neubauer-Juric, A., Blicke, J., Mettenleiter, T. C., & Beer, M.
(2022). Serological screening in wild ruminants in Germany, 2021/2022⊡: No evidence of
SARS-CoV-2, bluetongue virus or pestivirus spread but high seroprevalences against
Schmallenberg virus. *Transboundary and Emerging Diseases*, *69*(5), e3289⊡e3296.
https://doi.org/10.1111/tbed.14600
Wickham, H., Hester, J., & Bryan, J. (2023). *readr⊡: Read rectangular text data* (Version 2.1.4)
[Logiciel]. https://readr.tidyverse.org, https://github.com/tidyverse/readr
739