**Title**: Local variations in L/M ratio influence the detection and color naming of small spots

**Authors**: Maxwell J. Greene[1], Vimal P. Pandiyan[2], Ramkumar Sabesan[2], William S. Tuten[1]*

**Affiliations:**

[1]Herbert Wertheim School of Optometry and Vision Science, University of California, Berkeley,

Berkeley, CA, USA

[2]Department of Ophthalmology, University of Washington, Seattle, WA, USA

*Corresponding author: wtuten@berkeley.edu

**Abstract:**

The distribution of long-wavelength sensitive (L) and middle-wavelength sensitive (M) cones in

the retina determines how different frequencies of incident light are sampled across space, and

has been hypothesized to influence spatial and color vision. We asked whether the detection

and color naming of small, short-duration increment stimuli depend on the relative numbers of L

and M cones illuminated. Stimuli were corrected for optical aberrations by an adaptive optics

system, and targeted to locations in the parafovea where cone spectral types were known. We

found that sensitivity to 680 nm light, normalized by sensitivity to 543 nm light, grew with the

proportion of L cones at the stimulated locus, though intra- and intersubject variability was

considerable. A similar trend was derived from a simple model of the achromatic (L+M)

pathway, as well as from photoreceptor-level ideal observers, suggesting that small spot

detection mainly relies on a non-opponent mechanism. Most stimuli were called achromatic,

with red and green responses becoming more common as stimulus intensity and local L/M ratio

symmetry increased. Our detection data confirm earlier reports that small spot psychophysics

can reveal information about local cone topography, and our color naming findings suggest that

chromatic sensitivity may improve when the L/M ratio approaches unity.

## INTRODUCTION

The relative numbers of long- (L) and middle- (M) wavelength sensitive cones in the eye can vary substantially between human trichromats. The existence of such diversity at the first stage of retinal processing was first supposed based on intersubject differences in luminous efficiencies (de Vries, 1946, 1949), and later confirmed by molecular analyses (Hagstrom et al., 1998, 2000), electroretinographic recordings (Brainard et al., 1999; Carroll et al., 2002), and *in vivo* adaptive optics imaging (Hofer, Carroll, et al., 2005; Pandiyan et al., 2022; Roorda & Williams, 1999; Zhang et al., 2019). In addition to variations in L/M ratio between subjects, the spatial distribution of L and M cones within an individual primate retina is highly non-uniform, resulting in pockets of the mosaic where the local proportion of L cones deviates from the overall average (Hofer, Carroll, et al., 2005; Mollon & Bowmaker, 1992; Roorda et al., 2001). How L/M ratio variability within and between individuals affects vision is not yet fully understood. This is because the topography of the human cone mosaic, and its effect on downstream visual processing, have historically been unamenable to direct study. Adaptive optics-based technology is well suited to advance our understanding in this area, since it enables the spectral classification of single cones, and in turn the examination of how perception depends on stimulating cones of known spectral type (Greene et al., 2024; Sabesan et al., 2016; Schmidt et al., 2018, 2019).

The retinal L/M cone ratio has been thought to influence various aspects of visual performance. Consider an observer whose photoreceptor mosaic contains a higher-than-average number of L cones (i.e., a high L/M ratio). Two hypotheses can be made about their visual system. First, compared to a typical observer with an L/M ratio of ~2:1 (Carroll et al., 2002), they should be relatively more sensitive to long-wavelength light. This hypothesis is broadly supported by sensitivity measurements obtained with both large-field stimuli (Kremers et al., 2000; Rushton & Baker, 1964) and small flashes (Cicerone & Nerger, 1989; Vimal et al., 1989), although inferring cone numbers solely from threshold data requires making assumptions about the properties of the receptors themselves (e.g., spectral sensitivity, optical density; cf. Bieber et al., 1998; He et al., 2021), as well as how their signals are processed by post-receptoral circuits (Stockman et al., 2008).

Second, a more homogeneous cone mosaic may be detrimental to color vision. Extracting spectral information from the retinal image depends on circuits that compare signals from more than one cone class. If one cone type outnumbers the other, the fidelity of cone-opponent signals may be limited by noise in the minority receptor class. Previous work suggests the detectability and discriminability of low-frequency (≤ 2 cpd) red-green gratings is reduced in

retinas with presumed asymmetric L/M ratios (Gunther & Dobkins, 2002; Hood et al., 2006).

Likewise, the ability to resolve finer-grained spatiochromatic patterns also appears to be

constrained by the spectral makeup of the photoreceptor layer (Danilova et al., 2013; Neitz et

al., 2020).

In the present study, we examine both hypotheses by measuring the detectability and color

appearance of small, brief, narrowband increments delivered via adaptive optics scanning light

ophthalmoscopy (AOSLO; Roorda et al., 2002). Advantages of the investigation described here

over older ones concerned with similar questions are (1) objective cone spectral classifications

via optoretinography were available for each subject (Pandiyan et al., 2022), and (2) the use of

AOSLO enabled the controlled delivery of stimuli to regions that varied in local L/M ratio. Our

threshold data and modeling suggest that the primary mechanism mediating detection of our

stimuli is one that sums L and M cones with a weight that does not depend on the local

arrangement of L and M cones, although our color appearance findings suggest a secondary

contribution from chromatic mechanisms. Furthermore, after equating for detectability, subjects

were more likely to categorize the stimuli as chromatic when they were sampled by similar

numbers of L and M cones, whereas stimuli landing on more homogenous retinal loci tended to

yield achromatic percepts.

**METHODS**

*Subjects*

Two authors of the present study volunteered as experimental observers (subjects 10001R

and 20217R). Prior to study enrollment, subjects gave informed written consent. All study

procedures adhered to the tenets of the 2008 Declaration of Helsinki and were approved by

institutional review boards at the University of California, Berkeley and the University of

Washington. Both subjects were males (ages 40 and 27) with normal color vision as assessed

by a Neitz anomaloscope and/or pseudoisochromatic plates viewed under standard illuminant

C. Both participants have extensive experience as adaptive optics psychophysics subjects. The

subjects' cone spectral identities were previously determined by optoretinography, a technique

based on adaptive optics optical coherence tomography (AO-OCT; Pandiyan et al., 2020). Cone

classification with AO-OCT is faster and more sensitive than AO-densitometry (Pandiyan et al.,

2022; Zhang et al., 2019). Cone types were known within a 1.37 x 1.1 deg patch of retina at 2

deg temporal eccentricity for subject 10001R and within a 0.86 x 0.93 deg patch of retina at 2.5

deg temporal eccentricity for subject 20217R. Subregions of these cone-classified mosaics are

shown in **Figure 1A**. Within his classified region, subject 10001R's L/M ratio is 1.47, while

subject 20217R's L/M ratio is 3.66.

***Retinal imaging and stimulation with adaptive optics***

We measured the relationship between increment sensitivity and cone topography by

stimulating retinal loci that differ in constituent L and M cone number. Deliberate stimulus

placement at designated points of the photoreceptor mosaic required taking high resolution

images of the retina, which we accomplished with an adaptive optics scanning laser

ophthalmoscope (AOSLO). Our AOSLO, which is described in greater detail elsewhere

(Mozaffari et al., 2020), raster scanned light from four channels onto the retina. These channels

were: a 940 nm channel for wavefront sensing, an 840 nm channel for imaging, and two

stimulation channels – 680 nm (± 14.9 nm) or "red" and 543 nm (± 13.5 nm) or "green".  The

vergences of each channel were adjusted to counteract the longitudinal chromatic aberration of

the typical human eye, as approximated by equation 5a of Atchison & Smith (2005). Two

scanning mirrors, a 16 kHz horizontal resonant scanner and a 30 Hz vertical galvanometer

scanner, produced the raster scan and hence set the 30 Hz frame rate of the display.

Photomultiplier tubes detected light returning from the retina, which was sampled at 20 MHz

during the middle 80% of the forward scan to generate 512 x 512 pixel images. In the present

experiment, the imaging field of view was about 0.9 degrees, and thus 1 pixel subtended ~0.11

arcmin. This is about a quarter of the imaging channel's diffraction-limited point spread function

bandwidth, assuming a 6.5 mm pupil.

To draw stimuli at desired locations on the retina, acousto-optic modulators varied the

intensity of the 543 nm or 680 nm light at appropriate points of the scan. Transverse chromatic

aberration (TCA), which would normally cause stimuli of different wavelengths to be laterally

offset from one another on the retina, was measured using an image-based method (Harmening

et al., 2012), and corrected by modulating the 543 nm and 680 nm light at different times within

laser scan. The subject used a bite bar during imaging which along with an X/Y/Z translation

stage served to align the subject's eye to the pupil plane of the system. To avoid changes in

TCA with pupil position (Boehm et al., 2019), the experimenter used the translation stage to

negate deviations of the pupil from its position at the time of TCA measurement, which was

marked in a video feed of the subject's 1$^{st}$ Purkinje image. Prior to imaging, the subject's eye

was instilled with 1% tropicamide and 2.5% phenylephrine to dilate the pupil and freeze

accommodation. A 1-second (= 30 frames) video of the retina was recorded for each trial of the

experiment.

*Stimuli*

All stimuli were squares, 2.25 arcmin in side length and 3 AOSLO frames (~67 ms) in

duration. Stimuli consisted of narrowband light centered on either 543 nm or 680 nm, and were

shown against a large achromatic field, which was made of light from a Maxwellian-view RGB

projector in addition to the infrared light used for imaging and wavefront-sensing. Light leak from

the stimulation channels contributed negligibly to the background. The projector furnished a 1.1

deg square patch, which the subject aligned with the imaging raster to render the infrared light

invisible, as well as a dimmer circular surround subtending 18 deg. The projector provided a

small black spot to aid the subject's fixation. The experimenter could move the fixation marker

so that the desired retinal location was centered in the imaging field of view. The luminance of

the square background was 1702 cd m$^{-2}$, and the luminance of the surround was 851 cd m$^{-2}$.

Both background and surround were approximately metameric to equal-energy white (MacLeod-

Boynton coordinates: r = 0.707, b = 1.06), and functioned to saturate rod activity and to

approximately equalize activity in L, M, and S cones. Before the start of the experiment, the

subject focused the projector display by adjusting the position of a Badal lens.

*Stimulus delivery*

Retinal locations for stimulation were chosen at the beginning of each experiment, and

stimuli were locked in place at these locations during presentation. Stimulus stabilization on the

retina entailed several steps. First, an image of the retina captured over a single frame interval

was designated to serve as the initial reference frame. All subsequent frames were registered to

the reference frame by a fast strip-based cross-correlation algorithm (Arathorn et al., 2007;

Yang et al., 2010). This yielded a live video of the photoreceptors in which displacements and

distortions of the retinal image due to fixational eye movements were removed. Next, two

seconds of stabilized video were recorded, and the frames of this video were averaged together

to generate a high signal-to-noise ratio image which supplanted the initial reference frame. The

new reference image was compared to a previously taken image of the retina in which L, M, and

S cones were labeled, and based on this comparison the experimenter chose coordinates in the

reference image at which to deliver stimuli. The same process for video stabilization produced

real-time eye motion estimates that were used to track the points in the reference image

selected for testing, so that stimuli could be reliably drawn at the desired loci. Stimulus delivery

locations were marked in videos recorded during each trial with small digital crosses. Only trials

in which both the stimulus x-position and y-position were ≤0.75 arcmin from the tracked x and y-

positions for every frame of the delivery sequence contributed to threshold estimates. Out of

6,012 trials with subject 10001R, 5,266 (87.59%) were included, and out of 5,200 trials with

subject 20217R, 4,348 (83.62%) were included. Of the included trials, the standard deviation of

the x-position of the stimulus was 0.09 arcmin, and the standard deviation of the y-position was

0.10 arcmin, less than the diameter of a cone photoreceptor at our parafoveal test eccentricity.

Thirty-one loci were targeted in subject 10001R's retina, and 26 loci were targeted in subject

20217R's retina. At one location for subject 10001R, thresholds were measured twice to confirm

repeatability. Thus, 32 pairs of thresholds (for 543 and 680 nm stimuli) were collected from

subject 10001R, and 26 from subject 20217R. The white boxes overlaid on the mosaic images

in **Figure 1A** mark the locations targeted within each subject's eye. We compared, by Monte

Carlo method, the distribution of tested L/M ratios with the distribution expected from a

completely random selection of test locations. For each subject, $n$ retinal locations were

randomly picked, where $n$ is the number of unique 543 (or 680) nm threshold measurements

made for the given subject. The L cone proportions at the random locations were determined

(see *Computation of proportion L* below) and allocated to 8 equal-width bins over the range 0 to

1. This process was repeated 10,000 times in total, and the mean number of locations in each

bin, as well as the bin count's standard deviation across all iterations, were computed. The

Monte Carlo results are graphed together with the observed distribution of proportion L in

**Figure 1B**.

*Psychophysics*

Subjects completed a self-paced seen/not seen task to determine thresholds for detecting

small flashes of 543 or 680 nm light on a white background. On each trial, the subject pressed a

key to initiate stimulus presentation, and indicated with an additional keyboard press whether

the stimulus appeared red, green, or achromatic, or if it was not seen. In the case of poor

stimulus delivery (which might be caused by a microsaccade, for instance), the experimenter

instructed the subject to repeat the stimulus presentation and respond anew. Trial-to-trial

stimulus intensities were generated from a Bayesian, adaptive staircase procedure, QUEST+

(Watson, 2017). In the typical measurement block, two discrete retinal locations were selected

for testing. At each test location, sensitivity and color appearance were assessed with two test

wavelengths: 543 nm and 680 nm. For each wavelength and location, two staircases ran in

parallel, with one staircase starting at 25% of the maximum intensity and the other at 75%.

Thus, each measurement block consisted of 8 independent staircases; these staircases were

randomly interleaved so that the subject could not predict the intensity, wavelength, or retinal

locus of stimulation based on the preceding trial. Except for a few early experimental sessions,

each staircase contained 25 trials and, on 20% of trials the stimulus intensity suggested by the

QUEST+ algorithm was ignored and stimulus intensity was instead drawn at random from a

uniform distribution on the interval between 0 and the maximum intensity. Because threshold

intensities were typically a very small fraction of the maximum power level, this was practically

equivalent to interspersing high intensity stimuli throughout the staircase. These random, high

intensity stimuli served to maintain the subject's attention, as well as to assess suprathreshold

color judgements. Generally, 100 trials per location were collected for each of the two stimulus

wavelengths, per session. Three to six locations were tested every session, corresponding to a

total of 600 to 1200 trials. Weibull functions were fit to the staircase data with the Palamedes

toolbox, and threshold was taken as the stimulus intensity that corresponded to a 50%

frequency of seeing.

***Computation of proportion L***

The assortment of L and M cones hit by the stimulus is characterized in this paper as

"proportion L": the number of illuminated L cones out of the total number of illuminated L and M

cones. A simple way to count the number of stimulated cones would be to tally the number of

cone centers that fall within the nominal boundary of the stimulus on the retina. However, due to

diffraction and any residual aberrations not corrected by the AOSLO, the boundary of the

proximal stimulus is ambiguous. Additionally, a single cone admits light over an extended area

and not just a single point. Our calculation of proportion L accounted for these nuances. We

represented the cone mosaic as an array of two-dimensional Gaussians, which approximated

the cone aperture function (MacLeod et al., 1992). Each Gaussian had a full width at half

maximum of 0.59 arcmin and a volume of one. The spatial distribution of light on the retina

during stimulation was computed by convolving an intensity-normalized distribution of light

defined by the shape of the stimulus with a diffraction-limited point spread function computed for

a 6.5-mm pupil and subsequently aberrated by 0.05 D of defocus to account for any residual

aberrations (Harmening et al., 2014). The point spread function's volume was normalized to

unity, and its width was determined by the central wavelength of the stimulus (i.e., 543 nm or

680 nm). The retinal light distribution was multiplied with the array of L, M, or S cone aperture

functions, and the integral of this product gave the number of cones of a particular spectral type

that received the stimulus. Therefore, the number of stimulated L (or M, or S) cones is related to

the amount of light accepted by the L (or M, or S) submosaic. The proportion L reported for a

given threshold measurement is the average proportion L across trials. In cases where not all

stimulated cones were classified, the range of possible L proportions (assuming all the

unclassified cones were either L or M) is given in addition to the proportion L computed from
only the classified cones.

***Modeling***

Sensitivity to briefly-presented small spots might be expected to grow with the aggregate
cone excitations produced by the light. Based on this idea, we computed theoretical 680/543
sensitivity ratios for every local L/M ratio encountered experimentally. Theoretical sensitivity
ratios were derived from a simple additive model, and from ideal observers, which are detailed
below.

*Additive model*

In our simple model, sensitivity is taken to be proportional to the response of a mechanism
that combines L and M cone excitations. The responses to 680 and 543 nm stimuli at threshold
are necessarily equal, and from this it can be shown that the ratio of 680 nm sensitivity ($S_{680}$) to
543 nm sensitivity ($S_{543}$) is given by the following expression:

$$\frac{S_{680}}{S_{543}} = \frac{p_L L_{680} + (1 - p_L) M_{680}}{p_L L_{543} + (1 - p_L) M_{543}}$$   **(1)**

where $p_L$ is proportion L, $L_{680}$ (or $M_{680}$) are the average L (or M) cone excitations per quantum
of the 680 nm primary at the cornea, and $L_{543}$ (or $M_{543}$) are the average L (or M) cone
excitations per quantum of the 543 nm primary at the cornea. Sensitivities are in terms of
spectral power at the cornea (quanta s$^{-1}$ nm$^{-1}$), integrated over wavelength. $L_{680}$, $M_{680}$, $L_{543}$, and
$M_{543}$ are computed from the Stockman & Sharpe (2000) cone fundamentals (quantal units),
accounting for filtering by the lens and macula, photopigment optical densities, and the
wavelengths at which cone absorptances peak (Kraft et al., 1998; Neitz & Neitz, 2011).

*Ideal observer analysis*

We used ideal observers to assess how well our results agree with optimal usage of
information available at the level of cone photoisomerizations. Our implementation of this
analysis followed that of Geisler (1984) and incorporated the advantages of adaptive optics.
That is, instead of ideal observers with simulated cone lattices and theoretically-based optics,
we specified ideal observers that possessed our subjects' cone mosaics and the near-diffraction
limited optics afforded by AOSLO. The ideal observer's task was to distinguish the stimulus and

background from the background alone. The background was specified to be identical, in terms

of size and spectral power distribution, to the one used in the real experiment. The stimuli

presented to the ideal observer differed from the real stimuli only in that they were perfectly

monochromatic, rather than narrowband, and displayed continuously rather than raster

scanned. For each targeted location within each subject's eye, we determined the stimulus

intensity that yielded a d' of 1.36 (75% correct in a yes-no task), where d' is given by:

270
$$d' = \frac{\sum_{i=1}^{n}(\beta_i - \alpha_i)\ln\left(\beta_i/\alpha_i\right)}{\left[0.5\sum_{i}^{n}(\beta_i + \alpha_i)\ln^2\left(\beta_i/\alpha_i\right)\right]^{1/2}}$$
**(2)**

Geisler derived this formula from the optimal decision rule of an ideal observer limited by

Poisson noise arising from the quantal nature of light. Here, $\alpha_i$ represents the number of cone

excitations elicited by the background, and $\beta_i$ represents the number of cone excitations elicited

by the test and background together. The ideal observer's 680/543 sensitivity ratio is the 543

nm intensity required to achieve the desired d', divided by the 680 nm intensity required to

achieve the same d'. We note that one advantage of expressing both our experimental and

modeled data as sensitivity ratios is that it eliminates the need to apply a scaling factor to

account for efficiency differences between the ideal and human observers.

***Color appearance***

As described above, the subject indicated whether the color appearance of each detected

stimulus was best categorized as red, green, or achromatic.  680 nm light was seldomly

categorized as "green", and 543 nm light was rarely called "red" (see Results). Because of this,

we opted to analyze the effect of stimulus intensity and cone topography on the probability of

giving a color response consistent with the large-field appearance of the stimulus wavelength –

that is, "green" for 543 nm and "red" for 680 nm. For convenience, we refer to these responses

as "expected" color responses.  In particular, we were interested in whether the subject was

more likely to give the expected color response when the stimulus landed on spectrally-

balanced regions of the cone mosaic, compared with regions dominated by a single cone class.

For this analysis, we characterized the population of cones under the stimulus by its

heterogeneity, defined as:

293
$$\text{heterogeneity} \ = \ 1 \ - \ \frac{|p_L - 0.5|}{0.5}$$
**(3)**

Where $p_L$ is proportion L. This expression yields a heterogeneity value of 1 when the local

L/M ratio is 1, and a value of 0 when the local receptor configuration is completely

homogeneous (i.e. all L or all M).

Using a simple linear regression to examine the relationship between local heterogeneity

and color naming was infeasible, because of the dichotomous response variable (on each trial,

the subject either gave the expected color response or did not) and the non-independence of

the observations (multiple trials per subject). Instead, we designed the following generalized

linear mixed-effects model:

304
$$logit(\pi_{ijk}) = \beta_0 + \beta_1 Log\ Intensity_{ijk} + \beta_2 Heterogeneity_{ijk}$$

305
$$+ \beta_3(Log\ Intensity_{ijk} \times Heterogeneity_{ijk}) + b_{0i} + b_{0j}$$

The probability of the expected color response from subject *i* at wavelength *k* is represented

by $\pi_{ijk}$. Here, *i* is the subject index, *j* is the stimulus wavelength index, and *k* is the trial index

for the *i*$^{th}$ subject and at the *k*$^{th}$ wavelength. Stimulus intensity, in terms of threshold multiple,

and heterogeneity (as defined above) are treated as fixed effects with coefficients $\beta_1$ and $\beta_2$,

respectively. We include an interaction term with coefficient $\beta_3$. In addition to the fixed intercept

$\beta_0$, we accounted for between-subject variability by including the random intercept $b_{0i}$, and for

differences in color responses between wavelengths with the random intercept $b_{0j}$. We fit the

model to our data with the Laplace method, using the *fitglme* routine (MATLAB 2024b; The

MathWorks, Inc., Natick, MA, USA). Since our grouping variables of subject and wavelength

contained only two levels each, we verified that the size and significance of each effect of

interest estimated from the mixed model were similar to those estimated from a model in which

subject and wavelength were treated as fixed effects. Additionally, we compared the above

model to a reduced model in which the fixed effect of heterogeneity, and the interaction between

heterogeneity and intensity, were removed.

**RESULTS**

***680/543 nm sensitivity vs. local L/M ratio***

Thresholds were measured for stimuli targeted to the retinal regions indicated in **Figure 1A**.

Each flash stimulated an average of 4.6 cones in subject 10001R's eye, and 4.0 cones in

**Figure 1 | Local variation in cone spectral topography in the trichromatic retina. (A)** Labeled cone
mosaic images acquired from the two participants in this study (0.65° x 0.65° field of view). L, M, and S
cones are labeled using red, green, and blue markers, respectively. The mean delivery coordinate for
each tested location is marked by a white box whose size represents the nominal stimulus area on the
retina (2.25 x 2.25 arcmin). The white scale bar corresponds to a visual angle of 5 arcmin. The dark
unlabeled region in subject 20217R is due to shadows cast by an overlying retinal blood vessel, which
preclude reliable spectral classification by optoretinography. **(B)** Histograms showing the distribution of
local proportion L selected for testing in each subject. The white-filled bars represent the distribution of
test sites selected manually by the experimenter. The red lines and pink shaded areas show the results
(mean ± 1 STD) of a Monte Carlo simulation (10,000 iterations) in which an equivalent number of test
sites in each subject (n = 32 for 10001R; n = 26 for 20217R) was selected at random and histograms
were recomputed following the same procedure.

subject 20217R's eye, with cone number calculated as described in the *Computation of*
*proportion L* section. The histograms in **Figure 1B** summarize the local L/M ratios sampled
throughout the study, in addition to the variety in L/M ratio that would arise from arbitrary
stimulus placement. The empirical distributions of local proportion L we obtained for each
subject were qualitatively similar to the simulated distributions that resulted from randomly

selecting test locations. In other words, although we manually selected locations for testing,

systematic sampling biases are unlikely to underlie the results we describe below. That being

said, we did make a deliberate effort to increase our sampling of majority-M cone regions in

subject 20217R, who has a higher-than-average global L/M ratio, and in the end targeted three

such locations. According to the Monte Carlo simulation, the probability of encountering three or

more majority M cone regions in subject 20217R's eye, given the number of locations tested,

was only ~23%.

The spread of sensitivities to 543 nm and 680 nm flashes is depicted in the box plots of

**Figure 2A**. The CIE physiologically relevant 2-deg luminosity function, scaled to fit our data, has

been plotted for comparison (CIE, 2006). For the standard observer, the quantal efficiency of

543 nm light is 1.83 $\log_{10}$ units higher than it is for 680 nm light. The difference between the

median sensitivities to 543 and 680 nm small spots was 1.68 $\log_{10}$ units for subject 10001R, and

1.80 $\log_{10}$ units for subject 20217R, which is roughly what would be expected based on V(λ).

The interquartile range (IQR) of 543 nm sensitivities spanned 0.12 $\log_{10}$ units for subject

10001R and 0.16 $\log_{10}$ units for subject 20217R, while the IQR of 680 nm sensitivities were

slightly broader, spanning 0.22 $\log_{10}$ units for subject 10001R, and 0.34 $\log_{10}$ units for subject

20217R. Since L and M cones have almost identical sensitivities to 543 nm light, the dispersion

of 543 nm thresholds must have been driven primarily by factors other than cone topography. In

contrast, L cones absorb more at 680 nm than M cones, and so the increased variability in our

observers' 680 nm thresholds likely originated, at least in part, from local fluctuations in L/M

ratio. In addition to this within-observer variability, we note that, for both stimulus wavelengths,

subject 10001R was more sensitive than subject 20217R, likely because the latter was tested at

a slightly more eccentric location where cone density is lower (**Figure 1A**). Thus, to examine the

impact of proportion L on sensitivity while discounting other factors, we normalized the 680 nm

sensitivity measured at each test locus by the corresponding 543 nm sensitivity.

The ratio of 680 to 543 nm sensitivity is plotted as a function of local proportion L in **Figure**

**2B**. Subject 10001R's data are represented by the blue circles, and subject 20217R's by the

orange circles. As expected, normalized sensitivity to 680 nm increased with relative L cone

numerosity, and significant correlations between these variables were found for both subjects

(10001R: Spearman's rho = 0.80, p < 0.001; 20217R: Spearman's rho = 0.73, p < 0.001). There

was considerable variability in thresholds measured for patches of retina with similar proportions

of L cones; this variability would preclude precise estimation of L/M ratio from detection

thresholds. Data from subject 10001R generally lie above data from subject 20217R, indicating

that the former subject is more sensitive to 680 nm stimuli than the latter, all else being equal.

**Figure 2 | Local cone topography influences small-spot spectral sensitivity.** (**A**) Boxplots of
sensitivity to 543 nm and 680 nm narrowband increments obtained from subjects 10001R (blue) and
20217R (orange). Boxes indicate the median sensitivity (middle line) and interquartile range (upper and
lower box edges); the whiskers span the full range of sensitivities measured with each wavelength. For
clarity, each subject's boxes have been shifted ±5 nm from the tested wavelength. Retinal locations
selected for testing are shown in **Figure 1**. The physiologically-relevant CIE 2-deg photopic luminous
efficiency function, V(λ), is shown in black (CIE, 2006). Sensitivities are defined as the reciprocal of the
detection threshold intensity expressed in units of quanta s$^{-1}$ deg$^{-2}$ at the cornea. (**B**) The ratio of 680 nm
sensitivity to 543 nm sensitivity is plotted against the local proportion L of the stimulated region (see
Methods). The blue and orange circles correspond to data from subjects 10001R and 20217R,
respectively. In instances where the stimulated region contained an unclassified cone, the range of
possible local L/M ratios (calculated assuming the missing cone is either L or M) is denoted by a black
horizontal line. The pair of points with a dashed red outline correspond to the locus selected for repeat
testing in subject 10001R. Predictions from the additive model described in the text are shown by solid
lines, where the line color denotes the λ$_{max}$ value of the L-cone pigment used to model sensitivity (Kraft et
al., 1998).

This intersubject variability may reflect differences in L cone photopigments. The solid red

curves give the 680/543 sensitivity functions predicted from **Equation 1**, assuming L cone

spectral peaks at 563.4, 558.9, or 555.5 nm. Data from subject 10001R mostly fall above the

additive model predictions, while data from subject 20217R fall near the curves for which the L

cone λ$_{max}$ is either 558.9 or 563.4 nm. This does not necessarily mean that subject 10001R's L

cone spectral peak lies further in the red than would be explained by known L cone pigment

variants. Instead, it may be that the additive model underestimates long-wavelength sensitivity
by failing to account for signals from non-additive (i.e., cone-opponent) mechanisms that may
contribute to detection.
405
***Comparison of predicted and observed data***
**Figure 3A** plots, for each targeted location, the 680/543 nm sensitivity ratios predicted by
the additive model (diamonds), and by the ideal observer (asterisks), against the observed ratio.
In this plot, predictions for subject 10001R (blue markers) were made assuming an L cone $\lambda_{max}$
of 563.4 nm, while a $\lambda_{max}$ of 558.9 nm was used for subject 20217R (orange markers). The use
of different spectral peaks was meant to account for the systematic differences between the two
subjects' data (**Figure 2B**), though it is entirely possible that both subjects have the same L
cone opsins, and the enhanced long-wavelength sensitivity of subject 10001R might be
explained by post-receptoral factors. The additive model and ideal observer yielded similar
sensitivity ratios, which was expected since both sets of predictions share the premise that
sensitivity increases with the sum of cone excitations. Most of the data in **Figure 3A** lie close to
the 1:1 line, suggesting that our empirical results are consistent with the hypothesis that small
spot detection is largely mediated by a mechanism that adds unweighted inputs from L and M
cones in its receptive field; however, the theoretical data tend to underestimate the observed
sensitivity ratios.
To understand how errors in the model predictions vary with the choice of L cone $\lambda_{max}$, root
mean square error (RMSE) is graphed as a function of $\lambda_{max}$ in **Figure 3B**. In this plot, the
continuous lines give the additive model RMSE over a range of wavelengths. The errors for the
additive model and ideal observer at three biologically plausible spectral peaks are plotted as
diamonds and asterisks, respectively. If selection is restricted to the three $\lambda_{max}$ values of interest,
the RMSE is similar between the two subjects when a wavelength of 563.4 nm is selected for
10001R, and a wavelength of 558.9 nm for 20217R. However, as the solid curves indicate, the
RMSE (at least for the additive model) approach minima at longer wavelengths than those just
stated: the minimum for 10001R occurs around 566.5 nm, and the minimum for 20217R occurs
around 562 nm. Thus, to best model the data for subject 10001R, the L cone sensitivity peak
chosen must exceed the maximum value reported in the literature (~563 nm). The difference
between the optimal additive model $\lambda_{max}$ values for the two subjects is ~4.5 nm, which is similar
to the difference between the two most commonly found L-cone spectral peaks (Kraft et al.,
1998; Neitz & Neitz, 2011).

**Figure 3 | Modeling the 680/543 nm sensitivity ratio.** (**A**) Predicted 680/543 nm sensitivity ratios are
plotted against their empirically-measured counterparts. Additive model and ideal observer predictions for
each retinal locus are indicated by diamonds and asterisks, respectively. For subject 10001R, an L-cone
$\lambda_{max}$ of 563.4 nm was used to generate the predicted data; for subject 20217R, a value of 558.9 nm was
chosen. (**B**) Root mean squared errors (RMSE) of the additive and ideal observer model predictions are
plotted as a function of L-cone $\lambda_{max}$ for subject 10001R (blue) and 20217R (orange). Markers are shown
at three physiologically-plausible values of L-cone $\lambda_{max}$ for both models. In addition, solid lines depict the
performance of the additive model evaluated with finer L-cone $\lambda_{max}$ sampling, revealing minima at or
beyond the longest values reported in the literature.

### *Color appearance*

447        Our color appearance results are summarized in **Figure 4**. The stacked bar plots in **Figure**
**4A** represent the proportion of seen trials categorized as red, green or achromatic. For both
stimulus wavelengths, detected flashes were most frequently categorized as achromatic (90.5%
for 543 nm; 58.2% for 680 nm). The long-wavelength stimulus was perceived as red at a higher
rate (47.7% for 10001R; 32.5% for 20217R) than the middle-wavelength stimulus was as green
(6.6% for 10001R; 9.1% for 20217R). "Non-expected" color responses were rare: 680 nm
flashes were almost never seen as green (0.06% for 10001R; 1.01% for 20217R), and the 543
nm stimuli were very seldom called red (0.44% for 10001R; 4.01% for 20217R). Of note, subject
20217R, whose retina is comparatively L-rich, was less likely to call 680 nm stimuli red.

**Figure 4 | Summary of small-spot color appearance results.** (**A**) Stacked bar plots showing the proportion of seen trials categorized as red, green, and achromatic as a function of stimulus wavelength. Results from subjects 100001R and 20217R are shown in the top and bottom rows, respectively. (**B**) Color-coded matrices represent the naming data from (A) binned by local mosaic demographics and stimulus intensity (relative to local threshold). Darker squares indicate intensity-mosaic combinations that were more likely to yield achromatic percepts, whereas brighter green and red squares indicate a higher likelihood of responding "green" and "red" to the 543 nm and 680 nm flashes, respectively. Note that "non-expected" color responses (e.g., categorizing a 680 nm stimulus as "green") occurred rarely and were thus excluded from these plots. To enhance visualization, bins containing fewer than 10 trials are not shown. Squares filled with diagonal lines denote empty bins.

To examine the effects of local cone topography and stimulus intensity on color appearance, the data were binned along the intensity and local proportion L dimensions, and for each bin, the fraction of trials on which the subject gave the expected color response was calculated. The matrices in **Figure 4B** give the frequency of expected color responses as a function of intensity and L cone proportion. Darker matrix elements correspond to high rates of achromatic percepts.

In these plots, intensities are expressed as multiples of the local threshold, which has

implications for the distribution of matrix bins. As described in the Methods, trial intensity was

primarily determined by an adaptive staircase, which concentrates most presentations near

threshold; however, on a random subset of trials, high-intensity stimuli were interspersed.

Because subject 20217R had lower absolute sensitivity to both stimulus wavelengths (**Figure**

**2A**), these high-intensity trials end up being allocated to intensity bins closer to threshold. In

addition, 680 nm sensitivity increased with local proportion L in both subjects (**Figure 2B**).

Consequently, 680 nm flashes of a fixed intensity would be assigned to intensity bins

increasingly farther from threshold as the local proportion L increases. Despite this asymmetry

in stimulus detectability across observers, wavelength, and retinal locus, **Figure 4B** suggests

the likelihood of calling 680 nm flashes red or 543 nm flashes green grew with normalized

stimulus intensity.

The effect of local L/M ratio on color naming was less obvious than that of intensity. For the

543 nm stimulus, green responses were almost never given for stimuli falling within the leftmost

or rightmost bins, which correspond to highly asymmetric L/M ratios (**Figure 4B**, left column).

Instead, green responses were concentrated near 50% L at higher stimulus intensities. If a

similar hotspot exists for 680 nm, it is more subtle. At near-threshold intensities, chromatic

responses appear to be more frequent for retinal loci with balanced L and M numbers, but the

color response matrix rows become more uniform as normalized stimulus intensity increases

(**Figure 4B**, right column). In subject 10001R, one retinal locus containing mainly M cones was

tested (**Figure 4B**, top row, leftmost matrix columns). With 543 nm light, predominantly

achromatic percepts were logged at this spectrally homogenous location across a range of

stimulus intensities, mirroring the pattern observed in more frequently encountered regions

dominated by L cones. For 680 nm light, however, the stimulus was reliably categorized as

"red". We suspect this may be a consequence of the higher absolute illuminance required to

detect 680 nm flashes landing in this M-rich region, leading to increased light scatter onto

nearby cones that might counteract the decline in chromatic sensitivity that would otherwise

occur as local proportion L approaches zero.

To test the hypothesis that L/M ratio symmetry would be advantageous for color vision

(Gunther & Dobkins, 2002; Hood et al., 2006), we categorized each stimulated retinal locus by

its spectral heterogeneity, applying an index similar to the one developed by Hofer, Singer, et al.

(2005) to quantify the symmetry of global L/M cone ratios (see **Equation 3**). A schematic

illustrating the relationship between local spectral composition and the heterogeneity index is

shown in **Figure 5A**. The distribution of heterogeneity indices encountered across all trials is

**Figure 5 | Chromatic responses increase with stimulus intensity and mosaic heterogeneity.** (**A**) Schematic depicting the relationship between local cone configurations and the heterogeneity index described in the methods. (**B**) Histograms showing the distribution of heterogeneity indices encountered across all seen trials (n = 5,552). Bars are color-coded to represent the bottom (dark gray), middle (light gray), and top (white) tertiles of the heterogeneity distribution. Data from the two study participants are combined in this analysis. (**C**) The percentage of trials in which the "expected" chromatic response ("red" to 680 nm or "green" to 543 nm) was given is plotted as a function of binned stimulus intensity and mosaic heterogeneity. As in **Figure 4**, stimulus intensity is expressed in multiples of local detection threshold; the tick marks on the abscissa indicate the edges of the intensity bins. Markers are color-coded according to the heterogeneity tertiles shown in (B). Marker area is scaled to the number of trials in each bin, and its x-position represents the median stimulus intensity within each bin.

depicted in **Figure 5B**. We divided this distribution into tertiles to visualize the joint effects of

mosaic heterogeneity and normalized stimulus intensity on the probability that the subject gives

a chromatic response. In **Figure 5C**, it is clear that increasing stimulus intensity (relative to

threshold) increases the likelihood of categorizing the stimulus as chromatic, reiterating the

trends we observed in the color response matrices of **Figure 4**. Comparing the intensity-

response curves for the three heterogeneity bins, it is also apparent that the likelihood of our

small test flash being judged as chromatic increases as the mosaic becomes more

heterogeneous – at least for stimulus intensities at and slightly above threshold. To determine

whether these apparent effects of intensity ($\beta_1$) and cone mosaic topography ($\beta_2$) on color

response reached statistical significance, we used the generalized linear mixed effects model

detailed in the Methods. The main effects of intensity ($\beta_1$= 1.36, p < 0.001) and heterogeneity

($\beta_2$ = 1.47, p < 0.001) were significant, while the interaction of the two ($\beta_3$ = 0.34, p = 0.072) was

not. The full model outperformed the reduced model in which the heterogeneity and interaction

terms were removed ($\lambda_{LR}$ = 139.61, $\Delta$df = 2, $\Delta$AIC = -135.6, p < 0.001).

**DISCUSSION**

Small spot detection, discrimination, and color naming data have been used before to make

inferences about the distribution of L, M, and S cones in the retina as well as the nature of

mechanisms that combine inputs from cones of different type (Chaparro et al., 1994; Cicerone &

Nerger, 1989; Finkelstein & Hood, 1984; King-Smith & Carden, 1976; Krauskopf, 1964; Otake et

al., 2000; Vimal et al., 1989). Distinguishing the influence of cone spectral demographics from

the effects of postreceptoral processing on behavioral data is, however, challenging in the

absence of objective measurements of cone numbers (e.g., Krauskopf, 2000). We studied the

direct link between cone topography and behavior by measuring small spot detectability and

color appearance in subjects whose cones were spectrally classified by an objective method.

Our stimuli were delivered through an AOSLO, which mitigated the effects of optical aberrations

and fixational eye movements, in turn enabling us to estimate the trichromatic inputs to post-

receptoral mechanisms on each experiment trial. Our results permit two conclusions to be

drawn about the relationship between local cone mosaic topography and vision. First, as

expected, spectral sensitivity to short-duration, retinally-stabilized small spots depended

strongly on local L/M cone ratio. Second, subjects were more likely to rate brief, monochromatic

flashes as chromatic if they were sampled by a balanced number of L and M cones. We discuss

these findings in turn below.

***Local L/M cone ratio and small-spot detection***

The dependence of visual sensitivity on cone spectral topography was relatively

straightforward: long-wavelength stimuli were overall more easily seen as the retinal patches

they landed on became increasingly L cone rich (**Figure 2B**). The relationship between the

680/543 nm sensitivity ratio and local L/M ratio was reasonably well-described by two variants of

a non-opponent detection model that included a characterization of early visual processing

through the photoisomerization stage (**Figure 3A**). Together, these results are compatible with

the idea that carefully collected behavioral data can reveal spatiochromatic variations in the

receptor array (Cicerone & Nerger, 1989; Krauskopf, 1964; Otake et al., 2000; Vimal et al.,

1989; Williams et al., 1981). However, despite the strong correlation between our

measurements of local spectral sensitivity and objectively-determined cone topography, several

factors evident in our data would complicate the precise determination of local L/M ratio solely

from detection thresholds as measured in this study.

First, as can been seen in **Figure 2B**, locations within an individual retina with similar L/M

numerosity are not always equally sensitive. The origins of this within-subject variability are not

entirely clear. It likely reflects a combination of neural and psychophysical factors. Regarding

the former, physiological evidence from the primate retina suggests that downstream neurons

may not sample L and M cones with uniform weights across small spatial scales, even within

retinal ganglion cell subclasses that do not systematically favor one cone type over the other

(Field et al., 2010; Li et al., 2014). Likewise, it seems plausible that perceptual judgments reliant

on the signals of fewer than a half-dozen cones will be inherently noisy. Although the one test

site we deliberately retested in separate sessions yielded near-identical sensitivity ratios (**Figure**

**2B**, outlined pair), understanding how much variance in our data might be attributed to

psychophysical noise would require a more comprehensive analysis of test-retest variability in

our experimental paradigm. Finally, an additional source of experimental noise could arise from

imperfect cone spectral classifications. In our measurements, most of the stimulus light fell on

fewer than five cones on any given trial, and so a single misclassified (or unclassified) cone in

the stimulated region would produce significant error in our estimate of the underlying L/M ratio.

Given that previous work has estimated the error rate of OCT-based cone classification

methods to be less than 0.5% (Pandiyan et al., 2022; Zhang et al., 2019), we expect spectral

misclassifications to contribute only negligibly to the variance present in our dataset.

Intersubject variability poses a second, arguably more substantial challenge to estimating

proportion L from increment sensitivity measurements. In our sample of two trichromats, one

subject (10001R) tended to exhibit higher 680/543 nm sensitivity ratios than the other (20217R)

when the tested locations had matched L/M ratios (**Figure 2B**). This aligns with previous work

articulating how factors other than cone numerosity can drive considerable between-subject

differences in spectral sensitivity (Bieber et al., 1998; He et al., 2021). One well-known source of

individual differences in the color-normal population originates from genetic variations in the L

opsin gene that alter its spectral tuning (Kraft et al., 1998; Neitz & Jacobs, 1986; Neitz & Neitz,

2011). In the non-opponent modeling of our observers' data, comparable amounts of variance in

the data could be accounted for by shifting 10001R's L-cone $\lambda_{max}$ ~4.5 nm beyond the value

used for 20217R (**Figure 3B**). In addition to differences in cone opsin genetics, trichromatic

encoding can be affected by pre-retinal filtering by the lens and macular pigment, as well as by

variation in photopigment optical density (Asano et al., 2016). While genetic sequencing can be

used to estimate the peak sensitivities of the L and M cone opsins (Neitz & Neitz, 2011),

psychophysical approaches based on color matching have been proposed to derive more

comprehensive estimates of cone action spectra in individual eyes (Shi et al., 2024; Thomas &

Mollon, 2004). Applying these techniques to build more realistic models of our subjects' early

visual systems is a goal of future work.

Finally, we note that, alongside pre-retinal and receptoral factors, the involvement of non-

additive post-excitation mechanisms could further obscure the link between cone numbers and

small-spot detection thresholds. Although the systematic increase in sensitivity to long-

wavelength light with local L cone density comports with a significant contribution from non-

opponent mechanisms to small-spot detection, both our additive model and ideal observer

predictions tended to underestimate relative sensitivity to 680 nm across a range of L cone $\lambda_{max}$

values examined (**Figure 2B**; **Figure 3A**). Similarly, although the majority of near-threshold

stimuli generated achromatic percepts (**Figure 4A**), slightly suprathreshold stimuli were often

categorized as "chromatic", particularly when the stimulus wavelength was 680 nm (**Figure 4B**).

These facts collectively point towards a secondary contribution from an opponent channel to our

sensitivity measurements.

Previous studies have analyzed detection and discrimination data collected with small-field

stimuli and arrived at similar conclusions. King-Smith & Carden (1976) found evidence that

spectral sensitivity to small (3 arcmin), brief (10 ms) increment flashes presented on an intense

neutral background resembles the photopic luminous efficiency function measured with larger

stimuli via heterochromatic flicker photometry. This concordance suggests the involvement of a

non-opponent, "achromatic" pathway in detecting narrowband increments of small angular

subtense. However, their results also indicated that, at longer wavelengths, near-threshold

stimuli could be readily discriminated from white, leading them to conclude that the so-called

"color" and "luminance" systems have similar sensitivities in this part of the spectrum. By

contrast, King-Smith and Carden's results indicated a substantial reduction in the sensitivity of

the chromatic system to minute flashes of intermediate wavelength, presumably due to the

similarity in L and M responses produced by mid-spectrum lights. Subsequent investigations

into small-spot detection and discrimination also support a role for "red-green" opponent

mechanisms in processing small flashes near threshold, particularly for longer wavelength

stimuli (Chaparro et al., 1994; Finkelstein & Hood, 1984).

The likely involvement of the L-M pathway in our detection data introduces another plausible

explanation for the sensitivity differences we observed between individuals. Suprathreshold

color appearance has been shown to be robust to large individual variations in L/M ratio

(Brainard et al., 2000; Miyahara et al., 1998; Neitz et al., 2002). One way to accomplish this

invariance would be to reduce the input gain of the more numerous cone type to downstream

cone-opponent circuits. In cases where L cones predominate, such sensory regulation would

reduce the relative contribution of the chromatic pathway to detection. The loss of chromatic

sensitivity could manifest as a general elevation in long-wavelength thresholds and a

corresponding increase in "achromatic" percepts near threshold. In line with this hypothesis, we

found the subject with a greater abundance of L cones (20217R; L/M ratio = 3.66) exhibited a

lower relative sensitivity to long-wavelength stimuli (**Figure 2**) and was less likely to categorize

680 nm flashes as "red" (**Figure 4**) in comparison to subject 10001R, whose L/M ratio of 1.47 is

closer to the typical value (Carroll et al., 2002). While the mechanistic basis of these trends

remains unclear, obtaining a more detailed characterization of the light encoding properties of

our subjects' cone mosaics would, as described above, enable the relative contributions of post-

receptoral processes to our measurements to be delineated more clearly.

***Local L/M cone ratio and small-spot color appearance***

In addition to measuring detection performance, our observers were asked to assert

whether the appearance of each seen trial was best described as "red", "green", or

"achromatic". While this tripartite categorization scheme may seem coarse in relation to the

richness of human color experience, previous color naming studies suggest these terms capture

the majority of sensations evoked by small middle- and long-wavelength flashes presented on a

neutral photopic background (Bouman & Walraven, 1957; Cicerone & Nerger, 1989; Krauskopf

& Srebro, 1965; Sabesan et al., 2016; Schmidt et al., 2018). As shown in **Figure 4A**, we found a

disparity in the chromatic response rates for the 543 nm and 680 nm flashes, with the latter

stimuli more frequently rated as "red" than the former were categorized as "green". These

differences likely arise from the relative insensitivity of the L-M pathway to middle-spectrum

lights of small angular subtense, as discussed in the preceding section (Chaparro et al., 1994;

King-Smith & Carden, 1976). Despite the asymmetry in overall response rates to our two

narrowband stimuli, our analysis demonstrates that flashes of either wavelength were more

likely to be categorized as "chromatic" when the retinal neighborhood they encountered had an

L/M cone ratio near unity (**Figure 5**). Because the mosaic dependence of color naming was

maintained when stimuli were equated for detectability, we interpret our appearance results as

evidence that the sensitivity of the "red-green" chromatic mechanism covaries with the

underlying distribution of L and M cones.

Prior studies have produced data in support of the idea that L/M ratio symmetry may be

beneficial for chromatic detection and appearance mechanisms. Gunther & Dobkins (2002)

found a relationship between L/M ratio estimated from heterochromatic flicker photometry and

sensitivity to red/green gratings, concluding that observers with ratios closer to one had superior

chromatic contrast sensitivity compared to observers with skewed ratios. In subjects whose

mosaics had been characterized using adaptive optics densitometry, Neitz et al. (2020) also

observed a positive correlation between L/M ratio symmetry and sensitivity to fine-grained

red/green gratings, mirroring earlier results suggesting the ability to localize the gap in cone-

isolating "Landolt C" targets may depend on the relative numbers of L and M cones (Danilova et

al., 2013). Finally, Hofer, Singer, et al. (2005) reported that individuals with more balanced L/M

ratios were more likely to rate cone-sized flashes as chromatic. Our results demonstrating a

direct relationship between local cone demographics and color appearance are in keeping with

these earlier findings, although it is worth noting that our study has the added advantage of

knowing both the spectral topography of the observer's cone mosaic and the landing position of

the stimulus on each trial.

A spectrally heterogeneous mosaic may not only be beneficial for chromatic sensitivity, but

also for discriminating color from luminance information. In the context of the present

experiment, the task of distinguishing a small, monochromatic light from an equally luminous

broadband (white) light must be accomplished using signals from a small number of cones. If

the stimulated receptor array was composed of a single spectral type, identical sets of cone

excitations would be generated by the monochromatic and white lights, and the two stimuli

should, in principle, be indistinguishable.[1] If one constituent in the local receptor neighborhood

was replaced with a cone of another type, then monochromatic and white lights could be

discriminated on the basis of the outlier cell's response, though presumably noise in the

underrepresented spectral class would limit performance (Vorobyev & Osorio, 1998).

Measurements made from carriers of X-linked color vision deficiencies indicate that red-green

chromatic discrimination is impaired individuals with skewed cone ratios (Hood et al., 2006). In a

similar vein, although the midpoint of the classical Rayleigh match should depend solely on the

spectral sensitivity of the L and M cones, it has been hypothesized that individuals with

asymmetric L/M ratios will accept matches over a broader range, presumably because their

discriminative capacity is diminished (Barbur et al., 2008; Zhaoping & Carroll, 2016). While we

did not formally measure chromatic discrimination thresholds in the present study, it is evident

from comparing the color response matrices in **Figure 4B** that our 543 nm and 680 nm stimuli

could readily be distinguished at most intensities and retinal loci. Elucidating whether the

apparent advantage of L/M symmetry for chromatic sensitivity also extends to chromatic

discrimination is a goal of ongoing work.

An alternative approach to identifying what, if any, L/M cone ratio is optimal for color vision
is to evaluate the degree to which the signals encoded by the cone mosaic can be used to infer
the properties of the distal stimulus. In addition to incorporating contributions from receptor
noise, such reconstruction frameworks enable the role of L/M numerosity and arrangement to
be evaluated in the presence of other factors thought to be important for perceptual inference,
including natural image statistics, physiological optics, and prior visual experience. Using a
Bayesian approach to reconstruct signals from a simple dichromatic retina, Manning & Brainard
(2009) concluded that a spectrally balanced cone mosaic produces optimal estimates of the
retinal image, provided the noise characteristics of the two receptor classes are similar.
Subsequent computational investigations into mosaic optimality found that the advantages
conferred by a symmetrical L/M ratio become less pronounced once longitudinal chromatic
aberrations are incorporated into the simulation (Garrigan et al., 2010; Zhang et al., 2022). The
empirical findings we report here may serve as a useful benchmark for future attempts to
identify the conditions under which the trichromatic cone mosaic does (or does not) constrain
human color perception.
Whether the mosaic-dependent variation in chromatic sensitivity we observed at small
spatial scales has implications for suprathreshold color perception remains uncertain. Prior
studies using large-field stimuli have generally found that certain aspects of color appearance,
including the spectral locus of unique yellow, are invariant to individual differences in L/M ratio
(Brainard et al., 2000; Miyahara et al., 1998; Neitz et al., 2002). This constancy could be
achieved if downstream cone-opponent neurons amplify diminished L-M signals caused by
asymmetric cone ratios. It has been shown that unique yellow can be manipulated by changing
the spectral characteristics of the external environment (Neitz et al., 2002; Welbourne et al.,
2015), lending credence to the notion that color-mechanisms calibrate themselves according to
the statistics of their input signals (Mollon, 2006). Likewise, considerable attention has been
paid in recent years to how such sensory regulation might account for the near-normal
suprathreshold color vision of individuals with anomalous trichromacy (Boehm et al., 2014,
2021; Emery et al., 2023; Knoblauch et al., 2020; Tregillus et al., 2021). If similar processes are
engaged by local variations in the spectral composition of the cone mosaic, one might expect
the topography of color appearance to become more uniform as stimuli become increasingly
suprathreshold.
732
733
734

**Acknowledgements**

The authors thank Austin Roorda and Pavan Tiruveedhula for technical assistance. This work was supported by the Air Force Office of Scientific Research (FA955020-1-0195, FA9550-21-1-0230), National Eye Institute (R01EY023591, R01EY029710, U01EY032055, P30EY003176, P30EY001730, T32EY007043), Alcon Research Institute, Hellman Fellows Program, Burroughs Wellcome Fund Careers at the Scientific Interfaces, and an unrestricted grant from Research to Prevent Blindness.

**Footnote**

[1]Caution should be exercised when drawing links between color discrimination and appearance, which can become increasingly decoupled at small spatial scales. In the case of stimuli landing on a spectrally-homogenous patch of receptors, the Principle of Univariance (Rushton, 1972) dictates that discriminating two equiluminant spectral lights should not be possible. However, this does not necessarily imply that said stimuli will appear achromatic. Indeed, minute cone-sized stimuli delivered to single receptors often appear colorful (Hofer, Singer, et al., 2005; Sabesan et al., 2016), even though the individual cells themselves exhibit univariance (Schnapf et al., 1987) and are commonly considered to be "color-blind".

**References**

Arathorn, D. W., Yang, Q., Vogel, C. R., Zhang, Y., Tiruveedhula, P., & Roorda, A. (2007). Retinally stabilized cone-targeted stimulus delivery. *Optics Express*, *15*(21), 13731–13744. https://doi.org/https://doi.org/10.1364/OE.15.013731

Asano, Y., Fairchild, M. D., & Blondé, L. (2016). Individual Colorimetric Observer Model. *PLOS ONE*, *11*(2), e0145671-. https://doi.org/10.1371/journal.pone.0145671

Atchison, D. A., & Smith, G. (2005). Chromatic dispersions of the ocular media of human eyes. *Journal of the Optical Society of America A*, *22*(1), 29–37. https://doi.org/10.1364/JOSAA.22.000029

Barbur, J. L., Rodriguez-Carmona, M., Harlow, J. A., Mancuso, K., Neitz, J., & Neitz, M. (2008). A study of unusual Rayleigh matches in deutan deficiency. *Visual Neuroscience*, *25*(3), 507–516. https://doi.org/DOI: 10.1017/S0952523808080619

Bieber, M. L., Kraft, J. M., & Werner, J. S. (1998). Effects of known variations in photopigments on L/M cone ratios estimated from luminous efficiency functions. *Vision Research*, *38*(13), 1961–1966. https://doi.org/https://doi.org/10.1016/S0042-6989(97)00302-7

Boehm, A. E., Bosten, J., & MacLeod, D. I. A. (2021). Color discrimination in anomalous trichromacy: Experiment and theory. *Vision Research*, *188*, 85–95. https://doi.org/https://doi.org/10.1016/j.visres.2021.05.011

Boehm, A. E., MacLeod, D. I. A., & Bosten, J. M. (2014). Compensation for red-green contrast
loss in anomalous trichromats. *Journal of Vision*, *14*(13), 19.
https://doi.org/10.1167/14.13.19
Boehm, A. E., Privitera, C. M., Schmidt, B. P., & Roorda, A. (2019). Transverse chromatic
offsets with pupil displacements in the human eye: sources of variability and methods for
real-time correction. *Biomedical Optics Express*, *10*(4), 1691–1706.
https://doi.org/10.1364/BOE.10.001691
Bouman, M. A., & Walraven, P. L. (1957). Some Color Naming Experiments for Red and Green
Monochromatic Lights. *Journal of the Optical Society of America*, *47*(9), 834–839.
https://doi.org/10.1364/JOSA.47.000834
Brainard, D. H., Calderone, J. B., Nugent, A. K., & Jacobs, G. H. (1999). Flicker ERG
Responses to Stimuli Parametrically Modulated in Color Space. *Investigative
Ophthalmology & Visual Science*, *40*(12), 2840–2847.
Brainard, D. H., Roorda, A., Yamauchi, Y., Calderone, J. B., Metha, A., Neitz, M., Neitz, J.,
Williams, D. R., & Jacobs, G. H. (2000). Functional consequences of the relative numbers
of L and M cones. *Journal of the Optical Society of America A*, *17*(3), 607–614.
https://doi.org/10.1364/josaa.17.001684
Carroll, J., Neitz, J., & Neitz, M. (2002). Estimates of L: M cone ratio from ERG flicker
photometry and genetics. *Journal of Vision*, *2*(8), 1.
https://doi.org/https://doi.org/10.1167/2.8.1
Chaparro, A., Stromeyer, C. F., Kronauer, R. E., & Eskew, R. T. (1994). Separable red-green
and luminance detectors for small flashes. *Vision Research*, *34*(6), 751–762.
https://doi.org/https://doi.org/10.1016/0042-6989(94)90214-3
Cicerone, C. M., & Nerger, J. L. (1989). The relative numbers of long-wavelength-sensitive to
middle-wavelength-sensitive cones in the human fovea centralis. *Vision Research*, *29*(1),
115–128. https://doi.org/https://doi.org/10.1016/0042-6989(89)90178-8
CIE. (2006). *CIE 2006 LMS cone fundamentals for 2° field size in terms of energy*.
https://doi.org/10.25039/CIE.DS.tijidesg
Danilova, M. V, Chan, C. H., & Mollon, J. D. (2013). Can spatial resolution reveal individual
differences in the L:M cone ratio? *Vision Research*, *78*, 26–38.
https://doi.org/https://doi.org/10.1016/j.visres.2012.12.006
de Vries, H. L. (1946). Luminosity Curve of Trichromats. *Nature*, *157*(3996), 736–737.
https://doi.org/10.1038/157736b0
de Vries, H. L. (1949). The heredity of the relative numbers of red and green receptors in the
human eye. *Genetica*, *24*(1), 199–212. https://doi.org/10.1007/BF01487206
Emery, K. J., Isherwood, Z. J., & Webster, M. A. (2023). Gaining the system: limits to
compensating color deficiencies through post-receptoral gain changes. *Journal of the
Optical Society of America A*, *40*(3), A16–A25. https://doi.org/10.1364/JOSAA.480035
Field, G. D., Gauthier, J. L., Sher, A., Greschner, M., Machado, T. a., Jepson, L. H., Shlens, J.,
Gunning, D. E., Mathieson, K., Dabrowski, W., Paninski, L., Litke, A. M., & Chichilnisky, E.
J. (2010). Functional connectivity in the retina at the resolution of photoreceptors. *Nature*,
*467*(7316), 673–677. https://doi.org/10.1038/nature09424

Finkelstein, M. A., & Hood, D. C. (1984). Detection and discrimination of small, brief lights: Variable tuning of opponent channels. *Vision Research*, *24*(3), 175–181. https://doi.org/10.1016/0042-6989(84)90119-6

Garrigan, P., Ratliff, C. P., Klein, J. M., Sterling, P., Brainard, D. H., & Balasubramanian, V. (2010). Design of a Trichromatic Cone Array. *PLOS Computational Biology*, *6*(2), e1000677-. https://doi.org/10.1371/journal.pcbi.1000677

Geisler, W. S. (1984). Physical limits of acuity and hyperacuity. *Journal of the Optical Society of America A*, *1*(7), 775–782. https://doi.org/10.1364/JOSAA.1.000775

Greene, M. J., Boehm, A. E., Vanston, J. E., Pandiyan, V. P., Sabesan, R., & Tuten, W. S. (2024). Unique yellow shifts for small and brief stimuli in the central retina. *Journal of Vision*, *24*(6), 2. https://doi.org/10.1167/jov.24.6.2

Gunther, K. L., & Dobkins, K. R. (2002). Individual differences in chromatic (red/green) contrast sensitivity are constrained by the relative number of L- versus M-cones in the eye. *Vision Research*, *42*(11), 1367–1378. https://doi.org/https://doi.org/10.1016/S0042-6989(02)00043-3

Hagstrom, S. A., Neitz, J., & Neitz, M. (1998). Variations in cone populations for red–green color vision examined by analysis of mRNA. *NeuroReport*, *9*(9). https://journals.lww.com/neuroreport/fulltext/1998/06220/variations_in_cone_populations_for_red_green_color.9.aspx

Hagstrom, S. A., Neitz, M., & Neitz, J. (2000). Cone pigment gene expression in individual photoreceptors and the chromatic topography of the retina. *Journal of the Optical Society of America A*, *17*(3), 527–537. https://doi.org/10.1364/JOSAA.17.000527

Harmening, W. M., Tiruveedhula, P., Roorda, A., & Sincich, L. C. (2012). Measurement and correction of transverse chromatic offsets for multi-wavelength retinal microscopy in the living eye. *Biomedical Optics Express*, *3*(9), 2066–2077. https://doi.org/https://doi.org/10.1364/BOE.3.002066

Harmening, W. M., Tuten, W. S., Roorda, A., & Sincich, L. C. (2014). Mapping the perceptual grain of the human retina. *Journal of Neuroscience*, *34*(16). https://doi.org/10.1523/JNEUROSCI.5191-13.2014

He, J., Taveras-Cruz, Y., & Eskew Jr., R. T. (2021). Modeling individual variations in equiluminance settings. *Journal of Vision*, *21*(7), 15. https://doi.org/10.1167/jov.21.7.15

Hofer, H., Carroll, J., Neitz, J., Neitz, M., & Williams, D. R. (2005). Organization of the human trichromatic cone mosaic. *Journal of Neuroscience*, *25*(42), 9669–9679. https://doi.org/10.1523/JNEUROSCI.2414-05.2005

Hofer, H., Singer, B., & Williams, D. R. (2005). Different sensations from cones with the same photopigment. *Journal of Vision*, *5*(5), 444–454. https://doi.org/10:1167/5.5.5

Hood, S. M., Mollon, J. D., Purves, L., & Jordan, G. (2006). Color discrimination in carriers of color deficiency. *Vision Research*, *46*(18), 2894–2900. https://doi.org/https://doi.org/10.1016/j.visres.2006.02.028

King-Smith, P. E., & Carden, D. (1976). Luminance and opponent-color contributions to visual detection and adaptation and to temporal and spatial integration. *Journal of the Optical Society of America*, *66*(7), 709–717. https://doi.org/https://doi.org/10.1364/JOSA.66.000709

Knoblauch, K., Marsh-Armstrong, B., & Werner, J. S. (2020). Suprathreshold contrast response
in normal and anomalous trichromats. *Journal of the Optical Society of America A*, *37*(4),
A133–A144. https://doi.org/10.1364/JOSAA.380088
Kraft, T. W., Neitz, J., & Neitz, M. (1998). Spectra of human L cones. *Vision Research*, *38*(23),
3663–3670. https://doi.org/https://doi.org/10.1016/S0042-6989(97)00371-4
Krauskopf, J. (1964). Color Appearance of Small Stimuli and the Spatial Distribution of Color
Receptors. *Journal of the Optical Society of America*, *54*(9), 1171.
https://doi.org/10.1364/JOSA.54.001171
Krauskopf, J. (2000). Relative number of long- and middle-wavelength-sensitive cones in the
human fovea. *Journal of the Optical Society of America A*, *17*(3), 510–516.
https://doi.org/10.1364/JOSAA.17.000510
Krauskopf, J., & Srebro, R. (1965). Spectral Sensitivity of Color Mechanisms: Derivation from
Fluctuations of Color Appearance near Threshold. *Science*, *150*(3702), 1477–1479.
https://doi.org/10.1126/science.150.3702.1477
Kremers, J., Scholl, H. P. N., Knau, H., Berendschot, T. T. J. M., Usui, T., & Sharpe, L. T.
(2000). L/M cone ratios in human trichromats assessed by psychophysics,
electroretinography, and retinal densitometry. *Journal of the Optical Society of America A*,
*17*(3), 517–526. https://doi.org/https://doi.org/10.1364/JOSAA.17.000517
Li, P. H., Field, G. D., Greschner, M., Ahn, D., Gunning, D. E., Mathieson, K., Sher, A., Litke, A.
M., & Chichilnisky, E. J. (2014). Retinal representation of the elementary visual signal.
*Neuron*, *81*(1), 130–139. https://doi.org/https://doi.org/10.1016/j.neuron.2013.10.043
MacLeod, D. I. A., Williams, D. R., & Makous, W. (1992). A visual nonlinearity fed by single
cones. *Vision Research*, *32*(2), 347–363. https://doi.org/https://doi.org/10.1016/0042-
6989(92)90144-8
Manning, J. R., & Brainard, D. H. (2009). Optimal design of photoreceptor mosaics: Why we do
not see color at night. *Visual Neuroscience*, *26*(1), 5–19. https://doi.org/DOI:
10.1017/S095252380808084X
Miyahara, E., Pokorny, J., Smith, V. C., Baron, R., & Baron, E. (1998). Color vision in two
observers with highly biased LWS/MWS cone ratios. *Vision Research*, *38*(4), 601–612.
https://doi.org/10.1016/S0042-6989(97)88334-4
Mollon, J. D. (2006). Monge: The Verriest Lecture, Lyon, July 2005. *Visual Neuroscience*, *23*(3–
4), 297–309. https://doi.org/DOI: 10.1017/S0952523806233479
Mollon, J. D., & Bowmaker, J. K. (1992). The spatial arrangement of cones in the primate fovea.
*Nature*, *360*(6405), 677–679. https://doi.org/https://doi.org/10.1038/360677a0
Mozaffari, S., LaRocca, F., Jaedicke, V., Tiruveedhula, P., & Roorda, A. (2020). Wide-vergence,
multi-spectral adaptive optics scanning laser ophthalmoscope with diffraction-limited
illumination and collection. *Biomedical Optics Express*, *11*(3), 1617–1632.
https://doi.org/10.1364/BOE.384229
Neitz, A., Jiang, X., Kuchenbecker, J. A., Domdei, N., Harmening, W., Yan, H., Yeonan-Kim, J.,
Patterson, S. S., Neitz, M., Neitz, J., Coates, D. R., & Sabesan, R. (2020). Effect of cone
spectral topography on chromatic detection sensitivity. *Journal of the Optical Society of
America A*, *37*(4), A244–A254. https://doi.org/10.1364/JOSAA.382384

Neitz, J., Carroll, J., Yamauchi, Y., Neitz, M., & Williams, D. R. (2002). Color perception is
mediated by a plastic neural mechanism that is adjustable in adults. *Neuron*, *35*(4), 783–
792. https://doi.org/10.1016/S0896-6273(02)00818-8
Neitz, J., & Jacobs, G. H. (1986). Polymorphism of the long-wavelength cone in normal human
colour vision. *Nature*, *323*(6089), 623–625. https://doi.org/10.1038/323623a0
Neitz, J., & Neitz, M. (2011). The genetics of normal and defective color vision. *Vision
Research*, *51*(7), 633–651. https://doi.org/https://doi.org/10.1016/j.visres.2010.12.002
Otake, S., Gowdy, P. D., & Cicerone, C. M. (2000). The spatial arrangement of L and M cones
in the peripheral human retina. *Vision Research*, *40*(6), 677–693.
https://doi.org/https://doi.org/10.1016/S0042-6989(99)00202-3
Pandiyan, V. P., Maloney-Bertelli, A., Kuchenbecker, J. A., Boyle, K. C., Ling, T., Chen, Z. C.,
Park, B. H., Roorda, A., Palanker, D., & Sabesan, R. (2020). The optoretinogram reveals
the primary steps of phototransduction in the living human eye. *Science Advances*, *6*(37),
eabc1124. https://doi.org/10.1126/sciadv.abc1124
Pandiyan, V. P., Schleufer, S., Slezak, E., Fong, J., Upadhyay, R., Roorda, A., Ng, R., &
Sabesan, R. (2022). Characterizing cone spectral classification by optoretinography.
*Biomedical Optics Express*, *13*(12), 6574–6594. https://doi.org/10.1364/BOE.473608
Roorda, A., Metha, A. B., Lennie, P., & Williams, D. R. (2001). Packing arrangement of the three
cone classes in primate retina. *Vision Research*, *41*(10), 1291–1306.
https://doi.org/https://doi.org/10.1016/S0042-6989(01)00043-8
Roorda, A., Romero-Borja, F., Donnelly III, W. J., Queener, H., Hebert, T. J., & Campbell, M. C.
W. (2002). Adaptive optics scanning laser ophthalmoscopy. *Optics Express*, *10*(9), 405–
412. https://doi.org/https://doi.org/10.1364/OE.10.000405
Roorda, A., & Williams, D. R. (1999). The arrangement of the three cone classes in the living
human eye. *Nature*, *397*(6719), 520–522. https://doi.org/10.1038/17383
Rushton, W. A. H. (1972). Review lecture. Pigments and signals in colour vision. *The Journal of
Physiology*, *220*(3), 1–31. https://doi.org/https://doi.org/10.1113/jphysiol.1972.sp009719
Rushton, W. A. H., & Baker, H. D. (1964). Red/green sensitivity in normal vision. *Vision
Research*, *4*(1), 75–85. https://doi.org/https://doi.org/10.1016/0042-6989(64)90034-3
Sabesan, R., Schmidt, B. P., Tuten, W. S., & Roorda, A. (2016). The elementary representation
of spatial and color vision in the human retina. *Science Advances*, *2*(9).
https://doi.org/10.1126/sciadv.1600797
Schmidt, B. P., Boehm, A. E., Foote, K. G., & Roorda, A. (2018). The spectral identity of foveal
cones is preserved in hue perception. *Journal of Vision*, *18*(11), 19.
https://doi.org/https://doi.org/10.1167/18.11.19
Schmidt, B. P., Boehm, A. E., Tuten, W. S., & Roorda, A. (2019). Spatial summation of
individual cones in human color vision. *PLoS ONE*, *14*(7).
https://doi.org/10.1371/journal.pone.0211397
Schnapf, J. L., Kraft, T. W., & Baylor, D. A. (1987). Spectral sensitivity of human cone
photoreceptors. *Nature*, *325*(6103), 439–441. https://doi.org/10.1038/325439a0
Shi, K., Luo, M. R., Rider, A. T., Huang, T., Xu, L., & Stockman, A. (2024). A multi-primary
trichromator to derive individual color matching functions and cone spectral sensitivities.
*Color Research & Application*, *49*(5), 449–464.
https://doi.org/https://doi.org/10.1002/col.22928

Stockman, A., Jägle, H., Pirzer, M., & Sharpe, L. T. (2008). The dependence of luminous
efficiency on chromatic adaptation. *Journal of Vision*, *8*(16), 1.
https://doi.org/10.1167/8.16.1
Stockman, A., & Sharpe, L. T. (2000). The spectral sensitivities of the middle- and long-
wavelength-sensitive cones derived from measurements in observers of known genotype.
*Vision Research*, *40*(13), 1711–1737. https://doi.org/https://doi.org/10.1016/S0042-
6989(00)00021-3
Thomas, P. B. M., & Mollon, J. D. (2004). Modelling the Rayleigh match. *Visual Neuroscience*,
*21*(3), 477–482. https://doi.org/DOI: 10.1017/S095252380421344X
Tregillus, K. E. M., Isherwood, Z. J., Vanston, J. E., Engel, S. A., MacLeod, D. I. A., Kuriki, I., &
Webster, M. A. (2021). Color Compensation in Anomalous Trichromats Assessed with
fMRI. *Current Biology*, *31*(5), 936-942.e4. https://doi.org/10.1016/j.cub.2020.11.039
Vimal, R. L. P., Pokorny, J., Smith, V. C., & Shevell, S. K. (1989). Foveal cone thresholds.
*Vision Research*, *29*(1), 61–78. https://doi.org/https://doi.org/10.1016/0042-6989(89)90174-
0
Vorobyev, M., & Osorio, D. (1998). Receptor noise as a determinant of colour thresholds.
*Proceedings of the Royal Society of London. Series B: Biological Sciences*, *265*(1394),
351–358. https://doi.org/https://doi.org/10.1098/rspb.1998.0302
Watson, A. B. (2017). QUEST+: A general multidimensional Bayesian adaptive psychometric
method. *Journal of Vision*, *17*(3), 10. https://doi.org/10.1167/17.3.10
Welbourne, L. E., Morland, A. B., & Wade, A. R. (2015). Human colour perception changes
between seasons. *Current Biology*, *25*(15), R646–R647.
https://doi.org/https://doi.org/10.1016/j.cub.2015.06.030
Williams, D. R., MacLeod, D. I. A., & Hayhoe, M. M. (1981). Punctate sensitivity of the blue-
sensitive mechanism. *Vision Research*, *21*(9), 1357–1375.
https://doi.org/https://doi.org/10.1016/0042-6989(81)90242-X
Yang, Q., Arathorn, D. W., Tiruveedhula, P., Vogel, C. R., & Roorda, A. (2010). Design of an
integrated hardware interface for AOSLO image capture and cone-targeted stimulus
delivery. *Optics Express*, *18*(17), 17841–17858.
https://doi.org/https://doi.org/10.1364/OE.18.017841
Zhang, F., Kurokawa, K., Lassoued, A., Crowell, J. A., & Miller, D. T. (2019). Cone
photoreceptor classification in the living human eye from photostimulation-induced phase
dynamics. *Proceedings of the National Academy of Sciences*, *116*(16), 7951–7956.
https://doi.org/https://doi.org/10.1073/pnas.1816360116
Zhang, L. Q., Cottaris, N. P., & Brainard, D. H. (2022). An image reconstruction framework for
characterizing initial visual encoding. *eLife*, *11*, 1–30. https://doi.org/10.7554/eLife.71132
Zhaoping, L., & Carroll, J. (2016). An analytical model of the influence of cone sensitivity and
numerosity on the Rayleigh match. *Journal of the Optical Society of America A*, *33*(3),
A228–A237. https://doi.org/10.1364/JOSAA.33.00A228
981