1 Titles

2 Chromatin retained *MUSHER* lncRNA integrates ABA and DOG1 signalling pathways to enhance
3 Arabidopsis seeds dormancy.

4 Authors

Sebastian Przemyslaw Sacharowski[1], Michal Krzyszton[1]*, Lien Brzezniak[1]*, Karol Jerzy
Rogowski[2], Miguel Montez[3], Katarzyna Rosol[1], Magdalena Wrona[1], Ruslan Yatusevich[1], Veena
Halale Manjunath[1], Lukasz Szewc[4], Artur Jarmolowski[4], Szymon Swiezewski[1]

1. Laboratory of Seed Molecular Biology, Institute of Biochemistry and Biophysics PAS, Warsaw 02-106,
Poland
2. ElementZero Biolabs, Invalidenstraße 65 C/O Edge, Berlin, Germany
3. Department of Cell and Developmental Biology, John Innes Centre, Norwich Research Park, Norwich NR4
7UH, UK
4. Department of Gene Expression, Institute of Molecular Biology and Biotechnology, Faculty of Biology,
Adam Mickiewicz University, Poznan 61-614, Poland

*Equal contribution

16 Correspondence

Szymon Świeżewski, sswiez@ibb.waw.pl,

ORCID:
Sebastian Sacharowski   0000-0001-8220-9649
Michal Krzyszton   0000-0002-9795-5588
Lien Brzezniak   0000-0002-6932-413X
Karol Jerzy Rogowski   0000-0003-4560-0715
Miguel Montez   0000-0002-1383-0885
Katarzyna Rosol   0000-0001-6663-9884
Magda Wrona   0000-0001-9619-9664
Ruslan Yatusevich   0000-0001-5382-4079
Veena Halale Manjunath   0009-0006-1514-9379
Lukasz Szewc   0000-0002-2219-6513
Artur Jarmolowski   0000-0002-1073-146X
Szymon Swiezewski   0000-0002-2007-2779
31

32 Abstract

Many plant lncRNAs regulate gene expression by binding to chromatin, but how they are
retained at the target *loci* is unclear. We identify a new, chromatin-localized lncRNA - *MUSHER*,
which activates two parallel regulatory pathways to increase Arabidopsis seed dormancy.
MUSHER is upregulated in response to high temperatures, contributing to the induction of
secondary dormancy. It promotes *DOG1* expression by recruitment of the CPSF complex to
enhance the proximal cleavage and polyadenylation at the *DOG1* gene. It also increases ABA
sensitivity in seeds by activating *PIR1* gene transcription. These genes, located on different
chromosomes, are both bound by *MUSHER,* despite lacking sequence homology. The
chromatin association of *MUSHER* enables the integration of the DOG1- and ABA pathways to
adjust seed germination timing. Additionally, *MUSHER* and other lncRNAs interact with U1
snRNP, which is required for their chromatin localisation, revealing a novel function of U1
snRNP in plants.

Introduction

Multiple long non-coding RNA (lncRNA) are implicated in gene expression regulation in the nucleus by binding to chromatin[1]. Well-known examples of chromatin-bound RNA include *XIST* spreading on the whole X-chromosome and involved in its inactivation in mammals[2,3], and *MALAT1*, tightly bound to chromatin and involved in gene expression and nuclear architecture regulation in humans[4]. There are also many reports of plant lncRNA functioning at the chromatin. This includes *APOLO*, involved in remodelling chromatin structure[5], *SVALKA* proposed to act through induction of RNA polymerase II (Pol II) transcriptional collisions[6] and *PUPPIES*, a lncRNA with a role in delaying Arabidopsis seed germination by controlling Pol II pausing at target gene[7]. *APOLO* was shown to be retained at *trans*-target *loci* via the formation of R-loops[5]. Probably the best-described plant lncRNA *COOLAIR*, involved in flowering time regulation[8,9], has been shown to form a cloud around its locus[10]. Many lncRNAs functioning in the nucleus are involved in environmental sensing in plants. *COOLAIR* regulates vernalisation[9], SVALKA is implicated in cold acclimation in seedlings[6], *ARTA* enhances ABA response[11], *1GOD* affects drought resistance in mature plants[12], *PUPPIES* regulates seed response to salt[7] and *HID1* promotes germination[13]. While an RNA structural element strongly influences *COOLAIR* chromatin association[14], a broader mechanism for lncRNA retention at chromatin remains elusive.

Recently, a U1-based mechanism of human and mouse lncRNA retention at the chromatin was reported[15]. This is based on the physical binding of U1 to a large fraction of lncRNAs[15]. U1 is a highly conserved, small nuclear RNA (snRNA) that, together with conserved proteins including U1C and U1K70, forms the U1 snRNP[16]. U1 snRNP is primarily involved in the first step of splicing by recognising 5' splice sites[17]. However, U1 is also engaged in several other chromatin-related activities: enhancement of Pol II promoters directionality[18]; boosting the speed of RNA polymerase II (Pol II)[19]; and, conserved in plants, suppression of premature cleavage and polyadenylation in introns, known as telescripting[20,21]. Whether U1 snRNP plays a role in chromatin retention in plants is unknown.

Here, we leverage seed biology regulation to explore the functions of lncRNAs. Two key players in seed dormancy are plant hormone abscisic acid (ABA) and a plant-specific protein – DELAY OF GERMINATION 1 (DOG1). Both DOG1 and ABA inhibit seed germination by promoting dormancy, an ability to postpone germination despite favourable external conditions[22]. This allows seeds to ignore short spells of permissive conditions and align germination with seasonal clues. Seed dormancy is initially established when seeds develop on the mother plant[22]. Once this dormancy, known as primary, is released, exposure of imbibed seeds to unfavourable conditions can result in the establishment of secondary seed dormancy[23]. In many plant species, including Arabidopsis, temperature is a key environmental factor required for the efficient establishment of secondary dormancy[24].

DOG1 protein has been suggested to act as part of a thermal-sensing mechanism inducing secondary dormancy[25], but the precise mechanism of how temperature is integrated into the *DOG1* gene regulation during secondary dormancy establishment is not clear[22]. *DOG1* expression is positively correlated with dormancy strength among Arabidopsis accessions, and it has been identified as a major QTL and GWAS for seed dormancy[26,27]. *DOG1* transcription is extensively regulated, including by two independent lncRNA. *DOG1* expression is suppressed

by an antisense transcript *1GOD*, and activated in response to salt by *PUPPIES*, a lncRNA expressed in the *DOG1* promoter[7,28]. In addition, the selection of the *DOG1* proximal polyA site results in the production of *shDOG1* (*short DOG1*), the predominant and functional form of the *DOG1* transcript, while the selection of the distal polyA site results in a presumably non-functional and much less abundant *lgDOG1* (*long DOG1*) transcript[29–31]. Multiple regulators of *DOG1* proximal/distal polyA site selection have been described[30,32], but *DOG1* alternative polyadenylation has so far not been linked to environmental responses.

DOG1 expression is, to a large extent, unaffected by ABA levels, and *dog1* mutations only marginally affect ABA levels in seeds[33]. Interestingly, seed sensitivity to ABA can be modified by both environmental factors and mutations in genes involved in the ABA signalling pathway, without affecting basal ABA levels[34]. ABA is sensed in plants by ABA receptors PYR/PYLs, which once bound by ABA, bind to and inactivate several PP2C phosphatases, including PP2CA[35]. One way plants can modify ABA sensitivity is by interfering with the ABA signal transduction pathway. For example, PP2CA interacting RING finger protein 1 (PIR1) and (PIR2) have been shown to mediate PP2CA degradation to enhance abscisic acid response[36]. Although *dog1* mutants do not exhibit altered ABA sensitivity[27], DOG1 protein has been shown to interact with and inhibit the enzymatic activity of PP2CA *in vitro*[37,38]. In summary, DOG1 and ABA pathways are closely intertwined at the protein level, but genetic and transcriptomic analyses indicate that DOG1 function independently from ABA[22,27,33].

Here, we describe *MUSHER*, a novel lncRNA that integrates ABA and DOG1 pathways on the transcriptional level by enhancing *DOG1* and *PIR1* expression. Through genetic analyses, we demonstrate that *MUSHER* acts as a positive regulator of dormancy. Our findings imply that *MUSHER* responds to increased temperature and is essential for high-temperature-induced secondary dormancy in Arabidopsis. At the molecular level, *MUSHER* promotes *DOG1* mRNA proximal polyadenylation by recruiting the CPSF complex. Additionally, through RNA Pol II recruitment, *MUSHER* facilitates the expression of *PIR1 in trans*, thereby enforcing ABA sensitivity. Finally, we show that *MUSHER*, as well as *COOLAIR* and *PUPPIES*, interact with chromatin through U1 snRNP, suggesting the existence of a conserved mechanism of lncRNA-chromatin retention across eukaryotes.

Results

Characterisation of *MUSHER* lncRNA as a regulator of *DOG1* gene expression and seed germination.

To identify new lncRNA controlling dormancy, we performed RT-qPCR analysis with primers spanning the *DOG1* locus in WT and *hen2* mutant seeds (Figs. 1a,b). HEN2 is an RNA helicase that interacts with the exosome complex responsible for the nuclear 3' to 5' degradation of various ncRNAs[39], and its mutation provides a good background to identify unstable lncRNA species[40,41]. We found a *hen2*-dependent signal near the *DOG1* gene 3'end, pointing to transcription of a previously uncharacterised transcript targeted by the exosome in seeds (Fig. 1b). We did not detect increased signals in regions corresponding to *PUPPIES* and *1GOD* transcripts [7,28], indicating that these *DOG1* locus-originating lncRNAs are not targeted by the exosome in seeds. 5' and 3'RACE-seq analyses (Fig. 1c) suggested the existence of an intergenic transcriptional unit between the *SnRK3.6* and *DOG1* genes. Due to its location downstream of *DOG1*, we named this putative lncRNA *MUSHER*.

Measuring this ncRNA level throughout seed development (Fig. 1g) revealed that *MUSHER*
transcripts accumulate during seed maturation concomitant to *DOG1* induction. To disrupt the
MUSHER transcriptional unit, we used two T-DNA insertion lines, *msh-1* and *msh-2* (Fig. 1a), a
Cas9 deletion mutant *Δmsh,* and two additional lines: *msh_dCas9-1* and *msh_dCas9-2* with
catalytically inactive Cas9 (dCas9). dCas9 acts as a transcriptional repressor by binding to target
DNA without inducing cleavage[42]. T-DNA insertion and dCas9 lines, but not the dCas9 controls
targeting two intergenomic regions localised 100 kb from *DOG1 locus* (Supplementary Fig. 1b)
showed reduced *MUSHER* levels in seeds (Supplementary Fig. 1a). Unexpectedly, *Δmsh* despite
deletion of mapped *MUSHER* start site showed ectopic expression an antisense transcript
downstream of *DOG1* (Supplementary Fig. 1c and d). Despite this, all tested *musher* mutants
demonstrated reduced *DOG1* expression in seeds and increased seed germination (Figs. 1e,f
and Supplementary Fig. 1e). The impact of *msh_dCas9-1* on *DOG1* expression is observed
throughout seed maturation (Fig. 1f). Because of the observed *MUSHER*-like transcripts in *Δmsh*
we decided to exclude this line from downstream analysis.
Since *MUSHER* and *DOG1* are co-expressed during seed maturation (Fig. 1g), we questioned
whether *DOG1* expression can have a feedback effect on *MUSHER* transcription. We observed
reduced *MUSHER* expression in *dog1* insertional mutants, *dog1-3* and *dog1-4,* but no effect in
a *DOG1* large deletion mutant generated using CRISPR/Cas9, dog*1-Δ19* (Supplementary Fig. 1a).
This suggested that while *MUSHER* regulates *DOG1* expression, *MUSHER* is not dependent on
DOG1 expression. We speculated that the downregulation of *MUSHER* expression in *dog1*
insertional mutants could result from local chromatin state changes caused by T-DNA
insertion[43]. Furthermore, *MUSHER* expression was unaffected by mutations in the downstream
gene *SnRK3.6* (Supplementary Fig. 1a), confirming that *MUSHER* is an independent
transcriptional unit.

Fig. 1 | *MUSHER* lncRNA acts as a positive regulator of seed dormancy and *DOG1* expression.

a Schematic representation of the *DOG1-MUSHER* genomic region of chromosome 5 in *A. thaliana*. The direction of transcription is indicated with arrowheads. Grey rectangles represent coding exons and introns with thick black lines. The green arrow represents the newly annotated *MUSHER* transcriptional unit. Numbers along the x-axis on the top indicate genome coordinates (TAIR version 10). Triangles represent T-DNA insertions of two T-DNA lines (*msh-1* and *msh-2*). The green vertical lines show the positions of multiple guides for *msh_dCas9-1* and *msh_dCas9-2* dCas9 lines. The green horizontal line corresponds to the deleted region of MUSHER using Cas9 *(Δmsh)*. **b** Primer walking RT-qPCR analysis across the *DOG1* locus. The position of amplicons corresponds to the *DOG1* locus representation above. Expression levels in the *hen2* mutant were normalised to WT and *UBC21*. Error bars represent the standard deviation of four biological replicates. **c** Positions of *MUSHER* transcript ends found in WT seeds using 3' and 5' RACE-seq, coloured in dark and light green, respectively. **d** Expression profiles of *DOG1* and *MUSHER* transcripts during seed maturation in the WT. Expression levels were normalised to *UBC21* mRNA and 16 DAP. Error bars

represent the standard deviation of three biological replicates. **e** *DOG1* relative expression levels, *musher* mutants relative to WT in freshly harvested seeds, normalised to *UBC21*. *p < 0.05 from two-tailed Student's *t*-test. **f** Freshly harvested seeds of WT and different *musher* mutants were scored for germination. Germination of tested seeds was scored as radical protrusion and counted 4d after sowing. A two-tailed *t-test p*-value was calculated compared to WT (*p < 0.05). **g** *DOG1* expression profile during silique development in WT and *msh_dCas9-1* plants. Expression levels were normalised to *UBC21* and shown at the indicated number of days after pollination (DAP); error bars represent the standard deviation of three biological replicates. **h** Subcellular localisation of selected transcripts in maturing WT Arabidopsis seeds. Each transcript was tested in 5 biological replicates represented as separate bars. Black horizontal lines represent the mean of chromatin and nucleoplasmic fractions contribution of the total amount of indicated transcript.

Next, we tested if *MUSHER* accumulates in the cytoplasm or chromatin, which could provide insight into its mode of action. To test *MUSHER* localisation, we used developing seeds as seed dormancy is established during seed maturation, as a result of the expression of *DOG1* and other genes. We isolated cytoplasmic, nucleoplasmic and chromatin fractions and performed RT-qPCR analysis. Mitochondrial and chloroplast-localised RNA, used as negative controls, are nearly exclusively found in the cytoplasmic fraction. Several tested nuclear-encoded mRNAs showed similar levels in cytoplasm, nucleoplasm and chromatin. U6, a spliceosome component, was detected mainly in chromatin and nucleoplasmic fractions. *MUSHER*, *CHR1*- a transposon-derived RNA and U3.3, involved in co-transcriptional pre-rRNA modification, were predominantly detected in chromatin fraction (Fig. 1h).

In summary, our data showed that *MUSHER* is a lncRNA transcribed downstream of the *DOG1* gene that increases *DOG1* mRNA levels to delay the germination of freshly harvested Arabidopsis seeds. *MUSHER* chromatin association suggests that it may control *DOG1* expression by binding to its chromatin.

Supplementary Fig. 1| *MUSHER* expression is partially dependent on *DOG1*.

a *MUSHER* relative expression level in developing seeds of WT and indicated mutants, normalised to *UBC21* and relative to WT. *p*-value was calculated in comparison with WT. *p < 0.05. Error bars represent the standard deviation of four biological replicates. **b** *MUSHER* relative expression level in freshly harvested seeds of WT and dCas9 mutants, normalised to *UBC21* and relative to WT. *p*-value was calculated in comparison with WT. *p < 0.05. Error bars represent the standard deviation of four biological replicates. *MUSHER*-associated 2nd (c) and 3rd (d) transcript relative expression level in WT and indicated mutants seeds, normalised to *UBC21* and relative to WT. *p*-value was calculated in comparison with WT. *p < 0.05. Error bars represent

254 the standard deviation of four biological replicates. e Freshly harvested seeds of WT and Δ*musher*
255 mutant were scored for germination. **e** Germination of tested seeds was scored as radical protrusion
256 and counted 4d after sowing. A two-tailed *t-test p*-value was calculated compared to WT (*p < 0.05). **f**
257 *DOG1* relative expression levels in freshly harvested seeds of Δ*musher* mutant, relative to WT,
258 normalised to *UBC21*. *p < 0.05 from two-tailed Student's *t*-test.

MUSHER lncRNA is required for *DOG1* regulation by CPL1, a Pol II phosphatase

We showed previously that RNA Polymerase II Carboxyl Terminal Domain (CTD) Phosphatase-
Like1 (CPL1)-mediated suppression of *DOG1* expression (Supplementary Fig. 2a) requires the 3'
end of the *DOG1* locus but not the antisense *1GOD* lncRNA expression from this region[32].
Consequently, the molecular mechanism of CPL1 control of *DOG1* expression remains
incompletely understood. Given the effect of *MUSHER* on *DOG1* expression and dormancy, we
speculated that CPL1 could act through *MUSHER* regulation. Indeed, in the *cpl1-7* mutant, the
MUSHER transcript level is elevated (Supplementary Fig. 2b), suggesting that *cpl1* is a *MUSHER*
lncRNA over-expressor.

To test the requirement of *MUSHER* for CPL1's effect on *DOG1*, we constructed a full-length
DOG1-luciferase (LUC) fusion with or without *MUSHER* (*pDOG1::LUC-DOG1*)[30] (*pDOG1::LUC-
DOG1ΔMUSHER*). Transient expression in Arabidopsis seedlings showed that *MUSHER* deletion
suppressed *DOG1* upregulation in the *cpl1* plants (Supplementary Fig. 2c).

273
Supplementary Fig. 2| *MUSHER* overexpression activates *DOG1*.

a *DOG1* and **b** *MUSHER* relative expression level in seeds of WT and *cpl1-7* mutant, normalised to *UBC21*
and relative to WT. *p*-value was calculated in comparison with WT. *p*-value < 0.05. Error bars represent
the standard deviation of four biological replicates. **c** Luciferase (LUC) gene reporter assay was used to
measure *MUSHER's* ability to activate the *DOG1* gene expression. Schematic diagram showing regions
included in *LUC* gene fusion constructs used for transient transformation of Arabidopsis WT and *cpl1-7*
mutant seedlings. In boxplots, middle lines show the medians (solid line) and mean (dotted line); box
limits indicate the 25th and 75th percentiles; whiskers extend 1.5 times the interquartile ranges. * *p*-
value < 0.05. **d** Luciferase activity assay on transgenic seeds containing the *LUC* gene fused with the
MUSHER promoter. In boxplots, middle lines show the median (solid line) and mean (dotted line); box
limits indicate the 25th and 75th percentiles; whiskers extend 1.5 times the interquartile ranges. * *p*-

value < 0.01. **e** The ratio of *shDOG1* and *lgDOG1* isoforms in freshly harvested seeds of WT and *msh-1,*
cpl1-7 and double *msh-1 cpl1-7* mutants, normalised to *UBC21* and relative to WT. *p*-value from the two-
tailed t-test, **p*-value < 0.05. Error bars show standard deviations.

To cross-validate this, we checked the effect of CPL1 on *MUSHER* locus activity. We generated
transgenic plants expressing an internal ribosomal entry site-luciferase gene (IRES-LUC) reporter
cassette fused with *MUSHER* and crossed them to *cpl1-7* mutant. The resulting
pMUSHER::MUSHER-IRES-LUC showed stronger activity in *cpl1-7* when compared to WT,
indicating that *MUSHER* is a direct target of CPL1 (Supplementary Fig. 2d). Our data showed
that *MUSHER* lncRNA is required for CPL1 to control *DOG1* expression.
Interestingly, inactivation of CPL1 led to preferential expression of proximally over distally
polyadenylated *DOG1* mRNA isoform, known respectively as *shDOG1* (*short DOG1*) and *lgDOG1*
(*long DOG1*) (Supplementary Fig. 2e). Conversely, *MUSHER* inactivation resulted in reduced
shDOG1 isoform (Supplementary Fig. 2e). Thus, *MUSHER* and CPL1 showed antagonistic effect
on the production of *shDOG1*, encoding functional *DOG1* protein and complementing the *dog1*
mutant[30]. Importantly, mutating the *cpl1* gene in the *msh* background did not reverse *lgDOG1*
polyadenylation preference (Supplementary Fig. 2e), suggesting *DOG1* polyA site selection is
directly affected by *MUSHER* and not by CPL1 protein. Taken together, our data showed that
CPL1 indirectly affects *DOG1* polyadenylation site selection, by suppressing *MUSHER* expression
and activity.

Fig. 2| *MUSHER* enhances *DOG1* proximal polyA site selection.

a Snapshot of 3'RNA-seq reads for *DOG1* locus in WT and *msh-1* mutant (n=4). Schematic diagram showing *DOG1* gene: exons (grey rectangles), *shDOG1* 3'UTR region (light grey arrow), *lgDOG1* 3'UTR region (grey arrow), and *MUSHER* (green). The regions used to estimate the expression of mRNA isoforms are shown below chromosome coordinates. **b** Ratio of reads corresponding to short and long *DOG1* mRNA isoforms calculated for regions marked on the schematic. Data are the mean of four biological replicates; error bars show standard deviations. **c** The ratio of *shDOG1* and *lgDOG1* isoforms in freshly harvested seeds of WT and *musher* mutants, normalised to *UBC21* and relative to WT (n=4, **p*-value < 0.05). **d** CPSF73 occupancy was measured by ChIP-qPCR in WT, *msh-1* and *msh_dCas9-1* mutants. (n=3). **e** The ratio of *shDOG1* and *lgDOG1* isoforms in seeds of WT and *msh-1, esp1-2* and double *msh-1 esp1-2* mutants, normalised to *UBC21* and relative to WT (n=5, **p*-value < 0.05). **f** Pol II occupancy was measured by ChIP-qPCR in WT and

msh-1 mutant. (n=3). **g** *MUSHER* relative expression level in seeds of WT and *esp1-2*, normalised to
UBC21 and relative to WT (n=5, ns- not significant). **h** H3, **i** H3K4me3 and **j** H3K27me3 levels were
measured by ChIP-qPCR in WT and *msh-1* mutant. H3K4me3 and H3K27me3 values were normalised to
H3 and to *ACTIN7* (*AT5G09810*) gene body. For all ChIP-qPCR experiments presented in this figure: *p*-
value from the two-tailed t-test, *p*-value < 0.05, ns- not significant. Error bars represent the standard
deviation.

MUSHER promotes *DOG1* proximal polyadenylation by recruiting the CPSF complex

We used 3'RNA-seq to confirm that *shDOG1* is decreased in *msh-1* seeds, while *lgDOG1* remains
unchanged when compared to WT (Fig. 2a). Calculation of 3'RNA-seq read density assigned to
DOG1 isoforms revealed a significant reduction of *shDOG1/lgDOG1* ratio in *msh-1* (Fig. 2b).
Subsequent RT-qPCR analysis confirmed that *shDOG1/lgDOG1* ratio is significantly reduced in
all tested *musher* mutants (Fig. 2c).
We previously reported that mutations in pre-mRNA cleavage and polyadenylation factors result
in reduced *shDOG1* but not *lgDOG1* expression[30], similar to *MUSHER* knockouts shown here. To
confirm changes in *DOG1* mRNA 3'end formation, we performed the ChIP-qPCR analysis of the
CPSF73, a subunit of the Cleavage Stimulatory Factor complex[44]. We observed a localised
decrease in CPSF73 occupancy at the *DOG1* proximal but not distal termination site in *msh-1*
and *msh_dCas9-1* mutants (Fig. 2d). Because *cpsf73* mutants are sterile[44], for genetic analysis,
we used a mutant in an auxiliary component of Cleavage Stimulatory Factor complex encoding
ESP1 (CSTF64L) protein[45]. Analysis of the *msh-1 esp1-2* double mutant showed no additivity in
DOG1 termination site selection defects (Figs. 2e,f). This is supported by no change of *MUSHER*
expression in the *esp1* mutant (Fig. 2g). Our results implied that *MUSHER* increases *DOG1*
expression by enhancing CPSF complex recruitment to the proximal termination site, as
supported by findings in *cpl1-7* mutant showing *MUSHER* overexpression and enhanced
proximal polyA site selection (Supplementary Fig. 2).
Transcription termination can be coupled with Pol II occupancy changes[46]. We, therefore,
profiled Pol II occupancy along the *DOG1* locus in the *musher* mutant. This revealed a significant
decrease of Pol II level at the *DOG1* proximal termination region but not at the *DOG1*
transcription start site (TSS) or the distal polyA site (Fig. 2f). This suggested that *MUSHER*-
promoted proximal polyA site selection is not accompanied by changes in *DOG1* transcription
initiation but rather represents a result of *MUSHER* direct activity at *shDOG1* gene 3'end.
The localised decrease of Pol II and CPSF complexes occupancy at the *DOG1* proximal
termination site indicates that *MUSHER* promotes the proximal termination site selection rather
than by controlling transcription levels. To test this hypothesis, we applied single-molecule RNA
FISH (smFISH) to measure the level of *DOG1* mRNA transcripts and analyse transcription kinetics
by quantifying the intensity of *DOG1* transcription[7]. The brightness of dots identified as
transcription sites is generally considered to correlate with the number of RNA molecules
present at transcription sites, while the frequency of transcription sites reflects the proportion
of cells exhibiting detectable transcription activity[7]. We collected embryos from WT and *msh-1*
siliques at ten-time points, corresponding to the experiments showing dynamics of *DOG1* and
MUSHER expression during seed maturation (Fig. 1g). *DOG1* smFISH showed a consistent and
significant reduction in the number of both nuclear and cytoplasmic *foci* during seed
development in *msh-1* seeds when compared to WT (Supplementary Figs.3a,b). We observed
an increase in *DOG1* transcript number in WT siliques between the 16th and 21st day, which was
diminished in *msh-1* (Supplementary Figs.3a,b). Notably, the *msh* mutant showed no significant
change in the intensity of nuclear dots identified as transcription sites (Supplementary Fig. 3c)[7]

and only minor changes in the frequency of cells with active *DOG1* transcription
(Supplementary Fig. 3d). We therefore conclude that in agreement with Pol II ChIP (Fig. 2f),
smFISH indicated no major changes in *DOG1* transcription initiation in *msh-1* seeds.

Supplementary Fig. 3| *MUSHER* promotes *DOG1* mRNA abundance in the nucleus and cytoplasm but not burst frequency or intensity.
a Nuclear *DOG1* mRNA foci number per cell during silique development in WT and *msh-1.* **b** Cytoplasmic *DOG1* mRNA foci number per cell during silique development in WT and *msh-1.* **c** Intensity of *foci* corresponding to *DOG1* transcription sites (TS) during silique development in WT and *msh-1. Foci* with a fold-change ≥3 times the average intensity of all nuclear foci, were considered as TS. **d** Frequency of *DOG1* TS during silique development in WT and *msh-1.* For all box plots, the middle line represents the median; the upper and lower lines are the first and third quartiles (Q1 and Q3); whiskers extend 1.5 times the interquartile ranges. The x-axis represents days after pollination (DAP).

Our data suggest that *MUSHER* promotes *DOG1* proximal termination by locally affecting
cleavage and polyadenylation complexes and Pol II occupancy at the proximal transcription
termination site without affecting *DOG1* transcription levels. All modes of RNA Pol II activity,
including transcription termination, are accompanied and modulated by histone
posttranslational modifications. On the contrary, some lncRNAs may recruit specific histone
modelling complexes, including PRC2 involved in H3K27me3 deposition[47,48]. To address the
effects of *MUSHER* on the *DOG1* chromatin landscape, we analysed selected active (H3K4me3)
or repressive (H3K27me3) chromatin histone marks along the *DOG1* locus. Using ChIP-qPCR, we
detected no significant difference in both tested histone modification levels in the *msh-1*
mutant throughout the *DOG1* locus (Figs. 2h,j) compared to WT. This result showed that defects
at the proximal transcription termination site (TTS) are unrelated to chromatin modifications
and agree with no changes in Pol II recruitment for most of the *DOG1* locus. In summary, we
propose a model in which *MUSHER* lncRNA enhances *DOG1* proximal termination rather than
control Pol II recruitment.

MUSHER enhances secondary dormancy in response to high-temperature
DOG1 is a known regulator of both primary and secondary dormancy[49,50]. Based on the weak
dormancy in *musher* mutants and the requirement for high *DOG1* expression during primary
seed dormancy establishment (Fig.1), we hypothesised that *MUSHER* may also be involved in
secondary dormancy regulation. We used a well-established secondary dormancy induction
protocol that consists of prolonged incubation in the darkness at high temperatures followed

by germination analysis in permissive conditions[49]. In agreement with defects in primary dormancy, *msh-1* and *msh_dCas9-1* showed weaker secondary seed dormancy, and *cpl1-7*, a mutant with high *MUSHER* expression, showed stronger secondary dormancy when compared to WT (Figs.3a,b). Consistent with these phenotypes, we observed the down- and up-regulation of the *shDOG1* transcript in *musher* and *cpl1-7* mutants, respectively (Figs.3c,d).

Fig. 3| *MUSHER* activates *DOG1* expression during secondary dormancy induction in response to high temperature.

a Seed germination of WT and different *musher* mutants after 7 days of secondary dormancy (SD) induction at 35°C. Germination frequency was scored after 5 days. Error bars show the standard deviation of four biological replicates. *p < 0.05. **b** Seed germination of WT and *cpl1-7* mutant seeds after 7 days of SD induction. Germination was scored after 5 days. Error bars show the standard deviation of five biological replicates. *p < 0.05. **c** The expression profile of *DOG1* was analysed using RT-qPCR in seeds of WT, *msh-1* and *msh_dCas9-1* subjected to secondary dormancy induction. Error bars show the standard deviation of four biological replicates. *UBC21* mRNA was used as a reference gene. **d** The *DOG1* relative expression level in seeds of WT and *cpl1-7* mutant, subjected to SD induction. Data normalised to *UBC21* and related to WT. p-value from the two-tailed *t*-test, *p-value < 0.05. Error bars show standard deviations. **e** *DOG1* and **f** *MUSHER* expression profile in after-ripened seeds subjected to secondary dormancy induction at 25°C and 35°C. Expression was normalised to *UBC21* mRNA and dry seed levels. Bars represent the standard deviations from at least four biological replicates. **g** Schematic representation of constructs introduced to transgenic plants. RT-qPCR analysis of luciferase mRNA levels in seeds subjected to secondary dormancy induction under temperatures indicated on the right. *LUC* expression level measured after 5 days of SD induction, normalised to *UBC21* and related to 25°C for each construct. **h** Long and short *DOG1* expression analysis on the 5th day of secondary dormancy induction in WT at temperatures indicated. Error bars show the standard deviation of four biological replicates. *UBC21* mRNA was used as a reference gene. p-value from the two-tailed t-test, *p-value < 0.05.

Elevated temperature is a key environmental factor responsible for secondary dormancy establishment[51]. Consistent with previous studies[7,49,52,53], we showed that *DOG1* expression increases in response to high temperatures during secondary dormancy establishment (Fig. 3e). At 35°C *DOG1* expression dropped initially during imbibition but rose to levels observed in dry seeds on the second day of secondary dormancy induction and remained steady for the rest of the experiment (Fig. 3e). Similarly, high temperatures augmented *MUSHER* induction in imbibed

seeds (Fig. 3f). *MUSHER* levels were similarly reduced by imbibition and recovered at 2^{nd} day,
but, in contrast to *DOG1*, continued to rise during the entire experiment length without
reaching the plateau. This suggests that *MUSHER* is more responsive to secondary dormancy-
inducing conditions than *DOG1* (Fig. 3f).
Next, we tested if *MUSHER* could pass the temperature input for *DOG1* regulation during
secondary dormancy induction. We subjected luciferase reporter expressing seeds
(Supplementary Fig. 2) to secondary dormancy induction at 25, 28, and 35°C. We noted that
MUSHER and *DOG1* reporters showed similar gradual increases in expression in response to
increasing temperatures (Fig. 3g). Interestingly, *DOG1*-expressing construct devoid of *MUSHER*
region (*pDOG1::DOG1-LUC truncated*) showed no induction at 35°C (Fig. 3g). The
responsiveness of *MUSHER* to high temperatures, along with *MUSHER* requirement for high
temperature-driven *DOG1* upregulation, indicated that *MUSHER* conveys temperature signals
during the induction of secondary dormancy. In agreement with the findings that *MUSHER*
enhances the selection of the proximal termination site, we noticed a prominent increase in the
level of the short but much smaller increase in the long *DOG1* mRNA at high temperatures
during secondary dormancy induction (Fig. 3h).
491
492 In summary, our data are consistent with a model in which *MUSHER* lncRNA acts as an activator
493 of *DOG1* proximal polyA site selection both during primary and secondary dormancy. In
494 addition, we showed that during secondary dormancy induction, *MUSHER* lncRNA is required
495 for *DOG1* upregulation in response to high temperature.
496
MUSHER enhances the _PIR1_ gene expression independently of its role in _DOG1_ regulation.
3'RNA-seq analysis revealed 166 differentially expressed genes, including *DOG1,* between WT
and *msh-1* seeds (Fig. 4a). Among the downregulated genes, *PIR1* (*PP2CA INTERACTING RING*
FINGER PROTEIN 1) showed the strongest decrease (Fig. 4a). *PIR1* encodes an E3 ligase that
enhances the abscisic acid (ABA) response and is highly expressed in maturing siliques[36].
Inspection of 3'RNA-seq reads showed a strong reduction in reads corresponding to both *PIR1*
and an upstream localised *AT2G35320* gene (Fig. 4b). RT-qPCR confirmed the strong
downregulation of *PIR1* expression in *musher* mutant for mature (Fig. 4c) and secondary
dormancy-induced seeds (Fig. 4d). Interestingly, *PIR1* expression remained unchanged in the
dog1 mutant, indicating that *PIR1* is controlled by *MUSHER* independently from its effect on
DOG1 (Figs. 4c,d). Consistent with the conclusion that *MUSHER* activates *PIR1*, we observed
that increased *MUSHER* expression in *cpl1-7* led to higher *PIR1* levels (Fig. 4d). ChIP-qPCR
analysis revealed a significant decrease in Pol II occupancy throughout the entire *PIR1* locus in
msh, suggesting that in contrast to *DOG1*, *MUSHER* regulates transcription levels rather than
termination of *PIR1* gene (Fig. 4e). ChIP-qPCR analysis along *PIR1* locus showed a TSS-localised
decrease in H3K4me3, slight decrease in H3 and no change in H3K27me3 levels in *msh-1* (Figs.
4f-h). As H3K4me3 levels are highly correlated with Pol II transcriptional activity[54], this agrees
with the reduced expression of the *PIR1* gene.
515

Fig. 4| *MUSHER* enhances *PIR1* expression independently of its role in *DOG1* regulation.

a Volcano plot showing differentially expressed genes for WT and *msh-1* selected using DESeq2; absolute log2FC >log2 (1.5), FDR<0.05. *DOG1* and *PIR1* genes are highlighted. The number of downregulated (down) and upregulated (up) genes is provided on the plot. **b** Diagram of 3'RNA-seq reads for *PIR1* locus in WT and *msh-1* mutant. Schematic diagram showing *PIR1* locus: exons (grey rectangles), 3'UTR region (grey arrow) **c** The *PIR1* relative expression level in freshly harvested seeds (primary dormancy) of WT, *msh-1* and *dog1-4* mutants, normalised to *UBC21* and relative to WT. Error bars show the standard deviation of four biological replicates. *$p<0.05$. **d** The *PIR1* relative expression level in WT *msh-1*, *dog1-4* and *cpl1-7* mutant seeds, after 5 days of SD induction, normalised to *UBC21* and relative to WT. Error bars show the standard deviation of four biological replicates. *$p<0.05$. **e** Pol II **f** H3 **g** H3K4me3 and **h** H3K27me3 occupancy was measured by ChIP-qPCR in WT, *msh-1*. (n=3) H3K4me3 and H3K27me3 values were normalised to H3 and to *ACTIN7* (*AT5G09810*); Pol II and H3 were normalised to *ACTIN7* (*AT5G09810*). For all ChIP experiments *$P< 0.05$, ns - not significant (two-tailed t-test). Error bars represent the standard deviation.

Our results showed that *MUSHER* activates *PIR1* expression independently of its role in *DOG1* gene regulation. In addition, our data suggested that *MUSHER* regulates transcriptional activity at the *PIR1* gene, whereas at *DOG1*, it specifically promotes proximal termination site selection. Importantly, both *DOG1* and *PIR1* regulation by *MUSHER* is compatible with its chromatin association (Fig. 1).

MUSHER lncRNA activates *PIR1* expression through *PIR1-DOG1* chromatin interaction

We showed that *MUSHER* is chromatin-bound (Fig. 1h) and regulates gene expression at two *loci*. To identify *MUSHER* binding sites, we conducted a Chromatin Isolation by *MUSHER*-RNA Purification followed by DNA-seq (ChIRP-seq) in seeds[55]. This revealed 23 DNA regions bound by *MUSHER* lncRNA (Supplementary Table 1). The two most significantly enriched are *PIR1* and *DOG1 loci*. Quantifying the ChIRP signal at *DOG1* and *PIR1 loci* shows that target enrichment is greatly diminished in the *msh-1* mutant (Figs. 5a,b). In addition, ChIRP using *LacZ* probes, a commonly used negative control, also failed to recover significant signals at both targets (Figs. 5a,b). In agreement with our conclusion that *MUSHER* regulates *PIR1* transcription initiation,

MUSHER peaks centred around *PIR1* TSS. On the *DOG1* locus, *MUSHER* ChIRP-seq detected two strong peaks, one at a site where it is produced and one colocalising with the *DOG1* exon2-intron2 region, just upstream of the proximal polyA site, the selection of which is promoted by *MUSHER* (Fig. 5a).

Fig. 5 | *MUSHER* lncRNA integrates ABA and DOG1 pathways by interacting with *PIR1* TSS and *DOG1* proximal termination site.

Snapshot of ChIRP-seq in WT and *msh-1* mutant using *MUSHER*-specific and *LacZ*-specific probes on **a** *DOG1* and **b** *PIR1 loci*. Schematic diagram showing *DOG1* and *PIR* locus: exons (grey rectangles), *shDOG1* 3'UTR region (light grey arrow), *lgDOG1* 3'UTR region (grey arrow), and *MUSHER* (green). The lines below the scheme mark regions used to design *MUSHER* probes. **c** Snapshot of ChIDP-seq in WT using *PIR1*-specific probes on *DOG1* and **d** *PIR1 loci*. Data are the mean of two biological replicates. The lines below the scheme mark regions used to design *PIR1* probes. **e** The germination rate of WT, *dog1-3*, *at2g35320-1*, *pir1-1* and *msh-1* mutant seeds on MS media supplemented with various ABA concentrations. *P-value<0.05 from two-tailed Student t-test. Error bars show standard deviations from three biological replicates.

Our data showed that the *DOG1* proximal polyA site and *PIR1* TSS are bound by *MUSHER* lncRNA. Together with the lack of *PIR1* expression changes in the *dog1* mutant (Fig. 4c), this suggested that both *DOG1* and *PIR1* are direct and independent *MUSHER* targets. Of note, we could not find a common DNA motif or R-loop enrichment among *MUSHER* targets, indicating that *MUSHER* may not rely on a sequence-based mechanism for target specification. Whereas the *DOG1 locus* is localised next to *MUSHER* on chromosome 5, the *PIR1 locus* is localised on chromosome 2. This encouraged us to test the physical proximity of *PIR1* and *DOG1 locus* chromatin.

To this end, we used *PIR1* probes to fish out crosslinked DNA sequences followed by NGS. This assay is based on a similar principle as PiCh (Proteomics of isolated Chromatin segments)[56] but analyses DNA rather than proteins attached to a specific *locus*. This method was called chromatin-DNA precipitation assay (ChIDP). The use of glutaraldehyde combined with the covalent link between probes and beads allowed us to use more stringent conditions, resulting in a high signal-to-noise ratio (Figs. 5c,d, Supplementary Table 2). Surprisingly, we discovered that *PIR1* DNA interacts with the proximal polyA site of *DOG1* but not with *MUSHER*. This suggests that *MUSHER* activity on *PIR1* is mediated by long-range interaction established by the proximal polyA site of *DOG1*, which is also regulated by *MUSHER*.

DOG1 and ABA are the two main regulators of dormancy in Arabidopsis[57]. We confirmed the
published results that *pir1* mutants show decreased sensitivity to ABA-mediated germination
inhibition (Fig. 5e)[36]. Notably, a mutant in a gene directly upstream of *PIR1* (*at2g35320-1*)
showed no significant changes in ABA sensitivity compared to WT. In agreement with published
results, *dog1* mutant seed germination was suppressed by ABA in the same way as WT[27].
Importantly, *musher* seeds showed much weaker sensitivity to ABA-mediated inhibition of
germination across all tested ABA concentrations, similar to the *pir1* mutant (Fig. 5e). Therefore,
in contrast to the DOG1 protein, *MUSHER* lncRNA does modulate ABA responsiveness. Our data
are consistent with a model where *MUSHER* lncRNA enhances the ABA sensitivity by interacting
with and activating *PIR1* gene expression. *MUSHER's* ability to activate *DOG1* and *PIR1*
expression points to its role in integrating the DOG1 and ABA-dependent repression of seed
germination.

U1 snRNP mediates *MUSHER* association to chromatin
MUSHER binds to and regulates *DOG1* and *PIR1* gene expression, with most *MUSHER* transcripts
bound to chromatin (Fig. 1h). This raised the question of whether *MUSHER* association with
chromatin is required for its function and what is the mechanism of *MUSHER* binding to
chromatin.
In addition to its well-known splicing function, U1 snRNP has been shown in mammals to bind
various lncRNAs, facilitating their chromatin tethering[15]. Interestingly, we identified four
putative U1 binding sites within the *MUSHER* sequence even though we failed to detect
MUSHER splicing (Supplementary Table 3 and Fig. 1c). Using probes specific for *MUSHER*, but
not *LacZ* probes, we selectively recovered U1, but not U4 or U6 RNA using formaldehyde-
assisted RNA ImmunoPrecipitation (RIP) (Fig. 6a). Supporting the specificity of the approach, U1
recovery was strongly diminished when we repeated the experiment using *MUSHER* probes in
msh-1 lines (Fig. 6b). Next, we validated the *MUSHER*-U1 association by RNA
Immunoprecipitation (RIP) using the U1 snRNP component U1-70K. We used a recently
reported *u1-70k* complementing line expressing GFP tagged U1-70K driven by endogenous U1-
70K promoter (*pU1-70K::U1-70K-GFPu1-70k*)[21]. RT-qPCR showed that U1-70K
immunoprecipitation recovered spliced control RNA, failed to recover unspliced control mRNA,
and, importantly, retrieved a significant amount of *MUSHER* lncRNA (Fig. 6c).
639

Fig. 6| U1 is required for chromatin tethering of *MUSHER* and other lncRNA.

a *MUSHER* or *LacZ* probes were used in formaldehyde-assisted RIP to enrich associated RNA. U1, U4 and U6 recovery was analysed using RT-qPCR. Error bars show standard deviations. **b** *MUSHER* probes were used to enrich *MUSHER* from WT and as a negative control in *msh-1*. U1 and U6 recovery was analysed using RT-qPCR. Error bars show standard deviations. **c** U1A protein was precipitated side by side with a no-antibodies (NoAB) control from plants expressing *pU1-70K::U1-70K-GFP u1-70k,* and co-precipitated RNA was analysed using RT-qPCR for spliced (AT5G45830) and unspliced (AT5G45820) RNA controls and *MUSHER*. Error bars show standard deviations. **d** Subcellular localisation of mRNAs and lncRNAs in maturing Arabidopsis seeds of WT and *u1c* amiRNA line. Lines represent the mean of chromatin and nucleoplasmic fractions contribution of the total amount of indicated transcript. The blue-shaded colour refers to Col-0, while the grey-shaded colour corresponds to *u1c*. Each transcript was measured in 4 replicates represented as a separate bar. *p-value<0.01 from two-tailed Student t-test. **e** After ripened seeds of WT (WT), *u1c* amiRNA and *u1k-70k* mutants were induced into secondary dormancy, moved to permissive conditions and scored for germination ability. Germination (%) was scored as radical protrusion. A two-tailed t-test p-value was calculated compared to WT (*p-value < 0.05). Error bars show standard deviations. **f** *shDOG1* expression was analysed in WT, *u1c* amiRNA and *u1-70k* mutants seeds induced into secondary dormancy. A two-tailed t-test p-value was calculated compared to WT (*p-value < 0.05). Normalised to *UBC21* and relative to WT. Error bars show standard deviations.

Having shown that *MUSHER* is bound by U1 but not U6 or U4 snRNP, we asked if this contributes to *MUSHER* chromatin association. To do this, we used a recently described artificial microRNA (amiRNA) line that shows significantly downregulated *u1c* levels[21]. Subcellular localisation showed that *UBC21* and *DOG1* mRNA are nearly equally distributed between cytoplasmic, nucleoplasm and chromatin-bound fractions, as shown also in Fig. 1. In addition, neither *UBC21* nor *DOG1* mRNA distribution among the cellular fractions analysed was altered in the *u1c* mutant (Fig. 6d). Notably, *MUSHER* chromatin association was reduced in *u1c* mutant, a result that is expected given the presence of four U1 binding sites within the *MUSHER* transcriptional unit. Next, we asked whether the U1 requirement for *MUSHER* chromatin retention is a more general phenomenon in plants. We started by confirming that *COOLAIR*, shown to be expressed

in seeds[58], is predominantly chromatin-bound like *MUSHER* (Fig. 6d). In addition, unspliced *PUPPIES* isoform, a lncRNA involved in Pol II pausing regulation at *DOG1* in seeds[7] was also strongly enriched in chromatin fraction. Both *PUPPIES* and *COOLAIR* have multiple U1 binding sites (Supplementary Table 3) and we found that in contrast to the control mRNA, *MUSHER* and all other analysed lncRNAs showed reduced chromatin association and a corresponding increase in nucleoplasmic abundancy in the *u1c* mutant (Fig. 6d). We conclude that *MUSHER* binds U1 snRNP and that U1 facilitates *MUSHER* and other tested lncRNA-chromatin association and propose that the U1 role in chromatin retention of lncRNA is conserved in plants.

Finally, we asked if *MUSHER* chromatin association is required for its function. Both *u1c* and another amiRNA mutant in the U1 snRNP component U1-70K, showed pleiotropic phenotypes, including significant dwarfism and delayed development[21]. Thus, it would be difficult to distinguish between primary dormancy phenotype and developmental defects resulting in delayed germination. To circumvent this, we took advantage of the fact *MUSHER* shows defects in both primary and secondary dormancy induction. To synchronise the physiological state of seed , they were matured on mother plants, stored until they lost primary dormancy and then subjected to induce secondary dormancy. *u1c* and *u1-70k* mutant seeds showed low secondary dormancy phenotype with nearly all seeds germinating compared to less than 50% of germinated seeds in the WT (Fig. 6e). This agrees with the model where U1 is required for *MUSHER* association to chromatin and seed dormancy induction. The reduction in *u1c* and *u1-70k* secondary dormancy was accompanied by decreased *shDOG1* expression, as expected from *musher* mutants' analysis (Fig. 6f). In summary, U1 is required for multiple lncRNA chromatin associations and, at least in the case of *MUSHER,* this association is required for the lncRNA gene expression regulatory function.

Discussion

Our results indicate that *MUSHER* modulates seed dormancy through the transcriptional regulation of DOG1 and ABA sensitivity pathways (Fig. 7a). We show that *MUSHER* lncRNA is chromatin-bound and requires U1 snRNP for its retention and activity at chromatin (Fig. 7b).

Fig. 7| a *MUSHER* modulates seed dormancy through the transcriptional regulation of DOG1 and ABA pathways. b U1 is required for chromatin retention of lncRNA, including *MUSHER* in Arabidopsis.

MUSHER-Mediated Regulation of *DOG1* and *PIR1* Expression

Our analysis showed that *MUSHER* employs two different mechanisms to activate *DOG1* and *PIR1*. *MUSHER* promotes *DOG1* expression by enhancing CPSF complex recruitment to the proximal cleavage and polyadenylation site, resulting in enhanced production of a functional short *DOG1* alternative polyadenylation isoform (Fig. 2). In agreement, the *MUSHER* effect on *DOG1* is not accompanied by changes in Pol II recruitment or H3K4me3 levels at the *DOG1* gene (Fig. 2F-I and Supplementary Fig. 3). In contrast, *MUSHER* activates *PIR1* expression by enhancing Pol II recruitment and consequent H3K4me3 deposition at the *PIR1* promoter (Fig. 4). Similar to *HID1* lncRNA[13], and differently then *COOLAIR*[59] and *APOLO*[5], *MUSHER* promotes H3K4me3 deposition without affecting H3K27me3 repressive marks. Notably, *HID1* represses transcription by blocking the binding of ARABIDOPSIS TRITHORAX-RELATED7 (ATXR7), an H3K4me3 methyltransferase, to nucleosomes, leading to increased H3K4me3 levels when HID1 is inactivated[13]. In contrast, *MUSHER* inactivation reduces H3K4me3 deposition at the *PIR1* promoter, suggesting that *MUSHER* and *HID1* regulate H3K4me3 through distinct molecular mechanisms. Of note, we did not detect *PIR1* proximal polyadenylation either in our seed-specific 3'RNA-seq data (Figure 4B) or in the published long read-based transcript mapping dataset[39]. It is currently not clear why *MUSHER*-dependent *DOG1* and *PIR1* regulation are based on different modes of activity. It is possible that *MUSHER* acts in the same way on *DOG1* and *PIR1*, but the local chromatin environment of the target or *MUSHER* site of binding with respect to the target transcription start site results in seemingly different mechanisms. For example, the recruitment of CPSF complex to the gene promoter may enhance transcription initiation or transition to efficient elongation. We, however, favour a model where *MUSHER* employs two different mechanisms to enhance *DOG1* and *PIR1* expression. This view is supported by the fact that *DOG1* expression is reduced roughly by 50%, while *PIR1* expression is undetectable in multiple *msh* mutants tested.

Chromatin precipitation using *MUSHER* RNA (ChIRP) followed by sequencing revealed only a very limited number of *MUSHER* targets, with *DOG1* and *PIR1* being the most significantly enriched. The high selectivity of *MUSHER* is somehow surprising, as many of the published plant lncRNAs show binding to multiple sites spread across the genome[5,60]. *PIR1* DNA chromatin-DNA precipitation assay (ChIDP) followed by sequencing showed that *PIR1* DNA also has a very limited number of DNA targets, with, excluding *PIR1* bait, *DOG1* being the strongest. This suggests that high *MUSHER* selectivity could be guided by a spatial arrangement of DNA in seeds.

MUSHER as a Link Between ABA Sensitivity and *DOG1*

Interestingly, both *MUSHER* targets encoded proteins PIR1 and DOG1, interact with PP2CA protein, a regulator of ABA signalling[35]. PIR1 binding to PP2CA promotes its degradation, while DOG1 inhibits PP2CA phosphatase activity[35,37,61]. Both *dog1* and ABA synthesis deficient *nced2/5/6/9* mutants exhibit a failure to enter dormancy[62]. Simultaneously, *dog1* does not show major changes in ABA levels or sensitivity[27,63] and *nced2/5/6/9* shows unaffected *DOG1* induction kinetics during secondary dormancy establishment[53]. This suggests that DOG1 and ABA synthesis pathways function independently in dormancy regulation. We demonstrate that *MUSHER* inactivation leads to decreased transcription of *DOG1* and *PIR1*, which results in

decreased ABA sensitivity. We propose that *MUSHER* lncRNA facilitates the concurrent
activation of DOG1 and ABA pathways, integrating them into a shared regulatory mechanism.
High temperatures are known to enhance *DOG1* expression and ABA biosynthesis in seeds[51,67].
We observed altered ABA sensitivity and reduced secondary dormancy induction in response
to high temperature in *msh* mutants. Elevated temperatures contribute to two well-described
responses: 1) thermoinhibition, a high-temperature-mediated inhibition of germination in
light[68] and 2) secondary dormancy induction in non-dormant imbibed seeds exposed to high
temperatures in the darkness[31,49,51]. We hypothesise that *MUSHER*-mediated integration of ABA
and DOG1 pathways contributes not only to dormancy but also to other ABA/DOG1-related
phenotypes (e.g., drought tolerance or thermoinhibition). We conclude that, similar to other
long non-coding RNAs involved in temperature sensing, such as *COOLAIR*[8], *COLDWRAP*[69] and
SVALKA [6] in plants, and *Hsrω*[70] and *Heat*[71] in animals, *MUSHER* also plays a role in temperature
sensing in plants.

U1 role in chromatin retention of lncRNA in Arabidopsis.

In contrast to mature mRNAs, lncRNAs are often enriched in the nucleus[1,15]. Consistently, many
lncRNA have been shown to regulate gene expression by binding to chromatin[1,5,6,39]. The
molecular mechanism of human lncRNA retention in the nucleus has only recently started to
be explored [15,72,73], and those remain largely unknown in plants. Mechanisms ensuring human
lncRNA nuclear retention encompass the presence of specific sequences[74], including *Alu*
transposable elements binding the HNRNPK complex[75]. These protein-RNA interactions may
contribute to liquid-liquid phase separation (LLPS) that can also be responsible for the nuclear
retention of lncRNA. For example, LLPS by *NEAT1* lncRNA results in the formation of nuclear
paraspeckles[76] and *XIST* lncRNA was reported to undergo LLPS with different partners localised
at inactive X chromosome[77]. However, nuclear retention may not be enough to guide lncRNA
directly to chromatin, which has been proposed to be a result of direct RNA-DNA interaction
either in the form of R-loops or RNA:DNA triplex formation[15]. In addition, U1 snRNP, involved in
5' splice site recognition, has recently been shown to be responsible for the chromatin binding
of a large fraction of lncRNAs[15]. In contrast to *Alu* and other sequence elements identified in
humans, U1 binding sites are well conserved in plant genomes.

Plant lncRNA retention at chromatin has not been explored, but RNA-seq analysis indicates that
lncRNAs are enriched in the nucleus[78]. Also, like their human counterparts, plant lncRNAs often
regulate gene expression by chromatin binding[6], indicating that their chromatin retention may
be actively maintained. Multiple plant lncRNAs have been described, but few of them have been
reported to be expressed in seeds and to control gene expression by chromatin-related
mechanisms. This includes *COOLAIR* regulating the *FLC* gene chromatin state[9,79], *PUPPIES*
regulating Pol II pausing at the *DOG1* gene[7] and, described here, *MUSHER* promoting the *DOG1*
gene proximal cleavage and polyadenylation site selection. In seedlings, *COOLAIR* is retained at
chromatin, presumably by its ability to form R-loops[80] and a cloud around its transcription site[10].
Similarly, *APOLO* lncRNA has been shown to form R-loops at its targets located in *trans* to its
transcription site[5]. Here, we propose that U1-mediated chromatin binding of lncRNAs is a
largely sequence-independent mechanism used by plants to retain lncRNA at chromatin. We
showed that U1 snRNP is required for *MUSHER*, *COOLAIR* and *PUPPIES* to bind the chromatin

efficiently. As we did not test the interplay between U1 binding and LLPS, we cannot exclude
the possibility that U1 binding is required or otherwise interlinked with the phase separation of
lncRNA at chromatin. R-loops have not been reported in seeds, and the one shown in seedlings
at *PIR1* locus[81] is unlikely to be formed by *MUSHER* since *MUSHER* is very lowly expressed in
seedlings. Therefore, while we cannot exclude this possibility, *MUSHER* is unlikely to form R-
loops. As concluded for human U1 lncRNA targets[15], U1 also do not form direct R-loops with
the target DNA. This suggests that R-loops probably do not contribute to U1-mediated lncRNA
retention. The fact that the tested lncRNA have different modes of action implies that the U1
role in chromatin retention is not limited to a specific lncRNA activity. This, together with the
described role in chromatin retention of human and mouse lncRNA, strongly supports the idea
that the U1 role in chromatin retention of lncRNA is evolutionarily conserved in plants.

Methods

Plant material and growth conditions

WT used in this work is Columbia (Col-0) *Arabidopsis thaliana* accession. Plants were grown on soil in a
greenhouse under long-day conditions (16 h light/8 h dark) at 22°C/18°C. All mutants and transgenic
lines were created in Col-0 background and were homozygous for the indicated genotype. *MUSHER* lines
msh-1(SM_29297) and *msh-2*(SM_29310) were obtained from the NASC centre. Other mutants used in
this study have been described previously: *cpl1-7*[82], *cpl1-8* (GK-165H09)[32], *cpl1-9* (GK-849H10)[32], *dog1-*
3 (SALK_000867)[27], *dog1-4* (SM_3_20886)[30], *fy-2* (SAIL_ 657D4)[83], *esp1-2* (SALK_078793)[84], *pir1*
(SALK_078237)[36], *amiR-u1-70k* and *amiR-u1-c*[21].

Germination tests

Germination assays were performed as described previously to estimate primary dormancy levels[7,30].
Briefly, 100-200 freshly harvested seeds were sown on Petri plates with blue germination paper
(HoffmanMfg) and a layer of water-soaked fabric underneath to keep moisture. The plates were sealed
and placed in the Percival chamber under long-day conditions (16 h light/8 h dark) at 22°C/18°C.
Photographic documentation was performed once a day, and seeds with root protrusion were counted
as germinated.

To evaluate secondary dormancy induction levels, seeds were sown on Petri plates with blue
germination paper (HoffmanMfg) and a layer of water-soaked fabric underneath to keep moisture. The
plates were sealed and incubated in the darkness at varying temperatures (25°C, 28°C, 35°C) for 7 days.
Subsequently, plates were transferred to the Percival chamber under long-day conditions (16 h light/8 h
dark) at 22°C/18°C.

Cloning, Plant Transformation and CRISPR-dCas9 mutant generation

The generation of the dCas9 transgenic plants was performed according to Čermák et al.[42]. The specific
sgRNA to Arabidopsis TAIR10 genome were designed using the CRISPR-P 2.0
(http://crispr.hzau.edu.cn/cgi-bin/CRISPR2/CRISPR) to select the canonical NGG PAM motif. sgRNAs were
cloned to the dCas9 vector by Golden Gate assembly (*dCas9_control-1*, *dCas9_control-2*, *msh_dCas9-1*
and *msh_dCas9-2*) according to the protocols 3B and 5[42]. Primers used for cloning are listed in Table S4.
All constructs were verified by Sanger sequencing. Each construct was transformed into Agrobacterium
strain GV3101 and then transformed into Col-0 or an appropriate mutant background using the floral
dip method. T1 transformant plants were identified by selecting the resistance to glufosinate. The seeds
from the selected T1 plants were used for phenotypic analysis. At least two independent transgenic lines

were tested and propagated. As a control for potential indirect effects from dCas9 expression, we also
created two dCas9 lines with guides recognising different intergenic regions localised 100 kb from *DOG1*
locus: *dCas9_control_1* and *dCas9_control_2*, which showed no influence on *MUSHER* transcription.

Luciferase measurement

832   100 seeds placed on a microscope glass slide were squashed to release the embryo from the seed coat.
833   After adding D-Luciferin (Beetle luciferin potassium salt, Promega) and 20 minutes of incubation in
834   darkness, the light signal was measured using a NightSHADE camera (Berthold Technologies GmbH &
835   Co.KG) with a 10-minute exposure.

RNA extraction

RNA extraction protocol was adapted from Kowalczyk et al.[32] After grinding in liquid nitrogen, samples
were resuspended in 500 µl of homogenisation buffer (100 mM Tris pH 8.5, 5 mM EDTA, 100 mM NaCl,
0.5% SDS, 1% β-mercaptoethanol) and centrifuged for 5 min at room temperature at 8,000 g. The
supernatant was transferred to the new tubes containing 250 µl of chloroform. After 15 min of shaking,
250 µl of phenol was added, and shaking was repeated, followed by centrifugation for 10 min at
10,000 g. The upper phase was transferred to new tubes, and an equal volume of
phenol:chloroform:isoamyl alcohol was added. Shaking and centrifugation steps were repeated. The
upper phase was mixed with 10% of the volume of 3M sodium acetate (pH 5.2) and an equal volume of
isopropanol. After centrifugation for 30 min at 20,000 g at 4°C, the pellet was washed with 70% EtOH
and resuspended in water.

RT–qPCR

Trace DNA in the RNA samples was removed using a TURBO DNA-free kit (Thermo Fisher Scientific)
according to the manufacturer's protocol. The DNA removal was confirmed by PCR using primers to the
PP2A gene (AT1G13320). The RNA sample quality and quantity were checked by agarose gel
electrophoresis and using a Nanodrop 2000 spectrophotometer. Reverse transcription using 2.5 µg of
RNA was performed with a Superscript III cDNA Synthesis kit (Thermo Fisher Scientific) according to the
manufacturer's protocol. An equal mixture of oligo(dT) and hexamer primers was used for cDNA
synthesis according to the manufacturer's protocol. 5x diluted cDNA was used as a template in the qPCR
reaction (LightCycler 480; Roche) with SYBR Green mix (Roche). The relative transcript level was
determined with the $2^{-\Delta\Delta Ct}$ method [28] and normalised to mRNA of the *UBC21* (AT5G25760) gene. The
primers used are listed in Table S4.

Chromatin ImmunoPrecipitation (ChIP)

ChIP was performed as described[32] with the following modifications. 100 mg of fresh siliques (13-20
days after pollination) were ground in liquid nitrogen and resuspended in 20 ml of crosslink buffer (10
mM HEPES-KOH pH 7.4, 50 mM NaCl, 100 mM sucrose, 1% formaldehyde). After 10 min incubation on
a rotator at 4°C, the crosslinking agent was quenched by adding 1.25 ml of 2 M glycine. After another 5
min incubation on a rotator, samples were centrifuged at 4°C and 2,000 g for 15 min. The pellets were
resuspended in 20 ml of cold Honda Buffer (20 mM HEPES-KOH pH 7.4, 0.44 M sucrose, 1.25% Ficoll,
2.5% Dextran T40, 10 mM MgCl$_2$, 5 mM DTT, 0.5% Triton X-100, 10 mM β-mercaptoethanol, 1 mM
PMSF, 1x Complete protease inhibitors (Roche)) and filtered through Miracloth. Samples were
centrifuged for 15 min at 2,000 g at 4°C. The pellet was resuspended in ChIP lysis buffer (50 mM HEPES-
KOH pH 7.4, 10 mM EDTA, 1% SDS, 1x Complete protease inhibitors (Roche)) and sonicated using
Bioruptor (Diagenode) to fragments of 300-500 bp. Samples were next diluted 10 times with ChIP
Dilution Buffer (1.1% Triton X-100, 1 mM EDTA, 16.7 mM HEPES-KOH pH 7.4, 167 mM NaCl, 1 mM DTT,

1 mM PMSF, 1x Complete protease inhibitors (Roche)). Then antibodies against RNA polymerase II
subunit B1 (Agrisera, AS11 1804), H3 (Abcam, ab1791), H3K4me3 (Millipore, 07-473), H3K27me3
(Abcam, ab6002) or CPSF73 (PhytoAB, PHY1043A) were added followed by addition of the mixture of
Dynabeads Protein A and Protein G beads (Thermo Fisher Scientific). Next, samples were incubated
overnight on a rotator at 4°C. Two washing steps with low salt buffer (150 mM NaCl, 1 mM EDTA, 10
mM HEPES-KOH pH 7.4, 0.1% Triton X-100) and two washing steps using high salt buffer (500m M NaCl,
1 mM EDTA, 10 mM HEPES-KOH pH 7.4, 0.1% Triton X-100) were performed. DNA was released from
beads and reverse-crosslinked by incubation at 95°C with 10% Chelex 100 suspension for 15 min. After
Proteinase K treatment (55°C) for 90 min and repeated incubation at 95°C for 10 min, samples were
centrifuged for 5 min at RT at 10,000 g. Recovered DNA was used in qPCR reaction with primers listed
in Table S4.

RNA ImmunoPrecipitation (RIP)

For U1-70K-GFP RIP, we used 100 mg of fresh siliques (13-20 days after pollination), which were ground
in liquid nitrogen and resuspended in 20 ml of crosslink buffer (10 mM HEPES-KOH pH 7.4, 50 mM NaCl,
100 mM sucrose, 1% formaldehyde). After 10 min incubation on a rotator at 4°C, the crosslinking agent
was quenched by adding 1.25 ml 2 M glycine. After another 5 min incubation on a rotator, samples were
centrifuged at 4°C and 2,000 g for 15 min. The pellets were resuspended in 20 ml of cold Honda Buffer
(20 mM HEPES-KOH pH 7.4, 0.44 M sucrose, 1.25% Ficoll, 2.5% Dextran T40, 10 mM $MgCl_2$, 5 mM DTT,
0.5% Triton X-100, 10 mM β-mercaptoethanol, 1mM PMSF, 1x Complete protease inhibitors (Roche),
and 5 U/ml murine RNase inhibitor (NEB)) and filtered through Miracloth. Samples were centrifuged for
15 min at 2,000 g at 4°C. The pellet was resuspended in Nuclear Lysis Buffer (50 mM HEPES-KOH pH 7.4,
10 mM $MgCl_2$, 1% Triton X-100, 1x Complete protease inhibitors (Roche), 20 U/ml murine RNase
inhibitor (NEB)), followed by 20 min incubation at 37°C with TURBO DNAse (Thermo Fisher Scientific).
After adding SDS up to 1%, samples were sonicated using Bioruptor (Diagenode). Next, samples were
diluted 10 times with RIP Dilution Buffer (0.5 mM EDTA, 10 mM HEPES-KOH pH 7.4, 150 mM NaCl,
20 U/ml murine RNase inhibitor (NEB)) and incubated with GFP Agarose beads (Chromotek) for 1h on a
rotator at 4°C. After three washing steps with RIP Wash Buffer (150 mM NaCl, 1 mM EDTA, 10 mM
HEPES-KOH pH 7.4, 0.05% NP-40, 20 U/ml murine RNase inhibitor (NEB)), RNA was eluted (150 mM
NaCl, 1 mM EDTA, 10 mM HEPES-KOH pH 7.4, 1% SDS, 20 U/ml murine RNase inhibitor (NEB)) and
reverse-crosslinked by incubation at 95°C for 15 min. After 1h Proteinase K treatment (55°C), RNA was
isolated with Trisure (Bioline) according to producer protocol. After cDNA synthesis with SuperScript III
(Thermo Fisher Scientific), samples were used in qPCR reaction with appropriate primers listed in Table
S4.

Nuclei purification for ChIRP, RIRP and ChIDP

To uncover RNA-RNA (RNA Isolation by RNA Purification), RNA-DNA (Chromatin Isolation by RNA
Purification) and DNA-DNA (Chromatin Isolation by DNA Purification Assay) interactions, we used
400 mg of fresh siliques (13-20 days after pollination), which were ground in liquid nitrogen and
resuspended in 20 ml of crosslink buffer (10 mM HEPES-KOH pH 7.4, 50 mM NaCl, 100 mM sucrose, 1%
formaldehyde (for RNA-RNA and RNA-DNA) or 0.5% glutaraldehyde (for ChIDP)). After 10 min incubation
on a rotator at 4°C, the crosslinking reaction was quenched by adding 1.25 ml 2 M glycine. After an
additional 5 min incubation on a rotator, samples were centrifuged at 4°C and 2,000 g for 15 min. The
pellets were resuspended in 20 ml of cold Honda Buffer (20 mM HEPES-KOH pH 7.4, 0.44M sucrose,
1.25% Ficoll, 2.5% Dextran T40, 10 mM $MgCl_2$, 5 mM DTT, 0.5% Triton X-100, 10 mM
β-mercaptoethanol, 1 mM PMSF, 1x Complete protease inhibitors (Roche) and 5 U/ml murine RNase
inhibitor (NEB)) and filtered through Miracloth. Subsequently, samples were centrifuged for 15 min at

2,000 g at 4°C. According to downstream assays, the nuclei pellets were resuspended in appropriate
buffers (Element Zero Biolabs kit).

Chromatin Isolation by RNA Purification Assay (ChIRP)

To enrich specific RNA molecules or chromatin fragments, we used MagIC beads with custom probes
targeting those sequences (Element Zero Biolabs). Element Zero Biololabs designed probes are delivered
with kits and remain confidential as protected intellectual property.

For *MUSHER* probes RIRP and ChIRP with *MUSHER* and *lacZ* probes, nuclei pellets were resuspended in
1.1 ml of MagIC Lysis Buffer FA (Element Zero Biolabs kit), supplemented with 1x Complete protease
inhibitors (Roche) and 5 mM DTT. Next, samples were sonicated using a Bioruptor (Diagenode) and
diluted two times with MagIC Lysate Dilution Buffer FA, with 1% of the volume saved as an input sample.
MagIC beads were pre-cleared according to the producers' protocol and added to the samples that were
next incubated at 40°C for 30 minutes with occasional shaking. This was followed by three washes using
MagIC Wash Buffer I FA and nucleic acids elution by incubation at 92°C for 5 min with 10 mM Tris-HCl
pH 7.5. After Proteinase K treatment (55°C, 90 min) and phenol:chloroform:isoamyl alcohol extraction,
DNA and RNA were precipitated by adding 1 µl glycogen, 10% of the volume of 3 M LiCl, and an equal
volume of isopropanol. Samples were centrifuged for 30 min at 4°C at 20,000 g. Pellets were washed
with cold 80% ethanol, resuspended in water, and samples were split in half. The presence of specific
DNA fragments interacting directly or indirectly with *MUSHER* probes was validated by qPCR analysis. To
identify RNAs enriched with *MUSHER* probes, half of the sample was incubated with TURBO DNAse
(Thermo Fisher Scientific), extracted with phenol:chloroform: isoamyl alcohol as above, and used for
cDNA synthesis and qPCR analysis. For ChIRP library preparation, DNA fragments were eluted from
MagIC beads with RNase If (NEB). Libraries were prepared with the NEBNext® Ultra™ DNA Library Prep
Kit for Illumina using NEBNext Multiplex Dual Index Primers. Paired-end sequencing was performed on
the Illumina NovaSeq 6000.

Chromatin Isolation by DNA Purification Assay (ChIDP)

For ChIDP, the nuclei pellet was resuspended in 1.1 ml of MagIC Lysis Buffer FA (Element Zero kit)
supplemented with 1x Complete protease inhibitors (Roche) and 5 mM DTT. Samples were sonicated
using Bioruptor (Diagenode), diluted with MagIC Lysate Dilution Buffer FA, and split into halves (treated
with RNaseA at 37°C for 20 min and non-treated). MagIC beads recognising the *PIR1 locus* were mixed
with the lysates that were next incubated at 81°C for 5 min with occasional shaking to denature DNA
partially. Subsequently, samples were transferred to 40°C for 20 min to enable binding of the probes.
This was followed by six washing steps with the MagIC Lysate Wash buffer I and II, and DNA was eluted
with 10 mM Tris-HCl pH 7.5 at 92°C for 5 min. After adding an equal volume of MagIC Lysate Wash buffer
I to eluted samples, proteinase K treatment (55°C, 120 min) was performed. To extract DNA fragments,
an equal volume of phenol:chloroform:isoamyl alcohol was added. The upper phase was mixed with 1 µl
of glycogen, 10% of the volume of 3 M LiCl and an equal amount of isopropanol. After centrifugation for
30 min at 20000g at 4°C, the pellet was washed with 80% EtOH and resuspended in water. Libraries were
prepared with the NEBNext® Ultra™ DNA Library Prep Kit for Illumina using NEBNext Multiplex Dual
Index Primers. Paired-end sequencing was performed on the Illumina NovaSeq 6000.

ChIRP-seq and ChIDP-seq analysis

957 For ChIRP-seq and ChIDP-seq data analysis, Bowtie2 (v2.5.3)[85] was used to align the reads to the
958 reference genome (TAIR10, with default parameters). To identify significantly enriched regions, MACS2
959 was used with the default parameters: bandwidth:300; model fold of 5:50; q-value cutoff: 0.05
960 (v3.0.0a6)[86]. bamCoverage (v3.5.4)[87] with CPM normalisation was used to create bigwig files.

5'RACE-seq

Analysis was performed as described previously[7]. Shortly, 500 ng of total RNA, after DNAse treatment,
was used as the template for cDNA generation using SuperScript II (Thermo Fisher Scientific). The
resulting cDNA was purified using AMPure XP magnetic beads (Beckman Coulter) and amplified in a
series of three PCR reactions with Phusion polymerase and specific primers (1st PCR: only TSO_n1, 2nd
PCR: TSO_n2 and MUSHER_5RACE, 3rd PCR: Illumina indexing primers). The final PCR product was
sequenced using Illumina MiSeq. For data analysis, STAR (v2.7.8a)[88] was used to align the reads to the
reference genome, followed by UMI-based PCR duplicates removal using UMI-tools (v1.1.0)[89]. The
position of the first 5' nucleotide was extracted using bedtools (v2.30.0)[90].

3'RACE-seq

Analysis was performed as described previously[7] and based on the earlier procedure[91]. The ligation of
the pre-adenylated adaptor to the 3'end of the RNA was performed using truncated T4 RNA Ligase 2
(NEB). RNA ligated with RA3_15N adaptor (containing UMI) was cleaned with AMPure XP magnetic
beads and subjected to reverse transcription reaction with SuperScript III (Thermo Fisher Scientific).
After three rounds of PCR with specific primers (1st PCR: MUSHER_3RACE and RTPXT, 2nd PCR: mXTf and
mXTr,3rd PCR: Illumina indexing primers), each followed with AMPure beads purification, prepared
libraries were sequenced using Illumina MiSeq. For data analysis, STAR (v2.7.8a)[88] was used to align the
reads to the reference genome, followed by UMI-based PCR duplicates removal using UMI-tools
(v1.1.0)[89]. The position of the last 3' nucleotide was extracted using bedtools (v2.30.0).

3'RNA-seq

Analysis was performed as described previously[49]. Briefly, 0.5 μg of total RNA was reverse transcribed
with UMI-containing oligo(dT) primers. Differently barcoded cDNAs were pooled together, and second-
strand cDNA was synthesised using the nick translation method. Next, libraries were tagmented with
Tn5 transposase, and the 3'cDNA fragments were amplified with Illumina-compatible primers[49]. Paired-
end sequencing was performed on the Illumina NextSeq 500. Alignment and read demμltiplexing were
performed using STAR (v2.7.8a)[88] and BRBseqTools (v 1.6)[92], respectively. Differentially expressed genes
were identified using DESeq2 with a cutoff of FDR < 0.05 and FC > 1.

Fractionation

Chromatin RNA extraction was performed using the described method[7]. 100 mg of siliques were ground
in liquid nitrogen and resuspended in 20 ml of cold Honda Buffer (20 mM HEPES-KOH pH 7.4, 0.44 M
sucrose, 1.25% Ficoll, 2.5% Dextran T40, 10 mM $MgCl_2$, 5 mM DTT, 0.5% Triton X-100, 10 mM β-
mercaptoethanol, 1 mM PMSF, 1x Complete protease inhibitors (Roche), and 5 U/ml murine RNase
inhibitor (NEB)). After rotation at 4°C for 10 min and filtration through Miracloth, samples were
centrifuged at 2,000 g for 15 min at 4°C. The nuclei pellet was washed twice with 5 ml of Honda buffer.
The pellet was resuspended in 600 μl of Honda buffer and purified on a 40%-75% Percoll density
gradient by centrifugation at 10,000 g for 30 min at 4°C. Purified nuclei were collected from the Percoll
interface, washed with Honda buffer and resuspended in 500 μl of cold glycerol buffer (20 mM Tris-HCl
pH 8.0, 75 mM NaCl, 0.5 mM EDTA, 0.85 mM DTT, 50% glycerol, 1% Empigen, 10 mM
β-mercaptoethanol, 0.125 mM PMSF, 1x Complete protease inhibitors (Roche), and 5 U/ml murine
RNase inhibitor (NEB)). Samples were overlaid on urea lysis buffer (10 mM HEPES-KOH pH 7.4, 7.5 mM
$MgCl_2$, 0.2 mM EDTA,300 mM NaCl, 1 M urea, 1% NP-40, 10 mM β-mercaptoethanol, 0.5 mM PMSF, 1x
Complete protease inhibitors (Roche), and 5U/ml murine RNase inhibitor (NEB), vortexed for 2 s, and
incubated on ice for 30 min, followed by centrifugation at 20,000g for 2 min at 4°C. The chromatin pellet
was washed twice with 600 μl of urea lysis buffer for 30 min on a rotator at 4°C. Chromatin pellet was
used for RNA extraction and next for DNAse treatment as described above. The chromatin-attached

RNA quality was tested by agarose gel electrophoresis and quantified with a Nanodrop 2000
spectrophotometer. cDNA synthesis was performed using a SuperScript III kit (Invitrogen).

smFISH

smFISH assay was performed as described previously[7] using a manually dissected radicle from dry
embryos. The Stellaris probes used for smFISH target the full *shDOG1* sequence, including intron 1 and
are labelled with Quasar670 fluorophore (Biosearch Technologies). The embryos were fixed,
permeabilised and hybridised with the probes as described[7]. The smFISH signals were gathered using a
widefield fluorescence microscope Olympus IX81 (Olympus) and a Hamamatsu Orca-R2 (C10600) CCD
camera. The xCellence software (Olympus) was used for image acquisition. Cells were manually
segmented using Napari[93] and the PartSeq plugin, which allows foci identification and classification
based on subcellular location[94]. Dots number and intensity were analysed using R scripts described
earlier[7].
1018

Author contributions

1020 S.P.S., and S.S. designed the experiments. S.P.S. carried out most of the experiments, M.M. and
1021 V.H.M. performed smFISH, and L.B. performed ChIDP-seq. M.K. conducted RNA-seq, K.R.
1022 performed 3'RACE-seq and obtained *msh-1 x esp1* line, M.W. and R.Y. obtained CRISPR *DOG1*
1023 mutant, Ł.S. and A.J. developed *pU1-70K::U1-70K-GFP u1-70k* lines. S.P.S and S.S. wrote the
1024 manuscript.

Declaration of interests

1026 K.R. and S.S. have equity interests in ElementZero Biolabs UG. S.S. also serve on the Scientific Advisory
1027 Board for ElementZero Biolabs UG. All other authors declare no competing interests.

Declaration of generative AI and AI-assisted technologies in the writing process

1029 While preparing this work, the authors used Grammarly to improve readability and language. After using
1030 this tool, the authors reviewed and edited the content as needed and took full responsibility for the
1031 publication's content.

Data availability
1033 The 3'RNA-seq data generated for this study have been deposited at the Gene Expression Omnibus
1034 (GEO) under the accession code GSE274757.

Declaration of interests

Declaration of generative AI and AI-assisted technologies in the writing process

References
1043 1.   Statello, L., Guo, C.-J., Chen, L.-L. & Huarte, M. Gene regulation by long non-coding RNAs and its

1044   biological functions. *Nat Rev Mol Cell Biol* **22**, 96–118 (2021).

1045   2.   Dror, I. *et al.* XIST directly regulates X-linked and autosomal genes in naive human pluripotent

1046   cells. *Cell* **187**, 110-129.e31 (2024).

1047   3.   Li, J. *et al.* Long noncoding RNA XIST: Mechanisms for X chromosome inactivation, roles in sex-

1048   biased diseases, and therapeutic opportunities. *Genes Dis* **9**, 1478–1492 (2022).

1049   4.   Arun, G., Aggarwal, D. & Spector, D. L. MALAT1 Long Non-Coding RNA: Functional Implications.

1050   *Noncoding RNA* **6**, 22 (2020).

1051   5.   Ariel, F. *et al.* R-Loop Mediated trans Action of the APOLO Long Noncoding RNA. *Molecular Cell*

1052   **77**, 1055-1065.e4 (2020).

1053   6.   Kindgren, P., Ard, R., Ivanov, M. & Marquardt, S. Transcriptional read-through of the long non-

1054   coding RNA SVALKA governs plant cold acclimation. *Nat Commun* **9**, 4561 (2018).

1055   7.   Montez, M. *et al.* Promoter-pervasive transcription causes RNA polymerase II pausing to boost

1056   DOG1 expression in response to salt. *EMBO J* **42**, e112443 (2023).

1057   8.   Zhao, Y. *et al.* Natural temperature fluctuations promote COOLAIR regulation of FLC. *Genes Dev*

1058   **35**, 888–898 (2021).

1059   9.   Swiezewski, S., Liu, F., Magusin, A. & Dean, C. Cold-induced silencing by long antisense transcripts

1060   of an Arabidopsis Polycomb target. *Nature* **462**, 799–802 (2009).

1061   10.   Rosa, S., Duncan, S. & Dean, C. Mutually exclusive sense–antisense transcription at FLC facilitates

1062   environmentally induced gene repression. *Nat Commun* **7**, 13031 (2016).

1063   11.   Yang, J. *et al.* Long noncoding RNA ARTA controls ABA response through MYB7 nuclear trafficking

1064   in Arabidopsis. *Developmental Cell* **58**, 1206-1217.e4 (2023).

1065   12.   Yatusevich, R. *et al.* Antisense transcription represses Arabidopsis seed dormancy QTL DOG1 to

1066   regulate drought tolerance. *EMBO Rep.* **18**, 2186–2196 (2017).

1067   13.   Wang, Y. *et al.* The noncoding RNA *HIDDEN TREASURE 1* promotes phytochrome B-dependent

1068   seed germination by repressing abscisic acid biosynthesis. *The Plant Cell* **35**, 700–716 (2023).

1069   14.   Yang, M. *et al.* In vivo single-molecule analysis reveals COOLAIR RNA structural diversity. *Nature*

1070   **609**, 394–399 (2022).

15. Yin, Y. *et al.* U1 snRNP regulates chromatin retention of noncoding RNAs. *Nature* **580**, 147–150

(2020).

16. Kondo, Y., Oubridge, C., van Roon, A.-M. M. & Nagai, K. Crystal structure of human U1 snRNP, a

small nuclear ribonucleoprotein particle, reveals the mechanism of 5' splice site recognition. *Elife*

4, e04986 (2015).

17. Goguel, V., Liao, X. L., Rymond, B. C. & Rosbash, M. U1 snRNP can influence 3'-splice site

selection as well as 5'-splice site selection. *Genes Dev* **5**, 1430–1438 (1991).

18. Almada, A. E., Wu, X., Kriz, A. J., Burge, C. B. & Sharp, P. A. Promoter directionality is controlled by

U1 snRNP and polyadenylation signals. *Nature* **499**, 360–363 (2013).

19. Mimoso, C. A. & Adelman, K. U1 snRNP increases RNA Pol II elongation rate to enable synthesis

of long genes. *Mol Cell* **83**, 1264-1279.e10 (2023).

20. So, B. R. *et al.* A Complex of U1 snRNP with Cleavage and Polyadenylation Factors Controls

Telescripting, Regulating mRNA Transcription in Human Cells. *Mol Cell* **76**, 590-599.e4 (2019).

21. Mangilet, A. F. *et al.* The Arabidopsis U1 snRNP regulates mRNA 3'-end processing. *Nat. Plants* **10**,

1514–1531 (2024).

22. Carrillo-Barral, N., Rodríguez-Gacio, M. D. C. & Matilla, A. J. Delay of Germination-1 (DOG1): A

Key to Understanding Seed Dormancy. *Plants (Basel)* **9**, 480 (2020).

23. Auge, G. A. *et al.* Secondary dormancy dynamics depends on primary dormancy status in

Arabidopsis thaliana. *Seed Sci. Res.* **25**, 230–246 (2015).

24. Graeber, K. *et al. DELAY OF GERMINATION 1* mediates a conserved coat-dormancy mechanism for

the temperature- and gibberellin-dependent control of seed germination. *Proc. Natl. Acad. Sci.*

U.S.A. **111**, (2014).

25. Footitt, S., Douterelo-Soler, I., Clay, H. & Finch-Savage, W. E. Dormancy cycling in *Arabidopsis*

seeds is controlled by seasonally distinct hormone-signaling pathways. *Proc. Natl. Acad. Sci.*

U.S.A. **108**, 20236–20241 (2011).

26. Alonso-Blanco, C., Bentsink, L., Hanhart, C. J., Vries, H. B. & Koornneef, M. Analysis of Natural

Allelic Variation at Seed Dormancy Loci of *Arabidopsis thaliana*. *Genetics* **164**, 711–729 (2003).

27. Bentsink, L., Jowett, J., Hanhart, C. J. & Koornneef, M. Cloning of DOG1, a quantitative trait locus

controlling seed dormancy in Arabidopsis. *Proc. Natl. Acad. Sci. U.S.A* **103**, 17042–17047 (2006).

28. Fedak, H. *et al.* Control of seed dormancy in Arabidopsis by a cis-acting noncoding antisense

transcript. *Proc. Natl. Acad. Sci. U.S.A.* **113**, E7846–E7855 (2016).

29. Nakabayashi, K., Bartsch, M., Ding, J. & Soppe, W. J. J. Seed Dormancy in Arabidopsis Requires

Self-Binding Ability of DOG1 Protein and the Presence of Multiple Isoforms Generated by

Alternative Splicing. *PLoS Genet* **11**, e1005737 (2015).

30. Cyrek, M. *et al.* Seed Dormancy in Arabidopsis Is Controlled by Alternative Polyadenylation of

DOG1. *Plant Physiol.* **170**, 947–955 (2016).

31. Krzyszton, M. *et al.* Dormancy heterogeneity among Arabidopsis thaliana seeds is linked to

individual seed size. *Plant Communications* 100732 (2023) doi:10.1016/j.xplc.2023.100732.

32. Kowalczyk, J. *et al.* Alternative Polyadenylation of the Sense Transcript Controls Antisense

Transcription of DELAY OF GERMINATION 1 in Arabidopsis. *Molecular Plant* **10**, 1349–1352

(2017).

33. Kendall, S. L. *et al.* Induction of Dormancy in *Arabidopsis* Summer Annuals Requires Parallel

Regulation of *DOG1* and Hormone Metabolism by Low Temperature and CBF Transcription

Factors. *Plant Cell* **23**, 2568–2580 (2011).

34. Rodríguez-Gacio, M. D. C., Matilla-Vázquez, M. A. & Matilla, A. J. Seed dormancy and ABA

signaling: The breakthrough goes on. *Plant Signaling & Behavior* **4**, 1035–1048 (2009).

35. Park, S.-Y. *et al.* Abscisic Acid Inhibits Type 2C Protein Phosphatases via the PYR/PYL Family of

START Proteins. *Science* **324**, 1068–1071 (2009).

36. Baek, W., Lim, C. W., Luan, S. & Lee, S. C. The RING finger E3 ligases PIR1 and PIR2 mediate PP2CA

degradation to enhance abscisic acid response in Arabidopsis. *The Plant Journal* **100**, 473–486

(2019).

37. Née, G. *et al.* DELAY OF GERMINATION1 requires PP2C phosphatases of the ABA signalling

pathway to control seed dormancy. *Nat Commun* **8**, 72 (2017).

38. Nishimura, N. *et al.* Control of seed dormancy and germination by DOG1-AHG1 PP2C

phosphatase complex via binding to heme. *Nat Commun* **9**, 2132 (2018).

39. Zhang, S. *et al.* New insights into Arabidopsis transcriptome complexity revealed by direct

sequencing of native RNAs. *Nucleic Acids Research* **48**, 7700–7711 (2020).

40. Thomas, Q. A. *et al.* Transcript isoform sequencing reveals widespread promoter-proximal

transcriptional termination in Arabidopsis. *Nat Commun* **11**, 2589 (2020).

41. Thieffry, A. *et al.* Characterization of Arabidopsis thaliana Promoter Bidirectionality and Antisense

RNAs by Inactivation of Nuclear RNA Decay Pathways. *Plant Cell* **32**, 1845–1867 (2020).

42. Čermák, T. *et al.* A Multipurpose Toolkit to Enable Advanced Genome Engineering in Plants. *Plant

Cell* **29**, 1196–1217 (2017).

43. Jupe, F. *et al.* The complex architecture and epigenomic impact of plant T-DNA insertions. *PLoS

Genet* **15**, e1007819 (2019).

44. Xu, R. *et al.* The 73 kD Subunit of the cleavage and polyadenylation specificity factor (CPSF)

complex affects reproductive development in Arabidopsis. *Plant Mol Biol* **61**, 799–815 (2006).

45. Funck, D., Clauß, K., Frommer, W. B. & Hellmann, H. A. The Arabidopsis CstF64-like RSR1/ESP1

protein participates in glucose signaling and flowering time control. *Front Plant Sci* **3**, (2012).

46. Cortazar, M. A. *et al.* Control of RNA Pol II Speed by PNUTS-PP1 and Spt5 Dephosphorylation

Facilitates Termination by a "Sitting Duck Torpedo" Mechanism. *Molecular Cell* **76**, 896-908.e4

(2019).

47. Laugesen, A., Højfeldt, J. W. & Helin, K. Molecular Mechanisms Directing PRC2 Recruitment and

H3K27 Methylation. *Mol Cell* **74**, 8–18 (2019).

48. Blackledge, N. P. & Klose, R. J. The molecular principles of gene regulation by Polycomb

repressive complexes. *Nat Rev Mol Cell Biol* **22**, 815–833 (2021).

49. Krzyszton, M. *et al.* Single seeds exhibit transcriptional heterogeneity during secondary dormancy

induction. *Plant Physiology* **190**, 211–225 (2022).

50. Buijs, G. A Perspective on Secondary Seed Dormancy in Arabidopsis thaliana. *Plants* **9**, 749

(2020).

51. Ibarra, S. E. *et al.* Molecular mechanisms underlying the entrance in secondary dormancy of

Arabidopsis seeds: Mechanisms underlying secondary dormancy. *Plant Cell Environ.* **39**, 213–221

(2016).

52. Footitt, S., Huang, Z., Clay, H. A., Mead, A. & Finch-Savage, W. E. Temperature, light and nitrate

sensing coordinate A rabidopsis seed dormancy cycling, resulting in winter and summer annual

phenotypes. *The Plant Journal* **74**, 1003–1015 (2013).

53. Wrona, M. *et al.* The BRAHMA-associated SWI/SNF chromatin remodeling complex controls

Arabidopsis seed quality and physiology. *Plant Physiology* **197**, kiae642 (2024).

54. Wang, H. *et al.* H3K4me3 regulates RNA polymerase II promoter-proximal pause-release. *Nature*

615, 339–348 (2023).

55. Chu, C., Quinn, J. & Chang, H. Y. Chromatin isolation by RNA purification (ChIRP). *J Vis Exp* 3912

(2012) doi:10.3791/3912.

56. Déjardin, J. & Kingston, R. E. Purification of Proteins Associated with Specific Genomic Loci. *Cell*

136, 175–186 (2009).

57. Sajeev, N., Koornneef, M. & Bentsink, L. A commitment for *life:* Decades of unraveling the

molecular mechanisms behind seed dormancy and germination. *The Plant Cell* koad328 (2024)

doi:10.1093/plcell/koad328.

58. Chen, M. & Penfield, S. Feedback regulation of COOLAIR expression controls seed dormancy and

flowering time. *Science* **360**, 1014–1017 (2018).

59. Nielsen, M. *et al. COOLAIR*   and PRC2 function in parallel to silence   *FLC*   during

vernalization. *Proc. Natl. Acad. Sci. U.S.A.* **121**, (2024).

60. Jin, Y., Ivanov, M., Dittrich, A. N., Nelson, A. D. & Marquardt, S. LɴᴄRNA FLAIL affects alternative

splicing and represses flowering in *Arabidopsis*. *The EMBO Journal* **42**, e110921 (2023).

61. Née, G. *et al.* DELAY OF GERMINATION1 requires PP2C phosphatases of the ABA signalling

pathway to control seed dormancy. *Nat Commun* **8**, 72 (2017).

62. Krzyszton, M. *et al.* Single seeds exhibit transcriptional heterogeneity during secondary dormancy

induction. *Plant Physiol.* **190**, 211–225 (2022).

63. Nakabayashi, K. *et al.* The Time Required for Dormancy Release in *Arabidopsis* Is Determined by

DELAY OF GERMINATION1 Protein Levels in Freshly Harvested Seeds. *Plant Cell* **24**, 2826–2838

(2012).

64. Thatcher, L. F. *et al.* The Arabidopsis RNA Polymerase II Carboxyl Terminal Domain (CTD)

Phosphatase-Like1 (CPL1) is a biotic stress susceptibility gene. *Sci Rep* **8**, 13454 (2018).

65. Guan, Q., Yue, X., Zeng, H. & Zhu, J. The Protein Phosphatase RCF2 and Its Interacting Partner

NAC019 Are Critical for Heat Stress–Responsive Gene Regulation and Thermotolerance in

Arabidopsis. *Plant Cell* **26**, 438–453 (2014).

66. Qi, Y. *et al.* The over-expression of a chrysanthemum gene encoding an RNA polymerase II CTD

phosphatase-like 1 enzyme enhances tolerance to heat stress. *Hortic Res* **5**, 37 (2018).

67. Krzyszton *et al.* Single seeds exhibit transcriptional heterogeneity during secondary dormancy

induction. *Accepted in Plant Physiology* (2022).

68. Ali, F., Qanmber, G., Li, F. & Wang, Z. Updated role of ABA in seed maturation, dormancy, and

germination. *J Adv Res.* **35**, 199–214 (2022).

69. Kim, D.-H. & Sung, S. Vernalization-Triggered Intragenic Chromatin Loop Formation by Long

Noncoding RNAs. *Developmental Cell* **40**, 302-312.e4 (2017).

70. Guo, Y. *et al.* Long noncoding RNAs heat shock RNA omega nucleates TBPH and promotes

intestinal stem cell differentiation upon heat shock. *iScience* **27**, 109732 (2024).

71. Ji, Q., Zong, X., Mao, Y. & Qian, S.-B. A heat shock–responsive lncRNA *Heat* acts as a HSF1-

directed transcriptional brake via m^{6}A modification. *Proc. Natl. Acad. Sci. U.S.A.* **118**,

e2102175118 (2021).

72. Guo, C.-J. *et al.* Distinct Processing of lncRNAs Contributes to Non-conserved Functions in Stem

Cells. *Cell* **181**, 621-636.e22 (2020).

73. Bridges, M. C., Daulagala, A. C. & Kourtidis, A. LNCcation: lncRNA localization and function. *J Cell

Biol* **220**, e202009045 (2021).

74. Shukla, C. J. *et al.* High-throughput identification of RNA nuclear enrichment sequences. *EMBO J*

37, e98452 (2018).

75. Lubelsky, Y. & Ulitsky, I. Sequences enriched in Alu repeats drive nuclear localization of long RNAs

in human cells. *Nature* **555**, 107–111 (2018).

76. Grosch, M. *et al.* Nucleus size and DNA accessibility are linked to the regulation of paraspeckle

formation in cellular differentiation. *BMC Biol* **18**, 42 (2020).

77. Strehle, M. & Guttman, M. Xist drives spatial compartmentalization of DNA and protein to

orchestrate initiation and maintenance of X inactivation. *Current Opinion in Cell Biology* **64**, 139–

147 (2020).

78. Zhao, X. *et al.* Global identification of Arabidopsis lncRNAs reveals the regulation of MAF4 by a

natural antisense RNA. *Nat Commun* **9**, 5056 (2018).

79. Csorba, T., Questa, J. I., Sun, Q. & Dean, C. Antisense COOLAIR mediates the coordinated

switching of chromatin states at FLC during vernalization. *Proc. Natl. Acad. Sci. U.S.A.* **111**,

16160–16165 (2014).

80. Sun, Q., Csorba, T., Skourti-Stathaki, K., Proudfoot, N. J. & Dean, C. R-loop stabilization represses

antisense transcription at the Arabidopsis FLC locus. *Science* **340**, 619–621 (2013).

81. Xu, W. *et al.* The R-Loop Atlas of Arabidopsis Development and Responses to Environmental

Stimuli. *Plant Cell* **32**, 888–903 (2020).

82. Manavella, P. A. *et al.* Fast-Forward Genetics Identifies Plant CPL Phosphatases as Regulators of

miRNA Processing Factor HYL1. *Cell* **151**, 859–870 (2012).

83. Simpson, G. G., Dijkwel, P. P., Quesada, V., Henderson, I. & Dean, C. FY Is an RNA 3′ End-

Processing Factor that Interacts with FCA to Control the Arabidopsis Floral Transition. *Cell* **113**,

777–787 (2003).

84. Xing, D., Zhao, H., Xu, R. & Li, Q. Q. Arabidopsis PCFS4, a homologue of yeast polyadenylation

factor Pcf11p, regulates FCA alternative processing and promotes flowering time. *The Plant*

Journal **54**, 899–910 (2008).

85. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357–

359 (2012).

86. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

87. Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. deepTools: a flexible platform for

exploring deep-sequencing data. *Nucleic Acids Research* **42**, W187–W191 (2014).

88. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

89. Smith, T., Heger, A. & Sudbery, I. UMI-tools: Modelling sequencing errors in Unique Molecular

Identifiers to improve quantification accuracy. *Genome Res.* **27**, 491–499 (2016).

90. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.

Bioinformatics **26**, 841–842 (2010).

91. Warkocki, Z. *et al.* Uridylation by TUT4/7 Restricts Retrotransposition of Human LINE-1s. *Cell* **174**,

1537-1548.e29 (2018).

92. Alpern, D. *et al.* BRB-seq: ultra-affordable high-throughput transcriptomics enabled by bulk RNA

barcoding and sequencing. *Genome Biol* **20**, 71 (2019).

93. Sofroniew, N. *et al.* napari: a multi-dimensional image viewer for Python. Zenodo

https://doi.org/10.5281/ZENODO.6597502 (2022).

94. Bokota, G. *et al.* PartSeg: a tool for quantitative feature extraction from 3D microscopy images for

dummies. *BMC Bioinformatics* **22**, 72 (2021).

1247