Lysine Crotonylation Acts as an Epigenetic Switch for Glutamate Neurotransmission and Spatial Memory

Utsav Mukherjee[1,2,3], Budhaditya Basu[1,2], Pratyush Suryavanshi [2,7], Stacy E. Beyer[1,2], Saaman Ghodsi[1,2], Nathan Robillard[1,2], Yann Vanrobaeys[1,2,4], Eric B. Taylor[5,6], Joseph Glykys[2,7], Ted Abel[1,2*], and Snehajyoti Chatterjee[1,2*]

[1]Department of Neuroscience and Pharmacology, Carver College of Medicine, University of Iowa, Iowa City, IA 52242, United States

[2]Iowa Neuroscience Institute, University of Iowa, Iowa City, IA 52242, United States

[3]Interdisciplinary Graduate Program in Neuroscience, University of Iowa, Iowa City, IA 52242, United States

[4]Interdisciplinary Graduate Program in Genetics, University of Iowa, Iowa City, IA 52242, United States

[5]Fraternal Order of Eagles Diabetes Research Center, University of Iowa, Iowa City, IA 52242, United States

[6]Department of Molecular Physiology and Biophysics, Carver College of Medicine, University of Iowa, Iowa City, IA, 52242, United States

[7]Stead Family Department of Pediatrics, Carver College of Medicine, University of Iowa, Iowa City, IA, 52242, United States

*Corresponding authors: ted-abel@uiowa.edu and snehajyoti-chatterjee@uiowa.edu

Keywords: epigenetics, acylation, memory, chromatin plasticity, gene expression, glutamate

Running title: Histone crotonylation regulates long-term memory and glutamate neurotransmission.

Abstract

Histone post-translational modifications (PTMs), particularly lysine acetylation (Kac), are critical
epigenetic regulators of transcriptional programs underlying long-term memory consolidation. Beyond
Kac, several other non-acetyl acylations have been identified with the ability to regulate transcription;
however, their role in memory consolidation remains unknown. Here, we identify histone lysine
crotonylation (Kcr) as a molecular switch for long-term memory and glutamatergic neurotransmission.
We find that spatial learning induces distinct spatiotemporal patterns of Kcr across hippocampal
subregions, and that Kcr stimulates learning-induced gene expression. Through genetic and
pharmacological manipulations, we show that reducing hippocampal Kcr levels impairs memory, while
increasing Kcr enhances long-term memory. Single-nuclei transcriptome and chromatin accessibility
analyses reveals that Kcr facilitates activation of genes regulating glutamate signaling. Cell-cell
communication analysis further infers that Kcr enhancement strengthens glutamatergic signaling within
principal hippocampal neurons. Real-time fluorescence imaging with genetically encoded sensors
functionally validates our multiomics and computational findings—demonstrating the role of Kcr in
regulating presynaptic glutamate release and neuronal activity. In summary, our findings elucidate a
novel mechanism underlying long-term memory consolidation and excitatory neurotransmission, linking
epigenetic events to synaptic function and behavior.

Introduction

Histone PTMs play a key role in the epigenetic regulation of gene expression during memory
consolidation[1-5]. Short-chain acyl modifications on histone lysine residues are well-known PTMs that
regulate several biological processes[6-8]. Among these, histone lysine acetylation (Kac) has been
extensively studied for its role in memory storage. Altering histone Kac levels by selectively
manipulating the 'writers' (lysine acetyltransferases, KATs) and 'erasers' (lysine deacetylases, KDACs)
of histone Kac has been shown to impact hippocampal memory storage[2,9,10]. Interestingly, studies have
revealed the promiscuous nature of the majority of the writers and erasers for histone Kac—
demonstrating that these enzymes can also catalyze the incorporation or removal of non-acetyl acyl
moieties on histone lysine residues[6,11,12]. This raises the question of whether the impact of manipulating
KATs and KDACs on memory consolidation, as demonstrated in previous studies[13-18], could be
attributed to underlying changes in non-acetyl histone acylations besides Kac. Elucidating the functional
relevance of non-acetyl histone acylations in long-term memory is therefore essential for advancing our
conceptual understanding of epigenetic mechanisms underlying memory consolidation.

Advancements in mass spectrometry and biochemical approaches have led to the identification of
several novel histone lysine acylations that are structurally distinct from histone Kac[19,20]. Among these,
histone lysine crotonylation (Kcr), an evolutionarily conserved non-acetyl acylation, has gained
significant attention in recent years for its role in transcription regulation[20-26]. Kcr is enriched on active
gene promoters[20], and Kcr-mediated regulation of gene expression has been implicated in diverse
biological functions, including DNA damage repair[21], acute kidney injury[25], spermatogenesis[22], nerve-
injury-induced-neuropathic pain[26], and neural stem cell differentiation[23]. Interestingly, studies in cell-free
systems have demonstrated that the lysine acyltransferase p300 stimulates histone Kcr-mediated
transcription to a significantly higher magnitude than histone Kac[12]. Furthermore, several histone
'reader' proteins—initially characterized as Kac readers— exhibit a higher binding affinity for
crotonylated lysine residues compared to acetylated lysine moieties, and this preferential binding is
thought to underlie the ability of the readers to regulate transcription[27]. We and others have shown that
precisely timed transcriptional events, induced upon spatial learning, are critical for long-term memory
consolidation[28-31]. Although Kcr has been well established as a transcriptional activator, its physiological
relevance to memory consolidation—and signatures of Kcr-dependent gene expression programs that
underlie hippocampal memory storage—remain elusive.

The availability of metabolic precursors dictates the landscape of histone acylations[6]. Accordingly,
changes in intracellular crotonyl-CoA levels have been shown to alter histone Kcr levels, with direct

consequences in gene transcription. Increasing crotonyl-CoA levels through crotonate administration
elevates Kcr and stimulates gene transcription[12,23], whereas depletion of Kcr via manipulation of the
crotonyl-CoA hydratase chromodomain Y-like protein (CDYL) represses transcription[24]. Interestingly, in
addition to writers and erasers, histone Kac and histone Kcr share enzymes that synthesize their
respective metabolic precursors. The enzyme Acetyl-CoA synthetase 2 (ACSS2) synthesizes both
acetyl CoA and crotonyl-CoA[12,32]. We previously demonstrated that ACSS2 is a critical regulator of
neuronal histone Kac and spatial memory consolidation in the hippocampus[33]. Silencing ACSS2 levels
reduces histone Kac and impairs activity-dependent transcription of memory-related genes[33]. Notably,
knockdown of ACSS2 has also been shown to reduce Kcr levels and repress Kcr-regulated
inflammatory gene transcription[12]. However, whether ACSS2-mediated regulation of histone Kcr
constitutes a critical molecular mechanism underlying long-term memory consolidation remains
unknown. Altogether, these findings raise the possibility that metabolic coupling of crotonyl-CoA
availability to Kcr-dependent chromatin plasticity represents a previously unrecognized mechanism
linking neuronal metabolism to memory storage and learning-induced transcriptional events.

Here, we provide evidence demonstrating the role of Kcr as an epigenetic rheostat for hippocampal
memory storage and glutamatergic neurotransmission. We show that spatial learning in mice increases
Kcr levels in the dorsal hippocampus during the early temporal window of memory consolidation. Next,
we demonstrate that depleting histone Kcr levels in the hippocampus by overexpressing the crotonyl-
CoA hydratase CDYL leads to long-term spatial memory impairments. Conversely, increasing histone
Kcr levels by oral administration of crotonate, a precursor of crotonyl-CoA, enhances hippocampus-
dependent long-term memory. Using single-nuclei transcriptomic and chromatin accessibility analyses,
we define subregion-specific molecular signatures of Kcr regulation across the hippocampus—
revealing that Kcr enhancement selectively impacts genes encoding key regulators of glutamatergic
neurotransmission and identifying enriched transcription factor motifs at their promoters that may
govern their expression. We examine functional consequences of increasing histone Kcr by performing
ex vivo real-time fluorescent imaging using genetically encoded glutamate and calcium sensors—
demonstrating that Kcr enhancement augments presynaptic glutamate release and neuronal activity.
Together, our findings reveal, for the first time, the role of a non-acetyl acyl modification in long-term
memory and excitatory neurotransmission. More importantly, our study offers mechanistic evidence of
how epigenetic tuning of transcriptional programs by Kcr translates into changes in neuronal function
and behavioral outcomes—linking epigenetic events in the nucleus to synaptic activity.

Results

Spatial training induces lysine crotonylation in discrete hippocampal subregions

KATs catalyze the addition of crotonyl groups on histone lysine residues using crotonyl-CoA as a

cofactor, and crotonyl-CoA levels are regulated by the enzymes ACSS2 and CDYL (**Fig. 1a**). We first

investigated whether spatial learning induces changes in histone Kcr in the hippocampus, similar to the

well-established increases in histone Kac following spatial training[2,9,34]. We trained wild-type (WT) mice

in a spatial object recognition (SOR) task and examined Kcr levels in core histones extracted from the

dorsal hippocampus at different time-intervals (0.5-, 1-, and 2-hrs) post training (**Supplementary Fig.**

1a-b). We focused on the dorsal hippocampus in all our experiments, given extensive evidence for its

critical role in spatial memory consolidation[30,35-37]. Moreover, in the absence of prior knowledge

regarding specific crotonylated histone lysine residues that respond to learning, we measured pan-

lysine crotonylation (Kcr) to assess the global landscape of histone Kcr during spatial memory

consolidation. Using an antibody that has been shown to selectively detect crotonylated lysine

residues[12] and has been extensively used to characterize Kcr in several other studies[24,38-41], we

observed that histone Kcr levels significantly increased at the 1-hr timepoint after SOR training and

returned to baseline levels at the 2-hr timepoint (**Supplementary Fig. a-b**). Next, we performed

immunofluorescence using the pan-Kcr antibody to map the spatial profiles of Kcr induction in the

hippocampus following SOR training (**Fig. 1b-d**). Quantitative analysis of the nuclear Kcr levels

revealed significant increases in the CA1 pyramidal layer and the upper blade of dentate gyrus (DG)

(**Fig. 1c-d**), and in the subiculum (**Supplementary Fig. 2a-b**) at the 1-hr timepoint after training in the

SOR task, whereas no significant changes in the nuclear Kcr levels were observed in hippocampal

subregions CA3 and the lower blade of DG (**Fig. 1c-d**). These results identify histone Kcr to exhibit

distinct spatiotemporal dynamics in the dorsal hippocampus following spatial learning.

Enrichment of crotonylation on gene promoters correlates with higher magnitude of learning-

induced gene expression

To determine whether Kcr has a role in regulating learning-induced gene expression, we performed

Chromatin Immunoprecipitation followed by sequencing (ChIP-Seq) using a pan-Kcr antibody on dorsal

hippocampal tissue from mice that were either housed in their homecage or trained in the SOR task

(**Fig. 1b, e**). We found that spatial learning led to a significant enrichment of Kcr on gene promoters

(**Fig. 1e**), consistent with previous reports linking the presence of Kcr marks on gene promoters to

transcriptional activation. Next, we determined the relationship between Kcr and a transcriptionally

active Kac mark (H3K27ac) on learning-induced gene promoters. We compared the enrichment of Kcr

and H3K27ac[42] on the promoters of genes upregulated after learning in the dorsal hippocampus

following the SOR task[30]. Our analyses revealed that learning-induced genes enriched with H3K27ac

and pan-Kcr exhibit a significantly higher magnitude of expression (fold change) compared to genes

enriched with only H3K27ac **(Fig. 1f)**. These findings not only underscore the transcriptional activation

potential of Kcr but also reveal its role in regulating learning-dependent gene expression.

Reduction of hippocampal Kcr levels impairs long-term spatial memory

The crotonyl-CoA hydratase Chromodomain Y-like protein (CDYL) hydrolyzes crotonyl-CoA to beta-

hydroxy butyryl-CoA (**Fig. 1a**), thereby reducing histone Kcr levels[24]. To date, CDYL remains the only

bona fide negative regulator of Kcr that does not affect other histone acylations[39]—presenting us with a

unique strategy to selectively manipulate histone Kcr levels and examine its impact on memory

consolidation. Interestingly, neuronal activity has been shown to reduce CDYL levels[43], prompting us to

investigate whether spatial learning would similarly impact hippocampal CDYL levels. We performed

immunofluorescence on brain sections obtained from SOR-trained WT mice to examine spatial patterns

of CDYL protein levels following learning across hippocampal subregions (**Supplementary Fig. 3a-d**).

Hippocampal subregions CA1, subiculum, and the upper blade of DG exhibited significant

downregulation of nuclear CDYL levels 1-hr after SOR training, whereas nuclear CDYL levels remained

unchanged in the hippocampal subregions CA3 and the lower blade of DG (**Supplementary Fig. 3a-d).**

The downregulation of nuclear CDYL levels after learning in CA1, subiculum, and the upper blade of

DG offers a molecular explanation for the corresponding learning-induced increase in nuclear Kcr levels

observed within these hippocampal subregions.

Next, to determine whether histone Kcr has a role in hippocampal long-term memory, we designed a

genetic strategy to selectively modulate Kcr in the dorsal hippocampus by manipulating levels of CDYL.

We injected an adeno-associated virus (AAV) in WT mice to overexpress CDYL (AAV$_9$-CaMKIIα-CDYL-

V5) in excitatory neurons of the dorsal hippocampus (**Fig. 1g**). Immunofluorescence performed two

weeks after viral infusion confirmed the expression of CDYL-V5 across the dorsal hippocampal

principal neuronal layers (**Fig. 1h**). We observed significantly reduced Kcr levels in the hippocampal

areas CA1 and CA3 in CDYL-V5-expressing mice compared to EGFP controls (**Fig. 1i-j**), thus

validating the efficacy of our viral-based approach. Next, we evaluated long-term spatial memory in

mice infused with AAV-CDYL-V5 in the dorsal hippocampus. Mice were trained in the SOR task with

objects that were placed in specific spatial locations within a SOR arena. To test for long-term spatial

memory, we performed a retrieval task 24-hrs later with one of the objects displaced to a novel spatial

location (**Fig. 1k**). Control AAV-infused mice (AAV$_9$-CaMKIIα-EGFP) exhibited a higher preference

towards the displaced object during the test session, suggesting intact memory consolidation.

Conversely, CDYL-infused mice (AAV$_9$-CaMKIIα-CDYL-V5) failed to exhibit any preference towards the

displaced object during the test session (**Fig. 1I**). Collectively, these findings suggest a critical role of

hippocampal Kcr in spatial memory consolidation.

Pharmacologically increasing Kcr enhances long-term memory

Given that a reduction in hippocampal Kcr levels significantly impairs long-term memory, we next asked

whether increasing Kcr levels could enhance long-term memory. Administration of crotonate, a short-

chain fatty acid, has been shown to elevate histone Kcr[12,23] and stimulate gene expression[23] by

increasing Crotonyl-CoA levels[12,23]. Therefore, we implemented this strategy to pharmacologically

increase histone Kcr levels in mice. We administered two different doses of crotonate (50 and 200

mg/kg) to WT mice by oral gavage and examined histone Kcr levels in the dorsal hippocampus 1-hr

after drug administration (**Fig. 2a**). We observed a significant increase in histone Kcr following 200

mg/kg dose of crotonate, whereas the 50 mg/kg dose of crotonate failed to enhance histone Kcr levels

compared to vehicle (saline)-treated mice (**Fig. 2b**). Next, to study the impact of higher histone Kcr

levels on long-term memory, we trained WT mice in a weak-learning (sub-threshold) SOR paradigm[44]

to evaluate long-term memory enhancement. The subthreshold variant of the SOR task was designed

to assess enhancements in long-term memory, as weak SOR training protocols typically do not result in

long-term memory[44-46]. Mice were administered 50mg/kg of crotonate, 200 mg/kg of crotonate, or

vehicle via oral gavage immediately after the completion of the training session. When tested 24 hrs

later, both the vehicle-treated and 50 mg/kg crotonate-treated mice did not show any preference

towards the displaced object, whereas mice administered with 200 mg/kg crotonate showed a

significant preference towards the displaced object (**Fig. 2c**). Consistent with our SOR findings, we

observed a significantly enhanced freezing response upon crotonate administration when mice were

tested 24 hrs following a sub-threshold contextual fear conditioning (CFC) task (**Fig. 2d**). Our

behavioral findings demonstrate that increases in histone Kcr augments hippocampal long-term spatial

and contextual fear memories, further underscoring the functional relevance of Kcr in long-term

memory consolidation.

Neuronal ACSS2 is a key regulator of Kcr-dependent memory enhancement

The metabolic enzyme ACSS2 has been shown to synthesize acetyl-CoA. Previous work has

demonstrated that ACSS2 plays a critical role in long-term memory by coupling acetyl-CoA synthesis to

Kac-mediated transcription of learning-responsive genes[33]. However, ACSS2 can also synthesize

crotonyl-CoA from the short-chain fatty acid crotonate in mammalian cells[12] (**Fig. 1a, Supplementary**

Fig. 4a). This prompted us to investigate whether ACSS2 regulates the increases in histone Kcr and

enhancements in long-term memory seen after the administration of crotonate. We injected an AAV
vector (AAV$_9$-CaMKIIα-Cre-EGFP) into the dorsal hippocampus of male ACSS2$^{f/f}$ mice[47] to achieve a
conditional knockdown of ACSS2 expression selectively in the excitatory neurons of the dorsal
hippocampus (**Supplementary Fig. 4b**). Immunofluorescence performed on brain sections obtained
from the ACSS2 conditional knockout mice (ACSS2 cKO mice; ACSS2 $^{f/f}$:CaMKIIα-Cre-EGFP) revealed
the expression of Cre across hippocampal subregions (**Supplementary Fig. 4c**). Western blot
analyses confirmed the downregulation of ACSS2 in the dorsal hippocampus of ACSS2 cKO mice
(**Supplementary Fig. 4d-e**). We then examined the impact of crotonate administration on histone Kcr
levels in ACSS2 cKO mice. We administered crotonate (200 mg/kg) or vehicle (saline) via oral gavage
immediately after a sub-threshold SOR training session, and quantified histone Kcr levels in the dorsal
hippocampus 1-hr after training. We found that crotonate treatment failed to enhance hippocampal
histone Kcr levels in ACSS2 cKO mice (**Supplementary Fig. 4f-g**), despite administering a dose of
crotonate (200 mg/kg) that was sufficient to induce histone Kcr levels in WT mice (**Fig. 2b**). As
crotonate administration failed to enhance Kcr levels in ACSS2 cKO mice, we hypothesized that
crotonate treatment would also fail to elicit long-term memory enhancement in ACSS2 cKO mice. To
test this hypothesis, we trained ACSS2 cKO mice in a sub-threshold learning SOR paradigm,
administered vehicle or crotonate (200 mg/kg) immediately after training, and examined their long-term
memory 24 hrs after training (**Supplementary Fig. 4h**). Our behavioral assessment confirmed that
crotonate treatment in ACSS2 cKO mice had no impact in long-term memory compared to the vehicle-
treated group (**Supplementary Fig. 4i**). Taken together, these results demonstrate that ACSS2 is
critical in mediating the molecular impact of crotonate-mediated increases in Kcr on memory
consolidation.

Single-nuclei multiomics reveals the molecular signatures of Kcr underlying memory
enhancement

Next, we implemented a single-nuclei multiomics (snRNA-seq and snATAC-seq) strategy to elucidate
the molecular mechanisms underlying crotonate-mediated memory enhancement. Mice were trained in
SOR using a sub-threshold learning paradigm, and oral administration of crotonate (200 mg/kg) or
vehicle (saline) was performed immediately after training. 1-hr after crotonate administration, the dorsal
hippocampus was harvested, and nuclei were isolated for single-nuclei multiomics (**Fig. 2e**). Cell type
clustering identified well-distinguished clusters comprising excitatory and inhibitory neurons, as well as
clusters representing non-neuronal cell populations (**Fig. 2f**). Excitatory neurons were further classified
into CA1, CA3, subiculum (Sub), dentate gyrus, and excitatory cortical neurons based on cell-type-
specific marker gene expression (**Fig. 2f, Supplementary Fig. 5a**). Notably, crotonate and vehicle

treatment groups exhibited the same number of cell-type-specific clusters and comparable proportion of

cell-types (**Supplementary Fig. 5b-c, Supplemental table 1**). Next, we performed differential gene

expression (DEG) utilizing the snRNA data along with differential accessibility regions (DARs) on the

chromatin from the snATAC data of each cluster, comparing crotonate and vehicle treated groups. For

downstream analysis, we used the DEGs (FDR<0.05, and absolute $\log_2$ fold change > 0.2) exhibiting

concordant DARs (FDR<0.05, and absolute $\log_2$ fold change > 0.2) across their promoter and gene

body (**Fig. 2g, Supplemental table 2**). We found that excitatory neurons in the CA1 hippocampal area

showed the highest number of genes impacted by crotonate, with 205 upregulated genes showing

increased chromatin accessibility and 36 downregulated genes showing reduced chromatin

accessibility (**Fig. 2g**). Other cell types that showed a strong impact of crotonate on gene expression

and chromatin accessibility were CA3 excitatory neurons (118 upregulated genes with increased

chromatin accessibility and 44 downregulated genes with reduced chromatin accessibility), subiculum

(53 upregulated genes with increased chromatin accessibility and 18 downregulated genes with

reduced chromatin accessibility), and the DG (27 upregulated genes with increased chromatin

accessibility and one downregulated gene with reduced chromatin accessibility) (**Fig. 2g**). In addition to

the principal neuronal clusters (CA1, CA3, subiculum, and DG), our analysis identified discrete

molecular changes in inhibitory neurons (11 upregulated genes with increased DARs and 3

downregulated genes with reduced DARs) (**Fig. 2g**). Among the non-neuronal cells, oligodendrocytes

showed the highest number of overlapping DEGs and DARs (24 upregulated and 27 downregulated)

(**Fig. 2g**). Notably, we found a significant correlation between the significant differentially expressed

genes and genes with significant differential chromatin accessibility in hippocampal subregions CA1

and CA3 (**Supplementary Fig. 6**). Next, utilizing an Upset plot, we compared unique and overlapping

upregulated genes with concordant DARs in each cell-type. We found that CA1, CA3, subiculum, and

DG showed a common upregulation of 2 genes (**Fig. 2h**). Notably, CA1, CA3, and subiculum showed

17 commonly upregulated genes, CA1, CA3, and DG showed a common upregulation of 4 genes, and

CA1 and CA3 showed 22 commonly upregulated genes following crotonate treatment (**Fig. 2h**). Like

the cell-type expression patterns observed with upregulated DEGs with increased DARs, there was

some overlap in the expression of downregulated genes with reduced DARs between cell-types. We

identified 23 downregulated genes specific to CA1, 32 exclusively in CA3, and 12 selectively in the

subiculum (**Fig. 2i**). Four genes were commonly downregulated across CA1, CA3, and the subiculum,

while 8 downregulated genes were shared between CA1 and CA3 (**Fig. 2i**). Because the manipulation

of Kcr levels selectively in excitatory neurons was found to impact long-term memory (**Fig. 1g-l,**

Supplementary Fig. 4), our subsequent analyses focused on DEGs within the neuronal populations of

CA1, CA3, subiculum, and DG that also exhibit DARs.

To better understand the impact of Kcr enhancement on the molecular signatures of principal neuronal
sub-types in the dorsal hippocampus, we performed Gene Ontology (GO: Molecular Function)
overrepresentation analysis to identify the pathways enriched among the DEGs with DARs following
crotonate treatment. We found that crotonate treatment impacted the expression of genes that were
predominantly associated with regulating neurotransmission and synaptic function across the cell
clusters (**Fig. 3a-d)**. Among the most significant pathways found enriched following crotonate
administration, cell adhesion molecule binding was commonly enriched in subregions CA1, CA3, and
subiculum, whereas distinct pathways attributed to ion channel activity were commonly enriched in CA1
and CA3 (**Fig. 3a-d, Supplemental table 3**). In contrast, pathways attributed to PDZ domain binding
and calcium-dependent phospholipid binding were found enriched in the subiculum, whereas DG
exhibited an enrichment of pathways related to protein kinase C activity and serine/threonine kinase
activity. (**Fig. 3a-d**). Among the downregulated DEGs with reduced chromatin accessibility, we found
pathways attributed to ephrin receptor activity, calcium ion transmembrane transporter activity, and
phosphoric diester hydrolase activity downregulated in CA3, whereas pathways involved in cadherin
binding and beta-catenin binding were downregulated in DG (**Supplementary Fig. 7**).

To gain a mechanistic understanding of how crotonate-mediated changes in chromatin accessibility
activate gene transcription, we performed a Transcription Factor (TF) Motif Enrichment analysis on the
differentially accessible promoter regions of the upregulated DEGs (**Fig. 3e-f, Supplemental table 4**).
Among the top 15 most significant TF motifs enriched on the promoters of upregulated DEGs across
the cell clusters, TF motifs KLF15, ZNF148, and MAZ were found commonly enriched across
subregions CA1, CA3, and subiculum. Commonalities in the enrichment of TF motifs on upregulated
DEG promoters were observed across hippocampal subregions, such as EGR1 and CTCFL in CA1 and
CA3. No significant TF motifs were found enriched in the upregulated DEGs in the DG.

Cell-cell communication analysis reveals enhanced glutamate signaling following increases in Kcr

Activity-dependent synaptic transmission within the intrahippocampal circuit facilitates adaptive
behaviors for encoding spatial, episodic, and contextual memories[48-50]. Utilizing published databases of
ligand-receptor interactions[51], we constructed an intercellular communication network of the principal
neuronal subtypes to study the impact of Kcr enhancement on intrahippocampal signaling. Our analysis
revealed that crotonate administration primarily increases the strength of glutamatergic signaling
between the principal neuronal layers within the hippocampal circuit (**Fig. 4a**). Next, we investigated
the ligand-receptor pairs in each intrahippocampal connection that exhibit increase in their

communication probability following crotonate administration. We found that increasing Kcr levels

strengthened the communication between distinct glutamatergic ligand-receptor pairs within specific

neuronal networks of the hippocampal circuit. (**Fig. 4b**). These include the Slc1a2-Grik2/Grik4/Grik5,

and Slc1a2-Gria4 pairs in the DG-CA3, Slc1a2-Grik2/Grik5/Grm7, and Slc1a1-Grik2/Grik5/Grm7 pairs

in the CA3-CA1, and Slc1a1-Grik2/Grik5/Grm7, Slc1a2-Gria3/Gria4/Grik2/Grik5, and Slc17a7-

Grik2/Gria3/Grm7/Grm8 pairs in the CA1-subiculum neuronal connection (**Fig. 4b**). We computed the

communication probability of ligand-receptor interactions for glutamate signaling genes within principal

neuronal cell types, which further confirmed the enhanced strength of glutamatergic transmission after

crotonate administration (**Fig. 4c**). Collectively, our findings suggest that crotonate-dependent increase

in histone Kcr augments the strength of glutamatergic neurotransmission within the hippocampal circuit.

Real-time ex vivo fluorescent imaging reveals enhanced activity-dependent glutamate transients

following increases in Kcr

Our single-nuclei multiomics and cell–cell communication analyses indicated enhanced glutamatergic

signaling—a functional readout of which is presynaptic glutamate release. To test this, we transduced

WT mice with an AAV encoding the fluorescent glutamate sensor iGluSnFR (pAAV$_9$-hSyn-iGluSnFR) in

the dorsal hippocampus. Four weeks later, acute hippocampal slices were prepared and incubated in

either artificial CSF (aCSF) or aCSF containing 5 mM crotonate (**Fig. 4d**). We applied three stimulation

frequencies (10, 20, and 50 Hz) based on prior studies[52] and recorded stimulus-evoked glutamate

transients in CA1 following stimulation of CA3–CA1 Schaffer collateral afferents in the stratum radiatum

(**Fig. 4e–h**). Crotonate-treated slices exhibited a significant increase in peak glutamate amplitude

across all three stimulation frequencies (**Fig. 4g**), with a concomitant increase in area under the curve

(AUC) at 10 and 20 Hz stimulation (**Fig. 4h**). In contrast, glutamate clearance kinetics (τ_{fast} and τ_{slow};

depicted in a representative trace illustrated in **Supplementary Fig. 8**) were unchanged across

conditions (**Supplementary Fig. 9a-b**), suggesting that there is no difference in glutamate clearance.

Together, these results indicate that Kcr augments glutamatergic neurotransmission by facilitating

activity-dependent presynaptic glutamate release.

Real-time ex vivo fluorescent imaging reveals enhanced post-synaptic responsiveness to Kcr-

dependent increase in presynaptic glutamate release

Our iGluSnFR findings demonstrating increased presynaptic glutamate release complements our cell–

cell communication analyses, altogether indicating enhanced glutamatergic signaling strength with

elevated Kcr. This led us to hypothesize that Kcr-mediated enhancement of activity-dependent

presynaptic glutamate release at CA3–CA1 synapses would alter circuit responsiveness to increased

glutamatergic input. Such changes in circuit activity could be mediated, in part, by select postsynaptic

glutamate receptor genes that regulate activity-dependent calcium influx—the expression of which were

found upregulated following crotonate administration. As a proof of concept, we investigated the

expression profile of the AMPA receptor encoding gene *Gria4*[53], one of the significantly upregulated

glutamatergic signaling genes following crotonate treatment. Consistent with the single-nuclei

transcriptomic data (**Fig. 5a**), RNAscope analysis revealed a significant increase in *Gria4* expression in

hippocampal CA1 after crotonate administration (**Fig. 5b-c**).

We then employed an *ex vivo* fluorescence imaging approach using the genetically encoded calcium

indicator GCaMP6s to examine widefield CA1 circuit dynamics in response to Kcr-dependent increase

in activity-dependent presynaptic glutamate release from CA3 afferents (demonstrated earlier in **Fig.**

4d-h). Wild-type mice were transduced with an AAV encoding GCaMP6s (pAAV$_9$-hSyn-GCaMP6s-P2A-

mRuby3) in the dorsal hippocampus, and acute slices were prepared two weeks later. Slices were

incubated in either crotonate (5mM) or aCSF (**Fig. 5d**), and stimulus-evoked calcium transients were

recorded over time at three different frequencies (**Fig. 5e-f**). Crotonate-treated slices exhibited

significantly higher peak calcium transients across all three stimulation frequencies (**Fig. 5g**),

accompanied by increased AUC at 10 and 50 Hz (**Fig. 5h**). To confirm that these effects were driven by

transcriptional regulation rather than non-genomic actions of the drug, we performed a control

experiment in which slices were incubated with crotonate for only 10 minutes (**Fig. S10**). This brief

exposure failed to alter either peak amplitude or AUC (**Supplementary Fig. 10b-c**). Collectively,

these findings indicate that elevated histone Kcr enhances activity-dependent calcium signaling in

hippocampal neurons—an effect likely attributable to the transcriptional upregulation of postsynaptic

calcium-permeable glutamate receptor genes following Kcr elevation.

Discussion

The discovery of various non-acetyl histone acylations over the past decade has led to studies to

determine whether these modifications play a unique role in the epigenetic regulation of gene

expression. Histone Kac has long been regarded as a cornerstone of the epigenetic mechanisms

underlying long-term memory consolidation[2,4,9,54]. In this study, we demonstrate the critical role of lysine

crotonylation, a non-acetyl histone acylation associated with transcription activation[6,12,27], in regulating

hippocampal long-term memory consolidation. Through pharmacological and genetic approaches, we

demonstrate the bidirectional modulation of long-term memory by manipulating Kcr levels in the dorsal

hippocampus. Mechanistically, our findings reveal that memory enhancement observed upon

pharmacologically increasing Kcr levels is mediated through the function of ACSS2 in excitatory

neurons in the hippocampus. Single-nuclei transcriptomic and epigenomic analyses revealed that Kcr
primarily regulates the chromatin accessibility and expression of genes linked to synaptic function and
glutamatergic neurotransmission within principal neuronal populations of the dorsal hippocampus. We
then performed cell–cell communication analyses which predicted that increasing hippocampal Kcr
levels during the critical window of memory consolidation enhances the communication probability of
glutamatergic neurotransmission within intrahippocampal circuits. We performed *ex vivo* real-time
fluorescence-based imaging and found that Kcr indeed facilitates activity-dependent presynaptic
glutamate release and enhances neuronal excitability. Altogether, our study identified a novel
epigenetic mechanism critical for long-term memory storage and excitatory neurotransmission.

Histone acylation profiles are regulated not just by the enzymatic activities of their respective writers
and erasers, but also through changes in the nuclear concentrations of their metabolic precursors[6,12].
We found that bidirectional manipulation of hippocampal crotonyl-CoA levels had contrasting effects on
memory consolidation. Specifically, increasing crotonyl-CoA levels by administering crotonate
increased histone Kcr levels in the dorsal hippocampus and enhanced long-term memory. Conversely,
depletion of crotonyl-CoA levels by overexpressing CDYL reduced histone Kcr levels and led to deficits
in long-term memory. Together, these findings not just establish Kcr as a key regulator of long-term
memory but also identify CDYL as a potential memory suppressor gene[13]. Crotonyl-CoA levels are also
regulated by the metabolic enzyme ACSS2, which synthesizes crotonyl-CoA from crotonate[12,32]. In our
study, we found that crotonate fails to increase hippocampal histone Kcr levels or enhance long-term
memory in the ACSS2 cKD mice, indicating that ACSS2 is essential for maintaining the nuclear pool of
crotonyl-CoA. Collectively, these findings suggest that endogenous crotonate serves as a principal
source of nuclear crotonyl-CoA, and that ACSS2-mediated conversion of crotonate to crotonyl-CoA is a
critical molecular underpinning of Kcr-dependent enhancements in hippocampal memory.

Considering the promiscuity of histone acylation writers and erasers, the availability of metabolic
precursors becomes even more critical in defining the landscape of histone acylations. Emerging
evidence indicates that different acyl-CoA species compete for binding to KATs, such that changes in
the relative abundance of specific acyl-CoAs can alter the balance of histone acylation marks[6,12]. For
instance, depletion of cytoplasmic and nuclear acetyl-CoA pools in metazoans by knockdown of the
enzyme ATP citrate lyase (ACLY) led to an increase in p300-catalyzed histone Kcr[55]. Thus, increasing
the nuclear availability of less abundant acyl-CoA species enhances their access to shared
acyltransferases, which in turn facilitates the catalysis of histone acylation marks. Canonical Kac writers
such as p300, and readers such as the YEATS-domain proteins, catalyze and recognize Kcr more
efficiently[27,56], and Kcr itself serves as a potent activator of transcription[12]. Moreover, the thermal

instability of the carbon–carbon double bond in crotonyl-CoA facilitates the dissociation of the crotonyl

group from its CoA counterpart, thereby making it more convenient for the writers to efficiently deposit

crotonyl moieties onto histone lysine residues[24,57]. Together, these observations raise the possibility

that metabolically steering cells toward crotonyl-CoA utilization may represent an effective strategy to

regulate transcriptional events during memory consolidation. In line with this idea, our findings show

that pharmacologically increasing crotonyl-CoA levels to favor histone Kcr indeed offers a tractable and

physiologically efficient approach to enhance long-term memory—laying the conceptual foundation to

leverage Kcr as a "memory enhancer".

Our single-nuclei multiomics approach, aimed at elucidating the molecular underpinnings of Kcr-

mediated hippocampal memory enhancement, revealed unique signatures of differential gene

expression and corresponding chromatin accessibility profiles within the hippocampal principal neurons.

Surprisingly, most of the upregulated genes with increased chromatin accessibility following crotonate

administration are not the canonical 'Immediate Early Genes' (IEGs) that serve as defining markers of

engram ensembles activated by spatial learning[30,58-60]. Chromatin enrichment studies have shown that

an increase in histone Kcr correlates with a reduction in the transcriptionally repressive mark

(H3K27me3) and an increase in the transcription activation mark (H3K4me3) on gene promoters, with

no observed change in the occupancy of acetylated histones (H3K27ac)[23]. Such an interplay between

histone Kcr and other post-translational modifications, such as Kac and methylation, likely represents

the framework of a unique "histone code" [61-63]—the mechanistic framework of which remains to be

elucidated in regard to long-term memory consolidation.

An important advantage of our multiomics framework is that it enabled us to interrogate the chromatin

accessibility landscape and identify cell-type-specific signatures of TF motif enrichment that underlie

Kcr-dependent gene regulatory programs. Notably, binding motifs of several members of the Krüppel-

like family of transcription factors, such as KLF15 and ZNF148, were found enriched in hippocampal

subregions CA1, CA3, and subiculum following Kcr enhancement. These TFs—integral in regulating

diverse developmental and homeostatic functions—are increasingly being recognized for their role in

long-term memory and plasticity. For instance, *Klf15*-null mice show marked deficits in hippocampal-

dependent memory, including reduced contextual freezing after aversive learning[64]. Likewise, human

genetic studies have shown that heterozygous variant in ZNF148 gene causes intellectual disability,

whereas *de novo* mutations in ZNF148 have been associated with autism spectrum disorder, and

cognitive impairment[65]. Beyond the Krüppel-like family, we also observed enrichment of MAZ binding

across the gene bodies of Kcr-responsive hippocampal DEGs. MAZ is known to recognize a 27-bp GC-

rich region within the *GluN1* promoter, and binding of MAZ to *GluN1* is essential for its transcriptional

activation[66]. Another notable finding was the enrichment of EGR1 transcription factor motif in CA1 and

CA3. *Egr1* (also known as *Zif268*, *Krox-24*, and *NGFI-A*) is a critical regulator of long-term memory and

synaptic plasticity[67-70]. Beyond its canonical transcriptional regulatory function, recent studies have

highlighted a link between Egr1 and epigenetic plasticity. Egr1 interacts with the DNA demethylase

TET1 during postnatal brain development to program the brain methylome, with Egr1 binding sites

becoming hypomethylated in mature neurons to regulate downstream gene expression[71]. Additionally,

Egr1 knockdown increases histone acetylation at enhancer regions, regulating inflammatory gene

expression in human macrophages[72]. Our findings extend the role of Egr1 in transcriptional

regulation—indicating that it may play a previously unappreciated role in coordinating Kcr-mediated

chromatin remodeling and gene expression. Collectively, these motif enrichments point to a regulatory

architecture in which Kcr-mediated chromatin remodeling facilitates selective TF engagement with the

genome, thereby regulating the transcriptional states that facilitate memory enhancement.

Another major strength of our study derives from our findings that increases in Kcr augments the

dynamics of intrahippocampal glutamatergic neurotransmission. Our single-nuclei multiomics findings

revealed that crotonate administration alters chromatin accessibility to upregulate genes involved in

glutamatergic neurotransmission within excitatory neuronal populations. Cell-cell communication

analysis—a powerful computational tool to infer and predict intercellular communication strength

between neuronal populations within defined circuits[73], further revealed that Kcr-enhancement

augments glutamatergic neurotransmission within hippocampal circuit. Specifically, the analysis

identified strengthened glutamatergic ligand–receptor interactions following Kcr enhancement,

exemplified by an increased predicted communication probability between presynaptic vesicular

glutamate transporters (VGLUTs) and postsynaptic kainite (KARs) and AMPA receptors (AMPARs).

Together, these findings are suggestive of improved dynamics and fidelity of intrahippocampal

glutamatergic signaling. These predictions were corroborated by *ex vivo* real time imaging, which

demonstrated elevated activity-dependent presynaptic glutamate release in the CA3-CA1 Schaffer

collaterals. This led to enhanced activity-dependent CA1 responsiveness at the circuit-level, as

evidenced by amplified calcium transients *ex vivo* in response to increased presynaptic glutamate

release at CA3–CA1 synapses. Importantly, studies examining histone PTMs on genes involved in

glutamatergic neurotransmission are limited[74,75], and, to date, no work has directly elucidated how

epigenetic marks regulating glutamatergic gene expression bring about functional changes in synaptic

neurotransmission and neuronal excitability. Our study reveals a mechanistic blueprint of how

epigenetic regulation of glutamatergic neurotransmission shapes synaptic function and drives

hippocampal memory consolidation.

In summary, our work demonstrates the critical role of a non-acetyl histone acylation that has
previously not been studied in the context of long-term memory. We establish histone Kcr as a
molecular switch for long-term memory, redefining our conceptual understanding of epigenetic
mechanisms underlying long-term memory consolidation. We identify transcriptomic signatures of
learning in distinct hippocampal subregions regulated by Kcr and highlight the epigenetic control of
glutamatergic neurotransmission as a critical mechanism underlying Kcr-dependent long-term memory
enhancement. By demonstrating that histone Kcr lies at the molecular crossroads of learning-induced
transcriptional programs, synaptic neurotransmission, and behavior, our study provides a mechanistic
framework for how epigenetic mechanisms translates into functional outcomes—shedding light on the
elusive histone code of memory consolidation. Given that recent studies have demonstrated the role of
Kcr-mediated regulatory mechanisms in several neurological disorders[39,76], the broader relevance of
our work is that it lays the groundwork to develop more effective therapeutic interventions to treat
cognitive impairments associated with neurological, neurodevelopmental, and neuropsychiatric
disorders.

Methods

Data reporting: No statistical methods were applied to predetermine sample size.

Mouse lines: Adult male C57BL/6J mice were purchased from Jackson Laboratories. The ACSS2[f/f]
mice were generated at the University of Iowa Genome Editing Facility[47]. This mouse model was
created by inserting loxP sites flanking the Exon2 of ACSS2 using CRISPR/Cas9. The mice were 2-4
months of age at the time all the behavioral and biochemical experiments were performed. All mice had
ad libitum access to food and water, and lights were maintained on a 12-hr light/dark cycle. All
experiments were conducted according to US National Institutes of Health guidelines for animal care
and use and approved by the Institutional Animal Care and Use Committee of the University of Iowa,
Iowa.

Drug preparation: Sodium crotonate was freshly prepared on the day of the experiment. 2.5 gm of
Crotonic acid (Sigma) was dissolved in 30 ml of 0.9% saline. Thereafter, the pH of the solution was
adjusted to 7.4 using 5(N) NaOH, and additional 0.9% saline was added to adjust the final volume to 50
ml. The solution was then filtered through a 0.2 μm syringe filter. Appropriate volume of the drug
(calculated based on the body weight of the animals) was administered via oral gavage based on the
dosing regimen.

Histone extraction: Histone extraction was performed using a commercially available kit (Histone
Extraction Kit, Active Motif, #40028) according to the manufacturer's protocol. Briefly, flash frozen
hippocampi were mechanically homogenized in 300 µl of ice-cold Lysis Buffer AM8 using Dounce
homogenizers and incubated on ice for 30 mins. The homogenate was centrifuged at 2,644 x g for 2
minutes at 4°C, and the nuclear pellet was resuspended in 250 µl ice-cold Extraction buffer. Following
resuspension, the nuclear suspension was incubated on an end-to-end rotator overnight at 4°C. The
pellet insoluble material was centrifuged at 20,800 x g for 10 minutes at 4°C the next day, the
supernatant was collected, and the requisite volume of Complete Neutralization Buffer was added.
Samples were stored in -80°C prior to western blotting.

Western blot analysis: Western blotting was performed as previously described[30,58]. Whole cell
lysates were run on a 4–20% Tris-HCl Protein Gel (BIO-RAD, #3450033). Proteins from the gel were
then transferred to methanol-activated polyvinylidene difluoride membranes using the Trans-Blot Turbo
Transfer System (BIO-RAD). Membranes were blocked with the Odyssey® Blocking Buffer (LI-COR)
diluted in TBS and incubated overnight at 4°C with the following primary antibodies: pan-Kcr (1:1000,
PTM BIO, #PTM-501), Total H3 (1:5000, ACTIVE MOTIF, #61647), ACSS2 (1:1000, Cell Signaling,
#3658), and Actin (1:10,000, #MA1-91399). Post primary antibody incubation, membranes were
washed thrice in TBS for 5 mins and incubated with IRDye® 800CW Goat anti-Rabbit IgG Secondary
Antibody (1:5,000, LI-COR, #926-32211) and IRDye® 680LT Goat anti-Mouse IgG Secondary Antibody
(1:5000, LI-COR, #926-68020). Membranes were then washed thrice in TBS, 5 mins each. Images
were acquired using the Odyssey Infrared Imaging System (LI-COR). Quantification of western blot
bands was performed using Image Studio Lite ver5.2 (LI-COR).

Immunofluorescence: IHC experiments were performed as per earlier studies[30,58]. Animals were
perfused with 4% PFA. Whole brains were harvested, immersed in 4% PFA, and stored at 4°C. 24 hrs
after, brains were immersed in 30% sucrose and stored at 4°C. 20 µm coronal brain sections were
made in a cryostat (Leica). Free-floating sections were washed thrice with PBS, blocked in a blocking
buffer (0.1% PBST and 5% BSA) and incubated with the following primary antibodies for 48 hr: pan-Kcr
(1:2000, PTM BIO, #PTM-501), CDYL (1:500, Sigma, #HPA035578), V5 (1:500, ThermoFisher
Scientific, #37-7500), and GFP (1:4000, Abcam, #ab290). Post primary antibody incubation, sections
were washed in PBS thrice, followed by a 2 hr secondary antibody incubation with the following
antibodies: Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 546
(1:500, ThermoFisher Scientific, #A-11003), Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary
Antibody, Alexa Fluor™ 647 (1:500, ThermoFisher Scientific, #A-21244). Sections were then washed
thrice in PBS and mounted on Superfrost™ Plus microscope slides (Fisherbrand). This was followed by

coverslip mounting with ProLong™ Diamond Antifade Mountant with DAPI (ThermoFisher Scientific,
#P36962).

Confocal imaging and image analysis: Confocal images of IHC experiments were obtained in an
Olympus FV 3000 confocal microscope using a 20X NA = 0.4 achromat dry objective at 800 × 800-pixel
resolution and 1X optical zoom. All images (16 bit) were acquired with identical settings for laser power,
detector gain and pinhole diameter for each experiment and between experiments. Images from the
different channels were stacked and projected at maximum intensity using ImageJ (NIH). Mean
fluorescence intensity (MFI) was computed using ImageJ plugins. MFIs were then subtracted to
background intensities in each slice. Background subtracted MFIs in the experimental group were then
normalized to the average background subtracted MFIs in control. For MFI quantification, the entire
area corresponding to CA1, CA3, DG or subiculum were considered as region of interest.

Chromatin Immunoprecipitation (ChIP): Frozen hippocampal tissue samples (~20 mg) was
crosslinked in 1% formaldehyde for 10 min at room temperature with gentle rocking, followed by
quenching with 0.125 M glycine for 5 min. Samples were centrifuged at 4000 rpm for 5 min at 4°C and
washed three times in PBS supplemented with protease inhibitors and sodium butyrate to preserve
histone acylations. Crosslinked tissue was homogenized in Covaris ChIP lysis buffer using a type B
glass dounce (20 strokes per sample), followed by sequential washes and resuspension in SDS
shearing buffer. Chromatin was sonicated for 10 cycles using a Diagenode Bioruptor Pico to generate
DNA fragments of 200–500 bp. Following centrifugation at 10,000 × g for 5 min to remove insoluble
debris, supernatants were collected, and 50% Protein G slurry was added for preclearing at 4°C for 30
min. For immunoprecipitation, chromatin was incubated overnight at 4°C with anti-pan Kcr antibody
(PTM BIO, #PTM-501) or normal rabbit IgG (negative control; Diagenode, #C15410206) in ChIP
dilution buffer. The next day, antibody–chromatin complexes were captured with blocked Dynabeads
for 2 h at 4°C and subjected to sequential washes with low-salt, high-salt, LiCl, and TE buffers.
Chromatin was eluted twice with 100 μL of ChIP elution buffer at 65°C and pooled. Crosslinks were
reversed by 5 M NaCl and incubating at 65°C overnight, followed by RNAse A and proteinase K
digestions. DNA was purified using the QIAquick PCR purification kit (Qiagen) and eluted in nuclease-
free water.

ChIP-Seq library preparation, sequencing, and analysis: ChIP-Seq libraries were prepared using
KAPA library preparation kit. Indexed libraries were pooled and sequenced in illumina NovaSeq 6000.
Paired-end sequencing was performed for all input and ChIP samples, generating approximately 100
million reads per sample with a read length of 100 nucleotides. ChIP-Seq reads were trimmed from

both 5' and 3' ends using Trimmomatic paired end mode until the final base had a quality score > 30,

discarding reads with < 23 bp. ChIP-Seq trimmed reads were aligned to mm10 reference genome using

bwa aligner. Differential histone crotonylation between samples was assessed using the 'MAnorm'

algorithm[77] which quantitatively compares ChIP-seq signal intensities between two samples by

normalizing against the set of peaks shared in both datasets. Broad peaks were first called with

'MACS2' using lenient thresholds to capture histone mark enrichment. These peak sets were then

provided to MAnorm as input. The program merges peaks across both samples to define a unified

reference, identifies overlapping peaks, and computes read densities for each region. MAnorm then

models the relationship between samples by performing a robust linear regression on the $\log_2$ fold-

change versus mean intensity of common peaks, thereby normalizing systematic differences in

sequencing depth or background enrichment. For each peak, MAnorm reports a normalized M-value

($\log_2$ fold-change) and an associated p-value, which imply the degree and significance of differential

enrichment. To focus on regulatory elements, significant differential peaks were filtered to retain only

those located within ±2,000 bp of gene promoters (defined relative to the annotated TSS).

Comparative analysis of learning-induced gene targets regulated by H3K27ac and Kcr: We

compared gene targets of Kcr in dorsal hippocampus with gene promoters enriched with H3K27ac in

the cortex. Raw sequencing dataset focusing on H3K27ac modifications in cortical neurons

(GSM1629381, GSM1629397)[42] were downloaded from the Sequence Read Archive using fastq-dump

from sratoolkit. We then ran the script 'ChIP-seq_SE_pipeline.sh' for each raw dataset and normalized

results by their respective ChIP-seq inputs using the script 'subtractTwoWig.py'. Peak calling was

performed using 'MACS2', and identified peaks were overlapped with gene annotations within a region

spanning 2,000 base pairs upstream to 2,000 base pairs downstream of the transcription start site

(TSS). We analyzed the resulting gene lists by overlapping them with learning-induced genes from our

previously published RNA-seq study [28] to associate fold change values with different gene sets

harboring distinct epigenetic marks.

In situ hybridization (RNAscope): *In situ* hybridization was performed using the RNAscope™

Multiplex Fluorescent Reagent Kit v2 (Advanced Cell Diagnostics) according to the manufacturer's

protocol as previously described[30]. Briefly, 20 µm cryosections obtained from fixed frozen brains were

mounted on Superfrost™ Plus microscope slides (Fisherbrand). Slides then underwent serial

dehydration in Ethanol, followed by Hydrogen Peroxide treatment, Target Retrieval, and Protease III

treatment. Hybridization of probes was done at 40°C for 2 hr in an HybEZ oven using a probe

against *Gria4.* The probe signal was amplified with Pre-amplifiers (AMP 1-FL, AMP 2-FL, and AMP 3-

FL) and counterstained with OPAL 570 reagent (#FP1488001KT, Akoya Biosciences). Finally,

coverslip mounting was done on the slides using ProLong™ Diamond Antifade Mountant with DAPI

(ThermoFisher Scientific, P36962). The slides were stored in 4^0C until they were imaged.

Adeno-associated virus (AAV) constructs and stereotactic surgeries: AAV$_9$-CaMKIIα-EGFP and

AAV$_9$-CaMKIIα-CDYL-V5 were purchased from VectorBuilder (VectorBuilder Inc). AAV$_9$-CaMKIIα-GFP-

Cre (#105551), pAAV$_9$-hSyn-iGluSnFr (#98929), and pAAV$_9$-hSyn1-GCaMP6s-P2A-mRuby3 (#112005)

were purchased from Addgene. Mice were anesthetized using 5% isoflurane. A steady flow of 2.5%

isoflurane was maintained throughout the remainder of the stereotactic surgery. 1 µl of respective AAVs

were bilaterally injected into the dorsal hippocampus (coordinates: anteroposterior, −1.9 mm,

mediolateral, ±1.5 mm, and 1.5 mm below bregma). Following viral infusion, drill holes were closed with

bone wax (Lukens) and the incisions were sutured. Intraperitoneal (IP) injections of Meloxicam (5

mg/kg) were administered for 2 successive days after the surgery to manage the post-operative pain.

Spatial object recognition (SOR) task: All the behavioral experiments were performed based on

previously published studies[30,58] during the light cycle in between Zeitgeber time (ZT) 0 through 2. Mice

were individually housed for 7 days before the training. Animals were handled for 2 mins each day for 5

successive days before training. In the strong training paradigm, animals were habituated in an open

field for 6 minutes. This was followed by three 6-minute sessions inside the same arena containing

three different glass objects. These objects were placed at specific spatial coordinates with respective

to a spatial cue within the arena. An inter-trial interval of 5 minutes was set in-between the three

training sessions. In the sub-threshold training paradigm, mice were subjected to three 3-minute

training trials in the open field with two objects and an inter-trial interval of 5 minutes in between

sessions. 24 hr after training, the animals were returned to the arena with one of the objects displaced

to a novel spatial coordinate. Exploration time around all the objects were then manually scored.

Percent preference towards the displaced object was calculated using the following equation:

Percent preference for displaced object $= \dfrac{(exploration\ towards\ the\ displaced\ object)}{(total\ exploration\ towards\ all\ objects)}\ x\ 100$

For the histone Kcr profiling and the single-nuclei experiments, animals were euthanized by cervical

dislocation 1 hr after the last training trial in the SOR task. Hippocampal tissue was flash frozen and

stored at -80^0C. Home caged animals were euthanized within the same ZT window to eliminate the

possible confounding effects of circadian rhythmicity. To examine the learning-induced expression of

Kcr in hippocampal subregions, mice were perfused 1 hr after the training session, and whole brains

were harvested for IHC.

Contextual fear conditioning: Contextual fear conditioning was performed using a sub-threshold
learning paradigm. In brief, mice were handled daily for five days before conditioning. The conditioning
protocol involved a single 2-second, 0.75 mA footshock delivered 2.5 minutes after the mice were
introduced into the chamber. Mice remained in the chamber for an additional 30 seconds before being
returned to their home cage. 24-hrs later, they were reintroduced to the same chamber for 5 mins.
Freezing behavior was assessed using FreezeScan software (CleverSys Inc.).

Nuclei isolation: Nuclei isolation form frozen hippocampal tissue was performed according to the
manufacturer's protocol (Chromium Nuclei Isolation Kit with RNAse Inhibitor, 10x Genomics,
#1000494). Briefly, frozen tissue was homogenized in 500 µl of Lysis Buffer using Dounce
homogenizers. The lysate was transferred to a Nuclei Isolation Column and centrifuged at 16,000 x g
for 30 seconds at 4^{o}C. The pellet was then resuspended in 500 µl of Debris Removal Buffer and
centrifuged at 700 x g for 10 minutes at 4^{o}C. The nuclear pellet was then resuspended with 1 ml of
Wash Buffer and centrifuged at 500 x g for 5 minutes at 4^{o}C. Finally, the nuclear pellet was
resuspended in 50 µl of Resuspension Buffer. Nuclei count was manually done using a
Hemocytometer.

Single-nuclei multiomic data processing and analysis: Raw sequencing data were processed using
the 'Cell Ranger ARC' pipeline (v2.0.2) with the cell ranger-arc mm10 reference. Default parameters
were used to align reads, count unique fragments or transcripts, and filter high-quality nuclei. HDF5
files for each sample (Saline1, Saline2, Crotonate1, Crotonate2) containing barcoded RNA counts and
ATAC fragments per cell cluster were loaded into Seurat (Read10X_h5). This resulted in the generation
of four Seurat objects, each containing both RNA and ATAC assays. Nuclei with outliers within the
ATAC and RNA QC metrics (<200 and >100,000 ATAC read counts, <200 and >50,000 RNA read
counts, nucleosomal signal $>$ 4, TSS enrichment $<$ 3, %reads in peaks < 15 and percentage of
mitochondrial reads > 5) were removed.

To analyze the RNA component of the multiomics data, gene counts were normalized, and log
transformed (LogNormalize). The top 2,000 most variable features that distinguish each cell were
identified using 'FindVariableFeatures' (selection.method $=$ 'vst'). Features that are repeatedly
variable across cells and datasets were selected for integration ('SelectIntegrationFeatures'). We then
identified anchors ('FindIntegrationAnchors''), which took the list of 4 individual Seurat objects for each
sample as input. These anchors were used to integrate the four datasets together ('IntegrateData').
Linear dimensionality reduction was performed on the integrated Seurat object by principal component
analysis (runPCA, npcs $=$ 30). A k-nearest-neighbours graph was constructed based on Euclidean

distance in PCA space and refined ('FindNeighbors'), following which the nuclei were clustered using

the Louvain algorithm (FindClusters, resolution = 0.5). Clusters were visualized with UMAP

(runUMAP, dims = 30). Both RNA and ATAC assays were used to identify cell-type specific

signatures of biomarkers. Differentially expressed genes (DEGs) in individual clusters between saline

and crotonate treated groups were calculated (FindMarkers, test.use = 'wilcox,' Padj < 0.05,

absolute logFC.threshold of 0.2).

To analyze the ATAC component of the multiomics data, the default assay was switched to ATAC prior

to integrating the four Seurat objects, and peak calling was performed. The set of peaks identified by

'Cellranger' often merges distinct peaks that are close together - confounding the motif enrichment

analysis and peak-to-gene linkage. We were able to circumvent this concern and identify a more

accurate set of peaks by using the 'MACS2' (CallPeaks) peak calling feature on all cells together.

Peaks on nonstandard chromosomes and in genomic blacklist regions were removed

('keepStandardChromosomes' and 'subsetByOverlaps'). A frequency-inverse document frequency

normalization was performed across cells and peaks ('RunTFIDF'). Thereafter, a feature selection was

performed using all the peaks as input (FindTopFeatures, min.cutoff = 5). The selected peaks went

through a dimensional reduction on the TF-IDF normalized matrix using a singular value decomposition

('RunSVD'). To identify the differentially accessible regions (DAR) between crotonate versus saline

group, the Seurat function 'FindMarkers' was used using logistic regression (LR) as a method to test

significance. The DARs with adjusted p value < 0.05 and absolute log2foldchange threshold of above

0.2 were considered as significantly differentially accessible. Further, the DARs were annotated using

'Closestfeature' function from Signac package as well as 'annotatePeak' function from 'ChIPseeker'

package. The DEGs were furthered correlated with DARs where the DARs were selected only from the

promoter (+/- 2kb from transcription start site) and genebody regions, filtering out the peaks from distal

genomic regions and downstream of 3'UTR[78]. The genes found concordant in both DEG and DAR lists

were shortlisted for downstream gene ontology enrichment analysis. UpSet plots were generated using

UpSetR package.

Gene Ontology enrichment analysis: The concordant DARs and DEGs were analyzed for Molecular

Function (MF) enrichment analysis by using the 'clusterProfiler' package with the default criteria

(pvalueCutoff = 0.01 and qvalueCutoff = 0.05). Here, the significant upregulated DEGs (adj p value <

0.05 and log2FC > 0.2) concordant with the significant more accessible DARs (adj p value < 0.05 and

log2FC > 0.2) were used to generate the upregulated gene list, whereas the significant downregulated

DEGs (adj p value < 0.05 and log2FC < -0.2) concordant with the significant less accessible DARs (adj

p value < 0.05 and log2FC < -0.2) were used to generate the downregulated gene list. The DARs were

selected only from the promoter (+/- 2kb from transcription start site) and gene body regions, filtering

out the peaks from distal genomic regions and regions downstream of 3'UTR. All further data

visualizations were made using clusterProfiler package.

TF motif enrichment analysis: To identify cell-type specific regulatory sequences, we performed

transcription factor motif enrichment analysis on the DEGs that were found to have concordant

differentially accessible peaks in the promoter region. Here, we restricted TF motif enrichment only in

the promoter region (a window of 2000bp upstream and downstream of transcription start site). Motif

enrichment was performed using 'FindMotifs' function of signac package. The motifs that had adjusted

p value < 0.05 were considered significant. The top 15 significant motifs from the clusters were plotted

as heatmap using ComplexHeatmap package. Weight matrices for the top motifs were also plotted to

visualize the motif sequences.

Cell-cell communication analysis: The cell-cell communication analysis on the snRNA-seq data was

performed using CellChat (v2.1.2). The Saline and Crotonate RNA normalized counts were taken and

individual cellchat objects were created. Ligand-receptor (LR) interactions in each group were

calculated by identifying overexpressed ligands or receptors with a log fold change cutoff 0.1

(identifyOverExpressedGenes(thresh.fc = 0.1, thresh.p = 0.05)) followed by identifying interactions if LR

pairs are overexpressed (identifyOverExpressedInteractions). To assign each interaction with a

probability score, computeCommunProb function was used with a default statistical method called

'trimean'. Cell-cell communication was filtered to have minimum cell number 10 in each cell group

(filterCommunication(min.cells = 10)). After that communication probability on signaling pathway level

was calculated by summarizing the communication probabilities of all LR interactions associated with

each signaling pathway. Finally, the cell-cell communication network was aggregated by counting the

number of links or summarizing the communication probability amongst the cell groups. Thereafter, the

cellchat objects were merged and compared using netVisual_aggregate,

netAnalysis_signalingChanges_scatter, and netVisual_chord_gene functions.

Acute brain slice preparation: Acute hippocampal slices were prepared as described in published

studies[52,79]. 2-4-month-old male mice were deeply anaesthetized with isoflurane, following which their

brains were harvested for slice preparation. Coronal sections were obtained in ice-cold dissection buffer

(in mM: 220 sucrose, 3.3 KCl, 5 $MgSO_4$, 1.25 NaH_2PO4, 25 $NaHCO_3$, 11 D-glucose, and 0.2 $CaCl_2$)

and sections containing dorsal hippocampal representations were allowed to recover in a chamber

containing normal artificial CSF (aCSF) (in mM; 116 NaCl, 3.3 KCl, 1.3 $MgSO_4$, 1.25 NaH_2PO4, 25

$NaHCO_3$, 11 D-glucose, 1.3 $CaCl_2$) saturated with carbogen (95%O_2 / 5%CO_2) at 32°C. The slices were

incubated for at least 2 hrs either in normal aCSF or 5 mM Crotonate before being transferred to a

submerged chamber for imaging, constantly supplied with aCSF (flow rate: 2.5 ml/min) also maintained

at 32°C. For the experiment described in **Fig. S9**, the slices were incubated for 5-10 mins with the drug,

following which they were transferred to the submerged chamber containing aCSF for imaging.

Mesoscopic imaging and image analysis: Acute coronal sections containing hippocampi transduced

with either iGluSnFr or GCaMP6s were placed in a submerged chamber attached to the stage of an

upright microscope (Olympus MVX10), continuously perfused with aCSF saturated with carbogen and

maintained at 32-34°C. A monopolar stainless steel stimulation electrode (AM systems) was placed on

in the Stratum Radiatum to stimulate the Schaffer Collateral pathway. Optical excitation was delivered

by an LED light (X-Cite XYLIS XT720S). Fluorescent mesoscopic imaging of the entire brain slice was

performed using a 2x objective (NA 0.5). The LED light was band-passed through a 470/40 nm filter

and a dichroic 495 nm filter, while the emitted light was isolated using a 525/50 nm filter. For stimulus

trains, 10 stimuli (3mV, 500 µs) were delivered with variable inter-stimulus intervals (for 10 Hz: 100ms;

for 20Hz: 50ms; for 50Hz: 20ms), as done in previous studies[52]. Images were acquired with a CMOS

camera (Basler Ace acA4112-30µm) at a frame rate of 10 Hz (GCaMP6s imaging) or 90-100 Hz

(iGluSnFr imaging) with 4:1 binning. The camera acquisition and LED excitation were digitally

synchronized using Micro-manager (ImageJ). Raw images in time series were background-subtracted

and smoothed (median filter). Time series were baseline subtracted, scaled to baseline means ($\Delta F/F_0$,

baseline: ~2s before the stimulation / response onset), and then compressed into Max Intensity

Projections (MIPs). For Glutamate transients (iGluSnFr), responsive regions (ROIs) were generated by

converting the $\Delta F/F_0$ MIPs into binary (Threshold: mean + SD). For calcium transients, hippocampal

area that showed ectopic expression of GCaMP6s was selected as the ROI. Fluorescence traces

generated using the respective ROIs were analyzed in MATLAB (MathWorks). Ca^{2+}/ Glutamate

response parameters (peak amplitude, locations, width, prominence) were determined using the

'FindPeaks' function in MATLAB (minimum peak prominence: 0.025 for Glutamate, 0.05 for Ca^{2+}). Area

under the curve (AUC) was then calculated using trapezoidal numerical integration ('Trapz' function) in

MATLAB. For glutamate transients, the slow and fast components of the decay kinetics (Tau) were

determined by fitting two single-term exponential fits to the decay of the response (τ_{fast}: stimulus offset

to ~4s, τ_{slow}: ~4s to 10s).

Statistics: Behavioral and biochemical data were analyzed using unpaired two-tailed t-tests, one-way

ANOVA, or two-way ANOVAs (sometimes with repeated measures as the within-subject variable).

Sidak's multiple comparison tests or Dunnett's multiple comparison tests were used for post-hoc

analyses wherever required. In the iGluSnFR and GCaMP6s imaging data, normality of the distributions

was assessed using the D'Agostino–Pearson K2, Anderson-Darling tests, and QQ plots. If data was

log-normal, a Lognormal Welch's t test was used. Outliers were detected using Grub's or IQR tests (for

parametric and nonparametric data), although no outliers were detected or excluded. In all the

experiments, differences were considered statistically significant when $p<0.05$. As indicated for each

figure panel, all data were plotted in box plots.

Ethics: All procedures on mice in this study were conducted according to US National Institutes of

Health guidelines for animal care and use and were approved by the Institutional Animal Care and Use

Committee of the University of Iowa, Iowa.

Reporting summary: Further information on research design is available in the Nature Portfolio

Reporting Summary linked to this article.

Figure legends

Figure 1. Lysine crotonylation is a critical regulator of learning-induced gene expression and

long-term spatial memory consolidation. **a.** Depiction of molecular regulators of histone lysine

crotonylation (Kcr). **b.** Schematic of the experiments performed in **c-i**. **c.** Immunofluorescence using

anti-Kcr antibody showing levels of Kcr in different hippocampal sub-regions of HC and learning

(SOR+1hr) mice. **d.** Normalized Mean Fluorescent Intensity (MFI) of nuclear Kcr levels across the

groups. For CA1: unpaired t test: t (6)$\Box$= 3.810, **$p\Box=\Box$0.0089. Homecage (n$\Box=\Box$4) and SOR+1 hr

(n$\Box=\Box$4), males only. For CA3: unpaired t test: t (6) = 0.2749, p$\Box=\Box$0.7926. Homecage (n$\Box=\Box$4) and

SOR+1 hr (n$\Box=\Box$4), males only. For DG upper blade (U): unpaired t test: t (6)$\Box$= 3.118, *p$\Box=\Box$0.0206.

Homecage (n$\Box=\Box$4) and SOR+1 hr (n$\Box=\Box$4), males only. For DG lower blade (L): unpaired t test: t

(6)$\Box$= 0.5452, p$\Box=\Box$0.605. Homecage (n$\Box=\Box$4) and SOR+1 hr (n$\Box=\Box$4), males only. **e.** ChIP-seq

density heat maps in dorsal hippocampus harvested from homecage (left) vs 1 hour after SOR-trained

mice (right) for histone Kcr at ± 2 kb from the TSS. **f.** Box plots showing the extent of gene induction in

the hippocampus following learning[28] for genes that exhibit H3K27ac binding in the promoter region

compared to genes that exhibit binding of H3K27ac and Kcr in the promoter region. Unpaired two-tailed

t test was used to determine a significant change in mean log2FC between the groups. **g.** Schematic of

viral constructs transduced in the dorsal hippocampus of adult male mice to overexpress

Chromodomain Y-like protein (CDYL). AAV$_9$-CaMKIIα-EGFP served as vector control, while AAV$_9$-

CaMKIIα-CDYL-V5 was used to drive expression of CDYL in excitatory neurons. **h.**

Immunofluorescence using V5 antibody showing ectopic expression of CDYL-V5 in dorsal

hippocampus two weeks after viral infusion. **i.** Immunofluorescence image of the dorsal hippocampus

using pan-Kcr antibody in CDYL-V5 or control expressing mice. **j.** Normalized Mean Fluorescent

Intensity (MFI) of nuclear Kcr levels across the groups. For CA1: Unpaired t test: t (5) = 2.679,

*p = 0.439. EGFP (n = 3) and CDYL (n = 4), males only. For CA3: Unpaired t test: t (5) = 3.417,

*p = 0.0189. EGFP (n = 3) and CDYL (n = 4), males only. For DG: Unpaired t test: t (5) = 1.863,

p = 0.1215. EGFP (n = 3) and CDYL (n = 4), males only. **k**. Long-term memory (24 hr)

assessment of mice infused with AAV-EGFP or AAV-CDYL into dorsal hippocampus. **l.** Quantification

of **k**. Two-way ANOVA: significant session x virus interaction $F_{(1,13)}=6.323$, p=0.0259, Sidak's multiple

comparison tests, **p=0.0041 (EGFP 24 hr Test vs CDYL 24 hr Test) and **p=0.0051 (EGFP Training

vs EGFP 24 hr Test); AAV-EGFP (n=7), AAV-CDYL (n=8), males only. All box plots: the center line

represents the median, the box edges represent the top and bottom quartiles (25th to 75th percentiles),

and the minimum and maximum whiskers.

Figure 2. *Single-nuclei multiomics (RNA+ATAC-seq) reveals mechanistic insights underlying*

lysine crotonylation-mediated memory enhancement. **a.** Kcr Western blots of core histones

extracted from dorsal hippocampus 1 hr after crotonate treatment (50 mg/kg or 200 mg/kg dose).

Vehicle (saline) treated mice were used as controls. **b.** Quantification of **a.** One-way ANOVA: Kcr: $F_{(2, 14)}$

= 5.550, p=0.0168. Dunnett's multiple comparisons tests: *p= 0.0108 (saline versus crotonate

200mg/kg). Saline (n=6), Crotonate 50 mg/kg (n=6), Crotonate 200 mg/kg (n=5), males only. **c.** Long-

term memory enhancement in a sub-threshold SOR learning paradigm following crotonate treatment.

Two-way ANOVA: significant session x treatment interaction $F_{(2, 23)}$ = 5.081, p= 0.0149, Sidak's post

hoc tests, *p=0.0359 (200mg/kg crotonate 24 hr test vs saline 24 hr test), **p=0.0021 (200mg/kg

crotonate train vs 200mg/kg crotonate 24 hr test), **p=0.0025 (200mg/kg crotonate 24 hr test vs

50mg/kg crotonate 24 hr test). Saline (n=9), Crotonate 50 mg/kg (n=7), Crotonate 200 mg/kg (n=10),

males only. **d.** Long-term memory in a sub-threshold CFC learning paradigm following crotonate

treatment. Two-way ANOVA: Significant session x treatment interaction $F_{(2, 19)}$ = 4.946, *p=0.0187,

main effect of session (pre-shock and 24hr test): $F_{(1, 19)}$ = 140.9, p<0.0001. Sidak's post hoc tests,

*p=0.0130 (200mg/kg crotonate 24 hr test vs 50mg/kg crotonate 24 hr test), **p=0.0051 (200mg/kg

crotonate 24 hr test vs saline 24hr test). ****p<0.0001 (Saline pre-shock vs saline 24hr test),

****p<0.0001 (crotonate 50mg/kg pre-shock vs crotonate 50mg/kg 24hr test), and ****p<0.0001

(crotonate 200mg/kg pre-shock vs crotonate 200mg/kg 24hr test). Saline (n=8), Crotonate 50 mg/kg

(n=6), Crotonate 200 mg/kg (n=8), males only. **e.** Experimental scheme: Adult male C57BL/6J mice

were trained in a sub-threshold SOR paradigm and administered with crotonate (200 mg/kg, oral

gavage; n=4) or saline (oral gavage; n=4) immediately after the completion of training. 1-hr later, the

dorsal hippocampus was harvested, and nuclei were isolated for single-nuclei multiomics processing.

Hippocampi from two animals within the same group were pooled for each droplet capture, resulting in

a final sample size of n=2 per group for the single-nuclei multiomics analysis. **f.** UMAP plot showing

cell type-specific clusters of the dorsal hippocampus. **g.** Volcano plots depicting genes that exhibit

differential expression and chromatin accessibility following crotonate treatment. Genes labeled with

color are DEGs that also exhibit DARs, while depict genes that just exhibit differential expression but do

not exhibit DARs are labeled in gray. **h-i.** Upset plots depicting unique and overlapping signatures of

DEGs with DARs across all the hippocampal principal neurons. **h.** Upregulated DEGs with DARs. **i.**

Downregulated DEGs with DARs. All box plots: the center line represents the median, the box edges

represent the top and bottom quartiles (25th to 75th percentiles), and the minimum and maximum

whiskers.

Figure 3. *Increases in Kcr elicits distinct molecular signatures in principal hippocampal*

neuronal cell-types. **a-d.** Cnet plots depicting the top five enriched pathways (GO: Molecular Function)

and their corresponding DEGs with DARs in hippocampal subregions CA1 **(a)**, CA3 **(b)**, Subiculum **(c)**,

and Dentate gyrus **(d)** that are upregulated in response to crotonate administration. **e-f.** Single-nuclei

multiomics reveals transcription factor motif sites enriched on gene promoters that show differential

expression and differential chromatin plasticity following crotonate treatment. **e.** Transcription factor

(TF) motif enrichment analysis from upregulated genes with increased chromatin accessibility at the

promoter upon crotonate administration. **f.** Top 5 significant TF motifs enriched from CA1, CA3, and

subiculum.

Figure 4. *Increases in Kcr augments glutamatergic neurotransmission. a.* Differential incoming

interaction strength and differential outgoing interaction strength in crotonate-treated mice compared to

saline-treated mice in hippocampal subregions CA1, CA3, Subiculum, and DG. Differential changes in

interaction strength in 'Shared' (both incoming and outgoing connections in both the groups), 'Incoming

specific' (incoming connections in both the groups), 'Outgoing specific' (outgoing connections in both

the groups), and 'Incoming & Outgoing specific' (both incoming and outgoing connections in either of

the groups) clusters are depicted by symbols. **b.** Individual ligand-receptor interactions significantly

enhanced upon crotonate treatment in intrahippocampal projections of DG-CA3, CA3-CA1, and CA1-

subiculum. **c.** Interaction strength of ligand-receptor genes involved in glutamatergic signaling across

the principal hippocampal neuronal networks in saline-treated and crotonate-treated mice.

Communication probability, represented by numbers ranging from 0 to 10, is visualized through the

edge width of intrahippocampal connections, indicating the strength of communication between the

respective subregions. **d-h.** Kcr enhancement leads to larger stimulus-evoked glutamate transients in

acute hippocampal slices. **d.** Schematic of the experimental design. **e.** Representative traces of evoked

glutamate transient responses at 10, 20 and 50 Hz in control slices and ones incubated with crotonate

(5mM). **f.** Image panels showing stimulus-evoked glutamate transient response over time in the

hippocampus upon stimulating the Schaffer collaterals in the CA1 Stratum Radiatum. The black object

in each image depicts the stimulating electrode. t=0 denotes the time point corresponding to the peak

amplitude; t–500 represents the time point 500 ms prior to t=0. **g.** The peak amplitude of evoked

glutamate transients was significantly higher in crotonate-treated slices compared to control across all

the stimulation frequencies. Normalized Peak Amplitude across the groups. For 10 Hz (top): Lognormal

Welch's t test: t (13.96)□= 2.37, *p□=□0.0327. aCSF (n□=□9) and Crotonate (n□=□10). For 20 Hz

(middle): Lognormal Welch's t test: t (9.982) = 2.302, *p□=□0.0442. aCSF (n□=□9) and Crotonate

(n□=□10). For 50 Hz (bottom): Lognormal Welch's t test: t (14.72) = 2.274, *p□=□0.0384. aCSF

(n□=□9) and Crotonate (n□=□10). **h.** The area under the curve (AUC) of evoked glutamate transients

was significantly higher in crotonate-treated slices compared to control in the 10 Hz and 20 Hz

stimulation groups. For 10 Hz (top): Lognormal Welch's t test: t (10.34)□= 4.043, **p□=□0.0022. aCSF

(n□=□9) and Crotonate (n□=□10). For 20 Hz (middle): Lognormal Welch's t test: t (14.96) = 2.263,

*p□=□0.039. aCSF (n□=□9) and Crotonate (n□=□10). For 50 Hz (bottom): Lognormal Welch's t test: t

(16.92) = 1.625, p□=□0.1226. aCSF (n□=□9) and Crotonate (n□=□10). All scatter plots: the center line

represents the mean, error bars represent ±SEM.

Figure 5. ***Kcr enhancement amplifies post-synaptic responsiveness to increases in presynaptic***

glutamate release. **a**. Violin plot depicting expression of *Gria4* mRNA in hippocampal principal

neuronal cell types across saline- and crotonate-treated conditions. **b.** RNAscope validation of

crotonate-mediated increases in *Gria4* expression in hippocampal CA1. Mice were trained with a sub-

optimal SOR training protocol, administered saline or crotonate (200 mg/kg) immediately after training,

and perfused 1 hr later. Scalebar: 100 µm. **c.** Quantification of **b**. For CA1: Normalized Mean

Fluorescent Intensity (MFI) of *Gria4* levels across the groups. Unpaired t test: t (6)□= 2.613,

*p□=□0.04. Homecage (n□=□4) and SOR+1 hr (n□=□4), males only. **d-h.** Kcr enhancement leads to

larger stimulus-evoked calcium transients in acute hippocampal slices. **d.** Schematic of the

experimental design. **e.** Representative traces of evoked calcium transient responses at 10, 20 and 50

Hz frequencies in control and crotonate (5mM)-incubated slices. **f.** Image panels showing stimulus-

evoked calcium transients over time in the hippocampus upon stimulating the Schaffer collaterals in the

CA1 Stratum Radiatum. The black object in each image depicts the stimulating electrode. t=0 denotes

the time point corresponding to the peak amplitude; t–500 represents the time point 500 ms prior to t=0.

g. The peak amplitude of evoked calcium transients was significantly higher in crotonate-treated slices

compared to control across all the stimulation frequencies. Normalized Peak Amplitude across the

groups. For 10 Hz (top): Lognormal Welch's t test: t (17.84)□= 2.803, *p□=□0.118. aCSF (n□=□11)

and Crotonate (n□=□10). For 20 Hz (middle): Lognormal Welch's t test: t (17.18) = 2.734, *p□=□0.014.

aCSF (n□=□11) and Crotonate (n□=□10). For 50 Hz (bottom): Lognormal Welch's t test: t (18.98) =

3.134, **p□=□0.0055. aCSF (n□=□11) and Crotonate (n□=□10). **h.** The area under the curve (AUC) of

evoked calcium transients was significantly higher in crotonate-treated slices compared to control in the

10 Hz and 50 Hz stimulation groups. For 10 Hz (top): Lognormal Welch's t test: t (18.06)□= 2.260,

*p□=□0.0364. aCSF (n□=□11) and Crotonate (n□=□10). For 20 Hz (middle): Lognormal Welch's t

test: t (18.43) = 1.702, p□=□0.1056. aCSF (n□=□11) and Crotonate (n□=□10). For 50 Hz (bottom):

Lognormal Welch's t test: t (13.52) = 3.241, **p□=□0.0062. aCSF (n□=□11) and Crotonate (n□=□10).

All scatter plots: the center line represents the mean, error bars represent ±SEM. All box plots: the

center line represents the median, the box edges represent the top and bottom quartiles (25th to 75th

percentiles), and the minimum and maximum whiskers.

Acknowledgments

908   We thank the Neural Circuits and Behavior Core and the Iowa Institute of Human Genetics (IIHG) core

909   for using their facilities, Emily N. Walsh, Lucy Langmack, Quinlan Truax, Adam J Rauckhorst, and

910   Rebekah M. Peplinski for their technical assistance, and K. Peter Giese and Jacob Michaelson for their

911   valuable input on this work.

Funding

This work was supported by grants from the National Institute of Health R01 MH 087463 to T.A.,

Alzheimer's Association Research Grant AARG-23-1074289 to SC, and The National Institute of Health

R00 AG068306 to S.C. T.A. is also supported by the Roy J. Carver Charitable Trust. The sequencing

data presented herein were obtained at the Genomics Division of the Iowa Institute of Human Genetics

(RRID: SCR_023422), which is supported, in part, by the University of Iowa Carver College of

Medicine.

Author contributions

S.C. and T.A. conceived the study. S.C. and U.M. designed the experiments, interpreted the results,

and wrote the article with inputs from T.A. and E.B.T. U.M., S.C., and S.E.B. performed the behavioral

tasks, stereotactic surgeries, drug treatment, and biochemical experiments. S.G. and N.R. assisted with

the analysis of behavioral data and biochemical experiments. E.B.T. provided the ACSS2 mice. B.B.

performed the bioinformatic analyses with inputs from Y.V. P.S performed the functional

epifluorescence experiments with inputs from J.G. All the authors edited the manuscript.

Competing interests

927 T.A. serves on the Scientific Advisory Board of EmbarkNeuro and is a scientific advisor to Aditum Bio
928 and Radius Health. The other authors declare no conflicting interests.

Data availability

930 The datasets generated in this study have been deposited in the NCBI Gene Expression Omnibus
931 (GEO) database under accession code GSE281007. Sequencing files for the single-nuclei multiomics
932 (RNA-seq + ATAC-seq) and ChIP-Seq have been made publicly available through GSE281007.

Code availability

934 The code used for the analyses to generate the figures related to Single-nuclei multiomics data can be
935 accessed through GitHub (https://github.com/ChatterjeeEpigenetics/CrotonylationMultiomics2024).

References

1   Graff, J., Woldemichael, B. T., Berchtold, D., Dewarrat, G. & Mansuy, I. M. Dynamic histone
marks in the hippocampus and cortex facilitate memory consolidation. *Nat Commun* **3**, 991
(2012). https://doi.org/10.1038/ncomms1997
2   Levenson, J. M. *et al.* Regulation of histone acetylation during memory formation in the
hippocampus. *J Biol Chem* **279**, 40545-40559 (2004). https://doi.org/10.1074/jbc.M402229200
3   Maze, I., Noh, K. M. & Allis, C. D. Histone regulation in the CNS: basic principles of epigenetic
plasticity. *Neuropsychopharmacology* **38**, 3-22 (2013). https://doi.org/10.1038/npp.2012.124
4   Peixoto, L. & Abel, T. The role of histone acetylation in memory formation and cognitive
impairments. *Neuropsychopharmacology* **38**, 62-76 (2013). https://doi.org/10.1038/npp.2012.86
5   Zovkic, I. B., Guzman-Karlsson, M. C. & Sweatt, J. D. Epigenetic regulation of memory
formation and maintenance. *Learn Mem* **20**, 61-74 (2013).
https://doi.org/10.1101/lm.026575.112
6   Sabari, B. R., Zhang, D., Allis, C. D. & Zhao, Y. Metabolic regulation of gene expression through
histone acylations. *Nat Rev Mol Cell Biol* **18**, 90-101 (2017).
https://doi.org/10.1038/nrm.2016.140
7   Verdin, E. & Ott, M. 50 years of protein acetylation: from gene regulation to epigenetics,
metabolism and beyond. *Nat Rev Mol Cell Biol* **16**, 258-264 (2015).
https://doi.org/10.1038/nrm3931
8   Barnes, C. E., English, D. M. & Cowley, S. M. Acetylation & Co: an expanding repertoire of
histone acylations regulates chromatin and transcription. *Essays Biochem* **63**, 97-107 (2019).
https://doi.org/10.1042/EBC20180061
9   Bousiges, O. *et al.* Spatial memory consolidation is associated with induction of several lysine-
acetyltransferase (histone acetyltransferase) expression levels and H2B/H4 acetylation-
dependent transcriptional events in the rat hippocampus. *Neuropsychopharmacology* **35**, 2521-
2537 (2010). https://doi.org/10.1038/npp.2010.117

10 Peleg, S. *et al.* Altered histone acetylation is associated with age-dependent memory
impairment in mice. *Science* **328**, 753-756 (2010). https://doi.org/10.1126/science.1186088
11 Kaczmarska, Z. *et al.* Structure of p300 in complex with acyl-CoA variants. *Nat Chem Biol* **13**,
21-29 (2017). https://doi.org/10.1038/nchembio.2217
12 Sabari, B. R. *et al.* Intracellular crotonyl-CoA stimulates transcription through p300-catalyzed
histone crotonylation. *Mol Cell* **58**, 203-215 (2015). https://doi.org/10.1016/j.molcel.2015.02.029
13 Abel, T., Martin, K. C., Bartsch, D. & Kandel, E. R. Memory suppressor genes: inhibitory
constraints on the storage of long-term memory. *Science* **279**, 338-341 (1998).
https://doi.org/10.1126/science.279.5349.338
14 Bahari-Javan, S. *et al.* HDAC1 regulates fear extinction in mice. *J Neurosci* **32**, 5062-5073
(2012). https://doi.org/10.1523/JNEUROSCI.0079-12.2012
15 Guan, J. S. *et al.* HDAC2 negatively regulates memory formation and synaptic plasticity. *Nature*
459, 55-60 (2009). https://doi.org/10.1038/nature07925
16 Alarcon, J. M. *et al.* Chromatin acetylation, memory, and LTP are impaired in CBP+/- mice: a
model for the cognitive deficit in Rubinstein-Taybi syndrome and its amelioration. *Neuron* **42**,
947-959 (2004). https://doi.org/10.1016/j.neuron.2004.05.021
17 Korzus, E., Rosenfeld, M. G. & Mayford, M. CBP histone acetyltransferase activity is a critical
component of memory consolidation. *Neuron* **42**, 961-972 (2004).
https://doi.org/10.1016/j.neuron.2004.06.002
18 Chatterjee, S. *et al.* A novel activator of CBP/p300 acetyltransferases promotes neurogenesis
and extends memory duration in adult mice. *J Neurosci* **33**, 10698-10712 (2013).
https://doi.org/10.1523/JNEUROSCI.5772-12.2013
19 Shang, S., Liu, J. & Hua, F. Protein acylation: mechanisms, biological functions and therapeutic
targets. *Signal Transduct Target Ther* **7**, 396 (2022). https://doi.org/10.1038/s41392-022-01245-
y
20 Tan, M. *et al.* Identification of 67 histone marks and histone lysine crotonylation as a new type of
histone modification. *Cell* **146**, 1016-1028 (2011). https://doi.org/10.1016/j.cell.2011.08.008
21 Abu-Zhayia, E. R., Machour, F. E. & Ayoub, N. HDAC-dependent decrease in histone
crotonylation during DNA damage. *J Mol Cell Biol* **11**, 804-806 (2019).
https://doi.org/10.1093/jmcb/mjz019
22 Crespo, M. *et al.* Multi-omic analysis of gametogenesis reveals a novel signature at the
promoters and distal enhancers of active genes. *Nucleic Acids Res* **48**, 4115-4138 (2020).
https://doi.org/10.1093/nar/gkaa163
23 Dai, S. K. *et al.* Histone crotonylation regulates neural stem cell fate decisions by activating
bivalent promoters. *EMBO Rep* **22**, e52023 (2021). https://doi.org/10.15252/embr.202052023
24 Liu, S. *et al.* Chromodomain Protein CDYL Acts as a Crotonyl-CoA Hydratase to Regulate
Histone Crotonylation and Spermatogenesis. *Mol Cell* **67**, 853-866 e855 (2017).
https://doi.org/10.1016/j.molcel.2017.07.011
25 Ruiz-Andres, O. *et al.* Histone lysine crotonylation during acute kidney injury in mice. *Dis Model*
Mech **9**, 633-645 (2016). https://doi.org/10.1242/dmm.024455
26 Zou, Y. *et al.* Involvement of Histone Lysine Crotonylation in the Regulation of Nerve-Injury-
Induced Neuropathic Pain. *Front Immunol* **13**, 885685 (2022).
https://doi.org/10.3389/fimmu.2022.885685
27 Li, Y. *et al.* Molecular Coupling of Histone Crotonylation and Active Transcription by AF9 YEATS
Domain. *Mol Cell* **62**, 181-193 (2016). https://doi.org/10.1016/j.molcel.2016.03.028
28 Fernandez-Albert, J. *et al.* Immediate and deferred epigenomic signatures of in vivo neuronal
activation in mouse hippocampus. *Nat Neurosci* **22**, 1718-1730 (2019).
https://doi.org/10.1038/s41593-019-0476-2
29 Tyssowski, K. M. *et al.* Different Neuronal Activity Patterns Induce Different Gene Expression
Programs. *Neuron* **98**, 530-546 e511 (2018). https://doi.org/10.1016/j.neuron.2018.04.001

30 Vanrobaeys, Y. *et al.* Mapping the spatial transcriptomic signature of the hippocampus during
memory consolidation. *Nat Commun* **14**, 6100 (2023). https://doi.org/10.1038/s41467-023-
41715-7
31 Yap, E. L. & Greenberg, M. E. Activity-Regulated Transcription: Bridging the Gap between
Neural Activity and Behavior. *Neuron* **100**, 330-348 (2018).
https://doi.org/10.1016/j.neuron.2018.10.013
32 Li, L. *et al.* Inhibition of ACSS2-mediated histone crotonylation alleviates kidney fibrosis via IL-
1beta-dependent macrophage activation and tubular cell senescence. *Nat Commun* **15**, 3200
(2024). https://doi.org/10.1038/s41467-024-47315-3
33 Mews, P. *et al.* Acetyl-CoA synthetase regulates histone acetylation and hippocampal memory.
Nature **546**, 381-386 (2017). https://doi.org/10.1038/nature22405
34 Peixoto, L. L. *et al.* Memory acquisition and retrieval impact different epigenetic processes that
regulate gene expression. *BMC Genomics* **16 Suppl 5**, S5 (2015). https://doi.org/10.1186/1471-
2164-16-S5-S5
35 Broadbent, N. J., Squire, L. R. & Clark, R. E. Spatial memory, recognition memory, and the
hippocampus. *Proc Natl Acad Sci U S A* **101**, 14515-14520 (2004).
https://doi.org/10.1073/pnas.0406344101
36 Moser, M. B. & Moser, E. I. Functional differentiation in the hippocampus. *Hippocampus* **8**, 608-
619 (1998). https://doi.org/10.1002/(SICI)1098-1063(1998)8:6<608::AID-HIPO3>3.0.CO;2-7
37 Moser, M. B., Moser, E. I., Forrest, E., Andersen, P. & Morris, R. G. Spatial learning with a
minislab in the dorsal hippocampus. *Proc Natl Acad Sci U S A* **92**, 9697-9701 (1995).
https://doi.org/10.1073/pnas.92.21.9697
38 Abu-Zhayia, E. R. *et al.* CDYL1-dependent decrease in lysine crotonylation at DNA double-
strand break sites functionally uncouples transcriptional silencing and repair. *Mol Cell* **82**, 1940-
1955 e1947 (2022). https://doi.org/10.1016/j.molcel.2022.03.031
39 Liu, Y. *et al.* Chromodomain Y-like Protein-Mediated Histone Crotonylation Regulates Stress-
Induced Depressive Behaviors. *Biol Psychiatry* **85**, 635-649 (2019).
https://doi.org/10.1016/j.biopsych.2018.11.025
40 Yuan, H. *et al.* Lysine catabolism reprograms tumour immunity through histone crotonylation.
Nature **617**, 818-826 (2023). https://doi.org/10.1038/s41586-023-06061-0
41 Zheng, Y. *et al.* Modulation of cellular metabolism by protein crotonylation regulates pancreatic
cancer progression. *Cell Rep* **42**, 112666 (2023). https://doi.org/10.1016/j.celrep.2023.112666
42 Telese, F. *et al.* LRP8-Reelin-Regulated Neuronal Enhancer Signature Underlying Learning and
Memory Formation. *Neuron* **86**, 696-710 (2015). https://doi.org/10.1016/j.neuron.2015.03.033
43 Liu, Y. *et al.* CDYL suppresses epileptogenesis in mice through repression of axonal Nav1.6
sodium channel expression. *Nat Commun* **8**, 355 (2017). https://doi.org/10.1038/s41467-017-
00368-z
44 Chatterjee, S. *et al.* Pharmacological activation of Nr4a rescues age-associated memory
decline. *Neurobiol Aging* **85**, 140-144 (2020).
https://doi.org/10.1016/j.neurobiolaging.2019.10.001
45 McQuown, S. C. & Wood, M. A. HDAC3 and the molecular brake pad hypothesis. *Neurobiol
Learn Mem* **96**, 27-34 (2011). https://doi.org/10.1016/j.nlm.2011.04.005
46 Stefanko, D. P., Barrett, R. M., Ly, A. R., Reolon, G. K. & Wood, M. A. Modulation of long-term
memory for object recognition via HDAC inhibition. *Proc Natl Acad Sci U S A* **106**, 9447-9452
(2009). https://doi.org/10.1073/pnas.0903964106
47 Rauckhorst, A. J. *et al.* A hierarchical hepatic de novo lipogenesis substrate supply network
utilizing pyruvate, acetate, and ketones. *Cell Metab* (2024).
https://doi.org/10.1016/j.cmet.2024.10.013
48 Basu, J. & Siegelbaum, S. A. The Corticohippocampal Circuit, Synaptic Plasticity, and Memory.
Cold Spring Harb Perspect Biol **7** (2015). https://doi.org/10.1101/cshperspect.a021733

49   Basu, J. *et al.* A cortico-hippocampal learning rule shapes inhibitory microcircuit activity to
enhance hippocampal information flow. *Neuron* **79**, 1208-1221 (2013).
https://doi.org/10.1016/j.neuron.2013.07.001
50   Klausberger, T. & Somogyi, P. Neuronal diversity and temporal dynamics: the unity of
hippocampal circuit operations. *Science* **321**, 53-57 (2008).
https://doi.org/10.1126/science.1149381
51   Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat Commun* **12**,
1088 (2021). https://doi.org/10.1038/s41467-021-21246-9
52   Suryavanshi, P., Sawant-Pokam, P., Clair, S. & Brennan, K. C. Increased presynaptic
excitability in a migraine with aura mutation. *Brain* **147**, 680-697 (2024).
https://doi.org/10.1093/brain/awad326
53   Hale, W. D., Huganir, R. L. & Twomey, E. C. Architecture, Activation, and Conformational
Plasticity in the GluA4 AMPA Receptor. *bioRxiv* (2025).
https://doi.org/10.1101/2025.06.12.659357
54   Guan, Z. *et al.* Integration of long-term-memory-related synaptic plasticity involves bidirectional
regulation of gene expression and chromatin structure. *Cell* **111**, 483-493 (2002).
https://doi.org/10.1016/s0092-8674(02)01074-7
55   Wellen, K. E. *et al.* ATP-citrate lyase links cellular metabolism to histone acetylation. *Science*
324, 1076-1080 (2009). https://doi.org/10.1126/science.1164097
56   Zhao, D. *et al.* YEATS2 is a selective histone crotonylation reader. *Cell Res* **26**, 629-632 (2016).
https://doi.org/10.1038/cr.2016.49
57   Erb, T. J., Brecht, V., Fuchs, G., Muller, M. & Alber, B. E. Carboxylation mechanism and
stereochemistry of crotonyl-CoA carboxylase/reductase, a carboxylating enoyl-thioester
reductase. *Proc Natl Acad Sci U S A* **106**, 8871-8876 (2009).
https://doi.org/10.1073/pnas.0903939106
58   Chatterjee, S. *et al.* Endoplasmic reticulum chaperone genes encode effectors of long-term
memory. *Sci Adv* **8**, eabm6063 (2022). https://doi.org/10.1126/sciadv.abm6063
59   Guzowski, J. F., McNaughton, B. L., Barnes, C. A. & Worley, P. F. Environment-specific
expression of the immediate-early gene Arc in hippocampal neuronal ensembles. *Nat Neurosci*
2, 1120-1124 (1999). https://doi.org/10.1038/16046
60   Josselyn, S. A. & Tonegawa, S. Memory engrams: Recalling the past and imagining the future.
Science **367** (2020). https://doi.org/10.1126/science.aaw4325
61   Jenuwein, T. & Allis, C. D. Translating the histone code. *Science* **293**, 1074-1080 (2001).
https://doi.org/10.1126/science.1063127
62   Strahl, B. D. & Allis, C. D. The language of covalent histone modifications. *Nature* **403**, 41-45
(2000). https://doi.org/10.1038/47412
63   Turner, B. M. Cellular memory and the histone code. *Cell* **111**, 285-291 (2002).
https://doi.org/10.1016/s0092-8674(02)01080-2
64   Jeyaraj, D. *et al.* Klf15 orchestrates circadian nitrogen homeostasis. *Cell Metab* **15**, 311-323
(2012). https://doi.org/10.1016/j.cmet.2012.01.020
65   Miao, C., Du, L., Zhang, Y., Jia, F. & Shan, L. Novel de novo ZNF148 truncating variant causing
autism spectrum disorder, attention deficit hyperactivity disorder, and intellectual disability. *Clin
Genet* **103**, 364-368 (2023). https://doi.org/10.1111/cge.14272
66   Okamoto, S., Sherman, K., Bai, G. & Lipton, S. A. Effect of the ubiquitous transcription factors,
SP1 and MAZ, on NMDA receptor subunit type 1 (NR1) expression during neuronal
differentiation. *Brain Res Mol Brain Res* **107**, 89-96 (2002). https://doi.org/10.1016/s0169-
328x(02)00440-0
67   Mataga, N., Fujishima, S., Condie, B. G. & Hensch, T. K. Experience-dependent plasticity of
mouse visual cortex in the absence of the neuronal activity-dependent marker egr1/zif268. *J
Neurosci* **21**, 9724-9732 (2001). https://doi.org/10.1523/JNEUROSCI.21-24-09724.2001

Renaudineau, S., Poucet, B., Laroche, S., Davis, S. & Save, E. Impaired long-term stability of CA1 place cell representation in mice lacking the transcription factor zif268/egr1. *Proc Natl Acad Sci U S A* **106**, 11771-11775 (2009). https://doi.org/10.1073/pnas.0900484106

Veyrac, A. *et al.* Zif268/egr1 gene controls the selection, maturation and functional integration of adult hippocampal newborn neurons by learning. *Proc Natl Acad Sci U S A* **110**, 7062-7067 (2013). https://doi.org/10.1073/pnas.1220558110

Wei, F., Xu, Z. C., Qu, Z., Milbrandt, J. & Zhuo, M. Role of EGR1 in hippocampal synaptic enhancement induced by tetanic stimulation and amputation. *J Cell Biol* **149**, 1325-1334 (2000). https://doi.org/10.1083/jcb.149.7.1325

Sun, Z. *et al.* EGR1 recruits TET1 to shape the brain methylome during development and upon neuronal activity. *Nat Commun* **10**, 3892 (2019). https://doi.org/10.1038/s41467-019-11905-3

Trizzino, M. *et al.* EGR1 is a gatekeeper of inflammatory enhancers in human macrophages. *Sci Adv* **7** (2021). https://doi.org/10.1126/sciadv.aaz8836

Wilk, A. J., Shalek, A. K., Holmes, S. & Blish, C. A. Comparative analysis of cell-cell communication at single-cell resolution. *Nat Biotechnol* **42**, 470-483 (2024). https://doi.org/10.1038/s41587-023-01782-z

Huang, Y., Doherty, J. J. & Dingledine, R. Altered histone acetylation at glutamate receptor 2 and brain-derived neurotrophic factor genes is an early event triggered by status epilepticus. *J Neurosci* **22**, 8422-8428 (2002). https://doi.org/10.1523/JNEUROSCI.22-19-08422.2002

Stadler, F. *et al.* Histone methylation at gene promoters is associated with developmental regulation and region-specific expression of ionotropic and metabotropic glutamate receptors in human brain. *J Neurochem* **94**, 324-336 (2005). https://doi.org/10.1111/j.1471-4159.2005.03190.x

Wang, Z. *et al.* NEAT1 regulates neuroglial cell mediating Abeta clearance via the epigenetic regulation of endocytosis-related genes expression. *Cell Mol Life Sci* **76**, 3005-3018 (2019). https://doi.org/10.1007/s00018-019-03074-9

Shao, Z., Zhang, Y., Yuan, G. C., Orkin, S. H. & Waxman, D. J. MAnorm: a robust model for quantitative comparison of ChIP-Seq data sets. *Genome Biol* **13**, R16 (2012). https://doi.org/10.1186/gb-2012-13-3-r16

Stuart, T., Srivastava, A., Madad, S., Lareau, C. A. & Satija, R. Single-cell chromatin state analysis with Signac. *Nat Methods* **18**, 1333-1341 (2021). https://doi.org/10.1038/s41592-021-01282-5

Zonta, B. *et al.* A critical role for Neurofascin in regulating action potential initiation through maintenance of the axon initial segment. *Neuron* **69**, 945-956 (2011). https://doi.org/10.1016/j.neuron.2011.02.021

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

Figure 1

Figure 2

Figure 3

a CA1

metal ion transmembrane transporter activity
phosphatase binding
cell adhesion molecule binding
monoatomic ion gated channel activity
gated channel activity

b CA3

monoatomic ion channel activity
cell adhesion molecule binding
ligand−gated monoatomic ion channel activity
ligand−gated channel activity
monoatomic ion gated channel activity

c Subiculum

PDZ domain binding
cell adhesion molecule binding
structural constituent of synapse
ion channel regulator activity
calcium−dependent phospholipid binding

d Dentate gyrus

calcium−dependent protein kinase C activity
protein kinase C activity
calcium−dependent protein serine/threonine kinase activity
calcium−dependent protein kinase activity

e TF motif enrichment on upregulated gene promoters

KLF15, SP9, ZBTB14, MAZ, ZNF148, KLF16, SP8, CTCFL, EGR1, KLF6, SP3, KLF4, KLF2, TFAP2A(var.2), KLF14, KLF5, ZNF740, E2F6, PRDM1, ZNF460, TFAP2C(var.3), SNAI2, RBPJ, ZBTB33, SP1, Rbpjl, ZNF263, ZIC3, KLF10, SNAI3, Ahr::Arnt

−log10(P Value)

CA1 CA3 SUB

f

CA1	CA3	SUB
KLF15	KLF16	SNAI2
SP9	ZNF740	MAZ
ZBTB14	MAZ	RBPJ
SP3	ZBTB14	ZNF148
SP8	ZNF148	ZBTB33

Figure 4

Figure 5

a *Gria4*

Adj P val 1.12412E-32
Adj P val 3.13821E-14
Adj P val 5.60507E-10
Adj P val 4.4839E-10

Expression Level

CA1 CA3 Sub DG

Saline Crotonate

b *Gria4* RNAscope

Saline

CA1

Gria4 DAPI

Crotonate

CA1

Gria4 DAPI

250 μm

c *Gria4* (CA1)

Normalized Gria4 MFI (Fold saline)

*

Saline Crotonate

d ITR — Synapsin — GCaMP6s — SV40 polyA — ITR

2 Weeks

Stim Electrode

Mesoscope imaging (GCaMP)

aCSF/Crotonate

e
— aCSF
— Crotonate
■ Stimulus

0.25 ΔF/F₀, 2 S

10 Hz

20 Hz

50 Hz

f
10 Hz
aCSF Crotonate
t-500
t=0
-1 ΔF/F₀ 1

20 Hz
aCSF Crotonate
t-500
t=0
-1 ΔF/F₀ 1

50 Hz
aCSF Crotonate
t-500
t=0
-1 ΔF/F₀ 1

g
Peak (ΔF/F₀) (Fold aCSF)
*
aCSF Crotonate

Peak (ΔF/F₀) (Fold aCSF)
*
aCSF Crotonate

Peak (ΔF/F₀) (Fold aCSF)
**
aCSF Crotonate

h
AUC (ΔF/F₀ x S) (Fold aCSF)
*
aCSF Crotonate

AUC (ΔF/F₀ x S) (Fold aCSF)
aCSF Crotonate

AUC (ΔF/F₀ x S) (Fold aCSF)
**
aCSF Crotonate