The Adhesion GPCR ADGRL2 engages Ga13 to Enable Epidermal Differentiation

Xue Yang[1,2,10], Feng He[3,10], Douglas F. Porter[1], Krassimira Garbett[4], Robin M. Meyers[1,5], David L. Reynolds[1], Duy Lan Huong Bui[4], Audrey Hong[1], Luca Ducoli[1], Zurab Siprashvili[1], Vanessa Lopez-Pajares[1], Smarajit Mondal[1], Lisa Ko[1], Yuqing Jing[6], Shiying Tao[1], Bharti Singal[3], Richard Sando[4], Georgios Skiniotis[3,7,8]*, Paul A. Khavari[1,2,9,11]*

[1] Program in Epithelial Biology, Stanford University School of Medicine, Stanford, CA, USA.
[2] Program in Cancer Biology, Stanford University, Stanford, CA, USA.
[3] Department of Molecular and Cellular Physiology, Stanford University School of Medicine, Stanford, CA, USA
[4] Department of Pharmacology, Vanderbilt Brain Institute, Vanderbilt University, Nashville, TN 37240, USA
[5] Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA
[6] Departments of Pediatrics and Genetics, Stanford University, Stanford, CA, USA
[7] Department of Structural Biology, Stanford University School of Medicine, Stanford, CA, USA
[8] Present address: Center of Excellence for Structural Cell Biology, Department of Structural Biology, St Jude Children's Research Hospital, Memphis, TN 38120, USA
[9] Veterans Affairs Palo Alto Healthcare System, Palo Alto, CA, USA.
[10] These authors contributed equally.
[11] Lead Contact.
* Correspondence: khavari@stanford.edu and yiorgo@stanford.edu

Abstract

Homeostasis relies on signaling networks controlled by cell membrane receptors. Although G-protein-coupled receptors (GPCRs) are the largest family of transmembrane receptors, their specific roles in the epidermis are not fully understood. Dual CRISPR-Flow and single cell Perturb-seq knockout screens of all epidermal GPCRs were thus performed, uncovering an essential requirement for adhesion GPCR ADGRL2 (latrophilin 2) in epidermal differentiation. Among potential downstream guanine nucleotide-binding G proteins, ADGRL2 selectively activated Gα13. Perturb-seq of epidermal G proteins and follow-up tissue knockouts verified that Gα13 is also required for epidermal differentiation. A cryo-electron microscopy (cryo-EM) structure in lipid nanodiscs showed that ADGRL2 engages with Gα13 at multiple interfaces, including via a novel interaction between ADGRL2 intracellular loop 3 (ICL3) and a Gα13-specific QQQ glutamine triplet sequence in its GTPase domain. In situ gene mutation of this interface sequence impaired epidermal differentiation, highlighting an essential new role for an ADGRL2-Gα13 axis in epidermal differentiation.

16 Introduction

The epidermis is a self-renewing surface tissue composed primarily of specialized epithelial

keratinocytes that receive signals via a variety of cell surface receptors. The balance between

growth and differentiation in the epidermis represents a prototype for tissue homeostasis.

Dysregulation of epidermal homeostasis contributes to numerous human clinical disorders,

including cancer[1], psoriasis[2] and chronic wounds[3]. GPCRs comprise the largest family of

transmembrane receptors in mammals[4, 5]. Several studies have addressed the role of specific

GPCRs in epidermis. For example, activating mutations in the *SMO* GPCR gene are pathogenic

in basal cell carcinoma[6] and the SMO inhibitor, Vismodegib, prevents cancer progression[7]. Other

examples include the Frizzled GPCR, which regulates keratinocyte proliferation[8], as well as

LPAR GPCRs that enable epidermal differentiation[9]. The roles of the many other GPCRs

expressed in epidermis, however, are largely uncharacterized.

Traditional pooled CRISPR screening methods are of limited utility in characterizing the actions

of gene family members in epidermal homeostasis. Standard cell survival and proliferation

phenotypes within bulk cell populations can fail to detect genes with essential roles in

biomedically important cell transitions, such as occurs in epithelial differentiation[10]. Several

CRISPR screening approaches may overcome this limitation. For example, fluorescence

sorting-based pooled CRISPR screening (**CRISPR-Flow**), separates cells based on expression

of marker proteins representative of specific cell states[11, 12], such as KRT10 for epidermal

differentiation. Single cell Perturb RNA-sequencing (**Perturb-seq**) is another method that

quantifies a subset of the transcriptome in cells undergoing a variety of genetic manipulations[13, 14].

Integrating these two methods may enable systematic evaluation of the roles of members of

diverse gene families in the control of homeostatic gene expression programs.

Here we performed CRISPR-Flow and Perturb-seq knockout screens of all GPCRs expressed

in epidermis. These screens validated previously reported GPCR roles while also identifying a

new essential requirement for the ADGRL2 adhesion GPCR (**aGPCR**) in epidermal

differentiation. ADGRL2 expression was strongly upregulated during keratinocyte differentiation.

In epidermal tissue, ADGRL2 was required for induction of a key differentiation genes involved

in barrier formation. Among potential downstream G proteins, ADGRL2 selectively activated

$G\alpha 13$ ($G\alpha 13/G\beta/G\gamma$). Perturb-seq screening of genes encoding $G\alpha$ subunits expressed in

epidermis confirmed an essential requirement for Gα13 in epidermal differentiation. Cryo-
electron microscopy (**<u>Cryo-EM</u>**) of nanodisc embedded ADGRL2-Gα13 complexes facilitated
high-resolution structure determination. Structural analysis revealed an ADGRL2 ICL3 loop
interface with a QQQ sequence on Gα13, which appears to be a unique GPCR-G protein
interaction that may help explain the preferential engagement of ADGRL2 with Gα13. Mutation
of the Gα13 QQQ sequence diminished ADGRL2 binding and impaired epidermal differentiation,
underscoring the functional importance of this interface in newly uncovered pro-differentiation
roles for the ADGRL2 GPCR and its Gα13 protein target.

Results

CRISPR-Flow and Perturb-seq of epidermal GPCRs.

Analysis of dynamic epidermal differentiation RNA-seq[15] identified 101 GPCRs expressed in epidermal keratinocytes. These span the five known GPCR sub-families (**Fig. 1a**). A CRISPR library targeting these GPCRs was next generated. sgRNAs were included to target known positive controls essential for epidermal homeostasis as well as non-targeting negative controls, encompassing 620 sgRNAs in total (**Extended Data Fig. S1a-b, Supplementary Table 1**). Quality control assessments confirmed minimal skew and full coverage of the resulting plasmid library (**Extended Data Fig. 1c**), which was designed to support both CRISPR-Flow and Perturb-seq screening assays. The CRISPR-Flow screen employed flow cytometry to isolate populations expressing high and low levels of the KRT10 epidermal differentiation marker for sgRNA enrichment/depletion analysis to nominate pro-differentiation and pro-progenitor GPCRs while the single-cell Perturb-seq screen evaluated GPCR gene knockout effects on keratinocyte progenitor and differentiation mRNA expression signatures (**Fig. 1b**). The complementary nature of these screens, one based on the expression of a single canonical early differentiation protein representing entry into the differentiation pathway and the other on the levels of a panel of keratinocyte mRNAs, was designed to catalog GPCRs with non-redundant impacts on homeostatic gene expression in epidermal keratinocytes.

Flow cytometry benchmarking for the CRISPR-Flow effort demonstrated that the KRT10-high population increased to 13.5% by day 3 of calcium-induced differentiation in vitro compared to <0.05% in day 0 progenitors (**Extended Data Fig. 1d-e**). Known positive controls required for epidermal cell differentiation, such as *IRF6, NOTCH2, PRDM1, MAF, CEBPA* and *ZNF750,* as well as the *LPAR3* GPCR, were scored by CRISPR-Flow as required for KRT10 induction. Knockout of known progenitor genes, such as *ITGB1*, *FERMT1*, *TP63*, *EGFR*, and *YAP1*, in contrast, led to increased KRT10 protein levels. These findings confirmed prior work and validated the fidelity of the screen. Among GPCRs, ADGRL2 was nominated as the strongest novel pro-differentiation GPCR (**Fig. 1c-d, Supplementary Table 2**). CRISPR-Flow screening thus validated positive controls with known impacts on homeostatic gene expression in epidermal cells while also nominating specific GPCRs as new regulators of this process.

Single-cell RNA-seq of differentiating keratinocyte populations for Perturb-seq benchmarking
revealed a trajectory from progenitor-like cells to more differentiated states (**Fig. 1e**), consistent
with previous studies[16]. Representative differentiation genes, such as *KRT10*, *SBSN* and
KRTDAP, as well as undifferentiated progenitor population markers, such as *ITGB1*, *ITGA6* and
KRT14, exhibited expected expression dynamics across pseudotime (**Fig. 1f**), supporting
scRNA-seq dataset quality. As an additional quality check for CRISPR-Cas9-mediated gene
knockout effectiveness, expression of targeted genes was confirmed to be significantly reduced
compared to non-targeted controls (**Extended Data Fig. 1f**). Perturb-seq assigned known pro-
differentiation genes, *IRF6, NOTCH2/3, PRDM1, TFAP2A*, and *GRHL3*, as well as pro-
progenitor genes, *ITGB1*, *FERMT1*, *TP63*, and *YAP1* to their known roles, supporting the fidelity
of the screen (**Fig. 1g-h**). As in CRISPR-Flow, Perturb-seq identified specific LPAR genes as
pro-differentiation GPCRs, including *LPAR3, LPAR5 and LPAR6* (**Fig. 1g-h**), which are known
to be activated by Lysophosphatidic Acid (**LPA**) and which have been previously implicated in
keratinocyte differentiation[9]. ADGRL2 was again captured as a novel pro-differentiation GPCR
in Perturb-seq data (**Fig. 1g-h, Extended Data Fig. 1g, Supplementary Table 3**). Correlation
of the CRISPR-Flow and Perturb-seq dual screens identified ADGRL2 and LPAR3 among the
top GPCR candidates promoting differentiation (**Fig. 1i**). Integration of Perturb-seq and CRISPR-
Flow screens therefore nominated ADGRL2 as a novel pro-differentiation GPCR.

ADGRL2 is essential for epidermal differentiation

In keratinocyte differentiation RNA-seq datasets[15], mRNA levels of ADGRL2 were significantly
upregulated during differentiation (**Extended Data Fig. 2a, red labeled**). Consistent with this,
uniform manifold approximation and projection (**UMAP**) analysis performed on the Perturb-seq
dataset showed that ADGRL2 mRNA levels were low in progenitors but markedly elevated in
differentiated cells, concordant with *KRTDAP* differentiation gene expression and inversely
correlated to the *MKI67* proliferation marker (**Fig. 2a**). Further evaluation of ADGRL2 mRNA
levels showed significant upregulation during differentiation by qPCR analysis (**Fig. 2b**), along
with accumulation of the self-cleaved dual protein species C-terminal fragment (**CTF**) of
ADGRL2 (**Fig. 2c**). Both ADGRL2 mRNA and protein levels were therefore significantly
increased during differentiation, further suggesting a possible role for ADGRL2 in this process.

In vitro, keratinocyte ADGRL2 depletion was associated with reduced differentiation mRNA
levels as measured by both qPCR (**Fig. 2d, Extended Data Fig. 2b)** and RNA-seq (**Fig. 2e,**
Extended Data Fig. 2c, Supplementary Table 4). Gene Ontology (**GO**) analysis of RNA-seq
data showed that ADGRL2 depletion led to impaired induction of epidermal differentiation genes
in patterns linked to human clinical phenotypes characterized by abnormalities in this process
(**Fig. 2f**). Similarly, in 3-dimensional human skin organoid tissue regenerated from diploid human
keratinocytes treated with siRNAs to ADGRL2 or scrambled control, ADGRL2 depletion led to
decreases in differentiation proteins such as KRT10 and TGM1 in suprabasal epidermis, without
affecting proliferation markers Ki67 (**Fig. 2g, Extended Data Fig. 2d**). To investigate whether
ADGRL2 influences keratinocyte differentiation in a cell autonomous or non-cell autonomous
fashion, mosaic epidermal tissue was generated by mixing HA-KRT14 labeled ADGRL2
knockdown cells with Flag-KRT14 labeled control knockdown cells at 1:1 ratios. ADGRL2-
depleted cells displayed reduced KRT10 differentiation protein expression, in contrast to
adjacent control cells (**Fig. 2h, Extended Data fig. 2e-f)**. These findings nominate ADGRL2 as
a cell-autonomous pro-differentiation GPCR in epidermis.

ADGRL2 activates Gα13
GPCRs interact with Gα subunits of heterotrimeric G proteins to induce downstream signaling.
To identify specific Gα subunits activated by ADGRL2 during keratinocyte differentiation, a
TRUPATH assay[17] was employed. This approach uses the BRET2 sensor to measure the
attenuation of luminescence upon downstream G protein activation[17]. To avoid interference by
endogenous G proteins, a pan G protein protein-knockout (**GKO**) HEK293 cell line[18, 19] was used.
Exogenously expressed ADGRL3 co-transfected with Gα13/Gβ3/Gγ9 served as a positive
control and ADGRL3 co-expressed with GαoB/Gβ3/Gγ8 functioned as a negative control[20]. 12
Gα proteins are expressed across keratinocyte differentiation, as detected by RNA-seq[15]. A
panel of 12 putative G$\alpha\beta\gamma$ "epidermal transducerome" combinations was therefore evaluated in
cells expressing either the full-length ADGRL2 or the ADGRL2 CTF, both of which correctly
localized to the cell membrane (**Extended Data Fig. 3a**). Gα13 and Gα12 subunits displayed
the highest basal activation levels by full-length ADGRL2 and ADGRL2-CTF (**Fig. 3a-b**). The
Tethered Agonist (**TA**) peptide serves as an intrinsic ligand in aGPCRs; once exposed by
proteolytic cleavage, it activates the receptor to initiate intracellular signal transduction[21]. To
further dissect the role of the TA peptide in cleavage-dependent activation, therefore, two

independent systems, PAR1-ADGRL2 and EK-ADGRL2, were employed. In the PAR1-ADGRL2

system, the N-terminal extracellular portion of ADGRL2 was replaced with the N-terminal domain

of protease-activated receptor PAR1, creating a TA-protected variant[22]. Thrombin treatment

cleaves off the PAR1 N terminus, exposing a "scarred" TA peptide with a two amino acids

deletion[22]. Localization of the PAR1-ADGRL2 hybrid protein to the cell membrane was not

abolished due to these modifications (**Extended Data Fig. 3a**). Notably, despite generating a

non-native "scarred" TA peptide[20, 22], the receptor still predominantly activated Gα13 upon

cleavage (**Fig. 3c**). To further study this, an EK-ADGRL2 fusion was also generated. In this

approach, enterokinase-mediated cleavage selectively cleaves off the SNAP tag on the N

terminus of the protein and exposed a native TA peptide[20, 23]. Like the PAR1-ADGRL2 fusion,

localization of the EK-ADGRL2 hybrid protein remained unchanged compared to wild-type full

length ADGRL2 (**Extended Data Fig. 3a**). Upon enterokinase release of the TA peptide, the

induced receptor selectively activated Gα13, reaffirming the preferential activation of Gα13 by

ADGRL2 (**Fig. 3d**), and congruent with a finding reported by Pederick et al[24]. Gα13 activation

by these two TA peptide cleavage mimetics of ADGRL2 are consistent with the full length

ADGRL2 and ADGRL2-CTF results. As a negative control, we assessed the differential dose

responses of ADGRL2-CTF and its TA peptide deletion mutant (**Δ7TA**). This mutant has no

capacity to activate G proteins[25]. We also observed that the Δ7TA mutant failed to activate Gα13

in a dose-dependent fashion (**Fig. 3e**), despite similar expression and subcellular localization as

ADGRL2-CTF (**Extended Data Fig. 3a-b**). Further confirmation was obtained using a Gα13

reporter SRE-RF luciferase assay, which indicated that wild-type ADGRL2-CTF displayed an

enhanced ability to activate Gα13, while the Δ7TA mutant did not (**Extended Data Fig. 3c**).

These data suggest that Gα13 serves as the primary Gα protein subtype activated by ADGRL2.

Gα13 is essential for epidermal tissue differentiation

To investigate if Gα13 may act in a manner consistent with a role as a downstream effector of

ADGRL2 in keratinocytes, an epidermal G-protein Perturb-seq screening experiment was

conducted. The Perturb-seq data confirmed that ablation of *GNA13*, the gene encoding Gα13,

attenuated keratinocyte differentiation (**Extended Data Fig. 3d**). This observation was further

corroborated in 2D culture, where *GNA13* depletion was associated with a significant decrease

in expression of 299 differentiation-enriched genes that are also modulated by ADGRL2 (**Fig. 3f,**

Extended Data Fig. 3e-i, Supplementary Table 5). These findings were then further validated

in regenerated human skin organoid tissue where, similar to *ADGRL2* loss, *GNA13* depletion

reduced differentiation marker expression (**Fig. 3g, Extended Data Fig. 3j**). Gα13, like ADGRL2,

is therefore required for normal induction of epidermal differentiation genes.

Cryo-EM structure of the ADGRL2-Gα13 complex in lipid nanodiscs

To examine the structural basis of Gα13 activation by ADGRL2, human ADGRL2-CTF and a

miniGα13 heterotrimer were co-expressed in insect cells and reconstituted into lipid nanodiscs

for Cryo-EM analysis (**Extended Data Fig. 4a-b**). The Cryo-EM map of the ADGRL2- Gα13

complex was determined at a global nominal resolution of 3.1 Å, with local refinements yielding

further improved maps for the 7TM domain (2.9 Å) and G protein (2.9 Å) (**Extended Data Fig.**

4c-f, Supplementary Table 6). While previously reported aGPCR-G protein structures were

determined in detergent micelles[21, 26, 27, 28, 29, 30], this ADGRL2-Gα13 complex structure provides

an opportunity to examine how an aGPCR couples to G proteins in a lipid bilayer environment.

Of note, two cholesterol (**CLR**)-like molecules were observed bound to the receptor in the Cryo-

EM map, occupying hydrophobic cavities in between TM3 and TM4, and TM4 and TM5,

respectively (**Extended Data Fig. 4g-h**). From a global perspective, the structure of the

ADGRL2-Gα13 complex in lipid nanodiscs is not significantly different from the structure of the

ADGRL3-Gα13 complex in detergent micelles[26] and shows a deeply buried TA peptide in the

orthostatic 7TM binding site, bent TM6 and TM7 for G protein binding, and extensive interactions

between ICL2 and the α5 helix (**Extended Data Fig. 4i-j**). Mutations of key ICL2 residues, such

as the F943A and V942A/F943A mutants, dramatically reduced basal Gα13 activation by the

ADGRL2-CTF without affecting protein localization and stability (**Extended Data Fig. 4k-m**),

highlighting the broadly important role of ICL2 in the tethered agonist activation mechanism of

aGPCRs[30].

ADGRL2 ICL3 in Gα13 activation

In contrast to the previously described Gα13-coupled ADGRL3 structure[26], however, the Gα13-

coupled ADGRL2 structure in lipid nanodiscs revealed a distinct conformation of ICL3, which

protrudes from the lipid bilayer and contacts the GTPase domain of Gα13 (**Fig. 4a, Extended**

Data Fig. 4n-o). Additionally, subtle conformational differences were observed in the

extracellular regions (**Extended Data Fig. 4p**). Unexpectedly, and in contrast to ADGRL3, the

ICL3 of ADGRL2 appears to recognize three glutamine (**QQQ**) residues in the loop between

helix 4 and β-strand 6 of Gα13 (**Fig. 4b**), a region we observe relatively weak EM densities

suggesting the dynamic character of those elements (**Extended Data Fig. 4o**). The interactions

in this region likely facilitate Gα13 coupling by ordering ICL3 and promoting the outward

movement of TM6, which is positioned differently than in the ADGRL3 structure (**Fig. 4b**). Indeed,

3D variability analysis, as implemented previously[31], of the global Cryo-EM map revealed that

ICL3 of the receptor and this Gα13 loop flex together (**Supplementary Movie 1**). Notably, the

QQQ patch of Gα13 is unique across all Gα proteins, and is not even present in its close

subfamily member, Gα12, in spite of the fact that the surrounding sequences are conserved

between them (**Fig. 4c**). This uniqueness suggests the potential importance for these Gα13

residues in determining the ADGRL2 coupling specificity.

To evaluate the importance of the novel ICL3-Gα13 interaction interface, a Gα13 mutant (QQQ

to AAA) was generated and its basal activation by ADGRL2-CTF quantified. The AAA mutant

exhibited similar expression levels to wild-type Gα13 but showed significantly reduced activity

when activated by the wild-type ADGRL2-CTF (**Fig. 4d**). Moreover, the ICL2 mutant ADGRL2-

CTF-F943A could partially activate wild-type Gα13 (**Fig. 4d**) but failed to appreciably activate

the Gα13 AAA mutant (**Fig. 4d**), consistent with the premise that the ICL3-Gα13 interface

contributes to Gα13 activation by ADGRL2. To examine the functional impact of disrupting the

Gα13 QQQ interface with ADGRL2 on differentiation, the endogenous *GNA13* gene was edited

in normal human keratinocytes to substitute AAA for QQQ residues in Gα13. Control editing was

performed at the *GNA13* gene locus to alter codon sequences but preserve the QQQ residues

to generate two isogenic keratinocyte populations genetically identical except for this sequence

(**Fig. 4e**). AAA-edited keratinocytes displayed impaired differentiation gene induction while

control-edited cells showed no such defects (**Fig. 4f**). These findings suggest that intactness of

the newly identified ADGRL2 ICL3-Gα13 interface is required for normal epidermal

differentiation.

Discussion

The results of the CRISPR-Flow and Perturb-seq GPCR screens demonstrated a degree of concordance, not only in identifying a novel role for ADGRL2 but also in detecting LPAR GPCRs with previously reported actions in epidermal differentiation. Some differences, however, were observed between the two screens in the candidate GPCRs identified, likely due to differences in their design. The CRISPR-Flow screen relied on differential sorting based on levels of a single canonical differentiation protein, KRT10, whereas the single-cell RNA-seq approach used in Perturb-seq assessed broader transcriptomic outcomes that were then placed within a single cell differentiation trajectory. In addition to ADGRL2, each knockout screen nominated specific GPCRs for future study as candidate regulators of epidermal homeostasis. Specific known and novel epidermal GPCRs, including S1PR5, GPR157, ADRA2C, GPRC5B, PTGER2, ACKR3, ADRB1, and FZD6, were top hits in CRISPR-Flow. Additional known and novel epidermal GPCRs, such as F2RL1, GPR39, BDKRB2, and GPR153 scored highly in Perturb-seq. Several GPCRs, such as S1PR5 and EDNRA, were assigned contrasting functions between the screens, underscoring the need for single GPCR-focused follow-up functional studies, as performed here for ADGRL2. Combined analysis of both screening methods identified more known and novel GPCRs than either screen alone, suggesting that integrating both methods may be useful in future efforts to identify new regulators of complex genetic programs.

The role of the third intracellular loop in GPCR signaling has been elusive, largely due to its high sequence variability and structural flexibility[32, 33, 34]. While previous structural studies of aGPCR/G protein complexes typically relied on detergent purification methods and primarily highlighted interactions involving intracellular loop 2 (ICL2) of aGPCRs[21, 26, 27, 28, 29, 30], work here utilized lipid nanodiscs[35, 36], which incorporate lipid bilayers that may help recapitulate the native environment of transmembrane proteins. This may have helped to delineate the binding interface involving both ICL2 as well as ICL3 of ADGRL2 with $G\alpha13$. The present Cryo-EM structure-defined nanodisc embedded complex reveals that the ADGRL2 ICL3 engages a unique and functionally important QQQ-containing loop sequence in the GTPase domain of $G\alpha13$, an interaction which has never been observed in previous aGPCR structural studies. The QQQ site of $G\alpha13$ is unique across all $G\alpha$ proteins, and is not present in its close subfamily member, $G\alpha12$, in spite of the fact that the surrounding sequences are conserved between them. Despite the

noticeable conformational flexibility observed in the EM map, the ICL3 of ADGRL2 is positioned
to bind Gα13's glutamine residues, whose mutation attenuated receptor signaling and
differentiation. In agreement with a recent study showing that ICL3 modulates G protein
selectivity of the β2 adrenergic receptor[37], these findings suggest a regulatory role for the
ADGRL2 ICL3 in tuning receptor signaling specificity. The current data represent, to our
knowledge, the first reported instance of an adhesion GPCR ICL3 engaging with a Gα protein
in a structured fashion, and may thus enable future potential efforts designed to modulate
signaling by ADGRL2-Gα13 as well as other aGPCRs and their corresponding Gα subunit
effectors.
280
ADGRL1, ADGRL2, and ADGRL3 reside within the latrophilin subfamily of aGPCRs[38], which
have well characterized roles in synaptic function and maintenance[39, 40]. While ADGRL1 and
ADGRL3 are predominantly expressed in the brain[41, 42], ADGRL2 is expressed more widely[41, 42],
suggesting it plays roles in other tissues. Actions of ADGRL2 in stratified epithelia, however,
have been largely unexplored. Interestingly, both ADGRL2 mRNA and protein levels were
significantly increased during differentiation. ADGRL2 knockdown impaired induction of a host
of genes essential for epidermal differentiation, including a number whose mutation leads to
human clinical disorders of this process. Among these genes are *TGM1, CDSN, LOR, SPINK5*,
and *FLG*, each of whose mutation causes monogenic skin diseases characterized by abnormal
epidermal barrier function, such as autosomal recessive congenital ichthyosis[43], peeling skin
disease[44], keratoderma hereditaria mutilans[45], ichthyosis linearis circumflexa[46], and ichthyosis
vulgaris[47] respectively. The ADGRL2 GPCR is therefore necessary for the induction of
differentiation genes essential for formation of a healthy cutaneous barrier.

Reference

1. Ratushny V, Gober MD, Hick R, Ridky TW, Seykora JT. From keratinocyte to cancer: the pathogenesis and modeling of cutaneous squamous cell carcinoma. *J Clin Invest* **122**, 464-472 (2012).

2. Zhang X, Yin M, Zhang LJ. Keratin 6, 16 and 17-Critical Barrier Alarmin Molecules in Skin Wounds and Psoriasis. *Cells* **8**, (2019).

3. Pastar I*, et al.* Epithelialization in Wound Healing: A Comprehensive Review. *Adv Wound Care (New Rochelle)* **3**, 445-464 (2014).

4. Alhosaini K, Azhar A, Alonazi A, Al-Zoghaibi F. GPCRs: The most promiscuous druggable receptor of the mankind. *Saudi Pharm J* **29**, 539-551 (2021).

5. Weis WI, Kobilka BK. The Molecular Basis of G Protein–Coupled Receptor Activation. *Annual Review of Biochemistry* **87**, 897-919 (2018).

6. Gutzmer R, Solomon JA. Hedgehog Pathway Inhibition for the Treatment of Basal Cell Carcinoma. *Target Oncol* **14**, 253-267 (2019).

7. Sekulic A*, et al.* Efficacy and safety of vismodegib in advanced basal-cell carcinoma. *N Engl J Med* **366**, 2171-2179 (2012).

8. Ghahramani A, Donati G, Luscombe NM, Watt FM. Epidermal Wnt signalling regulates transcriptome heterogeneity and proliferative fate in neighbouring cells. *Genome Biol* **19**, 3 (2018).

9. Sumitomo A*, et al.* LPA Induces Keratinocyte Differentiation and Promotes Skin Barrier Function through the LPAR1/LPAR5-RHO-ROCK-SRF Axis. *J Invest Dermatol* **139**, 1010-1022 (2019).

10. Chen S*, et al.* Genome-wide CRISPR screen in a mouse model of tumor growth and metastasis. *Cell* **160**, 1246-1260 (2015).

11. Condon KJ*, et al.* Genome-wide CRISPR screens reveal multitiered mechanisms through which mTORC1 senses mitochondrial dysfunction. *Proc Natl Acad Sci U S A* **118**, (2021).

12. Fomicheva M, Macara IG. Genome-wide CRISPR screen identifies noncanonical NF-kappaB signaling as a regulator of density-dependent proliferation. *Elife* **9**, (2020).

13. Adamson B*, et al.* A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response. *Cell* **167**, 1867-1882 e1821 (2016).

338 14. Dixit A*, et al.* Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell RNA
339   Profiling of Pooled Genetic Screens. *Cell* **167**, 1853-1866 e1817 (2016).
340

341 15. Kim DS*, et al.* The dynamic, combinatorial cis-regulatory lexicon of epidermal
342   differentiation. *Nat Genet* **53**, 1564-1576 (2021).
343

344 16. Cockburn K*, et al.* Gradual differentiation uncoupled from cell cycle exit generates
345   heterogeneity in the epidermal stem cell layer. *Nat Cell Biol* **24**, 1692-1700 (2022).
346

347 17. Olsen RHJ*, et al.* TRUPATH, an open-source biosensor platform for interrogating the
348   GPCR transducerome. *Nat Chem Biol* **16**, 841-849 (2020).
349

350 18. Alvarez-Curto E*, et al.* Targeted Elimination of G Proteins and Arrestins Defines Their
351   Specific Contributions to Both Intensity and Duration of G Protein-coupled Receptor
352   Signaling. *J Biol Chem* **291**, 27147-27159 (2016).
353

354 19. Grundmann M*, et al.* Lack of beta-arrestin signaling in the absence of active G proteins.
355   *Nature Communications* **9**, (2018).
356

357 20. Bui DLH*, et al.* The adhesion GPCRs CELSR1-3 and LPHN3 engage G proteins via distinct
358   activation mechanisms. *Cell Rep* **42**, 112552 (2023).
359

360 21. Xiao P*, et al.* Tethered peptide activation mechanism of the adhesion GPCRs ADGRG2
361   and ADGRG4. *Nature* **604**, 771-778 (2022).
362

363 22. Mathiasen S*, et al.* G12/13 is activated by acute tethered agonist exposure in the
364   adhesion GPCR ADGRL3. *Nature Chemical Biology* **16**, 1343-1350 (2020).
365

366 23. Perry-Hauser NA, VanDyck MW, Lee KH, Shi L, Javitch JA. Disentangling autoproteolytic
367   cleavage from tethered agonist-dependent activation of the adhesion receptor ADGRL3.
368   *J Biol Chem* **298**, 102594 (2022).
369

370 24. Pederick DT, Perry-Hauser NA, Meng H, He Z, Javitch JA, Luo L. Context-dependent
371   requirement of G protein coupling for Latrophilin-2 in target selection of hippocampal
372   axons. *Elife* **12**, (2023).
373

374 25. Stoveken HM, Hajduczok AG, Xu L, Tall GG. Adhesion G protein-coupled receptors are
375   activated by exposure of a cryptic tethered agonist. *Proc Natl Acad Sci U S A* **112**, 6194-
376   6199 (2015).
377

378 26. Barros-Alvarez X*, et al.* The tethered peptide activation mechanism of adhesion GPCRs.
379   *Nature* **604**, 757-762 (2022).
380

381 27. Ping YQ*, et al.* Structural basis for the tethered peptide activation of adhesion GPCRs.
382   *Nature* **604**, 763-770 (2022).
383

384   28.   Qu X, *et al.* Structural basis of tethered agonism of the adhesion GPCRs ADGRD1 and
385   ADGRF1. *Nature* **604**, 779-785 (2022).
386

387   29.   Qian Y, *et al.* Structural insights into adhesion GPCR ADGRL3 activation and G(q), G(s),
388   G(i), and G(12) coupling. *Mol Cell* **82**, 4340-4352 e4346 (2022).
389

390   30.   Jones DTD, Dates AN, Rawson SD, Burruss MM, Lipper CH, Blacklow SC. Tethered agonist
391   activated ADGRF1 structure and signalling analysis reveal basis for G protein coupling.
392   *Nat Commun* **14**, 2490 (2023).
393

394   31.   Papasergi-Scott MM, *et al.* Time-resolved cryo-EM of G-protein activation by a GPCR.
395   *Nature* **629**, 1182-1191 (2024).
396

397   32.   Probst WC, Snyder LA, Schuster DI, Brosius J, Sealfon SC. Sequence alignment of the G-
398   protein coupled receptor superfamily. *DNA Cell Biol* **11**, 1-20 (1992).
399

400   33.   Katritch V, Cherezov V, Stevens RC. Diversity and modularity of G protein-coupled
401   receptor structures. *Trends Pharmacol Sci* **33**, 17-27 (2012).
402

403   34.   Venkatakrishnan AJ, Flock T, Prado DE, Oates ME, Gough J, Madan Babu M. Structured
404   and disordered facets of the GPCR fold. *Curr Opin Struct Biol* **27**, 129-137 (2014).
405

406   35.   Denisov IG, Sligar SG. Nanodiscs for structural and functional studies of membrane
407   proteins. *Nat Struct Mol Biol* **23**, 481-486 (2016).
408

409   36.   He F, *et al.* Allosteric modulation and G-protein selectivity of the Ca(2+)-sensing
410   receptor. *Nature* **626**, 1141-1148 (2024).
411

412   37.   Sadler F, Ma N, Ritt M, Sharma Y, Vaidehi N, Sivaramakrishnan S. Autoregulation of
413   GPCR signalling through the third intracellular loop. *Nature* **615**, 734-741 (2023).
414

415   38.   Purcell RH, Hall RA. Adhesion G Protein-Coupled Receptors as Drug Targets. *Annu Rev
416   Pharmacol Toxicol* **58**, 429-449 (2018).
417

418   39.   Sando R, Sudhof TC. Latrophilin GPCR signaling mediates synapse formation. *Elife* **10**,
419   (2021).
420

421   40.   Sando R, Jiang X, Sudhof TC. Latrophilin GPCRs direct synapse specificity by coincident
422   binding of FLRTs and teneurins. *Science* **363**, (2019).
423

424   41.   Krishnan A, Nijmeijer S, de Graaf C, Schioth HB. Classification, Nomenclature, and
425   Structural Aspects of Adhesion GPCRs. *Handb Exp Pharmacol* **234**, 15-41 (2016).
426

427   42.   Regan SL, Williams MT, Vorhees CV. Latrophilin-3 disruption: Effects on brain and
428   behavior. *Neurosci Biobehav Rev* **127**, 619-629 (2021).
429

430   43.   Herman ML*, et al.* Transglutaminase-1 gene mutations in autosomal recessive
431   congenital ichthyosis: summary of mutations (including 23 novel) and modeling of
432   TGase-1. *Hum Mutat* **30**, 537-547 (2009).
433
434   44.   van der Velden J, van Geel M, Engelhart JJ, Jonkman MF, Steijlen PM. Mutations in the
435   CDSN gene cause peeling skin disease and hypotrichosis simplex of the scalp. *J Dermatol*
436   **47**, 3-7 (2020).
437
438   45.   Pohler E*, et al.* Novel autosomal dominant mutation in loricrin presenting as prominent
439   ichthyosis. *Br J Dermatol* **173**, 1291-1294 (2015).
440
441   46.   Guerra L*, et al.* Ichthyosis Linearis Circumflexa as the Only Clinical Manifestation of
442   Netherton Syndrome. *Acta Derm Venereol* **95**, 720-724 (2015).
443
444   47.   Thyssen JP, Godoy-Gijon E, Elias PM. Ichthyosis vulgaris: the filaggrin mutation disease.
445   *Br J Dermatol* **168**, 1155-1166 (2013).
446
447

Figure Legends

Fig. 1. CRISPR-Flow and Perturb-seq screens in differentiating normal human keratinocytes. a, Pie chart of the 101 epidermal GPCRs and their classes. **b**, Schematic of experimental workflow for CRISPR screens. **c**, Scatterplot mapping guide enrichment (beta score) in low vs. high KRT10 cells from CRISPR-Flow screening. **d**, Bar graphs quantifying mean beta scores from top CRISPR-Flow identified candidates (red=GPCRs, black=controls). **e**, Principal Component Analysis (PCA) plot representing variance in gene expression between progenitor and differentiated epidermal states at day 3. **f**, Pseudotime trajectory analysis of progenitor markers ITGB1/KRT14/ITGA6 and differentiation markers KRTDAP/SBSN/KRT10 at day 3. **g**, Perturb-seq GPCR and positive control target effects plotted as pseudotime differences from non-target guides (Z score) *vs* Mann-Whitney p values for the comparison to cells with other guides. **h**, Enrichment of knock-out cells along the pseudotime trajectory (Mann-Whitney log10 P) compared to cells with safe target guides. A positive enrichment in low pseudotime cells indicates nominates a gene as necessary for differentiation. **i**, Correlation analysis of CRISPR-Flow beta scores plotted against Perturb-seq pseudotime measurements for each CRISPR target. ADGRL2 is highlighted in red. For all analyses *=p<0.05, **=p<0.01, ***=p<0.001.

Fig. 2. Requirement for ADGRL2 in epidermal differentiation. a, UMAP analysis depicting ADGRL2 expression in progenitor and differentiated states in scRNA-seq data. **b**, ADGRL2 mRNA levels across keratinocyte differentiation in vitro. **c**, ADGRL2 protein levels (C-terminal fragment dual bands) across differentiation. **d**, mRNA expression by qPCR of differentiation markers and progenitor markers with control versus ADGRL2 knockdown (L2 si1 and L2 si2) from four independent human skin donors evaluated. **e**, RNA-Seq analysis of ADGRL2 knockdown keratinocytes sourced from three unrelated skin donors, showing all genes with log2 fold change >0.2 or <-0.2, red/blue genes are the ones upregulated/downregulated during differentiation, white genes are unchanged ones. **f**, Gene Ontology analysis of all significantly altered mRNAs (FDR<0.05) following ADGRL2 knockdown based on the RNA-seq data. **g**, Immunostaining of KRT10, TGM1 and Ki67 in regenerated human skin organoid tissue with ADGRL2 depletion versus control. **h**, Immunostaining of KRT10 in mosaic skin organoids. In **b**

and **c**, one-way ANOVA with Dunnett's test against day 0 cells, n = 3. In **d**, one-way ANOVA

with Dunnett's test against control cells for each gene, n = 4.

Fig. 3. ADGRL2 activates Gα13. (**a, b**) Domain structures, the net BRET2 signals and

summarized BRET2 heatmap from full-length ADGRL2 and ADGRL2-CTF. (**c, d**) Domain

structures with drug treatment conditions, the induced BRET2 signals and summarized BRET2

heatmap from PAR1-ADGRL2 and EK-ADGRL2. **e**, Dose-dependent BRET2 signaling profiles

of ADGRL2-CTF and ADGRL2 TA peptide deficient mutants (Δ7TA) on Gα13. **f**, Gene Ontology

analysis of 299 significant altered mRNAs (FDR<0.1) shared between *GNA13* and *ADGRL2*

knockdown RNA-seq datasets. **g**, Immunostaining for KRT10 and TGM1 in human skin organoid

tissues with Gα13 depletion. In **e**, two-way ANOVA with Dunnett's test against each dosage of

wild-type ADGRL2-CTF condition, n=3.

Fig. 4. Structure of the ADGRL2-Gα13 complex. a, Cryo-EM map of the ADGRL2-Gα13

complex in a lipid nanodisc. The ADGRL2 ICL3 protrudes from the lipid bilayer and is highlighted

with a dashed blue box. **b**, Comparison of the Gα13-coupled ADGRL2 and ADGRL3 structures

showing conformational differences in ICL3 and TM6 (highlighted by arrows). In contrast to that

in the ADGRL3 structure, the ICL3 of ADGRL2 is more ordered and shifts towards Gα13 to

interact with a loop region containing three glutamine (QQQ) residues (shown as sticks). **c**,

Sequence alignment of different Gα subtypes (including Gαs-short, Gαi, Gαq, Gα12 and Gα13)

focusing on the distinct feature of three glutamine residues (QQQ) within the loop connecting

helix 4 and β-strand 6 in Gα13. **d**, Effects of alanine substitution of the Gα13 QQQ triplet on the

basal activity of ADGRL2-CTF WT and the ICL2 mutant (F943A). **e**, Gene editing tracks and

efficiency of AAA substitution for QQQ sequences at the endogenous *GNA13* gene in bulk

primary human keratinocyte populations. **f**, mRNA expression of early differentiation markers

with GNA13 QQQ and AAA edited cells evaluated by qPCR. In **d**, two-way ANOVA with

Dunnett's test against each dosage of wild-type ADGRL2-CTF + wild-type Gα13 condition, n=3

or 4. In **f**, One-way ANOVA with Dunnett's test against control Gα13 QQQ cells, n = 3.

Figure 1 Using CRISPR-Flow and Perturb-Seq to identify a functional role for the GPCRs in epidermal hemomeostasis.

Figure 2 ADGRL2 is essential for normal epidermal differentiation

a UMAP Analysis of ADGRL2 mRNA

b ADGRL2 mRNA

c ADGRL2-CTF protein

d qPCR of Differentiation mRNAs with *ADGRL2* Depletion

e RNA-seq from *ADGRL2* Depletion

f Top Gene Ontology(GO) terms with *ADGRL2* Depletion

BP:Biological Process; HP:human phenotype ontology; REAC: REACTOME

g *ADGRL2* Depletion in Human Skin Organoids

h

Figure 3 ADGRL2 activates Gα13 to guide epidermal differentiation

a Full length ADGRL2

b ADGRL2-CTF

c PAR1-ADGRL2

d EK-ADGRL2

e Gα13 Dosage Dependent Activation

f Shared GO terms with *GNA13/ADGRL2* Depletion

BP: Biological Process; HP: human phenotype ontology; REAC: REACTOME

g *GNA13* Depletion in Human Skin Organoid

Figure 4 ADGRL2 ICL3 in Gα13 activation

a

ADGRL2 · Nanodisc · ICL3 · Gα₁₃ · Gβ · Gγ

b

ADGRL2 · ADGRL3 · TM5 · TM6 · TM7 · H8 · α5 · ICL3 · Q · Q · Q · Gβ · Gα13

c

Gα13 Unique QQQ Patch Binding Sites

	h4s6.08			h4s6.11	h4s6.13		S6.01		S6.04			s6h5.01
Gαss	S	G	D	G	R	H	-	Y	C	Y	P	H	F	T
Gαi1	K	D	T	K	E	-	-	-	I	Y	T	H	F	T
Gαq	N	P	D	S	D	K	-	I	I	Y	S	H	F	T
Gα12	R	R	N	R	S	-	K	P	L	F	H	H	F	T
Gα13	R	R	D	Q	Q	Q	K	P	L	Y	H	H	F	T

d

Gα13 Dosage Dependent Activation

- ■ CTF-WT + Gα13-WT
- ▼ CTF-WT + Gα13-AAA
- ● CTF-F943A + Gα13-AAA

ΔNet BRET2 vs plasmid dosage (1µg, 2µg, 4µg, 6µg, 8µg)

Activation

e

GNA13 QQQ
G A C C **CAG C CAG C CAG** A A G C C C
QQQ>QQQ CAGCAACAG>CAGCAgCAG
editing efficiency 61.8%

GNA13 AAA
G A C C **CCT CCT GCT** A A G C C C
QQQ>AAA CAGCAACAG>gctgctgct
64.4%

f

qPCR of Early Differentiation Markers for *GNA13* Edited Cells

	KRT1			KRT10			MAFB			S100A8			S100A9		
Rep	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3
GNA13 QQQ															
GNA13 AAA															
QQQ vs. AAA		**			***			*			**			***	

log₂FC

Figure S1

a

Categories	Gene numbers	gRNAs
GPCR	101	404
Keratinocytes control genes	22	88
Progenitor control genes	22	88
Non-target gRNAs	NA	40
Total	145	620

b Keratinocyte differentiation associated genes

IRF6	GRHL3	MAF	MCM5	PKM	DUT
ZNF750	POU2F3	MAFB	ITGB1	PFN1	FEN1
KLF4	JUNB	VDR	FERMT1	DNMT1	ARPC1B
CEBPA	PRDM1	TP63	CD151	YWHAZ	DDIT4
EHF	OVOL1	YAP1	EGFR	S100A8	
NOTCH1	TFAP2A	PRMT1	DNAJC9	MGST1	
NOTCH2	HOPX	CALML3	UBE2S	ITGAV	
RBPJ	MPZL3	NOTCH3	SFN	TYMS	

c gRNA reads distribution 90%/10%=2.03

d Overall gating strategy

Scatter and singlet filter

mCherry positive (gRNA infected) cells compared to non-infected cells

Among mCherry+ cells, KRT10 high and low cell populations between day 0 and day 3 keratinocytes

non gRNA infected cells / gRNA infected cells

Day 0 keratinocytes / Day 3 keratinocytes

e KRT10 enrichment

f target mRNA level in perturb-seq

g Perturb-seq heatmap

Progenitor makers

Differentiation makers

Expression in KO cells vs all targets (Z score)

ADGRL2

Perturbed targets

Figure S2

Figure S3

a

ADGRL2-full length | ADGRL2-CTF | PAR-ADGRL2 | EK-ADGRL2 | ADGRL2-CTF-Δ7TA

HA DAPI

30μm

b

ADGRL2-CTF

β-actin

CTF | Δ7TA

c

SRE-RF Reporter Assay

Normalized luciferase

ns

***

EV | CTF | Δ7TA

d

Gα proteins gRNA Enrichment in Perturb Pseudotime

IRF6, GNA11, GNAO1, GNA13, GNA12, GNA15, GNAQ, GNAL, GNAI1, GNAI3, GNAS, GNAI2, SNAI2

gRNA enrichment
-log10(P)
gRNA depletion

Pseudotime
Progenitors → Differentiated

e

GNA13 Knockdown Efficiency

Control si | GNA13 pool si

Gα13

β-actin

f

RNA-seq from *GNA13* Depletion

Density

Genes ↑ during differentiatrion | Genes ↓ during differentiatrion

GNA13

Genes upregulated during differentiatrion
Genes downregulated during differentiatrion
Genes unchanged during differentiatrion

−log10 FDR

log2 FC (GNA13 siRNA/control siRNA)

g

Donor 1 1 2 2
Replicate 1 2 1 2

KRT1, SPRR3, SPRR2C, SPRR2E, DSG1, HOPX, KRTDAP, SPRR1A, FLG, LCE3D, SPRR4, LCE3E, IL20, LIPN, CYP26B1, LCE3B, LORICRIN

log2FC(GNA13i/ctrl)

h

Top GO terms with *GNA13* Depletion

REAC HP BP

tissue development
epithelium development
anatomical structure development
epithelial cell differentiation
developmental process
cell differentiation
cellular developmental process
animal organ development
epidermis development
cell population proliferation
positive regulation of cellular process
skin development
regulation of cell population proliferation
multicellular organismal process
anatomical structure morphogenesis
Abnormal epidermal morphology
Epidermal acanthosis
Corneal erosion
Palmoplantar hyperkeratosis
Hyperkeratosis
Plantar hyperkeratosis
Palmar hyperkeratosis
Palmar hyperlinearity
Palmoplantar keratoderma
Hypergranulosis
Abnormality of the plantar skin of foot
Formation of the cornified envelope
Keratinization
Condensation of Prometaphase Chromosomes

−log10(p_value)

intersection_size
100
200
300

BP:Biological Process; HP:human phenotype ontology; REAC: REACTOME

i

***ADGRL2* and *GNA13* Depletion RNA-seq Overlap Hits**

Genes upregulated during differentiatrion
Genes downregulated during differentiatrion
Genes unchanged during differentiatrion

log2FC(GNA13 si/control si)

R²=0.16

j

3D Organoid IF Quantification

**
***

KRT10 | TGM1

control siRNA
GNA13 smartpool siRNA

Figure S4

Supplemental Figure Legends

Fig. S1. CRISPR-Flow and Perturb-seq GPCR knockout screens of epidermal GPCRs in

differentiating keratinocytes. a, Numbers of gene targets and sgRNAs in the CRISPR plasmid

library. **b**, Selected keratinocyte progenitor and differentiation genes. **c**, sgRNA count distribution

in plasmid library. **d**, CRISPR-Flow gating strategy. **e**, Quantitative flow cytometric analysis of

KRT10 expression changes from progenitors to differentiated cells. **f**, Perturb-seq mRNA overall

knockout efficiency of target sgRNAs populations compared to non-target sgRNA populations.

g, Heatmap visualization of Perturb-seq results; columns correspond to mRNAs while rows

indicate the effect of individual gene perturbation on progenitor and differentiation genes.

Fig. S2. ADGRL2 affects keratinocyte differentiation in 2D and organoid models. a,

Temporal heatmap illustrating GPCR mRNA expression during across days 0 to 6 of calcium-

induced differentiation of primary human keratinocytes in vitro. **b**, ADGRL2 knockdown

evaluation by Western blot. **c**, Heatmap of significant altered genes, FDR<0.05 with fold

change >2 or <0.5, upon ADGRL2 knockdown. **d**, Quantification of KRT10, TGM1, Ki67

immunofluorescent signal in ADGRL2 knockdown organoid models, unpaired t Test. **e**, Western

blot of the mosaic samples. **f**, Quantification of KRT10 signal in mosaic tissue control and L2

siRNAs. In **f**, paired t Test, n = 11.

Fig. S3. ADGRL2 activates Gα13 subtype to enable epidermal differentiation. a, Cellular

localization of HA tagged ADGRL2 full-length, ADGRL2-CTF, PAR1-ADGRL2, EK-ADGRL2,

ADGRL2-CTF-Δ7TA proteins in HEK293T cells evaluated by immunostaining. **b**, Western blot

of ADGRL2-CTF and ADGRL2-CTF-Δ7TA protein expression. **c**, SRE-RF luciferase reporter

measurements of ADGRL2-CTF and ADGRL2 TA peptide-deficient mutants. **d**, Gα proteins

Perturb-seq showing enrichment of knock-out cells along the pseudotime trajectory (Mann-

Whitney log10 P) compared to cells with safe target guides. A positive enrichment in low

pseudotime cells indicates nominates a gene as necessary for differentiation. **e**, GNA13

knockdown evaluation by Western blot. **f**, RNA-Seq analysis of GNA13 knockdown samples

sourced from two independent skin donors, showing all genes with log2 fold change >0.2 or <-

0.2, red/blue genes are the ones upregulated/downregulated during differentiation, white genes

are unchanged ones. **g**, Heatmap of significant altered genes, FDR<0.05 with fold change >2 or

<0.5, upon GNA13 knockdown. **h**, Gene Ontology analysis of all significantly altered mRNAs

(FDR<0.05) by GNA13 knockdown, based on RNA-seq data. **i**, Correlation between GNA13 and

ADGRL2 knockdown RNA-seq results. **j**, Quantification of KRT10, TGM1 immunofluorescent

signal in *GNA13* depleted regenerated human skin organoid tissue, unpaired t Test.

Fig. S4. Cryo-EM analysis of the ADGRL2-Gα13 complex in lipid nanodiscs. a, The receptor

and Gα13 constructs used in the study. Sequence corresponding to the CTF of ADGRL2 (TA

and 7TM) was inserted after a hemagglutinin signal peptide (HA) and a methionine residue. The

plasmid for expressing the miniGα13heterotrimer is the same as for the ADGRL3- Gα13

complex[37]. **b**, Size-exclusion chromatography (SEC) profile and SDS-PAGE of purified

miniGα13-coupled ADGRL2. **c**, Representative 2D class averages of the ADGRL2- Gα13

complex. **d**, Cryo-EM data processing workflow for the ADGRL2-Gα13 complex in lipid

nanodiscs. **e**, Angular distribution heat map of particles used for the global 3D reconstruction of

the ADGRL3-Gα13 complex. **f**, Gold-standard Fourier shell correlation (FSC) curves of the

locally refined receptor and miniGα13 reconstructions. **g**, Cryo-EM map of the ADGRL2-Gα13

complex in lipid nanodiscs showing the resolved density for two cholesterol molecules (colored

in blue). The first methionine residue (M) used in the receptor construct was also resolved. **h**,

Cryo-EM density and models are shown for TMs 1-7, the tethered agonist (TA) peptide and the

CLR molecules of the ADGRL2-Gα13 complex. **i**, Binding mode of the TA peptide, and the bent

TM6 and TM7 (highlighted by dashed lines) in the miniGα13-coupled ADGRL2. **j**, Interactions

between ICL2 and the miniGα13 α5 helix of ADGRL2. Key residues are shown as sticks. **k**,

Comparison of the basal Gα13 activity of ADGRL2-CTF WT and ICL2 mutants (F943A and

V942/F943A) by titrating different amounts of transfected plasmids by BRET2 assay. **l**, Cellular

localization of HA tagged ADGRL2-CTF-F943A and V942/F943A mutants in HEK293T cells

evaluated by immunostaining. **m**, Western blot of empty vector, ADGRL2-CTF, ADGRL2-CTF-

F943A and ADGRL2-CTF-V942/F943A in HEK293T cells. **n**, Unsharpened Cryo-EM map of the

ADGRL2-Gα13 complex showing that ICL3 of the receptor protrudes from the lipid bilayer to

interact with miniGα13. **o**, The model and EM density for the interface between ICL3 of the

receptor and the QQQ patch of Gα13. **p**, Structural comparison of ADGRL2 and ADGRL3 when

coupled with miniGα13, highlighting extracellular regions. **q**, Western blot of Gα13 wild-type and

AAA mutant in HEK293T cells.

Methods

Keratinocyte culture and differentiation

Keratinocytes were grown in a 1:1 ratio of Keratinocyte-SFM and Medium 154, supplemented with human recombinant epidermal growth factor (rEGF), bovine pituitary extract, and Human Keratinocyte Growth Supplement along with Penicillin-Streptomycin and Antibiotic-Antimycotic. Discarded surgically excised skin samples were incubated overnight at 4°C in a solution containing 50% dispase and 50% culture medium. The epidermis was then mechanically separated from the dermis, followed by trypsinization for 5-10 minutes at 37°C with intermittent shaking. Cellular debris and membrane remnants were removed using sterile forceps, and the cell suspension was centrifuged for 5 minutes at 1000 rpm. The resulting cell pellet was resuspended and cultured as passage 0 primary human keratinocytes. For differentiation assays, keratinocytes were seeded at densities of 1.3×10^6 cells/well in 6-well plates or 300,000 cells/well in 24-well plates. The medium was fully replaced the following morning with 50:50 medium supplemented with 1.2mM $CaCl_2$. Subsequent half-medium changes were carried out daily until harvesting.

Regenerated human skin organoid tissue

Devitalized skin samples were purchased from New York Firefighter Skin Bank and subjected to an initial rinse in Phosphate-Buffered Saline (PBS). The samples were then incubated in 1x PBS containing 2x Penicillin-Streptomycin and Antibiotic-Antimycotic at 37°C for 15 days. Following incubation, the epidermal layer was separated from the underlying dermis, which was then stored at 4°C. Dermal tissue was divided into 1.5 cm square pieces and oriented within custom-designed Annular Dermal Supports (ADS) with the basement membrane matte side facing upwards and the shiny side facing downwards. Upon semi-drying, the orientation was inverted, followed by a uniform application of Matrigel to the shiny side. After Matrigel solidification, the dermal tissue was flipped. Keratinocyte Growth Media (KGM) was carefully added to the ADS plate. Keratinocytes were seeded on the basement membrane side at densities between 0.5×10^6 and 1×10^6 cells using a 50 µL aliquot of KGM. Media were replaced at 1–2 day intervals.

Lentivirus generation

8×10^6 Lenti-X cells were seeded in a 10 cm^2 dish. On the following day, 5 µg of target plasmid,

5 µg of 8.91 plasmid, and 1.6 µg of pUC-MDG plasmid were transfected using Lipofectamine

3000. A complete media change was performed 6 hours or overnight post-transfection, and

lentivirus was harvested 48 hours post-transfection. The viral supernatant was mixed with one-

third volume of Lenti-X™ Concentrator at 1500 g for 45 minutes at 4°C. The supernatant was

carefully removed, and the viral pellet was resuspended in 50 µL of cold PBS. Aliquots were

prepared in small volumes and stored at -80°C for long-term use.

Plasmid library generation

The sgRNA libraries were derived from a whole-genome sgRNA list provided by the Brunello

Lab. Each gene was targeted by four unique sgRNAs. The F+E parent vector, a lentiviral

expression construct, incorporates a U6 promoter followed by the sgRNA sequence. Capture

Sequence 1 from 10x Genomics was integrated into the F+E plasmid downstream of the sgRNA

sequence.   The   sgRNA   sequences,   flanked   by   the   adaptors

ATCTTGTGGAAAGGACGAAACACCG-sgRNAs-GTTTAAGAGCTAAGCTGGAAACAG,   were

synthesized by Agilent. A 200 nmol oligo library was ligated to Esp3I-digested F+E plasmid using

Gibson Assembly Master Mix. Transformation was executed using Stellar Competent Cells.

Transformed bacteria were plated on agar plate, and library coverage was calculated in 1:5000

dilution. All bacterial clones were harvested using 3 mL LB broth, rinsed with another 3 mL of

LB broth, and then purified using the QIAGEN Plasmid Plus Maxi Kit.

Plasmid library quality control

For the first PCR (PCR1), 50 ng of plasmid library was amplified over 5 cycles using 2x

PrimeSTAR HS DNA polymerase and primers

PCR1_F: GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCGTAATACGGTTATCCACGCGG

and PCR1_R: ACACGACGCTCTTCCGATCTNNNNNNNNNTGTGGAAAGGACGAAACACC to

introduce a 9 bp degenerate sequence (N's) at the beginning of read1, serving as UMIs. The

PCR product was purified using the Zymo Clean and Concentrator-5 kit and further cleaned with

1.1x SPRIselect beads, followed by two 80% ethanol washes, 1-2 minutes of air-drying, and

elution in elution buffer. For the second PCR (PCR2), full Illumina sequencing adapters on read1

were added using i7 and i5 primers. SYBR green was included in PCR2 and run on a Stratagene

Mx3000P (Agilent) qPCR machine to monitor amplification. The PCR2 reaction was stopped
before reaching plateau (around 4-5 cycles post-exponential growth). The product was cleaned
using the Zymo Clean and Concentrator-5 kit and run on a 4% E-Gel™ EX Agarose Gel. The
330 bp library band was gel-purified, and its concentration was quantified using the KAPA Library
Quant Kit. The plasmid library pool was run on Miseq machine (Illumina, SY-410-1003) using
the MiSeq Reagent Kit v3 150-cycle kit.

CellTiter-Blue cell viability assay

Cells were seeded at a density of 100,000 cells/well in 6-well plates. Cell Titer Blue media was
prepared by combining Cell Titer Blue reagent with 5x diluted 50:50 media. Cells were incubated
in 1 mL of CTB media for 45 minutes at 37°C in the dark. Fluorescence was measured at 560
nm excitation and 590 nm emission for 200 μL triplicate aliquots per sample.

Perturb-seq single cell CRISPR screening and analysis

MOIs for Cas9 and sgRNA library lentiviruses were quantified using Lenti-X™ GoStix™ Plus.
Primary cultured keratinocytes were transduced with Cas9 lentivirus and sgRNA library lentivirus
(MOI ≤ 0.3). Media was replaced with 50:50 media 24 hours post-transduction, and drug
selection with blasticidin (5 μg/ml) and puromycin (1 μg/ml) commenced 48 hours post-
transduction, continuing for a minimum of 3 days. Subsequent to drug selection, cell viability was
assessed using a Cell Titer Blue assay. Cells were seeded for differentiation 7–10 days post-
transduction to optimize CRISPR cutting efficiency. Progenitor day 0 cells and differentiation day
3 cells were harvested using the 10x Genomics protocol for Single Cell Suspensions from
Cultured Cell Lines for Single Cell RNA Sequencing. Cell number and viability were verified, and
40,000 cells/well were loaded onto 10x Genomics Chromium Chips for single-cell gem formation.
For library generation, protocols for Chromium Single Cell 3' Reagent Kits v3 (RevA) and
Chromium Next GEM Single Cell 3' Reagent Kits v3.1 Dual Index (RevB) were followed. Library
quality and quantity were evaluated using Bioanalyzer's high-sensitivity DNA assay.
Concentrations of gene expression and feature barcode libraries were determined using the
KAPA Library Quant Kit. All prepared libraries were pooled and sequenced on a NovaSeq S4
chip.

Raw 10x data was processed using Cellranger (v.3 for the larger screens, v.5 for the smaller
screens) to produce h5 files, which were converted to Seurat objects. Count data was normalized
with SCTransform11. Cells were filtered to only those with a single sgRNA, less than 5%
mitochondrial reads and at least 200 RNAs. Dimensionality reductions were performed with the
RunPCA and RunUMAP Seurat functions, except for the pseudotime analysis. Pseudotime
analysis was performed by passing the SCT normalized expression data to the
reduce_dimensionality function of SCORPIUS, using 10 dimensions and Pearson distances,
then trajectories were inferred with SCORPIUS12. For the day 3 larger screen, the data was
subset to the 2,000 most variable features (using the Seurat function FindVariableFeatures)
before dimensionality reduction to reduce memory usage. To test for effects on pseudotime
scores, single-sided Mann-Whitney U tests were performed on cells with guides for a target gene
vs all other cells with a single guide. To analyze cell enrichment at specific points along the
pseudotime axis, a Gaussian kernel density estimate (KDE) was applied to the pseudotimes of
all cells, as well as to those cells targeted by guides specific to the gene of interest, using the
'scipy.stats.gaussian_kde' function. This KDE was then used to estimate the density of cell
counts across the pseudotime axis at 40 evenly spaced points. A window of length 10 was rolled
along the pseudotime space and one-sided Mann-Whitney U tests were applied to the density
estimates from cells with the gene of interest vs all cells.

CRISPR-Flow and analysis
Two biological replicates of primary human keratinocytes from separate donors were infected
with lentivirus containing the CRISPR knockout library. Differentiated and progenitor cells were
harvested on day 3 using trypsin for subsequent staining. Following detachment, cells were
washed with PBS and fixed with 1x Fix/Perm Buffer for 20 minutes at room temperature. Cells
were washed twice with 1x Perm Wash Buffer and stained with 1 µl antibody/million cells,
followed by additional washes and resuspension in 1x Cell Staining Buffer. Cells were filtered
through a polystyrene test tube with a cell strainer snap cap and sorted using a FACS machine.
The sorted cells were then centrifuged at 800g for 5 minutes, washed with 1x PBS, and either
stored at -80°C or processed immediately for genomic DNA extraction. To extract DNA, cells
were resuspended in lysis buffer (50 mM Tris-HCl, pH 8.1, 10 mM EDTA, 1% SDS) and then
heated to 65°C for 10 minutes with shaking to reverse formaldehyde cross-linking. RNaseA was

added and samples were incubated at 37°C for 30 minutes with shaking. After that, Proteinase
K (20 mg/mL) was added, and the samples were further incubated for 2 hours at 37°C, then at
95°C for 20 minutes. DNA purification was performed using SPRI beads, using 1.8x sample
volume of beads. PCR amplifications were carried out using PrimeStar in 100 µL reaction
volumes with 500 ng of genomic DNA and 20 pmol of each primer per reaction. The PCR
program was set to an initial denaturation at 98°C for 1 minute, followed by 5 cycles of 98°C for
10 seconds, 56°C for 10 seconds, and 72°C for 15 seconds, with a final extension at 72°C. A
Zymo PCR cleanup kit was used to combined PCR reactions. Subsequently, a SPRI cleanup
was performed to eliminate primers from the samples. This purified product underwent a second
round of PCR using PrimeStar, with the addition of 20 pmol of each primer (TruSeq adapters
with dual i5/i7 indexing) and 33x SYBR Green for real-time monitoring. The PCR conditions were
the same as the first, except the final 72°C elongation time for 1 minute. The reaction was
stopped at cycle before plateau based on the amplification trace. PCR product was then run on
a 2% agarose gel and gel purified for sequencing. Sequencing was conducted using a NovaSeq
S4 chip, targeting 10 reads per cell.

Raw data was processed with a custom snakemake pipeline that counts the number of UMIs
per guide in each cell pool. The number of UMIs per guide were input to MageCK RRA analysis,
pairing biological samples together ("paired" option) and using the safe targeting guides as
controls. Counts in the "high KRT10" group were compared against all the other divisions of that
cell population (unsorted, low and medium KRT10 groups), and the reverse for the "low KRT10"
group. When the p-values for the enrichment in high KRT10 vs depletion in low KRT10 (and vice
versa) are combined, they are only pseudo-p-values (ψP), since each sorting group is part of
the background set for the other and the P values are therefore not independent.

siRNA reverse transfection
All ON-TARGETplus siRNAs were sourced from Horizon Discovery. A mixture of 250 pmol
siRNAs and 25 µl Lipofectamine RNAiMAX was prepared in Opti-MEM and incubated for 25
minutes. Following incubation, 1×10^6 keratinocytes were seeded in a 10 cm^2 dish and treated
with the siRNA-lipid complex for 24-48 hours.

RNA-seq library generation and analysis

RNA was isolated using the RNeasy Plus Kit. For RNA-seq library preparation, 500 ng of RNA

was processed according to the QuantSeq 3' mRNA-seq Library Prep Kit FWD for Illumina

protocol. Library quality was confirmed using Bioanalyzer's high-sensitivity DNA assay. Library

concentrations were quantified using the KAPA Library Quant Kit). Libraries were sequenced on

a Novaseq S4 chip, targeting 20–30 million reads per library. Adaptors and polyA tails were

removed using BBDuk (v.39.01). The processed data were aligned to the hg38 human genome

(Ensembl release 99) using STAR (v.2.7.1a). Gene read counts were calculated using rsem-

calculate-expression (v.1.3.1) and assembled into a data matrix via rsem-generate-data-matrix.

Normalization and analysis of the RNA read count matrix were performed using the Bioconductor

R package DESeq2 with default settings. The Benjamini-Hochberg method was applied for

multiple testing correction to assess the false discovery rate (FDR). Gene Ontology (GO)

analysis was performed using the gProfileR R package[1] according to the authors' tutorial and

identified significant GO terms associated with the most differentially expressed genes.

Reporter assay

The SRF-RE reporter was obtained from Promega (E1350). LentiX (HEK293T) cells were plated

at a density of 40,000 cells per well in 24-well plates. We transfected 100 ng of SRF-RE-Luc

plasmid, 10 ng each of ADGRL2 full-length, ADGRL2-CTF, and ADGRL2-CTF-Δ7TA plasmids,

along with 10 ng of pRL-Renilla plasmid (Promega) using Lipofectamine 3000. Cells were lysed

24 hours after transfection using 1x Passive Lysis Buffer (Promega). Dual-luciferase activity was

measured using the Dual-Luciferase Reporter Assay System (Promega) on a Tecan Infinite

M1000 instrument. Data analysis included normalizing firefly luciferase activity to renilla

luciferase, expressed in Relative Luciferase Units (RLU), and adjusting for protein expression

levels.

BRET2 assay

HEK G-protein K.O cells[2] were plated on a 12-well plate at a density of 3–4x 10^5 cells in 1 mL

per well. HEK G K.O. media contained 1x DMEM plus 10% FBS, 1x Penicillin-Streptomycin with

1x MEM Non-essential Amino Acid (NeA) Solution. After 20–24 hours, cells were co-transfected

with 1 µg different ADGRL2 receptors or increasing amounts ADGRL2 receptors (1, 2, 4, 6, and

8 µg) for dose-dependent BRET and TRUPATH plasmids at 1:1:1 DNA ratio (Gα13-
RLuc8:Gβ3:Gγ9-GFP2) via TransIT-2020 (Mirus, MIR5400). Each condition required 97 µL of
room-temperature 1x Opti-MEM. 1 µL each DNA TRUPAPH plasmid at 1 µg/µL concentration),
8 µL of 1µg/µL of the receptor plasmid/empty vector, and 3 µL of room temperature and gently
vortexed TransIT-2020 reagent. The volume of the reactions was maintained consistent with an
addition of empty vector. The TransIT-2020:DNA complexes mixture were gently mixed via
pipetting 10 times and incubated at room temperature for 20 mins before adding drop-wise in
the well. The plate was rocked gently side to side and incubated at 37oC for 24 hours before
harvesting as follows: In each well, media was aspirated, and cells were washed with 1 mL warm
PBS. Cells were detached with 300 µL warm Versene and incubated at 37oC for 5 mins then
resuspended via pipetting 10 times. Cells were plated in a Matrigel-coated 96-well assay plate
at a density of 30–50,000 cells per well in 200 µL complete DMEM containing 1x NeA. Each
experimental condition was plated into three separate wells within the 96-well assay plate.
BRET2 assays were performed 48 hours after transfection as follows: In each well, media was
aspirated and cells were incubated in 90 µL of 1x Hanks' balanced Salt Solution with 20 mM
HEPES pH7.4 and 10 µL 100 µM Coelenterazine-400a diluted in PBS for 5 minutes. For the
Thrombin treated BRET assay, followed by adding 10 µL of vehicle solution, Thrombin at 100
U/mL concentration for a final concentration of 10 U/mL, or 10 µL of Thrombin vehicle, which is
0.1% BSA in PBS. For Enterokinase treated BRET assay, followed by adding 10 µL of 5.5 U of
Enterokinase in PBS or 10 µL of vehicle PBS. BRET intensities were measured via BERTHOLD
TriStar2 LB 942 Multimode Reader with Deep Blue C filter (410 nm) and GFP2 filter (515 nm).
The BRET ratio was obtained by calculating the ratio of GFP2 signal to Deep Blue C signal per
well. The BRET2 ratio of the three wells per condition were then averaged. Net BRET2 was
subsequently calculated by subtracting the BRET2 ratio of cells expressing no receptor from the
BRET2 ratio of each respective experimental condition[3].
274

Cryo-EM Constructs
The sequence of human ADGRL2 (isoform 2, Uniprot ID: O95490-2) comprising the TA peptide
and the 7TM domain was cloned into pFastBac1 with an N-terminal haemagglutinin (HA) signal
peptide followed by an additional methionine residue, and a C-terminal Strep tag. The construct

of the miniGα13 heterotrimer was generated as previously described[4]. A schematic description

of the constructs is provided in Figure S4A.

Protein expression

The human ADGRL2 and miniGα13 were co-expressed in Sf9 insect cells using the Bac-to-Bac

Baculovirus Expression system (Invitrogen). Cells were grown in serum-free ESF 921 medium

(Expression Systems) to a density of 3.5-4 million cells per ml and then co-infected with 1%

culture volume of P2 baculoviruses for ADGRL2 and miniGα13. Cells were collected by

centrifugation 48 h post infection and the pellets were snap-frozen in liquid nitrogen and stored

at -80°C until use.

Purification and reconstitution of the ADGRL2-Gα13 complex into lipid nanodiscs

Cell pellets were thawed and resuspended by Dounce homogenization in buffer containing 20

mM HEPES pH 7.5, 100 mM NaCl, 10 mM MgCl$_2$, 10% glycerol, protease inhibitors and 0.1 mM

TCEP. Following the addition of 25 mU/ml apyrase, the cell suspension was incubated at room

temperature for 2 h. Membranes were solubilized in 1% n-dodecyl-β-maltoside (DDM, Anatrace),

0.1% cholesteryl hemisuccinate (CHS, Anatrace), and incubated for 2 h at 4°C. The supernatant

was isolated by centrifugation at 100,000g for 40 min, and the solubilized proteins was incubated

with Ni-NTA affinity resin in the presence of 20 mM imidazole for 1 h at 4°C with gentle stirring.

The resin was packed into a gravity column and washed with 30 column volumes of buffer

containing 20 mM HEPES pH 7.5, 100 mM NaCl, 10% glycerol, 5 mM MgCl$_2$, 0.02% DDM, 0.002%

CHS and 20 mM imidazole. The complex was eluted in buffer containing 250 mM imidazole, and

then incubated with Strep-Tactin®XT 4Flow® resin (IBA) for overnight at 4 °C with gentle stirring.

The resin was collected and washed with buffer containing 20 mM HEPES pH 7.5, 100 mM NaCl,

10 mM MgCl$_2$, 5% glycerol, 0.02% DDM and 0.002% CHS. Protein was eluted from the column

using Strep-Tactin®XT elution buffer (IBA) plus 0.02% DDM and 0.002% CHS. Eluted protein

was concentrated and injected onto a Superose 6 column with buffer containing 20 mM HEPES

pH 7.5, 100 mM NaCl, 2 mM MgCl$_2$, 0.02% DDM and 0.002% CHS. The peak fractions were

pooled and concentrated. The lipid mixture (DOPC: DOPG at a molar ratio of 3:2) used for

reconstituting the complex into lipid nanodiscs was prepared as previously described[5]. The

purified complex was mixed with the lipid mixture and the membrane scaffold protein (MSP1D1,

Sigma) at molar ratio 1:150:1.5 and incubated for 1 h on ice. Bio-Beads SM2 (Bio-Rad) were

then added into the mixture (0.35 g of beads per ml) and incubated with gentle rocking for

overnight at 4°C. The reconstitution mixture was spun down to remove the Bio-Beads, and the

supernatant was applied onto a Superose 6 column with buffer containing 20 mM HEPES pH

7.5, 100 mM NaCl, 1 mM $MgCl_2$. The peak fractions containing the complex were collected and

concentrated for preparing Cryo-EM grids with parallel examination by negative stain EM[6].

Cryo-EM data collection and processing

The ADGRL2-Gα13 complex (3 µl at 5.5 mg ml^{-1}) was applied to freshly glow-discharged 300-

mesh R1.2/R1.3 UltrAuFoil holey gold grids (Quantifoil) under 100% humidity at 4°C. Excess

sample was blotted away, and the grids were subsequently plunge-frozen into liquid ethane

using a Vitrobot Mark IV (Thermo Fisher Scientific) and stored in liquid nitrogen. Frozen grids

were imaged at cryogenic temperatures with a Titan Krios G2 (Thermo Fisher Scientific)

Transmission Electron Microscope with a Selectris X post-column energy filter operated at 300

kV on a Falcon 4i direct electron camera at a pixel size of 0.75 Å. Micrographs were recorded

with defocus values ranging from -0.5 µm to -1.5 µm using EPU (Thermo Fisher Scientific).

Micrographs were recorded for 4 s with a total exposure dose of 50 electrons·Å^{-2}.

For detailed breakdown of data processing and exact particle numbers, see Fig. S4C-F. Briefly,

dose-fractionated image stacks were imported into RELION[7] and subjected to motion correction

with MotionCor2[8]. Contrast transfer function parameter estimation was performed with

CTFFIND-4.1[9]. Particle selection and extraction was performed with the Laplacian autopicker

function in RELION. The extracted particles were imported into CryoSPARC[10] and subjected to

multiple rounds of reference-free 2D classification, followed by iterative rounds of 3D ab initio

reconstruction with multiple classes and 3D heterogeneous refinement to remove particles from

poorly defined 3D classes. Selected particles were re-imported to RELION and then subjected

to 3D refinement in RELION followed by Bayesian polishing. Polished particles were then

imported back to CryoSPARC for nonuniform refinement, global CTF refinement, and local

refinement with a soft mask around 7TM or miniGα13.

Model building and refinement

For model building and refinement, composite maps were generated in Chimera[11] by merging
local refinement maps of 7TM and miniGα13 using the 'vop maximum' command. The ADGRL3–
Gα13 structure (PDB ID: 7SF7) was used as the initial model for docking into the Cryo-EM maps
in Chimera. The model was then subjected to iterative rounds of manual adjustment in Coot[12]
and real-space refinement in Phenix[13]. The model statistics were validated in MolProbity[14]. The
refinement statistics are provided in Table S1. ChimeraX[15] was used for figure preparation.

Western blot

Proteins were lysed in RAPI buffer with protease inhibitor and quantified using the BCA assay.
10-20µg of cell lysates were loaded per lane on an SDS-PAGE gel and transferred to a
nitrocellulose membrane at 4°C. The resulting membrane was blocked with LI-COR blocking
buffer (TBS) at room temperature for 1 hour. The membrane was then incubated with primary
antibody at 4°C overnight. Membranes were washed with TBST and incubated with secondary
goat anti-mouse and goat anti-rabbit antibodies (LI-COR Biosciences) at a dilution of 1:15,000
for 1 hour at room temperature. After that membranes were washed with TBST and visualized
and quantified using the Odyssey CLx Infrared Imaging System (LI-COR Biosciences).

Immunostaining

7 µm tissue sections were prepared with Cryostat HM525 NX. The organoids tissues were fixed
in cold methanol. Cultured cell were fixed with 4% paraformaldehyde. The samples were
incubated with primary antibodies at 4°C overnight, and following by the secondary antibodies
room temperature for 1 hour. Samples were mounted in Duolink In Situ Mounting Medium with
DAPI. All images were taken and processed using Zeiss fluorescent microscope with
ApoTome.2.

Homology-directed repair (HDR) genome editing

The donor ssAAV vector used as a template for HDR was constructed by cloning 1965bp GNA13
genomic DNA sequence flanking the 5' and 3' end of the target mutation into the AAV transfer
plasmid between AAV ITR sequences. Control vector was generated with a synonymous
substitution preserve the same codon QQQ. During genomic amplification, the AAA mutation
was engineered: QQQ AAA (CAGCAACAG/GCTGCTGCT) within exon 4, and control

synonymous substitution QQQ (CAGCAACAG/CAGCAGCAG) was generated in parallel.
Additionally, in order to prevent endonuclease mediated re-cutting, the following PAM mutations
were engineered, resulting in synonymous amino acid substitutions: C/T (YY) in exon 4. Primers
contained homology arms to the AAV transfer vector allowing In-Fusion assembly into NheI/EoRI
digested AAV plasmid. After confirmation of the insert sequence integrity, constructs were used
with helper vectors from Cell Biolabs (VKP-400-DJ) for AAV virus production in Hek293t cells.
The viruses were purified using AAVpro® Purification Kit Midi.
379
The guide sequences targeting GNA13 for CRISPR/Cas9 genome editing were predicted using
the Feng Zhang lab website (Ref genome GRCh38, CRISPRko, SpyoCas9, Hsu design) and
were ordered from IDT as sgRNA. The sgRNA: 5'-TGATAGCAGTGGTGAAGTGG-3' was used
in this study. For CRISPR/Cas9 genome editing, 73pmol of sgRNA was complexed with 61pmol
recombinant Alt-RspCas9 protein in 8.31μL of PBS for 15 min and immediately used for
nucleofection of $8x10^5$ primary keratinocytes with the Amaxa nucleofection apparatus (Lonza)
using program T-018. After recovery, cells were mixed with AAV virus containing either control
wild type QQQ or mutant donor QQQ AAA template at MOI $8x10^6$, transferred into 10cm dishes
and cultured for minimal 72 hrs. Genomic DNAs were isolated and evaluated for the editing
efficiency using PCR amplification and sanger sequencing of the bulk cell population.
For genomic DNA amplification, the following primers were used:
F: 5'-CCCATTGCTTTTAATAGAGGAGCA-3'
R: 5'-GGACTGGACAGGACAGCAAA-3'
393
394 **Quantitative RT-PCR analysis and primers**
395 Total RNA was extracted using RNeasy Plus and further subjected to reverse transcription using
396 the iScript cDNA Synthesis Kit. qRT-PCR analysis was performed using the LightCycler 480 II
397 System (Roche) with the SYBR Green Master Mix. All Ct values are normalized to the levels of
398 60S ribosomal protein L32. The qRT-PCR primers are listed below.

Gene	Forward primer	Reverse primer
ADGRL2	TGTGGATCCAGCATTACAACTGA	GCTGCTCTGCTGAAACCTTC
GNA13	TTCTTGTCAAGGGCGTAGGG	CTCTCGAGCATCAACCAGCA
SBSN	TCGGTCAACACGCCTTTCAT	TGATGTGACAACGGCGACAT
CALML5	CTACGAGGAGTTCGCGAGG	GTTTCCCATCCACCGACCAG

CDSN	CCTTGAGCTGCCATCAGTCAG	GTCTGAGAAGGTGCCAATGCT
HOPX	ACAGCCCAGGAAAATGAGCA	ACAGCCCAGGAAAATGAGCA
FLG	AAAGAGCTGAAGGAACTTCTGG	AACCATATCTGGGTCATCTGG
TGM1	CTTCAAGAACCCCCTTCCCG	TGAGGATCTTGGGCCTCTGT
LOR	CTCTGTCTGCGGCTACTCTG	CACGAGGTCTGAGTGACCTG
Keratin 1	GAAGTCTCGAGAAAGGGAGCA	ATGGGTTCTAGTGGAGGTATCTA
Keratin 10	GCAAATTGAGAGCCTGACTG	CAGTGGACACATTTCGAAGG
LCE3D	GCTGCTTCCTGAACCAC	GGGAACTCATGCATCAAG
SPINK5	TTAGCAAGAGCTCCCAAGGC	CCATCTGTGCCACAAACAGC
KRTDAP	CTACCGCAACAGAGGGCCTTA	CAGGAGCGGCACTTCTCAGT
S100A8	GTCCTCAGTTTGTGCAGAATATAAA	GCCAGAAGCTCTGCTACTCC
RPL32	AGGCATTGACAACAGGGTTC	GTTGCACATCAGCAGCACTT

Statistical analysis

401 Statistical tests were performed using GraphPad version 10.2.3. Sample sizes, statistical
402 analyses methods, significance values are reported in the figure panel or figure legends. For
403 statistical analyses, $P > 0.05$ were considered not significant (ns), and asterisks denote the
404 following: * $0.01 < p \leq 0.05$; ** $0.001 < p \leq 0.01$; *** $p \leq 0.001$. Error bars represent standard
405 error mean (SEM) unless otherwise indicated.

Reagent sources

REAGENT	SOURCE	IDENTIFIER
Antibodies		
anti-ADGRL2	Thermo Fisher Scientific	PA565359
anti-GNA13	proteintech	67188-1-Ig
anti-Keratin 10	Thermo Fisher Scientific	MA1-06319
anti-HA	CST	3724
anti-Transglutaminase-1	proteintech	12912-3-AP
anti-Ki67	Abcam	15580
anti-beta-actin	Millipore Sigma	A2228
Anti Cytokeratin 10 Antibody (KRT10/844), Alexa Fluor™ 647	Novus Biologicals	NBP2-47825AF647
Cell Lines		
Human primary keratinocytes	Stanford University School of Medicine	Genomic data sharing (GDS)
LentiX 293T	Takara Bio	632180
GKO HEK293	Kind gift from Asuka Inoue, Tokyo University	Alvarez-Curto et al.,[16] J Biol Chem. Grundmann et al.,[17] Nat Commun.
Reagents and Materials		

DMEM	Gibco	11995-065
Keratinocyte-SFM	Gibco	17005-142
Medium 154	Gibco	M-154-500
HKGS	Thermo Fisher Scientific	S0015
rEGF	Gibco	10450-013
BPE	Gibco	13028-014
Antibiotic-Antimycotic	Thermo Fisher Scientific	15240-062
Penicillin-Streptomycin	Thermo Fisher Scientific	15140122
Dermis	New York Firefighter Skin Bank	N/A
Lipofectamine 3000	Invitrogen	L3000015
Lipofectamine RNAiMAX	Invitrogen	13778150
Lenti-X concentrator	Takara Bio	631231
RNeasy Plus Kit	QIAGEN	74136
iScript cDNA Synthesis Kit	Bio-Rad	1708891
SYBR Green Master Mix	Thermo Fisher Scientific	K0223
QuantSeq 3′ mRNA-seq Library Prep Kit FWD for Illumina	Lexogen	015.96
LDS sample loading buffer	Invitrogen	NP0008
Opti-MEM	Thermo Fisher Scientific	31985062
Dispase	Corning	354235
Antibiotic-Antimycotic	Thermo Fisher Scientific	15240-062
Stellar Competent Cells	Takara Bio	636763
Thrombin	Sigma	T4846
Enterokinase, light chain	NEB	P8070S
SPRIselect beads	Beckman Coulter	B23317
RIPA Lysis Buffer	Thermo Fisher Scientific	89900
cOmplete™, Mini Protease Inhibitor Cocktail	Roche	11836153001
Fetal Bovine Serum	Gibco	16000044
Gibson Assembly Master Mix	NEB	E2611S
4% E-Gel™ EX Agarose Gel	Thermo Fisher Scientific	G401004
KAPA Library Quant Kit	Kapa Biosystems	KK4854
AAVpro® Purification Kit Midi	Takara Bio	6675
AAV-DJ helper free packaging system	Cell Biolabs	VPK-400-DJ
MiSeq® Reagent Kit v3 150-cycle kit	Illumina	MS-102-3001
cell strainer snap cap	VWR	352235
RNAseA	Qiagen	19101
Passive Lysis Buffer	Promega	E1910
Proteinase K	Thermo Fisher Scientific	AM2546
PrimeStar	Takara	R045A

Dual-Luciferase Reporter Assay System	Promega	E1910
Non-essential Amino Acid Solution	Sigma	M7145
Versene	Gibco	15040066
Hanks' balanced Salt Solution	Gibco	14175095
Coelenterazine-400a	NanoLight Technologies	340
Thrombin	Sigma	T4648
Enterokinase	New England Biolabs	P8070S
BCA assay	Thermo Fisher Scientific	23225
4% paraformaldehyde	Alfa Aesar	43368
DAPI	Sigma	DUO82040-5ML
Alt-RspCas9 protein	IDT	10008100
Recombinant DNA		
pCMV-dR8.91	Addgene	Romain Zufferey et al.,[18] Nat. Biotechnol.
pMD2.G	Addgene	Romain Zufferey et al.,[18] Nat. Biotechnol.
TRUPATH kit	Addgene	1000000163
SRF-RE reporter plasmid	Promega	E1350
pRL-Renilla plasmid	Promega	E2261
pLEX CMV Cas9 blast	This study	N/A
F+E 10x cs1 sgRNA mCherry	This study	N/A
Plex Flag-Keratin14	This study	N/A
Plex HA-Keratin14	This study	N/A
Plex full-length ADGRL2-FHH	This study	N/A
Plex ADGRL2-CTF-FHH	This study	N/A
Plex PAR1-ADGRL2-CTF-FHH	This study	N/A
Plex SNAP-Flag-ADGRL2-CTF-FHH	This study	N/A
pEB PAR-Lphn3*tv7*	Gift from Richard Sando's lab	Duy Lan Huong Bui et al.,[3] Cell Rep.
Plex ADGRL2-CTF-F943A-FHH	This study	N/A
Plex ADGRL2-CTF-V943A/F943A-FHH	This study	N/A
Plex full-length rat ADGRL2-FHH	This study	N/A
Plex full-length rat ADGRL2 F943A-FHH	This study	N/A
Plex Gα13-BRET	This study	N/A
Plex Gα13-AAA-BRET	This study	N/A

409 **Data availability**

410 All processed and raw sequencing data have been deposited in Gene Expression Omnibus

411 (GEO), under the accession code: GSE281860. The Cryo-EM density maps and corresponding

coordinates have been deposited in the Electron Microscopy Data Bank (EMDB) and the Protein

Data Bank (PDB), respectively, under the following accession codes: EMD-47522 and 9E51.

Reference

1. Kolberg L, Raudvere U, Kuzmin I, Vilo J, Peterson H. gprofiler2 -- an R package for gene list functional enrichment analysis and namespace conversion toolset g:Profiler. *F1000Res* **9**, (2020).

2. Grundmann M*, et al.* Lack of beta-arrestin signaling in the absence of active G proteins. *Nature Communications* **9**, (2018).

3. Bui DLH*, et al.* The adhesion GPCRs CELSR1-3 and LPHN3 engage G proteins via distinct activation mechanisms. *Cell Rep* **42**, 112552 (2023).

4. Barros-Alvarez X*, et al.* The tethered peptide activation mechanism of adhesion GPCRs. *Nature* **604**, 757-762 (2022).

5. He F*, et al.* Allosteric modulation and G-protein selectivity of the Ca(2+)-sensing receptor. *Nature* **626**, 1141-1148 (2024).

6. Peisley A, Skiniotis G. 2D Projection Analysis of GPCR Complexes by Negative Stain Electron Microscopy. *Methods Mol Biol* **1335**, 29-38 (2015).

7. Zivanov J*, et al.* New tools for automated high-resolution cryo-EM structure determination in RELION-3. *Elife* **7**, (2018).

8. Zheng SQ, Palovcak E, Armache JP, Verba KA, Cheng Y, Agard DA. MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017).

9. Rohou A, Grigorieff N. CTFFIND4: Fast and accurate defocus estimation from electron micrographs. *J Struct Biol* **192**, 216-221 (2015).

10. Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).

11. Pettersen EF*, et al.* UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* **25**, 1605-1612 (2004).

12. Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501 (2010).

454 13.   Liebschner D*, et al.* Macromolecular structure determination using X-rays, neutrons and
455   electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861-877
456   (2019).
457
458 14.   Chen VB*, et al.* MolProbity: all-atom structure validation for macromolecular
459   crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12-21 (2010).
460
461 15.   Pettersen EF*, et al.* UCSF ChimeraX: Structure visualization for researchers, educators,
462   and developers. *Protein Sci* **30**, 70-82 (2021).
463
464 16.   Alvarez-Curto E*, et al.* Targeted Elimination of G Proteins and Arrestins Defines Their
465   Specific Contributions to Both Intensity and Duration of G Protein-coupled Receptor
466   Signaling. *J Biol Chem* **291**, 27147-27159 (2016).
467
468 17.   Grundmann M*, et al.* Lack of beta-arrestin signaling in the absence of active G proteins.
469   *Nat Commun* **9**, 341 (2018).
470
471 18.   Zufferey R, Nagy D, Mandel RJ, Naldini L, Trono D. Multiply attenuated lentiviral vector
472   achieves efficient gene delivery in vivo. *Nat Biotechnol* **15**, 871-875 (1997).
473
474

Acknowledgments

476 We thank T.C. Südhof, B.K. Kobilka, A.E. Oro, and H.Y. Chang for expert project advice. This
477 work was supported by a USVA Merit Review grant BX001409 to P.A.K. and by NIAMS/NIH
478 grants AR049737 to P.A.K. We thank Duy Thanh Nguyen, Ron Shanderson, Weili Miao, Ian
479 Ferguson, Mårten Winge, Jordan Meyers, Jerry Qu, Joy Ji for the help and consultation during
480 the project.
481

Author Contributions

X.Y. and P.A.K. conceived the project. F.H. and B.S. performed Cryo-EM, data collection and
analysis, D.F.P., R.M.M., L.D. performed and supported for data analysis. K.G., D.L.H.B.
performed TRUPATH assay. D.R., A.H. assisted with dual screening process. Z.S. assisted with
HDR experiment. S.M. performed the luciferease assay. V.L.-P assisted the organoid study. K.L.,
T.S., J.Y.Q assisted with cell culture and cloning experiments. Y.X., F.H., G.S., and P.A.K.
guided overall experiments and data analysis. X.Y. and P.A.K. wrote the manuscript with input
from all authors.

Competing interest

492 The authors declare no competing interests.