The Adhesion GPCR ADGRL2 engages Ga13 to Enable Epidermal Differentiation

Xue Yang[1,2,10], Feng He[3,10], Douglas F. Porter[1], Krassimira Garbett[4], Robin M. Meyers[1,5], David L. Reynolds[1], Duy Lan Huong Bui[4], Audrey Hong[1], Luca Ducoli[1], Zurab Siprashvili[1], Vanessa Lopez-Pajares[1], Smarajit Mondal[1], Lisa Ko[1], Yuqing Jing[6], Shiying Tao[1], Bharti Singal[3], Richard Sando[4], Georgios Skiniotis[3,7,8]*, Paul A. Khavari[1,2,9,11]*

[1] Program in Epithelial Biology, Stanford University School of Medicine, Stanford, CA, USA.
[2] Program in Cancer Biology, Stanford University, Stanford, CA, USA.
[3] Department of Molecular and Cellular Physiology, Stanford University School of Medicine, Stanford, CA, USA
[4] Department of Pharmacology, Vanderbilt Brain Institute, Vanderbilt University, Nashville, TN 37240, USA
[5] Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA
[6] Departments of Pediatrics and Genetics, Stanford University, Stanford, CA, USA
[7] Department of Structural Biology, Stanford University School of Medicine, Stanford, CA, USA
[8] Present address: Center of Excellence for Structural Cell Biology, Department of Structural Biology, St Jude Children's Research Hospital, Memphis, TN 38120, USA
[9] Veterans Affairs Palo Alto Healthcare System, Palo Alto, CA, USA.
[10] These authors contributed equally.
[11] Lead Contact.
* Correspondence: khavari@stanford.edu and yiorgo@stanford.edu

Abstract

Homeostasis relies on signaling networks controlled by cell membrane receptors. Although G-protein-coupled receptors (GPCRs) are the largest family of transmembrane receptors, their specific roles in the epidermis are not fully understood. Dual CRISPR-Flow and single cell Perturb-seq knockout screens of all epidermal GPCRs were thus performed, uncovering an essential requirement for adhesion GPCR ADGRL2 (latrophilin 2) in epidermal differentiation. Among potential downstream guanine nucleotide-binding G proteins, ADGRL2 selectively activated $G\alpha13$. Perturb-seq of epidermal G proteins and follow-up tissue knockouts verified that $G\alpha13$ is also required for epidermal differentiation. A cryo-electron microscopy (cryo-EM) structure in lipid nanodiscs showed that ADGRL2 engages with $G\alpha13$ at multiple interfaces, including via a novel interaction between ADGRL2 intracellular loop 3 (ICL3) and a $G\alpha13$-specific QQQ glutamine triplet sequence in its GTPase domain. In situ gene mutation of this interface sequence impaired epidermal differentiation, highlighting an essential new role for an ADGRL2-$G\alpha13$ axis in epidermal differentiation.

Introduction

The epidermis is a self-renewing surface tissue composed primarily of specialized epithelial keratinocytes that receive signals via a variety of cell surface receptors. The balance between growth and differentiation in the epidermis represents a prototype for tissue homeostasis. Dysregulation of epidermal homeostasis contributes to numerous human clinical disorders, including cancer[1], psoriasis[2] and chronic wounds[3]. GPCRs comprise the largest family of transmembrane receptors in mammals[4, 5]. Several studies have addressed the role of specific GPCRs in epidermis. For example, activating mutations in the *SMO* GPCR gene are pathogenic in basal cell carcinoma[6] and the SMO inhibitor, Vismodegib, prevents cancer progression[7]. Other examples include the Frizzled GPCR, which regulates keratinocyte proliferation[8], as well as LPAR GPCRs that enable epidermal differentiation[9]. The roles of the many other GPCRs expressed in epidermis, however, are largely uncharacterized.

Traditional pooled CRISPR screening methods are of limited utility in characterizing the actions of gene family members in epidermal homeostasis. Standard cell survival and proliferation phenotypes within bulk cell populations can fail to detect genes with essential roles in biomedically important cell transitions, such as occurs in epithelial differentiation[10]. Several CRISPR screening approaches may overcome this limitation. For example, fluorescence sorting-based pooled CRISPR screening (**CRISPR-Flow**), separates cells based on expression of marker proteins representative of specific cell states[11, 12], such as KRT10 for epidermal differentiation. Single cell Perturb RNA-sequencing (**Perturb-seq**) is another method that quantifies a subset of the transcriptome in cells undergoing a variety of genetic manipulations[13, 14]. Integrating these two methods may enable systematic evaluation of the roles of members of diverse gene families in the control of homeostatic gene expression programs.

Here we performed CRISPR-Flow and Perturb-seq knockout screens of all GPCRs expressed in epidermis. These screens validated previously reported GPCR roles while also identifying a new essential requirement for the ADGRL2 adhesion GPCR (**aGPCR**) in epidermal differentiation. ADGRL2 expression was strongly upregulated during keratinocyte differentiation. In epidermal tissue, ADGRL2 was required for induction of a key differentiation genes involved in barrier formation. Among potential downstream G proteins, ADGRL2 selectively activated $G\alpha 13$ ($G\alpha 13/G\beta/G\gamma$). Perturb-seq screening of genes encoding $G\alpha$ subunits expressed in

epidermis confirmed an essential requirement for Gα13 in epidermal differentiation. Cryo-
electron microscopy (**Cryo-EM**) of nanodisc embedded ADGRL2-Gα13 complexes facilitated
high-resolution structure determination. Structural analysis revealed an ADGRL2 ICL3 loop
interface with a QQQ sequence on Gα13, which appears to be a unique GPCR-G protein
interaction that may help explain the preferential engagement of ADGRL2 with Gα13. Mutation
of the Gα13 QQQ sequence diminished ADGRL2 binding and impaired epidermal differentiation,
underscoring the functional importance of this interface in newly uncovered pro-differentiation
roles for the ADGRL2 GPCR and its Gα13 protein target.

Results

CRISPR-Flow and Perturb-seq of epidermal GPCRs.

Analysis of dynamic epidermal differentiation RNA-seq[15] identified 101 GPCRs expressed in epidermal keratinocytes. These span the five known GPCR sub-families (**Fig. 1a**). A CRISPR library targeting these GPCRs was next generated. sgRNAs were included to target known positive controls essential for epidermal homeostasis as well as non-targeting negative controls, encompassing 620 sgRNAs in total (**Extended Data Fig. S1a-b, Supplementary Table 1**). Quality control assessments confirmed minimal skew and full coverage of the resulting plasmid library (**Extended Data Fig. 1c**), which was designed to support both CRISPR-Flow and Perturb-seq screening assays. The CRISPR-Flow screen employed flow cytometry to isolate populations expressing high and low levels of the KRT10 epidermal differentiation marker for sgRNA enrichment/depletion analysis to nominate pro-differentiation and pro-progenitor GPCRs while the single-cell Perturb-seq screen evaluated GPCR gene knockout effects on keratinocyte progenitor and differentiation mRNA expression signatures (**Fig. 1b**). The complementary nature of these screens, one based on the expression of a single canonical early differentiation protein representing entry into the differentiation pathway and the other on the levels of a panel of keratinocyte mRNAs, was designed to catalog GPCRs with non-redundant impacts on homeostatic gene expression in epidermal keratinocytes.

Flow cytometry benchmarking for the CRISPR-Flow effort demonstrated that the KRT10-high population increased to 13.5% by day 3 of calcium-induced differentiation in vitro compared to <0.05% in day 0 progenitors (**Extended Data Fig. 1d-e**). Known positive controls required for epidermal cell differentiation, such as *IRF6, NOTCH2, PRDM1, MAF, CEBPA* and *ZNF750,* as well as the *LPAR3* GPCR, were scored by CRISPR-Flow as required for KRT10 induction. Knockout of known progenitor genes, such as *ITGB1*, *FERMT1*, *TP63*, *EGFR*, and *YAP1*, in contrast, led to increased KRT10 protein levels. These findings confirmed prior work and validated the fidelity of the screen. Among GPCRs, ADGRL2 was nominated as the strongest novel pro-differentiation GPCR (**Fig. 1c-d, Supplementary Table 2**). CRISPR-Flow screening thus validated positive controls with known impacts on homeostatic gene expression in epidermal cells while also nominating specific GPCRs as new regulators of this process.

Single-cell RNA-seq of differentiating keratinocyte populations for Perturb-seq benchmarking
revealed a trajectory from progenitor-like cells to more differentiated states (**Fig. 1e**), consistent
with previous studies[16]. Representative differentiation genes, such as *KRT10*, *SBSN* and
KRTDAP, as well as undifferentiated progenitor population markers, such as *ITGB1*, *ITGA6* and
KRT14, exhibited expected expression dynamics across pseudotime (**Fig. 1f**), supporting
scRNA-seq dataset quality. As an additional quality check for CRISPR-Cas9-mediated gene
knockout effectiveness, expression of targeted genes was confirmed to be significantly reduced
compared to non-targeted controls (**Extended Data Fig. 1f**). Perturb-seq assigned known pro-
differentiation genes, *IRF6, NOTCH2/3, PRDM1, TFAP2A*, and *GRHL3*, as well as pro-
progenitor genes, *ITGB1*, *FERMT1*, *TP63*, and *YAP1* to their known roles, supporting the fidelity
of the screen (**Fig. 1g-h**). As in CRISPR-Flow, Perturb-seq identified specific LPAR genes as
pro-differentiation GPCRs, including *LPAR3, LPAR5 and LPAR6* (**Fig. 1g-h**), which are known
to be activated by Lysophosphatidic Acid (**LPA**) and which have been previously implicated in
keratinocyte differentiation[9]. ADGRL2 was again captured as a novel pro-differentiation GPCR
in Perturb-seq data (**Fig. 1g-h, Extended Data Fig. 1g, Supplementary Table 3**). Correlation
of the CRISPR-Flow and Perturb-seq dual screens identified ADGRL2 and LPAR3 among the
top GPCR candidates promoting differentiation (**Fig. 1i**). Integration of Perturb-seq and CRISPR-
Flow screens therefore nominated ADGRL2 as a novel pro-differentiation GPCR.
105

ADGRL2 is essential for epidermal differentiation
In keratinocyte differentiation RNA-seq datasets[15], mRNA levels of ADGRL2 were significantly
upregulated during differentiation (**Extended Data Fig. 2a, red labeled**). Consistent with this,
uniform manifold approximation and projection (**UMAP**) analysis performed on the Perturb-seq
dataset showed that ADGRL2 mRNA levels were low in progenitors but markedly elevated in
differentiated cells, concordant with *KRTDAP* differentiation gene expression and inversely
correlated to the *MKI67* proliferation marker (**Fig. 2a**). Further evaluation of ADGRL2 mRNA
levels showed significant upregulation during differentiation by qPCR analysis (**Fig. 2b**), along
with accumulation of the self-cleaved dual protein species C-terminal fragment (**CTF**) of
ADGRL2 (**Fig. 2c**). Both ADGRL2 mRNA and protein levels were therefore significantly
increased during differentiation, further suggesting a possible role for ADGRL2 in this process.
117

In vitro, keratinocyte ADGRL2 depletion was associated with reduced differentiation mRNA
levels as measured by both qPCR (**Fig. 2d, Extended Data Fig. 2b)** and RNA-seq (**Fig. 2e,**
Extended Data Fig. 2c, Supplementary Table 4). Gene Ontology (**GO**) analysis of RNA-seq
data showed that ADGRL2 depletion led to impaired induction of epidermal differentiation genes
in patterns linked to human clinical phenotypes characterized by abnormalities in this process
(**Fig. 2f**). Similarly, in 3-dimensional human skin organoid tissue regenerated from diploid human
keratinocytes treated with siRNAs to ADGRL2 or scrambled control, ADGRL2 depletion led to
decreases in differentiation proteins such as KRT10 and TGM1 in suprabasal epidermis, without
affecting proliferation markers Ki67 (**Fig. 2g, Extended Data Fig. 2d**). To investigate whether
ADGRL2 influences keratinocyte differentiation in a cell autonomous or non-cell autonomous
fashion, mosaic epidermal tissue was generated by mixing HA-KRT14 labeled ADGRL2
knockdown cells with Flag-KRT14 labeled control knockdown cells at 1:1 ratios. ADGRL2-
depleted cells displayed reduced KRT10 differentiation protein expression, in contrast to
adjacent control cells (**Fig. 2h, Extended Data fig. 2e-f)**. These findings nominate ADGRL2 as
a cell-autonomous pro-differentiation GPCR in epidermis.

ADGRL2 activates Gα13

GPCRs interact with Gα subunits of heterotrimeric G proteins to induce downstream signaling.
To identify specific Gα subunits activated by ADGRL2 during keratinocyte differentiation, a
TRUPATH assay[17] was employed. This approach uses the BRET2 sensor to measure the
attenuation of luminescence upon downstream G protein activation[17]. To avoid interference by
endogenous G proteins, a pan G protein protein-knockout (**GKO**) HEK293 cell line[18, 19] was used.
Exogenously expressed ADGRL3 co-transfected with Gα13/Gβ3/Gγ9 served as a positive
control and ADGRL3 co-expressed with GαoB/Gβ3/Gγ8 functioned as a negative control[20]. 12
Gα proteins are expressed across keratinocyte differentiation, as detected by RNA-seq[15]. A
panel of 12 putative G$\alpha\beta\gamma$ "epidermal transducerome" combinations was therefore evaluated in
cells expressing either the full-length ADGRL2 or the ADGRL2 CTF, both of which correctly
localized to the cell membrane (**Extended Data Fig. 3a**). Gα13 and Gα12 subunits displayed
the highest basal activation levels by full-length ADGRL2 and ADGRL2-CTF (**Fig. 3a-b**). The
Tethered Agonist (**TA**) peptide serves as an intrinsic ligand in aGPCRs; once exposed by
proteolytic cleavage, it activates the receptor to initiate intracellular signal transduction[21]. To
further dissect the role of the TA peptide in cleavage-dependent activation, therefore, two

independent systems, PAR1-ADGRL2 and EK-ADGRL2, were employed. In the PAR1-ADGRL2

system, the N-terminal extracellular portion of ADGRL2 was replaced with the N-terminal domain

of protease-activated receptor PAR1, creating a TA-protected variant[22]. Thrombin treatment

cleaves off the PAR1 N terminus, exposing a "scarred" TA peptide with a two amino acids

deletion[22]. Localization of the PAR1-ADGRL2 hybrid protein to the cell membrane was not

abolished due to these modifications (**Extended Data Fig. 3a**). Notably, despite generating a

non-native "scarred" TA peptide[20, 22], the receptor still predominantly activated Gα13 upon

cleavage (**Fig. 3c**). To further study this, an EK-ADGRL2 fusion was also generated. In this

approach, enterokinase-mediated cleavage selectively cleaves off the SNAP tag on the N

terminus of the protein and exposed a native TA peptide[20, 23]. Like the PAR1-ADGRL2 fusion,

localization of the EK-ADGRL2 hybrid protein remained unchanged compared to wild-type full

length ADGRL2 (**Extended Data Fig. 3a**). Upon enterokinase release of the TA peptide, the

induced receptor selectively activated Gα13, reaffirming the preferential activation of Gα13 by

ADGRL2 (**Fig. 3d**), and congruent with a finding reported by Pederick et al[24]. Gα13 activation

by these two TA peptide cleavage mimetics of ADGRL2 are consistent with the full length

ADGRL2 and ADGRL2-CTF results. As a negative control, we assessed the differential dose

responses of ADGRL2-CTF and its TA peptide deletion mutant (**Δ7TA**). This mutant has no

capacity to activate G proteins[25]. We also observed that the Δ7TA mutant failed to activate Gα13

in a dose-dependent fashion (**Fig. 3e**), despite similar expression and subcellular localization as

ADGRL2-CTF (**Extended Data Fig. 3a-b**). Further confirmation was obtained using a Gα13

reporter SRE-RF luciferase assay, which indicated that wild-type ADGRL2-CTF displayed an

enhanced ability to activate Gα13, while the Δ7TA mutant did not (**Extended Data Fig. 3c**).

These data suggest that Gα13 serves as the primary Gα protein subtype activated by ADGRL2.

Gα13 is essential for epidermal tissue differentiation

To investigate if Gα13 may act in a manner consistent with a role as a downstream effector of

ADGRL2 in keratinocytes, an epidermal G-protein Perturb-seq screening experiment was

conducted. The Perturb-seq data confirmed that ablation of *GNA13*, the gene encoding Gα13,

attenuated keratinocyte differentiation (**Extended Data Fig. 3d**). This observation was further

corroborated in 2D culture, where *GNA13* depletion was associated with a significant decrease

in expression of 299 differentiation-enriched genes that are also modulated by ADGRL2 (**Fig. 3f,**

Extended Data Fig. 3e-i, Supplementary Table 5). These findings were then further validated

in regenerated human skin organoid tissue where, similar to *ADGRL2* loss, *GNA13* depletion

reduced differentiation marker expression (**Fig. 3g, Extended Data Fig. 3j**). Gα13, like ADGRL2,

is therefore required for normal induction of epidermal differentiation genes.

Cryo-EM structure of the ADGRL2-Gα13 complex in lipid nanodiscs

To examine the structural basis of Gα13 activation by ADGRL2, human ADGRL2-CTF and a

miniGα13 heterotrimer were co-expressed in insect cells and reconstituted into lipid nanodiscs

for Cryo-EM analysis (**Extended Data Fig. 4a-b**). The Cryo-EM map of the ADGRL2- Gα13

complex was determined at a global nominal resolution of 3.1 Å, with local refinements yielding

further improved maps for the 7TM domain (2.9 Å) and G protein (2.9 Å) (**Extended Data Fig.**

4c-f, Supplementary Table 6). While previously reported aGPCR-G protein structures were

determined in detergent micelles[21, 26, 27, 28, 29, 30], this ADGRL2-Gα13 complex structure provides

an opportunity to examine how an aGPCR couples to G proteins in a lipid bilayer environment.

Of note, two cholesterol (**CLR**)-like molecules were observed bound to the receptor in the Cryo-

EM map, occupying hydrophobic cavities in between TM3 and TM4, and TM4 and TM5,

respectively (**Extended Data Fig. 4g-h**). From a global perspective, the structure of the

ADGRL2-Gα13 complex in lipid nanodiscs is not significantly different from the structure of the

ADGRL3-Gα13 complex in detergent micelles[26] and shows a deeply buried TA peptide in the

orthostatic 7TM binding site, bent TM6 and TM7 for G protein binding, and extensive interactions

between ICL2 and the α5 helix (**Extended Data Fig. 4i-j**). Mutations of key ICL2 residues, such

as the F943A and V942A/F943A mutants, dramatically reduced basal Gα13 activation by the

ADGRL2-CTF without affecting protein localization and stability (**Extended Data Fig. 4k-m**),

highlighting the broadly important role of ICL2 in the tethered agonist activation mechanism of

aGPCRs[30].

ADGRL2 ICL3 in Gα13 activation

In contrast to the previously described Gα13-coupled ADGRL3 structure[26], however, the Gα13-

coupled ADGRL2 structure in lipid nanodiscs revealed a distinct conformation of ICL3, which

protrudes from the lipid bilayer and contacts the GTPase domain of Gα13 (**Fig. 4a, Extended**

Data Fig. 4n-o). Additionally, subtle conformational differences were observed in the

extracellular regions (**Extended Data Fig. 4p**). Unexpectedly, and in contrast to ADGRL3, the

ICL3 of ADGRL2 appears to recognize three glutamine (**QQQ**) residues in the loop between

helix 4 and β-strand 6 of Gα13 (**Fig. 4b**), a region we observe relatively weak EM densities

suggesting the dynamic character of those elements (**Extended Data Fig. 4o**). The interactions

in this region likely facilitate Gα13 coupling by ordering ICL3 and promoting the outward

movement of TM6, which is positioned differently than in the ADGRL3 structure (**Fig. 4b**). Indeed,

3D variability analysis, as implemented previously[31], of the global Cryo-EM map revealed that

ICL3 of the receptor and this Gα13 loop flex together (**Supplementary Movie 1**). Notably, the

QQQ patch of Gα13 is unique across all Gα proteins, and is not even present in its close

subfamily member, Gα12, in spite of the fact that the surrounding sequences are conserved

between them (**Fig. 4c**). This uniqueness suggests the potential importance for these Gα13

residues in determining the ADGRL2 coupling specificity.

To evaluate the importance of the novel ICL3-Gα13 interaction interface, a Gα13 mutant (QQQ

to AAA) was generated and its basal activation by ADGRL2-CTF quantified. The AAA mutant

exhibited similar expression levels to wild-type Gα13 but showed significantly reduced activity

when activated by the wild-type ADGRL2-CTF (**Fig. 4d**). Moreover, the ICL2 mutant ADGRL2-

CTF-F943A could partially activate wild-type Gα13 (**Fig. 4d**) but failed to appreciably activate

the Gα13 AAA mutant (**Fig. 4d**), consistent with the premise that the ICL3-Gα13 interface

contributes to Gα13 activation by ADGRL2. To examine the functional impact of disrupting the

Gα13 QQQ interface with ADGRL2 on differentiation, the endogenous *GNA13* gene was edited

in normal human keratinocytes to substitute AAA for QQQ residues in Gα13. Control editing was

performed at the *GNA13* gene locus to alter codon sequences but preserve the QQQ residues

to generate two isogenic keratinocyte populations genetically identical except for this sequence

(**Fig. 4e**). AAA-edited keratinocytes displayed impaired differentiation gene induction while

control-edited cells showed no such defects (**Fig. 4f**). These findings suggest that intactness of

the newly identified ADGRL2 ICL3-Gα13 interface is required for normal epidermal

differentiation.

Discussion

The results of the CRISPR-Flow and Perturb-seq GPCR screens demonstrated a degree of concordance, not only in identifying a novel role for ADGRL2 but also in detecting LPAR GPCRs with previously reported actions in epidermal differentiation. Some differences, however, were observed between the two screens in the candidate GPCRs identified, likely due to differences in their design. The CRISPR-Flow screen relied on differential sorting based on levels of a single canonical differentiation protein, KRT10, whereas the single-cell RNA-seq approach used in Perturb-seq assessed broader transcriptomic outcomes that were then placed within a single cell differentiation trajectory. In addition to ADGRL2, each knockout screen nominated specific GPCRs for future study as candidate regulators of epidermal homeostasis. Specific known and novel epidermal GPCRs, including S1PR5, GPR157, ADRA2C, GPRC5B, PTGER2, ACKR3, ADRB1, and FZD6, were top hits in CRISPR-Flow. Additional known and novel epidermal GPCRs, such as F2RL1, GPR39, BDKRB2, and GPR153 scored highly in Perturb-seq. Several GPCRs, such as S1PR5 and EDNRA, were assigned contrasting functions between the screens, underscoring the need for single GPCR-focused follow-up functional studies, as performed here for ADGRL2. Combined analysis of both screening methods identified more known and novel GPCRs than either screen alone, suggesting that integrating both methods may be useful in future efforts to identify new regulators of complex genetic programs.

The role of the third intracellular loop in GPCR signaling has been elusive, largely due to its high sequence variability and structural flexibility[32, 33, 34]. While previous structural studies of aGPCR/G protein complexes typically relied on detergent purification methods and primarily highlighted interactions involving intracellular loop 2 (ICL2) of aGPCRs[21, 26, 27, 28, 29, 30], work here utilized lipid nanodiscs[35, 36], which incorporate lipid bilayers that may help recapitulate the native environment of transmembrane proteins. This may have helped to delineate the binding interface involving both ICL2 as well as ICL3 of ADGRL2 with Gα13. The present Cryo-EM structure-defined nanodisc embedded complex reveals that the ADGRL2 ICL3 engages a unique and functionally important QQQ-containing loop sequence in the GTPase domain of Gα13, an interaction which has never been observed in previous aGPCR structural studies. The QQQ site of Gα13 is unique across all Gα proteins, and is not present in its close subfamily member, Gα12, in spite of the fact that the surrounding sequences are conserved between them. Despite the

noticeable conformational flexibility observed in the EM map, the ICL3 of ADGRL2 is positioned
to bind Gα13's glutamine residues, whose mutation attenuated receptor signaling and
differentiation. In agreement with a recent study showing that ICL3 modulates G protein
selectivity of the β2 adrenergic receptor[37], these findings suggest a regulatory role for the
ADGRL2 ICL3 in tuning receptor signaling specificity. The current data represent, to our
knowledge, the first reported instance of an adhesion GPCR ICL3 engaging with a Gα protein
in a structured fashion, and may thus enable future potential efforts designed to modulate
signaling by ADGRL2-Gα13 as well as other aGPCRs and their corresponding Gα subunit
effectors.
280
ADGRL1, ADGRL2, and ADGRL3 reside within the latrophilin subfamily of aGPCRs[38], which
have well characterized roles in synaptic function and maintenance[39, 40]. While ADGRL1 and
ADGRL3 are predominantly expressed in the brain[41, 42], ADGRL2 is expressed more widely[41, 42],
suggesting it plays roles in other tissues. Actions of ADGRL2 in stratified epithelia, however,
have been largely unexplored. Interestingly, both ADGRL2 mRNA and protein levels were
significantly increased during differentiation. ADGRL2 knockdown impaired induction of a host
of genes essential for epidermal differentiation, including a number whose mutation leads to
human clinical disorders of this process. Among these genes are *TGM1, CDSN, LOR, SPINK5*,
and *FLG*, each of whose mutation causes monogenic skin diseases characterized by abnormal
epidermal barrier function, such as autosomal recessive congenital ichthyosis[43], peeling skin
disease[44], keratoderma hereditaria mutilans[45], ichthyosis linearis circumflexa[46], and ichthyosis
vulgaris[47] respectively. The ADGRL2 GPCR is therefore necessary for the induction of
differentiation genes essential for formation of a healthy cutaneous barrier.

Reference

1. Ratushny V, Gober MD, Hick R, Ridky TW, Seykora JT. From keratinocyte to cancer: the pathogenesis and modeling of cutaneous squamous cell carcinoma. *J Clin Invest* **122**, 464-472 (2012).

2. Zhang X, Yin M, Zhang LJ. Keratin 6, 16 and 17-Critical Barrier Alarmin Molecules in Skin Wounds and Psoriasis. *Cells* **8**, (2019).

3. Pastar I*, et al.* Epithelialization in Wound Healing: A Comprehensive Review. *Adv Wound Care (New Rochelle)* **3**, 445-464 (2014).

4. Alhosaini K, Azhar A, Alonazi A, Al-Zoghaibi F. GPCRs: The most promiscuous druggable receptor of the mankind. *Saudi Pharm J* **29**, 539-551 (2021).

5. Weis WI, Kobilka BK. The Molecular Basis of G Protein–Coupled Receptor Activation. *Annual Review of Biochemistry* **87**, 897-919 (2018).

6. Gutzmer R, Solomon JA. Hedgehog Pathway Inhibition for the Treatment of Basal Cell Carcinoma. *Target Oncol* **14**, 253-267 (2019).

7. Sekulic A*, et al.* Efficacy and safety of vismodegib in advanced basal-cell carcinoma. *N Engl J Med* **366**, 2171-2179 (2012).

8. Ghahramani A, Donati G, Luscombe NM, Watt FM. Epidermal Wnt signalling regulates transcriptome heterogeneity and proliferative fate in neighbouring cells. *Genome Biol* **19**, 3 (2018).

9. Sumitomo A*, et al.* LPA Induces Keratinocyte Differentiation and Promotes Skin Barrier Function through the LPAR1/LPAR5-RHO-ROCK-SRF Axis. *J Invest Dermatol* **139**, 1010-1022 (2019).

10. Chen S*, et al.* Genome-wide CRISPR screen in a mouse model of tumor growth and metastasis. *Cell* **160**, 1246-1260 (2015).

11. Condon KJ*, et al.* Genome-wide CRISPR screens reveal multitiered mechanisms through which mTORC1 senses mitochondrial dysfunction. *Proc Natl Acad Sci U S A* **118**, (2021).

12. Fomicheva M, Macara IG. Genome-wide CRISPR screen identifies noncanonical NF-kappaB signaling as a regulator of density-dependent proliferation. *Elife* **9**, (2020).

13. Adamson B*, et al.* A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response. *Cell* **167**, 1867-1882 e1821 (2016).

338 14. Dixit A*, et al.* Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell RNA
339   Profiling of Pooled Genetic Screens. *Cell* **167**, 1853-1866 e1817 (2016).
340

341 15. Kim DS*, et al.* The dynamic, combinatorial cis-regulatory lexicon of epidermal
342   differentiation. *Nat Genet* **53**, 1564-1576 (2021).
343

344 16. Cockburn K*, et al.* Gradual differentiation uncoupled from cell cycle exit generates
345   heterogeneity in the epidermal stem cell layer. *Nat Cell Biol* **24**, 1692-1700 (2022).
346

347 17. Olsen RHJ*, et al.* TRUPATH, an open-source biosensor platform for interrogating the
348   GPCR transducerome. *Nat Chem Biol* **16**, 841-849 (2020).
349

350 18. Alvarez-Curto E*, et al.* Targeted Elimination of G Proteins and Arrestins Defines Their
351   Specific Contributions to Both Intensity and Duration of G Protein-coupled Receptor
352   Signaling. *J Biol Chem* **291**, 27147-27159 (2016).
353

354 19. Grundmann M*, et al.* Lack of beta-arrestin signaling in the absence of active G proteins.
355   *Nature Communications* **9**, (2018).
356

357 20. Bui DLH*, et al.* The adhesion GPCRs CELSR1-3 and LPHN3 engage G proteins via distinct
358   activation mechanisms. *Cell Rep* **42**, 112552 (2023).
359

360 21. Xiao P*, et al.* Tethered peptide activation mechanism of the adhesion GPCRs ADGRG2
361   and ADGRG4. *Nature* **604**, 771-778 (2022).
362

363 22. Mathiasen S*, et al.* G12/13 is activated by acute tethered agonist exposure in the
364   adhesion GPCR ADGRL3. *Nature Chemical Biology* **16**, 1343-1350 (2020).
365

366 23. Perry-Hauser NA, VanDyck MW, Lee KH, Shi L, Javitch JA. Disentangling autoproteolytic
367   cleavage from tethered agonist-dependent activation of the adhesion receptor ADGRL3.
368   *J Biol Chem* **298**, 102594 (2022).
369

370 24. Pederick DT, Perry-Hauser NA, Meng H, He Z, Javitch JA, Luo L. Context-dependent
371   requirement of G protein coupling for Latrophilin-2 in target selection of hippocampal
372   axons. *Elife* **12**, (2023).
373

374 25. Stoveken HM, Hajduczok AG, Xu L, Tall GG. Adhesion G protein-coupled receptors are
375   activated by exposure of a cryptic tethered agonist. *Proc Natl Acad Sci U S A* **112**, 6194-
376   6199 (2015).
377

378 26. Barros-Alvarez X*, et al.* The tethered peptide activation mechanism of adhesion GPCRs.
379   *Nature* **604**, 757-762 (2022).
380

381 27. Ping YQ*, et al.* Structural basis for the tethered peptide activation of adhesion GPCRs.
382   *Nature* **604**, 763-770 (2022).
383

384 28. Qu X, *et al.* Structural basis of tethered agonism of the adhesion GPCRs ADGRD1 and
385   ADGRF1. *Nature* **604**, 779-785 (2022).
386

387 29. Qian Y, *et al.* Structural insights into adhesion GPCR ADGRL3 activation and G(q), G(s),
388   G(i), and G(12) coupling. *Mol Cell* **82**, 4340-4352 e4346 (2022).
389

390 30. Jones DTD, Dates AN, Rawson SD, Burruss MM, Lipper CH, Blacklow SC. Tethered agonist
391   activated ADGRF1 structure and signalling analysis reveal basis for G protein coupling.
392   *Nat Commun* **14**, 2490 (2023).
393

394 31. Papasergi-Scott MM, *et al.* Time-resolved cryo-EM of G-protein activation by a GPCR.
395   *Nature* **629**, 1182-1191 (2024).
396

397 32. Probst WC, Snyder LA, Schuster DI, Brosius J, Sealfon SC. Sequence alignment of the G-
398   protein coupled receptor superfamily. *DNA Cell Biol* **11**, 1-20 (1992).
399

400 33. Katritch V, Cherezov V, Stevens RC. Diversity and modularity of G protein-coupled
401   receptor structures. *Trends Pharmacol Sci* **33**, 17-27 (2012).
402

403 34. Venkatakrishnan AJ, Flock T, Prado DE, Oates ME, Gough J, Madan Babu M. Structured
404   and disordered facets of the GPCR fold. *Curr Opin Struct Biol* **27**, 129-137 (2014).
405

406 35. Denisov IG, Sligar SG. Nanodiscs for structural and functional studies of membrane
407   proteins. *Nat Struct Mol Biol* **23**, 481-486 (2016).
408

409 36. He F, *et al.* Allosteric modulation and G-protein selectivity of the Ca(2+)-sensing
410   receptor. *Nature* **626**, 1141-1148 (2024).
411

412 37. Sadler F, Ma N, Ritt M, Sharma Y, Vaidehi N, Sivaramakrishnan S. Autoregulation of
413   GPCR signalling through the third intracellular loop. *Nature* **615**, 734-741 (2023).
414

415 38. Purcell RH, Hall RA. Adhesion G Protein-Coupled Receptors as Drug Targets. *Annu Rev*
416   *Pharmacol Toxicol* **58**, 429-449 (2018).
417

418 39. Sando R, Sudhof TC. Latrophilin GPCR signaling mediates synapse formation. *Elife* **10**,
419   (2021).
420

421 40. Sando R, Jiang X, Sudhof TC. Latrophilin GPCRs direct synapse specificity by coincident
422   binding of FLRTs and teneurins. *Science* **363**, (2019).
423

424 41. Krishnan A, Nijmeijer S, de Graaf C, Schioth HB. Classification, Nomenclature, and
425   Structural Aspects of Adhesion GPCRs. *Handb Exp Pharmacol* **234**, 15-41 (2016).
426

427 42. Regan SL, Williams MT, Vorhees CV. Latrophilin-3 disruption: Effects on brain and
428   behavior. *Neurosci Biobehav Rev* **127**, 619-629 (2021).
429

430   43.   Herman ML*, et al.* Transglutaminase-1 gene mutations in autosomal recessive
431   congenital ichthyosis: summary of mutations (including 23 novel) and modeling of
432   TGase-1. *Hum Mutat* **30**, 537-547 (2009).
433

434   44.   van der Velden J, van Geel M, Engelhart JJ, Jonkman MF, Steijlen PM. Mutations in the
435   CDSN gene cause peeling skin disease and hypotrichosis simplex of the scalp. *J Dermatol*
436   **47**, 3-7 (2020).
437

438   45.   Pohler E*, et al.* Novel autosomal dominant mutation in loricrin presenting as prominent
439   ichthyosis. *Br J Dermatol* **173**, 1291-1294 (2015).
440

441   46.   Guerra L*, et al.* Ichthyosis Linearis Circumflexa as the Only Clinical Manifestation of
442   Netherton Syndrome. *Acta Derm Venereol* **95**, 720-724 (2015).
443

444   47.   Thyssen JP, Godoy-Gijon E, Elias PM. Ichthyosis vulgaris: the filaggrin mutation disease.
445   *Br J Dermatol* **168**, 1155-1166 (2013).
446
447

Figure Legends

Fig. 1. CRISPR-Flow and Perturb-seq screens in differentiating normal human
keratinocytes. a, Pie chart of the 101 epidermal GPCRs and their classes. **b**, Schematic of
experimental workflow for CRISPR screens. **c**, Scatterplot mapping guide enrichment (beta
score) in low vs. high KRT10 cells from CRISPR-Flow screening. **d**, Bar graphs quantifying mean
beta scores from top CRISPR-Flow identified candidates (red=GPCRs, black=controls). **e**,
Principal Component Analysis (PCA) plot representing variance in gene expression between
progenitor and differentiated epidermal states at day 3. **f**, Pseudotime trajectory analysis of
progenitor markers ITGB1/KRT14/ITGA6 and differentiation markers KRTDAP/SBSN/KRT10 at
day 3. **g**, Perturb-seq GPCR and positive control target effects plotted as pseudotime differences
from non-target guides (Z score) *vs* Mann-Whitney p values for the comparison to cells with
other guides. **h**, Enrichment of knock-out cells along the pseudotime trajectory (Mann-Whitney
log10 P) compared to cells with safe target guides. A positive enrichment in low pseudotime
cells indicates nominates a gene as necessary for differentiation. **i**, Correlation analysis of
CRISPR-Flow beta scores plotted against Perturb-seq pseudotime measurements for each
CRISPR target. ADGRL2 is highlighted in red. For all analyses *=p<0.05, **=p<0.01,
***=p<0.001.

Fig. 2. Requirement for ADGRL2 in epidermal differentiation. a, UMAP analysis depicting
ADGRL2 expression in progenitor and differentiated states in scRNA-seq data. **b**, ADGRL2
mRNA levels across keratinocyte differentiation in vitro. **c**, ADGRL2 protein levels (C-terminal
fragment dual bands) across differentiation. **d**, mRNA expression by qPCR of differentiation
markers and progenitor markers with control versus ADGRL2 knockdown (L2 si1 and L2 si2)
from four independent human skin donors evaluated. **e**, RNA-Seq analysis of ADGRL2
knockdown keratinocytes sourced from three unrelated skin donors, showing all genes with log2
fold change >0.2 or <-0.2, red/blue genes are the ones upregulated/downregulated during
differentiation, white genes are unchanged ones. **f**, Gene Ontology analysis of all significantly
altered mRNAs (FDR<0.05) following ADGRL2 knockdown based on the RNA-seq data. **g**,
Immunostaining of KRT10, TGM1 and Ki67 in regenerated human skin organoid tissue with
ADGRL2 depletion versus control. **h**, Immunostaining of KRT10 in mosaic skin organoids. In **b**

and **c**, one-way ANOVA with Dunnett's test against day 0 cells, n = 3. In **d**, one-way ANOVA

with Dunnett's test against control cells for each gene, n = 4.

Fig. 3. ADGRL2 activates Gα13. (**a, b**) Domain structures, the net BRET2 signals and

summarized BRET2 heatmap from full-length ADGRL2 and ADGRL2-CTF. (**c, d**) Domain

structures with drug treatment conditions, the induced BRET2 signals and summarized BRET2

heatmap from PAR1-ADGRL2 and EK-ADGRL2. **e**, Dose-dependent BRET2 signaling profiles

of ADGRL2-CTF and ADGRL2 TA peptide deficient mutants (Δ7TA) on Gα13. **f**, Gene Ontology

analysis of 299 significant altered mRNAs (FDR<0.1) shared between *GNA13* and *ADGRL2*

knockdown RNA-seq datasets. **g**, Immunostaining for KRT10 and TGM1 in human skin organoid

tissues with Gα13 depletion. In **e**, two-way ANOVA with Dunnett's test against each dosage of

wild-type ADGRL2-CTF condition, n=3.

Fig. 4. Structure of the ADGRL2-Gα13 complex. a, Cryo-EM map of the ADGRL2-Gα13

complex in a lipid nanodisc. The ADGRL2 ICL3 protrudes from the lipid bilayer and is highlighted

with a dashed blue box. **b**, Comparison of the Gα13-coupled ADGRL2 and ADGRL3 structures

showing conformational differences in ICL3 and TM6 (highlighted by arrows). In contrast to that

in the ADGRL3 structure, the ICL3 of ADGRL2 is more ordered and shifts towards Gα13 to

interact with a loop region containing three glutamine (QQQ) residues (shown as sticks). **c**,

Sequence alignment of different Gα subtypes (including Gαs-short, Gαi, Gαq, Gα12 and Gα13)

focusing on the distinct feature of three glutamine residues (QQQ) within the loop connecting

helix 4 and β-strand 6 in Gα13. **d**, Effects of alanine substitution of the Gα13 QQQ triplet on the

basal activity of ADGRL2-CTF WT and the ICL2 mutant (F943A). **e**, Gene editing tracks and

efficiency of AAA substitution for QQQ sequences at the endogenous *GNA13* gene in bulk

primary human keratinocyte populations. **f**, mRNA expression of early differentiation markers

with GNA13 QQQ and AAA edited cells evaluated by qPCR. In **d**, two-way ANOVA with

Dunnett's test against each dosage of wild-type ADGRL2-CTF + wild-type Gα13 condition, n=3

or 4. In **f**, One-way ANOVA with Dunnett's test against control Gα13 QQQ cells, n = 3.

Figure 1 Using CRISPR-Flow and Perturb-Seq to identify a functional role for the GPCRs in epidermal hemomeostasis.

Figure 2 ADGRL2 is essential for normal epidermal differentiation

a UMAP Analysis of ADGRL2 mRNA
(MKI67, KRTDAP, ADGRL2; Day 0 Progenitor, Day 3 Differentiation)

b ADGRL2 mRNA
Normalized ADGRL2 mRNA; day0, day3, day6; Differentiation; **, ***

c ADGRL2-CTF protein
Normalized ADGRL2-CTF protein; day0, day3, day6; *; ADGRL2-CTF; β-actin

d qPCR of Differentiation mRNAs with *ADGRL2* Depletion

Differentiation Markers — ADGRL2, SBSN, CDSN, S100A8, HOPX, FLG, TGM1, LOR, KRT1, LCE3D, SPINK5
Progenitor Markers — ANCK, CDK1, CCNB1, DNMT1, FN1, TFAP2B

	ADGRL2	SBSN	CDSN	S100A8	HOPX	FLG	TGM1	LOR	KRT1	LCE3D	SPINK5	ANCK	CDK1	CCNB1	DNMT1	FN1	TFAP2B
L2si1 vs. Ctrlsi	***	*	***	*	ns	*	**	*	**	**	***	ns	ns	ns	ns	ns	ns
L2si2 vs. Ctrlsi	***	**	***	ns	**	*	**	**	***	**	***	ns	ns	ns	ns	ns	ns

(Foreskin donor 1 2 3 4; Control si; L2 si1; L2 si2) log₂FC

e RNA-seq from *ADGRL2* Depletion
Density; Genes ↑ during differentiation; Genes ↓ during differentiation; -log10 FDR; ADGRL2; log2 (L2 siRNA/control siRNA); Genes upregulated during differentiation; Genes downregulated during differentiation; Genes unchanged during differentiation

f Top Gene Ontology(GO) terms with *ADGRL2* Depletion
(skin development, epidermal cell differentiation, keratinization, keratinocyte differentiation, epidermis development, epithelial cell differentiation, anatomical structure development, developmental process, tissue development, cell differentiation, cellular developmental process, animal organ development, multicellular organismal process, epithelium development, positive regulation of biological process, Abnormal epidermal morphology, Epidermal acanthosis, Hyperkeratosis, Congenital ichthyosiform erythroderma, Parakeratosis, Congenital nonbullous ichthyosiform erythroderma, Palmar hyperkeratosis, Thickened skin, Epidermal thickening, Alopecia, Palmoplantar hyperkeratosis, Abnormality of the plantar skin of foot, Orthokeratosis, Plantar hyperkeratosis, Palmoplantar keratoderma, Formation of the cornified envelope, Keratinization, Melanin biosynthesis)
REAC HP BP; BP:Biological Process; HP:human phenotype ontology; REAC: REACTOME; -log₁₀(P-value); intersection_size 50 100 150

g *ADGRL2* Depletion in Human Skin Organoids
Control siRNA; ADGRL2 siRNA1; ADGRL2 siRNA2; KRT10 DAPI; TGM1 DAPI; Ki67 DAPI; 40μm

h 3xFlag-K14; scramble si; Primary KCs; HA-K14; ADGRL2 si1; Primary KCs; Mosaic skin organoid; 1:1 mix
Control siRNA; Control i : L2 i = 1:1; ADGRL2 siRNA2; KRT10 HA DAPI; Rep1; Rep2; 50μm

Figure 3 ADGRL2 activates Gα13 to guide epidermal differentiation

Figure 4 ADGRL2 ICL3 in Gα13 activation

a ADGRL2, Nanodisc, ICL3, Gα13, Gβ, Gγ

b ADGRL2, ADGRL3, TM5, TM6, TM7, H8, α5, ICL3, Q, Q, Q, Gα13, Gβ

c Gα13 Unique QQQ Patch Binding Sites

	h4s6.08			h4s6.11		h4s6.13		S6.01			S6.04		s6h5.01
Gαss	S	G	D	G	R	H	-	Y	C	Y	P	H	F	T
Gαi1	K	D	T	K	E	-	-	-	I	Y	T	H	F	T
Gαq	N	P	D	S	D	K	-	I	I	Y	S	H	F	T
Gα12	R	R	N	R	S	-	K	P	L	F	H	H	F	T
Gα13	R	R	D	**Q**	**Q**	**Q**	K	P	L	Y	H	H	F	T

d Gα13 Dosage Dependent Activation

- CTF-WT + Gα13-WT
- CTF-WT + Gα13-AAA
- CTF-F943A + Gα13-AAA

ΔNet BRET2 vs plasmid dosage (1µg, 2µg, 4µg, 6µg, 8µg)

Activation

e

GNA13 QQQ: G A C C A G C A G C A G A A G C C C
QQQ>QQQ CAGCAACAG>CAGCAgCAG
editing efficiency 61.8%

GNA13 AAA: G A C C C T C C T G C T A A G C C C
QQQ>AAA CAGCAACAG>gctgctgct
64.4%

f qPCR of Early Differentiation Markers for *GNA13* Edited Cells

	KRT1			KRT10			MAFB			S100A8			S100A9		
Rep	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3
GNA13 QQQ															
GNA13 AAA															
QQQ vs. AAA	**			***			*			**			***		

log₂FC

Figure 1S

a

Categories	Gene numbers	gRNAs
GPCR	101	404
Keratinocytes control genes	22	88
Progenitor control genes	22	88
Non-target gRNAs	NA	40
Total	145	620

b

Keratinocyte differentiation associated genes

IRF6	GRHL3	MAF	MCM5	PKM	DUT
ZNF750	POU2F3	MAFB	ITGB1	PFN1	FEN1
KLF4	JUNB	VDR	FERMT1	DNMT1	ARPC1B
CEBPA	PRDM1	TP63	CD151	YWHAZ	DDIT4
EHF	OVOL1	YAP1	EGFR	S100A8	
NOTCH1	TFAP2A	PRMT1	DNAJC9	MGST1	
NOTCH2	HOPX	CALML3	UBE2S	ITGAV	
RBPJ	MPZL3	NOTCH3	SFN	TYMS	

c gRNA reads distribution 90%/10%=2.03

d Overall gating strategy

e KRT10 enrichment

f target mRNA level in perturb-seq

g Perturb-seq heatmap

ADGRL2

Figure 2S

b ADGRL2 Knockdown Efficiency

Control si   si1   si2
— ADGRL2-CTF
— β-actin

d 3D Organoid IF Quantification

- control siRNA
- ADGRL2 siRNA1
- ADGRL2 siRNA2

e Mosaic Western

Control si   ADGRL2 si2
Flag
HA
ADGRL2-CTF
β-actin

f Mosaic Tissue KRT10 Quantification

Figure 3S

a ADGRL2-full length | ADGRL2-CTF | PAR-ADGRL2 | EK-ADGRL2 | ADGRL2-CTF-Δ7TA

HA DAPI

30μm

b ADGRL2-CTF, β-actin (CTF, Δ7TA)

c SRE-RF Reporter Assay — Normalized luciferase — EV, CTF, Δ7TA — ns, ***

d Gα proteins gRNA Enrichment in Perturb Pseudotime

IRF6, GNA11, GNAO1, GNA13, GNA12, GNA15, GNAQ, GNAL, GNAI1, GNAI3, GNAS, GNAI2, SNAI2

Pseudotime
Progenitors → Differentiated

gRNA enrichment 3/2; -log10(P); gRNA depletion -2/-3

e GNA13 Knockdown Efficiency

Control si GNA13 pool si

Gα13, β-actin

f RNA-seq from *GNA13* Depletion

Density

Genes ↑ during differentiatrion | Genes ↓ during differentiatrion

Genes upregulated during differentiatrion
Genes downregulated during differentiatrion
Genes unchanged during differentiatrion

GNA13

-log10 FDR

log2 FC (GNA13 siRNA/control siRNA)

g Donor / Replicate: 1 1 2 2 / 1 2 1 2

KRT1, SPRR3, SPRR2C, SPRR2E, DSG1, HOPX, KRTDAP, SPRR1A, FLG, LCE3D, SPRR4, LCE3E, IL20, LIPN, CYP26B1, LCE3B, LORICRIN

log2FC(GNA13i/ctri)

h Top GO terms with *GNA13* Depletion

REAC HP BP

tissue development
epithelium development
anatomical structure development
epithelial cell differentiation
developmental process
cell differentiation
cellular developmental process
animal organ development
epidermis development
cell population proliferation
positive regulation of cellular process
skin development
regulation of cell population proliferation
multicellular organismal process
anatomical structure morphogenesis
Abnormal epidermal morphology
Epidermal acanthosis
Corneal erosion
Palmoplantar hyperkeratosis
Hyperkeratosis
Plantar hyperkeratosis
Palmar hyperkeratosis
Palmar hyperlinearity
Palmoplantar keratoderma
Hypergranulosis
Abnormality of the plantar skin of foot
Formation of the cornified envelope
Keratinization
Condensation of Prometaphase Chromosomes

-log10(p_value): 40, 30, 20, 10

intersection_size: 100, 200, 300

BP:Biological Process; HP:human phenotype ontology; REAC: REACTOME

i *ADGRL2* and *GNA13* Depletion RNA-seq Overlap Hits

Genes upregulated during differentiatrion
Genes downregulated during differentiatrion
Genes unchanged during differentiatrion

log2FC(GNA13 si/control si)

$R^2=0.16$

log2FC(L2 si/control si)

j 3D Organoid IF Quantification

control siRNA
GNA13 smartpool siRNA

KRT10, TGM1 — **, ***

Figure 4S