A gene-based model of fitness and its implications for genetic variation: Genetic and inbreeding loads

Parul Johri[1,2,3,*] and Brian Charlesworth[4,*]

[1]Department of Biology, University of North Carolina at Chapel Hill, NC 27599, United States
[2]Department of Genetics, University of North Carolina at Chapel Hill, NC 27599, United States
[3]Integrative Program for Biological and Genome Sciences, University of North Carolina at Chapel Hill, NC 27599, United States
[4]Institute of Ecology and Evolution, School of Biological Sciences, University of Edinburgh, EH9 3FL, United Kingdom

*correspondence may be addressed to these authors: pjohri@unc.edu; brian.charlesworth@ed.ac.uk

Keywords: complementation, dominance, epistasis, fitness models, genetic load, inbreeding load

Running title: Fitness models and mutational load

Abstract

In the companion paper to this, we examined the consequences for patterns of linkage disequilibrium of the "gene" model of fitness, which postulates that the effects of recessive or partially recessive deleterious mutations located at different sites within a gene fail to complement each other. Here, we examine the consequences of the gene model for the genetic and inbreeding loads, using both analytical and simulation methods, and contrast it with the frequently used "sites" model that allows allelic complementation. We show that the gene model results in a slightly lower genetic load, but a much smaller inbreeding load, than the sites model, implying that standard predictions of mutational contributions to inbreeding depression may be overestimates. Synergistic epistasis between pairs of mutations was also modeled, and shown to considerably reduce the inbreeding load for both the gene and sites models. The theoretical results are discussed in relation to data on inbreeding load in

Drosophila melanogaster. The widespread assumption that inbreeding depression is largely

due to deleterious mutations should be re-examined in the light of our findings.

Introduction

A large volume of recent work on the properties of diploid populations subject to the input of

deleterious mutations has been stimulated by concerns about their implications for the

survival of small, endangered populations, and by the availability of population genetic data

on non-model organisms that shed light on the prevalence of deleterious mutations in natural

populations (e.g., Teixeira and Huber 2021; Kardos et al. 2021; Bertorelli et al. 2022; Peréz-

Pereira et al. 2002; Kyriazis, Robinson and Lohmueller 2023; Robinson et al. 2023). There

are, however, sharp disagreements about the interpretation of the data (Kyriazis et al., 2021,

2023; García-Dorado and Hedrick 2023), emphasizing the need for a secure theoretical basis

for data analyses.

In the companion paper to this (Johri and Charlesworth 2025), we analysed the

consequences for patterns of linkage disequilibrium (LD) of the differences between two

models of the fitness effects of deleterious mutations located within the same gene, the

"gene" and "sites" models. The gene model assumes that two heterozygous mutations in

trans have a larger effect on fitness than two mutations in *cis*, due to a lack of

complementation, whereas the sites model assumes that there is no such difference. Many

theoretical predictions about the properties of deleterious mutations in populations, and their

effects on features such as population mean fitness, genetic variance in fitness and inbreeding

depression have been based on the sites model, together with the assumption of multiplicative

fitnesses when the effects of mutations at different sites are combined (for a recent review,

see Kyriazis, Robinson and Lohmueller 2023).

But the difference between the gene and sites models can affect the effective level of

recessivity experienced by deleterious mutations, as can be seen from the following, highly

simplified, example. Consider a single coding sequence that is segregating for mutations at

two different nucleotide sites (1 and 2) with the same selection coefficient s against

homozygotes at each site. Assume that the double mutant haplotypes are sufficiently rare that

they can be ignored. Let the frequency of mutant variants at site i be q_i. Then haplotypes

carrying a mutation at site 1 will meet a haplotype that is wild-type at site 1 and mutant at site

2 with an approximate frequency of p_1q_2; the fitness of this genotype is $1 - s$ if there is no

allelic complementation (*i.e*, under the gene model), but the sites model would assign it a

fitness of $1 - 2hs$ (assuming additive effects across sites), where h is the dominance

coefficient. It will meet haplotypes that are mutant at site 1 but wild-type at site 2 with a

frequency of approximately $q_1 p_2$. Both models would assign fitness $1 - s$ to this genotype. A

similar argument applies to a haplotype carrying a mutation at site 2, with the appropriate

switch between subscripts 1 and 2.

All of the double mutant genotypes are classed as homozygotes under the gene model,

which is biologically realistic for the majority of genes, for which sites with allelic

complementation are relatively rare (Fincham et al. 1979, pp.348-353; Hawley and Gilliland

2006). The mean difference between the gene and sites models in the fitness reduction

assigned to haplotypes that each carry a mutation at one or other site is thus:

80
$$- \delta w = (1 - 2h)s \frac{2(q_1 p_2 p_1 q_2)}{(q_1 p_2 + p_1 q_2)} \qquad (1)$$

If the frequencies of the deleterious mutations at the two sites are similar, so that the

subscripts can be dropped, $- \delta w \approx 2(1 - 2h)spq$, which is positive if $h << 0.5$. If $h > 0$ and

the population is at deterministic mutation-selection balance, $q \approx u/(hs)$, (Haldane 1927), and

$- \delta w \approx 2(1 - 2h)u/h$. For completely recessive deleterious mutations at equilibrium, $q \approx$

$\sqrt{u/s}$ and $-\delta w \approx 2(1 - 2h)\sqrt{us}$. This implies that the sites model can underestimate the

true homozygous fitness effects of deleterious mutations, although the effect is small for a

single pair of rare deleterious variants. It follows that deleterious mutations are more likely to

segregate at higher allele frequencies under the sites model than the gene model, leading to a

lower mean fitness of the population, a higher genetic load, and a larger effect of inbreeding

on fitness. In addition to this effect, our previous paper showed that the gene model can

generate positive LD among deleterious variants under circumstances where the sites model

causes zero or slightly negative LD (Johri and Charlesworth 2025). Since positive LD

enhances the efficacy of selection (e.g., Barton 1995; Roze 2021), this effect would also help

to reduce the frequency of deleterious mutations expected under the gene model compared

with the sites model.

The purpose of the present paper is to examine the consequences of the differences

between the gene and sites models for the load statistics, using deterministic and stochastic

simulation models of a single coding sequence subject to deleterious mutations, which were

described in Johri and Charlesworth (2025). We show that the difference between the two

models has especially significant consequences for the level of inbreeding depression caused

by deleterious mutations, suggesting that predictions based on the sites model could greatly

overestimate the expected level of inbreeding depression contributed by deleterious

mutations. We also show that a model of pairwise synergistic epistasis between mutations

within the same gene results in even lower levels of inbreeding depression than in its

absence. We also note that the sites model is used in current methods for estimating the

distribution of fitness effects of deleterious mutations, and that this will lead to

overestimation of the mean strength of selection against such mutations if the gene model is

more appropriate.

Methods

Fitness models and simulation methods

The fitness models and simulation results used here are described in the Methods section of

Johri and Charlesworth (2025). Recursion relations for haplotype frequencies for the two-site

deterministic model are described in the Appendix of that paper; use of these expressions

allows the numerical values of the equilibrium haplotype frequencies to be determined for a

given set of mutation and fitness parameters. In addition, the multi-site approximation for the

gene model without epistasis with the fixed selection and dominance coefficients at all sites,

described in the penultimate section of the Appendix to Johri and Charlesworth (2025),

provides expressions for the load statistics that can be compared with the simulations when

selection is strong relative to genetic drift. Standard formulae for mutation-selection balance

can be used for the corresponding calculations with the sites model (see section 2 of the

Appendix to the present paper).

Estimating the load statistics from the simulations

The state of a population with N diploid individuals (where N was usually equal to 1000)

with respect to the load statistics was summarized using four quantities – the genetic load (L),

the inbreeding load (B), the variance in fitness between individuals (V), and the mean

frequency of mutant alleles per site ($\bar{q}$). The genetic load for each simulation replicate was

calculated as $L = 1 - \bar{w}$, where $\bar{w}$ is the mean fitness of the population for all individuals in

the simulated population, relative to a value of 1 for individuals homozygous for the wild-

type allele at all sites; $\bar{w}$ was calculated using all individuals sampled after burn-in. The

variance V was calculated in a similar way. Allele frequencies (q) were estimated from a

sample of 100 genomes and included monomorphic sites.

135   To calculate B, which is defined here as the mean difference in fitness between

136 completely outbred and completely inbred individuals, 100 genomes were sampled randomly

137 from the population, from which 100 diploid inbred individuals were artificially created (*i.e,*

138 each haploid genome was duplicated from a single diploid individual). The inbreeding load

139 was calculated as $B = \bar{w} - \bar{w}_I$, where $\bar{w}_I$ is the mean fitness of the subsampled inbred

140 population. Because L and B are both small, this expression for B differs little from the usual

141 expression for inbreeding depression, $1 - \bar{w}_I/\bar{w}$. Under the assumption of small

142 multiplicative fitness effects across sites, it is equivalent to the negative of the regression

143 coefficient of the natural logarithm of fitness on inbreeding coefficient (Morton et al. 1956).

144 Each simulation replicate yields a single data point for each variable, and the distributions,

145 means and standard errors shown in the figures and tables are derived from 1000 replicate

146 simulations.

148 **Data availability**

149 The scripts used to determine the properties of two-site equilibrium populations, perform all

150 the simulations, and calculate population genetic statistics are provided at

151 https://github.com/paruljohri/Gene_vs_sites_model/tree/main.

153 **Results**

154 **Deterministic results for two selected sites**

155 To aid the reader, Table 1 of Johri and Charlesworth (2025) is reproduced here. The fact that

156 the two sites have identical fitness effects means that the four haplotype frequencies, x_1 (+ +),

157 x_2 (− +), x_3 (+ −) and x_4 (− −) (where + and – indicates wild-type and mutant alleles,

158 respectively) can be replaced with $x = x_2 = x_3$, $y = x_4$, and $z = 1 - 2x - y$. The frequency of the

159 – allele at a site is $q = x + y$. The load statistics L, B, and V (the genetic load, the inbreeding

160 load, and the genetic variance, respectively) can easily be determined once the equilibrium

161 haplotype frequencies are known, using the formulae presented in the Appendix, Equations

162 (A1) – A(8). As in Johri and Charlesworth (2025), we assume no recombination between the

163 two sites, which should provide a good approximation to what is expected for a single gene.

164   Table 2 shows the results for the case of $u = 0.0001$, $s = 0.01$, which was previously

165 used for the LD statistics presented in Johri and Charlesworth (2025). As noted there, these

166 are not realistic parameter values, but produce relatively large equilibrium allele frequencies,

167 enhancing the contrast between the two models. The frequency of deleterious mutations (q_r in

Table 2) is represented by the ratio of the exact deterministic equilibrium value to the

approximate value for a single site subject to mutation and selection, given by Equation (A4)

of Johri and Charlesworth (2025). Since the denominator q_r is independent of the strength of

epistasis, and is the same for the gene and sites model, differences in q_r between the two

models and between different values of the epistasis coefficient ϵ for a given h and s reflect

differences in the absolute deleterious allele frequency, q.

L and B are represented by the ratios of their values to those for an additive model of

two independent loci, when $h > 0$ and $u << hs$. In this case, $L \approx 4u$ (see Haldane 1937) and

$B \approx 2u[\ (1/(h) - 2]$ (see Morton et al. 1956); if u is sufficiently large relative to hs or drift

is operating, these formulae no longer hold. To avoid large values when h is small, the

variance is represented by $Var = hV/(4uhs)$, the denominator being the equilibrium value of

V for the purely additive case with no LD and $u << hs$. This means that results for the

variances with different h values are not directly comparable, but the sites and gene models

with the same h can be compared.

As before, we focus attention on cases with zero epistasis ($\epsilon = 0$) and synergistic

epistasis ($\epsilon > 0$). Over the range of values of ϵ considered here, its magnitude has relatively

little effect on L and B, for both the sites and gene models. Its effect on q_r is more complex.

For both models, q decreases with ϵ, but the decline is faster for the sites model when h is

close to ½, with the result that in this case q is smaller for the sites model than the gene model

when ϵ is large, despite their having the same q with no epistasis. For h sufficiently smaller

than ½, q is smaller for the gene model.

The gene model, as might be expected, is associated with lower values of L and B

than the sites model when $h <$½, with a much lower value of B than the sites model when h is

small. The higher values of V under the gene model with small h reflect its larger values of

LD compared with the sites model, with positive associations ($D > 0$) between deleterious

variants when ϵ is sufficiently small, especially with small h (Johri and Charlesworth 2025).

With $h =$ ½, $B = 0$ (so that the ratios in Table 2 are set to one) and L is the same for the two

models, as expected. V increases more quickly with ϵ for the sites model than the gene

model, the reverse of the pattern for q. This can result in a slightly higher V for the sites

model with large ϵ and h, probably reflecting the larger contributions of terms in ϵhs under

the sites model, as can be seen in the fitness matrices in Table 1.

Simulation results for multiple sites

Load statistics with no epistasis

The analytical results described above suggest that, relative to the sites model, the gene

model should result in smaller mean frequencies of the deleterious alleles, causing a

a lower genetic load and inbreeding load. The multiple site simulations with scaled

parameters that are appropriate for *Drosophila* populations, *i.e*, 1000 selected sites, $Nu =$

0.005, and $Nr = 0.01$ (for details, see the *Methods* section of Johri and Charlesworth 2025),

show that the equilibrium frequencies of deleterious alleles are much smaller for the gene

than the sites model, except when selection is weak relative to drift ($\gamma = 2$ for the case of a

fixed selection coefficient (Table 3) and $\bar{\gamma} = 2$ for the case of a gamma distribution of

selection coefficients (Table 4), in which case there is little difference between the two

models. There is also a drastic difference between the estimated inbreeding load (B) between

the two models: B is much smaller for the gene than the sites model when $h < ½$, even with

weak selection. There is a much smaller effect on the genetic load (L), which is noticeable

only when mutations are fully recessive.

The equations for the load statistics derived in section 2 of the Appendix were used to

provide the predicted values in Table 3. These are expected to be accurate only for $\gamma \gg 1$;

accordingly, it can be seen that the predictions of the genetic and inbreeding loads mostly fit

much worse for $\gamma = 2$ than for $\gamma = 20$ and $\gamma = 100$, although the mean frequencies of

deleterious alleles are always remarkably close to the predicted equilibrium values when $h =$

0.2 and $h = 0.5$. For the sites model, however, there are strong departures from the predicted

equilibrium values with $h = 0$, even for $\gamma = 100$. This reflects the fact that the mean allele

frequency for a fully recessive deleterious mutation in a finite population is expected to be

considerably lower than the equilibrium value, as a result of the elimination of deleterious

homozygotes produced by drift (Nei 1968). This also causes the inbreeding load for the sites

model to be greatly overpredicted when $h = 0$.

For $\gamma = 20$ and $\gamma = 100$, the predictions of the values of the genetic and inbreeding

loads with $h = 0.2$ and 0.5 are accurate for both the sites and gene models. In contrast, the

predicted variances in fitness with $h = 0$ and $h = 0.2$ for these cases are substantially smaller

than the variances found in the simulations. The main cause of this difference is likely to be

the positive covariance in allelic effects between different sites resulting from LD, which has

been ignored in these predictions, although there may also be some contribution from the

dominance variance. This effect can be assessed quantitatively, by means of the predicted

values of mean D at all sites, given by Equations (A19a) of Johri and Charlesworth (2025).

Using the formulae of Bulmer (1980, p.158), with equal allelic effects at all selected sites,

and assuming that non-additive variance in fitness is negligible, the expected component of

variance arising from LD is $C_L = 2G^2\overline{D}V_g$; the net predicted fitness variance is $V = V_g + C_L$.

Using the predicted values of V_g in Table 3, this formula yields predictions of 6.85 x 10^{-5} and

6.25 x 10^{-5} for the variances with $\gamma = 20$ and $h = 0$ and 0.2, respectively; for $\gamma = 100$, the

corresponding values are 1.47 x 10^{-4} and 1.73 x 10^{-4}. The corrections are quite substantial for

$\gamma = 20$, and bring the predicted values close to the simulated values, but are negligibly small

with $\gamma = 100$. No contribution from LD is expected for $h = 0.5$, and the predicted variances in

fitness match the simulation results in this case. LD is also expected to have minimal effects

on V for the sites model.

The effect of LD on V are counteracted by lower allele frequencies under the gene

model, an effect that is especially strong for fully recessive mutations, leading to a much lower

V under the gene model than the sites model when $h = 0$. However, LD causes V to be

significantly higher for the gene model than the sites model when $h = 0.2$ and $\gamma = 20$. With h

$= \frac{1}{2}$, there is no difference between the two models, as expected. The difference in V between

the two models thus has a non-linear relation with h.

Because the results discussed above are most relevant to *D. melanogaster*-like

population parameters, we also conducted simulations with parameters appropriate for human

populations (for details, see Johri and Charlesworth 2025), with much smaller values of Nu

(0.00025) and Nr (0.0002). The overall patterns are very similar to those for *Drosophila*-like

populations (Table S2 in Supplementary File S1). The mean frequencies of deleterious alleles

are significantly smaller with the gene than the sites model; genetic load is slightly lower, while

inbreeding load is again drastically smaller for the gene model than the sites model when $h <$

$\frac{1}{2}$.

Effects of rescaling of population genetic parameters

To evaluate the possible effects of rescaling to keep the products of N_e with the deterministic

population genetic parameters such as selection coefficients and mutation rates consistent with

their natural population values for *D. melanogaster* (see Dabi and Schrider 2024; Ferrari et al.

2024; Johri and Charlesworth 2025), simulations were also performed with 5000 diploid

individuals, where Nu, Nr, and Ns were the same as for $N = 1000$. The mean equilibrium

frequencies of deleterious alleles were found to be identical in both cases (compare Table 3

with Table S1 in Supplementary File S1), as is expected from the general theory of population

genetic processes when all evolutionary forces are weak and their strength can be represented

by single variables such as u, s and r (e.g., Ewens 2004, Chap.5). The genetic and inbreeding

loads involve products of the haplotype frequencies and s, whereas the variance involves

products of the haplotype frequencies and s^2 (see section 1 of the Appendix). If the haplotype

frequencies are preserved after multiplying the population size by a factor C and keeping the

scaled parameters constant, the simulated values of the two load statistics should thus be

multiplied by C and the variance by C^2, in order to produce values that are comparable with

the results for the smaller N value ($C = 5$ in the present cast). As can be seen from Tables 3 and

S1, the load statistics for $N = 5000$ after this correction agree well with those for $N = 1000$,

implying that we can have confidence in the rescaling procedure. These results imply that

studies interested in evaluating the extent of mutational load in endangered populations using

forward simulations need to apply similar corrections to the load statistics if rescaling is

employed at the single gene level, where recombination rate is almost linearly related to map

distance.

Load statistics with epistasis

With synergistic epistasis between deleterious mutations, a decrease in their mean

equilibrium frequencies is expected compared with the case with no epistasis, due to the

reduced fitness of genotypes with multiple mutations that is caused by synergistic epistasis

(Crow 1970; Kondrashov 1988; Charlesworth 1990) In the simulations with multiple linked

sites under a DFE with a relatively large mean scaled selection coefficient ($\bar{\gamma} = 2N\bar{s} = 100$)

and varying degrees of epistasis (with ϵ ranging from 0.02 to 0.32), there is a severe decrease

in the mean equilibrium frequency of deleterious mutations with increasing ϵ, for both the

gene and sites models (Figure 1). Even with the lowest strength of epistasis ($\epsilon = 0.02$), there

is no significant difference in mean equilibrium frequencies of deleterious alleles between the

gene and sites models. There is a corresponding decrease in the inbreeding load (B) with

increasing ϵ for both the gene and sites model (Figure 2) when $h < ½$.

B is still significantly and drastically higher with the sites model than the gene model

for both $h = 0$ and $h = 0.2$. There is no difference between the two models with $h = 0.5$, as

expected; unexpectedly, however, B becomes increasingly slightly negative as ϵ increases.

One possible explanation is that this is an artefact of the way B in which was estimated; the

fitness of inbred individuals was calculated by sampling 100 haploid genomes, whereas the

mean fitness was calculated using the entire population. However, even if we calculate the

fitness of outbred individuals by sampling 100 diploid individuals, B remains negative for

these parameter sets (see Figure S1 in Supplementary File S1), so that this explanation can be

ruled out.

The probable cause of this effect is as follows. With the parameter set in question, the

frequencies of deleterious alleles are sufficiently high that essentially all haplotypes carry at

least one deleterious mutation. In the two-site case with $h = 0.5$, a $-+/+-$ individual has

fitness reduction $s(1 + \epsilon)$, whereas $-+$ or $+-$ homozygotes have a fitness reduction s (there

is no effect of epistasis). Thus, the heterozygote has a lower fitness than the homozygote. In

other words, the outbred population has more opportunities for pairwise fitness interactions

that reduce fitness than the inbred population, resulting in slightly negative inbreeding

depression. Epistatic interactions of this kind have been studied theoretically and empirically

in connection with the biometrical genetics of crosses between inbred lines (e.g., Wright

1968, Chap. 15, pp. 391-403; Jinks 1983). A lower value of a fitness component in the F_1

compared with the mean of the two parents of the kind expected under this model appears to

be uncommon in such crosses, with a prevalence of the opposite pattern (heterosis),

consistent with the effects of largely recessive effects of deleterious alleles or heterozygote

advantage outweighing epistatic effects of this kind.

There is a much smaller effect of epistasis on the genetic load (L in Figure 2), with a

minor and steady decrease in L with an increase in ϵ for both fitness models. As before, the

difference in the genetic load between the two models is relatively small and only significant

when mutations are completely recessive. The variance in fitness (V) also increases with an

increase in the epistasis coefficient when $h = 0.2$ or 0.5 (and only very slightly when

mutations are fully recessive). However, as was also found without epistasis, the variance is

much larger for the gene model compared to the sites model, especially when mutations have

increased recessivity and intermediate/low levels of epistasis (Figure 1). With high levels of

epistasis ($\epsilon = 0.32$), however, there is no difference in V between the two fitness models. In

summary, if there were widespread epistasis between deleterious mutations, there would be

no substantial difference in mean equilibrium allele frequencies between the two models, but

the inbreeding and genetic loads would be lower with the gene model than the sites model if

mutations have moderate to high recessivity. Note that with much lower rates of crossing

over ($0.1 \times$ the standard rate used here), the behavior of the load statistics remains the same

as described above (Figures S1 and S2 of Supplementary File S1).

Discussion

Load statistics with no epistasis

Differences between the gene and sites models.

The main conclusion from the analytical and simulation results for the case of no epistasis is that the inbreeding load B is always much smaller under the gene model than the sites model, being effectively zero for $\bar{\gamma} \leq 20$, unless the dominance coefficient h is close to 0.5; in this case B is close to zero for both models, as expected from basic theory (Morton, Crow and Muller 1956). A similar pattern is seen with synergistic epistasis, but B becomes very small for both the gene and sites models, and even slightly negative, when epistasis is strong (Figure 2).

These results raise the question of why B should be so much smaller under the gene model than the sites model. One factor is the smaller mean frequency of deleterious alleles ($\bar{q}$) for the gene model than the sites model, at least when $\gamma >> 1$. However, the relation of $\bar{q}$ to the strength of selection does not track the behavior of B. For example, for the case of a gamma distribution with $h = 0.2$, B and $\bar{q}$ for the gene model with $\bar{\gamma} = 100$ are approximately 6% and 30% of the values for the sites model, respectively; when $\bar{\gamma} = 1000$, B and $\bar{q}$ for the gene model are 30% and 12% of the sites model values, respectively (see Table 4).

This discrepancy probably arises from the fact that haplotypes carrying deleterious mutant alleles can be relatively common when the number of selected sites in a gene (G) is large. For example, with $\gamma = 20$ and $h = 0.2$ the equilibrium frequency of deleterious alleles is approximately 0.001 (see Table 3) so that the mean number of mutant alleles per haploid genome with $G = 1000$ is $1000 \times 0.001 = 0.1$. Genotypes with deleterious mutations in *trans* are thus relatively common, so that a heterozygous mutation at a given site behaves more like a dominant mutation than under the sites model, reducing its contribution to inbreeding depression. It is surprising that such a large difference in the expectation for the level of inbreeding depression caused by a single gene can be produced by the lack of complementation between mutations in *trans*.

Another way of looking at this effect is to note that B for the additive gene model when $\gamma >> 1$ is approximately $(1 - 2h)\mu e^{-\mu}s$ (see Equation A8b), where $\mu = G\hat{q}$ is the mean number of mutations per haplotype and $\hat{q}$ is the deterministic equilibrium frequency of mutant alleles. In contrast, B for the additive sites model in this case is equal to $Gu(1 - 2h)/h$ when $h > 0$ (Morton et al. 1956; Charlesworth and Hughes 2000). When $h > 0$ and $\mu << h$, Equation (A21b) of Johri and Charlesworth (2025) shows that $\hat{q}$ approaches the sites value of

$u/(hs)$. The values of B for the two models thus converge when selection is sufficiently

strong, and G is sufficiently small that $e^{-\mu} \approx 1$. A similar result holds when $h = 0$.

There is a lack of information on whether or not the lack of complementation

assumed in the gene model applies to deleterious mutations with minor effects on fitness, as

opposed to the loss-of-function mutations that have generally been studied in relation to

complementation. There are, however, a number of studies of the phenotypes of

heterozygotes for hypomorphic mutations in the same gene, suggesting that partial

impairment of functionality is usually associated with lack of complementation. Wright

(1968, p.70) gives example of multiple alleles at coat color loci of the guinea pig that show

this behavior, and Chandler et al. (2017) describe similar effects for two wing size loci of *D.*

melanogaster. Quantitative studies of dominance effects on fitness in diploids such as yeast

(*e.g.*, Matsui et al. 2022), with a focus on measuring complementation, are needed to shed

further light on this question.

Other models of deleterious mutations

We now consider the relation of the gene model to the widely used model that treats a gene

as a single non-recombining unit subject to mutation from wild-type alleles to mutant alleles,

each of which causes a fitness reduction of s when homozygous or in combination with each

other, and hs when heterozygous with wild-type (e.g. Charlesworth et al. 1992; Wang et al.

1999; Kamran-Disfani and Agrawal 2014; Bersabé et al. 2016). This type of model is

equivalent to a single locus with mutation at rate Gu from wild-type to a mutant allele with

selection coefficient s.

At first sight, this situation seems equivalent to the gene model with identical

selection coefficients for each mutation. However, with the relatively weak selection against

many deleterious mutations suggested by population genomics data (see section 1 of

Supplementary File S2), the ratio of Gu to hs is likely to be such that most haplotypes

segregating in the population carry at least one mutation. Even with relatively strong

selection ($\bar{\gamma} = 1000$ and $h = 0.2$ in Table 4), the mean number of deleterious mutations per

haploid genome for a sequence of 1000 selected sites is 1000 x 0.00091= 0.91.

The assumption that all deleterious mutations are only one step away from wild-type

may thus be unsafe. This has important consequences for the predicted values of the genetic

and inbreeding loads. Consider the case of a fixed selection coefficient with $\gamma = 20$ described

in Table 3. In this case, the mean deleterious allele frequency per site is 0.00098 under the

gene model, with a very small standard error. With the weak LD associated with these

parameters (see Figure 3 of Johri and Charlesworth 2025), the distribution of the number of

mutations per haploid genome closely follows a Poisson distribution, so that (disregarding

variability between replicate simulations) the probability of a mutant-free haplotype is

approximately $\exp(-0.98) \approx 0.38$. The predicted frequency of haplotypes containing at least

one mutation is thus 0.62. If this is used as the deleterious allele frequency q in the standard

formulae for L and B ($L = [2h + (1 - 2h)q]sq$; $B = sq - L$), we obtain $L = 0.0048$ and $B =$

0.0014. These are substantially lower than the predicted values of $L = 0.0078$ and $B = 0.0022$

under the gene model, reflecting the fact that this calculation ignores the contributions of

haplotypes carrying more than one mutation. One-step mutational models can therefore

seriously underpredict the load statistics, producing a bias in the opposite direction to that

described in the previous section.

Implications for data on inbreeding load

Recent studies of genetic and inbreeding loads that have used population genomics data to

simulate populations subject to deleterious mutations and to predict genetic and inbreeding

loads (reviewed by Kyriazis, Robinson and Lohmueller 2023) have mostly used the sites

model. Multiplicative fitnesses across selected sites and genes have also usually been

assumed in models of inbreeding load – but see Charlesworth (1998) and García-Dominguez

et al. (2018) for exceptions. Our results are for a single gene, so it is important to consider

what happens if they are extrapolated to the whole genome. For convenience, we will assume

purely autosomal inheritance, leading to a probable overestimation of the load statistics for

the entire genome, given that the X chromosome is associated with lower loads than

autosomes due to selection against hemizygous mutations in males (Wilton and Sved 1979).

To scale up to multiple genes, we assume that the sites model applies to between-gene

effects for both models of single genes, due to complementation between non-allelic

mutations. As in previous studies, starting with Haldane (1937), we assume multiplicative

fitness effects across different genes; with small selective effects the differences from

additive effects will be small in most cases. Our simulation results were based on a scaling of

population size from 10^6 to 10^3; to generate realistic results for a *Drosophila*-like system, we

would need to rescale back to the $N = 10^6$ values; this means dividing linear quantities in s by

10^3 and quadratic quantities such as the variance by 10^6 (see the discussion of rescaling in the

section *Load statistics with no epistasis*). If we assume 14000 genes in the *Drosophila*

genome (Misra et al. 2002), the individual gene linear quantities (the loads and inbreeding

loads) in the tables should thus be multiplied by 14 and the variance by 0.014. This ignores

any contributions from mutations in non-coding sequences, which will be considered below.

In general, let the linear and quadratic multipliers reflecting rescaling be f and f^2, and

the number of genes be n. For the inbreeding load, defined as the difference between the

natural logarithms of outbred mean fitness and fully inbred mean fitness, the B values in the

tables need only be multiplied by nf to obtain the total inbreeding load, B_T. If the mean load L

for each gene is small, the total genetic load with multiplicative effects across n genes is

given by the following expression:

442
$$L_T \approx 1 - \exp(-nfL) \qquad (2)$$

444   Similarly, if there is no LD between different genes, using the rule that the

445   expectation of a product of a set of independent variables is the product of their expectations,

446   the total variance with multiplicative effects across genes is given by:

448
$$V_T \approx \exp(-2nfL)[\exp(-nf^2V_g) - 1] \qquad (3)$$

Given n and f, the values of the relevant statistics in the tables can easily be converted

into values for the whole genome by using these relations. Using the mean values of the load

statistics in Table 4 with $\bar{\gamma} = 1000$, Table 5 shows the values obtained with $n = 14000$ and $f =$

14; σ_T is the standard deviation corresponding to V_T. These results show that the difference

between the two models has only a small effect on the expected genome-wide genetic load

and genetic variance/standard deviation (mean s is only 5×10^{-4} after rescaling N), but that

the gene model has a drastically smaller total inbreeding load than the sites model with $h =$

0.2 (0.050 versus 0.165), despite the fact that the sites model applies to between-locus fitness

effects.

Our results suggest that the sites model of mutation and selection is likely to

considerably *overpredict* the value of B, unless the mean strength of selection is so large and

the distribution of selection coefficients is so tight that the population can be treated as fully

deterministic. Population genomics studies of the DFE for nonsynonymous mutations in

organisms such as humans (Kim et al. 2017) and *Drosophila melanogaster* (Campos et al.

2017; Johri et al. 2020) suggest that a substantial fraction of deleterious mutations are subject

to the joint effects of drift and selection (see section 1 of Supplementary File S2).

This conclusion is important, because reconciling observed and predicted values of B

has been the subject of much debate, with some workers asserting that the observed values

can be explained by mutation-selection balance with multiplicative fitnesses (e.g., Pérez-

Pereira et al. 2021; Robinson et al. 2023), while others have proposed that additional factors

(such as variation due to balancing selection) are likely to be involved (e.g., Charlesworth

2015). If it can be shown that empirical estimates of B are incompatible with values predicted

on the basis of the sites model, using well-grounded estimates of mutation rates, dominance

coefficients and realistic demographic histories, then contributions from balancing selection

or synergistic epistasis of the type modeled by Charlesworth (1998) must be involved.

A difficulty with this approach is that the currently used methods for inferring the

DFE from population genomic data assume the sites model for predicting variant frequency

distributions at sites under selection. As pointed out in section 1 of Supplementary File S2,

this means that the mean strength of selection as measured by $\bar{\gamma}$ is likely to be overestimated,

given that the gene model predicts lower deleterious variant frequencies than the sites model.

Further research into the magnitude of this problem needs to be carried out, but the fact that it

is likely to inflate the predicted sizes of the genetic and inbreeding loads needs to be borne in

mind. Moreover, interpreting empirical estimates of the extent of inbreeding is problematic

because reliable estimates of B for net fitness in natural populations are hard to obtain, with

substantial disagreements among different workers concerning the typical magnitude of B

(Robinson et al. 2023). This makes it difficult to compare the data on inbreeding load with

theoretical predictions.

Inbreeding load in Drosophila

In order to avoid the problems associated with interpreting measurements of fitness in wild

populations, we here compare theoretical estimates of levels of inbreeding load with

estimates obtained from laboratory experiments on *D. melanogaster*. An ingenious method

for estimating B for net fitness under laboratory conditions for a single major chromosome of

some *Drosophila* species was devised by John Sved, known as the balancer equilibration

(BE) technique (Sved and Ayala 1970). This is based on using population cages to compete

wild-type chromosomes extracted from natural populations (chosen to be homozygous viable

and fertile) against homozygous lethal balancer chromosomes that suppress crossing over

when heterozygous. The equilibrium frequencies of wild-type chromosomes can be used to

estimate the ratio R of the mean fitness of chromosomal homozygotes to the mean fitness of

chromosomal heterozygotes (reviewed by Frankham 2023). The mean value of R for

chromosome 2 of *D. melanogaster* over two well-replicated experiments was 0.18±0.03,

giving an estimate of $B = -\ln(R)$ of 1.7 (Frankham 2023). The BE results on the data for all

three major chromosomes imply that B for the whole genome of *D. melanogaster* (including

the contribution from homozygous lethal mutations) is approximately 5 (Frankham 2023).

504   The analysis of population genetic data for *D. melanogaster* presented in section 3 of

the Appendix yields a prediction of $R = 0.66$ and $B = 0.42$ for chromosome 2 under the sites

model. Here, possible effects of purifying selection at non-coding sites as well

nonsynonymous sites and strongly selected (but non-lethal) mutations are taken into account,

using fairly liberal assumptions with respect to the magnitude of B, so that the resulting

values are likely to overestimate B. This value of B uses the deterministic expressions for R

and B given by Equations (A9) and (A10), and assumes an unrealistically low h value of

0.125. In addition, an approximate correction for the effects of drift was obtained by

multiplying u by the proportion of mutations with $2N_e hs < 2.5$ under a gamma distribution

with shape parameter 0.3, thereby removing the contribution of nearly neutral mutations (see

section3 of the Appendix).

515   Given that the gene model predicts a substantially lower value of B than the sites

model, $B = 0.42$ for chromosome 2 is almost certainly an overestimate, but is still less than

25% of the observed value. There thus appears to be much more inbreeding load due to

deleterious mutations under laboratory conditions in *D. melanogaster* than can be accounted

for purifying selection under a multiplicative fitness model, especially considering the

various biases towards overpredicting the inbreeding load caused by deleterious mutations.

Why is the inbreeding load reduced when $N_e s$ is small?

Tables 3 and 4 show that, other things being equal, smaller values of B are associated with

smaller values of $N_e s$. Numerical methods for obtaining an exact single-locus prediction for B

in a finite population have been developed by Charlesworth (2018), and extensive multi-locus

simulation results using a variety of parameter values have been described in the literature

(Kyriazis et al. 2023). Both of these approaches show that the level of inbreeding load caused

by deleterious mutations is positively related to $N_e s$ under a wide range of model

assumptions. As has been widely discussed in the literature (see review by Robinson et al.

2023), there are two possible reasons for this effect – the "purging" of deleterious, strongly

recessive variants from small populations as a result of their exposure to selection against

homozygotes (Nei 1968), and the loss of variability, including the (temporary) fixation of

deleterious variants (Kimura et al. 1963). The analysis presented in sections 2 and 3 of

Supplementary File S2 shows that the second factor is responsible for the effects seen in our

simulations.

The weak effects of drift on the genetic load and variance in fitness seen in Table 4

are at first sight surprising, given that with $\bar{\gamma} = 2$ a substantial proportion of the distribution of

selection coefficients falls below the nearly neutral threshold of $h\gamma < 2.5$ described above,

where large departures from the deterministic values are expected. However, this part of the

distribution involves very small selection coefficients, and thus makes only a small

contribution to the genetic load and the variance in fitness. This mitigates the effect of drift in

causing an increase in load. In contrast, with a fixed scaled selection coefficient of $\gamma = 2$,

there is a substantial increase in load compared with the higher value of $\gamma = 20$ (Table 3).

The role of epistasis

Epistasis and inbreeding load

One important aspect of our simulation results with synergistic epistasis is that B for the gene

and sites models is strongly reduced by even a small amount of epistasis when $h < \frac{1}{2}$, despite

only a small reduction in load; there is a much smaller effect on B under the gene model,

where it is always small (Figure 2). This reduction in B with increasing epistasis is in sharp

contrast to the model of Charlesworth (1998), which predicts that synergistic epistasis causes

an *increase* in B when the logarithm of fitness is a quadratic, decreasing function of the

number of deleterious mutations. As shown in section 4 of the Appendix, when only

homozygous genotypes are considered, the pairwise epistasis model used here is identical to

the quadratic model. The opposite behavior of the two models is thus surprising at first sight.

The explanation lies in the fact that the model of Charlesworth (1998) assumes that h

> 0 and the quadratic term for heterozygous mutations is proportional to the negative of the

product of h^2 and the square of the number of heterozygous mutations (n^2). In contrast, the

pairwise epistasis model assumes that this term is proportional to $-hn^2$. This means that the

fitness of the outbred population is relatively insensitive to the quadratic fitness term under

the first model, since most mutations are present in heterozygotes. However, the fully inbred

genotypes have strongly reduced fitnesses due to the quadratic term, resulting in an increase

in B. Under the pairwise model, there is a sharp decline in $\bar{q}$ with increasing ϵ, while the load

is relatively unaffected (compare Figures 1 and 2). Since B is strongly determined by $\bar{q}$, the

net result is that synergistic epistasis reduces B, even when $h = 0$.

The consequences of epistasis for inbreeding depression are thus highly model dependent, and it cannot be assumed that synergistic epistasis necessarily leads to increased inbreeding depression compared with the multiplicative fitness model, as has been done in the past (Crow 1970; Charlesworth 1998; García-Dominguez et al. 2018). The pairwise model has the advantage of greater flexibility in modeling the fitnesses of heterozygotes and differences in selection coefficients between sites, whereas the quadratic model is forced to make somewhat arbitrary assumptions about the effects of dominance (for details, see Charlesworth 1998). However, it is important to note that we have only developed this model for within-gene interactions, and for pairwise interactions within genes; the nature of epistatic interactions between deleterious mutations in different genes and their consequences for the load statistics, as well as the effects of higher order interactions, remain to be explored.

Conclusions

Our results show that the gene model predicts much lower levels of inbreeding depression than are expected from current mutational models, and that a revised model of synergistic epistasis between mutations within genes leads to even smaller values of inbreeding depression, making it hard to explain observed values of the inbreeding load on the purely mutational hypothesis. It should, of course, be borne in mind that complex demography, including strong bottlenecks and hidden population structure, may also play a role in the observed mismatch between expected and observed estimates of B. More organism-specific simulations that take demographic history into account, as well as the differences between the gene and sites models, are likely needed to understand the respective contributions of deleterious mutations and balancing selection to inbreeding depression.

We note that much of the current literature on inbreeding depression assumes that this is caused almost entirely by deleterious mutations (e.g., Bertorelli et al. 2022; Peréz-Pereira et al., 2022; Robinson et al. 2023), despite evidence to the contrary dating back over many years (e.g., Charlesworth and Hughes 2000; Charlesworth 2015). The contribution of balancing selection to inbreeding depression should thus be further examined (*e.g.*, González-Castellano et al. (2025).

Acknowledgments

We thank Kirk Lohmueller, Denis Roze and three anonymous reviewers for their helpful comments on an earlier version of the manuscript. This research was conducted using

599   computational resources provided by ITS Research Computing at the University of North

600   Carolina at Chapel Hill. PJ was funded by the National Institute of General Medical Sciences

601   of the National Institutes of Health under the award number R35GM154969. We

Literature Cited

604 Assaf ZJ, Tilk S, Siegal ML, Petrov DA. 2017. Deep sequencing of natural and experimental
605   populations of *Drosophila melanogaster* reveals biases in the spectrum of new mutations.
606   Genome Res. 27:1988-2000.
607

608 Barton NH. 1995. A general model for the evolution of recombination. Genet. Res. 65:123-144.
609

610 Bertorelli G, Raffini M, Bortoluzzi C, Ianucci A, Trucchi E, Morales HE, Van Oosterhout C. 2022.
611   Genetic load: genomic estimates and applications in non- model animals. Nat. Rev. Genet.
612   23:492-503.
613

614 Bersabé D, Caballero A, Pérez-Figueroa A, García-Dorado A. 2016. On the consequences of purging
615   and linkage on fitness and genetic diversity. G3 6:171-181.
616

617 Bulmer MG. 1980. The Mathematical Theory of Quantitative Genetics. Oxford: Oxford University
618   Press.
619

620 Campos JL, Halligan DL, Haddrill PR, Charlesworth B. 2014. The relation between recombination
621   rate and patterns of molecular evolution and variation in *Drosophila melanogaster*. Mol. Biol.
622   Evol. 31:1010-1028.
623

624 Campos JL, Zhao L, Charlesworth B. 2017. Estimating the parameters of background selection and
625   selective sweeps in *Drosophila* in the presence of gene conversion. Proc Natl Acad Sci USA.
626   114(24):E4762–E4771.
627

628 Campos JL, Charlesworth B. 2019. The effects on neutral variability of recurrent
629   selective sweeps and background selection. Genetics 212:287-303.
630

631 Casillas S, Barbadilla A, Bergman CM. 2007. Purifying selection maintains highly conserved
632   noncoding sequences in Drosophila. Mol. Biol. Evol. 24:2222-2234.
633

634 Chandler CH, Chari S, Kowalaki A, Choi L, Tack D, DeNieu M, al. e. 2017. Complex effects of
635   genetic background on expressivity, complementation, and ordering of allelic effects. PloS Genet.
636   13:e1007075.
637

638 Charlesworth B. 1990. Mutation-selection balance and the evolutionary advantage of sex and
639   recombination. Genet Res. 55(3):199-221.
640

641 Charlesworth B. 1998. The effect of synergistic epistasis on the inbreeding load. Genet. Res. 71:85-
642   89.
643

644 Charlesworth B. 2015. Causes of natural variation in fitness: evidence from studies of Drosophila
645   populations. Proc. Natl. Acad. Sci. USA 12:1662-1669.

646
647 Charlesworth B. 2018. Mutational load, inbreeding depression and heterosis in subdivided
648   populations. Mol. Ecol. 24:4991-5003.
649
650 Charlesworth B, Hughes KA. 2000. The maintenance of genetic variation in life-history traits. In:
651   Singh RS, Krimbas CB, editors. Evolutionary Genetics from Molecules to Morphology.
652   Cambridge: Cambridge University Press. p. 369-392.
653
654 Charlesworth D, Morgan MT, Charlesworth B. 1992. The effect of linkage and population size on
655   inbreeding depression due to mutational load. Genet. Res. 59:49-61.
656
657 Crow JF. 1970. Genetic loads and the cost of natural selection. In: Kojima K, editor. Mathematical
658   Topics in Population Genetics. Berlin: Springer-Verlag. p. 128-177.
659
660 Dabi A, Schrider DR. 2025. Population size rescaling significantly biases outcomes of forward-in-
661   time population genetic simulations. Genetics 229:iyae180.
662
663 Ewens WJ. 2004. Mathematical Population Genetics. 1. Theoretical Introduction. New York:
664   Springer.
665
666 Eyre-Walker A, Keightley PD. 2009. Estimating the rate of adaptive mutations in the presence of
667   slightly deleterious mutations and population size change. Mol. Biol. Evol. 26:2097-2108.
668
669 Ferrari T, Feng S, Zhang X, Mooney J. 2024. Towards simulation optimization: An examination of
670   the impact of scaling on coalescent and forward simulations. bioRxiv 2024.04.27.591463.
671
672 Fincham JRS, Day PR, Radford A. 1979. Fungal Genetics. Berkeley, CA: University of California
673   Press.
674
675 Frankham R. 2023. Effects of genomic homozygosity on total fitness in an invertebrate: lethal
676   equivalent estimates for *Drosophila melanogaster*. Cons. Genet. 24:193-201.
677
678 García-Dominguez S, García C, Quesada H, Caballero A. 2018. Accelerated inbreeding depression
679   suggests synergistic epistasis for deleterious mutations in *Drosophila melanogaster*. Heredity
680   123:709-722.
681
682 García-Dorado A, Hedrick PW. 2023. Some hope and many concerns on the future of the vaquita.
683   Heredity 130:179-182.
684
685 González-Castellano I, Ordás P, Caballero A. 2025. Estimation of inbreeding depression from
686   overdominant loci using molecular markers. Evol. Appl. 18:e70085.
687
688 Haerty W, Ponting CP. 2014. No gene in the genome makes sense except in the light of evolution.
689   Ann. Rev. Genomics Hum. Genet. 15:71-92.
690
691 Haldane JBS. 1937. The effect of variation on fitness. Am. Nat. 71:337-349.
692
693 Halligan DL, Keightley PD. 2006. Ubiquitous selective constraints in the Drosophila genome
694   revealed by a genome-wide sequence comparison. Genome Res. 16:875-884.
695

696 Hawley RS, Gilliland WD. 2006. Sometimes the result is not the answer: The truths and the lies that
697   come from using the complementation test. Genetics 174:5-15.
698

699 Jackson BC, Campos JL, Haddrill PR, Charlesworth B, Zeng K. 2017. Variation in the intensity of
700   selection on codon bias over time causes contrasting patterns of base composition evolution in
701   Drosophila. Genome Biol Evol. 9:102–123.
702

703 Jinks JL. 1983. Biometrical genetics of heterosis. In: Frankel R, editor. Heterosis: Reappraisal of
704   Theory and Practice. Berlin: Springer-Verlag. p. 1-46.
705

706 Johri P, Charlesworth B. 2025. A gene-based model of fitness and its implications for genetic
707   variation: linkage disequilibrium. (*companion paper*).
708

709 Johri P, Charlesworth B, Jensen JD. 2020. Toward an evolutionarily appropriate null model: Jointly
710   inferring demography and purifying selection. Genetics 215:173-192.
711

712 Kamran-Disfani A, Agrawal AF. 2014. Selfing, adaptation and background selection in finite
713   populations. J. Evol. Biol. 27:1360-1371.
714

715 Kardos M, Armstrong EE, Ftzpatrick SW, S. H, Hedrick PW, Miller DE, Tallmon DA, Funk WC.
716   2021. The crucial role of genome-wide genetic variation in conservation. Proc. Natl. Acad. Sci.
717   USA 118:e2104642118.
718

719 Kim BY, Huber CD, Lohmueller KE. 2017. Inference of the distribution of selection coefficients for
720   new nonsynonymous mutations using large samples. Genetics. 206:345–361.
721

722 Kimura M, Maruyama T, Crow JF. 1963. The mutation load in small populations. Genetics 48:1303-
723   1312.
724

725 Kondrashov AS. 1988. Deleterious mutations and the evolution of sexual reproduction. Nature

726   336:435-440.

727 Kyriazis C. C., R. K. Wayne, and K. E. Lohmueller, 2021 Strongly deleterious mutations are
728   a primary determinant of extinction risk due to inbreeding depression. Evolution Letters 5:
729   33–47.
730

731 Kyriazis CC, Robinson JA, Nigenda-Morales SF, Beichman AC, Rojas-Bracho L, Robertson KM, et
732   al. 2023. Models based on best-available information support a low inbreeding load and potential
733   for recovery in the vaquita. Heredity 130:183-187.
734

735 Kyriazis CC, Robinson JA, Lohmueller KE. 2023. Using computational simulations to model
736   deleterious variation and genetic load in natural populations. Am. Nat. 302:737-752.
737

738 Manna F, Martin G, Lenormand T. 2011. Fitness landscapes: An alternative theory for the
739   dominance of mutation. Genetics 189:923-937.
740

741 Matsui T, Mullis MN, Roy KR, Hale JJ, Schell R, Levy SF, Ehrenreich IM. 2022. The interplay of
742   additivity, dominance, and epistasis on fitness in a diploid yeast cross. Nat Commun. 13:1463.
743

McVean GAT, Charlesworth B. 1999. A population genetic model for the evolution of synonymous codon usage: patterns and predictions. Genet. Res. 74:145-158.

Misra S, Crosby MA, Mungall CJ, Matthews BB, Campbell KS, et al. 2002. Annotation of the *Drosophila melanogaster* euchromatic genome: a systematic review. Gen. Biol. 3:Research0083.

Morton NE, Crow JF, Muller HJ. 1956. An estimate of the mutational damage in man from data on consanguineous marriages. Proc. Natl. Acad. Sci. USA 42:855-863.

Nei M. 1968. The frequency distribution of lethal chromosomes in finite populations. Proc. Natl. Acad. Sci. USA 60:517-524.

Pérez-Pereira N, Pouso R, Rus A, López-Cortegano E, García-Dorado A, Quesada H, Caballero A. 2021. Long-term exhaustion of the inbreeding load in *Drosophila melanogaster*. Heredity 127:373-383.

Pérez-Pereira N, Caballero A, García-Dorado A. 2022. Reviewing the consequences of genetic purging on the success of rescue programs. Cons. Genet. 2022:1-17.

Robinson JA, Kyriazis CC, Yuan SC, Lohmueller KE. 2023. Deleterious variation in natural populations and implications for conservation genetics. Annu. Rev. Animal Biosci. 11:93-114.

Roze D. 2021. A simple expression for the strength of selection on recombination generated by interference among mutations. Proc. Natl. Acad. Sci. USA 118:e2022805118.

Sved JA, Ayala FJ. 1970. A population cage test for heterosis in *Drosophila pseudoobscura*. Genetics 66:97-113.

Teixeira JC, Huber CD. 2021. The inflated significance of neutral genetic diversity in conservation genetics. Proc. Natl. Acad. Sci. USA 118:e2015096118.

Wang J, Hill WG, Charlesworth D, Charlesworth B. 1999. Dynamics of inbreeding depression due to deleterious mutations in small populations: mutation parameters and inbreeding rate. Genet. Res. 74:165-178.

Wang Y, McNeil P, Abdulazeez R, Pascual M, Johnston SE, Keightley PD, Obbard DJ. 2023. Variation in mutation, recombination, and transposition rates in *Drosophila melanogaster* and *Drosophila simulans*. Genome Res. 33:587-598.

Wilton AN, Sved JA. 1979. X-chromosomal heterosis in *Drosophila melanogaster*. Genet. Res. 34:303-315.

Wright S. 1968. Evolution and the Genetics of Populations. Vol.1. Genetic and Biometric Foundations. Chicago, IL: University of Chicago Press.

Appendix

1. The load statistics for the two-site deterministic models

Using the fitness model of Table 1, and the notation for the haplotype frequencies described in the main text, the genetic load under the gene model is given by:

$$L_g = \{4zh[x + y(1 + \epsilon)] + 4x[x + \tfrac{3}{2}y(1 + \epsilon)] + 2y^2(1 + \epsilon)\}s \qquad (A1)$$

The inbreeding load is:

$$B_g = 2[x + y(1 + \epsilon)]s - L_g \qquad (A2)$$

and the variance in fitness is:

$$V_g = \{4zh^2[x + y(1 + \epsilon)^2] + 4x[x + \tfrac{9}{4}y(1 + \epsilon)^2] + 4y^2(1 + \epsilon)^2\}s^2 - L_g^2$$

$$(A3)$$

Under the sites model, the corresponding formulae are:

$$L_s = \{4zh[x + y(1 + \epsilon)] + 2x[x + 2xh(1 + \epsilon) + 2y(1 + h)(1 + \epsilon)] + 2y^2(1 + \epsilon)\}s$$

$$(A4)$$

$$B_s = 2[x + y(1 + \epsilon)]s - L_s \qquad (A5)$$

$$V_s = \{4zh^2[x + y(1 + \epsilon)^2] + 2x[x + 4xh^2(1 + \epsilon)^2 + 2y(1 + h)^2(1 + \epsilon)^2] + 4y^2(1 + \epsilon)^2\}s^2 - L_s^2$$

$$(A6)$$

In the case of complete recessivity, Equations (A1) and (A4) for the genetic loads under the gene and sites models without epistasis reduce to the following expressions:

$$L_g = \{4x(x + y) + 2y^2\}s \qquad (A7a)$$

$$L_s = \{2x(x + 2y) + 2y^2\}s \qquad (A7b)$$

Since $y \ll x$, the dominant terms in these equations are $4x^2s$ and $2x^2s$, respectively.

It follows that the load is higher for the gene than the sites mode when $h = 0$. By continuity,

this will also be true when h is sufficiently small, as can be seen in Table 2 for the case of $h =$

0.01.

2. The load statistics for the multi-site deterministic models without epistasis

The formulae for the additive multi-site gene model derived in the last section of the Appendix of

Johri and Charlesworth (2025) can be used to obtain expressions for the load statistics for the case

of a large number (G) of sites with the same s and h, when $N_e s$ is sufficiently large that

deterministic results should apply. We first note that the equilibrium frequency of a deleterious

mutation at a given site ($\hat{q}$) can be found using numerical iteration of Equation (A21a) of Johri and

Charlesworth (2025), with the mean number of deleterious mutations per haploid gene given by

$\mu = G\hat{q}$. Under the assumptions used to derive this equation (including the condition $\hat{q} \ll 1$), the

equilibrium genetic load for the gene model is given by the following expression:

$$L_g = 1 - \bar{w} \approx 1 - w_z = [\,\mu - (1 - 2h)\mu e^{-\mu}\,]s \qquad (A8a)$$

The inbreeding load is given by:

$$B_g = \mu s - L_g \approx (1 - 2h)\mu e^{-\mu}s \qquad (A8b)$$

The variance in fitness contributed by the sum of the variances at individual sites (the genic

variance) is given by:

$$V_g \approx 2G\hat{q}(\bar{w} - w_x)^2 \approx \tfrac{1}{2}\mu[1 + (\mu - 1)(1 - 2h)e^{-\mu}]^2 \qquad (A8c)$$

The corresponding expressions for the sites model can be found from the products of

G with the standard single-locus expressions for deterministic mutation-selection equilibrium

when $\hat{q} \ll 1$ (see Charlesworth and Hughes 2000), on the assumption that LD is negligible.

In this case, $\hat{q} \approx \sqrt{u/s}$ when $h = 0$, and $\hat{q} \approx u/(hs)$ when h is bounded away from zero

(Haldane 1937).

3. Relating population genomics data to the classical expression for inbreeding

depression

Here we attempt to relate inferences from population genomics data on *D. melanogaster* to

the empirical results on the effect on fitness of homozygosity for chromosome 2 of this

species, described in the section of the Discussion *Inbreeding load in Drosophila.* The ratio

of the mean fitnesses of chromosomal homozygotes and heterozygotes under the

multiplicative fitness model with deterministic mutation selection balance for mutations with

$h > 0$ is:

$$R = \exp - n_s \, \mathrm{E}\{qs(1 - 2h)\} \qquad (A9)$$

866 where n_s is the mean number of sites on the chromosome that are subject to significant

867 selection (provisionally identified as nonsynonymous sites in coding sequences), q is the

868 frequency of a deleterious variant at a given nonsynonymous (NS) site, s and h are the

869 selection and dominance coefficients, and $\mathrm{E}\{\}$ denotes the expectation over all NS sites.

870 We have $q \approx u/(hs)$ for strongly selected mutations, so that:

$$R \approx \exp - n_s \, \mathrm{E}\{u[(1/h) - 2]\} \qquad (A10)$$

This is the exponential of the negative of the inbreeding load (B), under mutation-

selection balance with partial recessivity, assuming multiplicative fitnesses across loci

(Morton et al. 1956). This formula applies even when there are significant effects of drift on

q, provided that $2N_ehs$ is sufficiently large and mating is random, since the distribution of q

then follows a gamma distribution with mean $u/(hs)$ and variance $u/[4N_e(hs)^2]$ (Nei 1968).

In section 1 of Supplementary File S2, we present an analysis of relevant population

genomics data on *Drosophila melanogaster,* where it was proposed that 91% of NS sites are

under significant selection (such that $2N_ehs \geq 2.5$); it can be assumed that the remainder are

nearly neutral, with negligible effects on fitness. Given that chromosome 2 is 37% of the

genome, that there are approximately 14000 genes in the fly genome (Misra et al. 2002).

Assuming a mean of 1500 exonic sites per gene and that 70% of exonic sites are NS (Campos

and Charlesworth 2019), n_s for chromosome 2 is 0.37 x 0.91 x 14000 x 0.70 x 1500 = 4.95 x

10^6. With the mutation rate of $u = 4 \times 10^{-9}$ proposed in section 1 of Supplementary File S2,

and the value of $h = 0.25$ suggested by Manna *et al.* (2011), this gives $R = \exp - 0.0396 =$

0.961. If we multiply n_s by three, to include selection on non-coding sites, $R = 0.888$. The

strength of selection is irrelevant to this calculation, so that the mean of $2N_ehs$ could be very

different from that used above but the value of R would be the same.

Variation in h around 0.25 would, of course, increase these estimates. If there were a

two-fold increase in $1/h$, corresponding approximately to a squared coefficient of variation of

h of 1 (as under an exponential distribution), or to a fixed value of $h = 0.125$, then $B = -\ln R$

would be increased to three times that corresponding to the above value, giving $B = 0.356$, R

$= 0.700$ (this is likely to be an underestimate, since small values of h will reduce the

proportion of significantly selected mutations). If the above estimate of the contribution from

major effect mutations is also included, R becomes 0.656, $B = 0.421$.

The remaining question is whether nearly neutral sites, which are ignored in these

calculations, contribute significantly to inbreeding depression. As just described, the effect of

drift away from mutation-selection equilibrium is to reduce B, so that use of the deterministic

formula will overestimate it for nearly neutral sites. If 9% of new mutations fall into this

category, their maximal contribution to B would be $0.09/0.91 = 0.10$ of the values calculated

above, a fairly trivial amount.

4. Relating two models of epistasis

A widely used model of epistasis among deleterious mutations assumes that either fitness or

the logarithm of fitness has a quadratic relation to the number of deleterious mutations (Crow

1970; Charlesworth 1990, 1998). This raises the question of its relation to the pairwise

interaction model used in the present paper, which we now examine. For simplicity, consider

the fitness of homozygotes only, with fixed selective effects across sites. Under the quadratic

model, the fitness of an individual carrying n mutations can be described by a function of the

following form:

$$w_n = 1 - (\alpha n + \beta n^2) \qquad\qquad (A14)$$

Under the pairwise model, we have:

$$w_n = ns + n(n-1)\epsilon s = ns(1-\epsilon) + n^2\epsilon s \qquad\qquad (A15)$$

Equating the coefficients of n and n^2 in these two equations gives the relations

$$\alpha = (1 - \epsilon)s, \quad \beta = \epsilon s \qquad (\epsilon \leq 1) \qquad \qquad \text{(A16)}$$

This result implies that the quadratic model inherently involves pairwise epistasis, at least in

its simplest realization, when only homozygotes are considered. The pairwise model has the

advantage of greater flexibility in modeling the fitnesses of heterozygotes and differences in

selection coefficients between sites, whereas the quadratic model is forced to make somewhat

arbitrary assumptions about the effects of dominance (Charlesworth 1998).

FIGURES AND TABLES

Table 1. Models of the fitness effects of two sites with a fixed selection coefficient

The "sites" model

	++	+ -	- +	- -
+ +	1	$1 - hs$	$1 - hs$	$1 - 2hs - e_1$
- +	$1 - hs$	$1 - 2hs - e_1$	$1 - s$	$1 - s - hs - e_2$
+ -	$1 - hs$	$1 - s$	$1 - 2hs - e_1$	$1 - s - hs - e_2$
- -	$1 - 2hs - e_1$	$1 - s - hs - e_2$	$1 - s - h s - e_2$	$1 - 2s - e_3$

$e_1 = 2\epsilon hs$; $e_2 = \epsilon(1 + h)s$; $e_3 = 2\epsilon s$.

The "gene" model

	+ +	+ -	- +	- -
+ +	1	$1 - hs$	$1 - hs$	$1 - 2hs - e_1$
- +	$1 - hs$	$1 - s$	$1 - s$	$1 - 1.5s - e_2$
+ -	$1 - hs$	$1 - s$	$1 - s$	$1 - 1.5s - e_2$
- -	$1 - 2hs - e_1$	$1 - 1.5s - e_2$	$1 - 1.5s - e_2$	$1 - 2s - e_3$

$e_1 = 2\epsilon hs$; $e_2 = 1.5\epsilon s$; $e_3 = 2\epsilon s$.

Note that the parameter k used in the gene model in Table 1 of Johri and Charlesworth (2025) is here set to ½.

Table 2. Population genetic statistics for the gene and sites models with different values of the dominance coefficient (h) and epistasis coefficient (ϵ)

$h = 0.01$

ϵ	Gene model				Sites model			
	q_r	L	Var	B	q_r	L	Var	B
0	0.768	0.535	0.496	0.0577	1.045	0.548	0.456	0.0821
0.02	0.767	0.535	0.498	0.0577	1.045	0.548	0.458	0.0821
0.04	0.765	0.535	0.500	0.0577	1.043	0.548	0.459	0.0821
0.08	0.763	0.535	0.503	0.0578	1.039	0.548	0.462	0.0821
0.16	0.759	0.535	0.511	0.0578	1.032	0.548	0.468	0.0820
0.32	0.751	0.535	0.525	0.0578	1.021	0.548	0.480	0.0820
0.64	0.742	0.535	0.554	0.0578	1.004	0.548	0.504	0.0818

$h = 0.2$

ϵ	Gene model				Sites model			
	q_r	L	Var	B	q_r	L	Var	B
0	0.919	0.904	0.281	0.716	1.005	0.944	0.241	0.811
0.02	0.919	0.904	0.281	0.716	1.002	0.941	0.242	0.810
0.04	0.918	0.904	0.282	0.716	1.000	0.940	0.242	0.809
0.08	0.916	0.904	0.283	0.716	0.996	0.939	0.243	0.806
0.16	0.914	0.904	0.284	0.716	0.988	0.936	0.246	0.800
0.32	0.910	0.904	0.288	0.716	0.974	0.932	0.250	0.789
0.64	0.904	0.904	0.294	0.716	0.949	0.924	0.262	0.768

$h = 0.5$

ϵ	Gene model				Sites model			
	q_r	L	Var	B	q_r	L	Var	B
0	1.000	1.000	0.480	1.000	1.000	1.000	0.480	1.000
0.02	0.999	1.000	0.481	1.000	0.999	0.999	0.481	1.000
0.04	0.999	1.000	0.481	1.000	0.997	0.999	0.482	1.000
0.08	0.998	1.000	0.481	1.000	0.995	0.998	0.483	1.000
0.16	0.997	1.000	0.482	1.000	0.991	0.997	0.486	1.001
0.32	0.995	1.000	0.484	1.000	0.983	0.994	0.492	1.001
0.64	0.992	1.000	0.487	1.000	0.969	0.988	0.506	1.001

The mutation rate is $u = 0.0001$ and the selection coefficient is $s = 0.01$ ($u/s = 0.01$),

The genetic load L and inbreeding load B are expressed as their ratios with respect to their

single-locus deterministic values when $u << hs$ ($L \approx 4u$ and $B \approx 2u[\,(1/(h) - 2])$ i.e.

when the frequency of A_2 is $u/(hs)$; note that the deterministic value of B for $h = 0.5$ is $B = 0$.

The genetic variance (Var) is expressed as the ratio of hV/s^2 to $4u/s$, to ensure that it lies

between 0 and 1. The equilibrium frequency of deleterious mutations is expressed as its ratio

to the value for a single locus with general u and hs (Equation A4 of Johri and Charlesworth

2025), denoted here by q_r.

Table 3. The load statistics for the gene and sites models with three fixed values of $\gamma = 2Ns$.

1000 selected sites were simulated, with $Nu = 0.005$ and $Nr = 0.01$, where N is the population

size ($N = 1000$), u is the mutation rate per site/generation, and r is the rate of crossing over

between adjacent sites. Details of how to estimate the load statistics are given in the Methods

section. The means and standard errors (SEs) for 1000 replicate simulations are shown. The

predicted values obtained by the methods described in the text are also shown.

$\gamma = 2$		h=0.0		h=0.2		h=0.5	
		Gene model	Sites model	Gene model	Sites model	Gene model	Sites model
genetic load	mean	0.02238	0.01826	0.02260	0.02122	0.02289	0.02273
	SE	0.00012	0.00010	0.00012	0.00011	0.00012	0.00012
	predicted	0.0100	0.00500	0.00100	0.0100	0.01000	0.0100
inbreeding load	mean	0.00000	0.01201	-0.00001	0.00539	-0.00001	0.00000
	SE	0.00001	0.00007	0.00001	0.00004	0.00001	0.00001
	predicted	1.46×10^{-7}	0.0657	2.75×10^{-7}	0.0150	0	0
variance in fitness	mean	2.98×10^{-6}	2.15×10^{-5}	2.84×10^{-6}	5.97×10^{-6}	2.93×10^{-6}	2.98×10^{-6}
	SE	4.16×10^{-8}	2.53×10^{-7}	3.72×10^{-8}	7.97×10^{-8}	4.32×10^{-8}	4.23×10^{-8}
	predicted	5.00×10^{-6}	7.12×10^{-4}	5.00×10^{-6}	2.00×10^{-6}	5.00×10^{-6}	5.00×10^{-6}
allele frequency	mean	0.01100	0.02246	0.01122	0.01639	0.01115	0.01124
	SE	0.00008	0.00015	0.00009	0.00012	0.00008	0.00009
	predicted	0.0100	0.0707	0.0100	0.0250	0.0100	0.0100
$\gamma = 20$ genetic load	mean	0.00563	0.01296	0.00741	0.01140	0.01068	0.01077
	SE	0.00006	0.00016	0.00007	0.00012	0.00009	0.00009
	predicted	0.00632	0.00500	0.00779	0.0100	0.0100	0.0100
inbreeding load	mean	0.00385	0.05479	0.00220	0.01256	0.00001	-0.00004
	SE	0.00004	0.00041	0.00003	0.00010	0.00003	0.00003

	predicted	0.00368	0.219	0.00221	0.0150	0	0
variance in fitness	mean	7.20×10^{-5}	2.89×10^{-4}	5.67×10^{-5}	3.87×10^{-5}	4.88×10^{-5}	4.95×10^{-5}
	SE	7.27×10^{-7}	4.62×10^{-6}	5.52×10^{-7}	4.69×10^{-7}	4.11×10^{-7}	4.56×10^{-7}
	predicted	5.00×10^{-5}	2.74×10^{-4}	5.00×10^{-5}	2.00×10^{-5}	5.00×10^{-5}	5.00×10^{-5}
allele frequency	mean	0.00096	0.00707	0.00098	0.00241	0.00109	0.00107
	SE	0.00001	0.00006	0.00001	0.00002	0.00001	0.00001
	predicted	0.00100	0.0224	0.00100	0.00250	0.00100	0.0010
$\gamma = 100$ genetic load	mean	0.00521	0.00638	0.00788	0.00993	0.01001	0.01002
	SE	0.00006	0.00008	0.00007	0.00007	0.00007	0.00007
	predicted	0.00546	0.00500	0.00808	0.01000	0.0100	0.0100
inbreeding load	mean	0.01255	0.09448	0.00645	0.01423	0.00004	0.00009
	SE	0.00011	0.00067	0.00009	0.00014	0.00007	0.00007
	predicted	0.01257	0.495	0.00655	0.0150	0	0
variance in fitness	mean	3.18×10^{-4}	5.78×10^{-4}	2.15×10^{-4}	1.17×10^{-4}	2.46×10^{-4}	2.48×10^{-4}
	SE	3.45×10^{-6}	8.93×10^{-6}	2.41×10^{-6}	1.08×10^{-6}	1.88×10^{-6}	1.91×10^{-6}
	predicted	1.39×10^{-4}	3.50×10^{-4}	1.71×10^{-4}	1.00×10^{-4}	2.50×10^{-4}	2.50×10^{-4}
allele frequency	mean	0.00036	0.00205	0.00029	0.00049	0.00020	0.00021
	SE	0.00000	0.00002	0.00000	0.00000	0.00000	0.00000
	predicted	0.00036	0.0100	0.00029	0.00050	0.02000	0.0020

Table 4. The load statistics under the gene and sites models for a gamma distribution of
scaled selection coefficients with shape parameter 0.3 and mean $\bar{\gamma} = 2N\bar{s}$. 1000 selected sites
were simulated, with $Nu = 0.005$ and $Nr = 0.01$, where N is the population size ($N = 1000$), u
is the mutation rate per site/generation, and r is the rate of crossing over between adjacent
sites. Details of how to estimate the load statistics are given in the Methods section. The
means and standard errors (SEs) for 1000 replicate simulations are shown.

$\bar{\gamma} = 2$		h=0.0		h=0.2		h=0.5	
		Gene model	Sites model	Gene model	Sites model	Gene model	Sites model
genetic load	mean	0.00976	0.00860	0.00978	0.00951	0.00980	0.00978
	SE	0.00008	0.00008	0.00008	0.00008	0.00008	0.00008
inbreeding load	mean	0.00001	0.00758	0.00001	0.00314	0.00000	0.00002
	SE	0.00001	0.00007	0.00001	0.00003	0.00001	0.00001
variance in fitness	mean	4.33×10^{-6}	1.15×10^{-5}	4.26×10^{-6}	4.21×10^{-6}	4.20×10^{-6}	4.15×10^{-6}
	SE	7.64×10^{-8}	2.31×10^{-7}	7.55×10^{-8}	8.27×10^{-8}	8.02×10^{-8}	6.79×10^{-8}
allele frequency	mean	0.01340	0.01933	0.01347	0.01600	0.01356	0.01352
	SE	0.00011	0.00014	0.00011	0.00012	0.00010	0.00011
$\bar{\gamma} = 20$							
genetic load	mean	0.01098	0.01063	0.01109	0.01157	0.01141	0.01121
	SE	0.00009	0.00010	0.00009	0.00010	0.00009	0.00009
inbreeding load	mean	0.00002	0.02960	-0.00003	0.00714	0.00003	-0.00002
	SE	0.00003	0.00026	0.00003	0.00007	0.00003	0.00003
variance in fitness	mean	4.89×10^{-5}	1.05×10^{-4}	4.85×10^{-5}	2.75×10^{-5}	4.93×10^{-5}	4.84×10^{-5}
	SE	7.22×10^{-7}	2.02×10^{-6}	7.24×10^{-7}	4.53×10^{-7}	7.88×10^{-7}	7.43×10^{-7}
allele frequency	mean	0.00739	0.01231	0.00748	0.00907	0.00741	0.00745
	SE	0.00007	0.00010	0.00007	0.00008	0.00007	0.00007

$\bar{\gamma} = 100$							
genetic load	mean	0.00768	0.00968	0.01026	0.01110	0.01089	0.01087
	SE	0.00009	0.00010	0.00008	0.00009	0.00009	0.00009
inbreeding load	mean	0.00158	0.06449	0.00006	0.00977	-0.00008	-0.00008
	SE	0.00010	0.00062	0.00007	0.00013	0.00006	0.00006
variance in fitness	mean	2.57×10^{-4}	3.78×10^{-4}	2.48×10^{-4}	1.13×10^{-4}	2.43×10^{-4}	2.45×10^{-4}
	SE	4.28×10^{-6}	8.76×10^{-6}	3.96×10^{-6}	1.87×10^{-6}	4.01×10^{-6}	3.85×10^{-6}
allele frequency	mean	0.00298	0.00777	0.00471	0.00568	0.00482	0.00470
	SE	0.00007	0.00007	0.00005	0.00006	0.00005	0.00005
$\bar{\gamma} = 1000$							
genetic load	mean	0.00563	0.00851	0.00828	0.01014	0.01061	0.01066
	SE	0.00008	0.00013	0.00008	0.00008	0.00009	0.00009
inbreeding load	mean	0.01992	0.25320	0.00359	0.01176	-0.00005	0.00035
	SE	0.00059	0.00332	0.00031	0.00035	0.00027	0.00026
variance in fitness	mean	0.00398	0.00754	0.00264	0.00102	0.00345	0.00347
	SE	0.00020	0.00045	0.00011	0.00002	0.00013	0.00012
allele frequency	mean	0.00027	0.00394	0.00091	0.00299	0.00249	0.00260
	SE	0.00000	0.00004	0.00003	0.00004	0.00004	0.00004

958
959

960   **Table 5.** The values of the load statistics obtained from Table 4 with $\bar{\gamma} = 1000$ when scaled

961   up to the whole *D. melanogaster* genome.

Variable	$h = 0$		$h = 0.2$		$h = 0.5$	
	Gene model	Sites model	Gene model	Sites model	Gene model	Sites model
L_T	0.0752	0.112	0.109	0.132	0.138	0.139
B_T	0.279	3.54	0.0503	0.165	0	0
V_T	4.76×10^{-5}	8.34×10^{-5}	2.93×10^{-5}	1.08×10^{-5}	3.59×10^{-5}	3.60×10^{-5}
σ_T	0.00690	0.00913	0.00541	0.00329	0.00628	0.00603

Figure 1: The variance in fitness (Var) and mean deleterious allele frequency ($\bar{q}$) for the

gene and sites models with the *Drosophila*-like simulation parameters, and varying values of

the epistasis coefficient (ϵ). Scaled selection coefficients followed a gamma distribution with

$\bar{\gamma} = 100$ and shape parameter 0.3. 1000 selected sites were simulated with $Nu = 0.005$ and Nr

$= 0.01$, where N is the population size ($N = 1000$), u is the mutation rate per site and r is the

rate of crossing over between adjacent sites. The means and standard errors (SEs) for 1000

replicate simulations are shown.

Figure 2: The means across replicate simulations of the genetic load (L) and inbreeding load

(B) for the gene and sites fitness models, with varying values of the epistasis coefficient (ϵ).

The selection coefficients follow a gamma distribution with $\bar{\gamma} = 100$ and shape parameter

0.3. 1000 selected sites were simulated, with $Nu = 0.005$ and $Nr = 0.01$, where N is the

population size ($N = 1000$), u is the mutation rate per site/generation, and r is the rate of

crossing over between adjacent sites. The error bars represent the SEs across 1000 replicate

simulations.

983