RNA-guided nucleases enable a gene drive of insertion sequences in plasmids

Kepler S. Mears[1,*], Fernando W. Rossine[1,*], Natalia Quinones-Olvera[1], Célia Souque[1], Sophia S.

Wiesenfeld[1], Kesther D. C. Jean[1,2], Shreyas V. Pai[1], Michael Baym[1†]

1. Department of Biomedical Informatics and Laboratory of Systems Pharmacology, Harvard

Medical School, Boston, MA, USA.

2. Roxbury Community College, Boston, MA, USA

*These authors contributed equally

†To whom correspondence should be addressed: baym@hms.harvard.edu

Abstract

Mobile genetic elements (MGEs) are diverse, self-replicating DNA molecules that can

reside within cellular hosts and integrate into one another. This co-occurrence imposes distinct

evolutionary pressures. Plasmids often contain insertion sequences (ISs). However, the multi-

copy nature of plasmids should hinder IS introduction and spread, disfavoring inheritance of

nascent plasmid variants through genetic drift. Mechanisms by which ISs overcome these barriers

remain unidentified. Here we find that the RNA-guided nuclease TnpB enables such a mechanism

to bias its inheritance in plasmids. We show that TnpB, the likely ancestor to Cas12, enables a

gene drive to spread the IS within multicopy plasmids and functions as a primitive anti-self defense

system in conjugative plasmids. The gene drive between TnpB-bearing ISs and plasmids

promotes the spread of both MGEs beyond the ability of either individually. The nested existence

between MGEs is not an incidental result of selfish spread, but a driver of it.

Introduction

Mobile genetic elements (MGEs) are a diverse group of DNA molecules that can replicate

independently of the host chromosome, and therefore face evolutionary pressures of their own

[1,2]. For example, plasmids are circular, extrachromosomal, self-replicating DNA molecules, often

capable of engaging in horizontal gene transfer (HGT) between hosts [3]. In contrast, transposons

(TNs) are linearly integrated segments of DNA that can jump between genomic contexts and

rarely engage in HGT directly [4,5]. Notably, MGEs are a major source of evolutionary novelty in

bacteria: plasmids can transfer large amounts of genetic material between bacterial lineages while

TNs can disrupt genes upon insertion and modulate transcription. Despite their different structures

and functions, MGEs are often found in nested configurations. Plasmids can carry TNs and are

particularly enriched in insertion sequences (ISs) (**Figure 1A**), minimal TNs carrying only the

necessary genes for their transposition and maintenance [4–7]. The intimate relationship between

ISs and plasmids raises the question of what evolutionary processes lead to their association and

whether this relationship drives bacterial genome plasticity beyond the abilities of either MGE

individually.

The integration of ISs into plasmids enhances the evolvability of both MGEs and their

bacterial hosts. ISs enable plasmids to acquire genes by providing recombination homology

regions, diversifying the plasmid gene repertoire [8–10]. Conversely, plasmids allow ISs to mobilize

into different hosts and spread into new genomic contexts [11]. Altogether, these MGE activities

open new evolutionary paths for their bacterial hosts, regardless of any direct fitness effects.
However, this increased evolvability is a second-order selective benefit which may not, by itself,
explain the spread of IS-bearing (IS+) plasmids [12,13]. Importantly, theory predicts that genes that
increase the evolvability of an organism are unlikely to sweep through a population if they are
initially present at a low density [14]. Therefore, the origin of the high prevalence of IS on plasmids
remains unexplained.
IS+ plasmids face distinct challenges spanning many scales of biological organization as
they spread across a bacterial population. First, an IS must transpose into an IS- plasmid, which
is stoichiometrically disfavored by the higher proportion of chromosomal DNA of most bacterial
cells. Second, because plasmids often exist in multiple copies per cell, the nascent IS+ plasmid
is in direct intracellular competition with the IS- copies of the plasmid (**Figure 1B, top**). Moreover,
recent work suggests that the acquisition of an IS might impair the within-cell replicative fitness of
the plasmid [15]. This decreased replication across cell cycles reduces the probability of a newly
divided cell inheriting only IS+ plasmids (i.e. fixing the IS+ plasmid). Finally, any cells that may
have fixed the IS+ plasmid must compete against most of the bacterial population that carries
only the ancestral IS- plasmid. This competition between cells may further disfavor the spread of
the IS+ plasmids, due to a general lack of a host fitness benefit conferred by the IS acquisition.
However, these theoretical challenges to the spread of passenger ISs are in stark contrast with
the high IS prevalence observed in plasmids.
The discrepancy between the high IS+ plasmid prevalence and the obstacles to their
spread can be reconciled since ISs are not passive plasmid passengers. TNs and ISs often
encode systems known to improve their chromosomal retention [16,17], which could also promote
their stability in plasmids. For instance, widely distributed IS families carry TnpBs, which are
programmable, RNA-guided endonucleases. These enzymes can cleave DNA sequences that
lack the IS but that are homologous to where the IS has inserted. In cases where the TnpB
nuclease activity was targeted towards plasmids, TnpB-based systems could favor the fixation
probability of IS+ plasmids (**Figure 1B, bottom**). Generally, many such inheritance-biasing
mechanisms exist in nature and are a fundamental part of gene drives. Yet, to date, no such gene
drive activity, TnpB-dependent or otherwise, has been described for IS+ plasmids.
Here we find both a potential general mechanism to increase IS acquisition by plasmids,
and a specific mechanism for ISs to infiltrate plasmid populations. We find a strong association
between the chromosomal copy number of an IS and residence on plasmids, suggesting that
most IS families rely on high copy numbers to invade plasmids. However, we identify some
families that break the trend and appear to possess mechanisms which allow them other means
of spread. In particular we find that the IS605 family has a specific mechanism enabled by the
TnpB's RNA-guided nuclease that biases plasmid invasion. The IS605 family utilizes RNA-guided
nucleases to enable an intracellular gene drive in multicopy plasmids, promoting the spread of
the IS within the plasmid population. This mechanism additionally enables protection of plasmids
from other external invaders competing for hosts. The unique association between IS605 and
plasmids is the first known instance of a specific mechanism that overcomes the obstacles of
plasmid residency by transposons.
85
86   **Results**
87

Insertion sequence chromosomal copy number is predictive of plasmid carriage

Insertion sequences can exist in multiple copies within a bacterial genome. The number
of identical copies of an IS in a chromosome, the chromosomal copy number (CCN), is highly
flexible and can change as the IS transposes or is lost. More copies of an IS should theoretically
result in more potential transposition events, effectively modulating the IS activity which can be
further regulated by specific IS mechanisms [4,5]. We hypothesized that greater IS activity (i.e.
greater CCN) would allow for more transposition events from the chromosome into plasmids and
therefore result in a greater likelihood of IS co-occurrence (presence of the IS on a plasmid and
the chromosome)
To determine IS co-occurrence we searched 19,658 complete contiguous genomes from
the PLSDB database of plasmids for insertion sequences and determined their replicon location
(chromosome or plasmid) [18,19] (**Supplemental Figure 1A**). ISs were very common in plasmids.
Out of the total 59,895 plasmids within the dataset, 32,641 contained or more IS elements.
Plasmids were also more enriched in ISs having roughly an order of magnitude greater per base
frequency compared to the chromosome (**Figure 1A**). We found IS3 to be the most abundant IS
with over 250,000 instances in the dataset, (**Supplemental Figure 1B**) while IS21 was the most
commonly occurring in genomes with over 90% of all genomes containing one or more instances
(**Figure 1C, top**). The majority of IS elements were identified in the chromosome, aside from IS91
and ISKRA4 which were predominantly located on plasmids (**Figure 1C, bottom**).
IS co-occurrence is positively correlated with IS chromosomal copy number (**Figure 1D**).
For ISs within a genome we determined the chromosomal copy number (number of identical
copies of an IS) and the replicon location (chromosome or plasmid). We then asked whether that
same IS also occurred in any of the plasmids within the genome and classified these events as
IS co-occurrence (**Figure 1D**, dots bottom and top respectively). We calculated the co-occurrence
frequencies at binned chromosomal copy numbers and observed a clear positive association
which was reasonably modeled by a logistic regressor (**Figure 1D**).
The majority of IS families followed a similar trend of positive association between
chromosomal copy number and IS co-occurrence. To summarize the association across IS
families we extracted the rate parameter, β_1, derived from the sigmoid function with $\beta_1<0$ indicating
negative association with chromosomal copy number and $\beta_1>0$ a positive association. As
expected, (**Figure 1D**), the majority of IS families had positive rate parameters (**Figure 1E**) which
could be readily observed in their individual sigmoid fits, (**Supplemental Figure 1C**) with a
handful of outliers having $\beta_1<0$ or $\beta_1\approx0$. A few outliers, notably ISH3 and ISKRA4, were likely due
to the minimal data and abundance of these IS preventing an association being derived with
confidence (**Supplemental Figure 1C**). Increasing chromosomal copy number, and thus IS
activity, is a potentially common method for IS to bias their enrichment in plasmids.

TnpB breaks the positive association between plasmid co-occurrence and IS chromosomal copy
number

Among the associations in different IS families we noticed a difference between two highly
related families IS200 and IS605 [20,21]. The IS200 family are some of the most minimal insertion

sequences containing only a transposase TnpA while the IS605 (and the similar IS607) family
differ by the presence of TnpB, a second gene. The IS200 subfamily followed the general trend
with $\beta_1 > 0$ while IS605 did not with a $\beta_1 \approx 0$. The difference was readily apparent in the respective
sigmoid fits for each with IS (**Figure 1F**). Specifically at low chromosomal copy number IS605
had much higher co-occurrence frequency than IS200 implying that IS605 elements have
increased fixation rate at low chromosomal copy number compared to IS200 (**Supplemental
Figure 1D**). As IS200 and IS605 utilize the same transposition mechanism, we can infer that any
difference in interactions between IS and plasmids is likely due to the activity of TnpB.
The mechanism of TnpB has recently been described to aid in genomic retention of the IS
[22]. The TnpA of IS200/605 elements typically target specific, short (4-6 nucleotide), A/T rich motifs
and transposes via a scarless cut-and-paste mechanism triggered by genome replication [20]. The
result of excision leaves one daughter chromosome without the IS, recreating the original IS
insertion site. This non-replicative mechanism leads to a potential extinction of the IS if
transposition is not accomplished [21]. To prevent this IS loss, IS605 (and IS607) elements use the
RNA-guided nuclease TnpB, the likely ancestor to Cas12. TnpB records the sequence of the right
end flanking region in its guide RNA and in a similar manner to Cas12, utilizes the guide to
programmably cleave specific sequences. Like the PAM of Cas12, TnpB requires an additional
Target Adjacent Motif (TAM) to enable cleavage, that is the same as TnpA's preferred insertion
motif. Upon IS excision the resulting scar brings together the TAM and flanking region to produce
the ideal substrate for TnpB cleavage. The resulting dsDNA break triggers homologous
recombination to copy the IS to the now IS- daughter strand to ensure the IS is never lost from
the specific insertion site, akin to homing endonucleases [22]. Functionally, TnpB records the prior
insertion site of the IS then cleaves any such instance it encounters.
To measure the overall impact of TnpB we calculated the total co-occurrence frequency
of IS605 and IS200 elements. IS605 had double the co-occurrence frequency compared to IS200
(0.022 for IS200 vs 0.054 for IS605) (**Figure 1G**). This increase in co-occurrence persisted when
correcting for multiple potential sources of bias. The PLSDB database is derived from publicly
available genomes on NCBI and is skewed towards *E. coli*, where the majority TnpB occurred
(**Supplemental Figure 2A**). The trend was maintained when eliminating *E. coli* and highly similar
plasmid sequences (**Supplemental Figure 2B**). To isolate potential strain effects, we repeated
the analysis on all genomes from *E. faecium* due to the high density of MGEs and found the same
trend (**Supplemental Figure 2A/B**). The effect was also independent from any evolutionary bias
of TnpA (**Supplemental Figure 2C**). Our results indicate that the addition of TnpB fundamentally
changes the interaction between the IS and plasmids, biasing the fixation rate to increase overall
co-occurrence of the IS in plasmids.

Figure 1: Positive association between chromosomal copy number and plasmid co-occurrence is a potential general mechanism to overcome limits in stochastic fixation for insertion sequences.

A) Per base frequency of insertion sequences in plasmids and chromosomes in the PLSDB database, the dotted line indicates the median. B) Model of purely stochastic IS fixation in plasmids vs that of an IS with increased fixation rate. IS+ plasmids are indicated in blue and IS- in grey C) Prevalence of insertion sequences across genomes in the PLSDB database (top) and distribution across replicon types (bottom). Insertion sequence families are ordered from highest overall

abundance to lowest (Supplemental Figure 1B). D) Co-occurrence frequency of
insertion sequence in plasmids and chromosomes vs IS chromosomal copy
numbers. Dots on the bottom represent an IS found only in the chromosome of a
genome while dots on top represent IS that co-occur in the chromosome and
plasmid. Binned and calculated frequencies at specified chromosomal copy
number are plotted with the expected beta-posterior beta distribution in light blue
and medians indicated with a horizontal line. The sigmoid fit determined from the
raw data is plotted overtop in solid blue. E) Rate parameters from the sigmoid fit
of all IS families analyzed with 95% confidence intervals determined from a chi-
squared approximation from a profile likelihood of the rate parameter. F) Individual
co-occurrence frequency plots with overlaid sigmoid fit for IS200 in blue, IS605 in
red, with corresponding binned frequencies, expected posterior and fit. G) Overall
co-occurrence frequency of plasmid presence for TnpA and TnpA/B IS. 95%
confidence intervals determined from bootstrapping (N=1000) with each replicate
in light blue/red.

TnpB functions as a gene drive to allow IS+ plasmids to overcome stochastic fixation

To examine our observation that TnpB is beneficial for plasmid occupancy we sought to
experimentally measure whether a plasmid containing an active IS605 had an advantage in
replacing IS-negative copies of itself. We developed a plasmid displacement assay where a
mixture of two fluorescently labeled plasmids are electroporated into a test cell line harboring a
non-fluorescent target plasmid (**Figure 2A**) with a validated insertion/excision site from an IS605
[23,24]. By counting the resulting fluorescent colonies, we can measure the relative advantage of
target plasmid replacement by the two labeled plasmids. For example, by adding an IS605 to the
blue plasmid and leaving the red empty we can measure the advantage IS605 provides to a
plasmid in replacing the resident compared to no IS (**Figure 2A**). In parallel, the plasmid mix was
electroporated into plasmid negative cells as a normalization control. From fluorescent images
we counted colonies and compared the ratio of IS+ vs IS- colonies in the test and normalization
plates to determine the log odds ratio (**Supplemental Figure 3A/B**), reflective of the each
plasmids comparative ability to replace the resident (defined as positive if the mWatermelon
plasmid had the advantage and negative if mScarlet did). Introduction of a plasmid containing an
IS605 with a matched insertion site to the target plasmid containing cells mimics a transposition
event into one plasmid within a population, in which one member of an otherwise isogenic
population gains a copy of the IS.
IS605+ ColE1 plasmids displayed a significant advantage displacing the target plasmid
compared to IS605- plasmids, consistent with our bioinformatic predictions (**Figure 2B**). Matched
plasmids, either both IS605+ or IS605-, produced near zero log odds ratio indicating neither
plasmid had an advantage and the assay was functioning as expected (**Figure 2B**). Importantly,
this advantage was not due to a copy number difference between plasmids as all ColE1 variants
had similar plasmid copy numbers from qPCR measurements (**Supplemental Figure 3C**).
The presence of IS605 on its own was detrimental to intracellular plasmid competition. In
a within cell plasmid competition experiment plasmids containing an IS605 were strongly
outcompeted by those without, regardless of the IS location (**Figure 3C**) [15]. The disadvantage is

likely caused by the added transcription imposed by the IS, inhibiting plasmid replication as
recently described [15]. Thus, the advantage seen in our plasmid displacement experiment was not
due to a fitness advantage provided to the plasmid by IS605. Moreover, the results in figure 2B
are significant despite the large detriment of the IS to intracellular plasmid competition.
TnpB's nuclease activity and RNA guide are responsible for the displacement advantage.
Catalytic inactivation of TnpA (dTnpA) did not greatly affect the ability of a plasmid to replace the
target, indicating TnpA, and subsequent transposition activity, is not responsible for the
advantage. TnpB cleaves dsDNA through its RuvC domain which can be inactivated by a point
mutation to create a dead TnpB (dTnpB) [23,25]. Competition of two IS605+ plasmids but with one
harboring a dTnpB resulted in a log-odds favoring the unmodified IS605, thus dTnpB removed
the advantage provided by IS605 (**Figure 2B**). As the RuvC domain is required for DNA cleavage,
the nuclease activity of TnpB is likely driving the observed difference. Moreover, scrambling the
guide region of the IS605 (mutRNA) to prevent recognition of the target sequence produced a
similar result with the defective guide RNA plasmid losing any replacement advantage.
The advantage provided by TnpB is maintained over different multicopy plasmid types.
We repeated the plasmid displacement experiment with the lower copy number pSC101 origin
(~5 copies per cell compared to ColE1's ~30) and produced a similar result with IS605 plasmids
having a displacement advantage compared to IS605- (**Figure 2B)**. A lower copy number would
increase the rate of stochastic fixation for any novel plasmid, thus any advantage given to a low
copy number plasmid in our assay would be predicted to have a lessened effect. We did observe
a diminished magnitude of TnpB advantage with the pSC101 origin confirming that TnpB's effect
is maintained over different replicons and is sensitive to copy number.
As further support that the IS605 plasmids were actively displacing the target plasmid
population we observed diminished fluorescence of IS605- plasmids in test plates. Total
fluorescence in the normalization plates was similar regardless of IS605 presence, suggesting
the IS does not affect plasmid copy number which could account for the observed advantage
(**Supplemental Figure 3D**). In the target plates however, IS605+ plasmids resulted in brighter
colonies than IS605- indicating the proportion of IS605+ fluorescent plasmids expands within the
cell and thus colonies reflective of an increased fixation rate (**Supplemental Figure 3D**).
The results of our displacement experiment are largely insensitive to image thresholding
effects although there exists a slight bias towards mWatermelon (**Figure 2B**, **Supplemental Fig
3E**). We attribute the mWatermelon bias to a greater fluorescence intensity in our equipment, as
exemplified in the double negative condition where the bias is most magnified as brighter colonies
are easier to detect. However, this did not affect the results as the IS+ plasmid conditions were
insensitive to thresholding effects (**Supplemental Fig 3E**).
We conclude that TnpB's RNA-guided nuclease activity enables the effective
displacement of the resident plasmid population and results in an increased fixation rate. We
propose that the programmability of TnpB's guide enables a specific mechanism of the IS to fix
within the plasmid population through a gene drive. When the IS inserts into a plasmid, TnpB will
be reprogrammed to the unique plasmid insertion site. Since plasmids are initially genetically
identical, this insertion site will exist on all other plasmid copies (**Figure 2D**), providing a guide
matched substrate for TnpB cleavage. Cleavage by TnpB will result in destruction of a plasmid
and bias the replication of IS+ plasmids, regardless of any fitness disadvantage the IS may
impose on the plasmid.

The described displacement experiment mimics a transposition event with a target
plasmid designed to specifically match the TnpB guide in the IS. This design allows isolation of
the variables to affect displacement advantage but does not incorporate reprogramming of TnpB's
guide nor the transposition activity of TnpA. To test whether this ability is enabled by TnpB's
programmable RNA guide, we sought to evaluate whether this effect holds when an existing
plasmid gains a new IS.

Figure 2: TnpB enables IS+ plasmids to more efficiently displace IS-plasmids

A) Experimental workflow for plasmid displacement assay and example replicates. Fluorescent images are representations of the fluorescence signal that have been enhanced for visualization. The TAM (yellow) and unique plasmid site (orange) together make up the matched target for TnpB. B) Log-odds ratio of colony counts for ColE1 plasmid displacement assays. mWatermelon plasmid variants are labeled on top and mScarlet on bottom. Individual replicates are plotted with the expected posterior in light grey and 95% confidence intervals. Dotted line is 95% confidence interval from the empty-empty plasmid condition. The right box contains selected data for pSC101 origin plasmids. C) Plasmid competition between IS+ and IS- plasmids. IS+ plasmids are highly outcompeted by IS-regardless of placement on the plasmid. The circle indicates the inoculation line and the arrow the final distance measured. D) Proposed gene drive model for the advantage of IS+ plasmids. Insertion sites/TAM is indicated in yellow with the genomic flanking region in green and plasmid site in orange.

Fixation of novel transpositions into plasmids is enabled by TnpB

To measure the effect of new insertion events in plasmids we devised a transposon trap like those designed to discover novel transposon activity [26]. We integrated an IS607 with validated high levels of transposition [27] and active TnpB into the chromosome along with a resistance cassette under the control of the vanillin operator. We then introduced a target plasmid expressing the vanillin repressor into the cells, resulting in repression of the chromosomal resistance gene and preventing survival on selective media (**Figure 3A**). We designed the repressor coding region to be enriched with potential TnpA insertion sites, i.e. TAMs, setting up a circuit where transposition by TnpA would disrupt the repressor coding region, producing a non-functional repressor (**Figure 3A**). However, so long as a single copy of an intact repressor plasmid exists enough protein will be produced to effectively repress the resistance, preventing survival. Only when a disrupted repressor plasmid fixes within a cell will no repressor be synthesized, enabling survival. By counting colonies on selective plates, we can quantify transposition-and-fixation rates. As mutations in the operator could cheat the circuit, we incorporated fluorescent reporters to mitigate non-IS mediated mutation events (**Supplemental Figure 4 TBS**)

An active IS in the chromosome produced significant survival after incubation and selection with a ColE1 target plasmid in the transposon trap (**Figure 3B, left**). The result of 10 individual replicates followed an expected long tailed Luria-Delbrück distribution, common to similar mutation accumulation experiments [28]. Catalytic inactivation of TnpA resulted in extremely minimal survival, which was expected as the inability to transpose would prevent any disruption of the repressor and completion of the circuit (**Figure 3B, left**). Together these two results indicate the circuit works effectively and any decrease in survival would be indicative of an impaired ability to disrupt the plasmid population.

Inactivation of TnpB resulted in no detectable survival in the ColE1 condition, demonstrating that the programmable nuclease ability of TnpB was critical to fixing the plasmid population (**Figure 3B**). We repeated the experiment with the lower copy pSC101 origin in the

target plasmid and found a similar level of survival with an active IS compared to ColE1. However,
for the dTnpA and dTnpB conditions we observed detectable levels of survival with some
replicates within the range of the fully active condition (**Figure 3B, right**). As with the
superinfection experiment, we expected the low copy number to decrease the relative importance
of the effect of TnpB. The low copy number of pSC101 allows for a greater stochastic advantage
of any novel plasmid, whether that be from an insertion event or a mutation. Some survival in the
dTnpA and dTnpB conditions was likely due to background mutations that we could not eliminate.
There was a slight increase in survival for the dTnpB condition which we did expect to see as the
transposase is still active and able to disrupt the repressor. The pSC101 result demonstrates that
the effect of TnpB is magnified with the increased copy number of the target plasmid. Taken
together, the transposon trap reveals that TnpB improves fixation of newly-introduced IS in
multicopy plasmids, to the point of being effectively required for higher copy plasmids.

327
Figure 3: TnpA/B ISs are captured at higher rates than TnpA ISs in a plasmid
targeted transposon trap
A) Experimental workflow for transposon trap assay. Vanillin repressor in purple,
resistance cassettes in pink, transposon in grey. Yellow indicates potential
insertion sites for TnpA. B) colony counting results for insertion sequence variants
and target plasmid pairs (indicated on bottom). 95% confidence intervals are
indicated with bars around the median from 10 individual replicates as determined
from an expectation maximization of the Luria-Delbrück distribution. Outliers are
indicated with their colony counts above the dotted line.

337
338 *TnpB provides an advantage to intercellular competition of conjugative plasmids*
339
While we find that the strength of the TnpB gene is increased with a higher copy number
(**Figure 2B**), ISs are much more common in larger plasmids, likely due to the increased cargo
capacity (**Supplemental Figure 5A**) [4,5]. We determined the copy number relative to the
chromosome of all plasmids with short read data in the PLSDB dataset and found that indeed

IS200 and IS605 elements tended to be found in large, low copy plasmids (**Supplemental Figure**
5B, Figure 4A). We hypothesized that the location of IS605 elements on plasmids may not be
random as TnpB's targeting mechanism depends on the insertion site; the specific location could
enable targeting or other plasmids. As low copy plasmids are often conjugative, coding for
machinery to enable HGT, we moreover posited that TnpB could provide an intercellular
advantage.
TnpB-containing ISs clustered in conserved plasmid regions. We identified transfer and
mobilization genes (MOB) with MOB-suite to conduct a neighborhood analysis of IS200 vs IS605
elements in plasmids as a proxy for an IS residing in a conserved plasmid region (**Supplemental**
Figure 5C) as MOB genes are highly conserved between plasmids [29] (**Figure 4B**). IS605
elements had a significantly increased total fraction of MOB gene neighbors, strongly suggesting
that IS605 clustered in more conserved plasmid regions, enabling TnpB to target more conserved
genetic sequences (**Figure 4C**). This enrichment held across a range of neighborhood windows
(**Supplemental Figure 5D**) and was not due to any evolutionary bias (**Supplemental Figure**
5E/F). While we did not identify the specific TnpB guides to determine targeting potential, our
results strongly suggest that it is possible for TnpB to target a range of plasmids through residence
in a conserved region.
We created a system to measure competitive invasion advantage of conjugative plasmids.
Similar to the superinfection experiment we integrated fluorescent labels (sfGFP and
mTurquoise2) onto the F plasmid of *E. coli* with either an IS605 or TnpB target site to create
invader strains. We created a recipient strain with a potential TnpB target site within the F plasmid
along with a fluorescent marker and a chromosomally integrated resistance cassette to select for
transconjugants (**Figure 4D**). A mixture of two invader strains was incubated with the target for
24 hours before selecting for transconjugants and taking fluorescent images (**Supplemental**
Figure 6A). At the same time the invader mix was incubated with empty for normalization enabling
quantification of the advantage to invasion granted by the IS through a similar log odds ratio as
the displacement experiment (**Figure 4D, middle**). As a plasmid containing an IS could also be
the target of invasion, we also swapped the resident target plasmid for one containing the IS.
Thus, the assay enables us to measure the defensive advantage an IS grants a conjugative
plasmid in addition to the invasion benefit (**Figure 4D, bottom**).
IS605 provided an advantage to both plasmid invasion and defense. Competition of
matched invader strains (both with or without IS605) produced near zero log odds in both
experimental conditions as expected (**Figure 4E**). IS605 provided a modest, but significant
advantage in plasmid invasion (**Figure 4E, left**), indicating TnpB was clearing the resident target
plasmid to enable invasion. When selecting for superinfected cells, those where the invader and
target plasmid were both present (**Supplementary figure 6B**), we only observed colonies where
the invading TnpB was catalytically inactive, further supporting that TnpB was cleaving the target
plasmid (**Supplementary figure 6C**).
In the defense condition IS605 provided a much larger advantage (**Figure 4E, right**). We
reasoned the difference in magnitude between invasion and defense to be a result of differing
conjugation and transcription kinetics in relation to genetic drift. For the advantage to be realized
in the invasion condition the invader plasmid must enter as ssDNA, re-circularize then transcribe
TnpB, a relatively slow process compared to the genetic drift of low copy plasmids in actively
dividing cells. However, in the defense scenario TnpB is constantly transcribed by the resident

cells, able to immediately cleave the invading plasmid, analogous to the function of CRISPR.
Given recent work on the leading region of conjugative plasmids and TnpB's ability to target
ssDNA, we propose that the orientation and location of the IS could further modulate the strength
of the defensive advantage [25,30].

Figure 4: Invasion and defense of conjugative plasmids is benefitted by the presence of TnpB

A) Copy number vs length of plasmids from PLSDB. Short read sequencing was downloaded and aligned to the reference genomes for each available plasmid. The coverage of the chromosome and each plasmid was used to determine the copy number. B) Neighborhood analysis of IS in plasmids and examples of a conserved (top) and variable (bottom) neighborhood around an IS. For each plasmid in the PLSDB database the transfer and mobilization genes (Tra/MOB genes) are identified by MOB-suite (green). For each gene neighborhood of an IS (indicated by the black arch) the fraction of MOB genes is determined. The closer the IS is to the transfer and mobilization operons, the greater the fraction of MOB genes in the neighborhood and the conservation of IS location. C) Total fraction of MOB neighbors for IS200 and IS605 in the PLSDB dataset. Error bars are 95% confidence intervals determined from N=1000 bootstrapping of the data. D) Experimental set up for conjugation replacement assay. E) Log-odds quantification of invasion (left) and defense (right) conditions from the conjugation invasion assay with the *E. coli* W1485 F plasmid.

Discussion

In this work we uncovered a positive association between IS chromosomal copy number and plasmid co-occurrence across the majority of IS families (**Figure 1D-E, Supplementary Figure 1C**). This association points towards two complementary factors favoring the emergence of IS-bearing plasmids. First, high IS chromosomal copy number might provide greater transposition opportunities resulting in increased transposition rates to plasmids. Second, high IS chromosomal copy number may be indicative of a more permissive cellular environment to transposition, such as decreased regulation. However, a few IS families deviated from this general trend, suggesting mechanisms beyond a numerical advantage to increased plasmid carriage. Plasmid-specific mechanisms of IS invasion could fulfill a powerful role in the spread of their ISs across bacterial populations by facilitating the ability of ISs to rapidly increase their prevalence, and hitchhike on mobile plasmids. In searching for such a mechanism, we focused on the closely related IS200 and IS605 families that displayed distinct patterns of association with plasmids. The co-occurrence of IS200 in plasmids was strongly dependent on high chromosomal copy number. Yet IS605 elements, which are distinguished from IS200 by the presence of single additional protein TnpB, co-occurred in plasmids much more frequently, particularly at low chromosomal copy number, breaking the general trend. Therefore, we hypothesized that TnpB could be enabling a specific mechanism favoring the acquisition or spread of ISs in plasmids.

We found that TnpB facilitates the invasion of plasmid populations via a unique mechanism. The RNA guided nuclease activity of TnpB was originally characterized as promoting chromosomal IS retention by cleaving the excision scar left from transposition [22]. Our work revealed a complementary but distinct role for the programmable ability of TnpB's nuclease activity. In the context of plasmids, a nascent transposition event results in the reprogramming of TnpB's guide to target IS- plasmids, effectively biasing replication and inheritance towards plasmids that are IS+ (**Figure 2B, Figure 3B**). This biased inheritance is the defining feature of a

gene drive. Although usually considered a phenomenon tied to eukaryotic sexual reproduction,
gene drives in principle only require a high ploidy cellular environment. Our findings showcase
that even the simplest instances of high ploidy, such as plasmids, can support naturally occurring
gene drives.

Plasmid gene drives display complexities absent from their eukaryotic counterparts. While
eukaryotic gene drives usually occur in diploids with one chromosome acting as the driver and its
homolog the target, plasmid gene drives occur in truly polyploid environments with multiple
targets. The existence of plasmids in varying copy numbers allows us to investigate the role of
varying target numbers on the gene drive. We found that the higher the plasmid copy number
(i.e. ploidy), the greater the effect of the gene drive (**Figure 2B**), in line with theoretical predictions
on multi-scale evolution [31]. This observation can be understood by the lesser impact of stochastic
fixation with higher copy number and the consequently increased effect of the gene drive.
However, a refinement of our method is required to confirm the exact modulation effect by copy
number, as our results are potentially confounded by protein dilution effects particularly with the
transposon trap. Nevertheless, plasmid copy number is a key factor in the success of the IS
invasion. Overall, the emergence of an IS+ plasmid is not simply the result of a transposition
event, but part of a complex interaction network that is jointly modulated by plasmid ploidy and IS
chromosomal copy number.

The plasmid gene drive mechanism provides a plausible explanation for the broad
distribution of TnpB itself. TnpB is a common protein occurring in roughly 30% of all bacterial
genomes, despite their respective ISs having lower predicted mobility than other IS families[32,33].
The described gene drive could have promoted the spread of TnpB in two distinct scales. First,
by favoring the initial emergence of IS+ plasmids harboring TnpB (**Figure 3**). Second, by providing
IS+ conjugative plasmids with invasion and defense advantages against similar IS- plasmids
competing for the same hosts (**Figure 4**). The enhanced spread of TnpB across genomes has
likely enabled the functional diversification of these proteins as transcriptional modulators[34,35] and
most notably as the Cas12 effector in CRISPR[23,32]. In particular, the defense advantage given to
conjugative plasmids is analogous to the classically described function of CRISPR, providing
protection against other MGEs [32].

Previous work detailing the evolutionary relationship between Cas effectors and
transposon nucleases has found independent instances of guide RNA accumulation, sometimes
even resembling CRISPR arrays [32,33]. These events were proposed as evolutionary bridges
between transposon-bound nucleases and the modern defense CRISPR-Cas systems. Although
these observations demonstrate how the structure of CRISPR-Cas systems might have
originated, they do not explain how such systems gained their defensive functions. Our
conjugation results suggest that at least some defensive functions emerged naturally from the
interaction of ISs and plasmids even prior to the development of arrays. In effect, IS integration
into plasmids and other MGEs might have provided their nucleases with an ancestral evolutionary
path between transposon retention and defense activities.

While TnpB's gene drive mechanism has likely benefitted the spread of TnpB itself, ISs
broadly enable the spread of genes including those outside the bounds of the IS. ISs frequently
serve as recombination hot-spots to allow mobilization of normally immobile elements, enable
adaptation of bacteria to environments, and allow plasmid reorganization of genes through
composite transposons [36–38]. Antimicrobial resistance (AMR) is particularly subject to spread by

plasmids and is heavily modulated by ISs that serve to distribute AMR genes across plasmids
[8,9,39]. Absent from the understanding of the role ISs play in AMR acquisition by plasmids, however,
is how an IS comes to reside on the plasmid in the first place. For AMR to spread through plasmids
via an IS, a plasmid must first acquire the IS itself. The spread of AMR on plasmids is thus a
consequence of IS presence, but not the reason for their existence. TnpB's promotion of IS
incorporation into plasmids is a specific mechanism to achieve IS presence on a plasmid, resulting
in potentially powerful subsequent diversification from incorporation of new genetic material
besides the IS itself.

Plasmids steer bacterial evolution both by direct fitness effects and indirect evolvability
effects. Typically, the spread of genes in bacteria is associated with direct, first-order host-
advantageous traits such as AMR. In contrast, the acquisition of ISs by plasmids is a case where
gene dissemination regularly occurs in the absence of any direct host fitness benefits. Despite
this lack of a first-order host advantage, IS acquisition provides critical second-order benefits to
hosts by enabling gene exchanges across the bacterial pangenome and boosting the evolutionary
flexibility of the host cells. However, it is generally unclear to which extent second-order effects
actively drive the fixation of novel variants in the absence of a first-order benefit, as is the case
for IS-bearing plasmids [12,14]. Here we have shown that within-cell dynamics can drive the fixation
of some classes of IS-bearing plasmids, bypassing the need for first-order fitness benefits and
empowering the evolvability of host cells. This unexpected connection between multi-scale
selection and the evolvability of bacteria highlights the importance of the nonlinearities introduced
by MGE interactions. Crucially, IS-bearing plasmids spread beyond the ability of either MGE
individually. Far from being passive passengers on other MGEs, ISs can encode active
mechanisms to bias their carriage and spread. We argue that the nested existence and
subsequent interactions between MGEs are not an incidental result of selfish spread, but a driver
of it.
506

Acknowledgements
508 We thank Ákos Nyerges and the laboratory of George Church for supplying the MDS42 cells,
509 Arya Kaul, Siân Owen, Suchita Nety and Soumya Kannan for helpful discussions. This work was
510 supported by the NIGMS of the National Institutes of Health (R35GM133700 and R35GM156320),
511 the Pew Charitable Trusts and NSF grant MCB2426105. K.S.M. was supported by an NDSEG
512 Fellowship.

References

1. Horne, T., Orr, V. T. & Hall, J. P. How do interactions between mobile genetic elements

affect horizontal gene transfer? *Curr. Opin. Microbiol.* **73**, 102282 (2023).

2. Lang, A. S., Buchan, A. & Burrus, V. Interactions and evolutionary relationships among

bacterial mobile genetic elements. *Nat. Rev. Microbiol.* **23**, 423–438 (2025).

3. Rodríguez-Beltrán, J., DelaFuente, J., León-Sampedro, R., MacLean, R. C. & San Millán,

Á. Beyond horizontal gene transfer: the role of plasmids in bacterial evolution. *Nat. Rev.*

Microbiol. **19**, 347–359 (2021).

4.   Siguier, P., Gourbeyre, E. & Chandler, M. Bacterial insertion sequences: their genomic

impact and diversity. *FEMS Microbiol. Rev.* **38**, 865–891 (2014).

5.   Siguier, P., Gourbeyre, E., Varani, A., Ton-Hoang, B. & Chandler, M. Everyman's guide to

bacterial insertion sequences. in *Mobile DNA III* 555–590 (ASM Press, Washington, DC,

USA, 2015).

6.   Babakhani, S. & Oloomi, M. Transposons: the agents of antibiotic resistance in bacteria. *J.*

Basic Microbiol. **58**, 905–917 (2018).

7.   Leclercq, S., Gilbert, C. & Cordaux, R. Cargo capacity of phages and plasmids and other

factors influencing horizontal transfers of prokaryote transposable elements. *Mob. Genet.*

Elements **2**, 115–118 (2012).

8.   Cazares, A. *et al.* Pre- and postantibiotic epoch: The historical spread of antimicrobial

resistance. *Science* eadr1522 (2025).

9.   Kaul, A., Souque, C., Holland, M. & Baym, M. Genomic resistance in historical clinical

isolates increased in frequency and mobility after the age of antibiotics. *Microb. Genom.* **11**,

(2025).

10. Cohen, S. N. Transposable genetic elements and plasmid evolution. *Nature* **263**, 731–734

(1976).

11. Loftie-Eaton, W. *et al.* Evolutionary paths that expand Plasmid host-range: Implications for

spread of antibiotic resistance. *Mol. Biol. Evol.* **33**, 885–897 (2016).

12. Woods, R. J. *et al.* Second-Order Selection for Evolvability in a Large *Escherichia coli*

Population. *Science* **331**, 1433–1436 (2011).

13. Tenaillon, O., Taddei, F., Radman, M. & Matic, I. Second-order selection in bacterial

evolution: selection acting on mutation and recombination rates in the course of adaptation.

Res. Microbiol. **152**, 11–16 (2001).

14. Tenaillon, O., Toupance, B., Le Nagard, H., Taddei, F. & Godelle, B. Mutators, population

size, adaptive landscape and the adaptation of asexual populations of bacteria. *Genetics*

152, 485–493 (1999).

15. Rossine, F., Sanchez, C., Eaton, D., Paulsson, J. & Baym, M. Intracellular competition

shapes plasmid population dynamics. *bioRxivorg* (2025) doi:10.1101/2025.02.19.639193.

16. Meers, C. *et al.* Transposon-encoded nucleases use guide RNAs to selfishly bias their

inheritance. *bioRxiv* (2023) doi:10.1101/2023.03.14.532601.

17. Lima-Mendez, G. *et al.* Toxin-Antitoxin Gene Pairs Found in Tn Family Transposons

Appear To Be an Integral Part of the Transposition Module. *mBio* **11**, (2020).

18. Schmartz, G. P. *et al.* PLSDB: advancing a comprehensive database of bacterial plasmids.

Nucleic Acids Res. **50**, D273–D278 (2022).

19. Xie, Z. & Tang, H. ISEScan: automated identification of insertion sequence elements in

prokaryotic genomes. *Bioinformatics* **33**, 3340–3347 (2017).

20. Barabas, O. *et al.* Mechanism of IS200/IS605 family DNA transposases: activation and

transposon-directed target site selection. *Cell* **132**, 208–220 (2008).

21. He, S. *et al.* The IS200/IS605 family and 'Peel and Paste' single-strand transposition

mechanism. *Microbiol. Spectr.* **3**, (2015).

22. Meers, C. *et al.* Transposon-encoded nucleases use guide RNAs to promote their selfish

spread. *Nature* **622**, 863–871 (2023).

23. Karvelis, T. *et al.* Transposon-associated TnpB is a programmable RNA-guided DNA

endonuclease. *Nature* (2021) doi:10.1038/s41586-021-04058-1.

24. Gohil, K., Wu, S.-Y., Takahashi-Yamashiro, K., Shen, Y. & Campbell, R. E. Biosensor

optimization using a FRET pair based on mScarlet red fluorescent protein and an mScarlet-

derived green fluorescent protein. *bioRxiv* (2022) doi:10.1101/2022.06.20.496847.

25. Nety, S. P. *et al.* The Transposon-Encoded Protein TnpB Processes Its Own mRNA into

ωRNA for Guided Nuclease Activity. *CRISPR J* **6**, 232–242 (2023).

26. Tansirichaiya, S., Moyo, S. J., Al-Haroni, M. & Roberts, A. P. Capture of a novel, antibiotic

resistance encoding, mobile genetic element from Escherichia coli using a new entrapment

vector. *J. Appl. Microbiol.* **130**, 832–842 (2021).

27. Žedaveinytė, R. *et al.* Antagonistic conflict between transposon-encoded introns and guide

RNAs. *Science* **385**, eadm8189 (2024).

28. Luria, S. E. & Delbrück, M. Mutations of bacteria from virus sensitivity to virus resistance.

Genetics **28**, 491–511 (1943).

29. Robertson, J. & Nash, J. H. E. MOB-suite: software tools for clustering, reconstruction and

typing of plasmids from draft assemblies. *Microb. Genom.* **4**, (2018).

30. Samuel, B., Mittelman, K., Croitoru, S. Y., Ben Haim, M. & Burstein, D. Diverse anti-

defence systems are encoded in the leading region of plasmids. *Nature* **635**, 186–192

(2024).

31. Cooney, D. B., Rossine, F. W., Morris, D. H. & Levin, S. A. A PDE Model for Protocell

Evolution and the Origin of Chromosomes via Multilevel Selection. *Bull Math Biol* **84**, 109

(2022).

32. Altae-Tran, H. *et al.* Diversity, evolution, and classification of the RNA-guided nucleases

TnpB and Cas12. *Proc. Natl. Acad. Sci. U. S. A.* **120**, e2308224120 (2023).

33. Altae-Tran, H. *et al.* The widespread IS200/IS605 transposon family encodes diverse

programmable RNA-guided endonucleases. *Science* **374**, 57–65 (2021).

34. Wiegand, T. *et al.* TnpB homologues exapted from transposons are RNA-guided

transcription factors. *Nature* **631**, 439–448 (2024).

35. Hoffmann, F. T. *et al.* Exapted CRISPR-Cas12f homologs drive RNA-guided transcription.

bioRxivorg 2025.06.10.658865 (2025) doi:10.1101/2025.06.10.658865.

36. Chen, K., Xie, M., Chan, E. W.-C. & Chen, S. Delineation of ISEcp1 and IS26-mediated

Plasmid fusion processes by MinION single-molecule long-read sequencing. *Front.*

Microbiol. **12**, 796715 (2021).

37. Sastre-Dominguez, J. *et al.* Plasmid-encoded insertion sequences promote rapid

adaptation in clinical enterobacteria. *Nat. Ecol. Evol.* (2024) doi:10.1038/s41559-024-

02523-4.

38. He, S. *et al.* Insertion sequence IS26 reorganizes plasmids in clinically isolated multidrug-

resistant bacteria by replicative transposition. *MBio* **6**, e00762 (2015).

39. Che, Y. *et al.* Conjugative plasmids interact with insertion sequences to shape the

horizontal transfer of antimicrobial resistance genes. *Proc. Natl. Acad. Sci. U. S. A.* **118**,

e2008731118 (2021).

40. Nyerges, A. *et al.* A swapped genetic code prevents viral infections and gene transfer.

Nature **615**, 720–727 (2023).

41. Hamon, A. & Ycart, B. Statistics for the Luria-Delbrück distribution. *Electron. J. Stat.* **6**,

1251–1272 (2012).