Herring roe PLs promote SPM biosynthesis in macrophages and a keratinocyte/fibroblast co-culture as model of psoriasis

Thomas A. Ringheim-Bakka[1,*], Amitis Saliani[2], Tone-Kari K. Østbye[3], Jennifer Mildenberger[4], Matthew Dooley[2], Maftuna Busygina[1], Mona E. Pedersen[3], Nina T. Solberg[3], Jesmond Dalli[2,†], Runhild Gammelsaeter[1,†]

[1]Arctic Bioscience AS, Industrivegen 42, 6155 Ørsta, Norway

[2]William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, Charterhouse Square, London. UK. EC1M 6BQ.

[3]Nofima AS, Osloveien 1, 1433 Ås, Norway

[4]Møreforsking AS, Borgundveien 340, 6009 Ålesund, Norway

[†]These authors share last authorship

* Correspondence:

Thomas Ringheim-Bakka

thomas@arctic-bioscience.com

Short title: Herring roe lipids promotes *in vitro* SPM biosynthesis

Funding sources: This study was supported by a grant from the Norwegian Research Council (project no. 327953).

Abbreviations: CCL2 (chemokine ligand 2), DM (dry matter), DPA (docosapentaenoic acid), FID (flame-ionizing detector), GPCR (G protein-coupled receptor), HDHA (hydroxydocosahexaenoic acid), HDPA (hydroxydocosapentaenoic acid), HEPE (hydroxyeicosapentaenoic acid), HETE (hydroxyeicosatetraenoic acid), HRO (herring roe oil), ICSR (Immune Cell Serum Replacement), IHR (immature herring roe), IL (interleukin), IMID (immune-mediated inflammatory disease), LC-PUFA (long-chain polyunsaturated fatty acid), LLOQ (lower limit of quantification), LPS (lipopolysaccharide), MaR1 (maresin 1), MaR2 (maresin 2), MDM (monocyte-derived macrophages), MRM (multiple reaction monitoring), PASI (Psoriasis Area Severity Index), PBMC (peripheral blood mononuclear cells), PCTR (protectin conjugates in tissue regeneration), PD1 (protectin D1), PDX (protectin DX) PEHeRo (phospholipid esters from herring roe), PL (phospholipid), PLS-DA (partial least-squares discriminant analysis), PMN (polymorphonuclear leukocytes), RvD1 (resolvin D1), RvD2 (resolvin D2), RvD3 (resolvin D3), RvD4 (resolvin D4), RvD5 (resolvin D5), RvD6 (resolvin D6), RvE1 (resolvin E1), RvE2 (resolvin E2), RvE3 (resolvin E3), RvE4 (resolvin E4), RvT1 (resolvin T1), RvT2 (resolvin T2), RvT4 (resolvin T4), SPM (specialized pro-resolving mediators), SPE (solid-phase extraction), TAG (triacylglyceride), VIP (variable importance in projection)

Keywords: drug therapy, inflammation, skin, phospholipids, fish oil, lipid mediator, marine phospholipid esters, herring roe oil, psoriasis

Abstract

Psoriasis is a chronic immune-mediated inflammatory disease (IMID) affecting the skin which presents with both local and systemic inflammation as part of its pathophysiology. An oil rich in phospholipids extracted from herring roe has been shown to have immunomodulatory functions and to improve the clinical symptoms and impact inflammatory cytokine pathways in psoriasis in a clinical trial. The lipidic nature of herring roe oil (HRO) and its high content of polyunsaturated fatty acids suggests involvement of lipid mediator pathways for the observed alleviation of psoriatic inflammation. Of particular interest is the super-family of lipid mediators termed specialized pro-resolving mediators (SPMs), due to their involvement in resolution of inflammation and return to homeostasis. We therefore explored the influence of HRO and its phospholipids on lipid mediator and SPM biosynthesis in IFN-γ and LPS-stimulated human monocyte-derived macrophages and an IL-17A-stimulated keratinocyte/fibroblast co-culture. Lipid mediators including SPMs were quantified from cell supernatants using a validated LC-MS/MS protocol. In these experiments we observed broad SPM biosynthesis with dominant upregulation of RvE2 and RvE3 in both cell systems and upregulation of DHA-derived SPMs such as RvD2 and PDX. Observations of PCTR2 in macrophage cell supernatants also indicate activation of reparative pathways upon treatment with HRO. In conclusion, we observed promotion of SPM biosynthesis associated with a shift towards a protective and possibly reparative macrophage phenotype as well as promotion of biosynthesis of pro-resolving lipid mediators in a skin cell co-culture, thus demonstrating a possible mechanism for resolution of inflammation in the skin niche using HRO.

Introduction

Inflammation is a physiological process that has been described by scholars for centuries, even millennia, where inflammation was famously described two thousand years ago by Aulus Cornelius Celsus (25 BC – 50 AD) by the four cardinal signs of redness, swelling, heat, and pain (1–3). With the progress of science, including growing access to modern powerful analytical instruments, an enormous amount of work has since then been put into expanding the knowledge of initiation and perpetuation of inflammation by describing the actions of cytokines and chemokines, as well as the pro-inflammatory eicosanoids (e.g. prostaglandins and leukotrienes) and their importance in driving inflammatory processes (4–6). And it has not been until the most recent decades that scientists have been able to elucidate the active processes such as the lipid mediator class switch for containing inflammation and restoring tissue homeostasis (4,7). Where before it was thought that resolution of inflammation was a passive process that took place through gradual dilution of chemoattractants, it has now been shown that these are active processes regulated through the expression of a super-family of lipid mediators termed specialized pro-resolving mediators (SPMs) (2,5,6,8). SPMs are potent and transient small molecule signaling agents that are endogenously biosynthesized from polyunsaturated fatty acids (PUFAs) and that, as implied by their name, promote the shift towards resolution of inflammation (4,6). The biosynthesis of SPMs from their parent fatty acids (FAs) is regulated by cell type and phenotype, enzyme expression, and substrate availability (2,9–11). In regulating this vital response, SPMs bind to and activate G protein-coupled receptors (GPCRs) (6,12,13) that result in, amongst other, the reduction of polymorphonuclear leukocyte (PMN) infiltration (6), promotion of efferocytosis/phagocytosis (6), and promotion of tissue regeneration (14,15).

The lack of, or incomplete, resolution of inflammation perpetuates what should have been appropriate and self-limiting inflammatory processes and has been linked to chronic inflammation and multiple disease states (16). A group of chronic inflammatory disorders with unknown etiology that share common inflammatory pathways has been collectively termed as immune-mediated inflammatory diseases (IMIDs) (17). IMIDs are a major public health issue with a global incidence of over 67 million cases (18). IMIDs can affect a number of organs and tissues, including the skin, and can have a profound impact on quality of life, often present with

comorbidities (19–21). In 2019, 6.84% of new IMID cases were psoriasis (18), a chronic inflammatory skin disease which presents with both local and systemic inflammation as part of its pathophysiology. Hippocrates (460 – 377 BC) described the symptoms of psoriasis as *psora* (itch), and *lopoi (*scaling) and recommended treatment with tar followed by sun exposure (22). It is now known that albeit IMIDs display various symptoms depending on the organ they are affecting, IMIDs share an underlying pathogenesis and, indeed, many of the same drugs are therefore used to treat several of these diseases (23,24). Insights in the molecular basis of inflammatory disease has led to therapies targeting specific IMID-associated immune and inflammatory pathways, with recent focus on developing drugs targeting the cytokines IL-17 and IL-23 (25,26). However, due to their intricate involvement in cessation of inflammation responses, imbalances in lipid-mediated regulation of resolution/inflammation due to dysregulated lipid mediator and SPM biosynthesis and signaling have also been proposed to play a role in the pathophysiology of IMIDs (27–29), and it has been shown that intervention with PUFAs (30) or SPMs (31,32) may hold promise in correcting these imbalances. Lack of immunoresolution due to dysregulated SPM biosynthesis may arise both from dysregulated enzyme expression as a consequence of the disease state (33) or substrate competition from pro-inflammatory precursors released from membrane phospholipids (PLs) (34).

Marine PUFA oils have traditionally been considered to have anti-inflammatory properties and have been reported to have potential in treatment of selected IMIDs (35,36). Furthermore, supplementation with marine oils has also been reported to promote SPM and SPM precursor biosynthesis (37–39), where biosynthesis and downstream SPM profiles changes according to different marine oils (39). The difference in conversion between oils implies that the pharmacology may be impacted by other parameters, such as lipid composition and oil source, rather than just the PUFA content/ratios. One marine oil of interest in this regard is herring (*Clupea harengus*) roe oil (HRO), where the nature of the raw material used for extraction causes HRO to have a high content of membrane phospholipid esters enriched with marine PUFAs (PEHeRo = phospholipid esters from herring roe). The benefits of marine PLs have been described previously in literature as having anti-inflammatory effects (40–42) and reports also imply that marine PLs could be superior to other marine

lipid forms with regards to bioavailability and active uptake across the blood-brain barrier (43–47), neuroprotective properties (48), and ability to reduce lesion progress in atherosclerosis (49).

The long-chain PUFAs (LC-PUFAs) provided by PEHeRo in HRO are primarily docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA) in a ratio of approximately 3:1. As reported for other marine PLs, the fatty acids in PEHeRo have also been shown to be readily bioavailable and supplementation has shown reduction of plasma and RBC PC n-6/n-3 ratios (50), as well as a higher acute uptake of DHA and EPA when compared to a triacylglycerol source in a head to head study (51). The bioavailability of lipids and their corresponding fatty acids upon administration is relevant for lipid mediator biosynthesis, as LC-PUFA supplementation has been shown to give increased blood and plasma levels of lipid mediators (52–55). Being able to increase blood levels of lipid mediators through supplementation coincides with observations of positive effects on inflammation through direct supplementation of lipid mediators (33,56,57).

In addition to the general reports on benefits of marine PLs pertaining to inflammation, HRO specifically has been shown to display effect on peripheral inflammation (58) and has been shown to dampen psoriasis *in vivo* in a clinical trial (59–61). The dampening effect of HRO on psoriasis has been shown in human trials where oral daily treatment with HRO gave statistically significant improvement of mean Psoriasis Area Severity Index (PASI) score (59,60). *Post hoc* analyses of blood samples from the study by Tveit *et al.* (59) further supports the beneficial effects of HRO for dampening of psoriasis by showing that HRO had a beneficial effect on immune cell activation and influenced the immune cell profile by, amongst others, reducing CCL2 over time and leading to an increase in IFN-γR1 (61). The *in vitro* experiments performed by Mildenberger *et al.* (62) further corroborates the potential benefit of HRO in the treatment of psoriasis by showing that HRO limits secretion of IL-23 in macrophages in a dose-dependent manner and limits secretion of IL-17 from CD^{4+} T-cells. Thus, showing that HRO positively impacts immune cells and skin cells impacted by psoriasis in multiple ways.

To understand how HRO exerts anti-inflammatory actions, we have studied its impact on lipid mediator biosynthesis in several cell systems. We have studied the effects of HRO on SPM biosynthesis in primary

human monocyte-derived-macrophages (MDMs), as these cells have a role in all inflammatory processes and are known to readily express SPMs as part of their signaling in inflammation regulation. Macrophages are also known to change their phenotype towards more anti-inflammatory phenotypes upon stimulation with lipid mediators (63), which is followed by a shift in lipid mediator biosynthesis. In addition, the effects of HRO on SPM biosynthesis in a skin cell model consisting of a co-culture of IL-17A-treated keratinocytes and fibroblasts, which mimics psoriasis, are studied (64). Current knowledge about SPM biosynthesis in skin cells is limited. Therefore, the study aims to elucidate the role of these cells in SPM and lipid mediator biosynthesis during resolution of inflammation. Studying both the immune cells and skin cells simultaneously may give an image of the cell-cell interplay between these cell types locally in the skin niche. Lastly, we have also mapped the lipid mediator content of HRO and isolated PEHeRo in order to have a greater understanding of which mediators are presented to the cells exogenously compared to the output after treatment, and if any of the identified lipid mediators has described roles in specific interleukin pathways relevant to the pathogenesis of inflammatory diseases including psoriasis.

Materials and methods

Preparation of HRO and PEHeRo

Immature herring roe (IHR) was dried under reduced pressure (until H_2O <20% w/w), then contacted with EtOH/H_2O to extract the lipids. Centrifugation of the resulting suspension yielded a clarified lipid extract which then was concentrated under reduced pressure, desalinated by crystallization in EtOH (abs.), and filtered to give a desalinated phospholipid concentrate. The final HRO was prepared by standardization and viscosity modification through addition of a reconstituted triglyceride oil and evaporation under reduced pressure.

IHR was contacted with EtOH/H_2O to extract lipids before the resulting extract was clarified by centrifugation and dried under reduced pressure. The dried crude extract was then desalinated by crystallization in EtOH

(abs.) and filtration. PEHeRo (>90% PLs w/w dry matter, 85% DM) was purified from the desalinated extract using normal-phase flash column chromatography (EtOH/H_2O 88:12, 200 nm).

Phospholipid analyses were performed using quantitative ^{31}P NMR (Spectral Service AG) and oil fatty acid analyses were performed using GC-FID according to Ph. Eur. 2.4.29 with modifications to accommodate for PL-rich oils (Mylnefield Lipid Analysis).

HRO and PEHeRo emulsions

HRO and PEHeRo were prepared as 5% emulsions in dH_2O by stirring at 800 rpm under inert atmosphere (N_2 flushing) at 37°C for 1 hour. Emulsions were then pasteurized under inert atmosphere by sonication at 70°C for 2 min in an ultrasonic water bath (Branson 2200, 40 kHz) or by an ultrasonic probe. Emulsions were stored at 4°C for maximum 4 days.

Macrophage cell culture

Buffy coats were obtained from the blood bank Ålesund (Helse Møre og Romsdal, Norway) with approval by the Regional Ethics Committee (REK, ref.: 230804) and in compliance with the Helsinki Declaration. Peripheral blood mononuclear cells (PBMC) were isolated from buffy coats by Lymphoprep™ (AXIS-SHIELD PoC AS, 1114547) density gradient and monocytes were selected by plastic adherence and differentiated into monocyte-derived macrophages (MDM) by culture in AIM V medium (Gibco, 12055083) with 5 % CTS™ Immune Cell Serum Replacement (ICSR) and 10 ng/ml M-Csf (Miltenyi Biotec, 130-093-963) for 5 d, followed by incubation without M-Csf.

MDMs from each donor were treated in 1.5 mL per well in a 6-well plate (3 wells per condition) with indicated final percentages (w/w) of HRO or PEHeRo, by dilution of the prepared emulsions in phenol- and serum-free CTS™ AIM-V™ Medium (Gibco, A3830801) supplemented with Pen-Strep. MDMs were either pretreated with HRO for 16 h, followed by change of medium with addition of 1 µg/mL lipopolysaccharide (LPS, Sigma-Aldrich, L2630) and 50 ng/mL IFN-γ (Miltenyi Biotec, 130-096-481) for further 24 h or co-

treated with either HRO or PEHeRo and 1 ng/mL LPS for 24 h. Supernatants were collected, pooled per condition, centrifuged and frozen at -80°C until shipping on dry ice for analysis.

Keratinocyte and fibroblast co-culture

HaCaT keratinocytes (CLS Cell line service GmbH, cat no 300493) were co-cultured with normal human primary dermal fibroblasts (ATCC PCS-201-012) in Dulbecco´s modified Eagle´s medium (DMEM) supplemented with 10% fetal bovine serum (FBS), 100 U/mL penicillin, 100 µg/mL streptomycin and 250 µg/mL fungizone (all purchased from Thermo Fisher Scientific) in T25 tissue culture flasks (Nunc™ EasYFlask™, VWR) and stimulated with and without 100 ng/mL IL-17A (Sigma-Aldrich, cat.no SRP0675) and 5 µg/mL HRO for 96 h, before media were collected and stored at -80 ^{o}C for further fatty acid and lipid mediator analysis. The co-culture was established by first seeding fibroblasts at $1x10^3$ cells per cm^2 and pre-cultured in the media described above for 2-days, before adding $4,3 x10^3$ keratinocytes per cm^2 and co-culturing additional 5 days before initiating stimulation with HRO with and without IL-17A for 96 h as described above.

Fatty acid composition of cells

Total lipids were extracted from the cells according to the protocol by Folch *et al*. (65). The $CHCl_3$ phase was dried under nitrogen gas, and thereafter the samples were methylated over night with benzene, 2% H_2SO_4 in MeOH, and dimethoxypropanate at 50 °C. The methyl esters of the fatty acids were separated using GC (Hewlett Packard 6890) with flame-ionizing detector (FID) using a BPX70 column (60 m, 0.25 mm i.d., 0.25 µm film) with a temperature program starting at 50 °C, and increasing at a rate of 4 °C /min until 170 °C, then raised to 0.5 °C/min until 200°C, and finally to 10 °C/min until 240 °C. The peaks were integrated using GC ChemStation software (Agilent Technologies, Rev. B.01.01 SR1, 2001-2005). The fatty acids were identified by an external standard (GLC 463, Nu-Chek Prep), and the concentration of the individual fatty acids was calculated based on an internal standard (C23:0, Larodan). For analysis of fatty acid composition in lipid classes, the $CHCl_3$ lipid extracts were first evaporated under nitrogen gas and re-dissolved in hexane. The lipid classes (phospholipids, triacylglycerides and free fatty acids) were separated by thin layer chromatography

using a mixture of petroleum ether, Et$_2$O and AcOH (113:20:2, by volume) as the mobile phase. The lipids were visualized by spraying the TLC plates with 0.2% (w/v) 2′,7′-dichlorofluorescein in MeOH, and then identified by comparing with known standards (Sigma Chemical Co.) under UV light. The spots corresponding to the different fractions were scraped off into glass tubes, trans-methylated following the aforementioned procedure and the fatty acid composition determined by GC-FID.

Lipid mediator analyses

Lipid mediator analysis was performed using methodologies and approaches recently published by Dooley *et al.* (66).

Snap frozen media was placed in 2 volumes of MeOH containing deuterated internal standards (IS) (d$_4$-LTB$_4$, d$_5$-MaR1, d$_5$-MaR2, d$_4$-PGE$_2$, d$_5$-LXA$_4$, d$_5$-RvD$_3$, d$_5$-RvD2, d$_4$-RvE1, d$_5$-17R-RvD1, d$_5$-LTE$_4$, d$_5$-LTD$_4$ and d$_5$-LTC$_4$), representing the chromatographic regions of interest, were added to facilitate lipid mediator identification and quantification. This was placed at -20 $^{\circ}$C for at least 45 minutes then centrifuged and supernatants were subjected to solid-phase extraction (SPE) using the ExtraHera system (Biotage) and Isolute C18 500 mg columns (Biotage).

Oils samples were placed in 2 volumes of MeOH containing 2.5 ng of deuterated IS (d$_4$-LTB$_4$, d$_5$-MaR1, d$_5$-MaR2, d$_4$-PGE$_2$, d$_5$-LXA$_4$, d$_5$-RvD$_3$, d$_5$-RvD2, d$_4$-RvE1, d$_5$-17R-RvD1, d$_5$-LTE$_4$, d$_5$-LTD$_4$ and d$_5$-LTC$_4$) and incubated in 2 volumes of KOH (1N) for 45 minutes at 45 $^{\circ}$C. At the end of the incubation the samples were immediately acidified and subjected to SPE using the ExtraHera system (Biotage) and Isolute C18 500 mg columns (Biotage). Oils analysed for unesterified lipid mediators were not saponified prior to being subjected to SPE.

Methyl formate and MeOH fractions from C18 SPE were collected, brought to dryness, and suspended in phase (MeOH/H$_2$O, 1:1. v/v) for injection on a Shimadzu LC-20AD HPLC and a Shimadzu SIL-20AC autoinjector, paired with a QTrap 6500+ (Sciex). Analysis of mediators isolated in the methyl formate fraction was conducted as follows: an Agilent Poroshell 120 EC-C18 column (100 mm × 4.6 mm × 2.7 μm) was kept at 50 °C and mediators eluted using a mobile phase consisting of MeOH/H$_2$O/AcOH of 20:80:0.01

(v/v/v) that was ramped to 50:50:0.01 (v/v/v) over 0.5 min and then to 80:20:0.01 (v/v/v) from 2 min to

11 min, maintained till 14.5 min and then rapidly ramped to 98:2:0.01 (v/v/v) for the next 0.1 min. This

was subsequently maintained at 98:2:0.01 (v/v/v) for 5.4 min, and the flow rate was maintained at

0.5 mL/min. In the analysis of mediators isolated in the MeOH fraction, the initial mobile phase was

MeOH/H_2O/AcOH of 20:80:0.5 (v/v/v) which was ramped to 55:45:0.5 (v/v/v) over 0.2 min and then to

70:30:0.5 (v/v/v) over 5 min and then ramped to 80:20:0.5 (v/v/v) for the next 2 min. The mobile phase was

maintained for 3 min and ramped to 98:2:0.5 (v/v/v) for 2 min. QTrap 6500+ was operated using a multiple

reaction monitoring (MRM) method. Each lipid mediator was identified using the following criteria: (1)

matching retention time to synthetic or authentic standards (±0.05 min), (2) a signal-to-noise ratio ≥ 5 for a

primary transition, and (3) a signal-to-noise ratio of ≥3 for a secondary transition. Data was analyzed using

validated protocols (66) and Sciex OS v3.0. Chromatograms were reviewed using the AutoPeak algorithm,

using 'low' smoothing setting and signal to noise ratios were calculated using the relative noise algorithm

(66). External calibration curves were used to quantify identified mediators. Where available calibration

curves were obtained for each mediator using synthetic compound mixtures that gave linear calibration curves

with R^2 values of 0.98–0.99. These calibration curves were then used to calculate the abundance of each

mediator. Where synthetic standards were not available for the construction of calibration curves, calibration

curves for mediators with similar physical properties (e.g. carbon chain length, number of double bonds,

number of hydroxyl groups and similar elution time) were used.

Statistical analysis

Statistical significances were determined using; the Mann-Whitney U test (one-tailed/two-tailed as

appropriate) for comparison of two groups, one-factor ANOVA and one-way/two-way Dunnett's multiples

comparison test or Tukey's HSD for comparison of >2 groups. Statistical significance criteria for difference

between groups was $P < 0.05$. All results are presented as mean values ± the standard error of the mean

(SEM). PLS-DA analyses were performed using MetaboAnalyst v6.0 (67), all other statistical analyses were

performed using Microsoft Excel for Microsoft 365 MSO.

Results

Oil composition and lipid mediator content in HRO and PEHeRo

In order to understand the effects of phospholipid ester rich oils on cellular systems, it is important to understand the composition of the oils both with regards to fatty acid profile and lipid class distribution, but also with regards to lipid mediators potentially present in the oil. We therefore started off with characterizing the composition of the oils used.

Herring roe oil (HRO) is a lipid extract from herring roe that has been standardized with regards to content of *phospholipid esters from herring roe* (PEHeRo) using a marine triacylglyceride (TAG) oil. The phospholipid esters in PEHeRo are predominantly phosphatidylcholine (>80%) that are enriched with LC-PUFAs. The fatty acid profiles in both HRO and PEHeRo display a high content of DHA with a DHA/EPA ratio of approximately 3:1, where DHA and EPA make up 50-60% of the total fatty acid profile in area %. The HRO and PEHeRo investigated in this work had a phospholipid content of 34% and 91% (per anhydrous substance) and 425 mgTAG/g and 318 mgTAG/g DHA+EPA, respectively. Lastly, HRO and PEHeRo were found to contain 6% area and 1% area of omega-3 docosapentaenoic acid (n-3 DPA), respectively.

After establishing the macroscopic lipid composition, the content of lipid mediators in HRO and PEHeRo was determined, as it is possible that the lipid mediator levels in the oils could influence the behavior and SPM biosynthesis in the cell systems. The oils were screened for SPMs and monohydroxylated fatty acids (including SPM precursors) originating from DHA, n-3 DPA, EPA, and arachidonic acid (AA). The qualitative lipid mediator profiles of the two oils were, unsurprisingly, found to be similar, with the predominant SPMs originating mostly from the DHA metabolome in form of RvD5, RvD6, PDX, 17R-PD1, MaR1, and MaR2 (Fig. 1A). The most similar relative abundance between HRO and PEHeRo was found for 18-HEPE, RvD5, MaR2, and RvD5 $_{\text{n-3 DPA}}$ where the groups rank sums indicated a $P > 0.1$ (two-way) as denoted with # in (Fig 1 A-C). Notable differences between the oils included a relative lower abundance of PDX and higher relative abundance of MaR1 and RvE4 in PEHeRo compared to HRO. Furthermore, there

were also notable contributions from SPMs originating from the n-3 DPA (RvT4 and RvD5 $_{\text{n-3 DPA}}$) and EPA (RvE4) metabolomes, whereas lipid mediators from the AA metabolome, as expected, only made up 1-2% of the total SPM content in the oils.

The key difference between HRO and PEHeRo pertaining to SPM content, was the >10 times higher concentration of overall SPMs in the less refined HRO compared to the concentrated phospholipid oil PEHeRo. The same was also observed for the monohydroxylated fatty acid levels in the two oils.

Furthermore, the monohydroxylated fatty acid (HDHAs, HEPEs, HDPAs, and HETEs) content was found to be surprisingly high in HEPEs given the overall fatty acid composition, with 5-HEPE and 18-HEPE being the two major contributors (Fig. 1B and Fig. 1C). The HEPE/HDHA ratio was found to be 1.8 and 1.4 for HRO and PEHeRo respectively (Fig. 1B), where a significant portion (>50%) of total HEPEs and HDHAs were found to be present as metabolically available free fatty acids not bound in lipids. The sum of HEPEs and HDHAs made up >95% of the total amount of monohydroxylated fatty acids in the two oils. Lastly, we found higher levels of unesterified EPA in both HRO and PEHeRo compared to DHA free fatty acid.

Supplemented HRO is taken up by macrophages and keratinocytes/fibroblasts

The uptake of fatty acids from HRO was analyzed by GC-FID, showing a significant increase in DHA, n-3 DPA, and EPA in total lipids in the MDMs (Fig. 2A and supplementary table S1). In the direct keratinocyte/fibroblast co-culture, it was further shown that the changes introduced by HRO could be observed in the phospholipid fractions of the cells with similarly significantly increased EPA and DHA and decreased AA (Fig. 2B and supplementary table S2).

HRO and PEHeRo promote SPM biosynthesis in monocyte-derived macrophages

The macrophage experiments conducted by Mildenberger *et al.* demonstrated reduction in cytokine secretion, and we therefore wanted to study SPM biosynthesis in the macrophages treated with HRO in order to better understand how this oil and its phospholipids (PEHeRo) elicits their anti-inflammatory and possibly proresolving activities (62).

We therefore quantified the SPMs present in the supernatants from likewise stimulated MDMs, which showed a significant upregulation of SPM concentrations when macrophages were pre-treated with HRO prior to stimulation with LPS+IFN-γ (Fig. 3A-D and supplementary table S3). The biosynthesis was found to be significantly upregulated for SPMs from the DHA, EPA, and AA metabolomes, including a non-significant trend for upregulation of n-3 DPA SPMs. The EPA SPM RvE3 was found in highest concentration in the treated samples. The overall SPM profiles of the treated macrophages showed activation of both ALOX5 and ALOX15 pathways.

Partial least squares discriminant analysis (PLS-DA) of the SPMs in the cell supernatants from the treated and untreated macrophages (including a fresh media control sample) as shown in Fig 2C also displayed a difference between the groups. This shift in SPM concentrations indicated by the separation of the clusters in green (2, untreated) and blue (3, treated with HRO) in the PLS-DA scores plot. The chart shown in Fig. 3D shows the variable importance in projection (VIP) scores of individual SPMs that contribute to the separation of the clusters in the PLS-DA with a VIP score >1 identifying those mediators that contribute to the observed separation to the greatest extent. VIP scores >1 were observed for 4 SPMs from the DHA metabolome (RvD1, RvD4, PD1, and PDX) and 2 SPMs from the n-3 DPA metabolome (RvD5$_{\text{n-3 DPA}}$ and PD1$_{\text{n-3 DPA}}$).

Considering the promotion of SPM biosynthesis, we observed in the macrophages when pretreated with HRO, we wanted to further explore this and see if we were able to optimize the experiment setup with regards to measured concentrations of SPMs in the supernatant after treatment/stimulation. Also, considering that we this far have only looked at the complete HRO, we also wanted to evaluate if there were any differences between HRO and a purified sample of PEHeRo in terms of ability to promote SPM biosynthesis in primary immune cells. So, we tested two concentrations of HRO (0.05% and 0.25%) and one concentration of PEHeRo (0.25%) in an experiment where the macrophages were co-stimulated with the oils and 1.0 ng/mL LPS for 24 hours before SPM analysis of the cell supernatants, similarly to the conditions employed by Sobrino *et al.* (39).

Similarly to the initial experiment, we also observed a significant upregulation of SPM biosynthesis in the macrophages treated with HRO and PEHeRo (Fig. 4A-D and supplementary table S4). The results also

showed, as expected, generally higher concentrations of SPMs in the cell supernatants, thus offering a potential for greater understanding of the pathways activated upon treatment with HRO and PEHeRo.

All three treatments conditions (0.05% HRO, 0.25% HRO, and 0.25% PEHeRo) gave significant upregulation of the sum of SPMs from the DHA and EPA metabolomes (Fig. 4A-B) when compared to media stimulated with LPS (control). We also observed a nonsignificant trend for upregulation of SPMs from the DPA metabolome and, in contrast to the first experiment, no upregulation of AA SPMs. Similarly to the first experiments, the most strongly expressed SPM was the EPA-derived SPM RvE3.When comparing the two treatments concentrations of HRO (0.05% and 0.25%), we observed a nonsignificant dose-dependent trend where the higher treatment concentration gave higher mean values than the lower concentration. The macrophage SPM biosynthetic response to the treatment did not reflect the 5-times increase in treatment concentration and could indicate saturation of the system with regards to accessible fatty acids for SPM biosynthesis.

Looking at individual SPMs through the PLS-DA analysis from the experiments, we also see here that PEHeRo elicits a different response from the treated macrophages compared to the two concentrations of HRO. This is well visualized through the clear separation of the cluster for PEHeRo and the two HRO clusters in Fig. 4C. The partial overlap of the two different doses of HRO is in alignment with the observation made for the abundance of SPMs from the different metabolomes, where a nonsignificant trend for dose-dependency was observed. The VIP scores shown in Fig. 4D indicate which individual SPMs are drivers for the separation of the clusters similarly to Fig. 3C. The VIP scoring shows also here that RvD1 is the biggest driver for separation of the clusters, followed by RvE3 which was found to be the most abundant SPM present in both experiments. Furthermore, we see that individual SPMs from all four metabolomes (DHA, n-3 DPA, EPA, and AA) contribute to the separation of the different clusters. Despite not observing significant upregulation of total n-3 DPA SPMs between the groups, the individual n-3 DPA SPMs with a VIP-score >1 were found to be significantly upregulated: $RvD1_{\text{n-3 DPA}}$ for HRO and PEHeRo at 0.25%, RvT2 for PEHeRo at 0.25%, and RvT1 for both concentrations of HRO, thus corroborating their significance in the separation of the clusters in the PLS-DA analysis.

Interestingly, the VIP scoring in Fig. 4D also shows the cysteinyl DHA-derived SPM PCTR2 (PCTR = protectin conjugates in tissue regeneration) from the DHA metabolome to be a significant contributor to the separation of the different clusters, thus indicating activation of reparative pathways and not only protective SPM pathways.

HRO promotes SPM biosynthesis in IL-17A stimulated fibroblast/keratinocyte co-culture

To study the potential effects of HRO on the IL-23/IL-17 cytokine axis in psoriasis, a co-culture of keratinocytes and fibroblasts that mimics psoriasis upon stimulation with IL-17A was developed. Mildenberger *et al.* (62) showed that HRO dampens psoriatic marker S100A7 (psoriasine) and the IL-17 response gene NFKBIZ in this co-culture. Considering the dampening effects observed in these co-culture experiments, we wanted to see if the SPM biosynthesis was upregulated from the co-culture in this *in vitro* psoriasis-mimicking cell model.

Quantification of SPMs and monohydroxylated fatty acids in the co-culture cell media showed significant upregulation of SPMs from the DHA and EPA metabolomes, as well as significant upregulation of monohydroxylated DHA, EPA, and n-3 DPA (Fig. 5A-D and supplementary table S5). It was encouraging to see a significant promotion of SPM biosynthesis taking place despite only treating the cells with 1/100 (0.0005%) of the amount of HRO compared to the 0.05% HRO used in the macrophages and reporting on 1/3 of the amount of media (3 mL vs 1 mL). The amounts measured in the media were of course affected by the decreased concentration of HRO, and the overall quantities are therefore lower compared to what was observed in the macrophage experiments. It should be noted that a parallel co-culture experiment (n=6) showed similar results albeit with somewhat more moderate levels of lipid mediators. This experiment did not show significant upregulation of sum DHA-derived SPMs (only PD1 in the HRO group). Nor was a significant increase in the sum of HDPAs observed, only a significant increase of 17-HDPA was observed compared to the stimulated control (see supplementary table S6 for more information).

The DHA SPMs that contributed to the significant upregulation of sum of DHA SPMs (Fig. 5A) in both the groups treated with HRO included RvD1, RvD2, RvD3, RvD5, RvD6, and PD1, of which RvD5 and RvD6

were significant in HRO+IL-17A compared to the stimulated control. The contributors to the overall activation of the DHA SPM metabolome coincide with observations made in the macrophage experiments where RvD1 and PD1 had VIP scores >1 when HRO was used to pretreat macrophages prior to stimulation with LPS+IFN-γ and RvD1 and RvD2 had VIP scores >1 when macrophages were costimulated with HRO/PEHeRo + LPS. Significant upregulation of monohydroxylated fatty acids (HDHAs and HDPAs) also could indicate that the HRO indeed promotes activation of the metabolic pathways for SPM biosynthesis from the DHA and DPA metabolomes. The HDHA profile measured in the media supports the observations made of the SPMs (Fig. 5C), as the main monohydroxylated fatty acid found was 17-HDHA where the 17S-HDHA isomer is a precursor for the D-series resolvins formed through ALOX15. 17-HDHA was significantly favored over 14-HDHA where the 14S isomer is a pathway marker for the biosynthesis of MaR1 and MaR2 through ALOX12. The apparent preference for 17-HDHA thus correspond with the observed levels of MaR1 and MaR2 where neither of which were measured above the LLOQ. The same also being the case for the experiments with the macrophages where no consistent or significant measurements of MaR1 and MaR2 were made. It should be noted that this was more pronounced for the HDPAs, as only 17-HDPA was measured in the media samples.

Furthermore, SPMs from the EPA metabolome were also significantly increased in both treatment groups compared to the stimulated control (Fig. 5B). This was primarily driven by RvE3 and to a lesser extent RvE2 (approximately 1:40 RvE2/RvE3), both of which were also separately significantly upregulated compared to the stimulated control. RvE2 and RvE3 being the main EPA SPMs measured is not surprising when looking at the HEPE profile of both treatment groups (Fig. 5D), where 18-HEPE was found in the, by far, highest concentration in the media. The 18R isomer of 18-HEPE is a SPM precursor biosynthesized via CYP450 and GPX4 and is the main intermediate before a one-step conversion via ALOX15 into RvE3 or the conversion to RvE2 via ALOX5. The abundance of 15-HEPE was far lower than the abundance of 18-HEPE, which may explain why RvE4 was not found above the LLOQ as 15S-HEPE is the main precursor for RvE4.

Including a group of "healthy" cells only treated with HRO shows, that there may be an impact to the lipid mediator biosynthesis when the cells are stimulated with IL-17A. The greatest impact of IL-17A stimulation was observed for the EPA metabolome, with a significant reduction of 18-HEPE, sum HEPEs, and EPA SPMs

when comparing the HRO group with the HRO+IL-17A group. Interestingly, only RvE3 was significantly reduced, whereas only a nonsignificant trend was seen for RvE2. An impact of IL-17A stimulation was also observed for SPMs in the DHA metabolome, albeit to a lesser extent. Here we observed that both RvD2 and PD1 were significantly upregulated in the HRO group, but not in the HRO+IL-17A group. The negating effect of IL-17A on lipid mediator biosynthesis was only observed in one out of two data sets, as can be seen from Table S6 in the supplementary data.

Discussion

An interesting finding from the macrophage experiments in the present study was the comparison of total SPMs across different metabolomes. Where notably, treatment with PEHeRo significantly increased the biosynthesis of SPMs from the DHA metabolome compared to HRO. In addition to the significant difference between PEHeRo and HRO in the DHA metabolome, a similar observation was made for SPMs from the EPA metabolome albeit as a nonsignificant trend. The observed increase in SPM biosynthesis in macrophages treated with PEHeRo cannot be explained by availability of metabolically available free fatty acid precursors or availability of mono-hydroxylated fatty acids, including lipid mediator precursors, as PEHeRo was shown to contain far less of these compounds compared to the HRO used in these experiments. Thus, any normalization based on availability of free EPA/DHA, bound EPA/DHA, or monohydroxylated fatty acids (18-HEPE and 17-HDHA) only increases the difference observed when comparing HRO and PEHeRo based on overall concentration. This indicates that the more purified PEHeRo more efficiently promotes upregulation of biosynthesis and that the oils' ability to upregulate these pathways depends on other factors such as matrix complexity or lipid class composition and not just the precursor availability. For instance, phospholipids have been indicated to give increased *in vivo* bioavailability of PUFAs compared to corresponding triacylglycerides (51,68), and their lysoforms may also themselves act as signaling molecules (69). The observations correspond with existing data that indicates that different oils elicit different ability to promote conversion of PUFAs and

SPM precursors into SPMs (39). Pinpointing the impacting factors for these observations needs further studies to be verified.

Consistent dominant upregulation of the EPA-derived SPMs RvE2 and RvE3 was observed in all experiments, with RvE3 being the most abundant of the two. The higher relative concentration of RvE3 compared to other SPMs can possibly be attributed to RvE3 being available in one enzymatic transformation from 18R-HEPE by ALOX15, which was likely abundant in both HRO and PEHeRo from the measured levels of 18-HEPE (achiral analysis). RvE2 is also biosynthesized from 18R-HEPE but through two steps by ALOX5 and a dehydrogenase. The availability of 18-HEPE in the oils does not fully explain the strong upregulation of RvE3, as more RvE3 was found in cell media when the macrophages were stimulated with PEHeRo which contained >10x less 18-HEPE than HRO. The consistent upregulation of RvE2 and RvE3 across both the macrophages and the skin cell co-culture experiments is particularly interesting given that RvE2 has been reported to be a regulator of neutrophil infiltration in addition to enhancing phagocytosis (70), and more importantly, RvE3 has been reported to attenuate inflammation via the IL-23/IL-17A-axis (71) which is highly relevant for psoriatic inflammation.

The most recurring DHA-derived SPM was found to be RvD2, which was observed consistently in the macrophage experiments and for the HRO treated group in the keratinocyte/fibroblast co-culture experiments. RvD2 takes part in regulating inflammation by increasing phagocytosis and reducing neutrophil infiltration (72), promoting macrophage polarization (73), and reducing expression of proinflammatory cytokines (72). The protectin PDX has been reported to be implicated in regulation of psoriasis, as it has been found to be elevated in psoriasis lesional skin (74), and was measured in all the macrophage experiments. By being upregulated in the immune cells treated with HRO and PEHeRo and involved in regulation of inflammation in psoriasis, PDX could be a promising biomarker for future studies in psoriasis using marine phospholipids.

Showing that we were able to upregulate SPM biosynthesis in the keratinocyte/fibroblast co-culture was intriguing, considering that there, to our knowledge, are few studies detailing the ability of lipid supplementation to regulate SPM production in these cells. The monohydroxylated fatty acid profiles could

imply that a key pathway for synthesis of lipid mediators from DHA and DPA goes through 17-HDHA, as there were found significantly higher concentrations of 17-HDHA compared to 14-HDHA. This was even more pronounced for n-3 DPA, where only 17-HDPA was found in the media upon analysis. Also, it was interesting to see activation of the DPA metabolome considering the much lower content of this PUFA in HRO, thus implicating the importance of this pathway in regulation of inflammation. The apparent preference for the 17-hydroxylated DHA and n-3 DPA are aligned with literature showing 17-HDHA biosynthesis through ALOX15 in normal human keratinocytes when treated with DHA (75). Morin *et al.* have shown that supplementation of a psoriatic skin cell model with pure EPA also led to increased epidermal production of 17-HDHA, besides PGE3 and 12-HEPE, but not in the presence of polarized T-cells (76). Future experiments in the keratinocyte/fibroblast co-cultures should include chiral analyses of the monohydroxylated fatty acids in order to further elucidate which enzymes that are involved. Our findings furthermore indicates that the SPM biosynthesis from DHA and DPA is maintained in an inflamed state, as the precursor levels were found to be very similar between the HRO and the HRO + IL-17A groups. However, where ALOX15 seems to be important for conversion of DHA and DPA, this does not seem to be the case for EPA. Biosynthesis via 18-HEPE seems to be the main pathway for EPA derived lipid mediators with 15-HEPE levels being comparatively lower. This could be due to substrate vs enzyme availability in the system, but the finding is also consistent with the SPMs found in both the skin cells and the macrophages, where the downstream product of 15S-HEPE RvE4 was not measured in any of the experiments.

The observed biosynthesis of the cysteinyl SPM PCTR2 in the macrophages may also indicate a shift towards a reparative phenotype as well as a protective one, as the PCTRs have been reported to be biosynthesized by macrophages and to be involved in macrophage phenotype regulation and tissue regeneration (14,77,78). The PCTRs are biosynthesized similarly to PD1 through ALOX15 conversion of 17S-HpDHA into 16S,17S-epoxy-DHA, but with a last step through a glutamyltransferase instead of an epoxide hydrolase to give PCTR1 instead of PD1. PCTR1 is then hydrolyzed to PCTR2 which may be taken through a final hydrolysis to PCTR3. The linear nature of their biosynthesis implies that PCTR1 also was produced by these cells prior to

being hydrolyzed to PCTR2, which is interesting given that PCTR1 has been shown to stimulate keratinocyte migration and facilitate host defense (14).

In addition to promoting lipid mediator biosynthesis, we also observed that HRO significantly increased the levels of EPA/DHA in the total fatty acids in the MDMs and significantly increased the levels of EPA/DHA and reduced the levels of AA in the PL-fraction of the fibroblast/keratinocyte co-culture. Allowing for potential replacement of AA in cell membranes could positively impact psoriasis, as overexpressed PLA2 enzymes liberating AA for conversion into pro-inflammatory lipid mediators is a driver for psoriatic inflammation (79).

The results presented in this work also shows that both HRO and PEHeRo contain SPMs, including: 1) PDX that has been shown to ameliorate inflammation and promote macrophage polarization in experimental models of infection (80). 2) MaR1 that has been reported to display immunoresolving effects in models of skin inflammation through regulation of IL-23 (81), and 3) RvE4 that increases efferocytosis of apoptotic cells (82). The impact of these existing SPMs in the oils was not elucidated in this work, but considering their reported effects, they may provide anti-inflammatory effects and could potentially impact SPM biosynthesis. The effect of exogenous administration of lipid mediators has been shown in humans, where supplementation has been shown to increase blood lipid mediator levels (83). However, as stated above, the treatment of MDMs indicated that the existing lipid mediator levels in HRO and PEHeRo were not the primary drivers for biosynthesis, as the PEHeRo contained far less SPMs and monohydroxylated fatty acids than HRO but resulted in significantly increased levels of DHA-derived SPMs in the cell media.

The results shown in this work indicates relevance for understanding more of the underlying mechanisms that allows for regulation of cell processes involved in the pathophysiology of psoriasis through lipid mediators and SPMs. The key limitation to the data, however, is this being *in vitro* experiments, where all fatty acids and lipid mediators are made very accessible in concentrations that may not reflect realistic *in vivo* concentrations. Thus, the effects of directly metabolically available PUFAs and, particularly, lipid mediators present in the oils

may therefore be over-emphasized. It is therefore important to take this into further *in vivo* trials for confirming the relevance of the findings.

The data on skin cells mimicking psoriasis indicates a possible mechanism of resolution of inflammation in the local skin niche which could provide a treatment modality for psoriasis, and it is plausible that these mechanisms also could contribute to resolution of inflammation in other organs and IMIDs. The findings will also contribute to the broader understanding of the pharmacological factors that are in play when administering exogenous sources of SPMs and SPM precursors into biological systems.

In summary, we observed that HRO promotes broad SPM and lipid mediator biosynthesis in macrophages and keratinocytes/fibroblasts, with a subset of SPMs that are consistently upregulated throughout the stimulation/treatment conditions and cell types. Thus, indicating that these pathways may carry significance for signaling and inflammation resolution across several cell types. When treating macrophages with HRO and PEHeRo we found that the macrophages exhibited a dose-dependent trend for biosynthesis of SPMs associated with resolution of inflammation and a shift towards a more protective and possibly reparative phenotype. Given the human interventions and preclinical data that are available pertaining to the use of HRO in the management of psoriasis, the presented results suggest as a model of inflammatory disease for studies of local cellular resolution of inflammation processes.

Data availability

Lipid mediator raw data to be shared upon request. Contact corresponding author Thomas Ringheim-Bakka (Arctic Bioscience AS, Industrivegen 42, 6155 Ørsta, Norway, thomas@arctic-bioscience.com) for access.

Conflict of Interest

Thomas Ringheim-Bakka, Maftuna Busygina, and Runhild Gammelsæter are employed by Arctic Bioscience AS.

The authors declare that all experiments performed were partially funded by Arctic Bioscience AS.

Jesmond Dalli is an inventor on patents related to the composition of matter and/or use of pro-resolving mediators some of which are licensed by Brigham and Women's Hospital or Queen Mary University of London for clinical development.

Acknowledgments

We would like to thank Daniele Mancinelli and Federico Petrucelli for contributing to providing HRO and PEHeRo and describing the scope of the project, Sissel B. Rønning for contributing to the cell co-culture experiments and scientific discussions, Elin Merete Wetterhus for the skillful technical assistance in analyzing the fatty acid composition of the cells, and Bodil Stige at "Medisinsk biokjemi og blodbank Ålesund" for the provision of buffy coats.

References

1. Cavaillon JM. Once upon a time, inflammation. J Venom Anim Toxins Trop Dis. 2021 Apr 9;27:e20200147.

2. Serhan CN, Yacoubian S, Yang R. Anti-inflammatory and proresolving lipid mediators. Annu Rev Pathol. 2008;3:279–312.

3. Recchiuti A, Serhan CN. Pro-Resolving Lipid Mediators (SPMs) and Their Actions in Regulating miRNA in Novel Resolution Circuits in Inflammation. Front Immunol. 2012 Oct 22;3:298.

4. Serhan CN. Novel Pro-Resolving Lipid Mediators in Inflammation Are Leads for Resolution Physiology. Nature. 2014 Jun 5;510(7503):92.

5. Serhan CN, Chiang N, Dalli J. New pro-resolving n-3 mediators bridge resolution of infectious inflammation to tissue regeneration. Mol Aspects Med. 2018 Dec;64:1–17.

6. Serhan CN, Chiang N, Dalli J, Levy BD. Lipid Mediators in the Resolution of Inflammation. Cold Spring Harb Perspect Biol. 2015 Feb 1;7(2):a016311.

7. Serhan CN. The resolution of inflammation: the devil in the flask and in the details. FASEB J. 2011 May;25(5):1441.

8. Serhan CN, Levy BD. Resolvins in inflammation: emergence of the pro-resolving superfamily of mediators. J Clin Invest. 2018 Jul 2;128(7):2657–69.

9. Basil MC, Levy BD. Specialized pro-resolving mediators: endogenous regulators of infection and inflammation. Nat Rev Immunol. 2016 Jan;16(1):51–67.

10. Jordan PM, Werz O. Specialized pro-resolving mediators: biosynthesis and biological role in bacterial infections. FEBS J. 2022;289(14):4212–27.

11. Bannenberg G, Serhan CN. Specialized Pro-Resolving Lipid Mediators in the Inflammatory Response: An Update. Biochim Biophys Acta. 2010 Aug 10;1801(12):1260.

12. Murakami M. Lipid Mediators in Life Science. Exp Anim. 2011;60(1):7–20.

13. Lim JY, Park CK, Hwang SW. Biological Roles of Resolvins and Related Substances in the Resolution of Pain. BioMed Res Int. 2015;2015:830930.

14. Sansbury BE, Li X, Wong B, Riley CO, Shay AE, Nshimiyimana R, et al. PCTR1 Enhances Repair and Bacterial Clearance in Skin Wounds. Am J Pathol. 2021 Jun 1;191(6):1049–63.

15. Chiang N, de la Rosa X, Libreros S, Pan H, Dreyfuss JM, Serhan CN. Cysteinyl-specialized proresolving mediators link resolution of infectious inflammation and tissue regeneration via TRAF3 activation. Proc Natl Acad Sci U S A. 2021 Mar 9;118(10):e2013374118.

16. Dalli J, Gomez EA, Jouvene CC. Utility of the Specialized Pro-Resolving Mediators as Diagnostic and Prognostic Biomarkers in Disease. Biomolecules. 2022 Mar;12(3):353.

17. Kuek A, Hazleman BL, Östör AJK. Immune-mediated inflammatory diseases (IMIDs) and biologic therapy: a medical revolution. Postgrad Med J. 2007 Apr 1;83(978):251–60.

18. Wu D, Jin Y, Xing Y, Abate MD, Abbasian M, Abbasi-Kangevari M, et al. Global, regional, and national incidence of six major immune-mediated inflammatory diseases: findings from the global burden of disease study 2019. eClinicalMedicine [Internet]. 2023 Oct 1 [cited 2025 Jan 6];64. Available from: https://www.thelancet.com/journals/eclinm/article/PIIS2589-5370(23)00370-X/fulltext

19. Conrad N, Verbeke G, Molenberghs G, Goetschalckx L, Callender T, Cambridge G, et al. Autoimmune diseases and cardiovascular risk: a population-based study on 19 autoimmune diseases and 12 cardiovascular diseases in 22 million individuals in the UK. The Lancet. 2022 Sep 3;400(10354):733–43.

20. El-Gabalawy H, Guenther LC, Bernstein CN. Epidemiology of Immune-Mediated Inflammatory Diseases: Incidence, Prevalence, Natural History, and Comorbidities. J Rheumatol Suppl. 2010 May 1;85:2–10.

21. Takeshita J, Grewal S, Langan SM, Mehta NN, Ogdie A, Van Voorhees AS, et al. Psoriasis and Comorbid Diseases Part I. Epidemiology. J Am Acad Dermatol. 2017 Mar;76(3):377–90.

22. Olejnik M, Adamski Z, Dorocka-Bobkowska B. Psoriasis: From antiquity to the present. Indian J Dermatol Venereol Leprol. 2023;89(1):149–53.

23. Patel DD, Lee DM, Kolbinger F, Antoni C. Effect of IL-17A blockade with secukinumab in autoimmune diseases. Ann Rheum Dis. 2013 Apr 1;72(suppl 2):iii116–23.

24. Curtis JR, Mariette X, Gaujoux-Viala C, Blauvelt A, Kvien TK, Sandborn WJ, et al. Long-term safety of certolizumab pegol in rheumatoid arthritis, axial spondyloarthritis, psoriatic arthritis, psoriasis and Crohn's disease: a pooled analysis of 11 317 patients across clinical trials. RMD Open. 2019 May 1;5(1):e000942.

25. Navarro-Compán V, Puig L, Vidal S, Ramírez J, Llamas-Velasco M, Fernández-Carballido C, et al. The paradigm of IL-23-independent production of IL-17F and IL-17A and their role in chronic inflammatory diseases. Front Immunol. 2023;14:1191782.

26. Krueger JG, Eyerich K, Kuchroo VK, Ritchlin CT, Abreu MT, Elloso MM, et al. IL-23 past, present, and future: a roadmap to advancing IL-23 science and therapy. Front Immunol. 2024 Apr 15;15:1331217.

27. Simard M, Morin S, Ridha Z, Pouliot R. Current knowledge of the implication of lipid mediators in psoriasis. Front Immunol. 2022 Aug 26;13:961107.

28. Coras R, Kavanaugh A, Kluzniak A, Holt D, Weilgosz A, Aaron A, et al. Differences in oxylipin profile in psoriasis versus psoriatic arthritis. Arthritis Res Ther. 2021 Jul 24;23(1):200.

29. Sorokin AV, Domenichiello AF, Dey AK, Yuan ZX, Goyal A, Rose SM, et al. Bioactive Lipid Mediator Profiles in Human Psoriasis Skin and Blood. J Invest Dermatol. 2018 Jul;138(7):1518–28.

30. Sorokin AV, Arnardottir H, Svirydava M, Ng Q, Baumer Y, Berg A, et al. Comparison of the dietary omega-3 fatty acids impact on murine psoriasis-like skin inflammation and associated lipid dysfunction. J Nutr Biochem. 2023 Jul 1;117:109348.

31. Sawada Y, Honda T, Nakamizo S, Otsuka A, Ogawa N, Kobayashi Y, et al. Resolvin E1 attenuates murine psoriatic dermatitis. Sci Rep. 2018 Aug 8;8(1):11873.

32. Xu J, Duan X, Hu F, Poorun D, Liu X, Wang X, et al. Resolvin D1 attenuates imiquimod-induced mice psoriasiform dermatitis through MAPKs and NF-κB pathways. J Dermatol Sci. 2018 Feb 1;89(2):127–35.

33. Neuhofer A, Zeyda M, Mascher D, Itariu BK, Murano I, Leitner L, et al. Impaired Local Production of Proresolving Lipid Mediators in Obesity and 17-HDHA as a Potential Treatment for Obesity-Associated Inflammation. Diabetes. 2013 May 20;62(6):1945–56.

34. Calder PC. Eicosapentaenoic and docosahexaenoic acid derived specialised pro-resolving mediators: Concentrations in humans and the effects of age, sex, disease and increased omega-3 fatty acid intake. Biochimie. 2020 Nov 1;178:105–23.

35. Simopoulos AP. Omega-3 Fatty Acids in Inflammation and Autoimmune Diseases. J Am Coll Nutr. 2002 Dec 1;21(6):495–505.

36. Calder PC. Marine omega-3 fatty acids and inflammatory processes: Effects, mechanisms and clinical relevance. Biochim Biophys Acta BBA - Mol Cell Biol Lipids. 2015 Apr 1;1851(4):469–84.

37. Nordgren TM, Anderson Berry A, Van Ormer M, Zoucha S, Elliott E, Johnson R, et al. Omega-3 Fatty Acid Supplementation, Pro-Resolving Mediators, and Clinical Outcomes in Maternal-Infant Pairs. Nutrients. 2019 Jan;11(1):98.

38. Grenon SM, Owens CD, Nosova EV, Hughes□Fulford M, Alley HF, Chong K, et al. Short□Term, High□Dose Fish Oil Supplementation Increases the Production of Omega□3 Fatty Acid–Derived Mediators in Patients With Peripheral Artery Disease (the OMEGA□PAD I Trial). J Am Heart Assoc. 2015 Aug 25;4(8):e002034.

39. Sobrino A, Walker ME, Colas RA, Dalli J. Protective activities of distinct omega-3 enriched oils are linked to their ability to upregulate specialized pro-resolving mediators. PLoS ONE. 2020 Dec 16;15(12):e0242543.

40. Ahmmed MK, Ahmmed F, Tian H (Sabrina), Carne A, Bekhit AED. Marine omega-3 (n-3) phospholipids: A comprehensive review of their properties, sources, bioavailability, and relation to brain health. Compr Rev Food Sci Food Saf. 2020;19(1):64–123.

41. Lordan R, Tsoupras A, Zabetakis I. Phospholipids of Animal and Marine Origin: Structure, Function, and Anti-Inflammatory Properties. Molecules. 2017 Nov;22(11):1964.

42. Shabana S, Hamouda HI, Hamadou AH, Ahmed B, Chi Z, Liu C. Marine phospholipid nanoliposomes: A promising therapeutic approach for inflammatory bowel disease: Preparation, safety, and efficacy evaluation. Colloids Surf B Biointerfaces. 2024 Feb 1;234:113702.

43. Liu L, Bartke N, Daele HV, Lawrence P, Qin X, Park HG, et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. J Lipid Res. 2014 Mar 1;55(3):531–9.

44. Sugasini D, Yalagala PCR, Goggin A, Tai LM, Subbaiah PV. Enrichment of brain docosahexaenoic acid (DHA) is highly dependent upon the molecular carrier of dietary DHA: lysophosphatidylcholine is more efficient than either phosphatidylcholine or triacylglycerol. J Nutr Biochem. 2019 Dec 1;74:108231.

45. Patrick RP. Role of phosphatidylcholine-DHA in preventing APOE4-associated Alzheimer's disease. FASEB J. 2019 Feb;33(2):1554–64.

46. Sugasini D, Thomas R, Yalagala PCR, Tai LM, Subbaiah PV. Dietary docosahexaenoic acid (DHA) as lysophosphatidylcholine, but not as free acid, enriches brain DHA and improves memory in adult mice. Sci Rep. 2017 Sep 12;7(1):11263.

47. Hachem M, Géloën A, Van AL, Foumaux B, Fenart L, Gosselet F, et al. Efficient Docosahexaenoic Acid Uptake by the Brain from a Structured Phospholipid. Mol Neurobiol. 2016 Jul;53(5):3205–15.

48. Wang D, Zhang L, Wen M, Du L, Gao X, Xue C, et al. Enhanced neuroprotective effect of DHA and EPA-enriched phospholipids against 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) induced oxidative stress in mice brain. J Funct Foods. 2016 Aug 1;25:385–96.

49. Zhang LY, Ding L, Shi HH, Xu J, Xue CH, Zhang TT, et al. Eicosapentaenoic acid in the form of phospholipids exerts superior anti-atherosclerosis effects to its triglyceride form in ApoE-/- mice. Food Funct. 2019 Jul 17;10(7):4177–88.

50. Bjørndal B, Strand E, Gjerde J, Bohov P, Svardal A, Diehl BW, et al. Phospholipids from herring roe improve plasma lipids and glucose tolerance in healthy, young adults. Lipids Health Dis. 2014 May 17;13:82.

51. Cook CM, Hallaråker H, Sæbø PC, Innis SM, Kelley KM, Sanoshy KD, et al. Bioavailability of long chain omega-3 polyunsaturated fatty acids from phospholipid-rich herring roe oil in men and women with mildly elevated triacylglycerols. Prostaglandins Leukot Essent Fatty Acids. 2016 Aug;111:17–24.

52. Ramirez JL, Gasper WJ, Khetani SA, Zahner GJ, Hills NK, Mitchell PT, et al. Fish Oil Increases Specialized Pro-resolving Lipid Mediators in PAD (The OMEGA-PAD II Trial). J Surg Res. 2019 Jun 1;238:164–74.

53. Marchand NE, Choi MY, Oakes EG, Cook NR, Stevens E, Gomelskaya N, et al. Over-the-counter fish oil supplementation and pro-resolving and pro-inflammatory lipid mediators in rheumatoid arthritis. Prostaglandins Leukot Essent Fatty Acids. 2023 Mar 1;190:102542.

54. Mas E, Croft KD, Zahra P, Barden A, Mori TA. Resolvins D1, D2, and Other Mediators of Self-Limited Resolution of Inflammation in Human Blood following n-3 Fatty Acid Supplementation. Clin Chem. 2012 Oct 1;58(10):1476–84.

55. Norris PC, Skulas-Ray AC, Riley I, Richter CK, Kris-Etherton PM, Jensen GL, et al. Identification of specialized pro-resolving mediator clusters from healthy adults after intravenous low-dose endotoxin and omega-3 supplementation: a methodological validation. Sci Rep. 2018 Dec 21;8(1):18050.

56. Mottola G, Werlin E, Wu B, Chen M, Chatterjee A, Schaller M, et al. Oral Resolvin D1 attenuates early inflammation but not intimal hyperplasia in a rat carotid angioplasty model. Prostaglandins Other Lipid Mediat. 2019 Dec 10;146:106401.

57. Callan N, Hanes D, Bradley R. Early evidence of efficacy for orally administered SPM-enriched marine lipid fraction on quality of life and pain in a sample of adults with chronic pain. J Transl Med. 2020 Oct 21;18(1):401.

58. Caputo MP, Radlowski EC, Lawson MA, Antonson AM, Watson JE, Matt SM, et al. Herring roe oil supplementation alters microglial cell gene expression and reduces peripheral inflammation after immune activation in a neonatal piglet model. Brain Behav Immun. 2019 Oct 1;81:455–69.

59. Tveit KS, Brokstad KA, Berge RK, Sæbø PC, Hallaråker H, Brekke S, et al. A Randomized, Double-blind, Placebo-controlled Clinical Study to Investigate the efficacy of Herring Roe Oil for treatment of Psoriasis. Acta Derm Venereol. 2020 May 28;100(10):adv00154.

60. Tveit KS, Brokstad KA, Berge RK, Saeboe PC, Hallaraaker H, Brekke S, et al. Long Term Efficacy and Safety of Herring Roe Oil in the Treatment of Psoriasis, a 39-week Open-label Extension Study. Int J Clin Exp Med Sci. 2021 Jan;7(1):13–20.

61. Petrovic A, Bueide I, Tveit KS, Hallaråker H, Bjørndal B, Holmes TD, et al. Herring roe oil in treatment of psoriasis – influence on immune cells and cytokine network. Front Immunol. 2023 Sep 8;14:1128986.

62. Mildenberger J, Solberg N, Østbye TKK, Høst V, Federico P, Gammelsæter R, et al. Phospholipid Esters from Herring Roe have immunomodulatory anti-psoriatic effects by affecting signaling on the IL-17/23 axis in immune cells and psoriatic skin cell models in vitro. In: EADV Congress, Conference poster 3243. Amsterdam; 2024.

63. Titos E, Rius B, González-Périz A, López-Vicario C, Morán-Salvador E, Martínez-Clemente M, et al. Resolvin D1 and its precursor docosahexaenoic acid promote resolution of adipose tissue inflammation by eliciting macrophage polarization toward an M2-like phenotype. J Immunol Baltim Md 1950. 2011 Nov 15;187(10):5408–18.

64. Furue M, Furue K, Tsuji G, Nakahara T. Interleukin-17A and Keratinocytes in Psoriasis. Int J Mol Sci. 2020 Feb 13;21(4):1275.

65. Folch J, Lees M, Stanley GHS. A SIMPLE METHOD FOR THE ISOLATION AND PURIFICATION OF TOTAL LIPIDES FROM ANIMAL TISSUES. J Biol Chem. 1957 May 1;226(1):497–509.

66. Dooley M, Saliani A, Dalli J. Development and Validation of Methodologies for the Identification of Specialized Pro-Resolving Lipid Mediators and Classic Eicosanoids in Biological Matrices. J Am Soc Mass Spectrom. 2024 Oct 2;35(10):2331–43.

67. Pang Z, Lu Y, Zhou G, Hui F, Xu L, Viau C, et al. MetaboAnalyst 6.0: towards a unified platform for metabolomics data processing, analysis and interpretation. Nucleic Acids Res. 2024 Jul 5;52(W1):W398–406.

68. Wang DD, Wu F, Wen M, Ding L, Du L, Xue CH, et al. Replenishment of Docosahexaenoic Acid (DHA) in Dietary n-3-Deficient Mice Fed DHA in Triglycerides or Phosphatidylcholines After Weaning. J Food Sci. 2018 Feb;83(2):481–8.

69. Meyer zu Heringdorf D, Jakobs KH. Lysophospholipid receptors: Signalling, pharmacology and regulation by lysophospholipid metabolism. Biochim Biophys Acta BBA - Biomembr. 2007 Apr 1;1768(4):923–40.

70. Oh SF, Dona M, Fredman G, Krishnamoorthy S, Irimia D, Serhan CN. Resolvin E2 Formation and Impact in Inflammation Resolution. J Immunol. 2012 May 1;188(9):4527–34.

71. Sato M, Aoki-Saito H, Fukuda H, Ikeda H, Koga Y, Yatomi M, et al. Resolvin E3 attenuates allergic airway inflammation via the interleukin-23-interleukin-17A pathway. FASEB J Off Publ Fed Am Soc Exp Biol. 2019 Nov;33(11):12750–9.

72. Spite M, Norling LV, Summers L, Yang R, Cooper D, Petasis NA, et al. Resolvin D2 is a potent regulator of leukocytes and controls microbial sepsis. Nature. 2009 Oct;461(7268):1287–91.

73. Shan K, Feng N, Cui J, Wang S, Qu H, Fu G, et al. Resolvin D1 and D2 inhibit tumour growth and inflammation via modulating macrophage polarization. J Cell Mol Med. 2020 Jul;24(14):8045–56.

74. Sorokin AV, Norris P, English J, Dey AK, Chaturvedi A, Baumer Y, et al. Identification of pro-resolving and inflammatory lipid mediators in human psoriasis. J Clin Lipidol. 2018 Apr 6;12(4):1047.

75. Hellmann J, Sansbury BE, Wong B, Li X, Singh M, Nuutila K, et al. Biosynthesis of D-Series Resolvins in Skin Provides Insights into their Role in Tissue Repair. J Invest Dermatol. 2018 Sep 1;138(9):2051–60.

76. Morin S, Tremblay A, Dumais E, Julien P, Flamand N, Pouliot R. Eicosapentaenoic Acid Influences the Lipid Profile of an In Vitro Psoriatic Skin Model Produced with T Cells. Biomolecules. 2023 Sep;13(9):1413.

77. Dalli J, Ramon S, Norris PC, Colas RA, Serhan CN. Novel proresolving and tissue-regenerative resolvin and protectin sulfido-conjugated pathways. FASEB J. 2015 May;29(5):2120–36.

78. Dalli J, Serhan CN. Identification and structure elucidation of the pro☐resolving mediators provides novel leads for resolution pharmacology. Br J Pharmacol. 2019 Apr;176(8):1024–37.

79. Shao S, Chen J, Swindell WR, Tsoi LC, Xing X, Ma F, et al. Phospholipase A2 enzymes represent a shared pathogenic pathway in psoriasis and pityriasis rubra pilaris. JCI Insight. 6(20):e151911.

80. Xia H, Chen L, Liu H, Sun Z, Yang W, Yang Y, et al. Protectin DX increases survival in a mouse model of sepsis by ameliorating inflammation and modulating macrophage phenotype. Sci Rep. 2017 Mar 7;7(1):99.

81. Saito-Sasaki N, Sawada Y, Mashima E, Yamaguchi T, Ohmori S, Yoshioka H, et al. Maresin-1 suppresses imiquimod-induced skin inflammation by regulating IL-23 receptor expression. Sci Rep. 2018 Apr 3;8(1):5522.

82. Libreros S, Shay AE, Nshimiyimana R, Fichtner D, Martin MJ, Wourms N, et al. A New E-Series Resolvin: RvE4 Stereochemistry and Function in Efferocytosis of Inflammation-Resolution. Front Immunol. 2021 Feb 10;11:631319.

83. Souza PR, Marques RM, Gomez EA, Colas RA, De Matteis R, Zak A, et al. Enriched Marine Oil Supplements Increase Peripheral Blood Specialized Pro-Resolving Mediators Concentrations and Reprogram Host Immune Responses: A Randomized Double-Blind Placebo-Controlled Study. Circ Res. 2020 Jan 3;126(1):75–90.

Figure legends

Fig. 1 A) Relative concentrations of select SPMs in HRO and PEHeRo (mean of 3 replicates ±SEM) as a percentage of total SPMs found in the oils. B) Relative concentrations of monohydroxylated fatty acids in HRO and PEHeRo (mean of 3 replicates ±SEM) as a percentage of the total amount of monohydroxylated fatty acids found in the oils. C) Relative distribution of monohydroxylated fatty acids in HRO and PEHeRo (mean of 3 replicates ±SEM) as a percentage of the sum monohydroxylated fatty caids from DHA, EPA, and n-3 DPA. SPMs were identified and quantified using LC-MS/MS. # $P > 0.1$ (Mann-Whitney U test, two-tailed) and indicating similarity between HRO and PEHeRo for the analyte, remaining parameters were $P < 0.1$.

Fig. 2 A) Relative concentrations from GC-FID of fatty acids as % of total fatty acids in MDMs (mean of 3 replicates ±SEM) when treated with HRO versus control. B) Relative concentrations from GC-FID of fatty acids as % of total fatty acids in the phospholipid fraction of the keratinocyte/fibroblast co-culture (mean of 9 replicates ±SEM) when treated with HRO versus control. *$P < 0.05$ (One-factor ANOVA with Tukey's HSD).

Fig. 3 A) SPM levels (mean of 4 donors/experiments ± SEM) in supernatants from macrophages incubated with HRO prior to stimulation with LPS+IFN-γ versus only stimulated with LPS+IFN-γ (control). B) Partial least squares discriminant analysis (PLS-DA) of supernatants from macrophages incubated with HRO prior to stimulation with LPS+IFN-γ versus only stimulated with LPS+IFN-γ (control). Results are compared to lipid mediator concentrations identified in fresh media. C) Variable importance in projection (VIP) scores of SPMs that contribute to the difference between the tested conditions. D) Figure showing the treatment conditions and the highlighted measured outcomes. SPMs were identified and quantified using LC-MS/MS. *$P < 0.05$ (Mann-Whitney U test, two-tailed).

Fig. 4 SPM levels (mean of 4 donors/experiments ± SEM) from DHA (A) and EPA (B) metabolomes in macrophage cell supernatants (pg/3 mL) when incubated with HRO+LPS, PEHeRo+LPS, LPS (control), or untreated. C) Partial least squares discriminant analysis (PLS-DA) of supernatants from macrophages incubated with HRO+LPS, PEHeRo+LPS, LPS (control), or untreated. D) Variable importance in projection

(VIP) scores of SPMs that contribute to the difference between the tested conditions. SPMs were identified and quantified using LC-MS/MS. *P < 0.05 and **P < 0.01 (One-factor ANOVA with one-way Dunnett's multiple comparisons test for comparison to control and Tukey's HSD for comparison between treatment groups).

Fig. 5 SPM and SPM precursor levels (mean of 3 biological replicates ± SEM) in fibroblast/keratinocyte co-culture cell supernatants (pg/mL) incubated with IL-17A (control), HRO, HRO+IL-17A, or untreated. SPMs were identified and quantified using LC-MS/MS. A) SPMs and monohydroxylated fatty acids from DHA and monohydroxylated fatty acids from DPA. B) SPMs and monohydroxylated fatty acids from EPA. C) Monohydroxylated fatty acids from DHA. D) Monohydroxylated fatty acids from EPA. *P < 0.05 and **P < 0.01 (One-factor ANOVA with one-way Dunnett's multiple comparisons test for comparison to control and Tukey's HSD for comparison between treatment groups and between different monohydroxylated fatty acids).