Neuronal differentiation enhances a cytoplasmic pool of tousled-like kinase 2 (TLK2)

Abbreviated title: Neuronal differentiation affects TLK2 localisation

Lubna Nuhu-Soso, Heidi Denton, Darren L Goffin, Ines Hahn, Gareth J.O. Evans[#]

Department of Biology and York Institute of Biomedical Sciences, University of York, Wentworth Way, Heslington, York, YO10 5DD, United Kingdom

[#]corresponding author: gareth.evans@york.ac.uk

Number of pages: 29
Number of figures: 5
Word counts:
    Abstract: 250
    Significance statement: 117
    Introduction: 577
    Discussion: 1389

Conflicts of interest
The authors declare no competing financial interests.

Acknowledgements
This work was part-funded by the Wellcome Trust (ref: 204829) through the Centre for Future Health (CFH) at the University of York and an Academy of Medical Science Springboard Award to IH (SBF008\1140). We are grateful to Prof Betsy Pownall and Dr Harry Isaacs (University of York) for valuable advice regarding the *in situ* hybridisation experiments. We thank the Bioscience Technology Facility at the University of York for providing access to confocal microscopes.

Abstract

Mental retardation autosomal dominant 57 (MRD57) is a rare neurodevelopmental disorder

characterised by delayed language and psychomotor development, intellectual disability,

hypotonia, gastrointestinal issues and facial dysmorphia. It is linked to genetic mutations in

the serine/threonine kinase TLK2, characterised by haploinsufficiency and in some cases, its

loss or impaired kinase function. TLK2 is an established cell cycle regulator that has been

extensively studied in mitotic cells. It is upregulated in cancers, driving tumour growth,

however, the role of TLK2 in postmitotic neurons is not understood. We therefore aimed to

gain insight into how TLK2 mutations cause MRD57 by determining where TLK2 is

expressed in the brain and its subcellular localisation during neuronal differentiation. Public

human and mouse brain transcriptomic data revealed splice variant diversity in the N-

terminus of TLK2, which contains its nuclear localisation sequence (NLS). Using splice-

specific *in situ* hybridisation probes, we observed expression of TLK2 isoforms that contain

and lack the NLS in the mouse hippocampus and cerebellum. We confirmed these findings in

human SH-SY5Y neuroblastoma cells, and found that neuronal differentiation of these cells

enhances a cytoplasmic pool of TLK2 by two mechanisms: nuclear export of full length TLK2

and increased expression of TLK2 splice variants lacking the NLS. Finally, acute stimuli that

mimic synaptic activity were sufficient to elicit nuclear export of TLK2. Our data highlight the

need to establish the neuronal cytoplasmic substrates of TLK2 and determine how the loss of

TLK2 activity in MRD57 might impact their function in the developing and mature brain.

Significance statement

Tousled-like kinase 2 (TLK2) regulates the cell cycle and genome maintenance, and is

inextricably linked with disease. TLK2 is upregulated in several cancers, but conversely,

reduced TLK2 activity is implicated in a neurodevelopmental disorder, MRD57. However, the

role of TLK2 in the brain is unknown and we therefore characterised neuronal TLK2

expression and localisation. Rodent and human brain possess long and short TLK2 splice

variants that include or exclude the N-terminal nuclear localisation sequence respectively.

Neuronal differentiation increased cytoplasmic TLK2 resulting from enhanced expression of

short-TLK2 and nuclear export of long TLK2. Furthermore, nuclear-cytoplasmic shuttling of

TLK2 is activity-dependent. These findings suggest that the pathogenic mechanisms of

MRD57 likely include non-nuclear functions of TLK2 in postmitotic neurons.

Introduction

Tousled-like kinase 1 and 2 are the mammalian homologues of the *Arabidopsis* mutant,

tousled, which was discovered in a developmental screen (Roe et al., 1993). Indeed, both

human TLK1 and TLK2 have been linked with rare developmental disorders (Segura-Bayona

and Stracker, 2019; Villamor-Payà et al., 2024). To date, there have been close to 50

documented patients worldwide with TLK2 mutations, classified as autosomal dominant

Mental Retardation Disorder 57 (MRD57; (Lelieveld et al., 2016; Reijnders et al., 2018; Töpf

et al., 2020; Pavinato et al., 2022; Woods et al., 2022), with a predicted incidence of

approximately 3 in 100,000 individuals (López-Rivera et al., 2020). The clinical phenotypes

associated with MRD57, like other neurodevelopmental disorders, are heterogeneous,

including intellectual disability, delayed psychomotor development in infancy or early

childhood, language delay, hypotonia, feeding problems, gastrointestinal issues, dysmorphic

facial features, microcephaly, behaviour problems and global developmental delays

(Reijnders et al., 2018; Töpf et al., 2020; Pavinato et al., 2022; Woods et al., 2022). The

majority of TLK2 mutations in MRD57 are *de novo* mutations, but seven are inherited within

four different families (Reijnders et al., 2018; Töpf et al., 2020; Pavinato et al., 2022). Like the

de novo cases, inherited TLK2 mutations are heterozygous dominant except one report of a

homozygous recessive inheritance (Reijnders et al., 2018; Töpf et al., 2020; Pavinato et al.,

2022). MRD57 patients have TLK2 mutations throughout the protein and all are predicted to

result in a reduction of TLK2 expression or its catalytic activity (Mortuza etc).

Functionally, TLK2 is an established regulator of DNA replication, cell cycle checkpoint

recovery and chromatin remodelling (Silljé et al., 1999; Bruinsma et al., 2016; Kim et al.,

2016a; Segura-Bayona et al., 2017). TLK2 facilitates cell cycle progression mainly by

enhancing histone supply via phosphorylation of its best characterised substrate, ASF1

(Klimovskaia et al., 2014). In-line with other cell cycle kinases, TLK activity and gene copy

number are increased in cancer, including breast, cervix, lung, liver, colon and kidney (Kim et

al., 2016b; Lee et al., 2018; Bhoir and De Benedetti, 2023). The chromosomal locus of TLK2,

17q23, contains other oncogenes, and is frequently amplified in more than 40% of breast

cancer tumours (Kelemen et al., 2009; Kim et al., 2016b). Moreover, single nucleotide

polymorphisms (SNPs) in TLK2 have been identified as risk factors for breast cancer

(Kelemen et al., 2009). Indeed, inhibition of TLK2 has been identified as a therapeutic

strategy and small molecule inhibitors are in development (Kim et al., 2016b; Lee et al.,

2018).

In contrast to its role in the cell cycle and cancer, the neuronal expression and cell biology of
TLK2 is poorly understood. Here we have characterised its neuronal transcript expression,
revealing a conserved splice diversity in rodents and humans. TLK2 alternative splicing
occurs in exons encoding the N-terminus, resulting in 'long' and 'short' isoforms, with the
latter lacking the nuclear localisation sequence (NLS). Expression of both long and short
TLK2 isoforms is prevalent in neurons of the mouse hippocampus and cerebellum. Using a
cell model of neuronal differentiation, we observed the upregulation of short TLK2 expression
during neuronal differentiation, with a concomitant shuttling of long TLK from the nucleus to
the cytoplasm. Pharmacological treatments mimicking neuronal activity also drive nuclear-
cytoplasmic shuttling of TLK2. These data highlight the likely importance of cytoplasmic
TLK2 in terminally differentiated neurons, which have exited the cell cycle, and a need to
determine the neuronal substrates of TLK2 in order to shed light on the neuronal phenotypes
of MRD57.
112
113   **Materials and Methods**
114
115   **Materials**
116   SH-SY5Y and Flp-In T-REx SK-N-SH neuroblastoma cell lines were kind gifts from Dr
117   Andrew Stoker, University College London and Dr Han-Jou Chen, University of York
118   respectively. The Flp recombinase expression vector, pOG44 was a kind gift from Dr Paul
119   Pryor, University of York.
120
Bioinformatics
TLK2 transcript IDs and exon structures were derived from their Ensembl gene annotations
(version 113) using human (GRCh38.p14) or mouse (GRCm39) genome builds (Harrison et
al., 2024). The human TLK2 RNAseq data used for the transcriptomic analyses described in
this manuscript were obtained from the GTEx Portal, accession number phs000424.v8.p2.
Mouse brain TLK2 RNAseq data was obtained from the Human Protein Atlas
(proteinatlas.org; (Sjöstedt et al., 2020)). TLK2 transcript expression values in TPM were
converted to percentages by tissue or brain region.
129
RNA extraction, cDNA synthesis and RT PCR
All work involving mice was approved by the University of York Animal Welfare and Ethical
Review Body and performed under UK Home Office legislation (project licence P38B2013E).
Adult wild-type C57BL/6 mouse forebrain and cerebellum were dissected and homogenised
in RIPA buffer (50 mM Tris, pH 8.0, 150 mM NaCl, 1% (v/v) Triton-X-100, 0.5% (w/v) sodium
deoxycholate, 0.1% (w/v) SDS, 1 mM EDTA supplemented with 1 mM Na_3VO_4, 0.1% (v/v) β-

mercaptoethanol, 1 mM PMSF and 1:200 protease inhibitor cocktail (Sigma)) using 5 ml

buffer/g of tissue. The homogenate was incubated at 4°C with agitation for 1 h and then

centrifuged at 20,000g for 30 min at 4°C. RNA was extracted from the supernatant.

SH-SY5Y cells undergoing neuronal differentiation were plated at 10^4 cells/ cm^2 onto

collagen coated 6-well tissue culture dishes. RNA was extracted from the differentiating cells

at two day intervals. For both mouse brain and SH-SY5Ys, RNA was extracted using a

NucleoSpin RNA isolation kit (Macherey-Nagel) according to the manufacturer's instructions.

Complementary DNA was synthesised from 1 µg of RNA using a SuperScript IV First-Strand

cDNA Synthesis Kit (Life Technologies) according to the manufacturer's instructions in the

presence of both oligo d(T)$_{18}$ and random hexamer primers. A no RT cDNA sample was

performed for each time point and cDNA was stored at -20°C. PCR reactions were

performed in 25 µl reactions containing 1 µl of cDNA, 1X GoTaq flexi buffer, 0.2 mM each of

dNTP mix, 0.4 µM of each primer pair (Table 1) and 1U Taq polymerase. Cycling conditions

consisted of an initial denaturing step at 98°C for 30s followed by 25-30 cycles of 95°C for

10s, 55°C for 30s, 72°C for 1 min/kb and a final extension at 68°C for 10 min. 5 - 10 µl of the

PCR products were separated on 1.5-2% (w/v) agarose gels to confirm amplification.

***In situ* hybridisation**

Mice were anaesthetised with isoflurane inhalation followed by an intraperitoneal injection of

100 mg/kg ketamine and 12.5 mg/kg xylazine and then transcardially perfused with 4%

paraformaldehyde (PFA) in 0.1 M sodium-potassium PBS. The brain dissected and post-

fixed in 4% PFA overnight at 4°C. Following a further 24 h incubation in 30% sucrose, brains

were embedded in OCT, frozen at -80°C, and 25 µm coronal or sagittal sections were cut

using a Leica CM1950 cryostat. Probes for *in situ* hybridisation were transcribed using 10X

digoxigenin (DIG) RNA labelling mix (Roche) and T7 polymerase. For *in vitro* transcription,

template DNA was synthesised as gBlocks™ (IDT) and PCR amplified to introduce the T7

promoter sequence (Table 1). *In situ* hybridisation was performed according to (Fisher et al.,

2002) with adaptations for sections mounted on slides. Briefly, the sections were first

incubated in 4% PFA for 10 min at room temperature (RT) followed by two 5 min PBST

washes. Sections were then treated with 4 µg/ml proteinase K (Roche) in PBST for 10 min

and then washed twice in PBST for 5 min. A second fix in 4% (w/v) PFA was performed for

10 min, followed by two 5 min PBST washes. Next, the sections were acetylated using 0.5%

(v/v) acetic anhydride in 0.1 M triethanolamine, pH 7.8 for 10 min followed by two 5 min

PBST washes and finally a 15 min wash in 5X saline-sodium citrate buffer (SSC; Sigma).

Pre-hybridisation was carried out by incubating the sections in hybridisation buffer (50% (v/v)

formamide (Ambion, AM9342), 5X SSC, 100 µg/ml heparin, 1X Denhardt's, 0.1% (v/v)

Tween-20, 0.1% (w/v) CHAPS, 10 mM EDTA, 1 mg/ml total yeast RNA) containing no probe

at 65°C for 2-5 h in a humidified box. Hybridisation followed by incubating the sections with

300 ng/ml of probe solution (DIG prep in hybridisation buffer) overnight at 65°C in a

humidified box.

Post-hybridisation, the sections were subjected to several hot wash steps. First a 10 min

wash in hybridisation buffer containing no probe at 65°C, then a 1 h wash in 2X SSC

followed by a 1 h wash in 0.2X SSC, both at 65°C. The sections were then washed twice for

5 min in MABT (100 mM maleic acid, 150 mM NaCl, 0.1% (v/v) Tween-20, pH 7.8) at RT and

then blocked in blocking solution (2% (w/v) Boehringer Mannheim Blocking Reagent (Roche,

11096176001), 20% (v/v) heat-treated lamb serum (Sigma), 1X MAB) for 2 h at RT in a

humidified box. Next, the sections were incubated with anti-DIG-AP (Table 2) in blocking

solution overnight at 4°C in a humidified box. Post-antibody washes were carried out at RT

by washing the sections three times for 1 h in MABT and once in alkaline phosphatase (AP)

buffer (100 mM Tris, pH 9.5, 50 mM $MgCl_2$, 100 mM NaCl) for 10 min. The colour reaction

was initiated by adding a 1:3 dilution of BM purple (Roche, 11442074001) in AP buffer to the

sections and incubation at RT in a humidified box for 1-5 days. The reaction was stopped by

washing the sections twice for 10 min in PBST, fixing in 4% (w/v) PFA for 1 h or overnight at

4°C and a 15 min wash in PBST. The sections were air dried, mounted in aqueous mounting

media (Sigma, 324590) and stored at 4°C. Images were acquired on a Leica DM2500

microscope with a 10X objective (sagittal) and a Zeiss Stemi 508 stereo microscope

(coronal).

Table 1. Primers and probes used in this study

Name	Sequence (5'-3')
PCR primers	
mTLK2-exon1 sense	GACCCACGAAGGCAGGAATTA
mTLK2-exon5/6 anti	CTCGACGCGGTAAGGGATTG
mTLK2-exon4/6 anti	CTGCTCGACGCTCAAAGTAA
mTLK2-exon3/6 anti	GCTCGACGCTTGGCTAGT
mTLK2-exon6 anti	GCTGCACTGCCATCTAAACC
mGAPDH sense	CATCACTGCCACCCAGAAGACTG
mGAPDH anti	ATGCCAGTGAGCTTCCCGTTCAG
hTLK2-exon1 sense	ATTGCATAGCCTGGACCCAC
hTLK2-exon9 anti	TCGTTCCTTGTATTTCTCTAGCA
hMAPT sense	ACACCATGCACCAAGACCAA
hMAPT anti	TCCTTCTGGGATCTCCGTGT
hGAPDH sense	GTCTCCTCTGACTTCAACAGCG

hGAPDH anti	ACCACCCTGTTGCTGTAGCC
ISH probes	
mTLK2-204	GGAACCAGCCCTGGCAGAAGTGTTCCACCAGTTGCACGATCCTCACCGCA
	ACATTCCTTATCCAATCCCTTACCGCGTCGAGCAGAACAGCCTCTGTATG
	GTTTAGATGGCAGTGCAGCAAAGGAGGCCTCAGAAGAGCAGTCTGCTCTG
mTLK2-201	AGGAAAGCCGAGCCATATGACACTAGCCAAGGGAAAGGCACTCCTAGGGG
	ACATAAAATTAGTGATTACTTTGAGCGTCGAGCAGAACAGCCTCTGTATG
	GTTTAGATGGCAGTGCAGCAAAGGAGGCCTCAGAAGAGCAGTCTGCTCTG
mTLK2-207	TAGAGACTCCTGAGAAAAAGCAGAATGACCAGCGAAATCGGAAAAGGAAA
	GCCGAGCCATATGACACTAGCCAAGCGTCGAGCAGAACAGCCTCTGTATG
	GTTTAGATGGCAGTGCAGCAAAGGAGGCCTCAGAAGAGCAGTCTGCTCTG
mTLK2-novel	CAACCAGAGTCTGTGCAGCGTGGGGTCGTTGAGTGATAAAGAAGTAGAGA
	CTCCTGAGAAAAAGCAGAATGACCGCGTCGAGCAGAACAGCCTCTGTATG
	GTTTAGATGGCAGTGCAGCAAAGGAGGCCTCAGAAGAGCAGTCTGCTCTG
mCalb1	GATGCTTTGCTGAAAGATCTGTGTGAGAAGAACAAACAGGAATTGGATAT
	TAACAATATTACTACATACAAGAAGAACATAATGGCCTTGTCGGATGGAG
	GGAAGCTGTACCGAACAGACCTTGCTCTTATTCTTTCTGCTGGAGACAAC

Culture, neuronal differentiation and pharmacological treatment of SH-SY5Y cells

SH-SY5Y cells were maintained at 37°C, 5% CO_2 in a humidified atmosphere in Dulbecco's

Modified Eagle Medium (DMEM) with high glucose, pyruvate and L-glutamine supplemented

with 10% foetal bovine serum (FBS) and 1% penicillin/streptomycin.

Neuronal differentiation of SH-SY5Y cells was performed according to a previously described

method (Encinas et al., 2000). Briefly, tissue culture dishes or coverslips were coated in 0.05

mg/ml collagen in 60% (v/v) ethanol. Cells were plated at 10^4 cells/cm^2 (for protein analysis)

or 5 x 10^3 cells/well (for microscopy) in DMEM supplemented with 15% (v/v) heat inactivated

FBS and 1% (v/v) penicillin/streptomycin. Twenty four hours after plating, the media was

changed and supplemented with 10 µM retinoic acid (RA; Sigma, R2625). Six days after

plating, the cells were washed three times in serum-free DMEM and then incubated with

serum-free DMEM containing 50 ng/ml brain-derived neurotrophic factor (BDNF; Qkine) and

1% (v/v) penicillin/streptomycin and cultured for a further seven days. Throughout the

differentiation process, the media was changed every two days and supplemented with the

appropriate differentiation agent.

For analysing protein expression in differentiating cells, lysates were collected at two day

intervals. Cells were washed twice in sterile ice-cold PBS and scraped in 100 µl of lysis

buffer (20 mM HEPES, pH 7.4, 10 mM KCl, 2 mM $MgCl_2$, 1 mM EDTA and 1 mM EGTA, 1

mM DTT and protease inhibitor cocktail). Lysates were incubated on ice for 15 min, passed

through a 25G needle ten times and protein concentration was determined by Bradford

assay. The lysates were diluted in Laemmli sample buffer, boiled for 10 min at 90°C and 10

µg of protein was separated by SDS-PAGE and analysed by Western blotting.

Pharmacological stimulation of undifferentiated SH-SY5Y cells was performed on cells plated

24 h prior to treatment on 13 mm coverslips in a 24-well plate at 5×10^4 cells/well. Cells were

washed twice in PBS and then incubated for 1 h at 37°C, 5% CO_2 in either control buffer (170

mM NaCl, 3.5 mM KCl, 0.4 mM KH_2PO4, 10 mM HEPES, 5 mM $NaHCO_3$, 5 mM glucose, 1.2

mM Na_2SO_4, 1.2 mM $MgCl_2$, 1.3 mM $CaCl_2$, pH 7.4, 0.2% (v/v) DMSO) or KCl/FSK buffer

(control buffer with 73.5 mM NaCl, 100 mM KCl, 10 µM forskolin, 100 µM IBMX, 0.2% (v/v)

DMSO). Cells were then fixed and immunocytochemistry was performed as described below

using anti-TLK2.

Inducible stable TLK2 cell lines

Tetracycline-inducible SK-N-SH cell lines expressing either full length TLK2-203 or short N-

terminally truncated TLK2-213 were generated using the Flp-In™ T-REx™ System

(Invitrogen). pcDNA5/FRT/TO-FLAG-TLK2 was purchased from the MRC PPU Reagents

and Services facility (College of Life Sciences, University of Dundee). TLK2-213 was made

by subcloning its ORF from full length TLK2 and ligating it back into the same backbone via

5' and 3' NotI restriction sites with the N-terminal FLAG tag intact. Cells were co-transfected

with the FRT backbone containing the gene of interest (TLK2-203 or TLK2-213) and a Flp-

recombinase expression vector (pOG44). Positive clones were selected by treating with 0.5

mg/ml hygromycin B and assaying for protein expression after the addition of 1 µg/ml

doxycycline.

Subcellular fractionation

Flp-In SK-N-SH cells were fractionated according to the Abcam subcellular fractionation

protocol. TLK2 expression was induced using 1 µg/ml doxycycline and left in culture for 48 h

prior to fractionation. Briefly, cells were scraped in 500 µl of lysis buffer (described above)

and incubated on ice for 15 min. The scraped cells were passed through a 25G needle ten

times, incubated on ice for 20 min and then centrifuged at 720g for 5 min at 4°C. The post-

nuclear supernatant was transferred to a fresh tube. The nuclear pellet was washed by first

resuspending in lysis buffer, passing the sample through a 25G needle ten times and

centrifuging at 720g for 10 min at 4°C. The washed pellet was resuspended in TBS

containing 0.1% (w/v) SDS, and sonicated on ice for 3 s. The protein concentration of each

fraction was determined by Bradford assay and 20 µg of each fraction was separated by

SDS-PAGE and analysed by Western blotting.

Western blotting

Proteins separated by SDS-PAGE (7.5-12%) gels were transferred onto PVDF membranes (Millipore). Membranes were incubated with primary antibodies (Table 2) overnight at 4°C with agitation. Membranes were washed and incubated with appropriate HRP-conjugated secondary antibodies and immunoreactivity was visualised by incubation with enhanced chemiluminescence reagent and imaged on an iBright FL1000 scanner (Invitrogen).

Immunocytochemistry and fluorescence microscopy

Cells were washed 3x in PBS and then fixed in 4% (w/v) paraformaldehyde (PFA) in PBS for 20 min at RT. Next cells were washed 3x in PBS then permeabilised and blocked in PBS containing 0.1% (v/v) TritonX-100 and 1% (w/v) bovine serum albumin (BSA). Primary antibodies were diluted in PBS containing 1% (w/v) BSA at appropriate dilutions (Table 2) and incubated on cells overnight at 4°C. Cells were washed 3x in PBS then incubated with secondary antibodies in PBS containing 1% (w/v) BSA for 1 h in the dark at RT. Finally, cells were washed 3x in PBS, 1x in dH_2O, air dried and mounted in Mowial containing 1 µg/ml 4′,6-diamidino-2-phenylindole (DAPI). Images were acquired in the Bioscience Technology Facility (University of York) on a Zeiss LSM 880 confocal microscope on a 40x/1.4 Oil DIC III objective.

Table 2. Antibodies used in this study.

Name	Dilution	Species	Catalogue no.	Supplier
Primary antibodies				
Anti-TLK2	1:1000 (WB) 1:500 (IF)	Rabbit	A301-257A	Bethyl
Anti-β-III-tubulin	1:1000 (WB) 1:500 (IF)	Mouse	801202	Biolegend
Anti-β-actin	1:50000 - 1:100000	Mouse	60008-1-Ig	Proteintech
Anti-Histone H3	1:5000	Mouse	819411	Biolegend
Anti-GAPDH	1:5000	Mouse	MAB374	Sigma
Anti-FLAG	1:1000	Mouse	F1804	Sigma

Anti-Digoxigenin	1:2000	Sheep	11093274910	Roche
Secondary antibodies				
Anti-mouse HRP	1:5000	Goat	A4416	Sigma
Anti-rabbit HRP	1:5000	Goat	A6154	Sigma
Anti-mouse Alexa 594	1:500	Goat	A11020	Invitrogen
Anti-rabbit Alexa 488	1:500	Goat	A11008	Invitrogen

Data analysis

Quantification of hippocampal *in situ* hybridization images was performed on 16 bit grayscale

images in FIJI (Schindelin et al., 2012). The oval selection tool was used to draw regions of

interest (ROI) in different locations within the hippocampal formation (CA1, CA3, DGs and

DGi). The intensity was then measured and normalised to DGs for each section and probe.

Three to four hippocampi were quantified for each probe and the values were plotted as a

heatmap in Morpheus (https://software.broadinstitute.org/morpheus).

The quantification of cytoplasmic/nuclear TLK2 ratios of immunofluorescent images was

conducted by measuring the TLK2 intensity in the nucleus and cytoplasm of the cells in FIJI

using the ROI 1-click plug-in (Thomas and Gehrig 2020). Three hundred cells for each

condition were analysed from three technical replicates (100 cells/replicate). Western blots

were quantified by densitometry analysis in FIJI. Statistical significance was assessed by t-

test or non-parametric ANOVA with Dunn's post-hoc test. Statistical analyses and graph

plotting was performed in RStudio (RStudio Team, 2020).

Results

TLK2 is alternatively spliced in human and mouse brain

Since the neuronal distribution and function of TLK2 has not been investigated, we first sought

to characterise TLK2 transcript expression in the brain. Previous studies have observed tissue-

specific transcripts of TLK2, in the mouse testis for example (Shalom and Don, 1999), and

TLK2 immunoblots often have multiple bands that vary between tissues and cell lines (Kim et

al., 2016b; Segura-Bayona et al., 2017). Such bands could result from post-translational

modifications or alternative splicing of TLK2 or a combination of these. We began by

interrogating publicly available human (GTEx) and mouse brain (Human Protein Atlas)

transcriptomic data to characterise neuronal TLK2 transcript expression (Fig. 1A,B), focussing

on the hippocampus and cerebellum. In both human and mouse brains, a full length transcript
corresponding to the canonical Ensembl TLK2 transcript was found to be the most abundant
transcript (Figs. 1B, S1 & S2). However, we identified evolutionarily conserved splice variant
diversity at the N-terminus of the protein, encoded by exons 1-9. Not surprisingly, there were
fewer alternative splicing events in exons encoding the middle coiled-coil domain and C-
terminal kinase domain of TLK2, which are essential for dimerisation and catalytic activity
respectively (Mortuza et al., 2018); Fig. 1B). Interestingly, splice variants lacking exon 3 in
which the nuclear localisation sequence (NLS, Fig. 1A) is located (Yamakawa et al., 1997) are
also abundant (Fig. 1B,S1). This suggests that a pool of TLK2 is constitutively resident outside
the nucleus. Henceforth in this study we shall refer to isoforms that include the NLS as 'long
TLK2', and those lacking the NLS as 'short TLK2'. In the human GTEx dataset it was observed
that transcript TLK2-207 represents nearly a third of TLK2 mRNA expression, but is predicted
to be subject to nonsense mediated decay due to an intron retention at exon 21 (Fig. 1B).

To confirm the existence of alternative N-terminal TLK2 splice variants in the brain, we
performed RT-PCR of mouse brain TLK2 mRNA. The annotated transcripts of mouse TLK2
have splice diversity within exons 1-6 (Fig. 2A and Fig S2) and also at exon 12. We first
amplified and sequenced all N-terminal splice variants in mouse forebrain or cerebellar cDNA
using transcript-specific primers and also a pair of 'pan isoform' primers located in exon 1
(forward) and exon 6 (reverse), which yielded four bands (Fig. 2A). Three of the bands
corresponded to the 5' of annotated isoforms, two of which are predicted to contain the NLS
(mTLK2-204 and -201) and one that lacks the NLS due to an alternative start site in exon 6
(mTLK2-207). We also sequenced a novel isoform that has a truncated exon 3, lacks exon 4
and 5 and would also not encode the NLS (mTLK2-novel; Fig. 2B and Fig. S2). We next used
in situ hybridisation to observe whether there is any cell type specificity in the expression of
these TLK2 isoforms in the brain. All mouse TLK2 transcripts express exon 6 and combinations
of exons 1-5, therefore we designed a suite of short 150 bp splice junction-spanning *in situ*
probes anchored at the 3' end by the first 75 bp complementary to exon 6 and containing a
variable 5' 75 bp complementary to exons 5, 4 or 3 (Fig. 2B).

In coronal and sagittal mouse brain sections we examined the expression of TLK2 isoforms in
the hippocampal formation and cerebellar cortex respectively. A 150 bp exon junction spanning
calbindin probe was used as a positive control, confirming calbindin expression in the Purkinje
neurons of the cerebellar cortex and dentate gyrus and CA1 neurons of the hippocampus (Fig.
2C,E). Low background staining was observed in the absence of an *in situ* probe (Fig. 2C,E).
In the coronal sections, hippocampal staining of all TLK2 isoform probes was detected in
excitatory neurons of the dentate gyrus, CA1 and CA3 (Fig. 2C). Although quantifying the

intensity of staining between probes was not appropriate, using regions of interest placed in
the DG, CA1 and CA3 fields, we quantified the relative intensity of staining of each TLK2
isoform in these hippocampal cell types and normalised the density to the signal in the
suprapyramidal blade of the dentate gyrus (DGs; Fig. 2C). The full length TLK2 (exons 1-6)
probe, corresponding to the most abundant splice variant mTLK2-204, gave strong staining in
the cell soma of the granule neurons in the DG and pyramidal neurons in CA1 and to a lesser
extent in CA3 (Fig 2B, C). The other long TLK2 isoform (mTLK2-201) was less readily detected
in the hippocampus, but quantification revealed differential expression in the two pyramidal
blades of the DG, with the suprapyramidal blade having the highest intensity (Fig. 2D). The
relative staining of the shorter variant, mTLK2-207, was highest in CA1 neurons. In-line with
the PCR (Fig. 2A), staining of the novel short TLK2 isoform appeared weakest, with the highest
intensity in CA1 and CA3 compared to the DG (Fig. 2C, D).

In sagittal sections of the cerebellar cortex, staining of the TLK2 probes was readily detectable
in the abundant granule neurons and to a lesser extent in Purkinje neurons (Fig. 2E). The
intensity of staining for the full length mTLK2-204 probe was similar to that of the shorter
mTLK2-207 in the granule cell and Purkinje layers, whereas the mTLK2-201 and mTLK2-novel
probes had weak granule layer staining and barely detectable staining of Purkinje neurons
(Fig. 2E). For all TLK2 probes, we failed to observe staining of inhibitory interneurons in the
hippocampal cortex or molecular layer of the cerebellar cortex. Taken together, these data
confirm the RNAseq data that alternative splice variants of TLK2 are expressed in neurons of
the mammalian brain. It is currently unclear as to the physiological relevance of the neuronal
cell-type specific expression of these TLK2 variants we have observed.

Alternative splicing of TLK2 is regulated during neuronal differentiation
To date, our knowledge of TLK2 is mainly limited to its nuclear functions (Segura-Bayona and
Stracker, 2019). The possibility of a non-nuclear pool of TLK2 in post-mitotic neurons has
interesting implications for the role of TLK2 in the developing and mature brain. To study TLK2
expression in neuronal differentiation we adopted a protocol for differentiating the human
neuroblastoma cell line SH-SY5Y. Based on a previously published method we used the
sequential application of retinoic acid (RA) and then BDNF with serum withdrawal over several
days (Encinas et al., 2000), which produced robust neuronal morphology (Fig. 3A) and
enhanced transcript expression of the neuronal marker tau (Fig. 3B). To capture the diversity
of 5' TLK2 splice variants we performed RT-PCR on SH-SY5Y cDNA samples from day 1-13
with primers located in exons 1 (forward) and 9 (reverse). This yielded six bands (Fig. 3B),
which were sequenced (Fig. S3). Fig. 3B summarises the exon structure of each TLK2 PCR
product, any corresponding annotated Ensembl transcripts and the predicted mass of the

protein they encode. The three longest PCR products mapped onto several Ensembl

transcripts with alternative splicing of combinations of exons 5, 7 and 8. The top two bands

each potentially correspond to two Ensmbl isoforms due to alternative splicing at exon 12,

which wasn't amplified in the PCR reaction. These bands were also the most prominent PCR

products in HEK293 cells. Sequencing the third band yielded a mix of sequences that we were

unable to deconvolve, but likely comprises up to three isoforms of the same mass (Fig. 3B &

Fig. S3). We were not able to detect expression of a transcript corresponding to TLK2-213, the

annotated TLK2 transcript that lacks the NLS (Fig. 1B). Instead, the three smallest PCR

products represented novel isoforms lacking exons 3, 4 and 5, with alternative splicing of exons

2, 7 and 8 (Fig. 3B). All three of these transcripts are predicted to be protein coding and lack

the NLS. HEK293 cells possessed very low levels of these PCR products.

We observed a striking regulation of TLK2 isoform expression during neuronal differentiation

of the SH-SY5Y cells (Fig. 3B). In undifferentiated cells the TLK2 isoforms are dominated by

the long transcripts (TLK-201/203; TLK-202/204; TLK2-209/227/230) and the longest of the

short isoforms that lacks the NLS (TLK2-novel-a). After the addition of RA, the expression of

the long TLK-202/204 isoform is rapidly downregulated (Fig. 3B,C). The subsequent addition

of BDNF in serum-free media induced an upregulation of the shortest variants (TLK2-novel-b

and -c) and a concomitant downregulation of the longer isoforms. At the end of the

differentiation protocol, the TLK2 isoform profile was a broader mix of long and short variants

with similar levels of expression (Fig. 3B).

To establish the corresponding pattern of TLK2 isoform protein expression during

differentiation, SH-SY5Y whole cell lysates from the timecourse experiment were subjected to

immunoblotting with a C-terminal reactive antibody, for which the epitope is predicted to be

present in all major splice variants of TLK2 (Fig. 4A). Based on the predicted molecular weight

of the proteins encoded by the TLK2 transcripts identified in SH-SY5Y cells (see Fig. 3B), we

expected that the long isoforms would migrate in the range 78-87 kDa, while the shorter

isoforms lacking the NLS would migrate at 68-75 kDa. Indeed, two groups of bands were

observed at approximately 85 kDa and 70 kDa, and their expression was regulated during SH-

SY5Y differentiation, broadly correlating with the expression of the long and short TLK2-

transcripts observed in Figure 3. Quantification of replicate immunoblots by densitometry

revealed that the 85 kDa band had a biphasic expression during differentiation, which reduced

during RA treatment and then stabilised during BDNF treatment (Fig. 4A,B). The ~70 kDa band

increased in expression during differentiation, with this band becoming the dominant isoform

in the fully differentiated cells (Fig. 4A,B). Taken together these data suggest that neuronal

differentiation enhances expression of a TLK2 isoform that lacks the NLS and is predicted to
be cytoplasmic.

Cytoplasmic localisation of TLK2 in differentiated neurons
Immunofluorescent staining of SH-SY5Y cells at 1, 4, 9 and 11 days of differentiation with the
TLK2 C-terminal antiserum, revealed a time-dependent relocalisation of TLK2 from the nucleus
to the cytoplasm (Fig. 4C). We quantified its subcellular localisation by measuring the intensity
of TLK2 staining in regions of interest within the cytoplasm and nucleus of the cells. The plot
of log2(cytoplasm/nucleus) in Figure 4D revealed that TLK2 is significantly enriched in the
cytoplasm at day 11 compared to day 1.

To confirm that TLK2 proteins encoded by splice variants lacking the NLS-containing exon 3
localise to the cytoplasm, we generated stable tet-inducible SK-N-SH T-REx-Flp-in cell lines
that express either TLK2(180-722)-FLAG, corresponding to the predicted protein encoded by
Ensembl isoform TLK2-213 or full length TLK2-FLAG corresponding to the the protein product
of the canonical Ensembl TLK2 full length isoform (TLK2-202). Following a 48 h induction of
TLK2 expression with doxycycline in these cell lines, we performed a crude subcellular
fractionation (Fig. 5B). The cells were subjected to hypertonic lysis and we isolated the pellet
(enriched in nuclei) and post-nuclear supernatant. Immunoblotting with antibodies raised to
compartment markers, histone H3 and GAPDH, confirmed enrichment of nuclear and
cytoplasmic proteins respectively (Fig. 5B). The immunostaining of anti-FLAG of the
subcellular fractions and in fixed cells demonstrated that the N-terminal truncated kinase was
exclusively localised to the cytoplasm, while the full length kinase is predominantly nuclear
(Fig. 5A,B). Taken together we propose that a cytoplasmic pool of TLK2 increases during
neuronal differentiation due to two phenomena: an increase in the expression of a short TLK2
splice variant lacking the NLS and the export of full length TLK2 from the nucleus.

Nuclear-cytoplasmic shuttling of TLK2
The nuclear-cytoplasmic transport of TLK2 has been previously reported by Zhang et al (1999)
who discovered a yeast 2-hybrid interaction between a 14-3-3ζ bait and a C-terminal clone of
TLK2 (amino acids 657-750). They found that 14-3-3ζ immunoprecipitated from cell lysates
with TLK2 and a dominant negative 14-3-3 promoted the nuclear localisation of TLK2 (Zhang
et al., 1999). Since 14-3-3 proteins act by binding a phosphorylated motif (Yaffe et al., 1997),
we hypothesised that during neuronal differentiation, phosphorylation of TLK2 recruits 14-3-3,
thus preventing nuclear import of TLK2 and restricting it to the cytoplasm. Using
undifferentiated SH-SY5Y cells, we attempted to mimic these signals to drive cytoplasmic
localisation of endogenous TLK2 by stimulating kinase signalling with 100 mM KCl and 10 uM

forskolin for 1 h. 100 mM KCl is known to depolarise SH-SY5Y cells, mimicking synaptic

activity, and should activate Ca^{2+}-dependent intracellular signalling in addition to the MAP

kinase pathway (Corrales et al., 2005). Forskolin, an adenylyl cyclase agonist, stimulates

cAMP production and hence activates signalling downstream of PKA (cAMP-dependent

protein kinase). Cells treated with KCl and forskolin together caused TLK2 to translocate from

the nucleus to the cytoplasm as visualised by immunofluorescence (Fig. 5C) and quantified by

log2(cytoplasm/nucleus) of TLK2 intensity (Fig. 5D). These data suggest the

nucleocytoplasmic shuttling of TLK2 that occurs during neuronal differentiation might also be

regulated in neurons by activity-dependent signalling.

Discussion

We have undertaken the first characterisation of neuronal TLK2 expression and localisation.

We found that TLK2 is expressed in the hippocampus and cerebellum, brain regions that are

affected in neurodevelopmental disorders linked with disrupted motor function and cognition.

In addition to canonical full length TLK2, we observed splice diversity at the N-terminus of the

protein, including variants that lack an NLS. Furthermore, cytoplasmic TLK2 increased during

neuronal differentiation, driven by both enhanced expression of TLK2 variants lacking the

NLS, and nuclear export of full length TLK2. We also found that the localisation of TLK2 can

be regulated acutely by mimicking synaptic activity. Our data suggest future studies into the

pathological mechanisms of MRD57 should consider cytoplasmic TLK2 signalling in

developing and mature neurons.

Expression and alternative splicing of neuronal TLK2

Our analysis of TLK2 transcripts in mouse brain, SH-SY5Y cells, and publicly available

RNAseq datasets revealed broad conservation of full length TLK2 and shorter transcripts

lacking N-terminal exons. Interestingly the canonical, and longest, human Ensembl isoform,

TLK2-201, comprising 22 exons is weakly expressed, except in aorta and testis (Supp Fig.

1), while the most abundant variant in brain and across the body is TLK-202, which lacks

exons 5 and 12. Indeed the other major isoforms that are expressed (TLK-203 and TLK-213)

also lack exons 5 and 12, calling into question the physiological significance of these exons.

TLK-213 is the major annotated short isoform that lacks the N-terminus and NLS and is the

human homologue of the short mTLK2-207 isoform that was first identified in mouse testis

(Shalom and Don, 1999). However, when we assessed the splice diversity of TLK2

transcripts in SH-SY5Y cells during neuronal differentiation we failed to observe TLK2-213,

but instead discovered several novel isoforms that lack the NLS encoded by exon 3. These

variants were only weakly detected in HEK293 cDNA, suggesting they might be upregulated

in neurons. When we immunoblotted TLK2 in SH-SY5Y cells using an antibody with a C-

terminal epitope, we observed bands that broadly correspond to long and short isoforms.

However, due to the minor differences in molecular mass arising from the various splicing

events it is not possible to correlate specific protein bands with specific splice variants.

Interestingly, human TLK1, which shares 78% homology with full length TLK2, also

possesses a short isoform lacking the N-terminus, and hence the NLS. Constitutive

expression of TLK1B is highest in testis, liver and lung, but low in brain (Li et al., 2001).

Furthermore, TLK1B expression can be induced by genotoxic stress, such as ionising

radiation, through translational regulation of an alternative start site (Li et al., 2001).

Surprisingly, in contrast to short TLK2, TLK1B localises to the nucleus where it coordinates

DNA repair, even in the absence of the TLK1 NLS (Li et al., 2001; Sunavala-Dossabhoy et

al., 2005), suggesting TLK1B is targeted to the nucleus by an alternative mechanism.

Using splice junction-specific *in situ* probes, we were able to examine the relative

distributions of long and short TLK2 isoforms in the mouse brain. TLK2 was expressed in the

major cell types of the hippocampus and cerebellum, although it was notable that expression

in excitatory neurons was easily observed, but inhibitory interneurons were not detectably

stained. This is in contradiction to single cell RNAseq analyses of mouse hippocampus and

cerebellum, which detected abundant TLK2 transcripts in various interneuron cell types

(Habib et al., 2016; Kozareva et al., 2021). We observed differential expression of TLK2

isoforms in excitatory cell types of the hippocampal formation, with intriguing differences

between CA1 and CA3 intensity (mTLK2-204) and the blades that comprise the dentate

gyrus (mTLK2-201). It is not surprising that there are transcriptional differences between

hippocampal cell types, which presumably define their function. However, a deeper

understanding of the role of TLK2 in the developing and mature brain is needed to predict the

significance of these data.

Neuronal differentiation- and activity-dependent nucleocytoplasmic shuttling of TLK2

We observed a striking loss of nuclear TLK2 staining in SH-SY5Y cells during neuronal

differentiation. Since we did not observe a significant concomitant reduction in protein

expression of longer TLK2 isoforms, we concluded that TLK2 was shuttling from the nucleus

to the cytoplasm during differentiation. The upregulation of short TLK2 isoform expression we

observed is also likely to contribute to the enhancement of cytoplasmic TLK2. This was

validated by creating stable SK-N-SH cells cell lines expressing an N-terminally truncated

TLK2 variant, which was confined to the cytoplasm. These data agree with the non-nuclear

localisation of a short TLK2 variant that was overexpressed in AD293 cells (Mortuza et al.,

2018).

Our observation of long TLK2 export from the nucleus during differentiation is consistent with

previous data on the cytoplasmic and nuclear distribution of TLK2. Using a C-terminal TLK2

monoclonal antibody, Zhang et al (1999) demonstrated cell cycle-dependent shuttling of

TLK2 in cultures of synchronised 3T3 cells. During G1 phase, TLK2 was predominantly

outside the nucleus and co-localised with vimentin, an intermediate filament protein. With the

onset of S-phase TLK2 was observed to localise to the nuclear periphery and then within the

nucleus during late G2 (Zhang et al., 1999). Thus, the nuclear export and cytoplasmic

accumulation of TLK2 during neuronal differentiation correlates with these cells exiting the

cell cycle. Mechanistically, Zhang et al. attributed the nucleocytoplasmic shuttling of TLK2 to

an interaction with 14-3-3 proteins, with binding preventing TLK2 nuclear import. They

identified a putative 14-3-3 interaction motif in the C-terminus (amino acids 747-753). Serine

phosphorylation of 14-3-3 motifs is required for binding, but the phosphorylation of that motif

in TLK2 has not been characterised. Since we observed translocation of TLK2 in response to

KCl-induced depolarisation and FSK, it is possible that phosphorylation of TLK2 by a Ca^{2+}-

and/or cAMP-dependent kinase facilitates 14-3-3 binding and hence retains it in the

cytoplasm. There are examples in the literature of transcription factors that require neuronal

activity-dependent Ca^{2+} and cAMP signalling for nucleocytoplasmic shuttling, such as the

transcriptional activator CRTC1 (Ch'ng et al., 2012). The functions of non-nuclear TLK2 have

not been investigated, although its colocalisation with vimentin might suggest regulation of

the cytoskeleton (Zhang et al., 1999). This is in-line with the Tousled homologue in

Drosophila, which has been shown to genetically interact with *rho1* (Yeh et al., 2015)*,* a

cytoskeletal regulator, and the *tlk* mutant has altered microtubule and actin filament density

in developing follicle cells (Gregory et al., 2007). Other cytoplasmic TLK2 interactors are

lacking. Previous TLK2 immunoprecipitation or proximal interactome studies have been

conducted in non-neuronal proliferating cell lines, where it might be expected that TLK2 is

predominantly nuclear, and these have tended to identify nuclear binding partners (Segura-

Bayona et al., 2017; Pavinato et al., 2022).

The implications of neuronal TLK2 splice variants for MRD57

Our study raises mechanistic questions about the pathology of the neurodevelopmental

disorder MRD57, in which patients are haploinsufficient for TLK2, predominantly through *de*

novo mutations (Lelieveld et al., 2016; Reijnders et al., 2018; Pavinato et al., 2022; Woods et

al., 2022). The important role of TLK2 in cell cycle regulation and DNA repair has led to the

hypothesis that the proliferation and differentiation of neural progenitors is affected in MRD57

patient brain development (Segura-Bayona and Stracker, 2019). MRD57 phenotypes are

presumably due to processes involving TLK2 that are not compensated by TLK1, such as

non-nuclear functions in post-mitotic neurons. Surprisingly, the TLK2 knockout mouse did not

display any cell cycle defects, due to compensation by TLK1, except in the placenta where

TLK1 expression is low (Segura-Bayona et al., 2017). However, the knockout mouse might

not be a good model for MRD57 as there were no reported brain or behavioural anomalies in

the TLK2 conditional heterozygous or homozygous mutant mice (Segura-Bayona et al.,

2017).

It is notable that MRD57 patient missense mutations have not been identified in exons 2, 4-7

or 12. This suggests that these regions are either not necessary for TLK2 function, or their

mutation is tolerated through compensation by alternative splice variants lacking these

exons. Relevant to our study, one patient has been identified with a homozygous substitution

mutation (K55E) adjacent to the NLS (Töpf et al., 2020), and it would be interesting to study

the localisation and nuclear shuttling of this TLK2 mutant in neurons. Furthermore, future

studies of neuronal TLK2 should focus on identifying its neuronal cytoplasmic substrates to

advance our understanding of TLK2 in developing and mature neurons and the pathology of

MRD57.

References

Bhoir S, De Benedetti A (2023) Targeting prostate cancer, the "tousled way." Int J Mol Sci 24:11100.

Bruinsma W, van den Berg J, Aprelia M, Medema RH (2016) Tousled-like kinase 2 regulates recovery from a DNA damage-induced G2 arrest. EMBO Rep 17:659–670.

Ch'ng TH, Uzgil B, Lin P, Avliyakulov NK, O'Dell TJ, Martin KC (2012) Activity-dependent transport of the transcriptional coactivator CRTC1 from synapse to nucleus. Cell 150:207–221.

Corrales A, Montoya G JV, Sutachan J-J, Cornillez-Ty G, Garavito-Aguilar Z, Xu F, Blanck TJJ, Recio-Pinto E (2005) Transient increases in extracellular K+ produce two pharmacological distinct cytosolic Ca2+ transients. Brain Res 1031:174–184.

Encinas M, Iglesias M, Liu Y, Wang H, Muhaisen A, Ceña V, Gallego C, Comella JX (2000) Sequential treatment of SH-SY5Y cells with retinoic acid and brain-derived neurotrophic factor gives rise to fully differentiated, neurotrophic factor-dependent, human neuron-like cells. J Neurochem 75:991–1003.

Fisher ME, Isaacs HV, Pownall ME (2002) eFGF is required for activation of XmyoD expression in the myogenic cell lineage of Xenopus laevis. Development 129:1307–1315.

Gregory SL, Shandala T, O'Keefe L, Jones L, Murray MJ, Saint R (2007) A Drosophila overexpression screen for modifiers of Rho signalling in cytokinesis. Fly 1:13–22.

Habib N, Li Y, Heidenreich M, Swiech L, Avraham-Davidi I, Trombetta JJ, Hession C, Zhang F, Regev A (2016) Div-Seq: Single-nucleus RNA-Seq reveals dynamics of rare adult newborn neurons. Science 353:925–928.

Harrison PW et al. (2024) Ensembl 2024. Nucleic Acids Res 52:D891–D899.

Kim J-A, Anurag M, Veeraraghavan J, Schiff R, Li K, Wang X-S (2016a) Amplification of TLK2 Induces Genomic Instability via Impairing the G2-M Checkpoint. Mol Cancer Res 14:920–927.

Kim J-A, Tan Y, Wang X, Cao X, Veeraraghavan J, Liang Y, Edwards DP, Huang S, Pan X, Li K, Schiff R, Wang X-S (2016b) Comprehensive functional analysis of the tousled-like kinase 2 frequently amplified in aggressive luminal breast cancers. Nat Commun

7:12991.

Klimovskaia IM, Young C, Strømme CB, Menard P, Jasencakova Z, Mejlvang J, Ask K,
Ploug M, Nielsen ML, Jensen ON, Groth A (2014) Tousled-like kinases phosphorylate
Asf1 to promote histone supply during DNA replication. Nat Commun 5:3394.

Kozareva V, Martin C, Osorno T, Rudolph S, Guo C, Vanderburg C, Nadaf N, Regev A,
Regehr WG, Macosko E (2021) A transcriptomic atlas of mouse cerebellar cortex
comprehensively defines cell types. Nature 598:214–219.

Lee S-B, Segura-Bayona S, Villamor-Payà M, Saredi G, Todd MAM, Attolini CS-O, Chang T-
Y, Stracker TH, Groth A (2018) Tousled-like kinases stabilize replication forks and show
synthetic lethality with checkpoint and PARP inhibitors. Sci Adv 4:eaat4985.

Lelieveld SH et al. (2016) Meta-analysis of 2,104 trios provides support for 10 new genes for
intellectual disability. Nat Neurosci 19:1194–1196.

Li Y, DeFatta R, Anthony C, Sunavala G, De Benedetti A (2001) A translationally regulated
Tousled kinase phosphorylates histone H3 and confers radioresistance when
overexpressed. Oncogene 20:726–738.

López-Rivera JA, Pérez-Palma E, Symonds J, Lindy AS, McKnight DA, Leu C, Zuberi S,
Brunklaus A, Møller RS, Lal D (2020) A catalogue of new incidence estimates of
monogenic neurodevelopmental disorders caused by de novo variants. Brain 143:1099–
1105.

Mortuza GB, Hermida D, Pedersen A-K, Segura-Bayona S, López-Méndez B, Redondo P,
Rüther P, Pozdnyakova I, Garrote AM, Muñoz IG, Villamor-Payà M, Jauset C, Olsen JV,
Stracker TH, Montoya G (2018) Molecular basis of Tousled-Like Kinase 2 activation. Nat
Commun 9:2535.

Pavinato L et al. (2022) Functional analysis of TLK2 variants and their proximal interactomes
implicates impaired kinase activity and chromatin maintenance defects in their
pathogenesis. J Med Genet 59:170–179.

Reijnders MRF et al. (2018) De Novo and Inherited Loss-of-Function Variants in TLK2:
Clinical and Genotype-Phenotype Evaluation of a Distinct Neurodevelopmental Disorder.
Am J Hum Genet 102:1195–1203.

Roe JL, Rivin CJ, Sessions RA, Feldmann KA, Zambryski PC (1993) The Tousled gene in A.
thaliana encodes a protein kinase homolog that is required for leaf and flower

development. Cell 75:939–950.

RStudio Team (2020) RStudio Integrated Development for R. Available at:
636   http://www.rstudio.com/. [Accessed November 5, 2024].

637   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
638   Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
639   Tomancak P, Cardona A (2012) Fiji: an open-source platform for biological-image
640   analysis. Nat Methods 9:676–682.

641   Segura-Bayona S, Knobel PA, González-Burón H, Youssef SA, Peña-Blanco A, Coyaud É,
642   López-Rovira T, Rein K, Palenzuela L, Colombelli J, Forrow S, Raught B, Groth A, de
643   Bruin A, Stracker TH (2017) Differential requirements for Tousled-like kinases 1 and 2 in
644   mammalian development. Cell Death Differ 24:1872–1885.

645   Segura-Bayona S, Stracker TH (2019) The Tousled-like kinases regulate genome and
646   epigenome stability: implications in development and disease. Cell Mol Life Sci
647   76:3827–3841.

648   Shalom S, Don J (1999) Tlk, a novel evolutionarily conserved murine serine threonine
649   kinase, encodes multiple testis transcripts. Mol Reprod Dev 52:392–405.

650   Silljé HH, Takahashi K, Tanaka K, Van Houwe G, Nigg EA (1999) Mammalian homologues
651   of the plant Tousled gene code for cell-cycle-regulated kinases with maximal activities
652   linked to ongoing DNA replication. EMBO J 18:5691–5702.

653   Sjöstedt E et al. (2020) An atlas of the protein-coding genes in the human, pig, and mouse
654   brain. Science 367:eaay5947.

655   Sunavala-Dossabhoy G, Balakrishnan SK, Sen S, Nuthalapaty S, De Benedetti A (2005) The
656   radioresistance kinase TLK1B protects the cells by promoting repair of double strand
657   breaks. BMC Mol Biol 6:19.

658   Töpf A, Oktay Y, Balaraju S, Yilmaz E, Sonmezler E, Yis U, Laurie S, Thompson R, Roos A,
659   MacArthur DG, Yaramis A, Güngör S, Lochmüller H, Hiz S, Horvath R (2020) Severe
660   neurodevelopmental disease caused by a homozygous TLK2 variant. Eur J Hum Genet
661   28:383–387.

662   Villamor-Payà M, Sanchiz-Calvo M, Smak J, Pais L, Sud M, Shankavaram U, Lovgren AK,
663   Austin-Tse C, Ganesh VS, Gay M, Vilaseca M, Arauz-Garofalo G, Palenzuela L,
664   VanNoy G, O'Donnell-Luria A, Stracker TH (2024) De novo TLK1 and MDM1 mutations

665   in a patient with a neurodevelopmental disorder and immunodeficiency. iScience

666   27:109984.

667  Woods E, Spiller M, Balasubramanian M (2022) Report of two children with global

668   developmental delay in association with de novo TLK2 variant and literature review. Am

669   J Med Genet A 188:931–940.

670  Yaffe MB, Rittinger K, Volinia S, Caron PR, Aitken A, Leffers H, Gamblin SJ, Smerdon SJ,

671   Cantley LC (1997) The structural basis for 14-3-3:phosphopeptide binding specificity.

672   Cell 91:961–971.

673  Yamakawa A, Kameoka Y, Hashimoto K, Yoshitake Y, Nishikawa K, Tanihara K, Date T

674   (1997) cDNA cloning and chromosomal mapping of genes encoding novel protein

675   kinases termed PKU-alpha and PKU-beta, which have nuclear localization signal. Gene

676   202:193–201.

677  Yeh T-H, Huang S-Y, Lan W-Y, Liaw G-J, Yu J-Y (2015) Modulation of cell morphogenesis

678   by tousled-like kinase in the Drosophila follicle cell. Dev Dyn 244:852–865.

679  Zhang S, Xing H, Muslin AJ (1999) Nuclear localization of protein kinase U-alpha is regulated

680   by 14-3-3. J Biol Chem 274:24865–24872.

Figure legends

Figure 1. TLK2 has splice variant diversity in the human brain. A, Exonic structure of

human TLK2, mapped onto a schematic representation of its protein domains. NLS, nuclear

localisation sequence; CC, coiled-coil domain **B**, Table of % TLK2 transcript abundance in the

human hippocampus and cerebellum derived from GTEx RNAseq data. Exons and transcripts

labelled in red are predicted to be non-protein coding. Only transcripts present >2% in either

brain region are included.

Figure 2. Multiple isoforms of TLK2 are expressed in mouse forebrain and cerebellum.

A, RT-PCR of TLK2 splice variants from mouse forebrain and cerebellum cDNA. Transcript

specific primer pairs or pan exon 1-6 primers. **B**, Left, lane from PCR of exon 1-6 primer pair.

The four bands were excised and sequenced. Right, exon structure of sequenced mouse

brain splice variants, and location of *in situ* hybridisation probes. *In situ* hybridisation was

performed on 20 µm frozen sections from coronal (**C**) or sagittal (**E**) P56 mouse brain

sections. The indicated 150 bp exon junction spanning probes were used to examine

calbindin or TLK2 transcript expression in the hippocampus (**C**) or cerebellar cortex (**E**).

Representative images are shown from n=3 sections from two mouse brains. Cerebellar

cortex images are shown at x10 and x40 magnification, indicated by the insert box. DG,

dentate gyrus; ML, molecular layer; GL, granule layer; PCL, Purkinje cell layer. Scale bars:

hippocampus 100 µm, cerebellar cortex x10, 200 µm; x40, 20 µm. **D**, Heatmap depicting

quantification of TLK2 splice variant expression within the dentate gyrus, CA1 and CA3 sub-

regions of the hippocampal formation. For each section, pixel intensity was measured in

regions of interest (ROI), as indicated in the cartoon and normalised to the ROI located in the

suprapyramidal blade of the dentate gyrus. Three sections were quantified for each probe.

Figure 3. Alternative splicing of TLK2 is regulated during neuronal differentiation. SH-

SY5Y cultures were differentiated by sequential treatment with retinoic acid (RA) and brain

derived neurotrophic factor (BDNF) as described in Materials and Methods. **A**, Brightfield

images of SH-SY5Y cultures at the indicated days of differentiation. Scale bar = 40 µm. **B**,

cDNA was prepared from SH-SY5Y cells at the indicated time points, and HEK293 cells

(control) and subjected to PCR with primers for exons 1 and 9 of human TLK2, Tau, and

GAPDH. The six TLK2 bands obtained from the cells were excised and subjected to Sanger

sequencing. The right hand schematic summarises the exon structure of each sequence and

any corresponding Ensembl transcripts that these align to and the predicted molecular mass

of the encoded protein. The bottom plot presents quantitative densitometry of each TLK2 RT-

PCR product, expressed as a percentage of the total intensity at each timepoint.

Figure 4. Neuronal differentiation alters the nuclear-cytoplasmic distribution of TLK2.

SH-SY5Y cells were lysed or fixed at the indicated timepoints during neuronal differentiation

and subjected to immunoblotting with anti-TLK2, anti-β3-tubulin or anti-actin (**A**) or

immunocytochemistry with anti-TLK2 (1:1000; green) and DAPI staining (purple) Scale bar =

20 µm. (**C**). **B**, Quantification of immunoblot in **A**. Plot of the intensity of TLK2 bands with

high (long TLK2) and low (short TLK2) mass, normalised to the intensity of actin and then

day 4 short TLK2 (n=3 independent experiments). **D**, TLK2 intensity was measured in

regions of interest in the nucleus and cytoplasm of 100 cells from each timepoint (n=3

independent experiments). Data are plotted as $\log_2$(nuclear/cytoplasmic) TLK2 intensity at

each timepoint (grey filled circles). Hollow black circles represent the mean of each replicate.

For B and D, data were analysed by Kruskal-Wallis two-tailed ANOVA and Dunn's post-hoc

test; * $p < 0.05$; ** $p < 0.01$ compared to day 1.

Figure 5. Activity-dependent redistribution of TLK2. Inducible Flp-in SK-N-SH cells

expressing FLAG-tagged long TLK2-203 or short TLK2-213 were treated with or without 1

µg/ml doxycycline (dox) for 48 h. **A**, Representative immunofluorescence images of dox-

treated cells stained with anti-FLAG (1:2000) and DAPI. Scale bar = 20 µm. **B**, Cells plated in

dishes were lysed and subject to subcellular fractionation to yield a nuclear pellet and

postnuclear supernatant. The samples were subjected to immunoblotting with anti-FLAG,

anti-GAPDH and anti-histone H3. **C**, SH-SY5Y cells equilibrated in KREBS buffer were

incubated for 1 h in KREBS buffer with or without (control) 100 mM KCl and 10 µM forskolin

(FSK). Fixed cells were stained with anti-TLK2 and DAPI. Scale bar = 20 µm. **D**, TLK2

intensity was measured in regions of interest in the nucleus and cytoplasm of 100 cells from

each treatment (n=3 experiments). Data are plotted as $\log_2$(nuclear/cytoplasmic) TLK2

intensity at each timepoint (grey filled circles). Hollow black circles represent the mean of

each replicate and were analysed by t- test; * $p < 0.05$ compared to control.

Figure 1

red exon/transcript = non-coding

Figure 2

750
751
752

753   **Figure 3**

756 **Figure 4**

759 **Figure 5**

761
762