Neuronal differentiation and activity drive nucleocytoplasmic shuttling of the intellectual disability kinase TLK2

Lubna Nuhu-Soso, Heidi Denton, Darren L Goffin, Ines Hahn, Gareth J.O. Evans[#]

Department of Biology and York Institute of Biomedical Sciences, University of York, Wentworth Way, Heslington, York, YO10 5DD, United Kingdom

[#]Corresponding author: gareth.evans@york.ac.uk

Keywords
kinase, neurodevelopmental disorder, nuclear localisation signal, neuronal differentiation, splice variant

Abstract

Mental retardation autosomal dominant 57 (MRD57) is a rare neurodevelopmental disorder characterised by delayed language and psychomotor development, intellectual disability, hypotonia, gastrointestinal issues and facial dysmorphia. It is linked to genetic mutations in the serine/threonine kinase TLK2, characterised by haploinsufficiency and in some cases, its loss or impaired kinase function. TLK2 is an established cell cycle regulator that has been extensively studied in mitotic cells. It is upregulated in cancers, driving tumour growth, however, the role of TLK2 in postmitotic neurons is not understood. We therefore aimed to gain insight into how TLK2 mutations cause MRD57 by determining where TLK2 is expressed in the brain and its subcellular localisation during neuronal differentiation. Public human and mouse brain transcriptomic data revealed splice variant diversity in the N-terminus of TLK2, which contains its nuclear localisation sequence (NLS). Using splice-specific *in situ* hybridisation probes, we observed expression of TLK2 isoforms that contain and lack the NLS in the mouse hippocampus and cerebellum. We confirmed these findings in human SH-SY5Y neuroblastoma cells, and found that neuronal differentiation of these cells enhances a cytoplasmic pool of TLK2 by two mechanisms: nuclear export of full length TLK2 and increased expression of TLK2 splice variants lacking the NLS. Finally, acute stimuli that mimic synaptic activity were sufficient to elicit nuclear export of TLK2. Our data highlight the need to establish the neuronal cytoplasmic substrates of TLK2 and determine how the loss of TLK2 activity in MRD57 might impact their function in the developing and mature brain.

21 **Introduction**
22
Tousled-like kinase 1 and 2 are the mammalian homologues of the *Arabidopsis* mutant,
tousled, which was discovered in a developmental screen (Roe et al., 1993). Indeed, both
human TLK1 and TLK2 have been linked with rare developmental disorders (Segura-Bayona
and Stracker, 2019; Villamor-Payà et al., 2024). To date, there have been close to 50
documented patients worldwide with TLK2 mutations, classified as autosomal dominant
Mental Retardation Disorder 57 (MRD57; (Lelieveld et al., 2016; Reijnders et al., 2018; Töpf
et al., 2020; Pavinato et al., 2022; Woods et al., 2022), with a predicted incidence of
approximately 3 in 100,000 individuals (López-Rivera et al., 2020). The clinical phenotypes
associated with MRD57, like other neurodevelopmental disorders, are heterogeneous,
including intellectual disability, delayed psychomotor development in infancy or early
childhood, language delay, hypotonia, feeding problems, gastrointestinal issues, dysmorphic
facial features, microcephaly, behaviour problems and global developmental delays
(Reijnders et al., 2018; Töpf et al., 2020; Pavinato et al., 2022; Woods et al., 2022). The
majority of TLK2 mutations in MRD57 are *de novo* mutations, but seven are inherited within
four different families (Reijnders et al., 2018; Töpf et al., 2020; Pavinato et al., 2022). Like the
de novo cases, inherited TLK2 mutations are heterozygous dominant except one report of a
homozygous recessive inheritance (Reijnders et al., 2018; Töpf et al., 2020; Pavinato et al.,
2022). MRD57 patients have TLK2 mutations throughout the protein and all are predicted to
result in a reduction of TLK2 expression or its catalytic activity (Mortuza etc).
42
Functionally, TLK2 is an established regulator of DNA replication, cell cycle checkpoint
recovery and chromatin remodelling (Bruinsma et al., 2016; Kim et al., 2016a; Segura-
Bayona et al., 2017; Silljé et al., 1999). TLK2 facilitates cell cycle progression mainly by
enhancing histone supply via phosphorylation of its best characterised substrate, ASF1
(Klimovskaia et al., 2014). In-line with other cell cycle kinases, TLK activity and gene copy
number are increased in cancer, including breast, cervix, lung, liver, colon and kidney (Bhoir
and De Benedetti, 2023; Kim et al., 2016b; Lee et al., 2018). The chromosomal locus of
TLK2, 17q23, contains other oncogenes, and is frequently amplified in more than 40% of
breast cancer tumours (Kelemen et al., 2009; Kim et al., 2016b). Moreover, single nucleotide
polymorphisms (SNPs) in TLK2 have been identified as risk factors for breast cancer
(Kelemen et al., 2009). Indeed, inhibition of TLK2 has been identified as a therapeutic
strategy and small molecule inhibitors are in development (Kim et al., 2016b; Lee et al.,
2018).
56
In contrast to its role in the cell cycle and cancer, the neuronal expression and cell biology of
TLK2 is poorly understood. Here we have characterised its neuronal transcript expression,
revealing a conserved splice diversity in rodents and humans. TLK2 alternative splicing
occurs in exons encoding the N-terminus, resulting in 'long' and 'short' isoforms, with the
latter lacking the nuclear localisation sequence (NLS). Expression of both long and short
TLK2 isoforms is prevalent in neurons of the mouse hippocampus and cerebellum. Using a
cell model of neuronal differentiation, we observed the upregulation of short TLK2 expression
during neuronal differentiation, with a concomitant shuttling of long TLK from the nucleus to
the cytoplasm. Pharmacological treatments mimicking neuronal activity also drive
nucleocytoplasmic shuttling of TLK2. These data highlight the likely importance of
cytoplasmic TLK2 in terminally differentiated neurons, which have exited the cell cycle. There

is hence a need to determine the neuronal substrates of TLK2 to shed light on the neuronal
phenotypes of MRD57.
70
71 **Materials and Methods**
72
73 **Materials**
74 SH-SY5Y and Flp-In T-REx SK-N-SH neuroblastoma cell lines were kind gifts from Dr
75 Andrew Stoker, University College London and Dr Han-Jou Chen, University of York
76 respectively. The Flp recombinase expression vector, pOG44 was a kind gift from Dr Paul
77 Pryor, University of York.
78
Bioinformatics
TLK2 transcript IDs and exon structures were derived from their Ensembl gene annotations
(version 113) using human (GRCh38.p14) or mouse (GRCm39) genome builds (Harrison et
al., 2024). The human TLK2 RNAseq data used for the transcriptomic analyses described in
this manuscript were obtained from the GTEx Portal, accession number phs000424.v8.p2.
Mouse brain TLK2 RNAseq data was obtained from the Human Protein Atlas
(proteinatlas.org; (Sjöstedt et al., 2020)). TLK2 transcript expression values in transcripts per
million (TPM) were converted to percentages by tissue or brain region.
87
RNA extraction, cDNA synthesis and RT PCR
All work involving mice was approved by the University of York Animal Welfare and Ethical
Review Body and performed under UK Home Office legislation (project licence P38B2013E).
Adult wild-type C57BL/6 mouse forebrain and cerebellum were dissected and homogenised
in RIPA buffer (50 mM Tris, pH 8.0, 150 mM NaCl, 1% (v/v) Triton-X-100, 0.5% (w/v) sodium
deoxycholate, 0.1% (w/v) SDS, 1 mM EDTA supplemented with 1 mM Na_3VO_4, 0.1% (v/v) β-
mercaptoethanol, 1 mM PMSF and 1:200 protease inhibitor cocktail (Sigma)) using 5 ml
buffer/g of tissue. The homogenate was incubated at 4°C with agitation for 1 h and then
centrifuged at 20,000g for 30 min at 4°C. RNA was extracted from the supernatant.
SH-SY5Y cells undergoing neuronal differentiation were plated at 10^4 cells/ cm^2 onto
collagen coated 6-well tissue culture dishes. RNA was extracted from the differentiating cells
at two day intervals. For both mouse brain and SH-SY5Ys, RNA was extracted using a
NucleoSpin RNA isolation kit (Macherey-Nagel) according to the manufacturer's instructions.
Complementary DNA was synthesised from 1 μg of RNA using a SuperScript IV First-Strand
cDNA Synthesis Kit (Life Technologies) according to the manufacturer's instructions in the
presence of both oligo $d(T)_{18}$ and random hexamer primers. A no RT cDNA sample was
performed for each time point and cDNA was stored at -20°C. PCR reactions were
performed in 25 μl reactions containing 1 μl of cDNA, 1X GoTaq flexi buffer, 0.2 mM each of
dNTP mix, 0.4 μM of each primer pair (Table 1) and 1U Taq polymerase. Cycling conditions
consisted of an initial denaturing step at 98°C for 30s followed by 25-30 cycles of 95°C for
10s, 55°C for 30s, 72°C for 1 min/kb and a final extension at 68°C for 10 min. 5 - 10 μl of the
PCR products were separated on 1.5-2% (w/v) agarose gels to confirm amplification.
110
***In situ* hybridisation**
Mice were anaesthetised with isoflurane inhalation followed by an intraperitoneal injection of
100 mg/kg ketamine and 12.5 mg/kg xylazine and then transcardially perfused with 4%
paraformaldehyde (PFA) in 0.1 M sodium-potassium PBS. The brain dissected and post-
fixed in 4% PFA overnight at 4°C. Following a further 24 h incubation in 30% sucrose, brains

were embedded in OCT, frozen at -80°C, and 25 μm coronal or sagittal sections were cut
using a Leica CM1950 cryostat. Probes for *in situ* hybridisation were transcribed using 10X
digoxigenin (DIG) RNA labelling mix (Roche) and T7 polymerase. For *in vitro* transcription,
template DNA was synthesised as gBlocks™ (IDT) and PCR amplified to introduce the T7
promoter sequence (Table 1). *In situ* hybridisation was performed according to (Fisher et al.,
2002) with adaptations for sections mounted on slides. Briefly, the sections were first
incubated in 4% PFA for 10 min at room temperature (RT) followed by two 5 min PBST
washes. Sections were then treated with 4 μg/ml proteinase K (Roche) in PBST for 10 min
and then washed twice in PBST for 5 min. A second fix in 4% (w/v) PFA was performed for
10 min, followed by two 5 min PBST washes. Next, the sections were acetylated using 0.5%
(v/v) acetic anhydride in 0.1 M triethanolamine, pH 7.8 for 10 min followed by two 5 min
PBST washes and finally a 15 min wash in 5X saline-sodium citrate buffer (SSC; Sigma).
Pre-hybridisation was carried out by incubating the sections in hybridisation buffer (50% (v/v)
formamide (Ambion, AM9342), 5X SSC, 100 μg/ml heparin, 1X Denhardt's, 0.1% (v/v)
Tween-20, 0.1% (w/v) CHAPS, 10 mM EDTA, 1 mg/ml total yeast RNA) containing no probe
at 65°C for 2-5 h in a humidified box. Hybridisation followed by incubating the sections with
300 ng/ml of probe solution (DIG prep in hybridisation buffer) overnight at 65°C in a
humidified box.
Post-hybridisation, the sections were subjected to several hot wash steps. First a 10 min
wash in hybridisation buffer containing no probe at 65°C, then a 1 h wash in 2X SSC
followed by a 1 h wash in 0.2X SSC, both at 65°C. The sections were then washed twice for
5 min in MABT (100 mM maleic acid, 150 mM NaCl, 0.1% (v/v) Tween-20, pH 7.8) at RT and
then blocked in blocking solution (2% (w/v) Boehringer Mannheim Blocking Reagent (Roche,
11096176001), 20% (v/v) heat-treated lamb serum (Sigma), 1X MAB) for 2 h at RT in a
humidified box. Next, the sections were incubated with anti-DIG-AP (Table 2) in blocking
solution overnight at 4°C in a humidified box. Post-antibody washes were carried out at RT
by washing the sections three times for 1 h in MABT and once in alkaline phosphatase (AP)
buffer (100 mM Tris, pH 9.5, 50 mM $MgCl_2$, 100 mM NaCl) for 10 min. The colour reaction
was initiated by adding a 1:3 dilution of BM purple (Roche, 11442074001) in AP buffer to the
sections and incubation at RT in a humidified box for 1-5 days. The reaction was stopped by
washing the sections twice for 10 min in PBST, fixing in 4% (w/v) PFA for 1 h or overnight at
4°C and a 15 min wash in PBST. The sections were air dried, mounted in aqueous mounting
media (Sigma, 324590) and stored at 4°C. Images were acquired on a Leica DM2500
microscope with a 10X objective (sagittal) and a Zeiss Stemi 508 stereo microscope
(coronal).
151
152   **Table 1. Primers and probes used in this study**

Name	Sequence (5'-3')
PCR primers	
mTLK2-exon1 sense	GACCCACGAAGGCAGGAATTA
mTLK2-exon5/6 anti	CTCGACGCGGTAAGGGATTG
mTLK2-exon4/6 anti	CTGCTCGACGCTCAAAGTAA
mTLK2-exon3/6 anti	GCTCGACGCTTGGCTAGT
mTLK2-exon6 anti	GCTGCACTGCCATCTAAACC
mGAPDH sense	CATCACTGCCACCCAGAAGACTG
mGAPDH anti	ATGCCAGTGAGCTTCCCGTTCAG
hTLK2-exon1 sense	ATTGCATAGCCTGGACCCAC

hTLK2-exon9 anti	TCGTTCCTTGTATTTCTCTAGCA
hMAPT sense	ACACCATGCACCAAGACCAA
hMAPT anti	TCCTTCTGGGATCTCCGTGT
hGAPDH sense	GTCTCCTCTGACTTCAACAGCG
hGAPDH anti	ACCACCCTGTTGCTGTAGCC

ISH probes

mTLK2-204	GGAACCAGCCCTGGCAGAAGTGTTCCACCAGTTGCACGATCCTCACCGCA ACATTCCTTATCCAATCCCTTACCGCGTCGAGCAGAACAGCCTCTGTATG GTTTAGATGGCAGTGCAGCAAAGGAGGCCTCAGAAGAGCAGTCTGCTCTG
mTLK2-201	AGGAAAGCCGAGCCATATGACACTAGCCAAGGGAAAGGCACTCCTAGGGG ACATAAAATTAGTGATTACTTTGAGCGTCGAGCAGAACAGCCTCTGTATG GTTTAGATGGCAGTGCAGCAAAGGAGGCCTCAGAAGAGCAGTCTGCTCTG
mTLK2-207	TAGAGACTCCTGAGAAAAAGCAGAATGACCAGCGAAATCGGAAAAGGAAA GCCGAGCCATATGACACTAGCCAAGCGTCGAGCAGAACAGCCTCTGTATG GTTTAGATGGCAGTGCAGCAAAGGAGGCCTCAGAAGAGCAGTCTGCTCTG
mTLK2-novel	CAACCAGAGTCTGTGCAGCGTGGGGTCGTTGAGTGATAAAGAAGTAGAGA CTCCTGAGAAAAAGCAGAATGACCGCGTCGAGCAGAACAGCCTCTGTATG GTTTAGATGGCAGTGCAGCAAAGGAGGCCTCAGAAGAGCAGTCTGCTCTG
mCalb1	GATGCTTTGCTGAAAGATCTGTGTGAGAAGAACAAACAGGAATTGGATAT TAACAATATTACTACATACAAGAAGAACATAATGGCCTTGTCGGATGGAG GGAAGCTGTACCGAACAGACCTTGCTCTTATTCTTTCTGCTGGAGACAAC

Culture, neuronal differentiation and pharmacological treatment of SH-SY5Y cells
SH-SY5Y cells were maintained at 37°C, 5% CO_2 in a humidified atmosphere in Dulbecco's
Modified Eagle Medium (DMEM) with high glucose, pyruvate and L-glutamine supplemented
with 10% foetal bovine serum (FBS) and 1% penicillin/streptomycin.
Neuronal differentiation of SH-SY5Y cells was performed according to a previously described
method (Encinas et al., 2000). Briefly, tissue culture dishes or coverslips were coated in 0.05
mg/ml collagen in 60% (v/v) ethanol. Cells were plated at 10^4 cells/cm^2 (for protein analysis)
or 5 x 10^3 cells/well (for microscopy) in DMEM supplemented with 15% (v/v) heat inactivated
FBS and 1% (v/v) penicillin/streptomycin. Twenty four hours after plating, the media was
changed and supplemented with 10 µM retinoic acid (RA; Sigma, R2625). Six days after
plating, the cells were washed three times in serum-free DMEM and then incubated with
serum-free DMEM containing 50 ng/ml brain-derived neurotrophic factor (BDNF; Qkine) and
1% (v/v) penicillin/streptomycin and cultured for a further seven days. Throughout the
differentiation process, the media was changed every two days and supplemented with the
appropriate differentiation agent.
For analysing protein expression in differentiating cells, lysates were collected at two day
intervals. Cells were washed twice in sterile ice-cold PBS and scraped in 100 µl of lysis
buffer (20 mM HEPES, pH 7.4, 10 mM KCl, 2 mM $MgCl_2$, 1 mM EDTA and 1 mM EGTA, 1
mM DTT and protease inhibitor cocktail). Lysates were incubated on ice for 15 min, passed
through a 25G needle ten times and protein concentration was determined by Bradford
assay. The lysates were diluted in Laemmli sample buffer, boiled for 10 min at 90°C and 10
µg of protein was separated by SDS-PAGE and analysed by Western blotting.
Pharmacological stimulation of undifferentiated SH-SY5Y cells was performed on cells plated
24 h prior to treatment on 13 mm coverslips in a 24-well plate at 5 x 10^4 cells/well. Cells were
washed twice in PBS and then incubated for 1 h at 37°C, 5% CO_2 in either control buffer (170
mM NaCl, 3.5 mM KCl, 0.4 mM KH_2PO4, 10 mM HEPES, 5 mM $NaHCO_3$, 5 mM glucose, 1.2
mM Na_2SO_4, 1.2 mM $MgCl_2$, 1.3 mM $CaCl_2$, pH 7.4, 0.2% (v/v) DMSO) or KCl/FSK buffer

(control buffer with 73.5 mM NaCl, 100 mM KCl, 10 µM forskolin, 100 µM IBMX, 0.2% (v/v)
DMSO). Cells were then fixed and immunocytochemistry was performed as described below
using anti-TLK2.
184

Inducible stable TLK2 cell lines
Tetracycline-inducible SK-N-SH cell lines expressing either full length TLK2-203 or short N-
terminally truncated TLK2-213 were generated using the Flp-In™ T-REx™ System
(Invitrogen). pcDNA5/FRT/TO-FLAG-TLK2 was purchased from the MRC PPU Reagents
and Services facility (College of Life Sciences, University of Dundee). TLK2-213 was made
by subcloning its ORF from full length TLK2 and ligating it back into the same backbone via
5' and 3' NotI restriction sites with the N-terminal FLAG tag intact. Cells were co-transfected
with the FRT backbone containing the gene of interest (TLK2-203 or TLK2-213) and a Flp-
recombinase expression vector (pOG44). Positive clones were selected by treating with 0.5
mg/ml hygromycin B and assaying for protein expression after the addition of 1 µg/ml
doxycycline.
196

Subcellular fractionation
Flp-In SK-N-SH cells were fractionated according to the Abcam subcellular fractionation
protocol. TLK2 expression was induced using 1 µg/ml doxycycline and left in culture for 48 h
prior to fractionation. Briefly, cells were scraped in 500 µl of lysis buffer (described above)
and incubated on ice for 15 min. The scraped cells were passed through a 25G needle ten
times, incubated on ice for 20 min and then centrifuged at 720g for 5 min at 4°C. The post-
nuclear supernatant was transferred to a fresh tube. The nuclear pellet was washed by first
resuspending in lysis buffer, passing the sample through a 25G needle ten times and
centrifuging at 720g for 10 min at 4°C. The washed pellet was resuspended in TBS
containing 0.1% (w/v) SDS, and sonicated on ice for 3 s. The protein concentration of each
fraction was determined by Bradford assay and 20 µg of each fraction was separated by
SDS-PAGE and analysed by Western blotting.
209

Western blotting
Proteins separated by SDS-PAGE (7.5-12%) gels were transferred onto PVDF membranes
(Millipore). Membranes were incubated with primary antibodies (Table 2) overnight at 4°C
with agitation. Membranes were washed and incubated with appropriate HRP-conjugated
secondary antibodies and immunoreactivity was visualised by incubation with enhanced
chemiluminescence reagent and imaged on an iBright FL1000 scanner (Invitrogen).
216

Immunocytochemistry and fluorescence microscopy
Cells were washed 3x in PBS and then fixed in 4% (w/v) paraformaldehyde (PFA) in PBS for
20 min at RT. Next cells were washed 3x in PBS then permeabilised and blocked in PBS
containing 0.1% (v/v) TritonX-100 and 1% (w/v) bovine serum albumin (BSA). Primary
antibodies were diluted in PBS containing 1% (w/v) BSA at appropriate dilutions (Table 2)
and incubated on cells overnight at 4°C. Cells were washed 3x in PBS then incubated with
secondary antibodies in PBS containing 1% (w/v) BSA for 1 h in the dark at RT. Finally, cells
were washed 3x in PBS, 1x in dH$_2$O, air dried and mounted in Mowial containing 1 µg/ml
4',6-diamidino-2-phenylindole (DAPI). Images were acquired in the Bioscience Technology
Facility (University of York) on a Zeiss LSM 880 confocal microscope on a 40x/1.4 Oil DIC III
objective.
228

229 **Table 2. Antibodies used in this study.**

Name	Dilution	Species	Catalogue no.	Supplier
Primary antibodies				
Anti-TLK2	1:1000 (WB) 1:500 (IF)	Rabbit	A301-257A	Bethyl
Anti-β-III-tubulin	1:1000 (WB) 1:500 (IF)	Mouse	801202	Biolegend
Anti-β-actin	1:50000 - 1:100000	Mouse	60008-1-Ig	Proteintech
Anti-Histone H3	1:5000	Mouse	819411	Biolegend
Anti-GAPDH	1:5000	Mouse	MAB374	Sigma
Anti-FLAG	1:1000	Mouse	F1804	Sigma
Anti-Digoxigenin	1:2000	Sheep	11093274910	Roche
Secondary antibodies				
Anti-mouse HRP	1:5000	Goat	A4416	Sigma
Anti-rabbit HRP	1:5000	Goat	A6154	Sigma
Anti-mouse Alexa 594	1:500	Goat	A11020	Invitrogen
Anti-rabbit Alexa 488	1:500	Goat	A11008	Invitrogen

Data analysis
Quantification of hippocampal *in situ* hybridization images was performed on 16 bit grayscale
images in FIJI (Schindelin et al., 2012). The oval selection tool was used to draw regions of
interest (ROI) in different locations within the hippocampal formation (CA1, CA3, DGs and
DGi). The intensity was then measured and normalised to DGs for each section and probe.
Three to four hippocampi were quantified for each probe and the values were plotted as a
heatmap in Morpheus (https://software.broadinstitute.org/morpheus).
The quantification of cytoplasmic/nuclear TLK2 ratios of immunofluorescent images was
conducted by measuring the TLK2 intensity in the nucleus and cytoplasm of the cells in FIJI
using the ROI 1-click plug-in (Thomas and Gehrig 2020). Three hundred cells for each
condition were analysed from three technical replicates (100 cells/replicate). Western blots
were quantified by densitometry analysis in FIJI. Statistical significance was assessed by t-

test or non-parametric ANOVA with Dunn's post-hoc test. Statistical analyses and graph
plotting was performed in RStudio (RStudio Team, 2020).
245
246 **Results**
247
TLK2 is alternatively spliced in human and mouse brain
Since the neuronal distribution and function of TLK2 has not been investigated, we first sought
to characterise TLK2 transcript expression in the brain. Previous studies have observed tissue-
specific transcripts of TLK2, in the mouse testis for example (Shalom and Don, 1999), and
TLK2 immunoblots often have multiple bands that vary between tissues and cell lines (Kim et
al., 2016b; Segura-Bayona et al., 2017). Such bands could result from post-translational
modifications or alternative splicing of TLK2 or a combination. We began by interrogating
publicly available human (GTEx) and mouse brain (Human Protein Atlas) transcriptomic data
to characterise neuronal TLK2 transcript expression (Figure 1A,B), focussing on the
hippocampus and cerebellum. In the mouse brain, a full length transcript corresponding to the
canonical Ensembl TLK2 transcript was the most abundant transcript (mTLK2-204; mouse
TLK2 transcripts will be referred to here as 'mTLK2'). In the human brain, surprisingly the most
abundant transcript was not canonical TLK2 (TLK2-203), but a transcript lacking exon 5 (TLK2-
202; Figs. 1B, S1 & S2). We also identified further evolutionarily conserved splice variant
diversity at the N-terminus of the protein, encoded by exons 1-9. Not surprisingly, there were
fewer alternative splicing events in exons encoding the middle coiled-coil domain and C-
terminal kinase domain of TLK2, which are essential for dimerisation and catalytic activity
respectively (Mortuza et al., 2018); Figure 1B). Interestingly, splice variants lacking exon 3 in
which the nuclear localisation sequence (NLS, Figure 1A) is located (Yamakawa et al., 1997)
are also abundant (Figure 1B,S1). This suggests that a pool of TLK2 is constitutively resident
outside the nucleus. Henceforth in this study we shall refer to isoforms that include the NLS as
'long TLK2', and those lacking the NLS as 'short TLK2'. In the human GTEx dataset it was
observed that transcript TLK2-207 represents nearly a third of TLK2 mRNA expression, but is
predicted to be subject to nonsense mediated decay due to an intron retention at exon 21
(Figure 1B).
273
To confirm the existence of alternative N-terminal TLK2 splice variants in the brain, we
performed RT-PCR of mouse brain TLK2 mRNA. The annotated transcripts of mouse TLK2
have splice diversity within exons 1-6 (Figure 2A and Figure S2). We first amplified and
sequenced all N-terminal splice variants in mouse forebrain or cerebellar cDNA using
transcript-specific primers and also a pair of 'pan isoform' primers located in exon 1 (forward)
and exon 6 (reverse), which yielded four bands (Figure 2A). Three of the bands corresponded
to the 5' of annotated isoforms, two of which are predicted to contain the NLS (mTLK2-204 and
-201) and one that lacks the NLS due to an alternative start site in exon 6 (mTLK2-207). We
also sequenced a novel isoform that has a truncated exon 3, lacks exon 4 and 5 and would
also not encode the NLS (mTLK2-novel; Figure 2B and Figure S2). We next used *in situ*
hybridisation to observe whether there is any cell type specificity in the expression of these
TLK2 isoforms in the brain. All mouse TLK2 transcripts express exon 6 and combinations of
exons 1-5, therefore we designed a suite of short 150 bp splice junction-spanning *in situ* probes
anchored at the 3' end by the first 75 bp complementary to exon 6 and containing a variable 5'
75 bp complementary to exons 5, 4 or 3 (Figure 2B).
289

In coronal and sagittal mouse brain sections we examined the expression of TLK2 isoforms in
the hippocampal formation and cerebellar cortex respectively. A 150 bp exon junction spanning
calbindin probe was used as a positive control, confirming calbindin expression in the Purkinje
neurons of the cerebellar cortex and dentate gyrus and CA1 neurons of the hippocampus
(Figure 2C,E). Low background staining was observed in the absence of an *in situ* probe
(Figure 2C,E). In the coronal sections, hippocampal staining of all TLK2 isoform probes was
detected in excitatory neurons of the dentate gyrus, CA1 and CA3 (Figure 2C). Although
quantifying the intensity of staining between probes was not appropriate, using regions of
interest placed in the DG, CA1 and CA3 fields, we quantified the relative intensity of staining
of each TLK2 isoform in these hippocampal cell types and normalised the density to the signal
in the suprapyramidal blade of the dentate gyrus (DGs; Figure 2C). The full length TLK2 (exons
1-6) probe, corresponding to the most abundant splice variant mTLK2-204, gave strong
staining in the cell soma of the granule neurons in the DG and pyramidal neurons in CA1 and
to a lesser extent in CA3 (Fig 2B, C). The other long TLK2 isoform (mTLK2-201) was less
readily detected in the hippocampus, but quantification revealed differential expression in the
two pyramidal blades of the DG, with the suprapyramidal blade having the highest intensity
(Figure 2D). The relative staining of the shorter variant, mTLK2-207, was highest in CA1
neurons. In-line with the PCR (Figure 2A), staining of the novel short TLK2 isoform appeared
weakest, with the highest intensity in CA1 and CA3 compared to the DG (Figure 2C, D).
309
In sagittal sections of the cerebellar cortex, staining of the TLK2 probes was readily detectable
in the abundant granule neurons and to a lesser extent in Purkinje neurons (Figure 2E). The
intensity of staining for the full length mTLK2-204 probe was similar to that of the shorter
mTLK2-207 in the granule cell and Purkinje layers, whereas the mTLK2-201 and mTLK2-novel
probes had weak granule layer staining and barely detectable staining of Purkinje neurons
(Figure 2E). For all TLK2 probes, we failed to observe staining of inhibitory interneurons in the
hippocampus or cerebellar cortex. Taken together, these data confirm the RNAseq data that
alternative splice variants of TLK2 are expressed in neurons of the mammalian brain. It is
currently unclear as to the physiological relevance of the neuronal cell-type specific expression
of these TLK2 variants we have observed.
320
Alternative splicing of TLK2 is regulated during neuronal differentiation
To date, our knowledge of TLK2 is mainly limited to its nuclear functions (Segura-Bayona and
Stracker, 2019). The possibility of a non-nuclear pool of TLK2 in post-mitotic neurons has
interesting implications for the role of TLK2 in the developing and mature brain. To study TLK2
expression in neuronal differentiation we adopted a protocol for differentiating the human
neuroblastoma cell line SH-SY5Y. Based on a previously published method we used the
sequential application of retinoic acid (RA) and then BDNF with serum withdrawal over several
days (Encinas et al., 2000), which produced robust neuronal morphology (Figure 3A) and
enhanced transcript expression of the neuronal marker tau (Figure 3B). To capture the diversity
of 5' TLK2 splice variants we performed RT-PCR on SH-SY5Y cDNA samples from day 1-13
with primers located in exons 1 (forward) and 9 (reverse). This yielded six bands (Figure 3B),
which were sequenced (Figure S3). Figure 3B summarises the exon structure of each TLK2
PCR product, any corresponding annotated Ensembl transcripts and the predicted mass of the
protein they encode. The three longest PCR products mapped onto several Ensembl
transcripts with alternative splicing of combinations of exons 5, 7 and 8. The top two bands
each potentially correspond to two Ensmbl isoforms due to alternative splicing at exon 12,
which wasn't amplified in the PCR reaction. These bands were also the most prominent PCR

products in HEK293 cells. Sequencing the third band yielded a mix of sequences that we were
unable to deconvolve, but likely comprises up to three isoforms of the same mass (Figure 3B,
S3). We were not able to detect expression of a transcript corresponding to TLK2-213, the
annotated TLK2 transcript that lacks the NLS (Figure 1B). Instead, the three smallest PCR
products represented novel isoforms lacking exons 3, 4 and 5, with alternative splicing of exons
2, 7 and 8 (Figure 3B). All three of these transcripts are predicted to be protein coding and lack
the NLS. HEK293 cells possessed very low levels of these PCR products.
345
We observed a striking regulation of TLK2 isoform expression during neuronal differentiation
of the SH-SY5Y cells (Figure 3B). In undifferentiated cells the TLK2 isoforms are dominated
by the long transcripts (TLK-201/203; TLK-202/224; TLK2-209/227/230) and the longest of the
short isoforms that lacks the NLS (TLK2-novel-a). After the addition of RA, the expression of
the long TLK-202/224 isoform is rapidly downregulated (Figure 3B,C). The subsequent
addition of BDNF in serum-free media induced an upregulation of the shortest variants (TLK2-
novel-b and -c) and a concomitant downregulation of the longer isoforms. At the end of the
differentiation protocol, the TLK2 isoform profile was a broader mix of long and short variants
with similar levels of expression (Figure 3B).
355
To establish the corresponding pattern of TLK2 isoform protein expression during
differentiation, SH-SY5Y whole cell lysates from the timecourse experiment were subjected to
immunoblotting with a C-terminal reactive antibody, for which the epitope is predicted to be
present in all major splice variants of TLK2 (Figure 4A). Based on the predicted molecular
weight of the proteins encoded by the TLK2 transcripts identified in SH-SY5Y cells (see Figure
3B), we expected that the long isoforms would migrate in the range 78-87 kDa, while the
shorter isoforms lacking the NLS would migrate at 68-75 kDa. Indeed, two groups of bands
were observed at approximately 85 kDa and 70 kDa, and their expression was regulated during
SH-SY5Y differentiation, broadly correlating with the expression of the long and short TLK2-
transcripts observed in Figure 3. Quantification of replicate immunoblots by densitometry
revealed that the 85 kDa band had a biphasic expression during differentiation, which reduced
during RA treatment and then stabilised during BDNF treatment (Figure 4A,B). The ~70 kDa
band increased in expression during differentiation, with this band becoming the dominant
isoform in the fully differentiated cells (Figure 4A,B). Taken together these data suggest that
neuronal differentiation enhances expression of a TLK2 isoform that lacks the NLS and is
predicted to be cytoplasmic.
372
Cytoplasmic localisation of TLK2 in differentiated neurons
Immunofluorescent staining of SH-SY5Y cells at 1, 4, 9 and 11 days of differentiation with the
TLK2 C-terminal antiserum, revealed a time-dependent relocalisation of TLK2 from the nucleus
to the cytoplasm (Figure 4C). We quantified its subcellular localisation by measuring the
intensity of TLK2 staining in regions of interest within the cytoplasm and nucleus of the cells.
The plot of log2(cytoplasm/nucleus) in Figure 4D revealed that TLK2 is significantly enriched
in the cytoplasm at day 11 compared to day 1.
380
To confirm that TLK2 proteins encoded by splice variants lacking the NLS-containing exon 3
localise to the cytoplasm, we generated stable tet-inducible SK-N-SH T-REx-Flp-in cell lines
that express either FLAG-TLK2(180-750), corresponding to the predicted protein encoded by
Ensembl isoform TLK2-213 or full length FLAG-TLK2 corresponding to the protein product of
the canonical Ensembl TLK2 full length isoform (TLK2-203). Following a 48 h induction of TLK2

expression with doxycycline in these cell lines, we performed a crude subcellular fractionation
(Figure 5B). The cells were subjected to hypotonic lysis and we isolated the pellet (enriched in
nuclei) and post-nuclear supernatant. Immunoblotting with antibodies raised to compartment
markers, histone H3 and GAPDH, confirmed enrichment of nuclear and cytoplasmic proteins
respectively (Figure 5B). The immunostaining of anti-FLAG of the subcellular fractions and in
fixed cells demonstrated that the N-terminal truncated kinase was exclusively localised to the
cytoplasm, while the full length kinase is predominantly nuclear (Figure 5A,B). Taken together
we propose that a cytoplasmic pool of TLK2 increases during neuronal differentiation due to
two phenomena: an increase in the expression of a short TLK2 splice variant lacking the NLS
and the export of full length TLK2 from the nucleus.
396
Nuclear-cytoplasmic shuttling of TLK2
The nuclear-cytoplasmic transport of TLK2 has been previously reported by Zhang et al (1999)
who discovered a yeast 2-hybrid interaction between a 14-3-3ζ bait and a C-terminal clone of
TLK2 (amino acids 657-750). They found that 14-3-3ζ immunoprecipitated from cell lysates
with TLK2 and a dominant negative 14-3-3 promoted the nuclear localisation of TLK2 (Zhang
et al., 1999). Since 14-3-3 proteins act by binding a phosphorylated motif (Yaffe et al., 1997),
we hypothesised that during neuronal differentiation, phosphorylation of TLK2 recruits 14-3-3,
thus preventing nuclear import of TLK2 and restricting it to the cytoplasm. Using
undifferentiated SH-SY5Y cells, we attempted to mimic these signals to drive cytoplasmic
localisation of endogenous TLK2 by stimulating kinase signalling with 100 mM KCl and 10 uM
forskolin for 1 h. 100 mM KCl is known to depolarise SH-SY5Y cells, mimicking synaptic
activity, and should activate Ca^{2+}-dependent intracellular signalling in addition to the MAP
kinase pathway (Corrales et al., 2005). Forskolin, an adenylyl cyclase agonist, stimulates
cAMP production and hence activates signalling downstream of PKA (cAMP-dependent
protein kinase). Cells treated with KCl and forskolin together caused TLK2 to translocate from
the nucleus to the cytoplasm as visualised by immunofluorescence (Figure 5C) and quantified
by log2(nucleus/cytoplasm) of TLK2 intensity (Figure 5D). These data suggest the
nucleocytoplasmic shuttling of TLK2 that occurs during neuronal differentiation might also be
regulated in neurons by activity-dependent signalling.
416
417
418

Discussion

We have undertaken the first characterisation of neuronal TLK2 expression and localisation. We found that TLK2 is expressed in the hippocampus and cerebellum, brain regions that are affected in neurodevelopmental disorders linked with disrupted motor function and cognition. In addition to canonical full length TLK2, we observed splice diversity at the N-terminus of the protein, including variants that lack an NLS. Furthermore, cytoplasmic TLK2 increased during neuronal differentiation, driven by both enhanced expression of TLK2 variants lacking the NLS, and nuclear export of full length TLK2. We also found that the localisation of TLK2 can be regulated acutely by mimicking synaptic activity. Our data suggest future studies into the pathological mechanisms of MRD57 should consider cytoplasmic TLK2 signalling in developing and mature neurons.

Expression and alternative splicing of neuronal TLK2

Our analysis of TLK2 transcripts in mouse brain, SH-SY5Y cells, and publicly available RNAseq datasets revealed broad conservation of full length TLK2 and shorter transcripts lacking N-terminal exons. Interestingly the canonical, and longest, human Ensembl isoform, TLK2-201, comprising 22 exons is weakly expressed, except in aorta and testis (Figure S1), while the most abundant variant in brain and across the body is TLK-202, which lacks exons 5 and 12. Indeed the other major isoforms that are expressed (TLK-203 and TLK-213) also lack exon 12. TLK-213 is the major annotated short isoform that lacks the N-terminus and NLS and is the human homologue of the short mTLK2-207 isoform that was first identified in mouse testis (Shalom and Don, 1999). However, when we assessed the splice diversity of TLK2 transcripts in SH-SY5Y cells during neuronal differentiation we failed to observe TLK2-213, but instead discovered several novel isoforms that lack the NLS encoded by exon 3. These variants were only weakly detected in HEK293 cDNA, suggesting they might be upregulated in neurons. When we immunoblotted TLK2 in SH-SY5Y cells using an antibody with a C-terminal epitope, we observed bands that broadly correspond to long and short isoforms. However, due to the minor differences in molecular mass arising from the various splicing events it is not possible to correlate specific protein bands with specific splice variants.

Interestingly, human TLK1, which shares 78% homology with full length TLK2, also possesses a short isoform, TLK1B, lacking the N-terminus and hence the NLS. Constitutive expression of TLK1B is highest in testis, liver and lung, but low in brain (Li et al., 2001). Furthermore, TLK1B expression can be induced by genotoxic stress, such as ionising radiation, through translational regulation of an alternative start site (Li et al., 2001). Surprisingly, in contrast to short TLK2, TLK1B localises to the nucleus where it coordinates DNA repair, even in the absence of the TLK1 NLS (Li et al., 2001; Sunavala-Dossabhoy et al., 2005), suggesting TLK1B is targeted to the nucleus by an alternative mechanism.

Using splice junction-specific *in situ* probes, we were able to examine the relative distributions of long and short TLK2 isoforms in the mouse brain. TLK2 was expressed in the major cell types of the hippocampus and cerebellum, although it was notable that expression in excitatory neurons was easily observed, but inhibitory interneurons were not detectably stained. This is in contradiction to single cell RNAseq analyses of mouse hippocampus and cerebellum, which detected abundant TLK2 transcripts in various interneuron cell types (Habib et al., 2016; Kozareva et al., 2021). We observed differential expression of TLK2 isoforms in excitatory cell types of the hippocampal formation, with intriguing differences

467   between CA1 and CA3 intensity (mTLK2-204) and the blades that comprise the dentate
468   gyrus (mTLK2-201). It is not surprising that there are transcriptional differences between
469   hippocampal cell types, which presumably define their function. However, a deeper
470   understanding of the role of TLK2 in the developing and mature brain is needed to predict the
471   significance of these data.
472
Neuronal differentiation- and activity-dependent nucleocytoplasmic shuttling of TLK2
We observed a striking loss of nuclear TLK2 staining in SH-SY5Y cells during neuronal
differentiation. Since we did not observe a significant concomitant reduction in protein
expression of longer TLK2 isoforms, we concluded that TLK2 was shuttling from the nucleus
to the cytoplasm during differentiation. The upregulation of short TLK2 isoform expression we
observed is also likely to contribute to the enhancement of cytoplasmic TLK2. This was
validated by creating stable SK-N-SH cells cell lines expressing an N-terminally truncated
TLK2 variant, which was confined to the cytoplasm. These data agree with the non-nuclear
localisation of a short TLK2 variant that was overexpressed in AD293 cells (Mortuza et al.,
2018).
483
Our observation of long TLK2 export from the nucleus during differentiation is consistent with
previous data on the cytoplasmic and nuclear distribution of TLK2. Using a C-terminal TLK2
monoclonal antibody, Zhang et al (1999) demonstrated cell cycle-dependent shuttling of
TLK2 in cultures of synchronised 3T3 cells. During G1 phase, TLK2 was predominantly
outside the nucleus and co-localised with vimentin, an intermediate filament protein. With the
onset of S-phase TLK2 was observed to localise to the nuclear periphery and then within the
nucleus during late G2 (Zhang et al., 1999). Thus, the nuclear export and cytoplasmic
accumulation of TLK2 during neuronal differentiation correlates with these cells exiting the
cell cycle. Mechanistically, Zhang et al. attributed the nucleocytoplasmic shuttling of TLK2 to
an interaction with 14-3-3 proteins, with binding preventing TLK2 nuclear import. They
identified a putative 14-3-3 interaction motif in the C-terminus (amino acids 747-753). Serine
phosphorylation of 14-3-3 motifs is required for binding, but the phosphorylation of that motif
in TLK2 has not been characterised. Since we observed translocation of TLK2 in response to
KCl-induced depolarisation and FSK, it is possible that phosphorylation of TLK2 by a Ca^{2+}-
and/or cAMP-dependent kinase facilitates 14-3-3 binding and hence retains it in the
cytoplasm. There are examples in the literature of transcription factors that require neuronal
activity-dependent Ca^{2+} and cAMP signalling for nucleocytoplasmic shuttling, such as the
transcriptional activator CRTC1 (Ch'ng et al., 2012). The functions of non-nuclear TLK2 have
not been investigated, although its colocalisation with vimentin might suggest regulation of
the cytoskeleton (Zhang et al., 1999). This is in-line with the Tousled homologue in
Drosophila, which has been shown to genetically interact with *rho1* (Gregory et al., 2007)*, a
cytoskeletal regulator, and the *tlk* mutant has altered microtubule and actin filament density
in developing follicle cells (Yeh et al., 2015). Other cytoplasmic TLK2 interactors are lacking.
Previous TLK2 immunoprecipitation or proximal interactome studies have been conducted in
non-neuronal proliferating cell lines, where it might be expected that TLK2 is predominantly
nuclear, and these have tended to identify nuclear binding partners (Pavinato et al., 2022;
Segura-Bayona et al., 2017).
511
512   **The implications of neuronal TLK2 splice variants for MRD57**
513   Our study raises mechanistic questions about the pathology of the neurodevelopmental
514   disorder MRD57, in which patients are haploinsufficient for TLK2, predominantly through *de*

novo mutations (Lelieveld et al., 2016; Pavinato et al., 2022; Reijnders et al., 2018; Woods et al., 2022). The important role of TLK2 in cell cycle regulation and DNA repair has led to the hypothesis that the proliferation and differentiation of neural progenitors is affected in MRD57 patient brain development (Segura-Bayona and Stracker, 2019). MRD57 phenotypes are presumably due to processes involving TLK2 that are not compensated by TLK1, such as non-nuclear functions in post-mitotic neurons. Surprisingly, the TLK2 knockout mouse did not display any cell cycle defects, due to compensation by TLK1, except in the placenta where TLK1 expression is low (Segura-Bayona et al., 2017). However, the knockout mouse might not be a good model for MRD57 as there were no reported brain or behavioural anomalies in the TLK2 conditional heterozygous or homozygous mutant mice (Segura-Bayona et al., 2017).

It is notable that MRD57 patient missense mutations have not been identified in exons 2, 4-7 or 12. This suggests that these regions are either not necessary for TLK2 function, or their mutation is tolerated through compensation by alternative splice variants lacking these exons. Relevant to our study, one patient has been identified with a homozygous substitution mutation (K55E) adjacent to the NLS (Töpf et al., 2020), and it would be interesting to study the localisation and nuclear shuttling of this TLK2 mutant in neurons. Furthermore, future studies of neuronal TLK2 should focus on identifying its neuronal cytoplasmic substrates to advance our understanding of TLK2 in developing and mature neurons and the pathology of MRD57.

Conflict of interest statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

Credit author statement

Lubna Nuhu-Soso: Writing – review & editing, Writing – original draft, Visualization, Methodology, Investigation, Formal analysis, Conceptualization. **Heidi Denton**: Investigation, Methodology, Formal analysis. **Darren L. Goffin**: Supervision, Writing – review & editing, Methodology, Conceptualization. **Ines Hahn**: Writing – review & editing, Supervision, Funding acquisition. **Gareth J.O. Evans**: Writing – review & editing, Writing – original draft, Visualization, Supervision, Project administration, Methodology, Funding acquisition, Formal analysis, Data curation, Conceptualization.

Funding

This work was part-funded by the Wellcome Trust (ref: 204829) to GJOE and DLG through the Centre for Future Health (CFH) at the University of York, and by an Academy of Medical Science Springboard Award to IH (SBF008\1140).

Acknowledgements

We are grateful to Prof Betsy Pownall and Dr Harry Isaacs (University of York) for valuable advice regarding the *in situ* hybridisation experiments. We thank the Bioscience Technology Facility at the University of York for providing access to confocal microscopes.

References

Bhoir, S., De Benedetti, A., 2023. Targeting prostate cancer, the "tousled way." Int. J. Mol. Sci. 24, 11100. https://doi.org/10.3390/ijms241311100

Bruinsma, W., van den Berg, J., Aprelia, M., Medema, R.H., 2016. Tousled-like kinase 2 regulates recovery from a DNA damage-induced G2 arrest. EMBO Rep. 17, 659–670. https://doi.org/10.15252/embr.201540767

Ch'ng, T.H., Uzgil, B., Lin, P., Avliyakulov, N.K., O'Dell, T.J., Martin, K.C., 2012. Activity-dependent transport of the transcriptional coactivator CRTC1 from synapse to nucleus. Cell 150, 207–221. https://doi.org/10.1016/j.cell.2012.05.027

Corrales, A., Montoya G, J.V., Sutachan, J.-J., Cornillez-Ty, G., Garavito-Aguilar, Z., Xu, F., Blanck, T.J.J., Recio-Pinto, E., 2005. Transient increases in extracellular K+ produce two pharmacological distinct cytosolic Ca2+ transients. Brain Res. 1031, 174–184. https://doi.org/10.1016/j.brainres.2004.10.031

Encinas, M., Iglesias, M., Liu, Y., Wang, H., Muhaisen, A., Ceña, V., Gallego, C., Comella, J.X., 2000. Sequential treatment of SH-SY5Y cells with retinoic acid and brain-derived neurotrophic factor gives rise to fully differentiated, neurotrophic factor-dependent, human neuron-like cells. J. Neurochem. 75, 991–1003. https://doi.org/10.1046/j.1471-4159.2000.0750991.x

Fisher, M.E., Isaacs, H.V., Pownall, M.E., 2002. eFGF is required for activation of XmyoD expression in the myogenic cell lineage of Xenopus laevis. Development 129, 1307–1315. https://doi.org/10.1242/dev.129.6.1307

Gregory, S.L., Shandala, T., O'Keefe, L., Jones, L., Murray, M.J., Saint, R., 2007. A Drosophila overexpression screen for modifiers of Rho signalling in cytokinesis. Fly 1, 13–22. https://doi.org/10.4161/fly.3806

Habib, N., Li, Y., Heidenreich, M., Swiech, L., Avraham-Davidi, I., Trombetta, J.J., Hession, C., Zhang, F., Regev, A., 2016. Div-Seq: Single-nucleus RNA-Seq reveals dynamics of rare adult newborn neurons. Science 353, 925–928. https://doi.org/10.1126/science.aad7038

Harrison, P.W., Amode, M.R., Austine-Orimoloye, O., Azov, A.G., Barba, M., Barnes, I., Becker, A., Bennett, R., Berry, A., Bhai, J., Bhurji, S.K., Boddu, S., Branco Lins, P.R., Brooks, L., Ramaraju, S.B., Campbell, L.I., Martinez, M.C., Charkhchi, M., Chougule, K., Cockburn, A., Davidson, C., De Silva, N.H., Dodiya, K., Donaldson, S., El Houdaigui, B., Naboulsi, T.E., Fatima, R., Giron, C.G., Genez, T., Grigoriadis, D., Ghattaoraya, G.S., Martinez, J.G., Gurbich, T.A., Hardy, M., Hollis, Z., Hourlier, T., Hunt, T., Kay, M., Kaykala, V., Le, T., Lemos, D., Lodha, D., Marques-Coelho, D., Maslen, G., Merino, G.A., Mirabueno, L.P., Mushtaq, A., Hossain, S.N., Ogeh, D.N., Sakthivel, M.P., Parker, A., Perry, M., Piližota, I., Poppleton, D., Prosovetskaia, I., Raj, S., Pérez-Silva, J.G., Salam, A.I.A., Saraf, S., Saraiva-Agostinho, N., Sheppard, D., Sinha, S., Sipos, B., Sitnik, V., Stark, W., Steed, E., Suner, M.-M., Surapaneni, L., Sutinen, K., Tricomi, F.F., Urbina-Gómez, D., Veidenberg, A., Walsh, T.A., Ware, D., Wass, E., Willhoft, N.L., Allen, J., Alvarez-Jarreta, J., Chakiachvili, M., Flint, B., Giorgetti, S., Haggerty, L., Ilsley, G.R., Keatley, J., Loveland, J.E., Moore, B., Mudge, J.M., Naamati, G., Tate, J., Trevanion, S.J., Winterbottom, A., Frankish, A., Hunt, S.E., Cunningham, F., Dyer, S., Finn, R.D., Martin, F.J., Yates, A.D., 2024. Ensembl 2024. Nucleic Acids Res. 52, D891–D899. https://doi.org/10.1093/nar/gkad1049

Kim, J.-A., Anurag, M., Veeraraghavan, J., Schiff, R., Li, K., Wang, X.-S., 2016a. Amplification of TLK2 Induces Genomic Instability via Impairing the G2-M Checkpoint. Mol. Cancer Res. 14, 920–927. https://doi.org/10.1158/1541-7786.MCR-16-0161

Kim, J.-A., Tan, Y., Wang, X., Cao, X., Veeraraghavan, J., Liang, Y., Edwards, D.P., Huang, S., Pan, X., Li, K., Schiff, R., Wang, X.-S., 2016b. Comprehensive functional analysis of the tousled-like kinase 2 frequently amplified in aggressive luminal breast cancers. Nat. Commun. 7, 12991. https://doi.org/10.1038/ncomms12991

Klimovskaia, I.M., Young, C., Strømme, C.B., Menard, P., Jasencakova, Z., Mejlvang, J., Ask, K., Ploug, M., Nielsen, M.L., Jensen, O.N., Groth, A., 2014. Tousled-like kinases

phosphorylate Asf1 to promote histone supply during DNA replication. Nat. Commun. 5,
3394. https://doi.org/10.1038/ncomms4394
Kozareva, V., Martin, C., Osorno, T., Rudolph, S., Guo, C., Vanderburg, C., Nadaf, N.,
Regev, A., Regehr, W.G., Macosko, E., 2021. A transcriptomic atlas of mouse cerebellar
cortex comprehensively defines cell types. Nature 598, 214–219.
https://doi.org/10.1038/s41586-021-03220-z
Lee, S.-B., Segura-Bayona, S., Villamor-Payà, M., Saredi, G., Todd, M.A.M., Attolini, C.S.-
O., Chang, T.-Y., Stracker, T.H., Groth, A., 2018. Tousled-like kinases stabilize
replication forks and show synthetic lethality with checkpoint and PARP inhibitors. Sci
Adv 4, eaat4985. https://doi.org/10.1126/sciadv.aat4985
Lelieveld, S.H., Reijnders, M.R.F., Pfundt, R., Yntema, H.G., Kamsteeg, E.-J., de Vries, P.,
de Vries, B.B.A., Willemsen, M.H., Kleefstra, T., Löhner, K., Vreeburg, M., Stevens,
S.J.C., van der Burgt, I., Bongers, E.M.H.F., Stegmann, A.P.A., Rump, P., Rinne, T.,
Nelen, M.R., Veltman, J.A., Vissers, L.E.L.M., Brunner, H.G., Gilissen, C., 2016. Meta-
analysis of 2,104 trios provides support for 10 new genes for intellectual disability. Nat.
Neurosci. 19, 1194–1196. https://doi.org/10.1038/nn.4352
Li, Y., DeFatta, R., Anthony, C., Sunavala, G., De Benedetti, A., 2001. A translationally
regulated Tousled kinase phosphorylates histone H3 and confers radioresistance when
overexpressed. Oncogene 20, 726–738. https://doi.org/10.1038/sj.onc.1204147
López-Rivera, J.A., Pérez-Palma, E., Symonds, J., Lindy, A.S., McKnight, D.A., Leu, C.,
Zuberi, S., Brunklaus, A., Møller, R.S., Lal, D., 2020. A catalogue of new incidence
estimates of monogenic neurodevelopmental disorders caused by de novo variants.
Brain 143, 1099–1105. https://doi.org/10.1093/brain/awaa051
Mortuza, G.B., Hermida, D., Pedersen, A.-K., Segura-Bayona, S., López-Méndez, B.,
Redondo, P., Rüther, P., Pozdnyakova, I., Garrote, A.M., Muñoz, I.G., Villamor-Payà,
M., Jauset, C., Olsen, J.V., Stracker, T.H., Montoya, G., 2018. Molecular basis of
Tousled-Like Kinase 2 activation. Nat. Commun. 9, 2535.
https://doi.org/10.1038/s41467-018-04941-y
Pavinato, L., Villamor-Payà, M., Sanchiz-Calvo, M., Andreoli, C., Gay, M., Vilaseca, M.,
Arauz-Garofalo, G., Ciolfi, A., Bruselles, A., Pippucci, T., Prota, V., Carli, D., Giorgio, E.,
Radio, F.C., Antona, V., Giuffrè, M., Ranguin, K., Colson, C., De Rubeis, S., Dimartino,
P., Buxbaum, J.D., Ferrero, G.B., Tartaglia, M., Martinelli, S., Stracker, T.H., Brusco, A.,
2022. Functional analysis of TLK2 variants and their proximal interactomes implicates
impaired kinase activity and chromatin maintenance defects in their pathogenesis. J.
Med. Genet. 59, 170–179. https://doi.org/10.1136/jmedgenet-2020-107281
Reijnders, M.R.F., Miller, K.A., Alvi, M., Goos, J.A.C., Lees, M.M., de Burca, A., Henderson,
A., Kraus, A., Mikat, B., de Vries, B.B.A., Isidor, B., Kerr, B., Marcelis, C., Schluth-
Bolard, C., Deshpande, C., Ruivenkamp, C.A.L., Wieczorek, D., Deciphering
Developmental Disorders Study, Baralle, D., Blair, E.M., Engels, H., Lüdecke, H.-J.,
Eason, J., Santen, G.W.E., Clayton-Smith, J., Chandler, K., Tatton-Brown, K., Payne, K.,
Helbig, K., Radtke, K., Nugent, K.M., Cremer, K., Strom, T.M., Bird, L.M., Sinnema, M.,
Bitner-Glindzicz, M., van Dooren, M.F., Alders, M., Koopmans, M., Brick, L., Kozenko,
M., Harline, M.L., Klaassens, M., Steinraths, M., Cooper, N.S., Edery, P., Yap, P.,
Terhal, P.A., van der Spek, P.J., Lakeman, P., Taylor, R.L., Littlejohn, R.O., Pfundt, R.,
Mercimek-Andrews, S., Stegmann, A.P.A., Kant, S.G., McLean, S., Joss, S.,
Swagemakers, S.M.A., Douzgou, S., Wall, S.A., Küry, S., Calpena, E., Koelling, N.,
McGowan, S.J., Twigg, S.R.F., Mathijssen, I.M.J., Nellaker, C., Brunner, H.G., Wilkie,
A.O.M., 2018. De Novo and Inherited Loss-of-Function Variants in TLK2: Clinical and
Genotype-Phenotype Evaluation of a Distinct Neurodevelopmental Disorder. Am. J.
Hum. Genet. 102, 1195–1203. https://doi.org/10.1016/j.ajhg.2018.04.014
Roe, J.L., Rivin, C.J., Sessions, R.A., Feldmann, K.A., Zambryski, P.C., 1993. The Tousled
gene in A. thaliana encodes a protein kinase homolog that is required for leaf and flower
development. Cell 75, 939–950. https://doi.org/10.1016/0092-8674(93)90537-z
RStudio Team, 2020. RStudio Integrated Development for R. http://www.rstudio.com/.
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J.,
Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source
platform for biological-image analysis. Nat. Methods 9, 676–682.
https://doi.org/10.1038/nmeth.2019
Segura-Bayona, S., Knobel, P.A., González-Burón, H., Youssef, S.A., Peña-Blanco, A.,
Coyaud, É., López-Rovira, T., Rein, K., Palenzuela, L., Colombelli, J., Forrow, S.,
Raught, B., Groth, A., de Bruin, A., Stracker, T.H., 2017. Differential requirements for
Tousled-like kinases 1 and 2 in mammalian development. Cell Death Differ. 24, 1872–
1885. https://doi.org/10.1038/cdd.2017.108
Segura-Bayona, S., Stracker, T.H., 2019. The Tousled-like kinases regulate genome and
epigenome stability: implications in development and disease. Cell. Mol. Life Sci. 76,
3827–3841. https://doi.org/10.1007/s00018-019-03208-z
Shalom, S., Don, J., 1999. Tlk, a novel evolutionarily conserved murine serine threonine
kinase, encodes multiple testis transcripts. Mol. Reprod. Dev. 52, 392–405.
https://doi.org/10.1002/(SICI)1098-2795(199904)52:4<392::AID-MRD8>3.0.CO;2-Y
Silljé, H.H., Takahashi, K., Tanaka, K., Van Houwe, G., Nigg, E.A., 1999. Mammalian
homologues of the plant Tousled gene code for cell-cycle-regulated kinases with
maximal activities linked to ongoing DNA replication. EMBO J. 18, 5691–5702.
https://doi.org/10.1093/emboj/18.20.5691
Sjöstedt, E., Zhong, W., Fagerberg, L., Karlsson, M., Mitsios, N., Adori, C., Oksvold, P.,
Edfors, F., Limiszewska, A., Hikmet, F., Huang, J., Du, Y., Lin, L., Dong, Z., Yang, L.,
Liu, X., Jiang, H., Xu, X., Wang, J., Yang, H., Bolund, L., Mardinoglu, A., Zhang, C., von
Feilitzen, K., Lindskog, C., Pontén, F., Luo, Y., Hökfelt, T., Uhlén, M., Mulder, J., 2020.
An atlas of the protein-coding genes in the human, pig, and mouse brain. Science 367,
eaay5947. https://doi.org/10.1126/science.aay5947
Sunavala-Dossabhoy, G., Balakrishnan, S.K., Sen, S., Nuthalapaty, S., De Benedetti, A.,
2005. The radioresistance kinase TLK1B protects the cells by promoting repair of double
strand breaks. BMC Mol. Biol. 6, 19. https://doi.org/10.1186/1471-2199-6-19
Töpf, A., Oktay, Y., Balaraju, S., Yilmaz, E., Sonmezler, E., Yis, U., Laurie, S., Thompson,
R., Roos, A., MacArthur, D.G., Yaramis, A., Güngör, S., Lochmüller, H., Hiz, S., Horvath,
R., 2020. Severe neurodevelopmental disease caused by a homozygous TLK2 variant.
Eur. J. Hum. Genet. 28, 383–387. https://doi.org/10.1038/s41431-019-0519-x
Villamor-Payà, M., Sanchiz-Calvo, M., Smak, J., Pais, L., Sud, M., Shankavaram, U.,
Lovgren, A.K., Austin-Tse, C., Ganesh, V.S., Gay, M., Vilaseca, M., Arauz-Garofalo, G.,
Palenzuela, L., VanNoy, G., O'Donnell-Luria, A., Stracker, T.H., 2024. De novo TLK1
and MDM1 mutations in a patient with a neurodevelopmental disorder and
immunodeficiency. iScience 27, 109984. https://doi.org/10.1016/j.isci.2024.109984
Woods, E., Spiller, M., Balasubramanian, M., 2022. Report of two children with global
developmental delay in association with de novo TLK2 variant and literature review. Am.
J. Med. Genet. A 188, 931–940. https://doi.org/10.1002/ajmg.a.62580
Yaffe, M.B., Rittinger, K., Volinia, S., Caron, P.R., Aitken, A., Leffers, H., Gamblin, S.J.,
Smerdon, S.J., Cantley, L.C., 1997. The structural basis for 14-3-3:phosphopeptide
binding specificity. Cell 91, 961–971. https://doi.org/10.1016/s0092-8674(00)80487-0
Yamakawa, A., Kameoka, Y., Hashimoto, K., Yoshitake, Y., Nishikawa, K., Tanihara, K.,
Date, T., 1997. cDNA cloning and chromosomal mapping of genes encoding novel
protein kinases termed PKU-alpha and PKU-beta, which have nuclear localization
signal. Gene 202, 193–201. https://doi.org/10.1016/s0378-1119(97)00495-2
Yeh, T.-H., Huang, S.-Y., Lan, W.-Y., Liaw, G.-J., Yu, J.-Y., 2015. Modulation of cell
morphogenesis by tousled-like kinase in the Drosophila follicle cell. Dev. Dyn. 244, 852–
865. https://doi.org/10.1002/dvdy.24292
Zhang, S., Xing, H., Muslin, A.J., 1999. Nuclear localization of protein kinase U-alpha is
regulated by 14-3-3. J. Biol. Chem. 274, 24865–24872.
https://doi.org/10.1074/jbc.274.35.24865

Figure legends

Figure 1. TLK2 has splice variant diversity in the human brain. (A) Exonic structure of human TLK2, mapped onto a schematic representation of its protein domains. NLS, nuclear localisation sequence; CC, coiled-coil domain. (B) Table of % TLK2 transcript abundance in the human hippocampus and cerebellum derived from GTEx RNAseq data. Exons and transcripts labelled in red are predicted to be non-protein coding. Only transcripts present >2% in either brain region are included.

Figure 2. Multiple isoforms of TLK2 are expressed in mouse forebrain and cerebellum. (A) RT-PCR of TLK2 splice variants from mouse forebrain and cerebellum cDNA. Transcript specific primer pairs or pan exon 1-6 primers. (B) Left, lane from PCR of exon 1-6 primer pair. The four bands were excised and sequenced. Right, exon structure of sequenced mouse brain splice variants, and location of *in situ* hybridisation probes. *In situ* hybridisation was performed on 25 μm frozen sections from coronal (C) or sagittal (E) P56 mouse brain sections. The indicated 150 bp exon junction spanning probes were used to examine calbindin or TLK2 transcript expression in the hippocampus (C) or cerebellar cortex (E). Representative images are shown from n=3 sections from two mouse brains. Cerebellar cortex images are shown at x10 and x40 magnification, indicated by the insert box. DG, dentate gyrus; ML, molecular layer; GL, granule layer; PCL, Purkinje cell layer. Scale bars: hippocampus 100 μm, cerebellar cortex x10, 200 μm; x40, 20 μm. (D) Heatmap depicting quantification of TLK2 splice variant expression within the dentate gyrus, CA1 and CA3 sub-regions of the hippocampal formation. For each section, pixel intensity was measured in regions of interest (ROI), as indicated in the cartoon and normalised to the ROI located in the suprapyramidal blade of the dentate gyrus. Three sections were quantified for each probe.

Figure 3. Alternative splicing of TLK2 is regulated during neuronal differentiation. SH-SY5Y cultures were differentiated by sequential treatment with retinoic acid (RA) and brain derived neurotrophic factor (BDNF) as described in Materials and Methods. (A) Brightfield images of SH-SY5Y cultures at the indicated days of differentiation. Scale bar = 40 μm. (B) cDNA was prepared from SH-SY5Y cells at the indicated time points, and HEK293 cells (control) and subjected to PCR with primers for exons 1 and 9 of human TLK2, Tau, and GAPDH. The six TLK2 bands obtained from the cells were excised and subjected to Sanger sequencing. The right hand schematic summarises the exon structure of each sequence and any corresponding Ensembl transcripts that these align to and the predicted molecular mass of the encoded protein. The bottom plot presents quantitative densitometry of each TLK2 RT-PCR product, expressed as a percentage of the total intensity at each timepoint.

Figure 4. Neuronal differentiation alters the nuclear-cytoplasmic distribution of TLK2. SH-SY5Y cells were lysed or fixed at the indicated timepoints during neuronal differentiation and subjected to immunoblotting with anti-TLK2, anti-β3-tubulin or anti-actin (A) or immunocytochemistry with anti-TLK2 (1:1000; green) and DAPI staining (purple) Scale bar = 10 μm. (C). (B) Quantification of TLK2 immunoblot band intensity from (A), data plotted as stacked bars and raw data points for long TLK2 (yellow circles) and short TLK2 (green circles) isoforms. Data were normalised to the intensity of actin and then day 4 short TLK2 (n=3 independent experiments). (D) TLK2 intensity was measured in regions of interest in the nucleus and cytoplasm of 100 cells from each timepoint (n=3 independent experiments). Data are plotted as $\log_2$(nuclear/cytoplasmic) TLK2 intensity at each timepoint (grey filled

771   circles). Hollow black circles represent the mean of each replicate. For (B) and (D), data
772   were analysed by Kruskal-Wallis two-tailed ANOVA and Dunn's post-hoc test; * p < 0.05; ** p
773   < 0.01 compared to day 1. For significance in (B), * = long TLK2 and # = short TLK2.
774
Figure 5. Activity-dependent redistribution of TLK2. Inducible Flp-in SK-N-SH cells
expressing FLAG-tagged long TLK2-203 or short TLK2-213 were treated with or without 1
µg/ml doxycycline (dox) for 48 h. (A) Representative immunofluorescence images of dox-
treated cells stained with anti-FLAG (1:1000) and DAPI. Scale bar = 10 µm. (B) Cells plated
in dishes were lysed and subject to subcellular fractionation to yield a nuclear pellet and
postnuclear supernatant. The samples were subjected to immunoblotting with anti-FLAG,
anti-GAPDH and anti-histone H3. (C) SH-SY5Y cells equilibrated in KREBS buffer were
incubated for 1 h in KREBS buffer with or without (control) 100 mM KCl and 10 µM forskolin
(FSK). Fixed cells were stained with anti-TLK2 and DAPI. Scale bar = 10 µm. (D) TLK2
intensity was measured in regions of interest in the nucleus and cytoplasm of 100 cells from
each treatment (n=3 experiments). Data are plotted as $\log_2$(nuclear/cytoplasmic) TLK2
intensity at each timepoint (grey filled circles). Hollow black circles represent the mean of
each replicate and were analysed by t- test; * p < 0.05 compared to control.
788

Figure 1

A

B

TLK2 exon structure	Ensembl transcript	human RNAseq (GTEx) % HC	CBM
	TLK2-201	2	2.1
	TLK2-202	37.2	39
	TLK2-203	2.5	5.6
	TLK2-207	29	28
	TLK2-213	22.5	12.8
	TLK2-214	1.4	6.1

red exon/transcript = non-coding

Figure 2

Figure 3

Figure 4

Figure 5