Persistent dopamine-dependent remodeling of the neural transcriptome in response to pregnancy and postpartum

Jennifer C Chan[1], Giuseppina Di Salvo[1,2], Ashley M Cunningham[1], Sohini Dutta[1], Elizabeth A Brindley[1], Ethan Wan[4], Cindy Zhang[1], Ian Maze[1,3,5*]

Affiliations

[1]Nash Family Department of Neuroscience, Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA

[2]Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience (MHeNs), Maastricht University, Maastricht, The Netherlands

[3]Department of Pharmacological Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA

[4]Department of Neurobiology, University of Alabama at Birmingham, Birmingham, AL 35294, USA.

[5]Howard Hughes Medical Institute, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA

*Corresponding author
Ian Maze
Email: ian.maze@mssm.edu

SUMMARY

Pregnancy and postpartum experiences represent transformative physiological states that impose lasting demands on the maternal body and brain, resulting in lifelong neural adaptations. However, the precise molecular mechanisms driving these persistent alterations remain poorly understood. Here, we used brain-wide transcriptomic profiling to define the molecular landscape of parity-induced neural plasticity, identifying the dorsal hippocampus (dHpc) as a key site of transcriptional remodeling. Combining single-cell RNA sequencing with a maternal-pup separation paradigm, we additionally demonstrated that chronic postpartum stress significantly disrupts dHpc adaptations by altering dopamine dynamics, leading to dysregulated transcription, altered cellular plasticity, and impaired behavior. We further established the sufficiency of dopamine modulation in the regulation of these parity-induced adaptations via chemogenetic suppression of dopamine release into dHpc, which recapitulated key transcriptional and behavioral features of parity in virgin females. In sum, our findings establish dopamine as a central regulator of parity-induced neuroadaptations, revealing a fundamental transcriptional mechanism by which female reproductive experiences remodel the maternal brain to sustain long-term behavioral adaptations.

INTRODUCTION

Matrescence – the physical, emotional, hormonal, and social transition to motherhood – is a period of profound transformation that reshapes the maternal body and brain to support pregnancy, birth, and offspring care. While extensive research has focused on acute maternal brain processes that are essential for the onset and maintenance of parenting, how precisely brain adaptations are sustained beyond the postpartum period to persistently influence behavior remains unclear. Recent human neuroimaging studies have revealed that pregnancy experience induces long-lasting alterations in brain connectivity and structure, persisting for years or even decades following birth[1–6]. Animal models examining parity – the state of previously carrying a pregnancy to term – similarly have demonstrated persistent alterations in synaptic remodeling, cell proliferation, and behavioral outcomes[7–14]. However, the specific mechanisms within the extensive repertoire of maternal physiological changes that drive these long-lasting brain adaptations remain poorly understood. With over 130 million pregnancies occurring globally each year, elucidating how female reproductive experiences shape long-term brain plasticity remains a fundamental question with widespread clinical implications. Given that parity may modulate risk and/or resilience to certain neurological and neuropsychiatric disorders[15–17], addressing this gap in knowledge will be essential for advancing our understanding of maternal brain health.

Here, we began by employing brain-wide transcriptional profiling to systematically characterize the persistent impact of parity on the maternal brain in mice, which provided important insights into the regulation of signaling pathways and cellular compositions across maternal circuits. These analyses surprisingly revealed the dorsal hippocampus (dHpc) as a key site of sustained plasticity. We then leveraged single-cell RNA sequencing, a postpartum stress paradigm, behavioral assays, and chemogenetic approaches to identify and validate a dopamine-dependent neuromodulatory

process that orchestrates these sustained maternal adaptations. Importantly, these brain-wide

datasets will also serve as a critical resource for future investigations into the reproductive state-

dependent adaptations that contribute to matrescence.

RESULTS

Parity promotes long-lasting transcriptional and behavioral adaptations in maternal brain

To first investigate the persistent effects of female reproductive experiences in maternal brain, we

established a timeline to compare primiparous dams (PP), which experienced breeding, pregnancy,

parturition, lactation, and pup interactions, to age-matched virgins that lacked reproductive

experiences (nulliparous, NP; **Fig. 1A**). Given that reproductive hormones return to baseline levels

~7-weeks post-conception[18], we then conducted brain-wide transcriptomic profiling 4-weeks

following cessation of pup rearing (10-weeks post-conception; i.e. 49 days postpartum; dpp), as

this time point would be predicted to reflect sustained changes. Following differential expression

analyses of the 11 brain regions selected based on prior evidence supporting their involvement in

maternal behaviors[19,20] (**Fig. S1A**), our data revealed a wide-range in the number of differentially

expressed genes (DEGs, adj. $p < 0.05$) across brain regions (**Fig. 1B, Table S1**). These differences

indicated varying levels of transcriptomic sensitivity to parity brain-wide, with few changes

observed in the medial prefrontal cortex (mPFC), for example, as compared to the dHpc, the region

that was found to be most robustly regulated in our analyses.

87
To next assess whether these differential expression patterns could be attributed to changes in

shared cell-types, we performed cell-type deconvolution using BRETIGEA, which compares

whole transcriptome data to validated brain cell type-specific marker gene sets[21]. We observed

changes in cell marker expression for neurons, astrocytes, microglia, oligodendrocyte precursor

cells (OPCs), oligodendrocytes, and endothelial cells across brain regions, regardless of the

number of DEGs (except for mPFC; **Fig. 1C**). These data suggested that cellular changes may

occur across brain regions independently of the extent of transcriptomic alterations, potentially

reflecting shifts in the proportion of neurons to non-neuronal/glial cells, modifications in synaptic

plasticity, or changes in cell identity in response to prior reproductive experiences. Notably, the

expression of neuronal markers was found to be downregulated in brain regions displaying the

most DEGs (**Fig. 1C**). Furthermore, a significant inverse correlation between reduced neuronal

marker expression and the number of DEGs across all brain regions implicated changes in neuronal

populations that may underlie the observed transcriptional responses (**Fig. 1D**); note that this

relationship was not observed for other cell-types (**Fig. S1B-F**). These findings nicely align with

human neuroimaging studies, which have reported reduced gray matter in mothers even years after

birth, a phenomenon that has been proposed to refine maternal neural circuits to optimize

caregiving behavior[1,2,4,22].

105
Based on these findings, we hypothesized that such shared neuronal marker alterations may

indicate common upstream mechanisms driving regional-specific transcriptional responses. To test

this, we first compared DEGs across brain regions to identify common gene sets that were altered

by parity status, which led to the observation that the greatest extent of overlap occurs in those

regions displaying the highest levels of transcriptional sensitivity (**Fig. 1E**). Based on these

analyses, we then grouped the 11 brain regions according to the number of DEGs, the degree of

overlap with other regions, and significant fold changes in neuronal marker expression, based on

the premise that these regions may be co-regulated to elicit convergent transcriptional responses

to parity (referred to as "high sensitivity" regions: dHpc, nucleus accumbens (NAc), medial

preoptic area (mPOA), locus coeruleus (LC), ventral hippocampus (vHpc)). Brain regions lacking

these criteria were classified as "low sensitivity" regions (dorsal raphe nucleus (DRN),

paraventricular nucleus of the hypothalamus (PVN), basolateral amygdala (BLA), ventral

tegmental area (VTA), olfactory bulb (OB), mPFC). Next, we explored the predicted upstream

regulators of overlapping DEGs in "High" versus "Low" sensitivity brain regions, which revealed

several upstream regulators that were grouped into common molecular classes, including

hormones, neurotransmitter/neurotrophin signaling molecules, transcriptional regulators, and

others (**Fig. 1F**). While both High and Low sensitivity regions were found to share upstream

regulators within these molecular classes, High sensitivity regions displayed greater enrichment

for estrogen, progesterone, testosterone, dopamine, and other ligands with similar receptor affinity

or structural homology. In contrast, regulators that were classified within the lipid category were

uniquely associated with Low sensitivity regions.

127
To further explore the regulatory networks that distinguish High versus Low sensitivity regions,

we performed weighted gene co-expression network analysis (WGCNA) across all 11 brain

regions[23], resulting in nine co-expression modules (**Fig. S1G-J**). Module-trait correlations, in

which the trait refers to a combination of group (NP *vs*. PP) and regional sensitivity (High *vs*.

Low), identified significant correlations between High sensitivity regions in PP females and the

brown, green, yellow, and magenta modules (**Fig. 1G;** henceforth labeled 'parity-sensitive

modules'). To identify the specific brain regions exhibiting changes in module gene expression,

we then examined enrichment of DEGs within each module, with an extended focus on the parity-

sensitive modules. Consistent with our module-trait correlational analysis, parity-sensitive

modules displayed significant enrichment for DEGs from dHpc (**Fig. 1H;** Fisher's exact tests:

brown, $p < 10^{-16}$; green, yellow, $p < 10^{-6}$), NAc (brown, green, $p < 10^{-3}$; magenta, $p < 0.01$; yellow,

$p < 0.05$), mPOA (green, $p < 0.05$; magenta, $p < 0.01$), LC (magenta, $p < 0.05$), and vHpc (magenta,

p < 0.05; yellow, p < 0.01). Functional annotation analysis of the genes from these modules

highlighted their significant enrichment in signaling pathways related to neuromodulators, such as

estrogen (yellow, brown) and dopamine (green, magenta), in agreement with our upstream

regulator analyses (**Fig. 1I, Fig. S1K-L**). In contrast, Low sensitivity regions were found to be

significantly associated with the black module, which was enriched for genes associated with RNA

and macromolecule processing pathways (**Fig. 1G, 1I**).

146
Building on these transcriptomic findings, we next investigated whether parity leads to sustained

functional alterations in PP females. To evaluate the extent of such persistence, we performed

behavioral assessments at 140 dpp, ~16-weeks after pup weaning (**Fig. 1A**). Behavioral tasks were

selected to specifically assess functions associated with the brain regions displaying robust

transcriptional changes, including maternal behavior, learning, and memory. To assess maternal

behavior, we tested pup retrieval in the home cage by placing two donor pups in opposite corners

away from the nest. PP dams were observed to retrieve pups with significantly reduced latency as

compared to NP females (**Fig. 1J**), which is consistent with previous findings, though at a more

extended timepoint in our studies[24]. Next, to investigate the functional impact of dHpc

transcriptional changes, we utilized a contextual fear conditioning paradigm. Female mice were

habituated to the training context before receiving five shocks (2-seconds each, 0.7 mA) with 90-

second intertrial intervals. Both NP and PP females demonstrated significant training acquisition,

as indicated by increased freezing compared to no-shock controls (**Fig. 1K**). Notably, PP females

exhibited enhanced acquisition, with significantly increased freezing observed by the third shock,

while NP females only displayed significant freezing at the fifth shock (**Fig. 1K**). During the

context test conducted 24-hours later, both groups demonstrated recall, with PP females showing

higher freezing responses than NP animals (**Fig. 2L**). Given the involvement of these brain regions

in anxiety- and depressive-like behaviors, we also assessed affective phenotypes using the open

field, light-dark box, and forced swim tests, however, no significant differences were observed

(**Fig. S2A-H**). These results suggested that the transcriptional changes observed in parity-sensitive

brain regions may promote maternal and dHpc-related behavioral outcomes, as has been reported

previously[7,10,12,14,25–31], though the mechanisms underlying such prolonged functional adaptations

remained unclear. To account for the potential influence of estrous cycling on behavioral changes,

vaginal swabs were collected immediately following testing, and behavioral analyses were

stratified by estrous stage. No significant effects of estrous stage were observed (**Fig. S2I-N**),

thereby supporting parity status as the primary driver of the observed behavioral differences.

173
Pregnancy and postpartum experiences promote persistent maternal neuroplasticity

Since our paradigm comparing NP *vs.* PP females could not distinguish between the discrete

reproductive event(s) that may be responsible for the sustained transcriptional alterations observed,

we next sought to isolate those key events throughout the female reproductive window. To do so,

we compared females that were successfully bred to males but did not become pregnant (Mating

Only), females that experienced pregnancy and parturition but did not transition to postpartum due

to pup removal on the day of birth (Pregnancy Only), and virgin females exposed to 21-days of

donor pup interactions (Pup Sensitized). These groups were analyzed in parallel with age-matched

NP and PP females to delineate the contributions of specific reproductive and maternal experiences

(**Fig. 2A**). Consistent with previous studies[32], four days of pup exposures sensitized virgin females

to maternal behavior, as evidenced by increased crouching over pups and decreased pup avoidance

by day 4 (**Fig. S3A**). To determine the processes conferring sustained transcriptional sensitivity,

we focused our investigations on the dHpc due to its pronounced gene expression response to

parity. When comparing NP *vs.* PP gene expression profiles, we observed that the Pregnancy Only

group most closely resembled that of the expression profile of PP females, suggesting that

pregnancy is a primary driver of dHpc neuroplastic changes (**Fig. 2B, Fig. S3B**). However, the

magnitude of fold changes did not match that of the PP group, indicating that additional

experiences, likely during postpartum, are also necessary to fully establish parity-induced changes

in the dHpc.

While visualization of differential expression profiles in the Mating Only and Pup Sensitized

groups did not mimic PP changes, comparisons to NP did identify significant DEGs across all

groups, with the most substantial changes observed in the Pup Sensitized group, followed by the

Pregnancy Only, and Mating Only groups (**Fig. S3C-E, Table S2**). Overlapping DEGs induced

by discrete reproductive events significantly intersected with DEGs in PP animals, again

suggesting that while pregnancy is the primary driver of parity-induced effects, each reproductive

event contributes, in part, to this process (**Fig. S3F**). Next, we overlapped DEGs from each

comparison with genes from parity-sensitive modules (from **Fig. 1**) to assess the contribution of

each reproductive event to parity programming. Significant overlap was observed across all

groups, with Pregnancy Only and Pup Sensitized groups displaying the strongest enrichment,

indicating that these experiences play a key role in shaping persistent dHpc alterations (**Fig. 2C**).

Thus, despite the lack of similarity between the overall expression profiles between Pup Sensitized

vs. PP, gene expression changes induced by pup interactions likely converge on similar regulatory

networks, albeit through different DEGs, which are critical for parity-driven transcriptional

alterations. To further explore these relationships, we performed threshold-free Rank-Rank

Hypergeometric Overlap (RRHO) analyses[33], which demonstrated transcriptional concordance

across experiences, with the strongest overlap observed in pregnancy (**Fig. 2D**). These findings

suggest that reproductive exposures converge, or act additively, to drive the full extent of brain

transcriptional changes. Furthermore, comparing GO term analyses of DEGs for each group

revealed significant enrichment in overlapping biological processes, with many related to synaptic

signaling and plasticity pathways (**Fig. 2E**). Notably, certain pathways were unique to the

combination of reproductive exposures, including changes in CREB activity, catecholamine

secretion, and myelination, indicating that these regulatory processes depend on multiple

reproductive events to achieve full induction (**Fig. 2E**). Although this study did not address the

specific contributions of parturition or lactation, our findings demonstrate that pregnancy

establishes the foundation for parity-induced transcriptional alterations in dHpc, with additional

postpartum exposures also required to fully promote these responses.

221
Given that postpartum experiences enhanced changes that were initiated during pregnancy, we

conducted a controlled time course study to examine these transcriptional dynamics across

gestational and postpartum periods in age-matched females (**Fig. 2F**). Pregnancy timepoints were

selected to represent the anabolic and catabolic phases of gestation, corresponding to early (7.5

days post-conception, dpc) and late (17.5 dpc) stages, respectively. Postpartum timepoints were

chosen to capture the immediate effects of hormonal withdrawal following parturition (2 dpp), as

well as a later stage prior to pup weaning (21 dpp). Based on the hypothesis that parity-induced

transcriptional alterations are progressively enhanced across reproductive stages, we first

compared DEGs at each timepoint (*vs.* NP) to genes from parity-sensitive modules identified

previously. DEGs from all comparisons significantly overlapped with the brown module (**Fig. 2G,**

left; Table S3), suggesting that pathways enriched in this module are central to dHpc

programming. Enrichment in the green and yellow modules was predominantly found to be

induced by postpartum exposures, potentially amplifying the effects of the pathways driven by the

brown module (**Fig. 2G, left**). To assess the regional specificity of these dynamic changes, we

analyzed bulk RNA-seq data from the vHpc, a region that displayed moderate sensitivity to parity

status (**Fig. 2G, middle; Table S4**), along with the mPFC, which exhibited few transcriptional

alterations (**Fig. 2G, right; Table S5**). These regions exhibited limited overlap of DEGs with

parity-sensitive module genes, further emphasizing the specificity of the pathways that drive long-

term adaptations in brain regions displaying heightened sensitivity to parity.

241
To then characterize cumulative transcriptional changes that occur across reproductive stages, we

performed a time course analysis using the ImpulseDE2 package to detect genes displaying

sustained or transient expression alterations[34]. This analysis revealed a prominent set of genes that

displayed persistent and progressive downregulation, with more pronounced alterations during

postpartum (**Fig. 2H**). These genes were significantly enriched for processes related to

glutamatergic synapses, dopaminergic signaling, and endocrine-related pathways (**Fig. 2I**).

Conversely, genes with persistent upregulation were found to be enriched for pathways associated

with cell junction dynamics (**Fig. 2J**), which is consistent with overall prolonged changes in

synaptic plasticity and transmission. Additionally, we identified gene sets displaying transient

alterations, including downregulation of neurogenesis-related pathways, and upregulation of

copper ion metabolism (**Fig. 2K-L**); these data are consistent with other studies reviewed here[35,36].

In total, these findings highlight postpartum as a key window for reinforcing parity-induced

alterations.

255
Chronic maternal stress during postpartum disrupts dHpc adaptations

We next examined whether postpartum perturbations may disrupt pathways that reinforce parity-

induced alterations in dHpc. As becoming a new parent represents a significant source of stress,

we implemented a maternal stress paradigm that robustly increases stress hormone levels and

disrupts key postpartum experiences, including nursing and pup interactions, from 10–20 dpp (**Fig.**

3A, Fig. S3G). During this period, dams were separated from their pups for 3-hours daily and were

provided with limited nesting materials until pup weaning (Stress PP). As litter size represents a

potential confounding factor, the time period selected reduces the impact of maternal stress on pup

mortality that can occur immediately following birth. By 10 dpp, pups were more durable, as

reflected by the absence of litter size differences between Control and Stress dams (**Fig. S3H**).

Accordingly, no differences in postpartum weights were observed, suggesting that this postpartum

stress paradigm does not induce major metabolic alterations in the dams (**Fig. S3I**). To assess the

extent of parity adaptations in Stress PP, we compared this group to NP and Control PP animals.

Principal components analysis (PCA) of dHpc transcriptional profiles revealed distinct clustering

of NP and Control PP groups (reproducing our prior analyses), while Stress PP samples were found

to cluster intermediately between the two groups (**Fig. 3B**). Similarly, hierarchical clustering of all

DEGs resulted in intermingling of NP and Stress PP samples (**Fig. 3C, Table S6**), suggesting that

postpartum stress disrupts the extent of parity alterations in dHpc. To determine whether genes

induced by parity are reversed by stress, we then compared DEGs with opposing directionality in

NP *vs.* Control PP and Control PP *vs.* Stress PP groups (**Fig. 3D**). Approximately 85% of genes

altered by maternal stress overlapped with parity-induced gene expression changes. KEGG

pathway analysis of DEGs revealed shared enrichment across several pathways in both

comparisons, with maternal stress reversing gene expression associated with long-term

potentiation, dopaminergic synapses, oxytocin signaling, and other processes (**Fig. 3E**).

280
Since our pathway analyses indicated changes in long-term potentiation, a form of neuronal

plasticity that is important for learning and memory behaviors[37], we further focused our behavioral

assessments on contextual fear conditioning. Consistent with prior results, Control PP females

exhibited enhanced acquisition and context recall, as demonstrated by significantly increased

freezing behavior compared to NP animals. In contrast, however, Stress PP females showed no

significant differences *vs.* NP mice (**Fig. 3F-G**). To further evaluate dHpc-dependent function, we

next assessed behavior using the object location task. In this test, animals were initially trained by

allowing them to explore two identical objects, followed by one-hour of removal from the arena.

During the test phase, one object was displaced, and the animals' ability to identify the moved

object was assessed. Given prior findings that adult females require 10 minutes of training for

reliable discrimination[38], we first used this duration to confirm that all groups could successfully

identify the moved object (**Fig. S3J**). To next assess whether Control PP females exhibit

enhancements in this task, we reduced the training time to 5 minutes, a duration that is sufficient

for learning in adult males and preadolescent female mice, but not in adult females[38]. Under these

conditions, Control PP females retained the ability to discriminate the novel location, whereas NP

and Stress PP females did not, reflecting the same pattern observed in contextual fear conditioning,

which was not attributed to estrous stage or locomotion (**Fig. S3K-L**).

298
299 ***Cell-type specific transcriptional alterations reveal dopaminergic modulation underlying***
300 ***maternal dHpc plasticity***
301
To elucidate the mechanisms underlying parity-induced dHpc plasticity, we next leveraged the

transcriptomic and behavioral signatures shared between NP and Stress PP groups. This

postpartum stress paradigm resolves key windows during which disruptions to parity adaptations

occur, thereby providing greater temporal resolution of the mechanisms required to sustain these

alterations. Building on our findings, we next performed single-nuclei (sn) RNA-sequencing on

these three groups to identify the dHpc cell-types that integrate hormone and neurotransmitter

signaling to drive parity-related plasticity. Following rigorous quality control assessments to

remove confounding sources of variation, such as mitochondrial mapping percentage, doublets,

and cell-type contamination **(Fig. S4A-F)**, we retained 109,334 nuclei for downstream analysis

(NP = 37,070; Control PP = 35,631; Stress PP = 36,633). Cell-type annotation was performed

using both manual curation and label transfer from validated hippocampal datasets from the Broad

Institute and Allen Brain Atlas[39,40]. This resulted in 16 distinct clusters (**Fig. 4A, Fig. S4G-I**),

including excitatory neuron subtypes (dentate gyrus, CA1, CA3, subiculum, mossy cells; 43.3%

of total), GABAergic neurons (GABA.1–GABA.4; 9.1% of total), neural progenitors (neural stem

cells, radial glia-like cells; 9.7% of total), and non-neuronal/glial populations (astrocytes,

oligodendrocytes, OPCs, microglia, immune cells; 37.9% of total).

318
Since our cell-type deconvolution analyses indicated downregulation of neuronal markers, we first

assessed proportional differences across major cell-types. A Monte Carlo permutation test

confirmed significant shifts in several populations beyond random variation, with GABA.2 and

GABA.4 clusters being reduced in Control PP dHpc compared to both NP and Stress PP (**Fig. 4B-**

C). We next used pseudobulk aggregation and differential expression analysis to identify DEGs

within each annotated cluster ($p < 0.05$ and logFC > $|1.5|$). We observed transcriptional changes

across both neuronal and non-neuronal subtypes, with excitatory neuron alterations predominantly

observed between NP *vs.* Control PP, particularly within the subiculum (**Fig. 4D**). This pattern

aligns nicely with enhanced contextual fear conditioning, as prior studies have implicated altered

dHpc excitatory outputs in improved learning outcomes in dams[27]. In contrast, the most

pronounced neuronal changes between Control and Stress PP were observed in the GABA.2 and

mossy cell populations, alongside widespread DEGs across non-neuronal cell-types (**Fig. 4D**);

note that differential expression analysis could not be performed for the GABA.4 cluster due to

insufficient sample representation in multiple groups. GO term analysis of DEGs from cell-types

displaying proportional differences revealed significant enrichment in pathways associated with

responses to endogenous stimuli (GABA.2, GABA.1, GABA.3, OPC), cell proliferation

(GABA.2, CA1, immune cells), synaptic function (GABA.3, subiculum), and metabolic processes

(Oligo.2, GABA.1) across both comparisons (**Fig. 4E-F**).

337
Probing deeper into these altered cell-types, we focused on the GABA.2 cluster owing to shared

proportional and transcriptional changes observed in comparison to both NP and Stress PP groups,

which suggested a common mechanism limiting dHpc plasticity. Marker gene analysis revealed

the significant enrichment of dopamine receptor D1 *(Drd1)* and D2 *(Drd2)* expression within this

cluster compared to other cell-types (**Fig. S4J**), aligning with our prior bioinformatic analyses

implicating dopaminergic regulation. To further resolve this population, we subclustered the

GABA.2 cells and identified distinct subpopulations expressing *Drd1* and *Drd2*, both of which

appeared to be altered in Control PP animals (**Fig. 4G**). In a separate cohort of mice, we validated

these changes using RNA fluorescence in situ hybridization (FISH) in dorsal CA1 (**Fig. 4H-I, Fig.**

S4K-L), where dopamine receptor-expressing interneurons have been previously described[41,42].

While dopaminoceptive inhibitory neurons were implicated in our data, we examined whether

such changes extend to other dopamine-sensitive neuronal populations. RNA FISH in the dentate

gyrus, which contains excitatory *Drd1* and *Drd2*-expressing neurons (in granule and hilar mossy

cells, respectively; **Fig. S7A**), revealed significant group differences, alongside expected regional

variations consistent with the sparser distribution of dopamine receptor-expressing neurons in

CA1[43] (**Fig. 4H-I, Fig. S4M-N).**

354
Since these findings established dopaminergic signaling as a potential key mediator of persistent

parity-induced adaptations, this prompted us to examine whether maternal stress disrupts this

process within its defined postpartum window. Given prior evidence that acute pup separation

elevates dopamine levels in NAc[44,45], we first investigated whether a similar response occurs in

dHpc. Brain tissues were collected from 10 dpp dams at baseline, 30-minutes, and 3-hours post-

separation, as well as 30-minutes post-reunion (**Fig. 4J).** We observed robust increases in

dopamine levels in both NAc and dHpc during separation, with expected regional differences

reflecting the strength of innervation to these regions (**Fig. 4K**). Next, we reasoned that this initial

pup separation may reflect a normal, adaptive maternal response to unexpected offspring removal.

Since acute and chronic stress produce different biological responses and behavioral outcomes,

this prompted us to next assess the impact of repeated separation stress on dopamine modulation

in these brain regions. Indeed, our data revealed distinct dopaminergic responses following acute

vs. chronic separations, with repeated stress leading to loss of dopamine increases observed in

response to pup separation, as well as elevated baseline dopamine in both dHpc and NAc tissues

(**Fig. 4L, Fig. S4O**). Loss of induction following chronic stress may stem from elevated baseline

levels, which facilitate engagement of autoreceptors and negative feedback mechanisms[46].

Similarly, increased release of dopamine in dHpc tissues were not observed in NP females 30-

minutes following separation from a littermate (**Fig. S4P**), though this likely lacks the same

salience as pup separation. These findings align well with our snRNA-seq observations, in which

the reduced dopamine receptor expression resulting from parity reflects an adaptive process to

reduce basal dopamine (supported by our transcriptional data in Figs. 2 and 3 showing

downregulation of dopamine processes), thereby enabling robust responsiveness to relevant

stimuli. In contrast, elevated receptor expression in NP and Stress PP animals may be a

downstream effect of greater basal dopamine tone, thereby dampening the efficiency of stimulus-

dependent signaling.

380
Chemogenetic suppression of dopamine release into dHpc recapitulates key features of persistent parity-induced plasticity

383
Based on these observations, we hypothesized that the maintenance of low basal dopamine tone

may drive adaptations of the dopamine-sensitive machinery in dHpc to promote long-term

behavioral plasticity. To assess this, we chemogenetically suppressed the dopaminergic VTA

projection to dHpc by bilaterally expressing a floxed inhibitory neuronal hM4Di-DREADD fused

to mCherry (*vs.* floxed mCherry controls) in the VTA, coupled with retrograde AAVs expressing

a tyrosine hydroxylase (TH)-dependent Cre recombinase into the dHpc (**Fig. 5A**). Selection of the

VTA was guided by our findings that pup separation stress drives coordinated regulation in the

dHpc and NAc, along with its established role as a key upstream modulator of dopaminergic

signaling in the maternal NAc[47] (also supported by our data; **Fig. S4Q**). We validated this approach

by confirming selective mCherry expression in TH+ neurons of the VTA, as well as reduced *Fos*

expression following administration of the selective DREADD agonist deschloroclozapine (DCZ;

Fig. 5B, Fig. S5A-B.) To minimize injection-related stress in this paradigm, we sought to restrict

the window of chronic DCZ administrations. Since maternal stress from 10–20 dpp was sufficient

to disrupt parity-induced plasticity, we posited that dHpc dopamine suppression during this period

may correspondingly be sufficient to promote neuronal adaptations in virgin NP (NP-mCherry,

NP-hM4Di; **Fig. 5C**). As our comparisons aimed to determine if chronic dopamine reduction in

virgin females can phenocopy parity-induced adaptations, we administered DCZ to postpartum

dams in parallel (PP-mCherry, PP-hM4Di).

402
After confirming that our approach effectively suppressed dHpc dopamine (**Fig. 5D**) – and did not

induce gross metabolic changes (**Fig. S5C**) – we assessed behavioral outcomes 1-month after the

final DCZ injection (49 dpp). In the pup retrieval test, NP-hM4Di females exhibited significantly

reduced latency to retrieve pups compared to NP-mCherry, with no differences observed relative

to PP dams (**Fig. 5E**). Notably, while 8/13 NP-mCherry females failed to retrieve either pup, this

occurred in only 1/15 NP-hM4Di animals, comparable to PP-mCherry (0/12) and PP-hM4di (1/11)

dams. In contextual fear conditioning, NP-hM4Di females demonstrated enhanced acquisition,

mirroring the heightened learning observed in PP dams (**Fig. 1F**). In the context recall test, NP-

hM4Di females showed no significant differences from PP dams (**Fig. S5D**), though subtle effects

were observed, suggesting that additional neuromodulatory mechanisms may interact with

dopamine to modulate the consolidation and retrieval of conditioned fear responses underlying

maternal behavioral plasticity.

415
Next, we examined whether such dopamine suppression influences cellular receptor expression in

the dHpc. Similar to Control PP, NP-hM4Di females exhibited sustained downregulation of *Drd1*

and *Drd2* in both CA1 and dentate gyrus (**Fig. 5G-H, Fig. S5E-H**), thus highlighting the lasting

impact of prior dopamine modulation on dHpc receptor dynamics. Finally, we conducted

transcriptional profiling of the dHpc in virally infected groups to assess the degree to which

dopamine suppression mirrors parity-associated gene expression patterns. Comparing NP-

mCherry *vs.* PP-mCherry expression profiles from bulk RNA-seq data, we observed an

intermediate expression pattern in NP-hM4Di dHpc tissues (**Fig. 5I, Table S7**). RRHO analysis

further supported transcriptional concordance between PP-mCherry and NP-hM4Di groups in

comparison to NP-mCherry (**Fig. 5J**). To next assess whether the long-term transcriptional effects

of prior baseline dopamine suppression align with changes in parity-associated transcriptional

programs, we compared NP-mCherry *vs.* NP-hM4Di DEGs to previously identified parity-

sensitive modules. Consistent with our behavioral findings, DEGs from NP-hM4Di overlapped

with the brown module but not other parity-sensitive modules (**Fig. 5K**), thereby highlighting a

specific mechanistic contributor to behavioral adaptations within the broader transcriptional and

neuromodulatory landscape shaped by parity. Functional annotation of these DEGs revealed

enrichment in pathways governing synaptic plasticity and responses to endogenous stimuli (**Fig.**

5J), which mirrored those found to be altered in Control PP dHpc and reinforced dopamine

modulation as a key driver of long-lasting maternal brain adaptations (**Fig. S5I**).

435
DISCUSSION

Despite decades of evidence that matrescence induces long-lasting behavioral adaptations across

species, the mechanisms underlying these phenomena remain poorly understood. Through

controlled brain-wide, time-course, and cell type-specific transcriptomic analyses – paired with

robust behavioral outputs, – we identified gene networks, reproductive events, and

neuromodulatory pathways that dynamically shape regional sensitivity across pregnancy and

postpartum experiences, thereby promoting plasticity long after these stages have ended.

Furthermore, using a maternal-pup separation paradigm during postpartum, we found that

modulation of dopamine dynamics is essential for sustaining long-term neural and behavioral

adaptations in maternal dHpc. Given dopamine's role in reinforcing maternal behavior and reward,

the observed downregulation of dopamine pathways in maternal brain was unexpected, though

supported by a prior study[48]. Interestingly, reducing baseline dopamine may enhance the signal-

to-noise ratio (SNR) of maternal responses to relevant cues, whereas elevated baseline dopamine

levels may impair this process. Indeed, in this study we established the necessity and sufficiency

of maintaining appropriate dopamine tone for parity-related dHpc plasticity, demonstrating that

chronic dopamine elevation through maternal stress, and chemogenetic suppression within virgin

dHpc, bidirectionally modulate parity-related changes at transcriptional, behavioral, and cellular

levels. Such findings thus implicate dopamine as a key driver of long-term maternal brain

remodeling. These findings expand upon the prevailing view that maternal dopamine function is

solely driven by transient surges during maternal behaviors, instead revealing a concurrent

downregulation of dopamine tone in critical regions, such as the dHpc, to optimize stimuli

sensitivity and behavioral responsiveness.

458
Ethological significance of maternal dHpc plasticity

Why the dHpc? Improvement of classic dHpc functions, such as spatial cognition, represents

evolutionarily advantageous modes for maternal success by enabling more efficient resource

foraging and nest navigation[49,50]. Additional dHpc functions, including novelty detection and

sensory integration, may also heighten sensitivity to salient environmental cues that are essential

for offspring survival[51,52]. Indeed, heightened hippocampal activation in response to one's own

infant's cry in human fMRI studies underscores the diverse functional adaptations of this region[53],

suggesting a broad range of dHpc modifications that may converge to enhance maternal success.

Although chronic maternal stress disrupts the normal trajectory of dHpc adaptations, it remains

unclear whether such disruptions are maladaptive or represent evolved coping mechanisms in

response to repeated separations. Furthermore, whether stress effects arise from reduced pup

interactions, chronic stress exposures, or increased licking, grooming, and/or nursing upon pup

reunion is difficult to disentangle within the current model[54]. Notably, unexpected parenting

disruptions and resulting adjustments generate significant stress, with parenting-related

psychosocial stress uniquely contributing to perinatal and postpartum disorders[55]. Moreover, as

pup separation only reduces total pup exposure by ~6.5%, it is unlikely that reduced pup

interactions alone account for these effects. Notably, chronic restraint stress during pregnancy

similarly disrupts parity-induced spatial memory enhancements[27]. Moreover, human studies show

that maternal early life stress alters hippocampal responses to infant cues[56,57], supporting chronic

stress as a common interferer in the normal trajectory of maternal dHpc neuroadaptations.

Interestingly, we observed natural variation in maternal responses to both acute and chronic stress,

as reflected by variability in dHpc transcriptional changes, behavioral outcomes, and dopamine

release following pup separation. This suggests that individual differences in maternal motivation

or caregiving may influence dopaminergic responses to pup separation, as previously reported[45,58].

Interestingly, while the spectrum of outcomes in Stress PP dams indicate variability in stress

effects, corticosterone levels were uniformly elevated across all dams following pup separation.

This suggests that while chronic stress perturbs parity adaptations, these disruptions are mediated

by individual dopamine dynamics downstream of maternal stress responses.

Dopaminergic regulation of parity adaptations

Our findings, which reveal largescale transcriptional re-programming across the maternal brain,

suggest that regions beyond the dHpc also undergo dopamine-mediated adaptations. Indeed, pup

separation similarly increases dopamine release in the NAc[44,45], while chronic separations

dysregulate this induction by elevating basal dopamine concentrations, thereby facilitating

engagement of negative feedback mechanisms[46]. Importantly, this perturbation offers insight into

normal maternal dopamine regulation, emphasizing the need for low postpartum dopamine tone to

enable signaling induction to relevant stimuli[27,59]. Moreover, as restraint stress enhances

dopaminergic burst firing, and chronic gestational restraint impairs spatial learning, we propose

that dopamine dysregulation serves as a common mode of dysregulating parity-related

neuroadaptations across reproductive stages.

How do postpartum alterations drive sustained behavioral changes? Our findings reveal that

downregulation of dopamine transcriptional networks in Control PP dHpc corresponds with

reduced *Drd1/Drd2* expression, whereas elevated baseline dopamine in Stress PP associates with

increased receptor expression, likely reflecting compensations to tonic dopamine dynamics during

matrescence and/or chronic stress, rather than loss of dopaminoceptive cells. Notably, receptor

downregulation may also contribute to structural plasticity, as parity is linked to dendritic

remodeling in CA1 and CA3[7], paralleling gray matter reductions observed in mothers years after

parturition[1,2,4,22]. Thus, our work points to two mechanisms potentially underlying maternal dHpc

adaptations: alterations in *Drd1/Drd2* receptor expression and shifts in baseline dopamine tone.

507
Given that *Drd1/Drd2*-expressing GABAergic neurons contribute to both aversive contextual

memory formation and maternal behavior[45,60–64], receptor expression changes likely intersect with

these functions. Alternatively, receptor shifts may not directly influence behavior but instead

reflect downstream consequences of altered dopamine tone. Within this framework, maintenance

of low dopamine tone in Control PP dHpc may serve to reduce background noise, enhancing the

processing of relevant cues by increasing SNR[65]. In addition to limiting negative feedback, low

dopamine tone also prolongs and amplifies phasic dopamine responses to stimuli[66]. As phasic

dopamine responses are essential for memory formation[67,68], this process may underlie parity-

induced enhancements in fear conditioning outcomes. While not traditionally considered a

maternal behavior-regulating region, the dHpc integrates sensory, novelty, and salience

information to shape pup retrieval responses. Consistent with our findings, lesion studies

corroborate hippocampal involvement in pup retrieval behavior[69,70]. Moreover, hippocampal

activity modulates downstream circuits via the polysynaptic VTA-hippocampal loop, which

projects to the NAc and ventral pallidum - key regulators of pup retrieval[52,63]. Enhanced SNR

within the dHpc may thus enable more efficient processing of offspring cues, facilitating robust

behavioral responses to both rewarding and aversive stimuli through signal processing across this

broader network.

These findings expand upon a broader maternal neuroadaptive process in which enhanced SNR

optimizes responsiveness to offspring-related stimuli. Similar mechanisms have been described

for oxytocin signaling, which enhances sensory salience by increasing SNR across multiple brain

regions[71]. For example, in the left auditory cortex, oxytocin disinhibits pyramidal neuron activation

in response to pup calls, enabling pup retrieval behaviors in previously non-maternal virgins[72]. In

humans, disinhibition of cognitive and affective regions has also been associated with more

responsiveness in maternal brains[73]. Similarly, estradiol and progesterone modulate SNR by

reducing baseline activity of galanin-expressing mPOA neurons, leading to a reduced

depolarization block that enhances responsiveness to pup stimuli[18]. Although these effects occur

acutely in response to pregnancy hormones, they persist beyond postpartum[18], mirroring the long-

term neural adaptations observed in our study.

Broader implications of maternal brain remodeling

While our study focused largely on dopamine, it is notable that our chemogenetic manipulations

recapitulated key transcriptional and behavioral outcomes of parity, albeit not to the same extent.

This underscores dopamine as a crucial but not exclusive driver of parity-induced adaptations.

Indeed, our data also implicate hormonal contributions, including estrogen, progesterone, and

oxytocin signaling, in shaping transcriptional programming. Consistent with this, previous studies

have linked parity to estradiol-driven neuronal alterations[12], while others have demonstrated that

blocking oxytocin signaling disrupts parity-induced spatial learning enhancements[30]. These

findings suggest that parity programming arises from integration of multiple neuromodulatory

pathways to induce long-term effects. However, given that estrogen signaling during postpartum

regulates oxytocin release, which in turn modulates VTA signaling, we propose that dopamine

adaptations serve as a key downstream mechanism within a broader network driving sustained

maternal neuroplasticity[74].

What upstream mechanisms may regulate dopamine adaptations in maternal brain? As discussed,

oxytocin regulates VTA dopamine release, yet whether it contributes to enduring changes in VTA-

dHpc tone remains unknown. Additionally, the mechanisms by which chronic maternal stress

alters this dopaminergic tone warrant further investigation. While this study focused on the VTA

as an upstream provider of dHpc dopamine, the LC also provides substantial dopaminergic input

to the dHpc, suggesting broader brain region involvement in dopamine-mediated adaptations[68,75].

Given its additional involvement in stress signaling, the LC, alongside the VTA, may contribute

to signaling processes underlying chronic maternal stress effects. Furthermore, other brain regions,

such as hypothalamic inputs, may modulate dHpc dopamine tone in response to stress, representing

important avenues for future investigation.

Together, this work demonstrates that transient neuromodulatory processes during pregnancy and

postpartum drive lasting maternal neuroplasticity. Importantly, this fundamental shift in brain state

may interact with future experience-dependent plasticity. As parity is implicated in both risk and

resilience to brain disorders[15], future research should account for parity status as a key variable

shaping differential outcomes, particularly in its interactions with other risk factors. In particular,

our findings highlight an interplay between parity and stress, a key risk factor for brain disorder

vulnerability, in shaping maternal brain outcomes. Thus, this study provides novel insights into

the gene networks, dynamics, and neuromodulatory pathways that drive long-lasting parity-induced adaptations, while underscoring the importance of stress mitigation during pregnancy and postpartum.

ACKNOWLEDGEMENTS

We would like to thank members of the Maze lab for their helpful discussion on this study, and members of the Nestler, Russo, and Kenny labs at the Icahn School of Medicine at Mount Sinai for providing materials and/or access to behavioral equipment. We thank Dr. Min Chen, Winnie Chen, Emma Andraka, and Rasika Iyer for experimental assistance. This work was partially supported by grants from the National Institutes of Health: R01 MH116900 (I.M.), F32 MH126534 (J.C.C.), F31 NS132558 (A.M.C.), F99 NS139541 (A.M.C.), as well as funds from the Brain and Behavior Research Foundation (J.C.C.), and the Howard Hughes Medical Institute (I.M.). All schematics were created with Biorender.com.

DECLARATION OF INTEREST

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

J.C.C. and I.M. conceived of the study, designed the experiments, and interpreted the data. J.C.C., G.D.S., A.M.C., and S.D. performed behavioral experiments. J.C.C., G.D.S., A.M.C., and S.D. performed surgeries. J.C.C., G.D.S., and C.Z. performed estrous cycle analyses. J.C.C., G.D.S., and S.D. conducted immunofluorescence and RNAscope analyses. J.C.C. and G.D.S. performed molecular experiments. J.C.C. and E.A.B. performed single-cell sequencing experiments. J.C.C., G.D.S., and E.W. performed bioinformatic analyses. J.C.C. and I.M. wrote the manuscript.

Figure 1. Pregnancy and postpartum states promote lasting transcriptional and behavioral adaptations in the maternal brain. A) Experimental timeline comparing primiparous (PP) *vs.* age-matched nulliparous (NP) female mice 1-month after offspring weaning (day postpartum, 49 dpp), and behavioral outcomes at 140 dpp. **B)** Differentially expressed genes (DEGs) per brain region. Bubble size represents total number of NP *vs.* PP DEGs (adj. p < 0.05). **C)** Cell-type deconvolution of bulk RNA-seq data (NP *vs.* PP normalized surrogate proportion variables, Student's t-test with multiple comparisons correction; *FDR < 0.05, **FDR < 0.01, ***FDR < 0.001, ****FDR < 0.0001). **D)** Downregulation of neuron proportion significantly correlated with number of DEGs per brain region ($R^2 = 0.4198$, p = 0.0311). **E)** Overlap of DEGs across brain regions (p < 0.05; gray indicates nonsignificant overlap). **F)** Select predicted upstream regulators of overlapping DEGs across regions with high (dHpc, NAc, mPOA, LC, vHpc) *vs.* low (DRN, PVN, BLA, VTA, OB, mPFC) sensitivity to parity. **G)** Trait heatmap correlating gene co-expression modules (arbitrary color names, identified by weighted gene correlation network analysis) with group x brain region classification (*p < 0.05, #p < 0.1). **H)** Circos plot for gene co-expression modules, with the size and arbitrary color name of each module indicated by the arc thickness along the perimeter. Enrichment for DEGs per brain region for each module is indicated by the inner rings, with color indicating significance of overlap. **I)** Select GO Biological Processes

and KEGG Pathways significantly enriched from gene co-expression modules (FDR < 0.05). **J)** PP dams retrieved both pups significantly faster compared to NP females at 140 dpp (two-way rmANOVA, main effect of group ($F(1,11) = 14.47$, $p = 0.0029$), main effect of pup number ($F(1,11) = 11.36$, $p = 0.0062$)), with significantly reduced latency for retrieval of pup 1 (Holm-Šídák's multiple comparisons test, $t(22) = 2.301$, *adj. $p = 0.0312$) and pup 2 (Holm-Šídák's multiple comparisons test, $t(22) = 4.098$, ***adj. $p = 0.0009$). N = 6-7 animals/group. **K)** PP dams exhibited increased freezing behavior during the acquisition phase of contextual fear conditioning (two-way rmANOVA, main effect of group ($F(2,21) = 4.259$, $p = 0.028$), main effect of preshock ($F(2.827, 59.36) = 26.59$, $p < 0.0001$), interaction ($F(8,84) = 5.091$, $p < 0.0001$)), compared to no shock controls (Dunnett's multiple comparisons test, *$p \leq 0.05$, **$p < 0.01$). **L)** During the context recall test, PP dams froze significantly more compared to NP (one-way ANOVA, $F(2,22) = 18.72$, $p < 0.0001$; Tukey's multiple comparisons test, $t(22) = 3.640$, *$p = 0.044$), with both groups demonstrating contextual learning compared to no shock controls (PP vs. NS, $t(22) = 8.541$, ****$p < 0.0001$; NP vs. NS, $t(22) = 5.538$, **$p = 0.0021$). N = 7-10 animals/group. Error bars represent mean ± SEM.

Figure 2. Parity adaptations in the dHpc transcriptome integrate pregnancy and postpartum experiences. A) Experimental design to examine the contribution of discrete female reproductive events. **B)** Heatmap of differential expression profiles of top 500 genes, by ascending p-values, between NP *vs.* PP (N=3-7/group). **C)** Overlap of DEGs (*vs.* NP, adj. p < 0.05) for each gene co-expression module, with color representing extent of significance (*p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001). High parity sensitivity modules are indicated in bold. **D)** Threshold-free comparison of indicated comparisons by rank-rank hypergeometric overlap. Pixels represent the overlap of differential expression profiles indicated, with color representing extent of significance. The lower left and upper right quadrants represent concordant gene regulation. **E)** Shared changes in biological processes (identified through GO term analysis) observed following individual reproductive exposures with PP, along with processes exclusive to the combined experiences in PP (FDR < 0.05). **F)** Experimental design to examine the trajectory of PP dHpc gene expression changes across pregnancy and postpartum periods (dpc: days post-conception; dpp: days postpartum). **G)** Enrichment of DEGs (*vs.* NP, adj. p < 0.05) with parity-sensitive gene co-expression modules across reproductive time points, examining extent of significant overlap in brain regions exhibiting high (left, dHpc), moderate (middle, vHpc), and low (right, mPFC) transcriptional sensitivity to parity status (*p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001). N=6/group. **H)** Impulse time course analysis of dHpc gene expression identified patterns of transient *vs.* persistent up- and downregulation across pregnancy and postpartum (FDR < 0.05). **I-L)** Select GO terms and KEGG pathways significantly enriched for dHpc genes exhibiting patterns

658   of **(I)** persistent downregulation, **(J)** persistent upregulation, **(K)** transient downregulation, and **(L)**
659   transient upregulation across reproductive stages (FDR < 0.05).
660

Figure 3. Postpartum stress disrupts long-term maternal dHpc adaptations. A) Postpartum stress paradigm, wherein dams were provided with limited nesting and subjected to pup separation 3 hours daily between 10-20 dpp, followed by behavioral and transcriptomic assessments at 49 dpp. **B)** Principal components analysis of dHpc transcriptomes, and **C)** heatmap of all significant DEGs (adj. p < 0.05) with hierarchical clustering. **D)** Venn Diagrams showing ~20% DEGs altered in Control PP (*vs.* NP) are significantly disrupted by postpartum stress (adj. p < 0.05). **E)** Parity-dependent enrichment in select KEGG pathways are significantly reversed in Stress PP (FDR < 0.05). **F)** Control PP dams froze more during the acquisition phase of contextual fear conditioning (two-way rmANOVA, main effect of group (F(2,31) = 4.662, p = 0.017), main effect of preshock (F(3.024,93.75) = 129.8, p < 0.0001), interaction (F(8,124) = 2.406, p = 0.091)), compared to NP and Stress PP (Tukey's multiple comparisons test, #p ≤ 0.1, *p < 0.05, **p < 0.01). **G)** During the context recall test, NP froze significantly less than Control PP (one-way ANOVA, F(2,31) = 3.151, p = 0.05; Tukey's multiple comparisons test, t(31) = 3.54, *p = 0.0456), but not Stress PP (ns, p > 0.05). N = 11-12 animals/group. **H)** Control PP, but not Stress PP, dams spent more time investigating the moved object in the object location task compared to NP (one way ANOVA,

678    $F_{(2,30)} = 4.045$, $p = 2.0278$; Holm-Šídák's multiple comparisons test: NP vs. Control PP, $t(30) =$
679   2.764, *adj. $p = 0.0288$; Control PP vs. Stress PP, $t(30) = 2.046$, #adj. $p < 0.1$) N=10-12
680   animals/group. Error bars represent mean $\pm$ SEM.
681

Figure 4. Parity-driven dHpc plasticity is suppressed by elevated dopamine signaling.

Figure 4. Parity-driven dHpc plasticity is suppressed by elevated dopamine signaling. A)
UMAP representation of cell clusters (109,334 total cells) from dHpc tissues, colored by major
cell types (N = 6 NP, 4 Control PP, 5 Stress PP). **B, C)** Point range plots depicting significant
changes in proportions of major cell-type clusters between **(B)** NP vs. Control PP, and **(C)** Stress
PP vs. Control PP (FDR < 0.05 and |log₂Differerence| > log₂(1.5)). **D)** Number of DEGs observed
for each major cell-type following pseudobulk analysis (p < 0.05 and |log₂(Fold Change)| >
log₂(1.5)). **E, F)** GO term analyses (Biological Process) for DEGs identified in cell clusters with

significant proportional differences between **(E)** NP vs. Control PP, and **(F)** Stress vs. Control PP
(FDR < 0.05). **G)** Subclustering of GABA.2 neuronal population revealed reductions in *Drd1* (top)
and *Drd2* (bottom) expression in Control PP. **H)** Representative images of RNAScope for *Drd1*
and *Drd2* mRNAs in dorsal CA1 (top) and dentate gyrus (DG, bottom). Scale bars, 50 μm. **I)**
Quantification of *Drd1* and *Drd2* mRNA puncta in CA1 and DG nuclei (two-way ANOVA, main
effect of group ($F_{(2,82142)}$ = 65.81, $p < 0.0001$), main effect of gene ($F_{(1,82142)}$ = 1771, $p <$
0.0001), interaction ($F_{(2,82142)}$ = 39.99, $p < 0.0001$); Tukey's multiple comparisons test, **p <
0.01, ***p < 0.001, ****p < 0.0001). N = 5 animals/group. **J)** Experimental design of pup
separation test to examine changes in brain tissue dopamine levels. **K)** Pup separation significantly
increases dopamine levels from baseline in dHpc and NAc tissues (two-way ANOVA, main effect
of time ($F_{(3,52)}$ = 4.287, $p = 0.0089$), main effect of brain region ($F_{(1,52)}$ = 225.7, $p < 0.0001$;
interaction ($F3,52$) = 1.086, $p = 0.3717$), with significant increases in dHpc dopamine levels
(Dunnett's multiple comparisons test, *p < 0.05, **p < 0.01). N = 5-9 animals/time point. **L)**
Chronic pup separations increases baseline dHpc dopamine concentrations (two-way ANOVA,
interaction ($F_{(3,53)}$ = 4.822, $p = 0.0048$; group $(1, 53)$ = 1.876), $p = 0.1708$, time $(3, 53)$ = 1.736,
$p = 0.1706$; Tukey's multiple comparisons test, *p < 0.05, #p = 0.06). N = 5-12 animals/time point.
Error bars represent mean ± SEM.
709

Figure 5. Chronic dopamine suppression is sufficient to phenocopy persistent maternal dHpc plasticity. A) Experimental design (left) and representative image (right) of AAV2.hSyn.DIO.mCherry targeting to VTA projection neurons following dHpc injection of

retrograde (rg) Cre under control of the tyrosine hydroxylase (TH) promoter. Scale bar, 100 μm.
B) Representative images showing overlap of virally-induced mCherry with TH+ cells in VTA.
Scale bar, 20 μm. **C)** Timeline of stereotaxic surgeries and s.c. deschloroclozapine (DCZ)
injections from 10-20 dpp in NP and PP female mice, followed by experimental assessments
starting at 49 dpp. **D)** Significant reduction in dHpc dopamine levels following inhibition of VTA
projection neurons (Student's t-test:t(8) = 2.477, *p = 0.0383). N=5/group. **E)** PP-mCherry, NP-
hM4Di, and PP-hM4Di retrieved pups faster than NP-mCherry (two-way rmANOVA, main effect
of group (F(3,47) = 14.39, p < 0.0001), main effect of pup number (F(1, 47) = 18.96, p < 0.0001;
Sidak's multiple comparisons test, ****p < 0.0001, ***p < 0.001). N =11-15/group. Bar
represents mean. **F)** PP-mCherry, NP-hM4Di, and PP-hM4Di freeze more compared to no shock
controls at earlier phases of contextual fear acquisition, compared to NP-mCherry (two-way
rmANOVA, main effect of group (F(4,41) = 4.739, p = 0.0031), main effect of preshock
(F(2.818,115.5) = 61.92, p < 0.0001), interaction (F(16,164) = 2.606, p = 0.0012; Tukey's multiple
comparisons test, *p ≤ 0.05, **p < 0.01, ****p < 0.0001). N = 9-12/group. Error bars represent
mean ± SEM. **G)** Representative images of RNAScope for *Drd1* and *Drd2* mRNAs in dorsal CA1
(top) and DG (bottom). Scale bars, 50 μm. **H)** Quantification of *Drd1* and *Drd2* mRNA puncta in
CA1 and DG nuclei (two-way ANOVA, main effect of group (F(2,33404) = 64.43, p < 0.0001),
main effect of gene (F(1,33404) = 2770, p < 0.0001), interaction (F(2,33404) = 12.52, p = 0.0004);
Sidak's multiple comparisons test, **p < 0.01, ****p < 0.0001). N = 3 animals/group. **I)** Bulk
dHpc differential expression profiles of top 500 genes (by ascending p-values) between NP-
mCherry and PP-mCherry. N=6/group. **J)** Threshold-free comparison by rank-rank
hypergeometric overlap, showing concordant gene regulation (bottom left and upper right
quadrants) between the comparisons indicated. **K)** Overlap of DEGs (NP-hM4Di vs. NP-mCherry,
adj. p < 0.05) with gene co-expression modules (****p < 0.0001). High parity sensitivity modules
are indicated in bold. **L)** Select GO terms and KEGG pathways significantly enriched for NP-
mCherry vs. NP-hM4Di DEGs (FDR < 0.05).
742

743 **SUPPLEMENTARY MATERIALS**
744
745 Materials and Methods
746 Figs. S1 to S5
747 Tables S1 to S7
748 Supplemental References
749

REFERENCES

1. Hoekzema, E. *et al.* Pregnancy leads to long-lasting changes in human brain structure. *Nature neuroscience* **20**, 287–296 (2017).
2. Martínez-García, M. *et al.* Do Pregnancy-Induced Brain Changes Reverse? The Brain of a Mother Six Years after Parturition. *Brain sciences* **11**, 168 (2021).
3. Voldsbekk, I. *et al.* A history of previous childbirths is linked to women's white matter brain age in midlife and older age. *Human Brain Mapping* **42**, 4372–4386 (2021).
4. Pritschet, L. *et al.* Neuroanatomical changes observed over the course of a human pregnancy. *Nat Neurosci* 1–8 (2024) doi:10.1038/s41593-024-01741-0.
5. de Lange, A.-M. G. *et al.* Population-based neuroimaging reveals traces of childbirth in the maternal brain. *Proceedings of the National Academy of Sciences* **116**, 22341 LP – 22346 (2019).
6. de Lange, A.-M. G. *et al.* The maternal brain: Region-specific patterns of brain aging are traceable decades after childbirth. *Human Brain Mapping* **41**, 4718–4729 (2020).
7. Pawluski, J. L. & Galea, L. A. M. Hippocampal morphology is differentially affected by reproductive experience in the mother. *Journal of neurobiology* **66**, 71–81 (2006).
8. Duarte-Guterman, P. *et al.* Cellular and molecular signatures of motherhood in the adult and ageing rat brain. *Open Biology* **13**, 230217 (2023).
9. Workman, J. L. *et al.* Parity modifies the effects of fluoxetine and corticosterone on behavior, stress reactivity, and hippocampal neurogenesis. *Neuropharmacology* **105**, 443–453 (2016).
10. Macbeth, A. H., Scharfman, H. E., Maclusky, N. J., Gautreaux, C. & Luine, V. N. Effects of multiparity on recognition memory, monoaminergic neurotransmitters, and brain-derived neurotrophic factor (BDNF). *Hormones and behavior* **54**, 7–17 (2008).
11. Kinsley, C. H. & Lambert, K. G. Reproduction-Induced Neuroplasticity: Natural Behavioural and Neuronal Alterations Associated with the Production and Care of Offspring. *Journal of Neuroendocrinology* **20**, 515–525 (2008).
12. Barha, C. K., Lieblich, S. E., Chow, C. & Galea, L. A. M. Multiparity-induced enhancement of hippocampal neurogenesis and spatial memory depends on ovarian hormone status in middle age. *Neurobiology of Aging* **36**, 2391–2405 (2015).
13. Meurisse, M. *et al.* Estradiol receptor-α expression in hypothalamic and limbic regions of ewes is influenced by physiological state and maternal experience. *Hormones and Behavior* **48**, 34–43 (2005).
14. Love, G. *et al.* Maternal Experience Produces Long-Lasting Behavioral Modifications in the Rat. *Behavioral Neuroscience* **119**, 1084–1096 (2005).
15. Deems, N. P. & Leuner, B. Pregnancy, postpartum and parity: Resilience and vulnerability in brain health and disease. *Frontiers in Neuroendocrinology* **57**, 100820 (2020).
16. Figueiredo, B. & Conde, A. Anxiety and depression symptoms in women and men from early pregnancy to 3-months postpartum: Parity differences and effects. *Journal of Affective Disorders* **132**, 146–157 (2011).
17. Martínez-Galiano, J. M., Hernández-Martínez, A., Rodríguez-Almagro, J., Delgado-Rodríguez, M. & Gómez-Salgado, J. Relationship between parity and the problems that appear in the postpartum period. *Sci Rep* **9**, 11763 (2019).
18. Ammari, R. *et al.* Hormone-mediated neural remodeling orchestrates parenting onset during pregnancy. *Science* **382**, 76–81 (2023).

19. Leuner, B., Glasper, E. R. & Gould, E. Parenting and plasticity. *Trends in Neurosciences*
33, 465–473 (2010).
20. Bridges, R. S. Neuroendocrine Regulation of Maternal Behavior. *Front Neuroendocrinol* **0**,
178–196 (2015).
21. McKenzie, A. T. *et al.* Brain Cell Type Specific Gene Expression and Co-expression
Network Architectures. *Scientific Reports* **8**, 8868 (2018).
22. Paternina-Die, M. *et al.* Women's neuroplasticity during gestation, childbirth and
postpartum. *Nat Neurosci* **27**, 319–327 (2024).
23. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics* **9**, 559 (2008).
24. Bridges, R. S. Long-term effects of pregnancy and parturition upon maternal
responsiveness in the rat. *Physiology & Behavior* **14**, 245–249 (1975).
25. Pawluski, J. L., Vanderbyl, B. L., Ragan, K. & Galea, L. A. M. First reproductive
experience persistently affects spatial reference and working memory in the mother and
these effects are not due to pregnancy or 'mothering' alone. *Behavioural Brain Research*
175, 157–165 (2006).
26. Pawluski, J. L., Walker, S. K. & Galea, L. A. M. Reproductive experience differentially
affects spatial reference and working memory performance in the mother. *Horm Behav* **49**,
143–149 (2006).
27. Lemaire, V. *et al.* Motherhood-induced memory improvement persists across lifespan in
rats but is abolished by a gestational stress. *The European journal of neuroscience* **23**,
3368–3374 (2006).
28. Gatewood, J. D. *et al.* Motherhood mitigates aging-related decrements in learning and
memory and positively affects brain aging in the rat. *Brain research bulletin* **66**, 91–98
(2005).
29. Macbeth, A. H., Gautreaux, C. & Luine, V. N. Pregnant rats show enhanced spatial
memory, decreased anxiety, and altered levels of monoaminergic neurotransmitters. *Brain*
research **1241**, 136–147 (2008).
30. Tomizawa, K. *et al.* Oxytocin improves long-lasting spatial memory during motherhood
through MAP kinase cascade. *Nature Neuroscience* **6**, 384–390 (2003).
31. Paris, J. J. & Frye, C. A. Estrous cycle, pregnancy, and parity enhance performance of rats
in object recognition or object placement tasks. *Reproduction* **136**, 105–115 (2008).
32. Stolzenberg, D. S. & Rissman, E. F. Oestrogen-independent, experience-induced maternal
behaviour in female mice. *J Neuroendocrinol* **23**, 345–354 (2011).
33. Cahill, K. M., Huo, Z., Tseng, G. C., Logan, R. W. & Seney, M. L. Improved identification
of concordant and discordant gene expression signatures using an updated rank-rank
hypergeometric overlap approach. *Scientific Reports* **8**, 9588 (2018).
34. Fischer, D. S., Theis, F. J. & Yosef, N. Impulse model-based differential expression
analysis of time course sequencing data. *Nucleic Acids Research* **46**, e119 (2018).
35. McArdle, H. J. The metabolism of copper during pregnancy —a review. *Food Chemistry*
54, 79–84 (1995).
36. Leuner, B. & Sabihi, S. The birth of new neurons in the maternal brain: Hormonal
regulation and functional implications. *Frontiers in neuroendocrinology* **41**, 99–113
(2016).
37. Lynch, M. A. Long-Term Potentiation and Memory. *Physiological Reviews* **84**, 87–136
(2004).

38. Le, A. A. *et al.* Prepubescent female rodents have enhanced hippocampal LTP and learning
relative to males, reversing in adulthood as inhibition increases. *Nat Neurosci* **25**, 180–190
(2022).
39. Habib, N. *et al.* Div-Seq: Single-nucleus RNA-Seq reveals dynamics of rare adult newborn
neurons. *Science* **353**, 925–928 (2016).
40. Yao, Z. *et al.* A taxonomy of transcriptomic cell types across the isocortex and hippocampal
formation. *Cell* **184**, 3222-3241.e26 (2021).
41. Gangarossa, G. *et al.* Characterization of dopamine D1 and D2 receptor-expressing neurons
in the mouse hippocampus. *Hippocampus* **22**, 2199–2207 (2012).
42. Puighermanal, E. *et al.* Anatomical and molecular characterization of dopamine D1
receptor-expressing neurons of the mouse CA1 dorsal hippocampus. *Brain Struct Funct*
222, 1897–1911 (2017).
43. Godino, A. *et al.* Dopaminoceptive D1 and D2 neurons in ventral hippocampus arbitrate
approach and avoidance in anxiety. 2023.07.25.550554 Preprint at
https://doi.org/10.1101/2023.07.25.550554 (2023).
44. Hansen, S., Bergvall, Å. H. & Nyiredi, S. Interaction with pups enhances dopamine release
in the ventral striatum of maternal rats: A microdialysis study. *Pharmacology Biochemistry*
and Behavior **45**, 673–676 (1993).
45. Champagne, F. A. *et al.* Variations in nucleus accumbens dopamine associated with
individual differences in maternal behavior in the rat. *J Neurosci* **24**, 4113–4123 (2004).
46. Ford, C. P. The Role of D2-Autoreceptors in Regulating Dopamine Neuron Activity and
Transmission. *Neuroscience* **282**, 13–22 (2014).
47. Shahrokh, D. K., Zhang, T.-Y., Diorio, J., Gratton, A. & Meaney, M. J. Oxytocin-
Dopamine Interactions Mediate Variations in Maternal Behavior in the Rat. *Endocrinology*
151, 2276–2286 (2010).
48. Afonso, V. M., Grella, S. L., Chatterjee, D. & Fleming, A. S. Previous maternal experience
affects accumbal dopaminergic responses to pup-stimuli. *Brain Research* **1198**, 115–123
(2008).
49. Pawluski, J. L., Lambert, K. G. & Kinsley, C. H. Neuroplasticity in the maternal
hippocampus: Relation to cognition and effects of repeated stress. *Hormones and Behavior*
77, 86–97 (2016).
50. Kinsley, C. H. *et al.* Motherhood Induces and Maintains Behavioral and Neural Plasticity
across the Lifespan in the Rat. *Arch Sex Behav* **37**, 43–56 (2008).
51. Jeffery, K. J. Integration of the sensory inputs to place cells: What, where, why, and how?
Hippocampus **17**, 775–785 (2007).
52. Lisman, J. E. & Grace, A. A. The Hippocampal-VTA Loop: Controlling the Entry of
Information into Long-Term Memory. *Neuron* **46**, 703–713 (2005).
53. Musser, E. D., Kaiser-Laurent, H. & Ablow, J. C. The neural correlates of maternal
sensitivity: An fMRI study. *Dev Cogn Neurosci* **2**, 428–436 (2012).
54. Orso, R. *et al.* How Early Life Stress Impact Maternal Care: A Systematic Review of
Rodent Studies. *Front Behav Neurosci* **13**, 197 (2019).
55. Batt, M. M., Duffy, K. A., Novick, A. M., Metcalf, C. A. & Epperson, C. N. Is Postpartum
Depression Different From Depression Occurring Outside of the Perinatal Period? A
Review of the Evidence. *Focus (Am Psychiatr Publ)* **18**, 106–119 (2020).

56.   Neukel, C. *et al.* The maternal brain in women with a history of early-life maltreatment: an
imagination-based fMRI study of conflictual versus pleasant interactions with children. *J*
Psychiatry Neurosci **43**, 273–282 (2018).

57.   Kim, P. *et al.* Perceived quality of maternal care in childhood and structure and function of
mothers' brain. *Dev Sci* **13**, 662–673 (2010).

58.   Champagne, F. A., Curley, J. P., Keverne, E. B. & Bateson, P. P. G. Natural variations in
postpartum maternal care in inbred and outbred mice. *Physiology and Behavior* **91**, 325–
334 (2007).

59.   Anstrom, K. K. & Woodward, D. J. Restraint Increases Dopaminergic Burst Firing in
Awake Rats. *Neuropsychopharmacol* **30**, 1832–1840 (2005).

60.   Heath, F. C. *et al.* Dopamine D1-like receptor signalling in the hippocampus and amygdala
modulates the acquisition of contextual fear conditioning. *Psychopharmacology (Berl)* **232**,
2619–2629 (2015).

61.   Espadas, I. *et al.* Dopamine D2R is Required for Hippocampal-dependent Memory and
Plasticity at the CA3-CA1 Synapse. *Cerebral Cortex* **31**, 2187–2204 (2021).

62.   Silva, M. R. P., Bernardi, M. M. & Felicio, L. F. Effects of dopamine receptor antagonists
on ongoing maternal behavior in rats. *Pharmacology Biochemistry and Behavior* **68**, 461–
468 (2001).

63.   Numan, M. *et al.* The effects of D1 or D2 dopamine receptor antagonism in the medial
preoptic area, ventral pallidum, or nucleus accumbens on the maternal retrieval response
and other aspects of maternal behavior in rats. *Behav Neurosci* **119**, 1588–1604 (2005).

64.   Sariñana, J., Kitamura, T., Künzler, P., Sultzman, L. & Tonegawa, S. Differential roles of
the dopamine 1-class receptors, D1R and D5R, in hippocampal dependent memory.
Proceedings of the National Academy of Sciences **111**, 8245–8250 (2014).

65.   Grace, A. A. Phasic versus tonic dopamine release and the modulation of dopamine system
responsivity: a hypothesis for the etiology of schizophrenia. *Neuroscience* **41**, 1–24 (1991).

66.   Kroener, S., Chandler, L. J., Phillips, P. E. M. & Seamans, J. K. Dopamine Modulates
Persistent Synaptic Activity and Enhances the Signal-to-Noise Ratio in the Prefrontal
Cortex. *PLoS One* **4**, e6507 (2009).

67.   Fadok, J. P., Dickerson, T. M. K. & Palmiter, R. D. Dopamine Is Necessary for Cue-
Dependent Fear Conditioning. *J. Neurosci.* **29**, 11089–11097 (2009).

68.   Wilmot, J. H. *et al.* Phasic locus coeruleus activity enhances trace fear conditioning by
increasing dopamine release in the hippocampus. *eLife* **12**, RP91465.

69.   Kimble, D. P., Rogers, L. & Hendrickson, C. W. Hippocampal lesions disrupt maternal, not
sexual, behavior in the albino rat. *Journal of Comparative and Physiological Psychology*
63, 401–407 (1967).

70.   Terlecki, L. J. & Sainsbury, R. S. Effects of fimbria lesions on maternal behavior in the rat.
Physiology & Behavior **21**, 89–97 (1978).

71.   Froemke, R. C. & Young, L. J. Oxytocin, Neural Plasticity, and Social Behavior. *Annu Rev*
Neurosci **44**, 359–381 (2021).

72.   Marlin, B. J., Mitre, M., D'amour, J. A., Chao, M. V. & Froemke, R. C. Oxytocin enables
maternal behaviour by balancing cortical inhibition. *Nature* **520**, 499–504 (2015).

73.   Orchard, E. R. *et al.* The maternal brain is more flexible and responsive at rest: effective
connectivity of the parental caregiving network in postpartum mothers. *Sci Rep* **13**, 4719
(2023).

74. Champagne, F. A. & Curley, J. P. Plasticity of the Maternal Brain Across the Lifespan:
Plasticity of the Maternal Brain Across the Lifespan. *New Directions for Child and*
Adolescent Development **2016**, 9–21 (2016).
75. Kempadoo, K. A., Mosharov, E. V., Choi, S. J., Sulzer, D. & Kandel, E. R. Dopamine
release from the locus coeruleus to the dorsal hippocampus promotes spatial learning and
memory. *Proceedings of the National Academy of Sciences* **113**, 14835–14840 (2016).

Supplementary Materials for

Persistent dopamine-dependent remodeling of the neural transcriptome in response to pregnancy and postpartum

Jennifer C Chan *et al.*

Corresponding author: Ian Maze, ian.maze@mssm.edu

MATERIALS AND METHODS

Animals

Wild-type C57BL6/J mice were purchased from Jackson Laboratories at 8-weeks old, and maintained on a 12-h/12-h light/dark cycle throughout the entirety of the experiments. Mice were provided with *ad libitum* access to water and food throughout the entirety of the experiments. All behavioral testing occurred during the animals' light cycle. Experimenters were blind to experimental group, and the order of testing was counterbalanced during behavioral experiments. All animal procedures were performed in accordance with NIH guidelines and with the approval of the Institutional Animal Care and Use Committee of the Icahn School of Medicine at Mount Sinai.

Breeding

Adult virgin female mice were pair bred in-house with age-matched males. Males were removed after a maximum of 5 days, and pregnant females were singly-housed at least 2 days before parturition. Pups were counted on the day of birth (0 dpp) and weaned at 21 dpp. Only dams with litters between 4-10 pups were used for all experiments. On the day of weaning, dams were group-housed into cages of 3-5 mice with animals of the same experimental condition. For timed breedings, copulation plugs were checked every morning within 1-hour after lights on, where confirmation of a plug was designated as E0.5, signaling the immediate removal of the female to

her own cage with a nestlet. Virgin NP females were age-matched for each experimental cohort.

Mating-experienced NP females were confirmed for the presence of a copulation plug. For

Pregnancy Only dams, pups were removed at 0 dpp within 3 hours after lights-on to minimize

maternal-offspring interactions. For comparison of PP dams to pup sensitized virgin females,

litters were culled to 4 pups to equate litter size with the number of pups used for each sensitized

female. In all other experiments, litters were not culled.

1013
Postpartum stress paradigm

Stress PP females were subjected to limited nesting and maternal separation from 10-20 dpp, as

previously described[1–3]. On each day of separation, the entire litter was removed to a clean cage

with Sani-Chip bedding for 3-4 hours. Separations occurred during the light cycle, and the timing

varied each day to minimize predictability and acclimatization. EnviroDri nesting material was

depleted to 1/3 of control cages during the days of separation. Following pup weaning on 21 dpp,

the nesting material was restored to normal levels, and dams were group-housed into cages of 3-5

with animals of the same experimental condition.

1022
Pup sensitization

Pup sensitization was conducted as previously described[4,5]. On the first day of pup sensitization,

virgin NP females were presented with four pups (postnatal day 5) from a cage consisting of a

lactating donor dam and her litter. Donor pups were exchanged for satiated pups from the same

litter every 8-12 hours for 21 days. Donor pups were weighed daily to ensure continual weight-

gain throughout the experiment, and only pups that exhibited consistent weight-gain were used.

Behavioral observations were conducted during the first four days of sensitization. Each dam was

observed for 30 minutes during the light phase for the following maternal behaviors:

licking/grooming, crouching, and nestbuilding. During these observation periods, females were

closely monitored to ensure that they did not display aggressive behavior toward the donor pups.

On the last day of sensitization, donor pups were weaned from the cage, and sensitized females

were group-housed into cages of 3-5 mice with animals of the same experimental condition.

1035
Brain tissue collections

Animals were sacrificed by rapid decapitation. Whole brains were flash frozen with cold 2-

methylbutane and stored at −80°C until further processing. Flash-frozen brains were sectioned at

−20°C using a 1 mm mouse coronal brain matrix (Stoelting Co.). Tissues enriched for the brain

region of interest were micropunched using a hollow needle (Ted Pella) according to the Allen

Brain Atlas (see **Fig. S1A**).

1042
1043 **Behavioral analyses**

1044 All animals (18-30 weeks, depending on the time from pup weaning) were handled for 2 minutes

1045 for two consecutive days prior to initial behavioral testing. Animals were habituated to the testing

1046 room for 1 hour prior to each behavioral assay. All testing occurred during the light phase.

1047
Pup retrieval: All animals were individually housed for 24-hours prior to testing. Following

habituation, 2 pups from a donor litter with a lactating dam (aged 4-6 dpp) were placed in different

corners opposite to the nest in the home cage. Pup-directed interactions were recorded from above

for 15-minutes or until both pups were successfully retrieved into the nest. Retrieval latency was

calculated as [time pup was first placed into the nest by animal - time first pup was placed into the

cage by experimenter].

1054

Contextual fear conditioning: On day 1, mice were habituated for 10 minutes to the testing

chamber, which consisted of a square plexiglass box with a metal grid inside a sound-attenuating

cabinet wiped with 70% ethanol (Med Associates, VT). On day 2, following a 3-minute baseline

measurement, mice were trained with five 2.0-second 0.7 mA foot shocks delivered with an

intertrial interval of 90-seconds. Testing for conditioned fear responses (freezing) for a total of 5-

minutes occurred 24-hours following training. Freezing was measured using ANY-maze software

connected to a camera positioned above the testing chamber. Freezing is expressed as a percentage

of the total test time or as a percentage of the 60-seconds prior to each shock during conditioning.

Animals with freezing levels exceeding 40% prior to the first shock were excluded to remove

potential confounding effects of heightened baseline responsivity that could interfere with

accurately assessing conditioned fear.

1066
1067   *Open Field:* Mice were placed in a 16x16 square arena under dim lighting for 5 minutes. A camera

1068   positioned overhead recorded the total distance and time spent in the center *vs.* periphery using

1069   Ethovision software.

1070
Object location task: For training, animals were allowed to freely explore two identical objects

placed equidistant from adjacent corners of a 16x16 cm square arena for 5 or 10 minutes, before

being returned to the home cage. Following a 1-hour delay, the animals were returned to the arena

for testing, during which time one object was moved to the opposing corner. During the test, the

animals were allowed to freely explore the objects for 5-minutes. A camera positioned overhead

recorded time spent exploring each object using Ethovision software. The discrimination score

was calculated as [(time spent with moved object) – (time spent with unmoved object)/[(time spent

with moved object) + (time spent with unmoved object)].

1079

1080 *Light-dark box:* Anxiety-like behavior was tested in an apparatus containing two interconnected

1081 20x20 cm compartments (Omnitech Electronics Inc.). One compartment was illuminated during

1082 the session ("light" side), while the other was covered by an opaque black perspex lid ("dark"

1083 side). The distance, time spent, and number of crossovers in each compartment were automatically

1084 recorded by Fusion software during the 10-minute test.

1085

1086 *Forced Swim Test:* Mice were placed in a 4 liter glass beaker with 2 liters of room temperature

1087 water for 8 minutes. Each session was recorded and scored by a blinded observer. The total number

1088 of seconds that mice were immobile during the last 5 minutes of the test were recorded, as

1089 previously described[6].

1090

Observation of pup-directed behaviors in the home cage: Confirmation of maternal behavior in

pup sensitized females was conducted based on prior studies[7]. Animals were observed from the

first to fourth day of pup sensitization, based on published work that maternal sensitization in

C57BL6/J females occurs following 4-days of pup exposures[8,9]. Observations occurred in the light

period in the first 30-minutes after pups were placed in the cage. Each animal was scored every 3-

minutes, with the observed behavior recorded as one or more of the following categories:

nestbuilding, grooming, sniffing, crouching, nursing, eating, drinking, self-grooming, no-contact.

PP dams were scored for the first 4-days following birth (0-3 dpp) concurrently for comparison.

1099
Estrous cycle testing

Vaginal samples were taken on each day of behavioral testing. 15 µL of sterile PBS was gently

pipetted into the vagina and mounted on a glass slide. Vaginal smears were stained with crystal

violet dye, washed twice with water, and cover slipped with glycerol. Three 20x images per sample

were acquired on a light microscope. Estrous stage was determined using the pretrained network

of EstrousNet[10], and confirmed afterwards by a trained experimenter based on cytology of

nucleated, cornified, or leukocytic cells.

1107
Viral Transduction

Mice were anesthetized with ketamine (100mg/kg) and xylazine (10 mg/kg) i.p. and positioned in

a stereotaxic frame (Kopf instruments). Given the widespread changes in dopamine receptor

expression across dHpc subregions, we did not restrict our viral manipulations to a specific

subregion. 2 µl of retrograde AAV.rTH.PI.Cre.SV40 (titer $\geq 7\times10^{12}$ vg/mL, Addgene #107788-

AAVrg) was infused bilaterally into the dHpc at 0.2 uL/min using the following coordinates: 7°

angle; anterior-posterior (AP) -2.2 mm, medial-lateral (ML) ±2.0 mm, dorsal-ventral (DV) -2.0

mm. 1 µl of pAAV-hSyn-DIO-mCherry (titer $\geq 4\times10^{12}$ vg/mL, Addgene #50459-AAV2) or

pAAV-hSyn-DIO-hM4D(Gi)-mCherry (titer $\geq 5\times10^{12}$ vg/mL; Addgene #44362-AAV2) was

bilaterally infused into the VTA using the following coordinates: 7° angle; AP -3.3 mm, ML ±0.9

mm; DV -4.6 mm. Needles remained in place for 7 minutes following injection to minimize virus

diffusion. Viral validations were conducted at least 21 days post-surgery to allow for optimal viral

expression and recovery.

1121
Chemogenetic manipulation

To directly manipulate dopamine signaling in dHpc, we selectively targeted the VTA-dHpc

projection during a window established as being critical for dopaminergic downregulation.

Following 7 days of recovery, surgerized female mice were randomly assigned to NP or PP groups.

PP females were pair bred with naïve male mice for a maximum of 5 days, and individually housed

prior to parturition. From 10-20 dpp, PP females were injected subcutaneously with

deschloroclozapine (DCZ, Tocris #7193) – to minimize off-target effects[11] - at 1ug/kg in 1%

DMSO or vehicle (saline). NP females were injected with DCZ concurrently. Following weaning

on 21 dpp, PP dams were group-housed with animals of the same experimental condition.

Behavioral testing occurred beginning at 28-days post-weaning (49 dpp), and brain tissues were

collected following contextual fear conditioning for further analyses.

1133
RNAscope in situ hybridization and analysis

Fresh frozen brains were cut into 12-16 μm thick slices in the coronal plane with a cryostat (Leica

CM3050-S), mounted on charged Superfrost Plus microscope slides (Thermofisher #P36934), and

stored at -80°C until processing. Sections were post-fixed with 4% PFA for 1.5 hours at 4°C, and

permeabolized with hydrogen peroxide (10 minutes RT) and Protease III (30 minutes RT). The

RNAscope Multiplex Fluorescent Reagent Kit v2 (ACD Bio) was used according to

manufacturer's instructions to sequentially stain sections with the following probes: Drd1a (Mm-

Drd1a-C1, #461901) and Drd2 (Mm-Drd2-C3, #406501-C3). RNAscope probes were visualized

using TSA Vivid Fluorophores (Tocris) at 1:750 dilution. Sections were counterstained with DAPI

(ACD Bio) and mounted using ProLong Gold Antifade Mountant (Thermo). Confocal images (3

images per animal, 1024 × 1024 pixels) were acquired on a Zeiss LSM 780 upright microscope

using a 40X objective with Zen Black software, with 5x1 tiled images. Images were averaged

across 8 consecutive acquisitions at a bit depth of 16 bits, with 2 z-stacks acquired per image.

Subcellular quantification of individual puncta per 100 μm nucleus - identified by cellular

detection of DAPI staining - was performed for each maximum intensity projected image using

QuPath software (v.0.5.1)[12]. Regions of interest (CA1 and dentate gyrus) were annotated and

determined by superimposing images onto the Allen Brain Atlas.

1151

Immunofluorescence and analysis

Mice were anesthetized with isoflurane and perfused with cold 1×PBS followed by 4% PFA. Brains were post-fixed in 4% PFA overnight and then transferred to a 30% sucrose/PBS solution for 2 days. Brains were sectioned at 40 μm thickness using a Leica CM3050-S cryostat, with serial sections collected from the VTA. For each subject, 2–3 brain slices were blocked for 2 hours (0.1% Triton X-100, 10% normal donkey serum), followed by overnight incubation at 4 °C with primary antibodies: chicken anti-tyrosine hydroxylase (1:500, Aves Labs #TYH) and rabbit anti-Fos (1:2000, Synaptic Systems #226-008). The next day, slices were incubated for 2 hours at RT with fluorescent-conjugated secondary antibodies (donkey anti-chicken Alexa Fluor 488, Invitrogen #A78948; donkey anti-goat Alexa Fluor 680, Invitrogen #A10043). Slices were counterstained with DAPI (1:10000, Thermo Scientific #62248) and mounted with ProLong Gold Antifade Mountant (Thermo Fisher #P36934). Confocal images (2-3 replicates per animal, 1024 x 1024 pixels) were acquired on a Zeiss LSM 780 upright microscope using a 40x objective with Zen Black software. Images were averaged across 8 consecutive acquisitions at a bit depth of 16 bits. Image analysis was conducted using FIJI software (NIH). The Fos and TH channels were thresholded using the MaxEntropy method to define regions of interest. Following identification of colocalized Fos+/TH+ signal, Fos intensity was measured and averaged from 4-6 images per animal. Brightness and contast were adjusted for representative images.

Dopamine ELISA

Brain tissue dopamine levels were assessed in response to 3 hours of pup separation in the home cage. Independent biological replicates were collected at each time point, with separate animals used for each measurement. Brains were rapidly harvested (see Brain Tissue Collection, above) at

baseline (prior to pup removal, 0 min), during pup removal (30 and 180 min), and 30 min after

pups were returned to the home cage (210 min). For assessment of NP females, samples were

collected in the home cage 30-min following removal of a littermate. Equal weights of

micropunched brain tissues were homogenized in lysis buffer (0.01 N HCl, 1 mM EDTA, 4 mM

sodium metabisulfite). Tissue dopamine levels were assessed using the Dopamine (Research)

ELISA Kit (ALPCO Diagnostics) according to manufacturer's instruction.

1182
HPA axis assessment

Plasma corticosterone levels were assessed in response to 3-hours of pup separation in the home

cage. Testing was initiated within 2-hours after lights on. Tail blood was collected prior to pup

removal (0 min), during pup removal (30 and 180 min), and 120 min after pups were returned to

the cage (300 min). Samples for control animals in the home cage, used to account for the effects

of handling-induced stress, were collected concurrently. Blood samples were immediately mixed

with 50 mM EDTA and centrifuged at 5000 rpm for 10 minutes. Plasma was collected and stored

at −80 °C until analysis. Corticosterone levels were quantified using a Corticosterone ELISA kit

(ENZO Life Sciences) according to manufacturer's instruction.

1192
Statistics

Statistical analyses for behavioral and immunoassay data were conducted using Prism software

(GraphPad, v.10.4.1). For experiments involving multiple conditions, one-way or two-way

ANOVAs were performed, followed by *post hoc* analyses when appropriate. For time course

analyses where multiple measurements were taken from the same animal, repeated measures

ANOVAs were performed. Two-tailed Student's t-tests were used for comparisons between two

conditions. Behavioral data derived from manual observations and pregnancy/litter outcomes were

analyzed using chi-square tests. Grubb's test (alpha = 0.05) was applied to detect outliers where

necessary. Statistical significance was defined as $p \leq 0.05$.

1202

Bulk RNA-seq and analysis

RNA isolation and library preparation: Total mRNA was extracted from frozen brain tissues after

homogenization in Trizol Reagent (Thermo Fisher) and cleaned using RNeasy Microcolumns

(Qiagen) following the manufacturer's instructions. For RNA-seq library preparation, 150 ng of

mRNA per sample was used with either the Illumina Stranded mRNA Prep Kit (Illumina,

#20040534) or the TruSeq RNA Library Prep Kit v2 (Illumina, #RS-122-2001), according to the

manufacturer's protocols. Library quality was assessed using a Qubit Fluorometer 2.0 (Thermo

Fisher) and a High Sensitivity D5000 TapeStation assay (Agilent) before sequencing on a NovaSeq

6000 or NovaSeq X system.

1212

Differential expression analysis: Raw fastq files, containing an average of 20–30 million reads per

sample, were processed for pseudoalignment and abundance quantification using Kallisto (v.

0.46.1) against the Ensembl Mus musculus reference (v. 79)[13]. To filter lowly-expressed genes,

only those with a total read count of at least 10 across all samples were retained. To account for

unwanted variation among samples within each sequencing experiment that could arise from

technical or biological factors unrelated to the conditions of interest (including litter size, estrous

stage, day of sample collection, etc.), RUVs (v1.32.0) was applied with a negative control gene set

derived from the total genes identified per sequencing experiment, after ensuring that unwanted

variation did not correlate with covariates of interest, as described previously[14,15]. Differential

expression analysis was performed using DESeq2 (v1.38.3)[16], with significant genes defined by

an adjusted p-value < 0.05. For brain-wide transcriptome comparisons in which samples were

processed across multiple sequencing runs and stemming from separate cohorts, pairwise

comparisons were performed independently for each brain region. For all other experiments, where

subjects came from the same cohort and were processed in a single sequencing run, all groups

were analyzed together to maintain consistent normalization within the experiment. Gene

expression time course analyses examining the periods before, during, and after pregnancy and

postpartum were performed on normalized count data using the ImpulseDE2 package (v0.99.10)[17]

for each brain region. Significant genes exhibiting transient regulation or monotonous changes in

expression were identified using case-only differential expression analysis, with a Q-value

threshold of 0.05.

1233
Weighted gene co-expression analysis: To identify brain-wide gene co-expression networks,

normalized count data for all brain regions were compiled and analyzed using the WGCNA

package (v1.73)[18]. Co-expression networks were constructed from the 7,500 most variable genes,

determined by ranking gene variance. A soft threshold power of 12 was identified with the

pickSoftThreshold function to ensure scale-free network properties. Modules were identified

based on dissimilarity of a signed topological overlap matrix (TOM), and named with an arbitrary

color. The "gray" module encompassed genes that did not segregate into any specific module, and

was therefore removed from further analyses. To assess the enrichment of differentially expressed

genes (DEGs) within each brain region per module, Fisher's exact tests were performed using the

fisher.test() function in R (v4.3.0). To analyze the correlation between gene modules and brain

regional sensitivity to parity, brain regions were categorized into two groups based on the top five

regions with significant DEG overlap. These regions were designated as "High Sensitivity"

regions, while all other regions were classified as "Low Sensitivity." Pearson correlation

coefficients were calculated to examine the relationship between each module and the trait

(regional sensitivity × parity status), with statistical significance determined by Student's t-tests (p

<0.05). Heatmaps were generated to visualize these module-trait correlations, and module-specific

gene lists were exported for pathway enrichment analysis.

1251

Pathway and predicted upstream regulator analyses: Functional annotation of DEGs was

conducted using ShinyGO (v0.81)[19], with all protein-coding genes in the mm10 genome used as

the background. Pathways with an FDR < 0.05 were considered significant. All significant

pathways and associated statistics are provided in the Supplementary Tables. For figures, relevant

pathways were selected from the top 10 or 1/3 of significant terms, ranked by FDR, to emphasize

processes consistent with hypotheses informed by published literature. For GO term selection,

Revigo[20] was used to reduce redundancy of overlapping GO terms when needed. Ingenuity

Pathway Analysis (IPA; Qiagen, Inc., v.01-23-01) was used to predict upstream regulators for

DEG lists[21]. For "High Sensitivity" and "Low Sensitivity" regions (see *WGCNA* section), DEGs

shared by at least two or three brain regions were extracted, irrespective of the direction of change.

The IPA software was used to identify upstream regulators associated with each DEG list, with

statistical significance defined as p < 0.05. To investigate the mechanisms underlying parity

programming of dHpc plasticity, significant upstream regulators were systematically prioritized

based on one or more of the following criteria: (1) multiple molecules involved in the same

signaling pathways (e.g., progesterone and its receptor, PGR); (2) structurally similar molecules

that engage analogous signaling cascades (e.g., levodopa and dopamine); (3) molecules known to

be influenced by reproductive exposures; (4) molecules demonstrated to play a role in hippocampal

plasticity; (5) molecules identified as significant in both "High Sensitivity" and "Low Sensitivity"

regulator lists; and (6) molecules with high statistical significance values. Following selection,

upstream regulators were categorized into general molecular classes (e.g., "Hormone,"

"Transcription Factor," "Lipid") and grouped for visualization. Data are presented as a bubble plot, with marker shape representing number of brain regions sharing DEGs and marker size corresponding to significance.

Gene expression overlap analyses: Jaccard indices were calculated for overlapping gene lists using the GeneOverlap package (v1.36.0)[22], with significance defined by $p < 0.05$. Fisher's exact tests were calculated using the base *fisher.test()* function in R (v. 4.3.0) following construction of contingency tables for each comparison. Transcriptome-wide, threshold-free gene expression overlap was visualized using Rank-Rank Hypergeometric Overlap (RRHO) heatmaps generated with the RRHO2 package (v1.0)[23]. Gene lists were ranked by signed p-values, calculated as the log10-transformed nominal p-value multiplied by the sign of the fold change, without applying differential expression thresholds.

Cell-type deconvolution: Brain cell-type proportion was estimated from normalized expression data using the BRETIGEA package (v.1.0.4)[24] with default markers. For comparisons between groups, surrogate proportion variables were normalized using the Normalize function in Prism (GraphPad, v. 10.4.1). Each sub column was normalized separately, with the smallest value set to 0% and the largest value set to 100%. To calculate $\log_2$(fold change) between groups, normalized values per cell-type per group were averaged and expressed as the log2 ratio of the group means.

Single nuclei RNA-seq

Nuclei isolation and library preparation: For each animal, 2 mm dHpc-enriched tissue micropunches were collected bilaterally from consecutive 1 mm slices from -0.80 to -2.80 mm relative to bregma for a total of 4 punches (Ted Pella). Samples were processed in batches of 4-6, with each group represented within each batch. Nuclei were isolated using a modified version of

a sucrose density gradient isolation protocol[25]. Briefly, thawed tissues were placed in 1 mL of lysis

buffer (0.32 M sucrose, 5 mM $CaCl_2$, 3 mM magnesium acetate, 0.1 mM EDTA, 10 mM Tris-HCl

pH 8, 1 mM DTT, 0.1% Triton X-100) with 50 μL of 25 U/mL RNase inhibitor (Takara #2313B)

in a dounce homogenizer (Wheaton #357538). Homogenization was performed with 20 strokes

using a tight pestle. Another 1 mL of lysis buffer was added, followed by an additional 10 strokes.

The resulting 2 mL homogenate was transferred to a 15 mL Open-Top Thinwall Polypropylene

Tube (Beckman-Coulter #361707). The homogenizer and pestle were rinsed with 2 mL of lysis

buffer, and this wash was combined with the homogenate for a total of 4 mL. The homogenate

was carefully underlaid with 9 mL of sucrose solution (1.8 M sucrose, 3 mM magnesium acetate,

1 mM DTT, 10 mM Tris-HCl pH 8) and ultracentrifuged at 24,000 rpm for 1 hour at 4 °C using a

Sorvall™ WX+ centrifuge. After centrifugation, the supernatant and debris at the interface were

gently removed. The nuclear pellet was resuspended in 1 mL of resuspension buffer (0.02% bovine

serum albumin in DPBS with 25 μL of 25 U/mL RNase inhibitor) and incubated on ice for 10

minutes. The suspension was passed through a 35 μm nylon mesh filter (Corning #352235) into a

1.5 mL RNase/DNase-free microcentrifuge tube and centrifuged at 2600 x g for 10 minutes at 4

°C. Supernatants were discarded, and nuclei were resuspended in 200 μL of resuspension buffer.

A 10 μL aliquot of the nuclei suspension was stained with Trypan Blue to assess quality and

concentration using a Countess 3 Automated Cell Counter. Nuclei suspensions were loaded onto

a Chromium Single Cell 3′ chip (10X Genomics, v3) and processed according to the manufacturer's

protocol, targeting 10,000 nuclei. Single-nuclei libraries were generated using the 10X Chromium

Next GEM Single Cell 3′ v3.1 (Dual Index) protocol (CG000315 Rev A). Libraries were pooled,

loaded onto a single 10B 100 Cycle Flowcell, and sequenced using an Illumina NovaSeq 6000

system to generate 25-30,000 paired-end 2 × 100 bp reads.

1320

Data analysis: FastQ files were processed with the 10X Genomics Cell Ranger pipeline (v7.1.0)

to demultiplex reads, align them to the mouse genome (mm10-2020-A), remove PCR duplicates,

and generate gene expression matrices. Cell Ranger filtered outputs were analyzed using Seurat

v4.3.0[26], and mitochondrial RNA content per cell was calculated using the GRCm39 (mm10)

genome annotation and regressed out using SCTransform normalization protocol included in the

Seurat toolkit with 20 principal components (PCs) and a resolution of 0.1. To estimate ambient

RNA and correct for background contamination, the SoupX (v1.6.2) package[27] was used for each

sample using raw and filtered feature matrices from the Cell Ranger output. Heterotypic doublets

were identified and removed using DoubletFinder (v2)[28] to ensure the integrity of singlet datasets.

Filtered singlet datasets were then re-normalized and integrated using the same Seurat

SCTransform v2 workflow mentioned above. Cell clusters were annotated using a combination of

expert curation based on published marker genes[29–35], and label transfer from hippocampal

reference datasets, including the Allen Brain Map and Broad Institute resources[34,36]. Clusters with

contaminant cell populations expressing markers for choroid plexus (*Ttr*), ependymal (*Tmem212*),

and vascular leptomeningeal cells (*Vtn, Col1a2*) were removed from the analysis[35,37–39].

Additionally, as the sequential 2mm micropunches encompassed portions of cortical, thalamic and

vHpc regions, clusters characterized by enrichment of published non-dHpc neuronal markers using

Seurat's FindMarkers function (layer 5/6 cortical: *Rorb, Foxp2*[39]; ventral granule neurons: *Tox3*[31])

were also removed from the analysis. Cell cluster proportion analyses were conducted using the

scProportionTest package[40], which employs a Monte Carlo permutation test to evaluate whether

observed differences result from random sampling. Proportional differences between conditions

were compared to a null distribution generated by resampling, and statistical significance was

determined by permutation-based p-values with confidence intervals estimated via bootstrapping.

Differential expression analysis was conducted using pseudobulk analysis, where gene counts

were summed across all cells within each sample for each cell type cluster using the

AggregateExpression() function. DESeq2 was then applied at the sample level to conduct

differential expression. To explore pathways underlying cluster-specific differences across

conditions, pathway analysis was conducted using ShinyGO on genes meeting the following

criteria: $log2FC > 1.5$ and $p < 0.05$.

1350
DATA AND MATERIALS AVAILABILITY

The RNA-seq data generated in this study have been deposited in the National Center for

Biotechnology Information Gene Expression Omnibus (GEO) database under accession numbers

GSE289991 and GSE289992. We declare that the data supporting findings for this study are

available within the article and Supplementary Information. Related data are available from the

corresponding author upon reasonable request. No restrictions on data availability apply.

1357

1358

Figure S1: Brain-wide transcriptomic analysis of cell markers and gene expression networks.
A) Brain regions selected for bulk transcriptional profiling: olfactory bulb (OB), medial prefrontal cortex (mPFC), nucleus accumbens (NAc), medial preoptic area (mPOA), paraventricular nucleus (PVN), dorsal hippocampus (dHpc), basolateral amygdala (BLA), ventral hippocampus (vHpc), ventral tegmental area (VTA), dorsal raphe nucleus (DRN), and locus coeruleus (LC). Yellow circles indicate area selected for tissue micropunching. **B-F)** Nonsignificant correlations between the fold change (NP *vs.* PP) in normalized surrogate proportion variables generated from cell-type deconvolution of bulk RNA-seq data for **B)** astrocytic, **C)** microglia, **D)** endothelial, **E)** mature oligodendrocyte, and **F)** oligodendrocyte precursor cell markers with the number of DEGs identified from each brain region. **G)** Scale independence plot for soft-threshold power selection for WGCNA analysis, illustrating the relationship between β and the scale-free topology fit index, identifying the optimal β value = 12. **H)** Mean connectivity plot illustrating the average network density for each evaluated soft-thresholding power. **I)** Dendrogram of gene clustering based on topological overlap, highlighting the hierarchical clustering process used for module detection and merging of similar modules. **J)** Dendrogram of eigengene network adjacency depicting similarity of modules. **K, L)** Significantly enriched **K)** GO Biological Processes and **L)** KEGG pathways, sorted by FDR values, from module gene sets. Bars are color-coded to indicate modules. The total number of significant terms for each analysis is provided.

1377

1378

1379
1380
Figure S2: Parity associated behavioral adaptations are not influenced by locomotion,
anxiety-/depressive-like behaviors, or estrous stage. A-D) In the open field test, there was no
change in total distance, time spent in the center, or latency to the center. While there was a
significant increase in time spent in corners for PP dams (Student's t-test, t(13) = 2.307, *p =
0.0382), this did not impact any other outcomes on the open field test. **E-G)** In the light-dark box,
there was no difference in time spent in lights *vs.* dark zones, latency to enter the light zone, or
transitions between zones. **H)** There was no difference in time spent immobile on the forced swim
test. **I-K)** Stratification of pup retrieval behaviors by estrous stage on the day of testing did not
reveal a significant effect of estrous stage. **L-N)** Similarly, stratification of contextual fear
conditioning outcomes did not identify a significant effect of estrous stage on acquisition or
context recall in NP or PP groups. Error bars represent mean ± SEM. N=6-11 animals/group.
1392

1393

1394
1395
Figure S3: Postpartum experiences modulate the extent of parity-associated dHpc
transcriptomic and behavioral adaptations. A) Pie charts showing significant shifts towards
pup-directed behaviors, representing maternal behavior initiation after four days of pup exposure
in Pup Sensitized virgin females ($\chi^2(7) = 33.52$, p = 2.11×10^{-5}). **B)** Principle components analysis
of NP *vs.* PP DEGs (adj p < 0.05) from dHpc, showing the Pregnancy Only group clusters most
closely with PP. **C)** Heatmaps of all significant DEGs comparing NP *vs.* Mating only, **D)** NP *vs.*
Pregnancy only, **E)** NP *vs.* Pup sensitized dHpc transcriptomes. **F)** Odds ratio analysis of DEG
overlap for all comparisons (*vs.* NP). Insert numbers indicate respective *p* values for each
association (N.S., *p* > 0.05). **G)** Pup separation increases maternal corticosterone levels over 3-

hours (RM two-way ANOVA, group: $F_{(2,15)} = 12.86$, $p = 0.0006$, time: $F_{(2.125, 31.87)} = 2.759$, $p = 0.0756$, interaction: $F_{(6,45)} = 3.294$, $p = 0.009$; Tukey's multiple comparisons test: Control *vs.* Stress PP, **$p < 0.01$). N = 6 animals/group. **H)** No effect of postpartum stress on litter size. N = 12/group. **I)** Postpartum stress did not alter maternal weights. **J)** All groups discriminated the novel location following 10-minutes of training on the object location task (one way ANOVA, $F_{(2,15)} = 0.06452$, $p = 0.9378$) N=6 animals/group. **K)** Stratification of object location test scores did not show a significant effect of estrous stage. **L)** There was no difference in the total distance travelled in the open field test across groups. Error bars represent mean ± SEM.

1414

1415
1416
Figure S4: snRNA-seq quality control assessments, cell marker gene expression, and
postpartum dopamine modulation. Violin plots showing the **(A)** number of detected genes, **(B)**
UMI number, and **(C)** mitochondrial gene % by group following regression for mitochondrial gene
mapping. **(D-F)** Breakdown of these metrics by annotated clusters show <10% mitochondrial gene
across all cell-types, except for a minority of neural stem cells. Unsupervised clustering of snRNA-
seq data with cells colored for **(G)** neuronal, excitatory, and inhibitory markers; **(H)** hippocampal
subregional markers; and **(I)** non-neuronal markers including for glial and neural progenitor
populations. **J)** Unsupervised clustering of snRNA-seq data with cells colored for *Drd1* (left) and
Drd2 (right) expression. **K)** Quantification of *Drd1* mRNA puncta in CA1 nuclei (one-way
ANOVA, (F(2,19204) = 3.963, p = 0.019). **L)** Quantification of *Drd2* mRNA puncta in CA1 nuclei
(one-way ANOVA, (F(2,19204) = 23.16, p < 0.0001). **M)** Quantification of *Drd1* mRNA puncta
in dentate gyrus nuclei (one-way ANOVA, (F(2,21864) = 42.09, p < 0.0001). **N)** Quantification
of *Drd2* mRNA puncta in dentate gyrus nuclei (one-way ANOVA, (2,21864) = 41.79, p < 0.0001).
Tukey's multiple comparisons test, *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. **O)**
Chronic postpartum stress resulted in higher baseline levels of dopamine in the nucleus accumbens
(NAc; Student's t-test, t(7) = 2.568, *p = 0.0371). N= 4-5 animals/group. **P)** Dopamine response
is not induced in NP dHpc 30-minutes following littermate separation (two-way ANOVA,
interaction of time x group (F(1,30) = 8.78, p = 0.0059, time (F(1,30) = 1.47, p = 0.2359, group,
(F(1,30) = 2.054, p = 0.1621); Tukey's multiple comparisons test, *p < 0.05). N = 6-11
animals/group. **Q)** Dopamine levels increased during pup separation in the VTA (left y-axis), but
not in the locus coeruleus (right y-axis), resembling patterns of NAc and dHpc dopamine dynamics
(two-way ANOVA, effect of brain region (F(1,46) = 32.32, p < 0.0001), time (F(3,46) = 2.311, p
= 0.0866), interaction (F(3,46) = 1.118, p = 0.1109); Dunnett's multiple comparisons test, *p <
0.05). N = 5-8 samples/group. Error bars represent mean ± SEM.
1441

Figure S5. Inhibition of VTA-dHpc projection mimics sustained changes in dopamine receptor dynamics. A) Representative images of c-Fos immunostaining in the VTA. Scale bars, 20 μm. **B)** Quantification of c-Fos immunoreactivity in VTA TH+ neurons (Student's t-test; t(4) = 3.299, *p = 0.03). N = 3 animals/group. **C)** Chronic DCZ injections did not alter weights at 21 dpp (two-way ANOVA; main effect of group (F(1,48) = 128, p < 0.0001), virus (F(1,48) = 0.0862, p = 0.7702); Tukey's multiple comparison's test, ****p < 0.0001). **D)** Parity induced significantly greater freezing on the context recall text in mCherry dams, with trending effects between hM4Di groups (two-way ANOVA (effect of group, (F(1,38) = 7.177, p = 0.0109, virus (F(1,38) = 1.631,

p = 0.2093, Fisher's LSD, *p £ 0.05). N = 9-12/group. **E)** Quantification of *Drd1* mRNA puncta
in CA1 nuclei (Student's t-test, ($t(4854) = 2.249$, *p = 0.00246). **F)** Quantification of *Drd2* mRNA
puncta in CA1 nuclei (Student's t-test, ($t(4854) = 1.502$, p = 0.1132). **G)** Quantification of *Drd1*
mRNA puncta in dentate gyrus nuclei (Student's t-test, ($t(11946) = 3.900$, ****p < 0.0001). **H)**
Quantification of *Drd2* mRNA puncta in dentate gyrus nuclei (Student's t-test, ($t(11946) = 6.034$,
****p < 0.0001). **I)** Schematic of working model.
1459

SUPPLEMENTAL REFERENCES

76. Rice, C. J., Sandman, C. A., Lenjavi, M. R. & Baram, T. Z. A novel mouse model for acute and long-lasting consequences of early life stress. *Endocrinology* **149**, 4892–4900 (2008).

77. Gilles, E. E., Schultz, L. & Baram, T. Z. Abnormal corticosterone regulation in an immature rat model of continuous chronic stress. *Pediatr Neurol* **15**, 114–119 (1996).

78. Peña, C. J. *et al.* Early life stress confers lifelong stress susceptibility in mice via ventral tegmental area OTX2. *Science* **356**, 1185–1188 (2017).

79. Fleming, A. S. & Rosenblatt, J. S. Maternal behavior in the virgin and lactating rat. *Journal of Comparative and Physiology Psychology* **86**, 957–972 (1974).

80. Krishnan, V. *et al.* Molecular Adaptations Underlying Susceptibility and Resistance to Social Defeat in Brain Reward Regions. *Cell* **131**, 391–404 (2007).

81. Carcea, I. *et al.* Oxytocin neurons enable social transmission of maternal behaviour. *Nature* **596**, 553–557 (2021).

82. Wolcott, N. S. *et al.* Automated classification of estrous stage in rodents using deep learning. *Sci Rep* **12**, 17685 (2022).

83. Nagai, Y. *et al.* Deschloroclozapine, a potent and selective chemogenetic actuator enables rapid neuronal and behavioral modulations in mice and monkeys. *Nat Neurosci* **23**, 1157–1167 (2020).

84. Secci, M. E., Reed, T., Quinlan, V., Gilpin, N. W. & Avegno, E. M. Quantitative Analysis of Gene Expression in RNAscope-processed Brain Tissue. *Bio Protoc* **13**, e4580 (2023).

85. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq quantification. *Nature biotechnology* **34**, 525–527 (2016).

86. Risso, D., Ngai, J., Speed, T. P. & Dudoit, S. Normalization of RNA-seq data using factor analysis of control genes or samples. *Nat Biotechnol* **32**, 896–902 (2014).

87. Peixoto, L. *et al.* How data analysis affects power, reproducibility and biological insight of RNA-seq studies in complex datasets. *Nucleic Acids Res* **43**, 7664–7674 (2015).

88. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

89. Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals and plants. *Bioinformatics* **36**, 2628–2629 (2020).

90. Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms. *PLOS ONE* **6**, e21800 (2011).

91. Krämer, A., Green, J., Pollard, J., Jr & Tugendreich, S. Causal analysis approaches in Ingenuity Pathway Analysis. *Bioinformatics* **30**, 523–530 (2014).

92. Shen, L. GeneOverlap: An R package to test and visualize gene overlaps.

93. Matevossian, A. & Akbarian, S. Neuronal Nuclei Isolation from Human Postmortem Brain Tissue. *J Vis Exp* 914 (2008) doi:10.3791/914.

94. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29 (2021).

95. Young, M. D. & Behjati, S. SoupX removes ambient RNA contamination from droplet-based single-cell RNA sequencing data. *Gigascience* **9**, giaa151 (2020).

96. McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell systems* **8**, 329-337.e4 (2019).

1504   97.   Artegiani, B. *et al.* A Single-Cell RNA Sequencing Study Reveals Cellular and Molecular
1505   Dynamics of the Hippocampal Neurogenic Niche. *Cell Reports* **21**, 3271–3284 (2017).
1506   98.   Zhou, Z., Zhi, C., Chen, D., Cai, Z. & Jiang, X. Single-nucleus RNA sequencing reveals
1507   cell type-specific transcriptome alterations of Down syndrome hippocampus using the
1508   Dp16 mouse model. *Genes Genomics* **45**, 1305–1315 (2023).
1509   99.   Cembrowski, M. S., Wang, L., Sugino, K., Shields, B. C. & Spruston, N. Hipposeq: a
1510   comprehensive RNA-seq database of gene expression in hippocampal principal neurons.
1511   *eLife* **5**, e14997.
1512   100.   Harris, K. D. *et al.* Classes and continua of hippocampal CA1 inhibitory neurons revealed
1513   by single-cell transcriptomics. *PLoS Biol* **16**, e2006387–e2006387 (2018).
1514   101.   Haslinger, A., Schwarz, T. J., Covic, M. & Lie, D. C. Expression of Sox11 in adult
1515   neurogenic niches suggests a stage-specific role in adult neurogenesis. *Eur J Neurosci* **29**,
1516   2103–2114 (2009).
1517   102.   Bandler, R. C. *et al.* Single-cell delineation of lineage and genetic identity in the mouse
1518   brain. *Nature* **601**, 404–409 (2022).
1519   103.   Olney, K. C. *et al.* Widespread choroid plexus contamination in sampling and profiling of
1520   brain tissue. *Mol Psychiatry* **27**, 1839–1847 (2022).
1521   104.   Xiang, M. *et al.* A Single-Cell Transcriptional Roadmap of the Mouse and Human Lymph
1522   Node Lymphatic Vasculature. *Front Cardiovasc Med* **7**, 52 (2020).
1523   105.   Tasic, B. *et al.* Adult Mouse Cortical Cell Taxonomy by Single Cell Transcriptomics. *Nat
1524   Neurosci* **19**, 335–346 (2016).
1525   106.   Miller, S. A. *et al.* LSD1 and aberrant DNA methylation mediate persistence of
1526   enteroendocrine progenitors that support BRAF mutant colorectal cancer. *Cancer Res* **81**,
1527   3791–3805 (2021).
1528
1529