<u>Dopaminergic Modulation of Short-Term Associative Memory in *Caenorhabditis elegans*</u>

Anna McMillen[1], Caitlin Minervini[1], Renee Green[1], Michaela E. Johnson[1], Radwan Ansaar[1], Yee

Lian Chew[1]

1 - Flinders Health and Medical Research Institute, College of Medicine and Public Health,

Flinders University, Bedford Park 5042, South Australia

* - to whom correspondence should be addressed: <u>yeelian.chew@flinders.edu.au</u>

<u>Abstract</u>

Forgetting, the inability to retrieve previously encoded memories, is an active process involving

neurotransmission, second messenger signalling, and cytoskeletal modifications. Forgetting is

thought to be essential to remove irrelevant memories and to increase the capacity to encode new

memories. Therefore, identifying key regulators of active forgetting is crucial to advance our

understanding of neuroplasticity. In this study, we utilised the compact and tractable

Caenorhabditis elegans model to investigate the role of the neurotransmitter dopamine in

forgetting. We conducted butanone associative learning assays based on an established protocol

and used mutant strains deficient in dopamine synthesis (tyrosine hydroxylase CAT-2 and

dopamine transporter DAT-1) and signalling (G protein-coupled receptors DOP-1, DOP-2, and

DOP-3) to assess the impact on learning and memory retention. Learning was measured

immediately post-training, and memory retention was evaluated every 0.5 hours up to 2 hours.

Our results show that animals lacking dopamine display a modest enhancement in learning

relative to wild-type, with the learned association persisting for at least 2 hours after training. We

also found that the D2-like receptors DOP-2 and DOP-3 function together to modulate the

forgetting process, with D1-like receptor DOP-1 having an antagonistic effect. Furthermore, re-

expression of CAT-2 tyrosine hydroxylase in ADE or CEP neurons showed that dopamine

synthesis in both neuron types is essential for short-term memory retention, albeit at different

time points after learning. These findings highlight the critical role of dopamine in forgetting,

consistent with findings in *Drosophila*, and suggest potential relevance for understanding

memory retention during healthy ageing and in conditions with dopamine imbalances such as

Parkinson's disease.

<u>Introduction</u>

The nematode *Caenorhabditis elegans* serves as an excellent model for studying the molecular

regulation of memory due to its genetic tractability, well-characterised nervous system, and

capacity to learn and remember in ways analogous to higher organisms (Ardiel & Rankin 2010).

In *C. elegans* and other animals, memory is categorised into short-term and long-term, based on

how it is encoded and its susceptibility to decay (Kandel *et al.* 2014). Conversely, forgetting

involves the inability to retrieve previously encoded memories. Research indicates that forgetting

is an active process requiring protein synthesis, and its regulation involves processes related to

neurotransmission, second messenger signalling, and cytoskeletal modifications (Stein &

Murphy 2014; Liu *et al.* 2022). Identifying the key modulators of active forgetting and

understanding their mechanisms, distinct from those involved in learning and short- or long-term

memory retention, is crucial to advance our knowledge of neuroplasticity.

Studies investigating the underlying mechanisms involved in memory regulation and forgetting

have reported diverse molecular pathways. Two major regulatory mechanisms have been

identified: 1) the TIR-1 and MAP kinase/JNK pathway, along with associated proteins

(membrane protein MACO-1 and receptor tyrosine kinase SCD-2 and its ligand HEN-1) play

roles in olfactory adaptation (Inoue *et al.* 2013; Kitazono *et al.* 2017; Arai *et al.* 2022), while 2)

the RNA-binding protein MSI-1 regulates forgetting by downregulating the Arp2/3 complex

(Hadziselimovic *et al.* 2014) in olfactory conditioning. Other actin regulatory proteins such as α-

adducin ADD-1 and Rho family G proteins are implicated in memory loss, albeit with RAC-2

acting independently of actin dynamics (Vukojevic *et al.* 2012; Bai *et al.* 2022).

Neurotransmitters such as serotonin and acetylcholine also regulate memory retention and

forgetting in *C. elegans* (Liu *et al.* 2022; Niu *et al.* 2022). In addition, dopamine plays a crucial

role in modulating short-term associative memory in *C. elegans* and other organisms (McMillen

& Chew 2023). Dopamine's role in forgetting is well-documented in studies of aversive olfactory

learning in *Drosophila melanogaster*: blocking dopaminergic neurons after learning enhances

memory retention, while stimulation decreases or eliminates it (Berry *et al.* 2012; Berry *et al.*

2015; Berry *et al.* 2018). To determine if dopamine's role in forgetting is conserved across

species, in this study, we utilised the compact and tractable *C. elegans* to dissect the effect of

dopamine on forgetting and to identify the neural circuits involved.

To do this, we conducted butanone associative learning assays according to (Kauffman *et al.*

2011) and tested mutants deficient in dopamine synthesis and signalling. Behavioural tests for

forgetting in *C. elegans* involve assessing learned behaviours at various time points post-

conditioning to observe when responses revert to baseline levels. Our data indicate that

dopamine-deficient animals exhibit a slight improvement in learning compared to wild-type, and

the memory of this learned association is retained for at least 2 hours post-training. Additionally,

we found that DOP-2 and DOP-3 D2-like receptors work together to modulate the forgetting of

this learned association. Furthermore, cell-specific re-expression of CAT-2 tyrosine hydroxylase

in dopaminergic neurons suggests that dopamine from both ADE and CEP neurons is required

for short-term memory retention, albeit at different times post-training. Taken together, our data

showing that dopamine signals are required for forgetting is consistent with findings in other

invertebrates (Berry *et al.* 2012). These findings highlight the critical role of dopamine in the

process of forgetting and suggest implications for understanding the pathological mechanisms

that contribute to cognitive deficits in neurodegenerative diseases such as Parkinson's disease.

Results

Worms lacking dopamine show extended short-term associative memory.

Dopamine regulates both non-associative and associative learning in *C. elegans* (McMillen &

Chew 2023). To determine if dopamine regulates forgetting, we used the olfactory associative

memory paradigm (**Figure 1A**) (Kauffman *et al.* 2011) to assess associative memory over time in

mutants deficient in dopamine signalling. Briefly, worms are conditioned with the odour

butanone (conditioned stimulus) and presence of food (unconditioned stimulus) for 1 hour on

solid media, before being placed onto 'hold' plates containing food but no butanone for 0.5 to 2

hours. Then, worm populations are assessed for odour preference in a two-choice chemotaxis

assay as in (Kauffman *et al.* 2011), where high chemotaxis indices (CIs) indicate strong butanone

attraction, and CIs closer to zero represent neutral butanone attraction. 'Naïve' worms do not

undergo conditioning and are placed onto chemotaxis plates immediately, while the 'Timepoint

0' (t=0) cohort are placed onto chemotaxis plates directly after conditioning. Conditioned worms

undergo a 1 hour 'pre-starvation' step to increase the value of the unconditioned stimulus (food)

(Kauffman *et al.* 2011) (**Figure 1Aii)**. To simplify the visualisation of behavioural assay data, we

only display comparisons between genotypes at each post-conditioning time point, and between

naïve and t=0 for wild-type and other genotypes to show if learning has occurred. Other

statistical comparisons are described in the text. After conditioning (t=0), wild-type worms show

an increased attraction to butanone (shown by a higher chemotaxis index, CI) compared with the

naïve cohort, indicating the capacity for associative olfactory learning. This memory decreases

over time, returning to baseline (naïve) levels after ~0.5 hours (comparison between CIs of naïve

and 0.5 h post-conditioning cohorts: p-value = 0.6754; one-way ANOVA with Dunnett's post-

test) (**Figure 1B)**.

We first tested the *cat-2* tyrosine hydroxylase mutant strain that lacks the enzyme required to

synthesise dopamine (Lints & Emmons 1999). These worms have substantially reduced

dopamine levels (Sanyal *et al.* 2004). Interestingly, *cat-2* mutant worms show a small but

significant improvement in learning, and an increased attraction to butanone for up to 2 hours

(comparison between CIs of *cat-2* naïve and 2 h post-conditioning cohorts: p-value <0.0001;

one-way ANOVA with Dunnett's post-test), suggesting they retain the memory of the learnt

association for a longer period compared with wild-type controls (**Figure 1B)**. As a

complementary method to compare the rate of memory loss (i.e., the rate of forgetting) between

genotypes, we performed linear regression analysis on the chemotaxis indices of conditioned

worms at post-conditioning time points (t= 0, 0.5, 1, 1.5, 2). We then compared the slopes of the

lines of best fit between the tested genotypes. This is shown for data from wild-type and *cat-2*

mutant worms in **Figure 1Bii**. Statistical comparisons from linear regression analysis of short-

term associative memory assays in **Figure 1** (and other Figures) are summarised in **Table 1**. We

found that the slopes for wild-type (-0.1575) and *cat-2* mutants (-0.0353) are significantly

different, with a p-value of 0.0049 (**Table 1**), consistent with *cat-2* worms showing a slower rate

of memory loss.

To determine if a milder reduction in dopamine has the same effect, we next tested *dat-1* mutant

worms with a mutation in the DAT-1 sodium-dependent dopamine transporter. These worms are

able to synthesise dopamine; however, dopamine reuptake from the synaptic cleft is impaired,

reducing the amount of residual dopamine in the nervous system (Formisano *et al.* 2020; Kindt *et*

al. 2007). *dat-1* mutant worms showed improved learning compared with wild-type controls

immediately after conditioning (t=0) and, unlike wild-type, retained memory of the learnt

behaviour at t=0.5 (comparison between naïve and t=0.5 cohorts: p-value <0.01 for *dat-1*, p-

value >0.05 for wild-type; one-way ANOVA with Dunnett's post-test). However, *dat-1* mutant

worms do not display an extended memory phenotype like *cat-2* mutants (comparisons between

naïve and time points after t=0.5: all p-values >0.05; one-way ANOVA with Dunnett's post-test)

(**Figure 1C**). In addition, linear regression analysis of the rate of memory loss for wild-type and

dat-1 showed that these slopes were not significantly different (**Table 1**). This suggests that

while excess dopamine in the synaptic cleft (as seen in *dat-1* mutants) leads to a memory

retention phenotype at 0.5 hours, newly synthesized dopamine is necessary for maintaining

memory retention up to 2 hours post-conditioning.

Finally, we tested if re-expression of CAT-2 in dopaminergic neurons (using the *dat-1* promoter)

rescues the extended memory phenotype in *cat-2* mutants. We found that transgenic worms re-

expressing CAT-2 phenocopied wild-type animals, with no improved learning compared with

wild-type at t=0, and memory returned to baseline levels from t=0.5 onwards (comparisons

between naïve and t=0.5: p-values >0.05 for both wild-type and CAT-2 rescue worms; one-way

ANOVA with Dunnett's post-test) (**Figure 1D**). Linear regression analysis of the rate of memory

loss for both strains showed that these slopes were not significantly different (**Table 1**). This

indicates that the enzyme required for dopamine synthesis, CAT-2, is required for forgetting and

acts in the dopaminergic neurons.

Figure 1: Worms lacking dopamine show extended short-term associative memory. A) (i) Schematic

of butanone associative learning and short-term memory assay, adapted from (Kauffman *et al.* 2011). Naïve

worms are placed directly onto chemotaxis assay plates, while conditioned worms are first starved for 1

hour in M9 buffer prior to conditioning on plates with food and butanone (streaked on lid). After

conditioning, worms are transferred to 'hold' plates with food (no odour) for 0.5–2 hours before being

placed onto chemotaxis plates. **(ii)** Layout of chemotaxis assay plate indicating where butanone (But),

ethanol (EtOH) and sodium azide are spotted. Worms are placed on the 'origin', as indicated. **B) (i)** Wild-

type and *cat-2(e1112)* mutant worms tested for butanone associative memory (n = 5 biological replicates).

(ii) Linear regression of mean CIs for each time point, compared between genotypes. Dashed line indicates

CI of 'naïve' cohort for both genotypes. **C)** Wild-type and *dat-1(ok157)* mutant worms tested for butanone

associative memory (n = 5 biological replicates). **D)** Wild-type and *cat-2* 'rescue' worms re-expressing

CAT-2 in the dopaminergic neurons tested for butanone associative memory (n = 4 biological replicates).

For butanone assays (**Bi, C, D**), each data point represents one biological replicate (which includes four

technical replicates). Each technical replicate consists of 50-250 worms. Behaviour was quantified using

chemotaxis indices (CI) for naïve/untrained worms and conditioned animals at 0, 0.5-, 1-, 1.5- and 2-hours

post-conditioning. Statistical analysis: a pairwise t-test was performed to compare genotypes for each time

point, and a one-way ANOVA with Tukey's multiple comparisons test was performed to compare post-

conditioning t=0 with naïve chemotaxis indices (CIs) between genotypes. Error bars = mean ± SD. p-values

are represented by **** ≤ 0.0001; *** ≤ 0.001, ** ≤ 0.01; * ≤ 0.05; ns = non-significant.

Dopamine receptors act synergistically to modulate short-term memory.

In *C. elegans*, dopamine receptors are categorised as ligand-gated ion channels (LGIC) or G

protein-coupled receptors (GPCR). Dopaminergic GPCRs are divided into D1-like and D2-like

receptors, which stimulate or inhibit adenylyl cyclase, respectively (Chase & Koelle 2007;

Girault & Greengard 2004; Mersha *et al.* 2013). DOP-1 is a D1-like receptor, while DOP-2 and

DOP-3 are D2-like receptors (Chase & Koelle 2007; Mersha *et al.* 2013) (**Figure 2A**). We first

tested *dop-1, dop-2* and *dop-3* single mutants to determine if they showed a memory retention

phenotype similar to *cat-2* mutant worms. We found that none of the single mutants showed

statistically significant learning or memory differences compared with wild-type controls

(**Figure 2B**): all mutants showed intact learning at t=0, and butanone preference had returned to

baseline (naïve) levels at t=0.5 (comparisons between naïve and t=0.5: p-values >0.05 for all

genotypes; one-way ANOVA with Dunnett's post-test). Linear regression analysis also showed

no statistically significant differences between slopes for single mutants compared with wild-

type controls for data in **Figure 2Bi, ii** and **iii** (**Table 1**). This indicates that dopamine does not

regulate forgetting by acting through DOP-1, DOP-2 or DOP-3 alone.

One possibility is that these receptors could be acting in a functionally redundant manner for

dopamine signalling to modulate forgetting in *C. elegans*, as in other behaviours such as bacterial

avoidance (Chou *et al.* 2022) or the escape response to olfactory cues (Baidya *et al.* 2014).

Therefore, we next tested a *dop-1;dop-2;dop-3* triple mutant strain and found that they displayed

similar memory compared with wild-type controls. Although their CIs were higher than wild-

type at 1,1.5 and 2 h post-conditioning, likely due to their increased preference for butanone in

naïve worms, comparisons between naïve and post-conditioning CIs for all time points within the

triple mutant strain were not statistically significant ($p > 0.05$, one-way ANOVA with Dunnett's

post-test) (**Figure 2C**). Moreover, linear regression analysis revealed that the slopes for wild-

type and *dop-1;dop-2;dop-3* triple mutants were not significantly different (**Table 1**). We

postulated that there could be an antagonistic effect between D1 and D2-like dopamine receptors

that could be masking the effects of individual receptors (Chase *et al.* 2004), and so we

subsequently tested double mutants for *dop-1;dop-2*, *dop-1;dop-3*, and *dop-2;dop-3* for their

capacity to retain memory over time (**Figure 2D**). As in other assays, we found that wild-type

controls returned to baseline (naïve) CI values by t=0.5 (or t=1 for **Figure 2Di**). This was also

the case for *dop-1;dop-2* (**Figure 2Di**) and *dop-1;dop-3* (**Figure 2Dii**) double mutants, which

returned to baseline CI levels by t=1 and t=0.5, respectively. In contrast, the *dop-2;dop-3* double

mutants showed elevated CIs post-conditioning that had not returned to baseline by t=2

(comparisons between naïve and t>0: p-values <0.01 for all time points; one-way ANOVA with

Dunnett's post-test) (**Figure 2Diii**). Linear regression analysis also showed that the slopes for

wild-type and *dop-2;dop-3* are close to significantly different (p-value =0.06) (**Table 1**). This

suggests that the D2-like receptors DOP-2 and DOP-3 act synergistically to modulate forgetting

of an olfactory associative memory, and that the D1-like receptor DOP-1 may antagonise this

effect.

Figure 2: Dopamine receptors act synergistically to modulate short-term memory. **A)** *C. elegans* dopamine receptors DOP-1, DOP-2, and DOP-3 are D1 or D2-like receptors (Chase & Koelle 2007;

Mersha *et al.* 2013). **B)** Butanone associative memory assay for wild-type worms and worms containing

single mutations in **(i)** *dop-1(vs100)* (*n* = 4 biological replicates)*,* **(ii)** *dop-2(vs105)* (*n* = 4 biological

replicates), and **(iii)** *dop-3(vs106)* (*n* = 4 biological replicates). **C)** Butanone associative memory assay for

wild-type and *dop-1;dop-2;dop-3* triple mutant worms (*n* = 4 biological replicates). **D)** Butanone

associative memory assay for wild-type and **(i)** *dop-1;dop-2* (*n* = 4 biological replicates)*,* **(ii)** *dop-1;dop-3*

(*n* = 3 biological replicates), and **(iii)** *dop-2;dop-3* (*n* = 4 biological replicates) double mutant worms. For

butanone assays (**B, C, D**), each data point represents one biological replicate (which includes four technical

replicates). Each technical replicate consists of 50-250 worms. Behaviour was quantified using chemotaxis

indices (CI) for naïve/untrained worms and conditioned animals at 0, 0.5-, 1-, 1.5- and 2-hours post-

conditioning. Statistical analysis: a pairwise t-test was performed to compare genotypes for each time point,

and a one-way ANOVA with Tukey's multiple comparisons test was performed to compare post-

conditioning t=0 with naïve chemotaxis indices (CIs) between genotypes. Error bars = mean ± SD. p-values

are represented by **** $\leq$ 0.0001; *** $\leq$ 0.001, ** $\leq$ 0.01; * $\leq$ 0.05; ns = non-significant.

Dopamine acts in dop-2/dop-3 expressing neurons to modulate short-term memory

Our data suggests that dopamine acts through D2-like dopamine GPCRs, DOP-2 and DOP-3, to

promote forgetting in *C. elegans*. To provide additional evidence for this, we re-expressed DOP-

2 and DOP-3 in *dop-2;dop-3* double mutants using a transgene that contains DOP-2 cDNA

(corresponding to DOP-2S in (Suo *et al.* 2009)) and DOP-3a cDNA, expressed under the control

of 5 kb of promoter sequence for both *dop-2* and *dop-3* (see Experimental Procedures for

details). Due to practical limitations on the number of groups that can be handled in a single

experiment, for these assays we tested only naïve and post-conditioning t=0, 0.5 and 2 h cohorts

for four genotypes: wild-type, *dop-2;dop-3* double mutants, and two versions of the transgenic

DOP-2/3 rescue line (**Figure 3A** and **B**). As the DOP-2/3 rescue transgenic lines contain

extrachromosomal arrays, there may be minor differences in transgene expression, so we tested

two versions to assess the role of DOP-2 and DOP-3 in forgetting. We found for data shown in

both **Figure 3A** and **3B** that, as expected, wild-type controls showed intact learning at t=0, as

indicated by higher CI compared with naïve worms, and that this had returned to baseline at t=2

(comparisons between naïve and t=0, 0.5, and 2 for all genotypes are indicated underneath each

graph). Consistent with **Figure 2Diii**, *dop-2;dop-3* mutants also displayed intact learning at t=0,

and the memory of this learnt association did not return to baseline levels at t=2 (comparison

between CIs for naïve and t=2: p-value <0.05; one-way ANOVA with Dunnett's post-test)

(**Figures 3A, 3B**), indicating an extended memory retention phenotype. We found that both

DOP-2/3 re-expression lines showed a partial rescue phenotype: transgenic worms showed intact

learning at t=0, and had returned to baseline levels at t=2 (comparison between CIs for naïve and

t=2: p-value >0.05; one-way ANOVA with Dunnett's post-test) (**Figures 3A, 3B**). Additionally,

linear regression analysis showed that slopes of wild-type and DOP-2/3 rescue lines (both

versions) are not significantly different (p-values 0.6889 and 0.4740 for **Figure 3A** and **3B**,

respectively), while slopes for wild-type vs *dop-2;dop-3* mutants are significantly different for

Figure 3A (p-value = 0.0188) and trend towards a difference for **Figure 3B** (p-value =0.1442)

(**Table 1**). Importantly, the slope for the DOP-2/3 rescue lines falls between wild-type and *dop-*

2;dop-3 mutants for both versions (**Figure 3A:** -0.2414 for wild-type, -0.1161 for *dop-2;dop-3*

double mutant, -0.2153 for DOP-2/3 rescue; and **Figure 3B:** -0.1792 for wild-type, -0.09202 for

dop-2;dop-3 double mutant, -0.1363 for DOP-2/3 rescue; **Table 1**). However, the CIs of DOP-

2/3 rescue worms were not significantly different from wild-type or *dop-2;dop-3* mutant controls

at t=2 (or in naïve worms), suggesting that their forgetting phenotype is intermediate between

wild-type and dopamine receptor mutants (**Figures 3A, 3B**). Taken together, these data indicate

that re-expression of DOP-2 and DOP-3 in a *dop-2;dop-3* double mutant is able to at least

partially rescue the forgetting phenotype observed in the double mutants.

Strain	Naïve vs t=0	Naïve vs t=0.5	Naïve vs t=2
Wild-type	0.0001	0.0116	0.8004
dop-2; dop-3	<0.0001	<0.0001	0.0352
dop-2; dop-3 rescue (v1)	0.0175	0.0882	0.8441

Strain	Naïve vs t=0	Naïve vs t=0.5	Naïve vs t=2
Wild-type	0.0221	0.1710	0.9999
dop-2; dop-3	<0.0001	0.0009	0.0288
dop-2; dop-3 rescue (v2)	0.0003	0.0028	0.9266

Figure 3: Dopamine acts through DOP-2 and DOP-3 to modulate short-term memory. A) Butanone associative memory assay for wild-type worms, *dop-2;dop-3* double mutant worms and *dop-2;dop-3* double mutant worms re-expressing DOP-2 and DOP-3 under the control of their endogenous promoters. DOP-2 in this transgene is a truncated version missing 5 amino acids when compared to the DOP-2S sequence (*n* = 5 biological replicates). **B)** Butanone associative memory assay for wild-type worms, *dop-2;dop-3* double mutant worms and *dop-2;dop-3* double worms re-expressing DOP-2 and DOP-3 under the control of their endogenous promoters. DOP-2 in this transgene exactly matches the published DOP-2S sequence (Suo *et al.* 2009) (*n* = 4 biological replicates). For rescue lines expressing extrachromosomal arrays, only CIs from transgenic worms are displayed here. Each data point represents one biological replicate (which includes four technical replicates). Each technical replicate consists of 50-250 worms. Behaviour was quantified using chemotaxis indices (CI) for naïve/untrained worms and conditioned animals at 0, 0.5-, and 2-hours post-conditioning. Statistical analysis: one-way ANOVA with Dunnett's multiple comparisons test was performed to compare post-conditioning time points with naïve chemotaxis indices (CIs) within a single genotype (shown in tables below each graph) and with Tukey's multiple comparisons test to compare CIs for naïve vs t=0 between genotypes. Error bars = mean ± SD. p-values are represented by **** ≤ 0.0001; *** ≤ 0.001, ** ≤ 0.01; * ≤ 0.05; ns = non-significant.

Dopamine release through ADE and CEP neurons promotes forgetting.

There are eight dopaminergic neurons in hermaphrodite *C. elegans*: two ADE neurons, four CEP neurons, and two PDE neurons (**Figure 4A**), each of which have specific synaptic and

neuromodulatory inputs/outputs to other parts of the nervous system. To investigate whether

specific connections are required for memory retention, we examined the role of individual

subsets of dopamine-releasing neurons in the process of forgetting. To do this, we assayed

transgenic lines in which CAT-2 is specifically re-expressed in the ADE (using an *exp-1*

promoter) or CEP neurons (using an *ace-1* promoter), in a *cat-2* mutant background (Chou *et al.*

2022). We first examined the role of ADE neurons in forgetting (**Figure 4B**): wild-type controls

displayed the expected phenotype of showing an increased CI (indicating intact learning) at t=0,

which had returned to baseline at t=0.5 (comparison between CIs for naïve and t=0.5: p-value

>0.05; one-way ANOVA with Dunnett's post-test). We found that ADE::CAT-2 rescue worms

showed intact learning at t=0, which had also returned to baseline at t=0.5 (comparison between

CIs for naïve and t=0.5: p-value >0.05; one-way ANOVA with Dunnett's post-test), suggesting

that expression of CAT-2 from the ADE neurons is sufficient to rescue the forgetting phenotype

observed in *cat-2* mutant worms (**Figure 1B**). However, we noticed that the CIs for the

ADE::CAT-2 rescue line at t=1.5 and t=2 post-conditioning were significantly higher than wild-

type controls, and the comparison between naïve and t=2 for this rescue line showed a trend

towards a significant difference (p-value 0.0971) (**Figure 4B**). Indeed, linear regression analysis

showed that slopes for wild-type and ADE::CAT-2 rescue worms are significantly different

(**Table 1**), suggesting incomplete rescue. Based on this data, we postulate that dopamine release

from ADE neurons may have a stronger effect on forgetting at the earlier (t=0.5,1) time points.

We next tested CEP::CAT-2 rescue worms (**Figure 4C**) in the butanone associative memory

assay. We found that wild-type worms showed the expected phenotype of intact learning at t=0

and return to baseline (naïve) CI at t=0.5. In contrast, CEP::CAT-2 worms showed significantly

higher CIs at t=0 (indicating intact learning), but this did not return to baseline until t=1.5,

suggesting a delayed forgetting phenotype (comparison between naïve and t=0.5 gives p-value of

0.0103; comparison between naïve and t=1 gives p-value of 0.0660 i.e. close to significance

threshold; one way ANOVA, Dunnett's post-test). However, the slope of the CEP::CAT-2 rescue

line (-0.2469) is similar to, and indeed is not significantly different from, that of wild-type

controls (-0.2518) (p-value = 0.9453; **Table 1**). These data suggest that dopamine release from

CEP neurons does not facilitate memory returning to naïve levels by t=0.5, but is sufficient to

promote a rate of memory loss that is similar to that of wild-type. Taken together with findings

from ADE::CAT-2 rescue lines, we propose that the CEPs regulate the rate of memory loss at

both early and late time points post-conditioning, but that this requires additional dopamine

release from other neurons such as the ADEs, which are critical for forgetting at early time

points.

Figure 4: **Dopamine release from ADE and CEP neurons promotes forgetting. A)** Dopaminergic neurons in *C. elegans* are the ADE, CEP (D/V = dorsal/ventral), and PDE neurons. **B)** Butanone associative memory assay for wild-type worms and *cat-2* mutant worms re-expressing CAT-2 in the ADE neurons (*n* = 4 biological replicates). **C)** Butanone associative memory assay for wild-type worms and *cat-2* mutant worms re-expressing CAT-2 in the CEP neurons (*n* = 4 biological replicates). Each data point represents one biological replicate (which includes four technical replicates). Each technical replicate consists of 50-250 worms. Behaviour was quantified using chemotaxis indices (CI) for naïve/untrained worms and conditioned animals at 0, 0.5-, 1-, 1.5- and 2-hours post-conditioning. Statistical analysis: a pairwise t-test was performed to compare genotypes for each time point, and a one-way ANOVA with Tukey's multiple

comparisons test was performed to compare post-conditioning t=0 with naïve chemotaxis indices (CIs)
between genotypes. Error bars = mean ± SD. p-values are represented by **** $\leq$ 0.0001; *** $\leq$ 0.001, **
$\leq$ 0.01; * $\leq$ 0.05; ns = non-significant. **D)** Schematic of potential mechanism, summarising data in **Figures**
1-4: our findings show that newly synthesised dopamine from both the ADEs and CEPs is required for
forgetting of olfactory associative memory, acting through DOP-2 and DOP-3 receptors.

Table 1: Linear regression analysis for short-term associative memory assays. Slopes from linear
regression are presented to 4 decimal places. *n* indicates the number of biological replicates. Statistical
comparisons are performed using a two-tailed test (GraphPad Prism). p-values (significance) indicated by
** $\leq$ 0.01; * $\leq$ 0.05; ns = non-significant.

Strain 1	Strain 2	Figure	*n*	Slope strain 1	Slope strain 2	p-value	Significance
Wild-type	*cat-2*	1B	5	-0.1575	-0.0353	0.0049	**
Wild-type	*dat-1*	1C	5	-0.2223	-0.2257	0.9512	ns
Wild-type	*cat-2;dat-1p::CAT-2*	1D	4	-0.2167	-0.2148	0.9775	ns
Wild-type	*dop-1*	2Bi	4	-0.3088	-0.2799	0.6808	ns
Wild-type	*dop-2*	2Bii	4	-0.2896	-0.2606	0.6897	ns
Wild-type	*dop-3*	2Biii	4	-0.3141	-0.3168	0.9628	ns
Wild-type	*dop-1, dop-2, dop-3*	2C	4	-0.2500	-0.2084	0.4982	ns
Wild-type	*dop-1, dop-2*	2Di	4	-0.1708	-0.2190	0.3154	ns
Wild-type	*dop-1, dop-3*	2Dii	3	-0.1726	-0.1386	0.5189	ns
Wild-type	*dop-2, dop-3*	2Diii	4	-0.1475	-0.05769	0.0661	ns (close to 0.05)
Wild-type	*dop-2;dop-3* rescue 1	3A	4	-0.2414	-0.2153	0.6889	ns
Wild-type	*dop-2; dop-3*	3A	4	-0.2414	-0.1161	0.0188	*
Wild-type	*dop-2;dop-3* rescue 2	3B	4	-0.1792	-0.1363	0.4740	ns
Wild-type	*dop-2; dop-3*	3B	4	-0.1792	-0.09202	0.1442	ns

Wild-type	*cat-2* ADE rescue	4B	4	-0.2278	-0.05045	0.0406	*
Wild-type	*cat-2* CEP rescue	4C	4	-0.2518	-0.2469	0.9453	ns

Discussion

Our study demonstrates that dopamine is essential for active forgetting in *C. elegans*.

Interestingly, the absence of a forgetting phenotype in the dopamine transporter DAT-1 mutant,

compared with the strong phenotype observed in CAT-2 tyrosine hydroxylase mutants, suggests

that the ability to synthesise and retain high levels of dopamine is necessary for this process.

Additionally, our findings indicate that D2-like receptors, specifically DOP-2 and DOP-3, are

required for forgetting. Finally, we observed that ADE and CEP dopaminergic neurons both

modulate forgetting, with ADE required at early time points, and CEP required at both early and

late time points after learning, highlighting the temporal complexity of this process (**Figure 4D**).

The role of dopamine in learning and memory is conserved across species. However, while

dopamine synthesis and signalling pathways are highly conserved, dopaminergic circuits

between *C. elegans* and other organisms differ. In *C. elegans*, eight dopaminergic neurons act

'extra-synaptically' on over 100 targets, mainly motor neurons. In mammals, dopaminergic

neurons are concentrated in the ventral midbrain (SNc and VTA) and project to the basal ganglia,

amygdala, and prefrontal cortex. In *Drosophila melanogaster*, small populations of dopaminergic

neurons are found throughout the fly brain, with projections innervating almost the entire brain,

with dense projections to the mushroom body lobes, which regulate olfactory learning

(Riemensperger *et al.* 2005; Marquis & Wilson 2022). Despite these differences, our finding that

dopamine is required for forgetting aligns with previous research on the role of dopamine in

forgetting in *Drosophila*. In flies, blocking synaptic output from dopaminergic neurons promotes

memory retention (i.e. reduces forgetting), whereas activating dopamine neurons decreases

memory (Berry *et al.* 2012). Indeed, dopamine is essential for both forgetting and memory

acquisition in flies, but this involves different receptors: DAMB for forgetting and dDA1 for

acquisition (Berry *et al.* 2012; Himmelreich *et al.* 2017). In our study, we observed a small but

robust learning improvement in *cat-2* and *dat-1* mutant worms, as well as in *dop-1;dop-2;dop-3*

triple mutants (double mutant combinations of these three receptors also showed a learning

improvement, albeit slightly above the alpha value of p = 0.05). This indicates that the

requirement for dopamine in learning and forgetting is conserved between flies and worms,

suggesting it is evolutionary important and likely also retained in higher organisms.

Our findings contrast with the work by Raj and Thekkuveettil (Raj & Thekkuveettil 2022), which

showed that dopamine deficient *cat-2* mutant *C. elegans* display both a learning defect and a

memory defect, i.e. they forget more quickly than wild-type controls. Although both our study

and the 2022 study use the butanone associative learning assay, worms in this research were

conditioned with agar plates seeded 1-2 days in advance with 100 µL OP50 (and thus already

dried on the agar), whereas the Raj and Thekkuveettil study involved conditioning in 400 µL of

OP50 pipetted onto unseeded agar plates immediately after worms are added. This means that the

worms are being conditioned in liquid, which may influence other locomotion pathways in which

dopamine plays an important role, such as in regulating "swimming" in liquid, a distinct form of

locomotion separate to crawling on solid media. Dopamine modulates swimming behaviour (Xu

et al. 2021), swimming-induced paralysis (McDonald *et al.* 2007), and swim-to-crawl transitions

when moving from liquid to solid media (Vidal-Gadea *et al.* 2011). As learning is a multifaceted

process modulated by multiple internal and external cues, it is not surprising that dopamine could

impact memory in different ways depending on the mode of conditioning, for example due to an

interaction between the response to the food and odour cues, and the swimming activity. This

underscores dopamine's multifaceted role in modulating behaviour and memory in *C. elegans*.

Our data show that both DOP-2 and DOP-3 D2-like receptors are required to modulate

forgetting. Single mutants of either receptor show memory retention capacity similar to wild-

type, while the *dop-2;dop-3* double mutant show enhanced memory retention similar to *cat-2*

mutant animals. Interestingly, the rate of memory loss in *dop-1;dop-2;dop-3* triple mutants

(indicated by the slope from linear regression) is similar between the triple mutant and wild-type

controls. This suggests that while DOP-2 and DOP-3 may promote memory retention (i.e. block

forgetting), the D1-like DOP-1 receptor may have an antagonistic effect in the absence of these

D2-like receptors. D1- and D2-like receptors are well-known to antagonise each other in signal

transduction, and their opposing effects in *C. elegans* have been previously reported in dietary

restriction-mediated lifespan extension (Jiang *et al.* 2022), swimming (Xu *et al.* 2021), and basal

slowing behaviour in response to a food lawn (Chase *et al.* 2004). We did not test the

invertebrate specific DOP-4 receptor or dopamine ligand-gated ion channels such as LGC-53;

therefore, a role for these receptors in dopamine-dependent short-term associative memory in *C.*

elegans remains unknown.

Short- and long-term memory differ by the mechanism for encoding and time taken to fully

decay. In *C. elegans*, short-term memory is typically considered to be relevant at 1-2 hours post-

learning, while long-term memory is evaluated 12-24 hours post-learning (Rahmani & Chew

2021; Kauffman *et al.* 2011). In our study, we tested short-term associative memory only up to 2

hours post-conditioning. At this time point, dopamine-deficient *cat-2* mutants, as well as *dop-*

2;dop-3 double mutants, still showed robust memory. Identifying when the memory decays to

baseline levels may provide further understanding of the role of dopamine in forgetting, such as

whether processes associated with long-term memory e.g. *de novo* protein synthesis (Fioriti *et al.*

2015), are required. In addition, we found that temporal regulation of dopamine release from

ADE neurons and CEP neurons may be required to modulate forgetting of olfactory memories.

The mechanistic details of this process, as well as the potential (untested) requirement of PDE

neurons, warrant further investigation in future studies.

In conclusion, our study leverages the worm's compact nervous system and genetic tractability to

provide new insights into the role of dopamine in forgetting. Other studies on olfactory

paradigms in worms have identified diverse regulators of forgetting, indicating that multiple

pathways and mechanisms are involved in this complex process. Our study adds to this body of

knowledge by highlighting the specific role of dopamine and its receptors. In addition, these

findings are consistent with research in *Drosophila*, suggesting a mechanism for forgetting that is

conserved in even more complex brains. Based on this study, future research could focus on

exploring the conserved mechanisms of forgetting across species and investigating potential

therapeutic targets for memory-related disorders.

<u>Experimental Procedures</u>

C. elegans *strain maintenance*

Young adult (day 1) hermaphrodite *C. elegans* were grown using standard conditions on

nematode growth medium (NGM) agar in petri dishes at 22 °C for all experiments (Brenner

1974). Worms were cultured at 22 °C for at least two generations prior to all learning assays. For

all experiments, animals were fed *Escherichia coli* (*E. coli*) strain OP50.

A list of strains and transgenic lines used, including full genotype information, is in **Table 2**.

Genotypes of mutant strains were confirmed via PCR, based on information in Wormbase

version WS295 (Sternberg *et al.* 2024). Transgenes were generated through standard

microinjection procedures as in (Chew *et al.* 2018a; Chew *et al.* 2018b; Gadenne *et al.* 2022).

Table 2: Genotype information for strains used in this study. For transgenic lines generated in this

study, numbers in square brackets refer to the concentration injected in ng/µL.

Strain name	Genotype	Source
Bristol N2	Wild-type	*Caenorhabditis* Genetics Centre (CGC)
CB1112	*cat-2(e1112)*	*Caenorhabditis* Genetics Centre (CGC)
RM2702	*dat-1(ok157)*	*Caenorhabditis* Genetics Centre (CGC)
LX645	*dop-1(vs100)*	*Caenorhabditis* Genetics Centre (CGC)
LX702	*dop-2(vs105)*	*Caenorhabditis* Genetics Centre (CGC)
LX703	*dop-3(vs106)*	*Caenorhabditis* Genetics Centre (CGC)
LX706	*dop-1(vs100); dop-2(vs105)*	*Caenorhabditis* Genetics Centre (CGC)
LX704	*dop-2(vs105); dop-3(vs106)*	*Caenorhabditis* Genetics Centre (CGC)
LX705	*dop-1(vs100); dop-3(vs106)*	*Caenorhabditis* Genetics Centre (CGC)
LX734	*dop-1(vs100); dop-2(vs105); dop-3(vs106)*	*Caenorhabditis* Genetics Centre (CGC)
YLC299	*cat-2(e1112); baIs4 [dat-1p::GFP + dat-1p::CAT-2]*	*cat-2(e1112)* crossed with UA57 from CGC
CLP1318	*cat-2(n4547);twnEx593[Pace-1::cat-2; Pgcy-8::mCherry]*	(Chou *et al.* 2022)
CLP1320	*cat-2(n4547);twnEx599[Pexp-1::cat-2; Pgcy-8::mCherry]*	(Chou *et al.* 2022)
YLC381	*dop-2(vs105)V; dop-3(vs106)X; pepEx040[Pdop-2(5kb)::dop-2S cDNA::let-858 UTR (pYLC102) [50];*	This study

	Pdop-3(5kb)::dop-3a cDNA::let-858 UTR (pYLC104) [50]; ccGFP [50]]	
YLC382	*dop-2(vs105)V; dop-3(vs106)X; pepEx043[Pdop-2(5kb)::dop-2S cDNA::let-858 UTR (pYLC103) [50]; Pdop-3(5kb)::dop-3a cDNA::let-858 UTR (pYLC104) [50]; ccGFP [50]]*	This study

Molecular biology

Transgenes were cloned using the NEB HiFi cloning (NEB #E2621). Promoters for the *dop-2*

and *dop-3* gene were both cloned 5 kb before the ATG, starting from

CAAGACGGACGCAGTCCG and CTGGAATATTCCAGAACTTTC, respectively. *dop-2* and

dop-3 cDNA were amplified from wild-type N2 RNA, extracted using the using the Monarch

Total RNA Kit (NEB #T2010S) and converted to cDNA using the Luna® Universal One-Step

RT-qPCR Kit (NEB #E3005S). Two forms of the *dop-2* cDNA sequence were cloned: one

aligned to the DOP-2S sequence published in (Suo *et al.* 2009) (pYLC103), the other is identical

to this sequence but lacks a AGTACGACGGCAAAT sequence from the 1385^{th} nucleotide

position, corresponding to amino acids STTAN (pYLC102). The *dop-3* cDNA sequence aligned

with the expected sequence for DOP-3, isoform a (T14E8.3a.1) (Sternberg *et al.* 2024).

Behavioural assays

The butanone appetitive learning assay (Kauffman *et al.* 2011; Rahmani *et al.* 2024) tests naïve

worms, conditioned worms that are tested directly after training (t=0), and conditioned worms

that go through a 'hold' period between training and testing (t=0.5–2 h). Briefly, *C. elegans*

populations were washed twice in M9 buffer (86 mM NaCl, 42 mM Na_2HPO_4, 22 mM K_3PO_4

(pH 6.0), 1 mM M_gSO_4) to constitute the 'naïve' group in butanone chemotaxis assays.

Remaining groups were then subjected to a 1-hour starvation period in M9 buffer on a shaker

(175 rpm at room temperature). Conditioned groups were then transferred onto NGM plates pre-

seeded with 100 µL *E. coli* OP50 and exposed to 10% (v/v) butanone (Sigma-Aldrich, #360473;

final concentration, diluted in 95% ethanol), streaked on plate lids, for 1 h. The conditioned

groups undergoing the 'hold' period were transferred onto NGM plates pre-seeded with 100 µL

E. coli OP50 for the assigned time periods. All groups were transferred onto salt-deficient agar in

6 cm petri dishes to perform butanone chemotaxis assays as described in Kauffman *et al.* (2010)

and Rahmani *et al.* (2024). Butanone chemotaxis assay plates were prepared with two equidistant

points both spotted with sodium azide, as well as 95% (v/v) ethanol on one position (control) and

10% (v/v) butanone diluted in 95% (v/v) ethanol on the other spot. Butanone chemotaxis

behaviour was quantified using the following equation:

$$\frac{\#\ worms\ on\ butanone - \#\ worms\ on\ ethanol}{\#\ total\ population - \#\ worms\ on\ origin}$$

The chemotaxis index (CI) ranges between 0 (neutral butanone attraction) to 1.0 (complete

butanone attraction). Each assay plate constituted one technical replicate; technical replicates

were generated from <50 total worms or >250 worms were excluded from the data. On very rare

occasions where negative CIs are observed, these data are also excluded as these populations

failed to show the standard naïve attraction to butanone. Each biological replicate constituted 3-4

technical replicates. At least three biological replicates were performed for each genotype tested,

the biological replicates are stated for each experiment in the appropriate figure legend.

Statistical analysis

Statistical analysis for all experiments was performed using GraphPad Prism version 8.0. In

general, where two groups were compared, a paired t-test was used. A one-way ANOVA with

Dunnett's multiple comparisons test was performed to compare post-conditioning time points

with naïve chemotaxis indices (CIs) within a single genotype, or with Tukey's multiple

comparisons test to compare CIs between genotypes for naïve and t=0. The alpha value for all

analyses is set at 0.05. Simple linear regression was performed to calculate the slope of the

change in chemotaxis index over time for conditioned worms, and to perform a two-tailed test to

determine if the slopes are significantly different between genotypes tested.

<u>Acknowledgements</u>

Many thanks go to our team members in the Worm Neuroscience lab (Flinders) for thoughtful

discussions, and to our colleagues at Flinders University – Dr Nicholas Eyre, Professor Damien

Keating, and Professor Kim Hemsley – for feedback and scientific advice. We especially thank

Pasquale Vitaro (Summer Scholar 2025) for technical assistance with behavioural assays, and

Aelon Rahmani for assistance with genotyping. We are grateful to Prof Chun-Liang Pan

(National Taiwan University School of Medicine) for strains CLP1318 and CLP1320 (cell-

specific rescue of *cat-2*), and Prof Satoshi Suo (Saitama Medical University, Japan) for plasmids

encoding *dop-2* cDNA. Finally, we gratefully acknowledge the *Caenorhabditis* Genetics Centre,

which is supported by the National Institutes of Health (P40 OD010440), for providing some of

the strains used in this study (see Table 2).

Funding: A.M is funded by a Flinders University Research Scholarship (Flinders University).

Y.L.C is funded by the National Health and Medical Research Council (NHMRC)

(GNT1173448), the Flinders University Parental Leave Research Support Scheme, the Flinders

University Impact Seed Funding Grant for Early Career Researchers, and the Flinders

Foundation Mary Overton Senior Research Fellowship.

Author contributions: Conceptualisation/Methodology– **AM, YLC**. Data curation/ Formal

analysis/ Investigation/ Resources – **AM, CM, MEJ, RG, RA, YLC.** Project administration/

Supervision/ Funding acquisition – **YLC.** Visualisation/ Writing – original draft - **AM, YLC**

Writing – review & editing – **AM, MEJ, YLC**

Conflict of interest: The authors declare that they have no conflicts of interest.

Data Availability Section

This study includes no data deposited in external repositories. Reagents including *C. elegans*

strains and expression plasmids are available from the corresponding author upon request.

References

Arai, M., Kurokawa, I., Arakane, H., Kitazono, T. and Ishihara, T. (2022) Regulation of
Diacylglycerol Content in Olfactory Neurons Determines Forgetting or Retrieval of
Olfactory Memory in Caenorhabditis elegans. *J Neurosci* **42,** 8039-8053.
Ardiel, E. L. and Rankin, C. H. (2010) An elegant mind: learning and memory in Caenorhabditis
elegans. *Learn Mem* **17,** 191-201.
Bai, H., Huang, H., Zhao, N., Gu, H., Li, Y., Zou, W., Wu, T. and Huang, X. (2022) Small G
protein RAC-2 regulates forgetting via the JNK-1 signalling pathway in Caenorhabditis
elegans. *Eur J Neurosci* **56,** 6162-6173.
Baidya, M., Genovez, M., Torres, M. and Chao, M. Y. (2014) Dopamine modulation of
avoidance behavior in Caenorhabditis elegans requires the NMDA receptor NMR-1.
PLoS One **9,** e102958.
Berry, J. A., Cervantes-Sandoval, I., Chakraborty, M. and Davis, R. L. (2015) Sleep Facilitates
Memory by Blocking Dopamine Neuron-Mediated Forgetting. *Cell* **161,** 1656-1667.
Berry, J. A., Cervantes-Sandoval, I., Nicholas, E. P. and Davis, R. L. (2012) Dopamine is
required for learning and forgetting in Drosophila. *Neuron* **74,** 530-542.
Berry, J. A., Phan, A. and Davis, R. L. (2018) Dopamine Neurons Mediate Learning and
Forgetting through Bidirectional Modulation of a Memory Trace. *Cell Rep* **25,** 651-662
e655.
Brenner, S. (1974) The genetics of Caenorhabditis elegans. *Genetics* **77,** 71-94.

Chase, D. L. and Koelle, M. R. (2007) Biogenic amine neurotransmitters in C. elegans. *WormBook: The Online Review of C. elegans Biology [Internet]*.

Chase, D. L., Pepper, J. S. and Koelle, M. R. (2004) Mechanism of extrasynaptic dopamine signaling in Caenorhabditis elegans. *Nat Neurosci* **7,** 1096-1103.

Chew, Y. L., Grundy, L. J., Brown, A. E. X., Beets, I. and Schafer, W. R. (2018a) Neuropeptides encoded by *nlp-49* modulate locomotion, arousal and egg-laying behaviours in *Caenorhabditis elegans* via the receptor SEB-3. *Philos Trans R Soc Lond B Biol Sci* **373**.

Chew, Y. L., Tanizawa, Y., Cho, Y. et al. (2018b) An Afferent Neuropeptide System Transmits Mechanosensory Signals Triggering Sensitization and Arousal in C. elegans. *Neuron* **99,** 1233-1246 e1236.

Chou, S. H., Chen, Y. J., Liao, C. P. and Pan, C. L. (2022) A role for dopamine in C. elegans avoidance behavior induced by mitochondrial stress. *Neurosci Res* **178,** 87-92.

Fioriti, L., Myers, C., Huang, Y. Y. et al. (2015) The Persistence of Hippocampal-Based Memory Requires Protein Synthesis Mediated by the Prion-like Protein CPEB3. *Neuron* **86,** 1433-1448.

Formisano, R., Mersha, M. D., Caplan, J., Singh, A., Rankin, C. H., Tavernarakis, N. and Dhillon, H. S. (2020) Synaptic vesicle fusion is modulated through feedback inhibition by dopamine auto-receptors. *Synapse* **74,** e22131.

Gadenne, M. J., Hardege, I., Yemini, E., Suleski, D., Jaggers, P., Beets, I., Schafer, W. R. and Chew, Y. L. (2022) Neuropeptide signalling shapes feeding and reproductive behaviours in male Caenorhabditis elegans. *Life Science Alliance* **5,** e202201420.

Girault, J.-A. and Greengard, P. (2004) The neurobiology of dopamine signaling. *Archives of neurology* **61,** 641-644.

Hadziselimovic, N., Vukojevic, V., Peter, F. et al. (2014) Forgetting is regulated via Musashi-mediated translational control of the Arp2/3 complex. *Cell* **156,** 1153-1166.

Himmelreich, S., Masuho, I., Berry, J. A., MacMullen, C., Skamangas, N. K., Martemyanov, K. A. and Davis, R. L. (2017) Dopamine Receptor DAMB Signals via Gq to Mediate Forgetting in Drosophila. *Cell Rep* **21,** 2074-2081.

Inoue, A., Sawatari, E., Hisamoto, N., Kitazono, T., Teramoto, T., Fujiwara, M., Matsumoto, K. and Ishihara, T. (2013) Forgetting in C. elegans is accelerated by neuronal communication via the TIR-1/JNK-1 pathway. *Cell Rep* **3,** 808-819.

Jiang, Y., Gaur, U., Cao, Z., Hou, S. T. and Zheng, W. (2022) Dopamine D1- and D2-like receptors oppositely regulate lifespan via a dietary restriction mechanism in Caenorhabditis elegans. *BMC Biol* **20,** 71.

Kandel, E. R., Dudai, Y. and Mayford, M. R. (2014) The molecular and systems biology of memory. *Cell* **157,** 163-186.

Kauffman, A., Parsons, L., Stein, G., Wills, A., Kaletsky, R. and Murphy, C. (2011) C. elegans positive butanone learning, short-term, and long-term associative memory assays. *J Vis Exp*.

Kauffman, A. L., Ashraf, J. M., Corces-Zimmerman, M. R., Landis, J. N. and Murphy, C. T. (2010) Insulin signaling and dietary restriction differentially influence the decline of learning and memory with age. *PLoS Biol* **8,** e1000372.

Kindt, K. S., Quast, K. B., Giles, A. C., De, S., Hendrey, D., Nicastro, I., Rankin, C. H. and Schafer, W. R. (2007) Dopamine mediates context-dependent modulation of sensory plasticity in C. elegans. *Neuron* **55,** 662-676.

Kitazono, T., Hara-Kuge, S., Matsuda, O., Inoue, A., Fujiwara, M. and Ishihara, T. (2017) Multiple Signaling Pathways Coordinately Regulate Forgetting of Olfactory Adaptation through Control of Sensory Responses in Caenorhabditis elegans. *J Neurosci* **37,** 10240-10251.

Lints, R. and Emmons, S. W. (1999) Patterning of dopaminergic neurotransmitter identity among Caenorhabditis elegans ray sensory neurons by a TGFβfamily signaling pathway and a Hox gene. *Development* **126,** 5819-5831.

Liu, H., Wu, T., Canales, X. G., Wu, M., Choi, M. K., Duan, F., Calarco, J. A. and Zhang, Y. (2022) Forgetting generates a novel state that is reactivatable. *Sci Adv* **8,** eabi9071.

Marquis, M. and Wilson, R. I. (2022) Locomotor and olfactory responses in dopamine neurons of the Drosophila superior-lateral brain. *Curr Biol* **32,** 5406-5414 e5405.

McDonald, P. W., Hardie, S. L., Jessen, T. N., Carvelli, L., Matthies, D. S. and Blakely, R. D. (2007) Vigorous motor activity in Caenorhabditis elegans requires efficient clearance of dopamine mediated by synaptic localization of the dopamine transporter DAT-1. *J Neurosci* **27,** 14216-14227.

McMillen, A. and Chew, Yee L. (2023) Neural mechanisms of dopamine function in learning and memory in Caenorhabditis elegans. *Neuronal Signaling* **8,** NS20230057.

Mersha, M., Formisano, R., McDonald, R., Pandey, P., Tavernarakis, N. and Harbinder, S. (2013) GPA-14, a Gαi subunit mediates dopaminergic behavioral plasticity in C. elegans. *Behavioral and Brain Functions* **9,** 1-8.

Niu, S., Zhou, W., Li, Y. and Huang, X. (2022) The signaling pathway of levamisole-sensitive-acetylcholine receptors involved in short-term forgetting of Caenorhabditis elegans. *Mol Genet Genomics* **297,** 1027-1038.

Rahmani, A. and Chew, Y. L. (2021) Investigating the molecular mechanisms of learning and memory using Caenorhabditis elegans. *J Neurochem* **159,** 417-451.

Rahmani, A., McMillen, A., Allen, E., Minervini, C. and Chew, Y. L. (2024) Behavioral Tests for Associative Learning in Caenorhabditis elegans. In: *Neurobiology: Methods and Protocols*, (S. Dworkin ed.)*, pp. 21-46. Springer US, New York, NY.

Raj, V. and Thekkuveettil, A. (2022) Dopamine plays a critical role in the olfactory adaptive learning pathway in Caenorhabditis elegans. *J Neurosci Res* **100,** 2028-2043.

Riemensperger, T., Voller, T., Stock, P., Buchner, E. and Fiala, A. (2005) Punishment prediction by dopaminergic neurons in Drosophila. *Curr Biol* **15,** 1953-1960.

Sanyal, S., Wintle, R. F., Kindt, K. S. et al. (2004) Dopamine modulates the plasticity of mechanosensory responses in Caenorhabditis elegans. *The EMBO journal* **23,** 473-482.

Stein, G. M. and Murphy, C. T. (2014) C. elegans positive olfactory associative memory is a molecularly conserved behavioral paradigm. *Neurobiol Learn Mem* **115,** 86-94.

Sternberg, P. W., Van Auken, K., Wang, Q. et al. (2024) WormBase 2024: status and transitioning to Alliance infrastructure. *Genetics* **227**.

Suo, S., Culotti, J. G. and Van Tol, H. H. (2009) Dopamine counteracts octopamine signalling in a neural circuit mediating food response in C. elegans. *EMBO J* **28,** 2437-2448.

Vidal-Gadea, A., Topper, S., Young, L. et al. (2011) Caenorhabditis elegans selects distinct crawling and swimming gaits via dopamine and serotonin. *Proc Natl Acad Sci U S A* **108,** 17504-17509.

Vukojevic, V., Gschwind, L., Vogler, C., Demougin, P., de Quervain, D. J., Papassotiropoulos, A. and Stetak, A. (2012) A role for alpha-adducin (ADD-1) in nematode and human memory. *EMBO J* **31,** 1453-1466.

605   Xu, Y., Zhang, L., Liu, Y. et al. (2021) Dopamine receptor DOP-1 engages a sleep pathway to
606   modulate swimming in C. elegans. *iScience* **24,** 102247.