Dopaminergic Modulation of Short-Term Associative Memory in *Caenorhabditis elegans*

Anna McMillen[1], Caitlin Minervini[1], Renee Green[1], Michaela E. Johnson[1], Radwan Ansaar[1], Yee

Lian Chew[1]

1 - Flinders Health and Medical Research Institute, College of Medicine and Public Health,

Flinders University, Bedford Park 5042, South Australia

* - to whom correspondence should be addressed: yeelian.chew@flinders.edu.au

Abstract

Forgetting, the inability to retrieve previously encoded memories, is an active process involving

neurotransmission, second messenger signalling, and cytoskeletal modifications. Forgetting is

thought to be essential to remove irrelevant memories and to increase the capacity to encode new

memories. Therefore, identifying key regulators of active forgetting is crucial to advance our

understanding of neuroplasticity. In this study, we utilised the compact and tractable

Caenorhabditis elegans model to investigate the role of the neurotransmitter dopamine in

forgetting. We conducted butanone associative learning assays based on an established protocol

and used mutant strains deficient in dopamine synthesis (tyrosine hydroxylase CAT-2 and

dopamine transporter DAT-1) and signalling (G protein-coupled receptors DOP-1, DOP-2, and

DOP-3) to assess the impact on learning and memory retention. Learning was measured

immediately post-training, and memory retention was evaluated every 0.5 hours up to 2 hours.

Our results show that animals lacking dopamine display a modest enhancement in learning

relative to wild-type, with the learned association persisting for at least 2 hours after training. We

also found that D2-like receptors DOP-2 and DOP-3 act together to modulate the forgetting

process, with D1-like receptor DOP-1 functioning redundantly. Furthermore, re-expression of

CAT-2 tyrosine hydroxylase in ADE and/or CEP neurons was unable to rescue the memory

retention phenotype observed in *cat-2* mutants, suggesting that dopamine release from all

dopaminergic neurons is required to modulate forgetting. These findings highlight the critical

role of dopamine in forgetting, consistent with findings in *Drosophila*, and suggest potential

relevance for understanding memory retention during healthy ageing and in conditions with

dopamine imbalances such as Parkinson's disease.

List of abbreviations

C. elegans: Caenorhabditis elegans

CI: Chemotaxis Index

Escherichia coli: E. coli

LI: Learning index

PCR: Polymerase Chain Reaction

Introduction

The nematode *Caenorhabditis elegans* serves as an excellent model for studying the molecular

regulation of memory due to its genetic tractability, well-characterised nervous system, and

capacity to learn and remember in ways analogous to higher organisms (Ardiel & Rankin 2010).

In *C. elegans* and other animals, memory is categorised into short-term and long-term, primarily

based on functional gene conservation with analogous processes as defined other organisms

(Kandel *et al.* 2014; Brandel-Ankrapp & Arey 2023; Rahmani & Chew 2021; Giles *et al.* 2006).

On the other hand, forgetting involves the inability to retrieve previously encoded memories.

Research indicates that forgetting is an active process requiring protein synthesis, and its

regulation involves processes related to neurotransmission, second messenger signalling, and

cytoskeletal modifications (Stein & Murphy 2014; Liu *et al.* 2022). Identifying the key

modulators of active forgetting and understanding their mechanisms, distinct from those

involved in learning and short- or long-term memory retention, is crucial to advance our

knowledge of neuroplasticity.

Studies investigating the underlying mechanisms involved in memory regulation and forgetting

have reported diverse molecular pathways. Two major regulatory mechanisms have been

identified: 1) the TIR-1 and MAP kinase/JNK pathway, along with associated proteins

(membrane protein MACO-1 and receptor tyrosine kinase SCD-2 and its ligand HEN-1) play

roles in olfactory adaptation (Inoue *et al.* 2013; Kitazono *et al.* 2017; Arai *et al.* 2022), while 2)

the RNA-binding protein MSI-1 regulates forgetting by downregulating the Arp2/3 complex

(Hadziselimovic *et al.* 2014) in olfactory conditioning. Other actin regulatory proteins such as α-

adducin ADD-1 and Rho family G proteins are implicated in memory loss, albeit with RAC-2

acting independently of actin dynamics (Vukojevic *et al.* 2012; Bai *et al.* 2022).

Neurotransmitters such as serotonin and acetylcholine also regulate memory retention and

forgetting in *C. elegans* (Liu *et al.* 2022; Niu *et al.* 2022). In addition, dopamine plays a crucial

role in modulating short-term associative memory in *C. elegans* and other organisms (McMillen

& Chew 2023). Dopamine's role in forgetting is well-documented in studies of aversive olfactory

learning in *Drosophila melanogaster*: blocking dopaminergic neurons after learning enhances

memory retention, while stimulation decreases or eliminates it (Berry *et al.* 2012; Berry *et al.*

2015; Berry *et al.* 2018). To determine if dopamine's role in forgetting is conserved across

species, in this study, we utilised the compact and tractable *C. elegans* to dissect the effect of

dopamine on forgetting and to identify the neural circuits involved.

To do this, we conducted butanone associative learning assays according to (Kauffman *et al.*

2011) and tested mutants deficient in dopamine synthesis and signalling. Behavioural tests for

forgetting in *C. elegans* involve assessing learned behaviours at various time points post-

conditioning to observe when responses revert to baseline levels. Our data indicate that

dopamine-deficient animals exhibit a slight improvement in learning compared to wild-type, and

the memory of this learned association is retained for at least 2 hours post-training. Additionally,

we found that DOP-2 and DOP-3 D2-like receptors work together to modulate the forgetting of

this learned association. Lastly, CAT-2 function is required across all dopaminergic neurons to

support memory retention at 2 hours post-conditioning, as re-expression in ADE alone, CEP

alone, or in both ADE and CEP together was insufficient to rescue the memory defect. Taken

together, our data showing that dopamine signals are required for forgetting is consistent with

findings in other invertebrates (Berry *et al.* 2012). These findings highlight the critical role of

dopamine in the process of forgetting and suggest implications for understanding the

pathological mechanisms that contribute to cognitive deficits in neurodegenerative diseases such

as Parkinson's disease.

Materials and Methods

C. elegans *strain maintenance*

Young adult (day 1) hermaphrodite *C. elegans* were grown using standard conditions on

nematode growth medium (NGM) agar in petri dishes at 22 °C for all experiments (Brenner

1974). Worms were cultured at 22 °C for at least two generations prior to all learning assays. For

all experiments, animals were fed *Escherichia coli* (*E. coli*) strain OP50.

A list of strains and transgenic lines used, including full genotype information, is in **Table 1**.

Genotypes of mutant strains were confirmed via PCR, based on information in Wormbase

version WS295 (Sternberg *et al.* 2024). Transgenes were generated through standard

microinjection procedures as in (Chew *et al.* 2018a; Chew *et al.* 2018b; Gadenne *et al.* 2022).

Table 1: Genotype information for strains used in this study. For transgenic lines generated in this

study, numbers in square brackets refer to the concentration injected in ng/µL.

Strain name	Genotype	Source
Bristol N2	Wild-type	*Caenorhabditis* Genetics Centre (CGC)
CB1112	*cat-2(e1112)*	*Caenorhabditis* Genetics Centre (CGC)
RM2702	*dat-1(ok157)*	*Caenorhabditis* Genetics Centre (CGC)
LX645	*dop□1(vs100)*	*Caenorhabditis* Genetics Centre (CGC)
LX702	*dop□2(vs105)*	*Caenorhabditis* Genetics Centre (CGC)
LX703	*dop□3(vs106)*	*Caenorhabditis* Genetics Centre (CGC)
LX706	*dop□1(vs100); dop□2(vs105)*	*Caenorhabditis* Genetics Centre (CGC)
LX704	*dop□2(vs105); dop□3(vs106)*	*Caenorhabditis* Genetics Centre (CGC)
LX705	*dop□1(vs100); dop□3(vs106)*	*Caenorhabditis* Genetics Centre (CGC)
LX734	*dop□1(vs100); dop□2(vs105); dop□3(vs106)*	*Caenorhabditis* Genetics Centre (CGC)
YLC299	*cat-2(e1112); baIs4 [dat-1p::GFP + dat-1p::CAT-2]*	*cat-2(e1112)* crossed with UA57 from CGC
CLP1318	*cat-2(n4547);twnEx593[Pace-1::cat-2; Pgcy-8::mCherry]*	(Chou *et al.* 2022)

CLP1320	*cat-2(n4547);twnEx599[Pexp-1::cat-2; Pgcy-8::mCherry]*	(Chou *et al.* 2022)
YLC381	*dop-2(vs105)V; dop-3(vs106)X; pepEx040[Pdop-2(5kb)::dop-2S cDNA::let-858 UTR (pYLC102) [50]; Pdop-3(5kb)::dop-3a cDNA::let-858 UTR (pYLC104) [50]; ccGFP [50]]*	This study
YLC382	*dop-2(vs105)V; dop-3(vs106)X; pepEx043[Pdop-2(5kb)::dop-2S cDNA::let-858 UTR (pYLC103) [50]; Pdop-3(5kb)::dop-3a cDNA::let-858 UTR (pYLC104) [50]; ccGFP [50]]*	This study
CLP833	*cat-2{n4547); twnEx610[Pace-1::cat-2, Pexp-1::cat-2, Pgcy-8::mCherry]*	Gift from Chun-Liang Pan, National Taiwan University School of Medicine

Molecular biology

Transgenes were cloned using the NEB HiFi cloning (NEB #E2621). Promoters for the *dop-2*

and *dop-3* gene were both cloned 5 kb before the ATG, starting from

CAAGACGGACGCAGTCCG and CTGGAATATTCCAGAACTTTC, respectively. *dop-2* and

dop-3 cDNA were amplified from wild-type N2 RNA, extracted using the using the Monarch

Total RNA Kit (NEB #T2010S) and converted to cDNA using the Luna® Universal One-Step

RT-qPCR Kit (NEB #E3005S). Two forms of the *dop-2* cDNA sequence were cloned: one

aligned to the DOP-2S sequence published in (Suo *et al.* 2009) (pYLC103), the other is identical

to this sequence but lacks a AGTACGACGGCAAAT sequence from the 1385^{th} nucleotide

position, corresponding to amino acids STTAN (pYLC102). The *dop-3* cDNA sequence aligned

with the expected sequence for DOP-3, isoform a (T14E8.3a.1) (Sternberg *et al.* 2024).

Behavioural assays

The butanone appetitive learning assay (Kauffman *et al.* 2011; Rahmani *et al.* 2024) tests naïve

worms, conditioned worms that are tested directly after training (t=0), and conditioned worms

that go through a 'hold' period between training and testing (t=0.5–2 h). Briefly, *C. elegans*

populations were washed twice in M9 buffer (86 mM NaCl, 42 mM Na_2HPO_4, 22 mM K_3PO_4

(pH 6.0), 1 mM M_gSO_4) to constitute the 'naïve' group in butanone chemotaxis assays.

Remaining groups were then subjected to a 1-hour starvation period in M9 buffer on a shaker

(175 rpm at room temperature). Conditioned groups were then transferred onto NGM plates pre-

seeded with 100 µL *E. coli* OP50 and exposed to 10% (v/v) butanone (Sigma-Aldrich, #360473;

final concentration, diluted in 95% ethanol), streaked on plate lids, for 1 h. The conditioned

groups undergoing the 'hold' period were transferred onto NGM plates pre-seeded with 100 µL

E. coli OP50 for the assigned time periods. All groups were transferred onto salt-deficient agar in

6 cm petri dishes to perform butanone chemotaxis assays as described in Kauffman *et al.* (2010)

and Rahmani *et al.* (2024). Butanone chemotaxis assay plates were prepared with two equidistant

points both spotted with sodium azide, as well as 95% (v/v) ethanol on one position (control) and

10% (v/v) butanone diluted in 95% (v/v) ethanol on the other spot. Butanone chemotaxis

behaviour was quantified using the following equation:

$$\frac{\#\ worms\ on\ butanone - \#\ worms\ on\ ethanol}{\#\ total\ population - \#\ worms\ on\ origin}$$

The chemotaxis index (CI) ranges between 0 (neutral butanone attraction) to 1.0 (complete

butanone attraction). Each assay plate constituted one technical replicate, which each contain 50-

250 worms. On very rare occasions where negative CIs are observed, these data are also

excluded as these populations failed to show the standard naïve attraction to butanone. Each

biological replicate constituted 3-4 technical replicates. At least three biological replicates were

performed for each genotype tested, the number of biological replicates is stated for each

experiment in the figure legend and are at least N>3. The Learning Index (LI) is calculated as LI

$_{\text{Time point}} = CI_{\text{Time point}} - CI_{\text{Naive}}$, where 'Time point' refers to the post-conditioning time point i.e.

t=0, 0.5, 1, 1.5 or 2 (Kauffman *et al.* 2011).

Statistical analysis

Statistical analysis for all experiments was performed using GraphPad Prism version 8.0. In

general, where two groups were compared, a paired t-test was used. A one-way ANOVA with

Dunnett's multiple comparisons test was performed to compare post-conditioning time points

with naïve chemotaxis indices (CIs) within a single genotype, or with Tukey's multiple

comparisons test to compare CIs between genotypes for naïve and t=0. The alpha value for all

analyses is set at 0.05.

<u>Results</u>

Worms lacking dopamine show extended short-term associative memory.

Dopamine regulates both non-associative and associative learning in *C. elegans* (McMillen &

Chew 2023). To determine if dopamine regulates forgetting, we used the olfactory associative

memory paradigm (**Figure 1A**) (Kauffman *et al.* 2011) to assess associative memory over time in

mutants deficient in dopamine signalling. Briefly, worms are conditioned with the odour

butanone (conditioned stimulus) and presence of food (unconditioned stimulus) for 1 hour on

solid media, before being placed onto 'hold' plates containing food but no butanone for 0.5 to 2

hours. Then, worm populations are assessed for odour preference in a two-choice chemotaxis

assay as in (Kauffman *et al.* 2011), where high chemotaxis indices (CIs) indicate strong butanone

attraction, and CIs closer to zero represent neutral butanone attraction. 'Naïve' worms do not

undergo conditioning and are placed onto chemotaxis plates immediately, while the 'Timepoint

0' (t=0) cohort are placed onto chemotaxis plates directly after conditioning. Conditioned worms

undergo a 1 hour 'pre-starvation' step to increase the value of the unconditioned stimulus (food)

(Kauffman *et al.* 2011) (**Figure 1Aii**). To simplify the visualisation of behavioural assay data, we

only display comparisons between genotypes at each post-conditioning time point, and between

naive and t=0 for wild-type and other genotypes to show if learning has occurred. Other

statistical comparisons, specifically those that compare chemotaxis behaviour WITHIN

genotypes (i.e. between naïve and post-conditioning time points) are described in the text. After

conditioning (t=0), wild-type worms show an increased attraction to butanone (shown by a

higher chemotaxis index, CI) compared with the naïve cohort, indicating the capacity for

associative olfactory learning. This memory decreases over time, returning to baseline (naïve)

levels after ~0.5 hours (comparison between CIs of naïve and 0.5 h post-conditioning cohorts: p-

value = 0.6754; one-way ANOVA with Dunnett's post-test) (**Figure 1B**).

We first tested the *cat-2* tyrosine hydroxylase mutant strain that lacks the enzyme required to

synthesise dopamine (Lints & Emmons 1999). These worms have substantially reduced

dopamine levels (Sanyal *et al.* 2004). Interestingly, *cat-2* mutant worms show a small but

significant improvement in learning, and an increased attraction to butanone for up to 2 hours

(within genotype comparison between CIs of *cat-2* naïve and 2 h post-conditioning cohorts: p-

value <0.0001; one-way ANOVA with Dunnett's post-test), suggesting they retain the memory of

the learnt association for a longer period compared with wild-type controls (**Figure 1B**). Thus,

for the purposes of our study, we consider forgetting to be the loss of the learnt phenotype

(increased attraction to butanone) at 2 hours post-conditioning. We also calculated the learning

index (LI), defined as the CI at the post-conditioning time point minus the CI of the naïve cohort,

for all time points tested. This is shown for data from wild-type and *cat-2* mutant worms in

Figure 1Bii. The *cat-2* mutant strain shows a higher LI for t=1, 1.5 and 2 compared with wild-

type controls, consistent with a longer retention of the learnt phenotype at later time points post-

conditioning.

To determine if a milder reduction in dopamine has the same effect, we next tested *dat-1* mutant

worms with a mutation in the DAT-1 sodium-dependent dopamine transporter. These worms are

able to synthesise dopamine; however, dopamine reuptake from the synaptic cleft is impaired,

reducing the amount of residual dopamine in the nervous system (Formisano *et al.* 2020; Kindt *et*

al. 2007). *dat-1* mutant worms showed improved learning compared with wild-type controls

immediately after conditioning (t=0) and retained memory of the learnt behaviour at t=0.5

(within-genotype comparison between naïve and t=0.5 cohorts: p-value <0.01 for *dat-1*; one-way

ANOVA with Dunnett's post-test). Comparing CIs indicates that *dat-1* mutant worms do not

display an extended memory phenotype like *cat-2* mutants (within-genotype comparisons

between naïve and time points after t=0.5: all p-values >0.05; one-way ANOVA with Dunnett's

post-test) (**Figure 1Ci**). Notably, the *dat-1* mutant strain shows a significantly higher LI at t=1.5,

and close to significantly different at t=1 and t=2 compared with wild-type controls (**Figure**

1Cii), suggesting that there is an effect on memory, albeit less pronounced than *cat-2* mutants.

This indicates that while excess dopamine in the synaptic cleft (as seen in *dat-1* mutants) leads to

a memory retention phenotype at 1.5 hours, dopamine synthesised by CAT-2 is required for

normal retention of the learnt phenotype (i.e. the loss of memory) at 2 hours post-conditioning.

Finally, we tested if re-expression of CAT-2 in dopaminergic neurons (using the *dat-1* promoter)

rescues the extended memory phenotype in *cat-2* mutants. We found that transgenic worms re-

expressing CAT-2 generally phenocopied wild-type animals: the re-expression strain rescued the

cat-2 phenotype to wild-type levels at t=2 post-conditioning, as shown by comparing CIs to the

cat-2 mutant (**Figure 1Ci**). Comparing LIs between the rescue strain and the mutant strain was

near-significant at t=2, with p-value 0.069 (**Figure 1Cii**). This indicates that the enzyme required

for dopamine synthesis, CAT-2, is required for forgetting and acts in the dopaminergic neurons.

Figure 1: Worms lacking dopamine show extended short-term associative memory. A) (i) Schematic of butanone associative learning and short-term memory assay, adapted from (Kauffman *et al.* 2011). Naïve worms are placed directly onto chemotaxis assay plates, while conditioned worms are first starved for 1 hour in M9 buffer prior to conditioning on plates with food and butanone (streaked on lid). After conditioning, worms are transferred to 'hold' plates with food (no odour) for 0.5–2 hours before being placed onto chemotaxis plates. **(ii)** Layout of chemotaxis assay plate indicating where butanone (But), ethanol (EtOH) and sodium azide are spotted. Worms are placed on the 'origin', as indicated. **B)** Wild-type and *cat-2(e1112)* mutant worms tested for butanone associative memory (*n* = 5 biological replicates) **(i)** chemotaxis index and **(ii)** learning index. **C) (i)** Wild-type and *dat-1(ok157)* mutant worms tested for butanone associative memory (*n* = 5 biological replicates) **(i)** chemotaxis index and **(ii)** Learning index. **D) (i)** Wild-type and *cat-2* 'rescue' worms re-expressing CAT-2 in the dopaminergic neurons tested for butanone associative memory (*n* = 4 biological replicates). **(ii)** Learning indices for wild-type, *cat-2(e1112)* mutant worms, and CAT-2 rescue worms, calculated from **D(i)**. For butanone assays (**B, C, D**), graphs show 4-5 biological replicates: each data point represents one biological replicate (which includes four technical replicates). Each technical replicate consists of 50-250 worms. Behaviour was quantified using chemotaxis indices (CI) for naïve/untrained worms and conditioned animals at 0, 0.5-, 1-, 1.5- and 2-hours post-conditioning. Statistical analysis: a pairwise t-test was performed to compare genotypes for each time point for **B** and **C**, and a one-way ANOVA with Tukey's multiple comparisons test was performed to compare post-conditioning t=0 with naïve chemotaxis indices (CIs) between genotypes for all graphs and to compare genotypes for each time point for **D**. Error bars = mean ± SD. p-values are represented by **** $\leq$ 0.0001; *** $\leq$ 0.001, ** $\leq$ 0.01; * $\leq$ 0.05; ns = non-significant.

Dopamine receptors act redundantly to modulate short-term memory.

In *C. elegans*, dopamine receptors are categorised as ligand-gated ion channels (LGIC) or G

protein-coupled receptors (GPCR). Dopaminergic GPCRs are divided into D1-like and D2-like

receptors, which stimulate or inhibit adenylyl cyclase, respectively (Chase & Koelle 2007;

Girault & Greengard 2004; Mersha *et al.* 2013). DOP-1 is a D1-like receptor, while DOP-2 and

DOP-3 are D2-like receptors (Chase & Koelle 2007; Mersha *et al.* 2013) (**Figure 2A**). We first

tested *dop-1, dop-2* and *dop-3* single mutants to determine if they showed a memory retention

phenotype similar to *cat-2* mutant worms. We found that none of the single mutants showed

statistically significant learning or memory differences compared with wild-type controls

(**Figure 2B, Figure S1**): CIs and LIs for all mutants showed intact learning at t=0, and butanone

preference had returned to baseline (naïve) levels at t=0.5 (within genotype comparisons for all

dopamine receptor mutants between naïve and t=0.5: p-values >0.05 for all genotypes; one-way

ANOVA with Dunnett's post-test). This indicates that dopamine does not regulate forgetting by

acting through DOP-1, DOP-2 or DOP-3 alone.

One possibility is that these receptors could be acting in a functionally redundant manner for

dopamine signalling to modulate forgetting in *C. elegans*, as in other behaviours such as bacterial

avoidance (Chou *et al.* 2022) or the escape response to olfactory cues (Baidya *et al.* 2014).

Therefore, we next tested a *dop-1;dop-2;dop-3* triple mutant strain and found that they displayed

similar memory compared with wild-type controls. Comparisons between naïve and post-

conditioning CIs and LIs for all time points within the triple mutant strain were not statistically

significant (for CIs, all within genotype comparisons are p>0.05, one-way ANOVA with

Dunnett's post-test) (**Figure 2Ci, 2Cii**). We postulated that there could be a synergistic or

redundant effect between D1 and D2-like dopamine receptors that could be masking the effects

of individual receptors (Chase *et al.* 2004), and so we subsequently tested double mutants for

dop-1;dop-2, *dop-1;dop-3*, and *dop-2;dop-3* for their capacity to retain memory over time

(**Figure 2D**). As in other assays, we found that wild-type controls returned to baseline (naïve) CI

values by t=0.5 (or t=1 for **Figure 2Di**). This was also the case for *dop-1;dop-2* (**Figure 2Di**) and

dop-1;dop-3 (**Figure 2Dii**) double mutants, which returned to baseline CI levels by t=1 and

t=0.5, respectively. In contrast, the *dop-2;dop-3* double mutants showed elevated CIs post-

conditioning that had not returned to baseline by t=2 (within genotype comparisons between

naïve and t>0: p-values <0.01 for all time points; one-way ANOVA with Dunnett's post-test)

(**Figure 2Diii**). Consistent with this, the LI for the *dop-2;dop-3* double mutant is significantly

higher than wild-type controls at t=2 (**Figure 2Div**). Taken together, these data suggest that the

D1-like receptor DOP-1 and D2-like receptors DOP-2 and DOP-3 act redundantly to modulate

forgetting of an olfactory associative memory.

Figure 2: Dopamine receptors act synergistically to modulate short-term memory. **A)** *C. elegans* dopamine receptors DOP-1, DOP-2, and DOP-3 are D1 or D2-like receptors (Chase & Koelle 2007; Mersha *et al.* 2013). **B)** Butanone associative memory assay for wild-type worms and worms containing single mutations in **(i)** *dop□1(vs100)* (*n* = 4 biological replicates), **(ii)** *dop□2(vs105)* (*n* = 4 biological replicates), and **(iii)** *dop□3(vs106)* (*n* = 4 biological replicates). **C)** Butanone associative memory assay for wild-type and *dop□1;dop□2;dop□3* triple mutant worms (*n* = 4 biological replicates) **(i)** chemotaxis index and **(ii)** learning index **D)** Butanone associative memory assay for wild-type and **(i)** *dop□1;dop-2* (*n* = 4 biological replicates), **(ii)** *dop□1;dop-3* (*n* = 3 biological replicates), and **(iii)** *dop□2;dop-3* (*n* = 4 biological replicates) double mutant worms. **(iv)** The learning index calculated from **(iii)**. For butanone assays (**B, C, D**), each data point represents one biological replicate (which includes four technical replicates). Each technical replicate consists of 50-250 worms. Behaviour was quantified using chemotaxis indices (CI) for naïve/untrained worms and conditioned animals at 0, 0.5-, 1-, 1.5- and 2-hours post-conditioning. Statistical analysis: a pairwise t-test was performed to compare genotypes for each time point, and a one-way ANOVA with Tukey's multiple comparisons test was performed to compare post-conditioning t=0 with naïve chemotaxis indices (CIs) between genotypes. Error bars = mean ± SD. p-values are represented by **** ≤ 0.0001; *** ≤ 0.001, ** ≤ 0.01; * ≤ 0.05; ns = non-significant.

Dopamine acts in dop-2/dop-3 expressing neurons to modulate short-term memory

Our data suggests that dopamine acts through D2-like dopamine GPCRs, DOP-2 and DOP-3, to promote forgetting in *C. elegans*. We re-expressed DOP-2 and DOP-3 in *dop-2;dop-3* double mutants using a transgene that contains DOP-2 cDNA (corresponding to DOP-2S in (Suo *et al.* 2009)) and DOP-3a cDNA, expressed under the control of 5 kb of promoter sequence for both *dop-2* and *dop-3* (see Materials and Methods for details). Due to practical limitations on the number of groups that can be handled in a single experiment, for these assays we tested only naïve and post-conditioning t=0, 0.5 and 2 h cohorts for four genotypes: wild-type, *dop-2;dop-3* double mutants, and two versions of the transgenic DOP-2/3 rescue line (**Figure 3A** and **B**). As the DOP-2/3 rescue transgenic lines contain extrachromosomal arrays, there may be minor

differences in transgene expression, so we tested two versions to assess the role of DOP-2 and

DOP-3 in forgetting. We found for data shown in both **Figure 3A** and **3B** that, as expected, wild-

type controls showed intact learning at t=0, as indicated by higher CI compared with naïve

worms, and that this had returned to baseline at t=2 (comparisons between naïve and t=0, 0.5,

and 2 for all genotypes are indicated underneath each graph). Consistent with **Figure 2Diii**, *dop-*

2;dop-3 mutants also displayed intact learning at t=0, and the memory of this learnt association

did not return to baseline levels at t=2 (comparison between CIs for naïve and t=2: p-value

<0.05; one-way ANOVA with Dunnett's post-test) (**Figures 3A, 3B**), indicating an extended

memory retention phenotype. When comparing CIs, we found that both DOP-2/3 re-expression

lines showed a trend towards a rescue phenotype: transgenic worms showed intact learning at

t=0, and had returned to naïve levels at t=2 (within-genotype comparison between CIs for naïve

and t=2: p-value >0.05; one-way ANOVA with Dunnett's post-test) (**Figures 3Ai, 3Bi**). Although

the mean CI for both rescue strains was intermediate between wild-type or *dop-2;dop-3* mutant

controls, the CIs of DOP-2/3 rescue worms were not significantly different from either of the

controls at t=2 (**Figures 3Ai, 3Bi**). Comparison of LIs at t=2 showed that the LIs of both rescue

strains were lower than the mutant (approaching wild-type levels), but this was not statistically

significant (**Figures 3Aii, 3Bii**); notably, the comparison between the rescue strain and mutant

was near-significant for v2 at p=0.07 (**Figure 3Bii**). Taken together, these data suggest a

consistent trend suggesting that re-expression of DOP-2 and DOP-3 in the *dop-2;dop-3* double

mutant background contributes towards restoring the forgetting phenotype. While the rescue

effect does not reach statistical significance, the observed effects support a potential role for

these receptors in modulating memory decay.

Figure 3: Dopamine acts through DOP-2 and DOP-3 to modulate short-term memory. A) Butanone associative memory assay for wild-type worms, *dop-2;dop-3* double mutant worms and *dop-2;dop-3* double mutant worms re-expressing DOP-2 and DOP-3 under the control of their endogenous promoters, **(i)** the chemotaxis index and **(ii)** the learning index graph. DOP-2 in this transgene is a truncated version missing 5 amino acids when compared to the DOP-2S sequence (*n* = 5 biological replicates). **B**) Butanone associative memory assay for wild-type worms, *dop-2;dop-3* double mutant worms and *dop-2;dop-3* double worms re-expressing DOP-2 and DOP-3 under the control of their endogenous promoters; **(i)** the chemotaxis index and **(ii)** the learning index graph. DOP-2 in this transgene exactly matches the published DOP-2S sequence (Suo *et al.* 2009) (*n* = 4 biological replicates). For rescue lines expressing extrachromosomal arrays, only CIs from transgenic worms are displayed here. Each data point represents one biological replicate (which includes four technical replicates). Each technical replicate consists of 50-250 worms. Behaviour was quantified using chemotaxis indices (CI) for naïve/untrained worms and conditioned animals at 0, 0.5-, and 2-hours post-conditioning. Statistical analysis: one-way ANOVA with Dunnett's multiple comparisons test was performed to compare post-conditioning time points with naïve chemotaxis indices (CIs) within a single genotype (shown in tables below each graph) and with Tukey's

331   multiple comparisons test to compare CIs for naïve vs t=0 between genotypes. Error bars = mean ± SD. p-

332   values are represented by **** $\leq$ 0.0001; *** $\leq$ 0.001, ** $\leq$ 0.01; * $\leq$ 0.05; ns = non-significant.

Dopamine release through ADE and CEP neurons promotes forgetting.

There are eight dopaminergic neurons in hermaphrodite *C. elegans*: two ADE neurons, four CEP

neurons, and two PDE neurons (**Figure 4A**), each of which have specific synaptic and

neuromodulatory inputs/outputs to other parts of the nervous system. To investigate whether

specific connections are required for memory retention, we examined the role of individual

subsets of dopamine-releasing neurons in the process of forgetting. To do this, we assayed

transgenic lines in which CAT-2 is specifically re-expressed in the ADE (using an *exp-1*

promoter) or CEP neurons (using an *ace-1* promoter), or both, in a *cat-2* mutant background

(Chou *et al.* 2022). We first examined the role of ADE neurons in forgetting (**Figure 4B**): wild-

type controls displayed the expected phenotype of showing an increased CI (indicating intact

learning) at t=0, which had returned to baseline at t=0.5 (within genotype comparison between

CIs for naïve and t=0.5: p-value >0.05; one-way ANOVA with Tukey's post-test). We found that

ADE::CAT-2 rescue worms showed intact learning at t=0, which had also returned to baseline at

t=0.5 (within genotype comparison between CIs for naïve and t=0.5: p-value >0.05; one-way

ANOVA with Tukey's post-test) (**Figure 4A**). To assess whether re-expression in ADE neurons

alone rescued the forgetting phenotype in *cat-2* mutants, we compared CIs and LIs at t=2

between the ADE rescue strain and mutant siblings. We found that there was no statistically

significant difference in CI (**Figure 4Ai**) or LI (**Figure 4Aii**), suggesting that CAT-2 expression

in ADE neurons is insufficient for normal forgetting.

We next tested CEP::CAT-2 rescue worms (**Figure 4B**) in the butanone associative memory

assay. We found that wild-type worms showed the expected phenotype of intact learning at t=0

and return to baseline (naïve) CI at t=0.5. In contrast, CEP::CAT-2 worms showed significantly

higher CIs at t=0 (indicating intact learning) and t=0.5 (within genotype comparison between

naïve and t=0.5 gives p-value of 0.0103; one way ANOVA, Tukey's post-test). Similarly, to

assess whether re-expression in CEP neurons alone rescued the forgetting phenotype in *cat-2*

mutants, we compared CIs and LIs at t=2 between the rescue strain and mutant siblings. There

was no statistically significant difference in CI (**Figure 4Bi**) or LI (**Figure 4Bii**) t=2, indicating

that CAT-2 expression in CEP neurons is insufficient for normal forgetting.

As re-expression of CAT-2 in ADE or CEP did not rescue the extended memory phenotype

observed in *cat-2* mutants, we tested a strain where CAT-2 was re-expressed in both ADE and

CEP neurons (referred to as ADE+CEP; **Figure 4C**). Similar to ADE and CEP alone, this

ADE+CEP rescue line did not show a statistically different CI (**Figure 4Ci**) or LI (**Figure 4Cii**)

compared to the *cat-2* mutant siblings at t=2. Taken together, this indicates that CAT-2

expression in ADEs and CEPs does not rescue the memory phenotype, and suggests that

dopamine release from all dopaminergic neurons (ADEs, CEPs and PDEs), or from PDE alone,

is required for normal forgetting.

Figure 4: **Dopamine release from ADE and CEP neurons promotes forgetting. A)** Butanone

associative memory assay for wild-type worms and *cat-2* mutant worms re-expressing CAT-2 in the ADE

neurons (*n* = 7 biological replicates); **(i)** the chemotaxis index and **(ii)** the learning index graphs. **B)**

Butanone associative memory assay for wild-type worms and *cat-2* mutant worms re-expressing CAT-2 in

the CEP neurons (*n* = 7 biological replicates); **(i)** the chemotaxis index and **(ii)** the learning index graphs.

C) Butanone associative memory assay for wild-type worms and *cat-2* mutant worms re-expressing CAT-

2 in both the ADE and CEP neurons (*n* = 4 biological replicates); **(i)** the chemotaxis index and **(ii)** the

learning index graphs. Each data point represents one biological replicate (which includes four technical

replicates). Each technical replicate consists of 50-250 worms. Behaviour was quantified using

chemotaxis indices (CI) for naïve/untrained worms and conditioned animals at 0, 0.5-, and 2-hours post-

conditioning. Statistical analysis: a pairwise t-test was performed to compare genotypes for each time

point, and a one-way ANOVA with Tukey's multiple comparisons test was performed to compare post-

conditioning t=0 with naïve chemotaxis indices (CIs) between genotypes. Error bars = mean ± SD. p-

values are represented by **** ≤ 0.0001; *** ≤ 0.001, ** ≤ 0.01; * ≤ 0.05; ns = non-significant. **D)**

Schematic of potential mechanism, summarising data in **Figures 1-4**: our findings suggest that newly

synthesised dopamine from all dopaminergic neurons is required for forgetting of olfactory associative

memory, acting primarily through DOP-2 and DOP-3 receptors.

Discussion

Our study demonstrates that dopamine is essential for active forgetting in *C. elegans*. Our

findings indicate that dopamine-deficient mutants, particularly *cat-2*, exhibit enhanced memory

retention following conditioning, consistent with a delayed forgetting phenotype. The *dat-1*

mutant, which retains dopamine synthesis but has impaired reuptake, shows a similar trend,

albeit less pronounced. These results suggest that even partial restoration of dopamine may

influence forgetting, consistent with the idea that boosting dopamine levels (such as via

pharmacological pretreatment) could restore the delayed forgetting phenotype in *cat-2* mutants.

Additionally, our findings indicate that D2-like receptors, specifically DOP-2 and DOP-3, are

required for forgetting. Finally, we observed that CAT-2 function is required across all

dopaminergic neurons to support memory retention at 2 hours post-conditioning. Re-expression

of *cat-2* in ADE alone, CEP alone, or in both ADE and CEP together was insufficient to rescue

the memory defect observed in the mutant background at this time point (**Figure 4**).

The role of dopamine in learning and memory is conserved across species. However, while

dopamine synthesis and signalling pathways are highly conserved, dopaminergic circuits

between *C. elegans* and other organisms differ. In *C. elegans*, eight dopaminergic neurons act

'extra-synaptically' on over 100 targets, mainly motor neurons (Bentley *et al.* 2016). In

mammals, dopaminergic neurons are concentrated in the ventral midbrain (SNc and VTA) and

project to the basal ganglia, amygdala, and prefrontal cortex. In *Drosophila melanogaster*, small

populations of dopaminergic neurons are found throughout the fly brain, with projections

innervating almost the entire brain, with dense projections to the mushroom body lobes, which

regulate olfactory learning (Riemensperger *et al.* 2005; Marquis & Wilson 2022). Despite these

differences, our finding that dopamine is required for forgetting aligns with previous research on

the role of dopamine in forgetting in *Drosophila*. In flies, blocking synaptic output from

dopaminergic neurons promotes memory retention (i.e. reduces forgetting), whereas activating

dopamine neurons decreases memory (Berry *et al.* 2012). Indeed, dopamine is essential for both

forgetting and memory acquisition in flies, but this involves different receptors: DAMB for

forgetting and dDA1 for acquisition (Berry *et al.* 2012; Himmelreich *et al.* 2017). In our study,

we observed a small but robust learning improvement in *cat-2* and *dat-1* mutant worms, as well

as in *dop-1;dop-2;dop-3* triple mutants (double mutant combinations of these three receptors also

showed a learning improvement, albeit slightly above the alpha value of $p = 0.05$). This indicates

that the requirement for dopamine in learning and forgetting is conserved between flies and

worms, suggesting it is evolutionary important and likely also retained in higher organisms.

Our findings contrast with the work by Raj and Thekkuveettil (Raj & Thekkuveettil 2022), which

showed that dopamine deficient *cat-2* mutant *C. elegans* display both a learning defect and a

memory defect, i.e. they forget more quickly than wild-type controls. Although both our study

and the 2022 study use the butanone associative learning assay, worms in this research were

conditioned with agar plates seeded 1-2 days in advance with 100 µL OP50 (and thus already

dried on the agar), whereas the Raj and Thekkuveettil study involved conditioning in 400 µL of

OP50 pipetted onto unseeded agar plates immediately after worms are added. This means that the

worms are being conditioned in liquid, which may influence other locomotion pathways in which

dopamine plays an important role, such as in regulating "swimming" in liquid, a distinct form of

locomotion separate to crawling on solid media. Dopamine modulates swimming behaviour (Xu

et al. 2021), swimming-induced paralysis (McDonald *et al.* 2007), and swim-to-crawl transitions

when moving from liquid to solid media (Vidal-Gadea *et al.* 2011). As learning is a multifaceted

process modulated by multiple internal and external cues, it is not surprising that dopamine could

impact memory in different ways depending on the mode of conditioning, for example due to an

interaction between the response to the food and odour cues, and the swimming activity. This

underscores dopamine's multifaceted role in modulating behaviour and memory in *C. elegans*.

Our data show that both DOP-2 and DOP-3 D2-like receptors are required to modulate

forgetting. Single mutants of either receptor show memory retention capacity similar to wild-

type, while the *dop-2;dop-3* double mutant show enhanced memory retention similar to *cat-2*

mutant animals. Taken together, our data suggest that the D1-like receptor DOP-1 and D2-like

receptors DOP-2 and DOP-3 act redundantly to modulate forgetting of an olfactory associative

memory. Although D1- and D2-like receptors are known to antagonise each other in signal

transduction, and their opposing effects in *C. elegans* have been previously reported in dietary

restriction-mediated lifespan extension (Jiang *et al.* 2022), swimming (Xu *et al.* 2021), and basal

slowing behaviour in response to a food lawn (Chase *et al.* 2004), this did not appear to be the

case in our study. We did not test the invertebrate specific DOP-4 receptor or dopamine ligand-

gated ion channels such as LGC-53; therefore, a role for these receptors in dopamine-dependent

short-term associative memory in *C. elegans* remains unknown. Importantly, although re-

expression of DOP-2 and DOP-3 in the mutant background showed trends toward restoring

normal forgetting, these effects did not reach statistical significance (**Figure 3**). Future studies

could refine the re-expression strategy, such as by modulating DNA dosage for microinjection or

selecting alternative isoforms, to enhance rescue efficacy and more clearly define the

contributions of these receptors to forgetting.

Short- and long-term memory differ by the mechanism for encoding and time taken to fully

decay. In *C. elegans*, short-term memory is typically considered to be relevant at 1-2 hours post-

learning, while long-term memory is evaluated 12-24 hours post-learning (Rahmani & Chew

2021; Kauffman *et al.* 2011). In our study, we tested short-term associative memory only up to 2

hours post-conditioning. At this time point, dopamine-deficient *cat-2* mutants, as well as *dop-*

2;dop-3 double mutants, still showed robust memory. Identifying when the memory decays to

baseline levels may provide further understanding of the role of dopamine in forgetting, such as

whether processes associated with long-term memory e.g. *de novo* protein synthesis (Fioriti *et al.*

2015), are required. The mechanistic details of this process warrant further investigation in future

studies.

In conclusion, our study leverages the worm's compact nervous system and genetic tractability to

provide new insights into the role of dopamine in forgetting. Other studies on olfactory

paradigms in worms have identified diverse regulators of forgetting, indicating that multiple

pathways and mechanisms are involved in this complex process. Our study adds to this body of

knowledge by highlighting the specific role of dopamine and its receptors. In addition, these

findings are consistent with research in *Drosophila*, suggesting a mechanism for forgetting that is

conserved in even more complex brains. Based on this study, future research could focus on

exploring the conserved mechanisms of forgetting across species and investigating potential

therapeutic targets for memory-related disorders.

<u>Acknowledgements</u>

Many thanks go to our team members in the Worm Neuroscience lab (Flinders) for thoughtful

discussions, and to our colleagues at Flinders University – Dr Nicholas Eyre, Professor Damien

Keating, and Professor Kim Hemsley – for feedback and scientific advice. We especially thank

Pasquale Vitaro (Summer Scholar 2025) for technical assistance with behavioural assays, and

Aelon Rahmani for assistance with genotyping. We are grateful to Prof Chun-Liang Pan

(National Taiwan University School of Medicine) for strains CLP1318, CLP1320, and CLP833

(cell-specific rescue of *cat-2*), and Prof Satoshi Suo (Saitama Medical University, Japan) for

plasmids encoding *dop-2* cDNA. Finally, we gratefully acknowledge the *Caenorhabditis*

Genetics Centre, which is supported by the National Institutes of Health (P40 OD010440), for

providing some of the strains used in this study (see **Table 1**).

Funding: A.M is funded by a Flinders University Research Scholarship (Flinders University).

Y.L.C is funded by the National Health and Medical Research Council (NHMRC)

(GNT1173448), the Flinders University Parental Leave Research Support Scheme, the Flinders

University Impact Seed Funding Grant for Early Career Researchers, and the Flinders

Foundation Mary Overton Senior Research Fellowship.

Author contributions: Conceptualisation/Methodology– **AM, YLC**. Data curation/ Formal

analysis/ Investigation/ Resources – **AM, CM, MEJ, RG, RA, YLC.** Project administration/

Supervision/ Funding acquisition – **YLC.** Visualisation/ Writing – original draft - **AM, YLC**

Writing – review & editing – **AM, MEJ, YLC**

Conflict of interest: The authors declare that they have no conflicts of interest.

Data Availability Section

This study includes no data deposited in external repositories. Reagents including *C. elegans*

strains and expression plasmids are available from the corresponding author upon request.

References

Arai, M., Kurokawa, I., Arakane, H., Kitazono, T. and Ishihara, T. (2022) Regulation of
Diacylglycerol Content in Olfactory Neurons Determines Forgetting or Retrieval of
Olfactory Memory in Caenorhabditis elegans. *J Neurosci* **42,** 8039-8053.
Ardiel, E. L. and Rankin, C. H. (2010) An elegant mind: learning and memory in Caenorhabditis
elegans. *Learn Mem* **17,** 191-201.
Bai, H., Huang, H., Zhao, N., Gu, H., Li, Y., Zou, W., Wu, T. and Huang, X. (2022) Small G
protein RAC-2 regulates forgetting via the JNK-1 signalling pathway in Caenorhabditis
elegans. *Eur J Neurosci* **56,** 6162-6173.
Baidya, M., Genovez, M., Torres, M. and Chao, M. Y. (2014) Dopamine modulation of avoidance
behavior in Caenorhabditis elegans requires the NMDA receptor NMR-1. *PLoS One* **9,**
e102958.
Bentley, B., Branicky, R., Barnes, C. L., Chew, Y. L., Yemini, E., Bullmore, E. T., Vertes, P. E. and
Schafer, W. R. (2016) The Multilayer Connectome of Caenorhabditis elegans. *PLoS*
Comput Biol **12,** e1005283.
Berry, J. A., Cervantes-Sandoval, I., Chakraborty, M. and Davis, R. L. (2015) Sleep Facilitates
Memory by Blocking Dopamine Neuron-Mediated Forgetting. *Cell* **161,** 1656-1667.

Berry, J. A., Cervantes-Sandoval, I., Nicholas, E. P. and Davis, R. L. (2012) Dopamine is
required for learning and forgetting in Drosophila. *Neuron* **74,** 530-542.
Berry, J. A., Phan, A. and Davis, R. L. (2018) Dopamine Neurons Mediate Learning and
Forgetting through Bidirectional Modulation of a Memory Trace. *Cell Rep* **25,** 651-662
e655.
Brandel-Ankrapp, K. L. and Arey, R. N. (2023) Uncovering novel regulators of memory using C.
elegans genetic and genomic analysis. *Biochem Soc Trans* **51,** 161-171.
Brenner, S. (1974) The genetics of Caenorhabditis elegans. *Genetics* **77,** 71-94.
Chase, D. L. and Koelle, M. R. (2007) Biogenic amine neurotransmitters in C. elegans.
WormBook: The Online Review of C. elegans Biology [Internet].
Chase, D. L., Pepper, J. S. and Koelle, M. R. (2004) Mechanism of extrasynaptic dopamine
signaling in Caenorhabditis elegans. *Nat Neurosci* **7,** 1096-1103.
Chew, Y. L., Grundy, L. J., Brown, A. E. X., Beets, I. and Schafer, W. R. (2018a) Neuropeptides
encoded by *nlp-49* modulate locomotion, arousal and egg-laying behaviours in
Caenorhabditis elegans via the receptor SEB-3. *Philos Trans R Soc Lond B Biol Sci* **373**.
Chew, Y. L., Tanizawa, Y., Cho, Y. et al. (2018b) An Afferent Neuropeptide System Transmits
Mechanosensory Signals Triggering Sensitization and Arousal in C. elegans. *Neuron* **99,**
1233-1246 e1236.
Chou, S. H., Chen, Y. J., Liao, C. P. and Pan, C. L. (2022) A role for dopamine in C. elegans
avoidance behavior induced by mitochondrial stress. *Neurosci Res* **178,** 87-92.
Fioriti, L., Myers, C., Huang, Y. Y. et al. (2015) The Persistence of Hippocampal-Based Memory
Requires Protein Synthesis Mediated by the Prion-like Protein CPEB3. *Neuron* **86,** 1433-
1448.
Formisano, R., Mersha, M. D., Caplan, J., Singh, A., Rankin, C. H., Tavernarakis, N. and
Dhillon, H. S. (2020) Synaptic vesicle fusion is modulated through feedback inhibition by
dopamine auto-receptors. *Synapse* **74,** e22131.
Gadenne, M. J., Hardege, I., Yemini, E., Suleski, D., Jaggers, P., Beets, I., Schafer, W. R. and
Chew, Y. L. (2022) Neuropeptide signalling shapes feeding and reproductive behaviours
in male Caenorhabditis elegans. *Life Science Alliance* **5,** e202201420.
Giles, A. C., Rose, J. K. and Rankin, C. H. (2006) Investigations of learning and memory in
Caenorhabditis elegans. *Int Rev Neurobiol* **69,** 37-71.
Girault, J.-A. and Greengard, P. (2004) The neurobiology of dopamine signaling. *Archives of
neurology* **61,** 641-644.
Hadziselimovic, N., Vukojevic, V., Peter, F. et al. (2014) Forgetting is regulated via Musashi-
mediated translational control of the Arp2/3 complex. *Cell* **156,** 1153-1166.
Himmelreich, S., Masuho, I., Berry, J. A., MacMullen, C., Skamangas, N. K., Martemyanov, K. A.
and Davis, R. L. (2017) Dopamine Receptor DAMB Signals via Gq to Mediate Forgetting
in Drosophila. *Cell Rep* **21,** 2074-2081.
Inoue, A., Sawatari, E., Hisamoto, N., Kitazono, T., Teramoto, T., Fujiwara, M., Matsumoto, K.
and Ishihara, T. (2013) Forgetting in C. elegans is accelerated by neuronal
communication via the TIR-1/JNK-1 pathway. *Cell Rep* **3,** 808-819.
Jiang, Y., Gaur, U., Cao, Z., Hou, S. T. and Zheng, W. (2022) Dopamine D1- and D2-like
receptors oppositely regulate lifespan via a dietary restriction mechanism in
Caenorhabditis elegans. *BMC Biol* **20,** 71.
Kandel, E. R., Dudai, Y. and Mayford, M. R. (2014) The molecular and systems biology of
memory. *Cell* **157,** 163-186.
Kauffman, A., Parsons, L., Stein, G., Wills, A., Kaletsky, R. and Murphy, C. (2011) C. elegans
positive butanone learning, short-term, and long-term associative memory assays. *J Vis
Exp*.

Kauffman, A. L., Ashraf, J. M., Corces-Zimmerman, M. R., Landis, J. N. and Murphy, C. T.
(2010) Insulin signaling and dietary restriction differentially influence the decline of
learning and memory with age. *PLoS Biol* **8,** e1000372.
Kindt, K. S., Quast, K. B., Giles, A. C., De, S., Hendrey, D., Nicastro, I., Rankin, C. H. and
Schafer, W. R. (2007) Dopamine mediates context-dependent modulation of sensory
plasticity in C. elegans. *Neuron* **55,** 662-676.
Kitazono, T., Hara-Kuge, S., Matsuda, O., Inoue, A., Fujiwara, M. and Ishihara, T. (2017)
Multiple Signaling Pathways Coordinately Regulate Forgetting of Olfactory Adaptation
through Control of Sensory Responses in Caenorhabditis elegans. *J Neurosci* **37,**
10240-10251.
Lints, R. and Emmons, S. W. (1999) Patterning of dopaminergic neurotransmitter identity among
Caenorhabditis elegans ray sensory neurons by a TGFβfamily signaling pathway and a
Hox gene. *Development* **126,** 5819-5831.
Liu, H., Wu, T., Canales, X. G., Wu, M., Choi, M. K., Duan, F., Calarco, J. A. and Zhang, Y.
(2022) Forgetting generates a novel state that is reactivatable. *Sci Adv* **8,** eabi9071.
Marquis, M. and Wilson, R. I. (2022) Locomotor and olfactory responses in dopamine neurons
of the Drosophila superior-lateral brain. *Curr Biol* **32,** 5406-5414 e5405.
McDonald, P. W., Hardie, S. L., Jessen, T. N., Carvelli, L., Matthies, D. S. and Blakely, R. D.
(2007) Vigorous motor activity in Caenorhabditis elegans requires efficient clearance of
dopamine mediated by synaptic localization of the dopamine transporter DAT-1. *J*
Neurosci **27,** 14216-14227.
McMillen, A. and Chew, Yee L. (2023) Neural mechanisms of dopamine function in learning and
memory in Caenorhabditis elegans. *Neuronal Signaling* **8,** NS20230057.
Mersha, M., Formisano, R., McDonald, R., Pandey, P., Tavernarakis, N. and Harbinder, S.
(2013) GPA-14, a Gαi subunit mediates dopaminergic behavioral plasticity in C. elegans.
Behavioral and Brain Functions **9,** 1-8.
Niu, S., Zhou, W., Li, Y. and Huang, X. (2022) The signaling pathway of levamisole-sensitive-
acetylcholine receptors involved in short-term forgetting of Caenorhabditis elegans. *Mol*
Genet Genomics **297,** 1027-1038.
Rahmani, A. and Chew, Y. L. (2021) Investigating the molecular mechanisms of learning and
memory using Caenorhabditis elegans. *J Neurochem* **159,** 417-451.
Rahmani, A., McMillen, A., Allen, E., Minervini, C. and Chew, Y. L. (2024) Behavioral Tests for
Associative Learning in Caenorhabditis elegans. In: *Neurobiology: Methods and*
Protocols, (S. Dworkin ed.)*,* pp. 21-46. Springer US, New York, NY.
Raj, V. and Thekkuveettil, A. (2022) Dopamine plays a critical role in the olfactory adaptive
learning pathway in Caenorhabditis elegans. *J Neurosci Res* **100,** 2028-2043.
Riemensperger, T., Voller, T., Stock, P., Buchner, E. and Fiala, A. (2005) Punishment prediction
by dopaminergic neurons in Drosophila. *Curr Biol* **15,** 1953-1960.
Sanyal, S., Wintle, R. F., Kindt, K. S. et al. (2004) Dopamine modulates the plasticity of
mechanosensory responses in Caenorhabditis elegans. *The EMBO journal* **23,** 473-482.
Stein, G. M. and Murphy, C. T. (2014) C. elegans positive olfactory associative memory is a
molecularly conserved behavioral paradigm. *Neurobiol Learn Mem* **115,** 86-94.
Sternberg, P. W., Van Auken, K., Wang, Q. et al. (2024) WormBase 2024: status and
transitioning to Alliance infrastructure. *Genetics* **227**.
Suo, S., Culotti, J. G. and Van Tol, H. H. (2009) Dopamine counteracts octopamine signalling in
a neural circuit mediating food response in C. elegans. *EMBO J* **28,** 2437-2448.
Vidal-Gadea, A., Topper, S., Young, L. et al. (2011) Caenorhabditis elegans selects distinct
crawling and swimming gaits via dopamine and serotonin. *Proc Natl Acad Sci U S A* **108,**
17504-17509.

Vukojevic, V., Gschwind, L., Vogler, C., Demougin, P., de Quervain, D. J., Papassotiropoulos, A.
and Stetak, A. (2012) A role for alpha-adducin (ADD-1) in nematode and human
memory. *EMBO J* **31,** 1453-1466.
Xu, Y., Zhang, L., Liu, Y. et al. (2021) Dopamine receptor DOP-1 engages a sleep pathway to
modulate swimming in C. elegans. *iScience* **24,** 102247.

 Supplementary Figure legends

 Figure S1: Dopamine receptors *dop-1*, *dop-2* and *dop-3* act redundantly to modulate short-term
 memory. Learning indices calculated for naïve/untrained worms and conditioned animals at 0, 0.5-, 1-,
 1.5- and 2-hours post-conditioning, for chemotaxis data shown in Figure 2: **A)** *dop-1* mutants, **B)** *dop-2*
 mutants, **C)** *dop-3* mutants, **D)** *dop-1;dop-2* double mutants, **E)** *dop-1;dop-3* double mutants. Graphs
 show 3-4 biological replicates: each data point represents one biological replicate (which includes four
 technical replicates). Each technical replicate consists of 50-250 worms. Statistical analysis: a one-way
 ANOVA with Tukey's multiple comparisons test was performed to compare between genotypes for all
 post-conditioning time points. Error bars = mean ± SD. p-values are represented by **** ≤ 0.0001; *** $\leq$
 0.001, ** ≤ 0.01; * ≤ 0.05; ns = non-significant.

637   **Figure S1**

Figure 1 (McMillen *et al.*)

Figure 2 (McMillen *et al.*)

Figure 3 (McMillen *et al.*)

A(i)

A(ii)

B(i)

B(ii)

Figure 4 (McMillen *et al.*)