Dehalobacter **converts 2,3-dichloroaniline to aniline via 2-chloroaniline in a methanogenic**

enrichment culture

Shuping Wang,[1] Sofia P. Araújo,[2,3] Line Lomheim,[2] E. Erin Mack,[4] Elizabeth A. Edwards,[2*] and

Elodie Passeport[1,2,5]

[1]Department of Civil & Mineral Engineering, University of Toronto

35 St. George Street, Toronto, Ontario, M5S 1A4, Canada

[2]Department of Chemical Engineering & Applied Chemistry, University of Toronto

200 College Street, Toronto, Ontario, M5S 3E5, Canada

[3]Departamento de Engenharia Civil e Ambiental, Universidade Federal de Pernambuco, Recife,

Pernambuco, 50740-530, Brazil

[4]Corteva Environmental Remediation, Corteva Agriscience, Wilmington, DE, 19805, USA

[5]Department of Environmental Sciences, Rutgers University, 14 College Farm Rd, New

Brunswick NJ 08901, USA

*Corresponding authors: elodie.passeport@rutgers.edu, +1 848-932-6740
elizabeth.edwards@utoronto.ca, +1 416-946-3506
24

1   ABSTRACT

2,3-dichloroaniline (2,3-DCA) has widespread use in chemical manufacturing and remains a

persistent groundwater contaminant. To better understand the mechanism and kinetics of its

anaerobic biotransformation, we conducted a laboratory kinetic experiment using an anaerobic

enrichment culture dominated by *Dehalobacter*. At an initial field-relevant concentration of 40

mg/L, complete and stoichiometric dechlorination of 2,3-DCA to aniline was achieved. The

intermediates, 2-chloroaniline (2-CA) and 3-chloroaniline, were transiently formed in a ratio of

8:1. The growth yields of *Dehalobacter* on 2,3-DCA and 2-CA were $1.2 \pm 0.1 \times 10^8$ and $1.3 \pm$

0.1×10^8 16S rRNA gene copies/µmol chloride, respectively. The maximum specific growth rate

for 2,3-DCA, μ_{max}, was 0.18 ± 0.03 day^{-1} with a half-saturation constant, K_s, at 45 ± 16 mg /L.

The first-order decay constant for *Dehalobacter* when starved of chlorinated electron acceptors

was estimated at 0.017 ± 0.001 day^{-1}. Lactate fermentation, acetogenesis from ethanol,

syntrophic propionate oxidation, and hydrogenotrophic methanogenesis were observed during

dechlorination. This work provides insights into the biotransformation pathway of 2,3-DCA to

aniline and advances the understanding of microbial interactions during anaerobic

dechlorination. These results offer guidance for developing stable dechlorinating microbial

ecosystems and key kinetic parameters for predictive modeling of groundwater 2,3-DCA fate

and transport.

2   Keywords

Biotransformation   Anaerobic Microbial Community

Reductive dechlorination   Syntrophy   Microbial Kinetics   Pesticide Metabolites

3   Synopsis

47   This study demonstrated stochiometric dechlorination of dichloroaniline to aniline, characterized

48   syntrophic interactions in a mixed anaerobic dechlorinating culture, and determined growth

49   yield, first-order decay constant, and Monod constants for *Dehalobacter*.

51   **Graphical abstract**

52
53   Created in BioRender. Wang, S. (2025) https://BioRender.com/f47t993
54

4 INTRODUCTION

Dichloroaniline (DCA) isomers are a class of amino-substituted chlorinated aromatic compounds that are widely used as intermediates in the manufacture of various chemical products, including herbicides, cosmetics, pigments, pharmaceuticals, and antimicrobials.[1–3] DCAs are primarily released into the environment through improper handling and disposal during industrial processes, they are also produced from natural attenuation of phenylurea herbicides and nitro-substituted aromatics.[4–6] Due to high chemical stability and toxicity, DCA contamination poses a significant environmental challenge, particularly in aquatic and soil ecosystems.[7–9]

Contaminated groundwater environments, which are often anoxic, provide conditions for reductive biotransformation processes. Such processes have proven successful for many halogenated pollutants, particularly chlorinated alkenes and alkanes.[10–12] However, studies on the anaerobic biotransformation of DCAs remain limited.

Reductive dechlorination of DCAs was first observed using pond sediment[13] and aquifer slurries[14]. The latter reported the biotransformation of 3,4-DCA (around 50 µM) to 3-chloroaniline (3-CA) during the 8-month experiment.[14] Similarly, Susarla et al. (1998) observed reductive dechlorination of 3,4-DCA (4 µM) using sulfidogenic estuarine sediment with 3-CA and aniline as the end products.[15] However, these early studies used low DCA concentrations and often required extended incubation periods (e.g., >200 days) for complete transformation. Moreover, the microorganisms responsible for these anaerobic transformations were not identified and the relatively poor mass balance obtained in these studies limited the understanding of the underlying biotransformation pathways.

More recent studies have identified microorganisms capable of anaerobic DCA biotransformation. Duc and Oanh (2019) isolated *Geobacter* sp. KT5 from contaminated river

sediment, which grew on a wide range of chlorinated anilines (50 µM) and transiently produced

dechlorinated metabolites such as 3-CA, aniline and 4-aminobenzoate.[16] Cooper et al. (2015)

showed that organohalide respiring *Dehalococcoides* strain CBDB1 could respire and

dechlorinate 2,3-DCA (but not 2,4- or 2,6-DCA) with H_2 as electron donor.[17] In a follow up

study contrasting dechlorinating mechanisms between *Dehalococcoides mccartyi* strain CBDB1

and *Dehalobacter* sp. strain 14DCB1 previously isolated on chlorinated benzenes, the authors

found that both strains could transform 2,3-DCA and 3,4-DCA with hydrogen as the electron

donor producing 3-CA (by CBDB1) or very minor amounts of 2-chloroaniline (2-CA, by

14DCB1) and 3-CA (by 14DCB1).[18] Despite these advances, no studies have focussed on

achieving complete and stoichiometric anaerobic transformation of DCAs to aniline at

concentrations relevant to those observed at a highly contaminated site[19,20] with average total

DCA concentrations above 200 µM in groundwater. Furthermore, although *Dehalobacter* has

been shown to transform a variety of halogenated organic contaminants (e.g., chlorinated

ethanes, toluene, benzenes, and chloroform),[21–23] reproducible and robust microbial kinetics of

Dehalobacter-mediated dechlorination are lacking, limiting the predictability and quantification

of natural or bioaugmented dechlorination at contaminated sites. The interactions between

Dehalobacter and other microorganisms in dechlorinating microbial consortia also remain poorly

understood. In a companion study, subtle aniline formation was observed after 2,3-DCA

dechlorination to 2-CA in culture 23DCA-T2.[24] We made a transfer from this culture and the

resulting subcultures surprisingly exhibited significantly more aniline formation. This prompted

us to further explore dechlorination of 2,3-DCA to aniline and *Dehalobacter* kinetics. The

objectives of our study were to: (1) demonstrate the complete anaerobic biotransformation of

2,3-DCA to aniline at a high initial concentration; (2) identify the associated transformation

pathway; (3) determine growth yield and kinetic parameters to improve the predictability of dechlorination and (4) investigate syntrophic interactions between cocultured microorganisms and *Dehalobacter*. These objectives were achieved by frequent (every 3 days) sampling of chemical and microbial concentrations in stable enrichment cultures derived from a contaminated site.

5 MATERIALS AND METHODS

5.1 Chemicals

All chemicals were obtained from Sigma Aldrich (St. Louis, Missouri, USA) unless otherwise noted. Cultures were amended with 2,3-DCA (99.6% purity). Aniline and 2-CA used as standards were both at 99.5% purity; whereas, 3-CA and 4-chloroaniline (4-CA) were 99% and 98% pure, respectively. Ethanol was HPLC grade. Sodium lactate (60% by mass) was purchased from Spectrum Chemical MFG Corp.

5.2 Microbial Enrichment Cultures

The culture used as an inoculum in this study was the third consecutive subculture (referred to as WANG-23DCA-T3) from an anaerobic methanogenic 2,3-DCA-dechlorinating enrichment culture (i.e., 23DCA-T2), as described in Araújo et al. (2025).[24] In brief, 23DCA-T2 originated from microcosms established in 2015 using groundwater and soil samples collected from a heavily contaminated industrial site in Brazil.[24] On May 4, 2022, the first transfer (WANG-23DCA-T1) was prepared by inoculating 90 mL of anaerobic medium with 10 mL 23DCA-T2 sample. After three-month cultivation, on August 9, 2022, a 50 mL portion of culture WANG-23DCA-T1 was transferred to establish a new subculture (WANG-23DCA-T2). On day 188 of WANG-23DCA-T2 (February 13, 2023), a 3% transfer of culture (12 mL) from WANG-

23DCA-T2 was made to establish subculture WANG-23-DCA-T3 (400 mL). All three

subcultures were repeatedly amended with 2,3-DCA and demonstrated activity in transforming

it, with WANG-23-DCA-T3 showing particularly strong conversion of 2,3-DCA to aniline.

These subcultures were topped up with medium multiple times to build up culture volume for

future experiments. On day 254 of WANG-23-DCA-T3 (October 25, 2023), the inoculum for

this study was taken from WANG-23-DCA-T3. Dechlorination profiles and associated

cultivation conditions for subcultures and transfers are detailed in Supporting Information (SI)

Section S1 and Figure S1-S3.

131 5.3 Experimental Setup

In this study, all experiments were performed in an anaerobic glovebox (Vinyl Anaerobic

Chamber glovebox, Coy Laboratory Products) at room temperature (21-22°C). The glovebox

was filled with a gas mix of 10% H_2, 10% CO_2 and 80% N_2, however owing to reaction with

diffusing oxygen, the H_2 concentration in the glovebox remained around 1-1.5% H_2.

To establish the biotransformation kinetics, four replicates with an initial concentration of 40

mg/L (i.e., 250 μM) 2,3-DCA and 3% (v/v) inoculum were prepared in autoclaved 250-mL

Boston round glass bottles sealed with Mininert valves, with an initial medium volume of 210

mL and headspace volume of 40 mL. They are referred to as "biotic bottles". The medium used

in this study was a pre-reduced mineral medium described in Edwards and Grbic-Galić (1992).[25]

To estimate the decay kinetics of *Dehalobacter* under conditions lacking electron acceptors,

triplicate decay bottles were established in the same manner as the biotic bottles, excluding the

addition of 2,3-DCA. To monitor the mass loss due to sampling and control for the possibility of

abiotic transformation processes, triplicate abiotic controls were set up similarly without

inoculation in autoclaved 250-mL Boston round bottles, each containing 210 mL of mineral

medium spiked at 40 mg/L of 2,3-DCA. Biotic and abiotic bottles were amended with 2,3-DCA

by dropping a feeding vial containing 2,3-DCA into the bottle. Details are provided in Section

S2. After spiking with 2,3-DCA, the concentration of 2,3-DCA was regularly monitored in all

bottles. Inoculation of the biotic bottles was not conducted until the concentration of 2,3-DCA

had stabilized at the target concentration (i.e., 40 mg/L), which took three days. Once complete

dissolution of 2,3-DCA was attained, 6.5 mL of homogenized, active inoculum was added to

four biotic bottles. The biotic and decay bottle were then amended with sodium lactate (0.6 mM)

and ethanol (3.3 mM). Abiotic controls did not receive any amendment of electron donors. The

inoculation defined the time zero of the experiment. Samples were collected immediately after

inoculation to characterize the initial concentration of methane (headspace), 2,3-DCA,

transformation products, volatile fatty acids, inorganic anions, *Dehalobacter* abundance, and

microbial community composition. During the experiment, all the bottles were covered with

black cloth to minimize exposure to light and possible photodegradation.

Aqueous samples (350 µL) were taken from biotic and abiotic bottles approximately three times

every week for chloroaniline and aniline analysis. A 4-mL sample was collected for DNA

extraction from biotic and decay bottles roughly every ten days. Samples (500 µL) for

quantifying inorganic anions and volatile fatty acids were taken weekly from biotic and abiotic

bottles. Headspace samples (300 µL) for determining methane production were withdrawn

approximately once every ten days from the biotic bottles. The total liquid volume withdrawn

did not exceed 20% of the initial volume during the first 50 days (i.e., dechlorination of 2,3-DCA

to 2-CA and 3-CA). Sample collection and processing are detailed in the SI.

5.4 Analytical procedures

Chloroanilines and aniline were measured using high-performance liquid chromatography coupled with a diode array detector (HPLC/DAD, Thermo Scientific Dionex Ultimate 3000 series) equipped with an Accucore C18 column (100 mm × 2.1 mm × 2.6 μm) and guard column (10 mm × 2.1 mm × 2.6 μm; Thermo Scientific, Waltham, MA, USA). Methane and other possible volatiles (e.g., benzene) were measured by a Hewlett-Packard 5890 Series II gas chromatography (GC) using a GSQ I. D. PLOT column (30 m × 0.53 mm; J & W Scientific, Folsom, USA) configured with a flame ionization detector (FID). Inorganic anions (sulfate, phosphate, nitrate, nitrite, and chloride) and volatile fatty acids (acetate, propionate, butyrate, lactate, formate, and pyruvate) were measured by ion chromatography (IC) using a Dionex™ Integrion™ high pressure IC coupled with an ASRS 500 suppressor and a Dionex™ AS-DV autosampler (Thermo Fisher Scientific). Separation was achieved using a Dionex™ IonPac™ 2 × 250 mm AS11-HC column. Analytical method details are provided in Sections S3-S5.

5.5 DNA sampling and extraction

Culture samples (4 mL) were withdrawn for DNA extraction and centrifuged (described in Section S6), supernatant discarded, and the cell pellets were stored at -80 °C. DNA was extracted using the DNeasy PowerSoil Pro kit (Qiagen) following manufacturer's guidelines. DNA was eluted in 50 μL of 10 mM Tris-HCl buffer and stored at -80 °C until analysis.

5.6 Quantitative PCR analysis

Quantitative polymerase chain reaction (qPCR) primer sets targeting *Dehalobacter* (Dhb) were used as previously described.[26] Primer sequences were Dhb477f (5'-GATTGACGGTACCTAACGAGG-3') and Dhb647r (5'-TACAGTTTCCAATGCTTTACG-

189 3').[26] Primers targeting most bacteria (Bac) were as follows: Bac1055f (5'-

190 ATGGCTGTCGTCAGCT-3') and Bac1392r (5'-ACGGGCGGTGTGTAC-3').[27] Temperature

191 profiles and reaction details are available in Section S7. The qPCR results with *Dehalobacter*

192 biomass estimation are provided in Table S1.

5.7 16S rRNA gene amplicon sequencing

194 16S rRNA gene amplicon sequencing targeting the V6-V8 region of bacteria and archaea was

195 carried out using the Illumina NovaSeq PE300 platform at Genome Québec (Montreal, QC,

196 Canada), using a set of modified "staggered end" primers: forward primer 926f (5'-

197 AAACTYAAAKGAATWGRCGG-3') and reverse primer 1392r (5'-ACGGGCGGTGWGTRC-

198 3', as previously described.[28] Original amplicon reads were processed using QIIME 2 (v.2021.2)

199 (Section S8).[29] Amplicon sequence variants (ASVs) were generated using DADA2 plugin [30] and

200 taxonomically classified with the SSU SILVA v.138 database.[31] Relative abundance of ASVs in

201 bacterial (Table S2) and archaeal (Table S3) communities were calculated. Sequence alignment

202 for *Dehalobacter* ASVs was performed in Geneious Prime (v.2023.1.2) using the MUSCLE

203 Alignment feature. Original sequence reads are available through NCBI (Accession:

204 PRJNA1040917).

5.8 Modeling dechlorination of 2,3-DCA using Monod kinetics

206 We applied the Monod equations[32,33] to estimate the kinetic parameters for *Dehalobacter* growth

207 on 2,3-DCA with two coupled non-linear differential equations that describe the growth of

208 *Dehalobacter* (Eq. 1) and the associated consumption of the limiting substrate (Eq. 2):

209
$$\frac{dX}{dt} = \mu_{max} X \left(\frac{S}{S + K_s} \right) - k_d X$$

210                               *Equation (1)*

211
$$-\frac{dS}{dt} = \frac{\mu_{max}X}{Y}\left(\frac{S}{S + K_s}\right)$$

212
Equation (2)

where S and X represent concentrations (g/L) of the substrate (2,3-DCA or 2-CA) and active

biomass (*Dehalobacter*) at time t, respectively; Ks is the half-saturation constant (g/L), and μ_{max}

is the maximum specific growth rate (day^{-1}); Y is the growth yield (g biomass /g substrate); k_d is

the first-order decay constant (day^{-1}) for biomass; dX/dt is the growth rate of biomass (g/L/day),

and -dS/dt is the substrate consumption rate (g/L/day).

Yield and decay parameters Y and k_d were independently determined in this study. Therefore, Ks

and μ_{max} were the only two Monod parameters obtained through the least squares optimization.

The time-course substrate depletion data were used to fit the Monod model. The coupled

differential equations were numerically solved using the "solve_ivp" function from the

"scipy.integrate" module in Python, with initial conditions set to the substrate and biomass

concentrations measured at time zero (S_0 and X_0, respectively). S_0 was well constrained, but X_0

being a less accurate measurement was allowed to vary within a narrow range during

optimization. The optimal initial biomass for modelling is referred to as X_0'. The combined

residual sum of squares (RSS) from all replicates was minimized. We employed the

"least_squares" function from the "scipy.optimize" module in Python to achieve the best fit by

adjusting optimization parameters, Ks, μ_{max}, and X_0' for 2,3-DCA dechlorination. The starting

biomass used for modeling the subsequent 2-CA dechlorination was calculated from the kinetics

of the preceding 2,3-DCA dechlorination step. Thus, for 2-CA dechlorination kinetics, only Ks

and μ_{max} were adjusted to minimize RSS. To accommodate the determination of Ks and μ_{max},

broad parameter ranges were applied ([0,1000] mg/L for Ks and [0,10] day^{-1} for μ_{max}). A range of

[X_0/2, 4X_0] for X_0' was set to account for potential estimation errors associated with the initial

biomass measurement using qPCR. The goodness-of-fit was assessed using RSS, reduced chi-

square, and root-mean-square errors (RMSE).

6 RESULTS AND DISCUSSION

6.1 Complete dechlorination of 2,3-DCA to aniline

We observed complete and stoichiometric transformation of 2,3-DCA to aniline via 2-CA in all

four biotic replicates. The data for Biotic Bottle I is shown in Figure 1a, with the results for the

other replicates available in Figure S4. In the abiotic controls, 2,3-DCA concentration remained

constant throughout the course of the experiment (Figure 1a) and no transformation products

were detected.

In the biotic bottles, between days 0 and 53, 2,3-DCA underwent quantitative dechlorination

without a significant lag phase (< 2 days), resulting in the formation of 2-CA and 3-CA at an

overall stoichiometric ratio of 7.6 ($\pm$ 1.9): 1 (n=88 using all time points). Our observations

indicate that *Dehalobacter* preferentially removes the chlorine substituent at the meta position

relative to the amino group to produce 2-CA. This finding aligns with a previous study showing

the dechlorination of 2,3-DCA to 2-CA by *Dehalobacter* Strain 14DCB1.[18] However, in that

study, the formation of 2-CA was minimal (<0.8 µM) over five successive 2,3-DCA feeding

cycles at 40 µM across 35 days.[18] In contrast, we observed a 1:1 stoichiometric conversion of

2,3-DCA to CAs (mainly 2-CA). A mole balance analysis for each biotic replicate showed that

92 ($\pm$ 4) % (n=4) of the reduction in 2,3-DCA can be explained by the production of

dechlorination intermediates 2-CA and 3-CA over the course of 50 days.

After day 50, 2-CA was further dechlorinated to aniline, whereas 3-CA persisted (Figures 1a and

S4). End products of this two-step dechlorination were predominantly aniline (89%), with minor

amounts of 3-CA. On day 103, complete biotransformation of 2-CA to aniline was observed. 2-

CA dechlorination is also a quantitative process as 88 (± 6) % (n=3) of 2-CA consumed can be

attributed to the corresponding increase in the aniline concentration. Dechlorination of 2,3-DCA

to aniline is a sequential process. Dechlorination of 2-CA to aniline did not start until 2,3-DCA

was depleted in the bottles. This observation aligns with the results from the parent culture

(Figure S3), which shows that aniline formation began when 2-CA concentration was much

higher than 2,3-DCA concentration.

The preferential utilization of 2,3-DCA as an electron acceptor by *Dehalobacter* over 2-CA is

consistent with the reported Gibbs free energy for dechlorination of 2,3-DCA to either 2-CA (-

135.0 kJ/reaction) or 3-CA (-137.7 kJ/reaction), which are more thermodynamically favorable

than the dechlorination of 2-CA to aniline (-126.8 kJ/reaction).[34] Moreover, this preferential

utilization may explain the negligible aniline formation in the earlier subcultures (Figures S1 and

S2), where 2,3-DCA was re-amended before its depletion and consistently available to

Dehalobacter.

6.2   *Dehalobacter* is responsible for dechlorination

A continuous and consistent rise in *Dehalobacter* absolute abundance (16S rRNA gene copies/L)

was observed across all biotic replicates during both the 2,3-DCA and 2-CA biotransformation

stages (Figure 1c). Biotic bottle II showed slightly lower *Dehalobacter* growth during 2-CA

dechlorination compared to other replicates, but this aligns with the limited biotransformation of

2-CA to aniline in this bottle (Figure S4II), confirming the correlation between 2-CA

dechlorination and *Dehalobacter* growth. The results also indicate that *Dehalobacter* can be

enriched through the supply of 2,3-DCA and 2-CA (Figure 1.e). From days 12 to 87, the relative

abundance of *Dehalobacter* in the bacterial community increased by 46.9 ± 8.4% (n=4),

becoming the most dominant bacterial genus. In line with stoichiometry, the results demonstrate

that *Dehalobacter* can grow on 2,3-DCA and 2-CA through reductive dechlorination. Two

predominant ASVs (ASV3 and ASV4) accounted for 99.9 ± 0.1% (n=32) of total *Dehalobacter*

reads (Table S2). *Dehalobacter* ASV3 and ASV4 shared >99.78% 16S rRNA gene sequence

identity (differing by one base pair in the 468-bp amplicon) and exhibited consistent abundance

trends throughout the experiment, maintaining a stable ASV3/ASV4 ratio of 3.1 ± 0.1:1 (n=32)

in all the samples. Therefore, the two ASVs likely belong to the same *Dehalobacter* strain.

Dehalobacter ASV3 and ASV4 were not the dominant *Dehalobacter* ASVs (ASV1 and ASV2)

in the original parent culture (23DCA-T2)[24] but the ASVs were present in other enrichment

cultures stemming from the original site material. With approximately 97% sequence identity

(Table S4), both ASV3 and ASV4 are genetically distinct from the dominant ASV1 and ASV2

found in 23DCA-T2,[24] where slow and low-level aniline formation was noticed. While these

genetic differences may account for the distinct capability of using 2-CA as an electron acceptor

to yield aniline, future research should investigate this further.

Figure 1. (a) Dechlorination of 2,3-DCA to aniline via 2-CA in biotic replicate I.

Concentrations (μmol/L) of 2,3-dichloroaniline (2,3-DCA, red squares), 2-chloroaniline (2-

CA, blue triangles), 3-chloroaniline (3-CA, green triangles), and aniline (orange diamonds)

over the experimental time (days). 4-CAwas not detected. The average 2,3-DCA

concentrations of triplicate abiotic controls are represented by black circles. (b)

Concentration (mmol/L) of lactate (purple pentagons), acetate (red circles), propionate

(blue squares), and formate (light teal triangles) in replicate I. Butyrate and pyruvate were

not detected and fatty acid concentrations in abiotic controls remained below the LOQ

(Table S5c). (c) Increase in *Dehalobacter* absolute abundance (16S rRNA gene copies/L) in

all biotic replicates. The absolute abundance was measured using qPCR with primers

targeting *Dehalobacter*. (d) Average methane production (µmol per bottle) of all biotic

replicates. Error bars represent the standard deviation among the four biotic replicates. (e)

Changes in bacterial composition during dechlorination in replicate I. The relative

abundance of bacteria at the genus level (unless otherwise specified) obtained from 16S

rRNA amplicon sequencing is presented in combination with total bacterial concentrations

(gene copies/L) determined using qPCR targeting general bacteria. Others: ASVs with

relative abundance < 1%. Results from all four biotic replicates are provided in Figures S4,

S5, and S6a.

6.3 Growth yields and microbial kinetics of *Dehalobacter*

6.3.1 Growth yields and first-order decay

Through the two-step dechlorination process, we measured *Dehalobacter* growth yields on 2,3-DCA and 2-CA (Figure 2). The growth yield for 2,3-DCA dechlorination was $1.2 \pm 0.1 \times 10^8$ 16S rRNA gene copies/μmol 2,3-DCA converted (Figure 2a). Following the depletion of 2,3-DCA, *Dehalobacter* continued to grow on 2-CA, with a growth yield of $1.3 \pm 0.1 \times 10^8$ 16S rRNA gene copies/μmol aniline formed (Figure 2b). An extensive review of *Dehalobacter* growth yields measured from various chlorinated organic compounds was conducted previously.[24] For an easier comparison with other reported values, measured yields from this study are converted into the same units (Table S7). Measured yields from this study are within the range reported previously for 2,3-DCA and 3,4-DCA,[24] and are close to those previously reported in the literature for other substrates including chlorinated ethanes and benzenes,[24] falling on the higher end within one order of magnitude. Many studies underestimate the true yield because the impact of cell decay is not considered. This impact becomes more significant when data are analyzed over long-term cultivation (e.g., hundreds of days) and with a limited number of samples. In this study, measurements of cell abundance were frequent and rates were generally faster than many published studies, improving the estimate of yield. Moreover, this study is the first to demonstrate the decay of *Dehalobacter* under stress conditions in the absence of chlorinated organic compounds. After 62 days of starvation, the *Dehalobacter* concentration in the triplicate decay bottles decreased to 35% of its initial level (Figure 2c). The first-order decay constant was calculated as 0.017 ± 0.001 (n=3) day^{-1}, corresponding to a half-life of 41 days for *Dehalobacter* without chlorinated electron acceptors.

We converted the measured yield into mass units (g/g) assuming a mass per cell and 4 16S rRNA

gene copies per cell, yielding $1.7 \pm 0.2 \times 10^{-2}$ g *Dehalobacter*/ g 2,3-DCA and $2.4 \pm 0.2 \times 10^{-2}$ g

Dehalobacter/ g 2-CA. The theoretical maximum growth yields were also calculated applying

the McCarty TEEM model based on microbial energetics,[35,36] assuming hydrogen as the sole

electron donor, ammonium as the sole nitrogen source, and acetate as the sole carbon source

(Section S9). The predicted yields at various energy transfer efficiencies, along with the

measured yields are compared in Table S8. The thermodynamically predicted yields of

Dehalobacter, matched the experimental measurements when the energy transfer efficiency was

between 45% and 50% (Table S8). The consistency between predicted and measured values

gives confidence in the accuracy of the measured yields. The energy transfer efficiency range (45

– 50%) for *Dehalobacter* used in this study are consistent with the reported range (45 – 65%) for

anaerobic heterotrophic growth.[36]

Figure 2. Growth yields (panels a and b) and decay constant (panel c) for *Dehalobacter*

(*Dhb*). Growth yields (Y) growing on 2,3-DCA and 2-CA are determined using Ordinary

Least Squares regression. Linear regressions are evaluated in Table S6. In panels a and b,

solid red lines represent the fitted linear regression, with shaded areas corresponding to the

95% confidence interval. *Dhb* growth (i.e., X_t-X_0) denotes the total increase in qPCR-

determined *Dehalobacter* absolute abundance at each sampling time relative to the initial

abundance at the start of each dechlorination stage. 2,3-DCA consumption (i.e., S_0-S_t)

represents the cumulative substrate (2,3-DCA) consumption at time t during 2,3-DCA

dechlorination, while aniline production (i.e., P_t-P_0) indicates the cumulative product

(aniline) formation at time t during 2-CA dechlorination. SE: standard errors associated

with the slope. The red line in panel c is the fitted first-order decay model, with dashed

lines representing the 95 % confidence interval. Error bars represent the standard

deviation in normalized *Dehalobacter* concentration (i.e., C_t/C_0) across decay bottles.

6.3.2 Modeling Monod kinetics to estimate K_s and μ_{max}

With the experimentally determined growth yields and decay constant, kinetics of dechlorination

of chloroanilines can be modelled to determine K_s and μ_{max} using the Monod equation, assuming

electron donor is not limiting. Optimal Monod constants and model evaluation are presented in

Table S9. In the case of *Dehalobacter* growth on 2,3-DCA, K_s and μ_{max} are determined as 45 ±

16 mg/L and 0.18 ± 0.03 day^{-1}, respectively. Due to ample initial electron donor amendment,

electron donor limitation unlikely occurred during 2,3-DCA dechlorination. Reproducible trends

of 2,3-DCA dechlorination were observed across all four replicates, and the minor variations

between replicates can be explained by slight differences (±25%) in the initial biomass (Figure

2). The optimal μ_{max} results in a doubling time for *Dehalobacter* growing on 2,3-DCA without

donor limitation ranging from 3.3 to 4.6 days. This range aligns with the doubling time estimated

using measured *Dehalobacter* abundance in the initial stage (days 0 to 12), which was 3.3 days.

For 2-CA dechlorination, similar constants (K_s = 35 ± 24 mg/L and μ_{max} = 0.14 ± 0.06 day^{-1})

were found using data from bottles I and IV, exhibiting complete dechlorination. Comparable

Monod constants were found for two dechlorination steps, as 2,3-DCA and 2-CA have the

similar chemical structure. The μ_{max} of *Dehalobacter* during 2-CA dechlorination was slightly

lower than during 2,3-DCA dechlorination, aligning with the thermodynamic predictions that

2,3-DCA dechlorination can yield more energy (Section S9). However, kinetics of 2-CA

dechlorination varied among individual replicates. Biotic bottles II and III showed that 2-CA

dechlorination had much slow kinetics, which may indicate that electron donor-limiting

conditions were reached during 2-CA dechlorination, as no additional donor was added after

T=0.

Figure 3. Monod kinetics of dechlorination of 2,3-DCA to aniline. Three simulations were

performed at different initial biomass of *Dehalobacter* (75% X_0', 100% X_0', and 125% X_0'),

using the measured yields and decay constant, along with the optimal values for K_s and

μ_{max}. Red squares and orange diamonds represent the observed depletion data of 2,3-DCA

(from all four biotic replicates) and 2-CA (from biotic replicates I and IV), respectively,

with error bars showing the standard deviation across the replicates.

6.4 Roles of other microbes in the consortium

Electron distribution and balances combining results from analyses of chloroanilines, fatty acids, and methane production were investigated to elucidate electron flows and syntrophic relationships within the anaerobic dechlorinating microbial consortium (Figures 4 and S7). This analysis is summarized in Section S11. Other than for biotic replicate IV that exhibited strangely slow acetogenesis, a good electron balance was achieved across replicates, at 84 ± 17 %.

Figure 4. Syntrophic relationships among representative microorganisms in the anaerobic dechlorinating microbial consortium (panel a) and temporal dynamics of major microbiological processes during the experiment (panel b). Electrons available for the whole system originated from the ethanol and lactate. Fermenting bacteria, possibly ***Clostridium sensu stricto 7*, *Sporomusa*, and *Sedimentibacter*, consumed ethanol, releasing electrons in the form of hydrogen gas (yellow), which supported both 2,3-DCA**

dechlorination (teal) and hydrogenotrophic methanogenesis (navy blue). However, a

fraction of ethanol was converted to acetate by *Sporomusa* (red) rather than to hydrogen.

In contrast, lactate was not initially used for hydrogen formation; instead, *Clostridium*

***sensu stricto 7* converted most of the amended lactate into acetate and propionate (orange).**

Subsequently, *Pelotomaculum* oxidized propionate to release hydrogen (purple), supporting

2-CA dechlorination (green) and the late-stage hydrogenotrophic methanogenesis (blue).

6.4.1 Lactate fermentation and acetogenesis from ethanol to produce acetate and hydrogen for *Dehalobacter*

We observed significant changes in the concentrations of volatile fatty acids between days 0 and

5 (Figures 1b and S5), especially production of acetate and propionate from added donors,

ethanol and lactate, indicating active fermentative processes early in the experiment. This finding

was also supported by 16S rRNA amplicon sequencing and total bacterial concentration data,

which revealed a notable growth of all fermenting bacteria, primarily *Clostridium sensu stricto 7*

and *Sporomusa*, dominating the bacterial community in the initial stage of the process (Figure 1e

and S6a). Additionally, methane production was observed during the same period (between day

0 and day 10, Figure 1d), indicating that methanogenesis aligned with fermentation. Microbial

data suggests that fermentative processes were hydrogen-yielding, as the dominant methanogenic

genus, *Methanobacterium*, is hydrogenotrophic (Figure S8).

Between days 0 and 5, lactate concentration, initially at 0.6 mM, decreased below the LOQ while

acetate, propionate, and formate concentrations increased (Figure 1e). Ethanol initially at 3.3

mM was presumably also depleted in the first 5 days because the electron balance is preserved

(Figure S7). The stoichiometry ratio of lactate consumed to propionate formed was $2.9 \pm 0.2 : 2$

(n=4; Section S10) for biotic replicates. This ratio is consistent with the theoretical stoichiometry

ratio of lactate: propionate: acetate = 3: 2: 1 during lactate fermentation to propionate and acetate

(Section S10).[37,38] Therefore, this observed ratio confirms that most of the lactate was fermented

to propionate and acetate, and that propionate was solely associated with lactate fermentation.

The identified *Clostridium sensu stricto 7* was likely responsible for lactate fermentation since

species of *Clostridium* were reported to convert lactate to propionate in a ratio of 3:2.[43,44]

Fermentative bacteria can alter their metabolic pathways to produce hydrogen in response to a

hydrogen demand.[41] *Clostridium sensu stricto* has also been documented to produce hydrogen

via fermentation.[42–44] Therefore, *Clostridium sensu stricto 7* along with other identified

fermenters, such as *Sedimentibacter* and *Sporomusa*, might be involved in hydrogen-yielding

fermentation from ethanol and fatty acids. Partnering with hydrogen consumers such as

Dehalobacter and methanogens, fermenting bacteria were able to oxidize ethanol to acetate, with

hydrogen gas as a product.[45–47] The electron balance results indicate that the electron demand for

complete dechlorination is negligibly low compared to the electrons supplied by ethanol (Figure

S7b). Consequently, the hydrogen demand of *Dehalobacter* can be readily met through

hydrogen-yielding fermentation of ethanol.

The ratio of initial ethanol addition to measured acetate production, excluding acetate

contribution from lactate fermentation, was calculated as 1.2 ± 0.4 (n=4). This stoichiometry is

close to fermentation of ethanol to acetate and hydrogen (ethanol: acetate = 1:1)[41] and

acetogenesis from ethanol via the Wood-Ljungdahl pathway (ethanol: acetate = 1:1.5)[48].

Fermentation reactions are detailed in Section S10. The stoichiometry indicates hydrogen

production, and measured products suggest the coexistence of hydrogen-, formate- and non-

hydrogen-yielding acetogenesis during fermentation. Acetogenesis was mediated by *Sporomusa*,

as indicated by a significant increase in their abundance (Figure 1e). *Sporomusa*, a known

homoacetogen, can consume ethanol and CO_2 to produce acetate through the Wood-Ljungdahl

pathway[48]. Furthermore, the electron balance results suggested that only small fraction of

electrons from ethanol was used for hydrogen production, as the majority was converted into

acetate (Figure S7b).

The acetate pattern in biotic bottle IV differed noticeably from the other replicates, with a slow

increase in acetate over the course of the experiment, whereas the other replicates showed an

increase mainly within the first 10 days (Figures S5 and S7). This discrepancy on acetate

production suggests that biotic bottle IV received an erroneously small amount of ethanol at time

zero, potentially due to errors associated with syringe addition. Surprisingly, this discrepancy

allows us to identify *Sporomusa* as the primary acetate producer. The limited acetogenesis from

ethanol in biotic bottle IV significantly impacted its bacterial composition. In biotic bottle IV,

Sporomusa was much less abundant in the early stages of the experiment compared to the other

replicates, but its abundance gradually increased as the acetate production occurred later (Figure

S7). This clear correlation between acetate production and *Sporomusa* growth confirms that

Sporomusa is responsible for acetogenesis.

Apart from biotic bottle IV, no notable increase in acetate and propionate was observed after day

10, indicating that lactate fermentation and acetogenesis were much faster processes compared to

reductive dechlorination, and were primarily active during the early stages of 2,3-DCA

dechlorination (Figure S5). This is further supported by the amplicon sequencing results, which

revealed a continuous and significant decline of *Clostridium sensu stricto 7* and *Sporomusa* as

the experiment progressed. Beyond day 10, fermentation reactions were much slower owing to

lack of readily fermentable substrates. Once amended lactate and ethanol were depleted, the

abundance of dominant fermenting bacteria decreased, while *Dehalobacter* became the most

abundant bacterial genus (Figures 1e and S6a).

A variety of fermenting bacteria are involved in interspecies electron transfer within the

microbial consortium. They converted fatty acids and alcohols, which *Dehalobacter* cannot

utilize for energy and cell synthesis, into hydrogen, formate, and acetate, all of which supported

dehalorespiration and growth by *Dehalobacter*. Previous studies have reported *Dehalobacter*

relies on hydrogen or formate as the electron donors for anaerobic respiration[49–51] and acetate as

a carbon source.[52] In this work, acetate uptake into biomass by *Dehalobacter*, estimated from the

overall energetic equation (Section S9), was small (< 0.1 mM) and not observable amidst the high

acetate levels.

6.4.2 Syntrophic propionate oxidation to provide hydrogen and acetate for *Dehalobacter*

During 2,3-DCA dechlorination to monochloroanilines, concentrations of both acetate and

propionate, the primary products from fermentation, were steady in the bottles. However,

following the depletion of 2,3-DCA and initiation of dechlorination of 2-CA to aniline,

propionate levels began to decrease (days $50 - 95$), while acetate concentrations exhibited

minimal decline (Figure 1b). A moderate amount of methane was also produced over the same

period, following a phase of negligible methanogenesis throughout 2,3-DCA dechlorination

(Figure 1d). This late-stage methanogenesis can be attributed to the decrease in propionate

concentrations (biotic bottles I, II, and IV), with a recovery of 78 ($\pm$ 11) % (n=3), calculated as

the electron equivalent ratio of the methane production to the reduction in propionate. Moreover,

this late-stage methanogenesis was associated with a rise in the bacterial abundance of

Pelotomaculum (Figure 1e). These are syntrophic propionate-oxidizing bacteria that are

obligately co-cocultured with methanogens.[53,54] Thus experimental evidence indicates that

propionate was consumed by *Pelotomaculum* via syntrophic oxidation. Subsequently, the

507 products from syntrophic propionate oxidation, formate and hydrogen, were used by

508 hydrogenotrophic methanogens to form methane and by *Dehalobacter* for 2-CA dechlorination.

509 The syntrophic oxidation of propionate had an important role in this anaerobic dechlorinating

510 system. Lactate fermentation observed in the initial stage of the experiment produced propionate

511 and acetate, which was not a hydrogen-yielding process. In addition, propionate cannot be

512 directly utilized by *Dehalobacter* or methanogens and therefore built up (days 10 – 50). Thanks

513 to *Pelotomaculum*, propionate was metabolized into formate and hydrogen, thus enhancing the

514 electron transfer efficiency from the amended lactate to hydrogen gas and keeping the propionate

515 concentration low.

516 Additionally, our results clearly demonstrate that syntrophic propionate oxidation and associated

517 hydrogenotrophic methanogenesis did not commence until the depletion of 2,3-DCA (Figure 1).

518 This relatively late onset of syntrophic propionate oxidation may result from the toxicity of 2,3-

519 DCA on *Pelotomaculum* or methanogens. Syntrophic propionate oxidation is energetically

520 unfavourable unless coupled with effective hydrogen-consuming processes (e.g., dechlorination

521 and methanogenesis) to maintain low hydrogen patrial pressures.[46] During 2,3-DCA

522 dechlorination, *Dehalobacter* likely outcompeted methanogens for hydrogen, as shown by the

523 consistent and reproducible dechlorination profiles, while methane production remained

524 minimal. However, the hydrogen demand for dechlorination, as previously discussed, was too

525 low to trigger the syntrophic propionate oxidation.

526 6.4.3 Acetate accumulation and potential inhibition of chloroanilines on
527     acetoclastic methanogens

528 The accumulation of acetate in the dechlorinating cultures was unexpected. Rather than methane,

529 most of electrons from added lactate and ethanol ended up in the form of acetate (Figure S7).

530 The acetate accumulation suggests a lack of effective acetate consumer in the community, which

is further confirmed by the 16S rRNA sequencing results (Figure S8). All identified

methanogens, such as *Methanobacterium*, which dominated the Archaea community with a

relative abundance greater than 97%, and *Methanoregula*, are hydrogenotrophic methanogens

that produce methane exclusively from hydrogen and CO_2 and not from acetate.[55,56] Thus, acetate

accumulation was due to the absence of acetoclastic methanogens. The absence of acetoclastic

methanogens suggests they were eliminated by elevated concentrations of chlorinated anilines,

and effectively diluted out of the culture after successive transfers. Previous studies have

reported that chlorinated compounds and amino-substituted aromatics can inhibit conversion of

acetate to methane.[57,58] Excessive acetate not only led to a low electron transfer efficiency from

the amended electron donors to hydrogen gas but may also exhaust the buffering capacity of the

mineral medium, leading to low pH conditions that could harm the microbial system. Thus, for

dechlorinating microbial systems lacking the acetoclastic methanogens, regular culture transfer

can be a preventive measure against the excessive acetate. Moreover, acetate accumulation can

serve as a potential indicator of methanogenesis inhibition in the groundwater.

Environmental Implications

This work presents the complete and stochiometric biotransformation of dichloroaniline to

aniline under anaerobic conditions. Thanks to a unique high-frequency sampling regime, we

estimated kinetic parameters and growth yields that are valuable for modeling the fate and

transport of chloroanilines in groundwater and to quantify *in situ* biotransformation. The

syntrophic interactions between *Dehalobacter* and its microbial community can be used as one

line of evidence to identify natural attenuation of chlorinated anilines at contaminated sites. In

addition, the achievement of complete dechlorination at a high substrate level (40 mg/L) with a

low inoculum (3% v/v) holds promise for the application of biostimulation (with electron

donors) and bioaugmentation strategies to address chloroaniline contamination. This study also

explored interspecies electron transfer as a function of amended electron donors over time,

deciphering some of the dynamics of successive microbial processes, including the issue of

acetate accumulation, offering insights into establishing and maintaining a stable anaerobic

dechlorinating system in culture or *in situ*.

7 ASSOCIATED CONTENT

560 The supporting information is available free of charge. Supplementary descriptions of subculture

561 establishment, analytical methods, and data analysis, along with the relevant figures and tables

562 discussed in the paper (PDF). Additionally, original data provided in spreadsheet format (.xlsx).

8 ACKNOWLEDGMENTS

This research was supported by the Natural Sciences and Engineering Research Council of

Canada (NSERC) Discovery (RGPIN-2015-06460), Alliance (ALLRP 575243-22), and Canada

Research Chair (950-230892) grants to E. Passeport, and a Mitacs (IT29192) grant to both E.

Passeport and E. Edwards. The authors declare no conflicts of interest. This publication is in

accordance with the Brazilian biodiversity law n.º 13.123, from May 20th, 2015. This is non-

profit research and has no commercial intention between the parties. This is a collaboration

project between the parties involved. Access Registration No. A3D9F11.

9 REFERENCES

(1) Rebelo, D.; Antunes, S. C.; Rodrigues, S. The Silent Threat: Exploring the Ecological and
Ecotoxicological Impacts of Chlorinated Aniline Derivatives and the Metabolites on the
Aquatic Ecosystem. *J. Xenobiotics* **2023**, *13* (4), 604–614.
https://doi.org/10.3390/jox13040038.

(2) Rasmussen, H. T.; Omelczenko, N.; Friedman, S. K.; McPherson, B. P. Determination of
Chloroanilines in Antibacterial Soaps Using Cation-Exchange Chromatography with UV
Detection. *J. Chromatogr. A* **1996**, *719* (2), 434–437. https://doi.org/10.1016/0021-
9673(95)00731-8.
(3) Vogt, P. F.; Gerulis, J. J. Amines, Aromatic. In *Ullmann's Encyclopedia of Industrial*
Chemistry; Wiley-VCH, Ed.; Wiley, 2000. https://doi.org/10.1002/14356007.a02_037.
(4) Batista, S.; Silva, E.; Galhardo, S.; Viana, P.; José Cerejeira, M. Evaluation of Pesticide
Contamination of Ground Water in Two Agricultural Areas of Portugal. *Int. J. Environ.*
Anal. Chem. **2002**, *82* (8–9), 601–609. https://doi.org/10.1080/03067310290009488.
(5) Berrada, H.; Font, G.; Moltó, J. C. Application of Solid-Phase Microextraction for
Determining Phenylurea Herbicides and Their Homologous Anilines from Vegetables. *J.*
Chromatogr. A **2004**, *1042* (1–2), 9–14. https://doi.org/10.1016/j.chroma.2004.05.017.
(6) Oanh, N. T.; Duc, H. D. Anaerobic Degradation of Propanil in Soil and Sediment Using
Mixed Bacterial Culture. *Curr. Microbiol.* **2021**, *78* (4), 1499–1508.
https://doi.org/10.1007/s00284-021-02419-7.
(7) Ibrahim, M. A.; Zulkifli, S. Z.; Azmai, M. N. A.; Mohamat-Yusuff, F.; Ismail, A. Embryonic
Toxicity of 3,4-Dichloroaniline (3,4-DCA) on Javanese Medaka (Oryzias Javanicus
Bleeker, 1854). *Toxicol. Rep.* **2020**, *7*, 1039–1045.
https://doi.org/10.1016/j.toxrep.2020.08.011.
(8) Philippe, C.; Hautekiet, P.; Grégoir, A. F.; Thoré, E. S. J.; Brendonck, L.; De Boeck, G.;
Pinceel, T. Interactive Effects of 3,4-DCA and Temperature on the Annual Killifish
Nothobranchius Furzeri. *Aquat. Toxicol.* **2019**, *212*, 146–153.
https://doi.org/10.1016/j.aquatox.2019.05.009.
(9) Rodrigues, A. D.; dos Santos Montanholi, A.; Shimabukuro, A. A.; Yonekawa, M. K. A.;
Cassemiro, N. S.; Silva, D. B.; Marchetti, C. R.; Weirich, C. E.; Beatriz, A.; Zanoelo, F. F.;
Marques, M. R.; Giannesi, G. C.; das Neves, S. C.; Oliveira, R. J.; Ruller, R.; de Lima, D.
P.; dos Anjos dos Santos, E. N-Acetylation of Toxic Aromatic Amines by Fungi: Strain
Screening, Cytotoxicity and Genotoxicity Evaluation, and Application in Bioremediation of
3,4-Dichloroaniline. *J. Hazard. Mater.* **2023**, *441*, 129887.
https://doi.org/10.1016/j.jhazmat.2022.129887.
(10) Klečka, G. M.; Carpenter, C. L.; Gonsior, S. J. Biological Transformations of 1,2-
Dichloroethane in Subsurface Soils and Groundwater. *J. Contam. Hydrol.* **1998**, *34* (1–2),
139–154. https://doi.org/10.1016/S0169-7722(98)00096-5.
(11) Major, D. W.; McMaster, M. L.; Cox, E. E.; Edwards, E. A.; Dworatzek, S. M.;
Hendrickson, E. R.; Starr, M. G.; Payne, J. A.; Buonamici, L. W. Field Demonstration of
Successful Bioaugmentation To Achieve Dechlorination of Tetrachloroethene To Ethene.
Environ. Sci. Technol. **2002**, *36* (23), 5106–5116. https://doi.org/10.1021/es0255711.
(12) Gieg, L. M.; Suflita, J. M. Detection of Anaerobic Metabolites of Saturated and Aromatic
Hydrocarbons in Petroleum-Contaminated Aquifers. *Environ. Sci. Technol.* **2002**, *36* (17),
3755–3762. https://doi.org/10.1021/es0205333.
(13) Struijs, J.; Rogers, J. E. Reductive Dehalogenation of Dichloroanilines by Anaerobic
Microorganisms in Fresh and Dichlorophenol-Acclimated Pond Sediment. *Appl. Environ.*
Microbiol. **1989**, *55* (10), 2527–2531. https://doi.org/10.1128/aem.55.10.2527-2531.1989.
(14) Kuhn, E. P.; Suflita, J. M. Sequential Reductive Dehalogenation of Chloroanilines by
Microorganisms from a Methanogenic Aquifer. *Environ. Sci. Technol.* **1989**, *23* (7), 848–
852. https://doi.org/10.1021/es00065a014.

(15) Susarla, S.; Yonezawa, Y.; Masunaga, S. Reductive Transformations of Halogenated
Aromatics in Anaerobic Estuarine Sediment: Kinetics, Products and Pathways. *Water Res.*
1998, *32* (3), 639–648. https://doi.org/10.1016/S0043-1354(97)00265-0.
(16) Duc, H. D.; Oanh, N. T. Anaerobic Degradation of Chloroanilines by Geobacter Sp. KT5.
Curr. Microbiol. **2019**, *76* (2), 248–257. https://doi.org/10.1007/s00284-018-1617-7.
(17) Cooper, M.; Wagner, A.; Wondrousch, D.; Sonntag, F.; Sonnabend, A.; Brehm, M.;
Schüürmann, G.; Adrian, L. Anaerobic Microbial Transformation of Halogenated
Aromatics and Fate Prediction Using Electron Density Modeling. *Environ. Sci. Technol.*
2015, *49* (10), 6018–6028. https://doi.org/10.1021/acs.est.5b00303.
(18) Zhang, S.; Wondrousch, D.; Cooper, M.; Zinder, S. H.; Schüürmann, G.; Adrian, L.
Anaerobic Dehalogenation of Chloroanilines by *Dehalococcoides Mccartyi* Strain CBDB1
and *Dehalobacter* Strain 14DCB1 via Different Pathways as Related to Molecular
Electronic Structure. *Environ. Sci. Technol.* **2017**, *51* (7), 3714–3724.
https://doi.org/10.1021/acs.est.6b05730.
(19) Suzana de Paula Queiroz Kraus. Aerobic and Anaerobic Biotransformation of
Chloroanilines, Chlorobenzenes, and Dichlonitrobenzenes at a Complex Industrial Site in
Brazil and Analysis of Associated Microbial Communities. Master Thesis, University of
Toronto, 2018. http://hdl.handle.net/1807/91484.
(20) Barreto, P.; Lemes, M.; Jimenez, J.; Mack, E. E.; Henderson, J.; Freedman, D. L.
Evaluation of Strategies to Remediate Mixed Wastes at an Industrial Site in Brazil.
Groundw. Monit. Remediat. **2023**, *43* (3), 93–107. https://doi.org/10.1111/gwmr.12607.
(21) Nelson, J. L.; Jiang, J.; Zinder, S. H. Dehalogenation of Chlorobenzenes, Dichlorotoluenes,
and Tetrachloroethene by Three *Dehalobacter* Spp. *Environ. Sci. Technol.* **2014**, *48* (7),
3776–3782. https://doi.org/10.1021/es4044769.
(22) Bulka, O.; Webb, J.; Dworatzek, S.; Mahadevan, R.; Edwards, E. A. A Multifunctional
Dehalobacter? Tandem Chloroform and Dichloromethane Degradation in a Mixed
Microbial Culture. *Environ. Sci. Technol.* **2023**, *57* (48), 19912–19920.
https://doi.org/10.1021/acs.est.3c06686.
(23) Justicia-Leon, S. D.; Higgins, S.; Mack, E. E.; Griffiths, D. R.; Tang, S.; Edwards, E. A.;
Löffler, F. E. Bioaugmentation with Distinct *Dehalobacter* Strains Achieves Chloroform
Detoxification in Microcosms. *Environ. Sci. Technol.* **2014**, *48* (3), 1851–1858.
https://doi.org/10.1021/es403582f.
(24) Sofia P. Araújo; Line Lomheim; Suzana P. Q. Kraus; E. Erin Mack; Jim C. Spain; Sávia
Gavazza; Elizabeth A. Edwards. Dehalobacter Dechlorinates Dichloroanilines and
Contributes to the Natural Attenuation of Dichloronitrobenzenes at a Complex Industrial
Site. *https://www.biorxiv.org/content/10.1101/2025.02.13.638192v1* **2025**.
(25) Edwards, E. A.; Grbić-Galić, D. Complete Mineralization of Benzene by Aquifer
Microorganisms under Strictly Anaerobic Conditions. *Appl. Environ. Microbiol.* **1992**, *58*
(8), 2663–2666. https://doi.org/10.1128/aem.58.8.2663-2666.1992.
(26) Grostern, A.; Edwards, E. A. Growth of Dehalobacter and Dehalococcoides Spp. during
Degradation of Chlorinated Ethanes. *Appl. Environ. Microbiol. 2006, 72, (1)*, 428–436.
(27) Amann, R. I.; Ludwig, W.; Schleifer, K. H. Phylogenetic Identification and in Situ
Detection of Individual Microbial Cells without Cultivation. *Microbiol. Rev.* **1995**, *59* (1),
143–169. https://doi.org/10.1128/mr.59.1.143-169.1995.
(28) Toth, C. R. A.; Luo, F.; Bawa, N.; Webb, J.; Guo, S.; Dworatzek, S.; Edwards, E. A.
Anaerobic Benzene Biodegradation Linked to the Growth of Highly Specific Bacterial

Clades. *Environ. Sci. Technol.* **2021**, *55* (12), 7970–7980. https://doi.org/10.1021/acs.est.1c00508.

(29) Bolyen, E.; Rideout, J. R.; Dillon, M. R.; Bokulich, N. A.; Abnet, C. C.; Al-Ghalith, G. A.; Alexander, H.; Alm, E. J.; Arumugam, M.; Asnicar, F.; Bai, Y.; Bisanz, J. E.; Bittinger, K.; Brejnrod, A.; Brislawn, C. J.; Brown, C. T.; Callahan, B. J.; Caraballo-Rodríguez, A. M.; Chase, J.; Cope, E. K.; Da Silva, R.; Diener, C.; Dorrestein, P. C.; Douglas, G. M.; Durall, D. M.; Duvallet, C.; Edwardson, C. F.; Ernst, M.; Estaki, M.; Fouquier, J.; Gauglitz, J. M.; Gibbons, S. M.; Gibson, D. L.; Gonzalez, A.; Gorlick, K.; Guo, J.; Hillmann, B.; Holmes, S.; Holste, H.; Huttenhower, C.; Huttley, G. A.; Janssen, S.; Jarmusch, A. K.; Jiang, L.; Kaehler, B. D.; Kang, K. B.; Keefe, C. R.; Keim, P.; Kelley, S. T.; Knights, D.; Koester, I.; Kosciolek, T.; Kreps, J.; Langille, M. G. I.; Lee, J.; Ley, R.; Liu, Y.-X.; Loftfield, E.; Lozupone, C.; Maher, M.; Marotz, C.; Martin, B. D.; McDonald, D.; McIver, L. J.; Melnik, A. V.; Metcalf, J. L.; Morgan, S. C.; Morton, J. T.; Naimey, A. T.; Navas-Molina, J. A.; Nothias, L. F.; Orchanian, S. B.; Pearson, T.; Peoples, S. L.; Petras, D.; Preuss, M. L.; Pruesse, E.; Rasmussen, L. B.; Rivers, A.; Robeson, M. S.; Rosenthal, P.; Segata, N.; Shaffer, M.; Shiffer, A.; Sinha, R.; Song, S. J.; Spear, J. R.; Swafford, A. D.; Thompson, L. R.; Torres, P. J.; Trinh, P.; Tripathi, A.; Turnbaugh, P. J.; Ul-Hasan, S.; Van Der Hooft, J. J. J.; Vargas, F.; Vázquez-Baeza, Y.; Vogtmann, E.; Von Hippel, M.; Walters, W.; Wan, Y.; Wang, M.; Warren, J.; Weber, K. C.; Williamson, C. H. D.; Willis, A. D.; Xu, Z. Z.; Zaneveld, J. R.; Zhang, Y.; Zhu, Q.; Knight, R.; Caporaso, J. G. Reproducible, Interactive, Scalable and Extensible Microbiome Data Science Using QIIME 2. *Nat. Biotechnol.* **2019**, *37* (8), 852–857. https://doi.org/10.1038/s41587-019-0209-9.

(30) Callahan, B. J.; McMurdie, P. J.; Rosen, M. J.; Han, A. W.; Johnson, A. J. A.; Holmes, S. P. DADA2: High-Resolution Sample Inference from Illumina Amplicon Data. *Nat. Methods* **2016**, *13* (7), 581–583. https://doi.org/10.1038/nmeth.3869.

(31) Quast, C.; Pruesse, E.; Yilmaz, P.; Gerken, J.; Schweer, T.; Yarza, P.; Peplies, J.; Glöckner, F. O. The SILVA Ribosomal RNA Gene Database Project: Improved Data Processing and Web-Based Tools. *Nucleic Acids Res.* **2012**, *41* (D1), D590–D596. https://doi.org/10.1093/nar/gks1219.

(32) Monod, J. The Growth of Bacterial Cultures. *Ann. Rev. Microbiol* **1949**.

(33) Pirt, S. J. *Principles of Microbe and Cell Cultivation*; Blackwell Scientific, 1975.

(34) Susarla, S.; Masunaga, S.; Yonezawa, Y. Redox Potential as a Parameter To Predict the Reductive Dechlorination Pathway of Chloroanilines in Anaerobic Environments. *Microb. Ecol.* **1997**, *33* (3), 252–256. https://doi.org/10.1007/s002489900028.

(35) Rittmann, B. E.; McCarty, P. L. *Environmental Biotechnology: Principles and Applications*, 2001st ed.; McGraw-Hill Education, 2001.

(36) McCarty, P. L. Thermodynamic Electron Equivalents Model for Bacterial Yield Prediction: Modifications and Comparative Evaluations. *Biotechnol. Bioeng.* **2007**, *97* (2), 377–388. https://doi.org/10.1002/bit.21250.

(37) Madigan, M. T.; Martinko, J. M.; Bender, K. S.; Buckley, D. H.; Stahl, D. A.; Brock, T. *Brock Biology of Microorganisms*, 14th ed.; Pearson Education, 2014.

(38) Seeliger, S.; Janssen, P. H.; Schink, B. Energetics and Kinetics of Lactate Fermentation to Acetate and Propionate via Methylmalonyl-CoA or Acrylyl-CoA. *FEMS Microbiology Letters* **2002**, *211* (1), 65–70.

(39) Leaver, F. W.; Wood, H. G.; Stjernholm, R. THE FERMENTATION OF THREE CARBON SUBSTRATES BY *CLOSTRIDIUM PROPIONICUM* AND

PROPIONIBACTERIUM. *J. Bacteriol.* **1955**, *70* (5), 521–530.
https://doi.org/10.1128/jb.70.5.521-530.1955.

(40) Stams, A. J. M.; Dijkema, C.; Plugge, C. M.; Lens, P. Contribution of 13C-NMR Spectroscopy to the Elucidation of Pathways of Propionate Formation and Degradation in Methanogenic Environments. **1998**.

(41) Fennell, D. E.; Gossett, J. M. Modeling the Production of and Competition for Hydrogen in a Dechlorinating Culture. *Environ. Sci. Technol.* **1998**, *32* (16), 2450–2460. https://doi.org/10.1021/es980136l.

(42) Litti, Yu. V.; Kovalev, D. A.; Kovalev, A. A.; Merkel, A. Yu.; Vishnyakova, A. V.; Russkova, Yu. I.; Nozhevnikova, A. N. Auto-Selection of Microorganisms of Sewage Sludge Used as an Inoculum for Fermentative Hydrogen Production from Different Substrates. *Int. J. Hydrog. Energy* **2021**, *46* (58), 29834–29845. https://doi.org/10.1016/j.ijhydene.2021.06.174.

(43) Tang, T.; Chen, Y.; Liu, M.; Du, Y.; Tan, Y. Effect of pH on the Performance of Hydrogen Production by Dark Fermentation Coupled Denitrification. *Environ. Res.* **2022**, *208*, 112663. https://doi.org/10.1016/j.envres.2021.112663.

(44) Yang, G.; Wang, J. Changes in Microbial Community Structure during Dark Fermentative Hydrogen Production. *Int. J. Hydrog. Energy* **2019**, *44* (47), 25542–25550. https://doi.org/10.1016/j.ijhydene.2019.08.039.

(45) Falkow, S.; Rosenberg, E.; Schleifer, K.-H.; Stackebrandt, E. *The Prokaryotes: Vol. 4: Bacteria: Firmicutes, Cyanobacteria*; Springer Science & Business Media, 2006; Vol. 4.

(46) Schink, B. Energetics of Syntrophic Cooperation in Methanogenic Degradation. *Microbiol. Mol. Biol. Rev.* **1997**, *61* (2), 262–280. https://doi.org/10.1128/mmbr.61.2.262-280.1997.

(47) Schink, B. Anaerobic Digestion: Concepts, Limits and Perspectives. *Water Science and Technology* **2002**, *45* (10), 1–8.

(48) Schuchmann, K.; Müller, V. Energetics and Application of Heterotrophy in Acetogenic Bacteria. *Appl. Environ. Microbiol.* **2016**, *82* (14), 4056–4069. https://doi.org/10.1128/AEM.00882-16.

(49) Fincker, M.; Spormann, A. M. Biochemistry of Catabolic Reductive Dehalogenation. *Annu. Rev. Biochem.* **2017**, *86* (1), 357–386. https://doi.org/10.1146/annurev-biochem-061516-044829.

(50) Wang, S.; Qiu, L.; Liu, X.; Xu, G.; Siegert, M.; Lu, Q.; Juneau, P.; Yu, L.; Liang, D.; He, Z.; Qiu, R. Electron Transport Chains in Organohalide-Respiring Bacteria and Bioremediation Implications. *Biotechnol. Adv.* **2018**, *36* (4), 1194–1206. https://doi.org/10.1016/j.biotechadv.2018.03.018.

(51) Sun, B.; Griffin, B. M.; Ayala-del-Río, H. L.; Hashsham, S. A.; Tiedje, J. M. Microbial Dehalorespiration with 1,1,1-Trichloroethane. *Science* **2002**, *298* (5595), 1023–1025. https://doi.org/10.1126/science.1074675.

(52) Holliger, C.; Hahn, D.; Harmsen, H.; Ludwig, W.; Schumacher, W.; Tindall, B.; Vazquez, F.; Weiss, N.; Zehnder, A. J. B. Dehalobacter Restrictus Gen. Nov. and Sp. Nov., a Strictly Anaerobic Bacterium That Reductively Dechlorinates Tetra- and Trichloroethene in an Anaerobic Respiration. *Arch. Microbiol.* **1998**, *169* (4), 313–321. https://doi.org/10.1007/s002030050577.

(53) Bok, F. A. M.; Harmsen, H. J. M.; Plugge, C. M.; de Vries, M. C.; Akkermans, A. D. L.; de Vos, W. M.; Stams, A. J. M. The First True Obligately Syntrophic Propionate-Oxidizing Bacterium, Pelotomaculum Schinkii Sp. Nov., Co-Cultured with Methanospirillum

Hungatei, and Emended Description of the Genus Pelotomaculum. *Int. J. Syst. Evol.*
Microbiol. **2005**, *55* (4), 1697–1703. https://doi.org/10.1099/ijs.0.02880-0.

(54) Imachi, H.; Sakai, S.; Ohashi, A.; Harada, H.; Hanada, S.; Kamagata, Y.; Sekiguchi, Y.
Pelotomaculum Propionicicum Sp. Nov., an Anaerobic, Mesophilic, Obligately Syntrophic,
Propionate-Oxidizing Bacterium. *Int. J. Syst. Evol. Microbiol.* **2007**, *57* (7), 1487–1492.
https://doi.org/10.1099/ijs.0.64925-0.

(55) Bräuer, S. L.; Cadillo-Quiroz, H.; Ward, R. J.; Yavitt, J. B.; Zinder, S. H. Methanoregula
Boonei Gen. Nov., Sp. Nov., an Acidiphilic Methanogen Isolated from an Acidic Peat Bog.
Int. J. Syst. Evol. Microbiol. **2011**, *61* (1), 45–52. https://doi.org/10.1099/ijs.0.021782-0.

(56) Schonheit, P.; Moll, J.; Thauer, R. K. Nickel, Cobalt, and Molybdenum Requirement for
Growth of Methanobacterium Thermoautotrophicum. *Arch. Microbiol.* **1979**, *123* (1), 105–
107. https://doi.org/10.1007/BF00403508.

(57) Donlon, B. A.; Razo-Flores, E.; Field, J. A.; Lettinga, G. Toxicity of N-Substituted
Aromatics to Acetoclastic Methanogenic Activity in Granular Sludge. *Appl. Environ.*
Microbiol. **1995**, *61* (11), 3889–3893. https://doi.org/10.1128/aem.61.11.3889-3893.1995.

(58) Yang, J.; Speece, R. E. The Effects of Chloroform Toxicity on Methane Fermentation.
Water Res. **1986**, *20* (10), 1273–1279. https://doi.org/10.1016/0043-1354(86)90158-2.