Short title: Mannanases necessary for glucomannan synthesis

Atypical endo-β-1,4-mannanases are necessary for normal
glucomannan synthesis in Arabidopsis

Aina Kikuchi[1, †], Eriko Sato[1, †], Yoshihisa Yoshimi[2,3], Hironori Takasaki[1], Naho
Nishigaki[1], Kimie Atsuzawa[4], Yasuko Kaneko[5], Masatoshi Yamaguchi[1], Daisuke
Takahashi[1], Paul Dupree[2], and Toshihisa Kotake[1,6,*]

Author affiliations:

[1]Graduate School of Science and Engineering, Saitama University, 255 Shimo-Okubo,
Sakura-ku, Saitama 338-8570, Japan

[2]Department of Biochemistry, University of Cambridge, Hopkins Building, The
Downing Site, Tennis Court Road, Cambridge CB2 1QW, UK

[3]Faculty of Biology-Oriented Science and Technology, Kindai University, 930
Nishimitani, Kinokawa 649-6433, Japan

[4]Comprehensive Analysis Center for Science, Saitama University, Saitama 338-8570,
Japan

[5]Faculty of Education, Saitama University, 255 Shimo-Okubo, Sakura-ku, Saitama 338-
8570, Japan

[6]Green Bioscience Research Center, Saitama University, 255 Shimo-Okubo, Sakura-ku,
Saitama 338-8570, Japan

*Author for correspondence: kotake@mail.saitama-u.ac.jp (T. K.)

[†]These authors contributed equally to this work.

The author responsible for distribution of materials integral to the findings presented in
this article in accordance with the policy described in the Instructions for Authors
(https://academic.oup.com/plphys/pages/General-Instructions) is Toshihisa Kotake
(kotake@mail.saitama-u.ac.jp).

Keywords: endo-β-1,4-mannanase, glucomannan, Golgi apparatus, membrane-
anchored enzyme, quality control

ABSTRACT

The molecular mechanisms underlying the synthesis of large cell wall polysaccharides

in plant cells are not fully understood. Here we report that two atypical endo-β-1,4-

mannanases (MANs), which are not secreted and do not degrade glucomannan in the

cell wall, play a novel role in glucomannan synthesis. Among the six MANs in

Arabidopsis, AtMAN2 and AtMAN5 contain a transmembrane domain at their N-

terminal region instead of a signal peptide. Subcellular localization using MAN protein

fused with fluorescent protein demonstrated that AtMAN2 localizes to the

endomembrane system including the Golgi apparatus in xylem and interfascicular fiber

cells. We found that an Arabidopsis *man2 man5* double mutant lost 65% of

glucomannan in the cell walls of the inflorescence stem. Immunostaining and

immunoelectron microscopic observation also revealed that the *man2 man5* double

mutant lost glucomannan in the cell walls to about the same extent as the *csla2 csla9*

double mutant, which lacks major glucomannan synthases. Gene complementation

experiments showed that the enzymatic activities of AtMAN2 and AtMAN5 are

important for the function in synthesis of glucomannan for the cell wall. Arabidopsis

possesses another atypical MAN, AtMAN6, with an HDEL retention signal at its C-

terminus. However, mutation of AtMAN6 did not affect glucomannan content in the cell

walls, suggesting distinct functions for these MANs. This study has identified AtMAN2

and AtMAN5 as novel factors necessary for the normal glucomannan synthesis in

Arabidopsis, along with GDP-mannose generating enzymes and CslAs, and suggests

that the glucomannan hydrolysis by these MANs contributes to the maintenance of

glucomannan synthesis.

INTRODUCTION

Plant cell wall polysaccharides constitute the largest biomass carbon reservoir on

earth. Understanding the mechanism for their synthesis is thus indispensable for their

efficient production and utilization (Bar-on et al. 2018). Large polysaccharides are

synthesized by glycosyltransferases from building block nucleotide sugars (Verbančič et

al. 2018), but the molecular mechanisms are not fully understood. It is well-known that

proteins undergo quality control during their synthesis, which prevents protein

aggregation and removes unfolded proteins from the endoplasmic reticulum (ER) to be

degraded in 26S proteasomes in the cytosol (Mori 2000; Mayer and Bulau 2005). On

the other hand, how the aggregation of large cell wall polysaccharides is prevented to

maintain the synthesis reaction in the Golgi apparatus is still unknown.

Mannan polysaccharides are cell wall polysaccharides widely distributed in land

plants. They commonly possess a backbone composed of β-1,4-mannose (β-1,4-Man)

and/or β-1,4-glucose (β-1,4-Glc), while the backbone sequences and decorations are

diverse (Scheller and Ulvskov 2010; Voiniciuc 2022; Yoshimi et al. 2025). The

backbones of mannan polysaccharides can take on a two-fold structure, in which the

sugar residues turn at 180° resulting in a straight ribbon structure suitable for cellulose

binding (Yu et al. 2018; 2022). Mannan polysaccharides without decoration tend to

become insoluble, probably as a result of intermolecular hydrogen bonding and

hydrophobic interactions (Chanzy et al. 1984; Zhou et al. 2018).

The backbones of mannan polysaccharides are synthesized from GDP-Man and

GDP-Glc by the action of specific glycosyltransferases, called cellulose-synthase like A

(CslA) proteins (Dhugga et al. 2004; Concklin et al. 1999; Liepman et al. 2005). In

Arabidopsis (*Arabidopsis thaliana*), CslA2 synthesizes the backbone for β-

galactoglucomannan (β-GGM) with alternating β-1,4-Man and β-1,4-Glc residues,

whereas ClsA9 makes the backbone for acetylated GGM (AcGGM) with a random

sequence of β-1,4-Man and β-1,4-Glc residues (Goubet et al. 2009; Yu et al. 2018; 2022,

in this paper, glucomannan is taken to include β-GGM and AcGGM). The reduced

glucomannan content in Arabidopsis *vitamin c defective1*, *konjac1*, *csla2,* and *csla9*

mutants is an indication of the importance of factors involved in the building block

generation and the backbone synthesis reactions (Conklin et al. 1999; 2000; Goubet et

al. 2009; Sawake et al. 2015; Nishigaki et al. 2021). These backbones subsequently

undergo decoration with α-galactose by mannan α-galactosyltransferase and/or acetyl

groups by mannan *O*-acetyltransferase, with which the solubility of glucomannan

increases (Reid et al. 2003; Voiniciuc et al. 2015; Zhong et al. 2018; Yoshimi et al.

2025). MANNAN SYNTHESIS-RELATED (MSR) proteins are other factors that have

been shown to affect the glucomannan content, although their molecular functions are

still obscure (Wang et al. 2013). The identification of additional factors is important for

understanding the molecular mechanism of glucomannan synthesis.

Endo-β-1,4-mannanase (MAN) is one of the conserved cell wall enzymes in plants,

belonging to the glycoside hydrolase (GH) 5 family (Aspeborg et al. 2021). The

physiological importance of MANs has been extensively studied in seed germination

and fruit ripening (Nonogaki et al. 2000; Schröder et al. 2004; Belotserkovsky et al.

2007; Iglesias-Fernández et al. 2011). Based on domains, motifs, and sequence

similarity, plant MANs can be classified into several types (Fig. 1A) (Wang et al. 2014).

Arabidopsis secreted type MANs, AtMAN3 and AtMAN7, hydrolyze cell wall

glucomannan to contribute to cadmium tolerance via activation of phytochelatin and

glutathione synthesis (Chen et al. 2015; Wu et al. 2023). AtMAN6 is a HDEL-type

MAN with a C-terminal HDEL signal, which is a retrograde transport signal back from

the Golgi apparatus to the ER (Alvim et al. 2023). AtMAN6 and its poplar (*Populus*

trichocarpa) orthologue, PtrMAN6, are specifically localized in vessel cells and

presumed to release GGM oligosaccharides as signaling molecules, which negatively

regulate secondary cell wall formation in the vessel and interfascicular fiber cells (Zhao

et al. 2013; Zhang et al. 2023). On the other hand, AtMAN2 and AtMAN5 are a

different type of MAN with a transmembrane (TM) domain instead of a signal peptide

in the N-terminal region, designated TM-type MANs here. Recombinant AtMAN2

expressed in *Escherichia coli* shows hydrolytic activity toward mannan polysaccharides

and lacks transglycosylation activity (Wang et al. 2015), but its physiological

importance and functions remain to be studied.

This study focused on the importance and functions of the TM-type MANs, AtMAN2

and AtMAN5, comparing them with another atypical MAN, AtMAN6, in glucomannan

synthesis. The analysis of Arabidopsis loss-of-function mutants suggested that the

hydrolysis of glucomannan by AtMAN2 and AtMAN5 is necessary for normal

glucomannan synthesis, where their roles must thus be different from that of AtMAN6.

These findings provide new insights into the synthesis of plant cell wall polysaccharides.

RESULTS

TM-type MANs are conserved in angiosperms

In Arabidopsis, two TM-type MANs, AtMAN2 (AT2G20680) and AtMAN5

(AT4G28320), are highly related in their amino acid sequences (identity, 83%), while

AtMAN6 (AT5G01930) is the sole HDEL-type MAN (Fig. 1A). TM-type MANs can be

found in other angiosperms including amborella (*Amborella trichopoda*) and rice

(*Oryza sativa*), forming a clade apart from the secreted type MANs in the phylogenetic

tree (Fig. 1B). On the other hand, *Physcomitrium patens* (a moss species) and

Marchantia polymorpha (a liverwort species) lack this type. The HDEL-type MANs

also formed a different clade, which did not include MANs from *P. patens* and *M.*

polymorpha. The distribution and phylogenetic relationships suggest that the TM-type

MANs have functions different from the secreted and the HDEL-type MANs.

Loss of glucomannan in *man2 man5* double mutant

Because high sequence similarity suggested functional redundancy of AtMAN2 and

AtMAN5, an Arabidopsis *man2-1 man5-1* double mutant was generated for functional

analysis. The double mutants did not show significant growth defects in the seedlings

(Supplementary Fig. S1). In addition, although the double mutants had slightly shorter

inflorescence stems than wild type (WT) plants as was also observed for the *csla2-2*

csla9-2 double mutants, they did not show severe growth defects (Yu et al. 2022)

(Supplementary Fig. S2). Based on the expression patterns (Winter et al. 2007),

AtMAN2 and *AtMAN5* are presumed to function in the inflorescence stem

(Supplementary Fig. S3).

To examine the importance of AtMAN2 and AtMAN5 for glucomannan accumulation,

we observed the cross sections of the bottom part of inflorescence stems of Arabidopsis

man2-1 single, *man2-2* single, *man5-1* single, and *man2-1 man5-1* double mutants and

compared them with WT with LM21 antibody specific to glucomannan (Thorne et al.

2023). Although MANs are hydrolytic enzymes acting on mannan polysaccharides

(Wang et al. 2015), the glucomannan signals were, unexpectedly, almost lost in the cell

walls of the *man2-1 man5-1* double mutant as they were in the *csla2-2 csla9-2* double

mutant, while the signals were observed in the single mutants as they were in the WT,

indicating the functional redundancy of AtMAN2 and AtMAN5 (Fig. 2). To examine the

functional relationships of AtMAN2 and AtMAN5 with AtMAN6, the cross sections of

the *man6-1* single, the *man2-1 man6-1* double, the *man5-1 man6-1* double, the *man2-1*

man5-1 man6-1 triple, and the *man2-2 man5-1 man6-1* triple mutants were also

observed. The cell wall glucomannan signals were almost lost in the triple mutants as in

the *man2-1 man5-1* double mutant but remained in the other double mutants (Fig. 2),

suggesting that the AtMAN2 and AtMAN5 contribution to the glucomannan

accumulation differs from that of AtMAN6. Xylan, together with cellulose, is a major

component of secondary cell walls, but immunostaining with LM10 antibody specific to

xylan did not point to compensatory changes in the mutants (Supplementary Fig. S4).

The TM-type MANs do not seem to influence the content of other cell wall

polysaccharides in the inflorescence stem.

Drastic decrease of cell wall glucomannan in the double and triple mutants

The changes in the glucomannan accumulation in the inflorescence stems of the *man2-1*

man5-1 double and the *man2-1 man5-1 man6-1* and the *man2-2 man5-1 man6-1* triple

mutants were further investigated by cell wall fractionation and subsequent sugar

composition analysis. The cell walls were fractionated into water-soluble fraction

(soluble fraction), alkali-extractable fraction (alkali fraction), and cellulose fraction (Fig.

3A; Supplementary Fig. S5), and the sugar composition of these fractions was

determined by high-performance anion-exchange chromatography with pulsed

amperometric detection (HPAEC-PAD) (Supplementary Fig. S6). Consistent with the

immunostaining of glucomannan with LM21 in the cross sections (Fig. 2), the *man2-1*

man5-1 double and the *man2-1 man5-1 man6-1* and *man2-2 man5-1 man6-1* triple

mutants lost more than 65% mannose in these cell wall fractions (Fig. 3B). As other

hemicelluloses and pectin rarely have mannosyl residues in Arabidopsis (Mohnen 2008;

Goubet et al. 2009; Scheller and Ulvskov 2010), this result indicates that the double and

triple mutants lost a large part of its glucomannan. It has been reported that AtMAN6

negatively regulates secondary cell wall thickening (Zhang et al. 2023), but neither

mannose content nor the sugar content proportion of cell wall fractions was changed in

the *man6-1* mutant (Fig. 3).

The Arabidopsis inflorescence stem has two different glucomannans, β-GGM and

AcGGM, which are synthesized by CslA2 and CslA9, respectively (Goubet et al. 2009;

Yu et al. 2018; 2022). The influence of the loss of TM-type MANs on these

glucomannans was evaluated by polysaccharide analysis using carbohydrate gel

electrophoresis (PACE) (Supplementary Fig. S7). Here, β-GGM and AcGGM included

in the stem cell walls were hydrolyzed into oligosaccharides by a microbial MAN,

CjMan26A, separated, and detected (Hogg et al. 2001). The *man2-1 man5-1* double and

the triple mutants with *man6-1* showed almost no β-Glc-(β-Gal-α-Gal-)β-Man-β-Glc-(α-

Gal-)β-Man-β-Glc-Man (GBGAGM) and β-Glc-(β-Gal-α-Gal-)β-Man-β-Glc-Man

(GBGM) oligosaccharides from β-GGM and no β-Glc-β-Glc-Man (GGM) and β-Glc-β-

Man-Man (GMM) from AcGGM (Supplementary Fig. S7), so they were similar to the

csla2-2 csla9-2 double mutant in this respect (Goubet et al. 2009) (Fig. 4). These results

demonstrate that the TM-type MANs are necessary for the normal synthesis of both β-

GGM and AcGGM in the inflorescence stem.

Retention of glucomannan in the vacuole

β-GGM and AcGGM are synthesized in the Golgi apparatus by CslA2 and CslA9,

respectively, then secreted to the cell walls. While the expression levels of these *CslA*

genes did not decrease (Supplementary Fig. S8), the glucomannan amount drastically

decreased (Figs. 3 and 4). To gain insight into the functions of AtMAN2 and AtMAN5,

the glucomannan in the cross sections of *man* mutants was observed at high

magnification. Although the *man2-1 man5-1* double and the *man2-1 man5-1 man6-1*

triple mutants lost most cell wall glucomannan as shown in Fig. 3, these mutants

contained relatively large (>1 μm) particles detected with LM21 antibody inside the

xylem and interfascicular fiber cells of the bottom part of inflorescence stems (Fig. 5A).

Importantly, glucomannans inside cells have also been observed in Arabidopsis and

other plants such as aloe (*Aloe vera*) (Ahl et al. 2019; Handford et al. 2003; Herburger

et al. 2022), although these glucomannans have not been identified. Because the

particles significantly decreased in the *csla2-2 csla9-2* double mutant, most of these

particles can be presumed to be β-GGM and AcGGM that were not secreted to the cell

walls. On the other hand, the particles that remained even in the *csla2-2 csla9-2* double

mutant may be glucomannan synthesized by different CslAs from CslA2 and Csla9.

Although significant differences from WT were not observed, these particles tended to

increase rather than decrease in the double and triple mutants (Fig. 5B).

To identify the glucomannan particles retained inside cells, immunoelectron

microscopy was conducted. In this experiment, the glucomannans in the ultrathin cross

sections from the bottom part of inflorescence stems of the *man2-1 man5-1* double and

the *man2-1 man5-1 man6-1* triple mutants were recognized by LM21 as the primary

antibody and visualized with a secondary antibody conjugated to gold particles.

Consistent with immunostaining and cell wall analysis, there were fewer gold particles

in the cell walls of the triple mutant, showing loss of glucomannan in the cell walls.

Relatively large clusters (>1 µm) of gold particles were found in the vacuole in the triple

mutant (Fig. 5C). These observations suggest that the glucomannan particles are formed

and retained in the vacuole in the absence of the TM-type MANs, AtMAN2 and

AtMAN5.

230 **Tissue and subcellular localization of AtMAN2 and AtMAN6 differ**

231 The tissue and subcellular localization of AtMAN2 and AtMAN5 was investigated

232 using transgenic Arabidopsis harboring *pMAN2::MAN2TM-mNeon* or

233 *pMAN5::MAN5TM-mNeon* in which the fluorescent protein mNeonGreen sequence was

234 inserted right behind the TM domain in the genomic sequences (Supplementary Fig. S9).

235 In this experiment, the gene constructs were introduced into the *man2-1 man5-1 man6-1*

236 or the *man2-2 man5-1 man6-1* triple mutants. Immunostaining of the cross sections of

237 the bottom part of inflorescence stems showed that AtMAN2TM-mNeon and

238 AtMAN5TM-mNeon at least partially restored impaired glucomannan accumulation in

239 the triple mutants (Supplementary Fig. S9). For microscopic observation, the top part of

240 the inflorescence stem was chosen because strong mNeonGreen signals were found

241 there. The AtMAN2TM-mNeon signal was widely observed in the xylem and

242 interfascicular fiber cells of inflorescence stems (Supplementary Fig. S10). The

243 AtMAN2TM-mNeon signal was not found in the cell walls but localized in the

244 compartments (Fig. 6A-E), suggesting that AtMAN2 functions in the endomembrane

245 including the Golgi apparatus. The AtMAN5TM-mNeon signal was also found inside

246 cells (Fig. 6F-J), although the localization was not clear because of low signal intensity.

247 To compare, we also generated *pMAN6::MAN6signal-mNeon* in which the

248 mNeonGreen sequence was inserted after the signal peptide into the genomic *AtMAN6*

249 sequence (Supplementary Fig. S9). Contrasting with AtMAN2TM-mNeon signals, the

250 AtMAN6signal fluorescence was limited to developing vessels (Supplementary Fig.

251 S10). The fluorescence was spread in the vessel cells (Fig. 6K-O), indicating its

localization in the ER or the vacuole. Previously it was suggested that AtMAN6

localizes on the plasma membrane (PM), but this was using a C-terminal fusion of

AtMAN6 with green fluorescent protein, which might have affected the localization.

But in that study, as in this, AtMAN2 and AtMAN6 were found in substantially different

locations, indicating that they have different molecular functions (Zhang et al. 2023).

The enzymatic activities of AtMAN2 and AtMAN5 is required

To determine the importance of the enzymatic activities of AtMAN2 and AtMAN5, we

performed gene complementation experiments on the *man2-1 man5-1* double mutant

using point-mutated genomic *AtMAN2* and *AtMAN5* genes in which the catalytic

residues were replaced with Ala (E215A and E335A in AtMAN2; E214A and E334A in

AtMAN5). The mutated *AtMAN2* and *AtMAN5* genes failed to restore the glucomannan

accumulation detected with the LM21 antibody in the inflorescence stem of the double

mutant (Supplementary Figs. S9, S11). This result suggests that for proper functioning

of the glucomannan synthesis in the Golgi apparatus, the enzymatic activity of these

MANs is necessary.

DISCUSSION

MANs are newly recognized as necessary for normal glucomannan synthesis

The present study has identified AtMAN2 and AtMAN5 as factors necessary for normal

glucomannan synthesis in Arabidopsis, to work along with VTC1 and KJC1 generating

GDP-Man, CslAs catalyzing the glucomannan backbone synthesis, and MSRs (Conklin

274 et al. 1999; Goubet et al. 2009; Wang et al. 2013; Yu et al. 2014; Sawake et al. 2015;

275 Nishigaki et al. 2021). Based on the distribution of orthologues, it is likely that many

276 angiosperms utilize TM-type MANs in glucomannan synthesis.

277  Because the expression levels of *CslA2* and *CslA9* did not decrease in the *man2-1*

278 *man5-1* double and the *man2-1 man5-1 man6-1* triple mutants (Supplementary Fig. S8),

279 we infer that the decreased glucomannan in the mutants was not caused by changes in

280 the expression of *CslA* genes and that AtMAN2 and AtMAN5 contribute to

281 glucomannan synthesis through their enzymatic activity. The requirement of

282 polysaccharide degrading enzymes for polysaccharide synthesis presents an apparent

283 paradox. Yet this is not the first example: among other plant glycoside hydrolases, the

284 membrane-associated GH9 endo-β-1,4-glucanase KORRIGAN1 (KOR1) is well known

285 for its role in cellulose synthesis (Nicol et al. 1996). The Arabidopsis *kor1* mutant

286 suffers severe defects in the synthesis of cellulose, demonstrating the importance of this

287 protein. The sea squirt *Ciona intestinalis* and bacteria such as *Agrobacterium*

288 *tumefaciens* and *Acetobacter xylinum* also require endo-β-1,4-glucanases for their

289 cellulose synthesis. These enzymes belong to the GH6 and 8 families and are thus

290 distinct from KOR1 (Standal et al. 1994; Matthysse et al. 1995; Li et al. 2023). These

291 observations imply that hydrolytic activity of endo-β-1,4-glucanases is important to

292 facilitate cellulose synthesis. The present study describes a second instance of a version

293 of this paradox in polysaccharide synthesis. We suggest that this paradox may represent

294 a general mechanism to prevent the aggregation of newly synthesized cell wall

295 polysaccharides in plant cells.

Not all CslAs require TM-type MANs for glucomannan synthesis

The TM-type MANs are not necessarily required for all glucomannan synthesis by

CslAs in plants. In the present study, the alkali fraction from the inflorescence stems of

man2-1 man5-1 double mutant still contained 30% of mannose compared with WT (Fig.

3B). Because *N*-glycosylated proteins with mannosyl residues would not have been

collected in the alkali fraction, there seems to be mannan polysaccharide synthesized

independently from the functions of AtMAN2 and AtMAN5. This unknown mannan

polysaccharide would have to be resistant to hydrolysis by CjMan26A used for PACE.

Many green algae and bryophytes such as *P. patens* and *M. polymorpha* have mannan

polysaccharides as major cell wall components (Kolkas et al. 2023; Chernova et al.

2024), although they lack genes orthologous to *AtMAN2* and *AtMAN5* (Fig, 1B). To

understand the molecular mechanism for the glucomannan synthesis involving TM-type

MAN, future research should try to clarify which types of mannan polysaccharides

require these MANs.

Putative function of AtMAN2 and AtMAN5 in quality control

AtMAN2TM-mNeon localized in the endomembrane including the Golgi apparatus in

the inflorescence stem (Fig. 6). In the gene complementation experiments, the genomic

AtMAN2 and *AtMAN5* genes with mutations at catalytic residues failed to restore cell

wall glucomannan in the *man2 man5* double mutant (Supplementary Fig. S11). Recently,

AtMAN2 and AtMAN5 were also shown to increase soluble mannan polysaccharides in

Pichia yeast (*Pichia pastoris*) expressing Arabidopsis *CslA2* (Jacobson et al. 2025). It

has been reported that AtMAN2 lacks transglycosylation activity (Wang et al. 2015).

Based on these facts, we suggest that glucomannan undergoes hydrolysis by AtMAN2

and AtMAN5 after the synthesis in the Golgi apparatus. The drastic decrease in the

glucomannan content in the double and triple mutants implies that the continuous

synthesis reaction by CslAs is not maintained in these mutants. The alternative, that

only secretion is impaired in the mutants, while the glucomannan synthesis by CslA2

and CslA9 is maintained, does not seem likely, because in this case the glucomannan

would have been collected in one of the fractions (Fig. 3B). We cannot specify the

molecular functions of these MANs in detail, but it is conceivable that large β-GGM

and AcGGM molecules synthesized by CslA2 and CslA9 before decoration easily

aggregate during synthesis, which may cause the termination of the chain elongation

reaction by these CslAs. This problem may be alleviated through removal by partial

hydrolysis with these MANs as polysaccharides generally become more soluble through

hydrolysis into shorter chains. These reactions to prevent aggregate formation and to

remove aggregates in the Golgi apparatus can be considered analogous to the protein

quality control in the ER (Mori 2000; Oda et al. 2003). It is also probable that AtMAN2

and AtMAN5 facilitate the release of glucomannan from CslA proteins by partial

hydrolysis during synthesis, enabling its proper loading into the secretion pathway. The

impairment of this process may also negatively affect glucomannan synthesis by CslAs.

In xyloglucan, the importance of galactose-side chains to avoid aggregation has been

shown (Tamura et al. 2005; Kong et al. 2015; Hoffmann and McFarlane 2024). The

Arabidopsis *murus3* mutant, which lacks galactose-side chains on its xyloglucan, suffers

intracellular aggregation of xyloglucan in the roots, resulting in growth defects. It is

probable that such intracellular aggregation was avoided through the termination of the

synthesis reaction in the *man2-1 man5-1* double mutant. Clarifying this mechanism will

presumably require the identification of as yet unknown factors involved in this reaction.

MATERIALS AND METHODS

Arabidopsis mutants and growth conditions

Arabidopsis thaliana ecotype Col-0 was used in this study. The T-DNA insertion lines

SALK_0383344 (*man2-1*), SALK_016450 (*man2-2*), SALK_059047 (*man5-1*), and

SALK_129977 (*man6-1*) were provided by the Arabidopsis Biological Resource Center.

The double and triple mutants were generated by crosses between these mutants. The

genotypes of the mutants were determined using specific primers (Supplementary Fig.

S12 and Table S1). The *csla2-2 csla9-2* and *msr1-1 msr2-2* double mutants had been

obtained in our previous studies (Goubet et al. 2007; 2009; Wang et al. 2013).

Arabidopsis seeds were sterilized and sown on Murashige and Skoog (MS) medium in

0.8% (w/v) agar. After stratification at 4 °C in the dark for 2 days, the seeds were

germinated and grown under continuous light (cold cathode fluorescent lamp;

Biomedical Science) at 23 °C for 10 days, then on rockwool fiber for 4 weeks under

continuous light at 23 °C.

For the gene complementation experiments, genomic DNA fragments for *AtMAN2*

and *AtMAN5* were amplified by PCR and subcloned into the binary vector pBGGN

(Inplanta Innovation, Yokohama, Japan). The point mutations at the catalytic sites were

introduced by PCR with specific primers (Supplementary Table S1). The *man2-1 man5-*

1 double mutant was transformed with mutated AtMAN2/pBGGN, WT

AtMAN2/pBGGN, mutated AtMAN5/pBGGN, and WT AtMAN5/pBGGN by an

Agrobacterium-mediated (*Rhizobium radiobacter*, EHA105 strain) method (Clough and

Bent 1998).

Phylogenetic analysis

The amino acid sequences of MANs from various plants were collected using Basic

Local Alignment Search Tool in NCBI (https://blast.ncbi.nlm.nih.gov/Blast.cgi)

(Supplementary Table S2). The signal peptide and TM region were detected using the

SignalP 6.0 and TMHMM-2.0 programs, respectively (Teufel et al. 2022; Krogh et al.

2001; Moller et al. 2001). The alignments of amino acid sequences were performed by

the MUSCLE program in MEGA software (version 11; Tamura et al. 2021) with default

parameters. The phylogenetic trees were generated using the maximum likelihood

method with 100 bootstraps.

Cell wall fractionation and sugar composition analysis

The bottom part of first inflorescence stem 2 weeks after bolting (approximately 50 mg)

was homogenized with mortar and pestle in water (Supplementary Fig. S4). After

centrifugation at 10,000 x g at 4°C for 5 min, the supernatant was collected as the

soluble fraction. The precipitate was suspended in 0.4 mL of 4 M potassium hydroxide

(KOH) containing 0.04% (w/v) sodium borohydride at 25°C. After centrifugation at

10,000 x g at 4°C for 5 min, the supernatant was collected. This extraction was repeated.

The combined supernatant (0.8 mL) was neutralized with 0.4 mL of glacial acetic acid

and collected as the alkali fraction. The precipitate was washed with 98% (v/v) ethanol,

2% (v/v) acetic acid twice, lyophilized for 24 h, hydrolyzed with 72% (v/v) sulfuric acid

at 4 °C for 1 h and 8% (v/v) sulfuric acid at 100 °C for 4 h, and then collected as the

cellulose fraction. Total sugar composition and quantity of the above fractions were

determined using the phenol sulfuric acid procedure (Dubois et al. 1956).

For sugar composition analysis, the alkali fraction was dialyzed against water at 4 °C

for 2 days. The soluble and alkali fractions were hydrolyzed into monosaccharides with

2 M trifluoroacetic acid (TFA) at 121°C for 1 h, then lyophilized to remove TFA. The

cellulose fraction after the hydrolysis with sulfuric acid was neutralized with barium

carbonate and desalted with Dowex beads (Dowex 50WX8, Dow Chemical).

Monosaccharide composition was determined by HPAEC-PAD using a Dionex ICS-

5000+ liquid chromatograph (Thermo Fisher Scientific) fitted with a CarboPac PA-1

column (Thermo Fisher Scientific) and a pulsed amperometric detector (Thermo Fisher

Scientific) as described previously (Sawake et al. 2015).

PACE

Alcohol-insoluble residue (AIR) was prepared from the bottom part of Arabidopsis

inflorescence stems (fresh weight, 100 mg) according to Yu et al. (2018). Glucomannans

were extracted from the AIR with 1 mL of 4 M KOH solution by shaking at 1,000 rpm

at room temperature for an hour. After centrifugation at 15,000 x g for 15 minutes, the

supernatant was collected. To collect remaining glucomannans, the pellet was

suspended in 1 mL of water. After centrifugation, the supernatant was collected. The

supernatants were combined and applied to a PD-10 column (Cytiva) to exchange the

solvent with 50 mM ammonium acetate buffer (pH 6.0). The eluted fraction was used as

the KOH fraction.

Glucomannans in the KOH fraction were hydrolyzed into specific oligosaccharides

with a commercial endo-β-1,4-mannanase, CjMan26A (1 unit, E-BMACJ, Megazyme),

from *Cellvibrio japonicus* in 50 mM ammonium acetate buffer (pH 6.0) at 37 °C

overnight (Supplementary Fig. S7) (Hogg et al. 2001). After the reaction, the enzyme

was denatured by heating at 100 °C for 15 minutes. The oligosaccharides were

derivatized with 8-aminonapthalene-1,3,6-trisulphonic acid, resuspended in urea, and

separated by electrophoresis as described (Goubet et al. 2009; Yoshimi et al. 2025). Gels

were visualized using a G-Box (Syngene, Cambridge, UK) with a transilluminator with

long-wave tubes emitting at 365 nm and a shortpass (500 to 600 nm) detection filter.

Mannose and β-1,4-mannooligosaccharides (Man_{2-6}, Megazyme) were used as standards.

Immunostaining of glucomannan in the mutants

The glucomannan and xylan in the inflorescence stems were observed using specific

antibodies. The bottom part of inflorescence stems (the lowest one-sixth portion) was

fixed with 4% (v/v) paraformaldehyde and 0.6% (v/v) glutaraldehyde at 4 °C for 3 h.

The fixed samples were dehydrated through a graded ethanol series, embedded in

Technovit 7100 (Nisshin EM, Tokyo, Japan) using a graded series of ethanol and

Technovit 7100 mixtures, and solidified with Technovit hardener (Nisshin EM) at 60 °C

for 12 h. Sections were sliced to a thickness of 10 μm using a microtome (RM2125RT;

Leica, Tokyo, Japan) equipped with a disposable steel blade (C35; Feather, Osaka,

Japan).

To remove acetyl groups from polysaccharides, the cross-sections were treated with

50 mM sodium hydroxide at 25 °C for 30 min. The sections were then washed with

phosphate-buffered saline (PBS) containing 137 mM NaCl, 10 mM disodium

hydrogenphosphate, 2.7 mM KCl, and 1.8 mM potassium dihydrogenphosphate (pH

7.4) and then blocked in PBS containing 0.3% (w/v) blocking milk (Sigma-Aldrich,

Tokyo, Japan) for 30 min. Then, the sections were incubated for 1.5 h with antibodies

against glucomannan (LM21, Kerafast, Shirley, MA; dilution 1:5) or xylan (LM10,

Kerafast; dilution 1:5) in PBS with blocking milk. After washing three times with PBS,

the sections were incubated for 1 h with Alexa Fluor 488-conjugated anti-mouse IgM

for LM21 (abcam, Cambridge, UK; dilution 1:100) or Alexa Fluor 488-conjugated anti-

mouse IgG for LM10 (abcam; dilution 1:100) in PBS with bovine serum albumin

(Thorne et al. 2023). The sections were washed three additional times with PBS. The

cross sections were imaged with a microscope (ECLIPSE Ci-L Plus, Nikon, Japan) at

488 nm. The dark field image was also acquired simultaneously.

Immunoelectron microscopy

Immunoelectron microscopy to detect the glucomannan localization was performed according to Atsuzawa et al. (2010). The bottom part (the lowest one-sixth portion) and upper part (the highest one-sixth portion) of the inflorescence stems of WT, the *man6-1* single, the *man2-1 man5-1* double, and the *man2-1 man5-1 man6-1* triple mutants were fixed with 2% (v/v) glutaraldehyde in 50 mM potassium phosphate (pH 6.8) at room temperature for 3 h. After washing with 100 mM lysine in 50 mM potassium phosphate (pH 6.8) the samples were dehydrated in a graded ethanol series. The tissues were embedded in LR White resin (Electron Microscopy Sciences, USA) and polymerized by heating at 60 °C. Ultrathin sections of 100 nm thickness were cut on an UltracutN ultramicrotome (Nissei) and collected on nickel grids coated with polyvinyl formvar membrane.

Ultra-thin sections were immersed in 50 mM sodium hydroxide at room temperature for 30 min and washed with Tris-HCl buffer (TBS: 50 mM Tris-HCl (pH 7.5), 150 mM sodium chloride) containing 0.05% Triton X-100 (TBS/Tx). They were treated with blocking solution (10 mg/ml bovine serum albumin in TBS) at room temperature for 60 min, and they were incubated with LM21 antibody at 1:500 dilution at 4 °C overnight. After washing with TBS/Tx, the sections were incubated with a solution of 10 nm gold particles conjugated with goat anti-rat IgG (BBI solutions, UK) as the secondary antibody at a 1:30 dilution for 1 h at room temperature and washed with TBS/Tx. Specimens were observed with a transmission electron microscope (H-7500, Hitachi,

Japan) at an accelerating voltage of 80 kV after staining with uranyl acetate and lead

citrate.

Transgenic plants expressing MAN-mNeonGreen

For the generation of transgenic Arabidopsis harboring *pMAN2::MAN2TM-mNeon*,

pMAN5::MAN5TM-mNeon, or *pMAN6::MAN6signal-mNeon*, the DNA fragments of

genomic *AtMAN2*, *AtMAN5*, and *AtMAN6* and mNeonGreen were amplified from

Arabidopsis genomic DNA and template mNeonGreen (Addgene, #189976),

respectively, using the primer sets listed in Supplementary Table S1. The DNA

fragments were assembled and subcloned into the binary vector pBGGN with DNA

assembly kit (Gibson) to yield the constructs, *pMAN2::MAN2TM-mNeon/pBGGN*,

pMAN5::MAN5TM-mNeon/pBGGN and *pMAN6::MAN6signal-mNeon/pBGGN*

(Supplementary Fig. S9). These constructs were introduced into the *man2-1 man5-1*

man6-1 or the *man2-2 man5-1 man6-1* triple mutants by an Agrobacterium-mediated

(*Rhizobium radiobacter*, EHA105 strain) method (Clough and Bent 1998).

Observation of tissue and subcellular localization of MAN-mNeonGreen proteins

Cross sections were excised from the top part of the inflorescence stems of

pMAN2::MAN2signal-mNeon (line #3), *pMAN2::MAN2signal-mNeon* (line #4), and

pMAN6::MAN2signal-mNeon (line #3) plants and stained with 6.34 μM fixable analog

of N-(3-triethylammoniumpropyl)-4-(6-(4-(diethylamino) phenyl) hexatrienyl)

pyridinium dibromide, FM 4-64FX (ThermoFisher). The confocal images were obtained

491 using a Zeiss LSM800 (Zeiss, Jena, Germany) with a Plan-Apochromat 63×/1.4 oil

492 immersion objective lens. FM 4-64FX was excited at 561 nm and detected between 400

493 and 645 nm. Fluorescence signals from mNeonGreen were detected after excitation at

494 488 nm and detected at 410-546 nm. Autofluorescence was excited at 488 nm and

495 detected at 550-700 nm. Transmitted photomultiplier tube (T-PMT) images were also

496 captured during confocal imaging.

498 **Author Contributions**: T.K. and P.D. designed the research. A.K., E.S., Y.Y., H.T.,

499 N.N., K.A., Y.K., D.T., and T.K. performed research. E.S., Y.Y., D.T., Y.K., and T.K.

500 analyzed the data. H.T., Y.K., D.T., P.D., and T.K. wrote the manuscript.

502 **Conflict of interest**

503 The authors declare no conflict of interest.

505 **Acknowledgments**: We thank Dr. Kiminori Toyooka from RIKEN and Prof. Masatsugu

506 Toyota and Dr. Hiraku Suda from Saitama University for valuable discussions. This

507 work was supported by MEXT KAKENHI Grant-in-Aid for Scientific Research to T.

508 Kotake [no. 18H05495], Grant-in-Aid for Scientific Research to T. Kotake [no.

509 23H04302 and no. 23H02134] and to D. Takahashi [no. 23K05144 and no. 24KK0273],

510 the ERC Advanced Grant EVOCATE to P. Dupree funded by the United Kingdom

511 Research and Innovation (UKRI) grant number EP/X027120/1 (www.ukri.org) and by

512 the grant OpenPlant (UKRI BBSRC, BB/L014130/1) to P. Dupree, and a Broodbank

513 Research Fellowship to Y. Yoshimi (reference no. PD16178).

Supplementary Data

Supplementary Table S1. Supplementary Table S1. Primers used for genotyping,

plasmid construction, point mutations, and RT-PCR.

Supplementary Table S2. List of MAN sequences in the phylogenetic analysis.

Supplementary Fig. S1. Growth phenotypes of seedlings of *man* mutants

Supplementary Fig. S2. Growth phenotypes in the inflorescence stems of *man* mutants

Supplementary Fig. S3. Expression patterns of *AtMAN2*, *AtMAN5*, and *AtMAN6* genes.

Supplementary Fig. S4. Xylan accumulation in *man* mutants.

Supplementary Fig. S5 Cell wall fractionation from inflorescence stems.

Supplementary Fig. S6. Monosaccharide composition of the soluble, alkali, and

cellulose fractions.

Supplementary Fig. S7. Oligosaccharides detected by PACE.

Supplementary Fig. S8. Expression levels of *CslA2* and *CslA9* in *man* mutants.

Supplementary Fig. S9. Gene constructs and complementation of *man* mutants.

Supplementary Fig. S10. Tissue localization of MAN-mNeon proteins.

Supplementary Fig. S11. Gene complementation experiment with mutated genomic

AtMAN2 and *AtMAN5* genes.

Supplementary Fig. S12. T-DNA insertion sites and genotyping of *man* mutants.

LITERATURE CITED

Ahl LI, Mravec J, Jørgensen B, Rudall PJ, Rønsted N, Grace OM. Dynamics of
intracellular mannan and cell wall folding in the drought responses of succulent
Aloe   species.   *Plant   Cell   Environ.*   2019:42(8):2458-2471.
https://doi.org/10.1111/pce.13560

Alvim JC, Bolt RM, An J, Kamisugi Y, Cuming A, Silva-Alvim FAL, Concha JO,
daSilva LLP, Hu M, Hirsz D, Denecke J. The K/HDEL receptor does not recycle
but   instead   acts   as   a   Golgi-gatekeeper.   *Nat   Commun.*   2023:14(1):1612.
https://doi.org/10.1038/s41467-023-37056-0

Aspeborg H, Coutinho PM, Wang Y, Brumer H 3rd, Henrissat B. Evolution,
substrate specificity and subfamily classification of glycoside hydrolase family 5
(GH5). *BMC Evol Biol.* 2012:12:186. https://doi.org/10.1186/1471-2148-12-186

Atsuzawa K, Nakazawa A, Mizutani K, Fukasawa M, Yamamoto N, Hashimoto T,
Usuda N. Immunohistochemical localization of mitochondrial fatty acid β-oxidation
enzymes in Müller cells of the retina. *Histochem Cell Biol.* 2010:134(6):565-579.
https://doi.org/10.1007/s00418-010-0752-4

Bar-On YM, Phillips R, Milo R. The biomass distribution on Earth. *Proc Natl Acad*
Sci USA. 2018:115(25):6506-6511. https://doi.org/10.1073/pnas.1711842115

Belotserkovsky H, Berger Y, Shahar R, Wolf S. Specific role of LeMAN2 in the
control of seed germination exposed by overexpression of the LeMAN3 gene in
tomato plants. *Planta.* 2007:227(1):199-209. https://doi.org/10.1007/s00425-007-
0607-y

Chanzy HD, Grosrenaud A, Vuong R, Mackie W. The crystalline polymorphism of
mannan in plant cell walls and after recrystallisation. *Planta.* 1984:161(4):320-329.
https://doi.org/10.1007/BF00398722

Chernova T, Ageeva M, Ivanov O, Lev-Yadun S, Gorshkova T. Characterization of
the fiber-like cortical cells in moss gametophytes. *Planta.* 2024:259(5):92.
https://doi.org/10.1007/s00425-024-04367-5

Chen J, Yang L, Gu J, Bai X, Ren Y, Fan T, Han Y, Jiang L, Xiao F, Liu Y, Cao S.
MAN3 gene regulates cadmium tolerance through the glutathione-dependent
pathway   in   *Arabidopsis   thaliana.   New   Phytol.*   2015:205(2):570-582.
https://doi.org/10.1111/nph.13101

Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-mediated
transformation   of   *Arabidopsis   thaliana.   Plant   J.*   1998:16(6):735-743.
https://doi.org/10.1046/j.1365-313x.1998.00343.x

Conklin PL, Norris SR, Wheeler GL, Williams EH, Smirnoff N, Last RL. Genetic
evidence for the role of GDP-mannose in plant ascorbic acid (vitamin C)
biosynthesis.   *Proc   Natl   Acad   Sci   USA.*   1999:96(7):4198–4203.
https://doi.org/10.1073/pnas.96.7.4198

Conklin PL, Saracco SA, Norris SR, Last RL. Identification of ascorbic acid-

deficient *Arabidopsis thaliana* mutants. *Genetics.* 2000:154(2):847–856. https://doi.org/10.1093/genetics/154.2.847

Dhugga KS, Barreiro R, Whitten B, Stecca K, Hazebroek J, Randhawa GS, Dolan M, Kinney AJ, Tomes D, Nichols S, Anderson P. Guar seed beta-mannan synthase is a member of the cellulose synthase super gene family. *Science.* 2004:303(5656):363-366. https://doi.org/10.1126/science.1090908

Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. Colorimetric method for determination of sugars and related substances. *Anal Chem.* 1956:28:350-356 https://doi.org/10.1021/ac60111a017

Goubet F, Barton CJ, Mortimer JC, Yu X, Zhang Z, Miles GP, Richens J, Liepman AH, Seffen K, Dupree P. Cell wall glucomannan in Arabidopsis is synthesised by CSLA glycosyltransferases, and influences the progression of embryogenesis. *Plant J.* 2009:60(3):527-538. https://doi.org/10.1111/j.1365-313X.2009.03977.x

Goubet F, Dupree P, Johansen KS. Carbohydrate gel electrophoresis. *Methods Mol Biol.* 2011:715:81-92. https://doi.org/10.1007/978-1-61779-008-9_5

Handford MG, Baldwin TC, Goubet F, Prime TA, Miles J, Yu X, Dupree P. Localisation and characterisation of cell wall mannan polysaccharides in *Arabidopsis thaliana. Planta.* 2003:218(1):27-36. https://doi.org/10.1007/s00425-003-1073-9

Herburger K, Głazowska S, Mravec J. Bricks out of the wall: polysaccharide extramural functions. *Trends Plant Sci.* 2022:27(12):1231-1241. https://doi.org/10.1016/j.tplants.2022.07.008

Hoffmann N, McFarlane HE. Xyloglucan side chains enable polysaccharide secretion to the plant cell wall. *Dev Cell.* 2024:59(19):2609-2625.e8. https://doi.org/10.1016/j.devcel.2024.06.006

Hogg D, Woo EJ, Bolam DN, McKie VA, Gilbert HJ, Pickersgill RW (2001) Crystal structure of mannanase 26A from Pseudomonas cellulosa and analysis of residues involved in substrate binding. *J Biol Chem.* 2001: 276(33):31186-31192. https://doi: 10.1074/jbc.M010290200

Iglesias-Fernández R, Rodríguez-Gacio MC, Barrero-Sicilia C, Carbonero P, Matilla A. Three endo-β-mannanase genes expressed in the micropylar endosperm and in the radicle influence germination of *Arabidopsis thaliana* seeds. *Planta.* 2011:233(1):25-36. https://doi.org/10.1007/s00425-010-1257-z

Jacobson T, Edwards M, Qiande M, Robert M, Moncrieff J, Voiniciuc C (2025) Intracellular mannanases sustain matrix polysaccharide biosynthesis. *bioRxiv.* https://doi.org/10.1101/2025.02.25.640093

Kolkas H, Burlat V, Jamet E. Immunochemical Identification of the Main Cell Wall Polysaccharides of the Early Land Plant *Marchantia polymorpha. Cells.* 2023:12(14):1833. https://doi.org/10.3390/cells12141833

Kong Y, Peña MJ, Renna L, Avci U, Pattathil S, Tuomivaara ST, Li X, Reiter WD, Brandizzi F, Hahn MG, et al. Galactose-depleted xyloglucan is dysfunctional and

leads to dwarfism in Arabidopsis. *Plant Physiol*. 2015:167(4):1296-306. https://doi.org/10.1104/pp.114.255943

Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol*. 2001:305(3):567-580. https://doi.org/10.1006/jmbi.2000.4315

Liepman AH, Wilkerson CG, Keegstra K. Expression of cellulose synthase-like (Csl) genes in insect cells reveals that CslA family members encode mannan synthases. *Proc Natl Acad Sci USA*. 2005:102(6):2221-2226. https://doi.org/10.1073/pnas.0409179102

Matthysse AG, White S, Lightfoot R. Genes required for cellulose synthesis in Agrobacterium tumefaciens. *J Bacteriol*. 1995:177(4):1069-1075. https://doi.org/10.1128/jb.177.4.1069-1075.1995

Mayer MP, Bukau B. Hsp70 chaperones: cellular functions and molecular mechanism. *Cell Mol Life Sci*. 2005:62(6):670-684. https://doi.org/10.1007/s00018-004-4464-6

Mohnen D. Pectin structure and biosynthesis. *Curr Opin Plant Biol*. 2008:11(3):266-277. https://doi.org/10.1016/j.pbi.2008.03.006

Möller S, Croning MD, Apweiler R. Evaluation of methods for the prediction of membrane spanning regions. *Bioinformatics*. 2001:17(7):646-53. https://doi.org/10.1093/bioinformatics/17.7.646

Mori K. Tripartite management of unfolded proteins in the endoplasmic reticulum. *Cell*. 2000:101(5):451-454. https://doi.org/10.1016/s0092-8674(00)80855-7

Li KL, Nakashima K, Hisata K, Satoh N. Expression and possible functions of a horizontally transferred glycosyl hydrolase gene, *GH6-1*, in *Ciona* embryogenesis. *Evodevo*. 2023:14(1):11. https://doi.org/10.1186/s13227-023-00215-x

Nicol F, His I, Jauneau A, Vernhettes S, Canut H, Höfte H. A plasma membrane-bound putative endo-1,4-beta-D-glucanase is required for normal wall assembly and cell elongation in Arabidopsis. *EMBO J*. 1998:17(19):5563-5576. https://doi.org/doi:10.1093/emboj/17.19.5563

Nishigaki N, Yoshimi Y, Kuki H, Kunieda T, Hara-Nishimura I, Tsumuraya Y, Takahashi D, Dupree P, Kotake T. Galactoglucomannan structure of Arabidopsis seed-coat mucilage in GDP-mannose synthesis impaired mutants. *Physiol Plant*. 2021:173(3):1244-1252. https://doi.org/10.1111/ppl.13519

Nonogaki H, Gee OH, Bradford KJ. A germination-specific endo-β-mannanase gene is expressed in the micropylar endosperm cap of tomato seeds. *Plant Physiol*. 2000:123(4):1235-1246. https://doi.org/10.1104/pp.123.4.1235

Oda Y, Hosokawa N, Wada I, Nagata K. EDEM as an acceptor of terminally misfolded glycoproteins released from calnexin. *Science*. 2003:299(5611):1394-1397. https://doi.org/10.1126/science.1079181

Reid JS, Edwards ME, Dickson CA, Scott C, Gidley MJ. Tobacco transgenic lines that express fenugreek galactomannan galactosyltransferase constitutively have structurally altered galactomannans in their seed endosperm cell walls. *Plant*

657 *Physiol.* 2003:131(3):1487-1495. https://doi.org/10.1104/pp.102.016840

658 **Sawake S, Tajima N, Mortimer JC, Lao J, Ishikawa T, Yu X, Yamanashi Y, Yoshimi**
659 **Y, Kawai-Yamada M, Dupree P, et al.** KONJAC1 and 2 are key factors for GDP-
660 mannose generation and affect L-ascorbic Acid and glucomannan biosynthesis in
661 Arabidopsis. *Plant Cell.* 2015:27(12):3397-3409.
662 https://doi.org/10.1105/tpc.15.00379

663 **Schröder R, Wegrzyn TF, Bolitho KM, Redgwell RJ.** Mannan transglycosylase: a
664 novel enzyme activity in cell walls of higher plants. *Planta.* 2004:219(4):590-600.
665 https://doi.org/10.1007/s00425-004-1274-x

666 **Scheller HV, Ulvskov P.** Hemicelluloses. *Annu Rev Plant Biol.* 2010:61:263-289.
667 https://doi.org/10.1146/annurev-arplant-042809-112315

668 **Standal R, Iversen TG, Coucheron DH, Fjaervik E, Blatny JM, Valla S.** A new gene
669 required for cellulose production and a gene encoding cellulolytic activity in
670 *Acetobacter xylinum* are colocalized with the *bcs* operon. *J Bacteriol.*
671 1994:176(3):665-672. https://doi.org/10.1128/jb.176.3.665-672.1994

672 **Tamura K, Shimada T, Kondo M, Nishimura M, Hara-Nishimura I.**
673 KATAMARI1/MURUS3 Is a novel golgi membrane protein that is required for
674 endomembrane organization in Arabidopsis. *Plant Cell.* 2005:17(6):1764-76.
675 https://doi.org/10.1105/tpc.105.031930

676 **Tamura K, Stecher G, Kumar S.** MEGA11: Molecular Evolutionary Genetics
677 Analysis version 11. *Mol Biol Evol.* 2021:38(7):3022-3027.
678 https://doi.org/10.1093/molbev/msab120

679 **Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD,**
680 **Winther O, Brunak S, von Heijne G, et al.** SignalP 6.0 predicts all five types of
681 signal peptides using protein language models. *Nat Biotechnol.* 2022:40(7):1023-
682 1025. https://doi.org/10.1038/s41587-021-01156-3

683 **Thorne K, Urbanowicz BR, Hahn MG** (2023) Plant cell wall glycan-directed
684 monoclonal antibodies. In: Plant cell walls - Research Milestones and Conceptual
685 Insights (Ed., Geitmann, A.). Boca Raton: CRC Press, Taylor & Francis Group, pp.
686 206-236. https://doi.org/10.1201/9781003178309

687 **Verbančič J, Lunn JE, Stitt M, Persson S.** Carbon supply and the regulation of cell
688 wall synthesis. *Mol Plant.* 2018:11(1):75-94.
689 https://doi.org/10.1016/j.molp.2017.10.004

690 **Voiniciuc C.** Modern mannan: a hemicellulose's journey. *New Phytol.*
691 2022:234(4):1175-1184. https://doi.org/10.1111/nph.18091

692 **Voiniciuc C, Schmidt MH, Berger A, Yang B, Ebert B, Scheller HV, North HM,**
693 **Usadel B, Günl M.** MUCILAGE-RELATED10 produces galactoglucomannan that
694 maintains pectin and cellulose architecture in Arabidopsis seed mucilage. *Plant*
695 *Physiol.* 2015:169(1):403-420. https://doi.org/10.1104/pp.15.00851

696 **Wang Y, Azhar S, Gandini R, Divne C, Ezcurra I, Aspeborg H.** Biochemical
697 characterization of the novel endo-beta-mannanase AtMan5-2 from *Arabidopsis*

thaliana. *Plant Sci*. 2015:**241**:151-63. https://doi.org/10.1016/j.plantsci.2015.10.002

Wang Y, Mortimer JC, Davis J, Dupree P, Keegstra K. Identification of an additional
protein involved in mannan biosynthesis. *Plant J*. 2013:73(1):105-117.
https://doi.org/10.1111/tpj.12019

Wang Y, Vilaplana F, Brumer H, Aspeborg H. Enzymatic characterization of a
glycoside hydrolase family 5 subfamily 7 (GH5_7) mannanase from *Arabidopsis*
thaliana. *Planta*. 2014:239(3):653-665. https://doi.org/10.1007/s00425-013-2005-y

Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ. An "Electronic
Fluorescent Pictograph" browser for exploring and analyzing large-scale biological
data sets. PLoS One. 2007:2(8):e718. https://doi.org/10.1371/journal.pone.0000718

Wu Q, Meng YT, Feng ZH, Shen RF, Zhu XF. The endo-beta mannase MAN7
contributes to cadmium tolerance by modulating root cell wall binding capacity in
Arabidopsis thaliana. *J Integr Plant Biol*. 2023:65(7):1670-1686.
https://doi.org/10.1111/jipb.13487

Yoshimi Y, Yu L, Cresswell R, Guo X, Echevarría-Poza A, Lyczakowski JJ, Dupree
R, Kotake T, Dupree P. Glucomannan engineering highlights the role of galactosyl
modification in fine-tuning cellulose-glucomannan interaction in Arabidopsis cell
walls. *Nat Commun*. 2025:16(1):1235. https://doi.org/10.1038/s41467-025-56626-y

Yu L, Lyczakowski JJ, Pereira CS, Kotake T, Yu X, Li A, Mogelsvang S, Skaf MS,
Dupree P. The patterned structure of galactoglucomannan suggests it may bind to
cellulose in seed mucilage. *Plant Physiol*. 2018:178(3):1011-1026.
https://doi.org/10.1104/pp.18.00709

Yu L, Yoshimi Y, Cresswell R, Wightman R, Lyczakowski JJ, Wilson LFL, Ishida
K, Stott K, Yu X, Charalambous S, Wurman-Rodrich J, Terrett OM, Brown SP,
Dupree R, Temple H, Krogh KBRM, Dupree P. Eudicot primary cell wall
glucomannan is related in synthesis, structure, and function to xyloglucan. *Plant*
Cell. 2022:34(11):4600-4622. https://doi.org/10.1093/plcell/koac238

Yu L, Shi D, Li J, Kong Y, Yu Y, Chai G, Hu R, Wang J, Hahn MG, Zhou G.
CELLULOSE SYNTHASE-LIKE A2, a glucomannan synthase, is involved in
maintaining adherent mucilage structure in Arabidopsis seed. *Plant Physiol*.
2014:164(4):1842-1856. https://doi.org/10.1104/pp.114.236596

Zhang R, Li B, Zhao Y, Zhu Y, Li L. An essential role for mannan degradation in both
cell growth and secondary cell wall formation. *J Exp Bot*. 2023:75(5):1407-1420.
https://doi.org/10.1093/jxb/erad463

Zhao Y, Song D, Sun J, Li L. *Populus* endo-beta-mannanase PtrMAN6 plays a role in
coordinating cell wall remodeling with suppression of secondary wall thickening
through generation of oligosaccharide signals. *Plant J*. 2013:74(3):473-85.
https://doi.org/10.1111/tpj.12137

Zhou Y, Jiang R, Perkins WS, Cheng Y. Morphology evolution and gelation
mechanism of alkali induced konjac glucomannan hydrogel. *Food Chem*.
2018:269:80-88. https://doi.org/10.1016/j.foodchem.2018.05.116

Zhong R, Cui D, Ye ZH. Members of the DUF231 family are *O*-acetyltransferases

catalyzing 2-*O*- and 3-*O*-acetylation of mannan. *Plant Cell Physiol.*

2018:59(11):2339-2349. https://doi.org/10.1093/pcp/pcy159

 FIGURE LEGENDS

Figure 1. Distribution of TM-type MANs in plants.

(A) Schematic models of AtMAN proteins. The signal peptide and TM region were

identified with the aid of SignalP 6.0 and TMHMM 2.0 programs, respectively. (B)

Phylogenetic tree of plant MANs. Phylogenetic relationships of MANs from

Arabidopsis and other plants including *S. moellendorffii*, *P. patens*, and *M. polymorpha*

were analyzed with MEGA software (version 11.0). Rice OsMAN5 and poplar

PtrMAN6 indicated with pink dots have HHEL and HDDL sequences at the C-terminus

instead of HDEL, respectively. SmMAN2 with the orange dot has a TM domain in the

N-terminal region that its neighbor, SmMAN3, lacks. MAN sequences from

angiosperms are shown in black whereas those from other organisms are in blue. Amt,

amborella; Lj, *Lotus japonicus*; Mp, *M. polymorpha*; Os, rice; Pp, *P. patens*; Sm, *S.*

moellendorffii. AnMAN5A from *Aspergillus nidulans* was used as the outgroup.

Numbers in the nodes represent bootstrap values. The accession numbers and

abbreviated names for these sequences are listed in Supplementary Table S2.

Figure 2. Reduced glucomannan accumulation in the cell walls of *man* mutants.

Cross sections from the bottom part of the first inflorescence stem of *man* mutants were

compared with WT and the *csla2-2 csla9-2* double mutant. The images for dark field

and glucomannan signals with LM21 antibody are shown. The bars indicate 100 μm.

Figure 3. Cell wall fractions and mannose content in *man* mutants.

(A) Relative sugar content of cell wall fractions in the *man* mutants. Sugar content of

soluble, alkali, and cellulose fractions from the bottom part of inflorescence stems was

quantified by the phenol-sulfuric acid method (Supplementary Fig. S4). (B) Mannose

content of cell wall fractions. The content was calculated based on the sugar content and

the monosaccharide composition of cell wall fractions (Supplementary Fig. S5). The

values are shown in weight per gram fresh weight (g fw). Data are mean values with

standard deviation (three biological replicates). Different letters indicate significant

differences determined with the Tukey HSD test ($p < 0.05$).

Figure 4. Change in β-GGM and AcGGM.

GGMs included in the cell walls of the bottom part of inflorescence stems were

hydrolyzed into specific oligosaccharides such as MGM, GBGM, $(GA)_2GM$

(Supplementary Fig. S6), which were labelled with a fluorescent compound then

separated in the gel. Here, *man* mutants were compared with WT, *csla2-1 csla9-1*

double mutants known to lack stem glucomannans, and the *msr1-1 msr2-1* double

mutant with reduced glucomannan (Goubet et al. 2009; Wang et al. 2013). The sugar

symbols   follow   the   international   glycan   code   nomenclature

(www.ncbi.nlm.nih.gov/glycans/snfg.html). The oligosaccharide structures are shown

on the right. The results for control reactions are shown in Supplementary Fig. S7.

Figure 5. Glucomannan particles inside cells.

(*A*) Magnified images of xylem and interfascicular fiber cells. The merged image of

calcofluor staining and glucomannan signals with LM21 antibody are shown. The bars

indicate 50 μm. Arrowheads indicate relatively large (>1 μm) glucomannan signals

found inside cells. (*B*) Numbers of glucomannan particles in the pith, xylem, and

interfascicular fiber cells. The values shown are the number of particles per cell. Data

are mean values with standard deviation (WT, n□=□6; *man6-1*, n□=□6; *man2-1 man5-*

1, n□=□6; *man2-1 man5-1 man6-1*, n□=□6; *csla2-2 csla9-2*, n□=□5). The asterisk

indicates significant difference from WT (Student's t test, *, P < 0.05). (C) Observation

by immunoelectron microscopy. Ultrathin sections prepared from the bottom part of the

inflorescence stem of WT, *man6-1* single, *man2-1 man5-1* double, and *man2-1 man5-1*

man6-1 triple mutants were observed. Glucomannan was detected with LM21 antibody.

The gold particles found in the cell walls are shown as red dots and those in the

cytoplasm including the vacuole are shown as blue dots. Representative images are

shown. CP, chloroplast; CW, cell wall; V, vacuole. The bars indicate 1 μm.

Figure 6. Subcellular localization of MAN-mNeon proteins.

Xylem and interfascicular fiber cells expressing AtMAN2TM-mNeon (A-E) and

AtMAN5TM-mNeon (F-J), and vessels expressing AtMAN6signal-mNeon (K-O) in the

top part of the inflorescence stem were observed with a confocal microscope. Images

for mNeonGreen (A, F, K), autofluorescence (B, G, L), FM 4-64FX staining (C, H, M),

and transmitted photomultiplier tube (D, I, N) were obtained and merged (E, J, O). The

bars indicate 5□μm. Schematic diagrams for each mNeonGreen (mNeon)-MAN protein

are shown above the merged images (E, J, O). The representative mNeonGreen signals

in the endomembrane system including the Golgi apparatus are indicated by yellow

arrowheads and those in vessels with by pink arrowheads. The gene constructs and the

complementation of the triple mutants with *pMAN2:MAN2TM-mNeon* and

pMAN5:MAN5TM-mNeon are shown in Supplementary Fig. S9. Images of the tissue

localization of these MAN-mNeon proteins and longitudinal sections for

AtMAN6signal-mNeon are shown in Supplementary Fig. S10.