Trait-environment interactions influence tree planting success in Peruvian tropical dry

forests

Running title Trait-environment interactions influence restoration success

Authors and affiliations

Goossens, W.[1*], Fremout, T.[1,2], Muys, B.[1,3], Soto Quispe, J.C. [4,5], Van Meerbeek, K.[1,3], Maes,

S.L.[1]

[1] Department of Earth & Environmental Sciences, Division of Forest, Nature and Landscape,
KU Leuven, Belgium
[2] Alliance Bioversity International - CIAT, Lima, Peru
[3] KU Leuven Plant Institute, KU Leuven, Belgium
[4] Universidad Nacional de Piura, Piura, Peru
[5] Servicio Nacional de Áreas Naturales Protegidas por el Estado (SERNANP), Piura, Peru

Corresponding author detail

Willem Goossens, willem.goossens@kuleuven.be

Department of Earth & Environmental Sciences, Division of Forest, Nature and Landscape, KU
Leuven
Celestijnenlaan 200E, box 2411
B-3001 Leuven, Belgium

Authors

Name	Mail	ORCID-ID
Willem Goossens	willem.goossens@kuleuven.be	0009-0006-4832-0902
Tobias Fremout	t.fremout@cgiar.org	0000-0002-0812-3027
Bart Muys	bart.muys@kuleuven.be	0000-0001-9421-527X
Juan Carlos Soto Quispe	jcarlos030119@gmail.com	0000-0002-6295-5764
Koenraad Van Meerbeek	koenraad.vanmeerbeek@kuleuven.be	0000-0002-9260-3815
Sybryn L. Maes	sybryn.maes@gmail.com	0000-0002-7168-2390

20 **Author contributions**

21 WG, SM, BM conceived and designed the research; WG, JCSQ carried out the fieldwork; WG,

22 SM, KVM analysed the data; WG drafted the manuscript; TF, BM, JCSQ, KVM, SM critically

23 reviewed the manuscript

Keywords

- Forest restoration
- Resource-use strategy
- Seedling growth and survival
- Trait-environment interactions
- Tropical dry forests

Abstract

Tropical dry forests are among the most endangered ecosystems globally, despite their critical socio-ecological importance. In recent decades, various tree-planting initiatives have been undertaken to restore these forests, yielding mixed results. To enhance restoration effectiveness, it is imperative to adopt a predictive approach, focussing on the drivers of seedling performance and their interactions. This study explores how tree species traits and environmental conditions interact to influence survival and growth rates in the Tumbes-Piura tropical dry forests of northwestern Peru. Our findings support the widely accepted *growth-mortality* hypothesis, indicating that acquisitive species characterised by a high specific leaf area and low wood density achieve faster growth but experience higher mortality compared to conservative species. According to the general resource-use theory, these acquisitive species are also more sensitive to variations in water and nutrient availability, showing steep declines in growth and survival rates as site conditions become less favourable. Interestingly, we found that the *growth-mortality* hypothesis holds along environmental gradients, with acquisitive species maintaining their high growth rates even under resource-poor conditions, but at the cost of greater vulnerability to mortality. In contrast, the growth rates of conservative species increase markedly when water and nutrients become less limiting. These insights underscore the importance of considering trait-environment interactions in planning tree-planting projects and adapting the management strategy to overall restoration objectives. By doing so, we can improve the outcomes and sustainability of restoration efforts in tropical dry forests.

Implications for practice

- The *growth-mortality* hypothesis holds along resource gradients in tropical dry forest restoration.
- Acquisitive species can exhibit high growth rates even in resource-poor sites, which comes at the cost of higher mortality. Increasing successful establishment through irrigation and fertilization will likely accelerate forest recovery in these restoration sites.
- Planting conservative species in less favourable sites is a more viable option when economic resources are limited, offering a more certain, albeit slower, forest establishment without the need for intensive management.
- Our findings highlight the importance of accounting for trait-environment interactions when evaluating restoration success in future studies.

INTRODUCTION

All around the globe, forests face significant threats from deforestation and degradation, leading to biodiversity loss, accelerating climate change and casting negative impacts on human well-being (HLPE 2017; FAO & UNEP 2020). In response to these alarming trends, there has been a surge in ambitious commitments to restore degraded forest ecosystems across all continents (Sewell et al. 2020). This restoration movement is further amplified by the current 'UN Decade on Ecosystem Restoration' (2021-2030), which underscores the urgency to revive damaged ecosystems (United Nations General Assembly 2019).

Many restoration initiatives involve the active planting of trees to speed up ecosystem recovery (Brancalion & Holl 2020; Lamb et al. 2005), which is particularly important in tropical dry forests (TDFs) that experience a dry season of several months (Mooney et al. 1995). Active planting can considerably accelerate forest establishment in these areas, where natural recovery is often slow and uncertain due to the highly fragmented state of the forests and harsh microclimatic conditions (Alvarez-Aquino & Williams-Linera 2012; Dimson & Gillespie 2020; Griscom & Ashton 2011).

Despite their global socio-ecological importance in housing numerous endemic species and supporting the livelihoods of some of the world's poorest communities (Blackie et al. 2014; Schröder et al. 2021; Banda et al. 2016), TDFs remain among the most threatened terrestrial ecosystems (Miles et al. 2006; Janzen 1988; Hoekstra et al. 2005). The urgency to restore these key systems has led to various TDF restoration initiatives (Campo et al. 2023; Dimson & Gillespie 2020; Cárdenas et al. 2022). However, tree planting as a means of TDF restoration often yields variable and disappointing outcomes in terms of growth and survival rates (Dimson & Gillespie 2020; Werden et al. 2023), despite substantial investments of resources (Birch et al. 2010; Griscom & Ashton 2011). Restoration ecology is therefore urged to become more predictive by studying the factors influencing seedling performance regarding growth and survival (Campo et al. 2023; Brudvig & Catano 2021; Lanuza et al. 2020). Both the identity of the planted species and the environmental conditions at the planting site play a crucial role in governing growth and survival rates (Rosell et al. 2022; Werden et al. 2020). Hence, understanding how these factors relate to each other is key to successfully adapting restoration designs to site-specific contexts and improving overall restoration success (Eviner

& Hawkes 2008; Zirbel & Brudvig 2020; Werden et al. 2023).

The functioning of species can be summarised by placing them along the *acquisitive-*
conservative functional trait spectrum (Díaz et al. 2004; Wright et al. 2010). By grouping
species with similar resource-use strategies, we can identify and predict their response to
environmental gradients (Pinho et al. 2019) and adapt management practices accordingly
(Eviner & Hawkes 2008). In TDFs, interactions between traits and environmental conditions
have already been used to study differences in the functionality of species assemblages along
various edaphic, climatic and topographic gradients (Méndez-Toribio et al. 2017; Comita &
Engelbrecht 2009; Pinho et al. 2019). Given that the composition of these assemblages
depends strongly on seedling performance (Pinho et al. 2019; Sterck et al. 2011), the observed
importance of trait-environment interactions indicates their significance for tree growth and
survival.

TDF species have developed distinct resource-use strategies to cope with the seasonally
limited water availability, which is widely regarded as the main limiting factor for seedling
growth and survival in these forests (Alvarez-Aquino & Williams-Linera 2012; Méndez-Toribio
et al. 2020; Markesteijn & Poorter 2009). On the one hand, drought-avoidant acquisitive
species are usually deciduous and characterised by a rapid growth response to changes in
water availability (Rosell et al. 2022). They possess relatively large transpirational tissues (i.e.
high specific leaf area (SLA)) and low wood density (WD), which enable high hydraulic
conductivity during the wet season and considerable water storage capacity to bridge short
dry periods (Poorter et al. 2019; Sterck et al. 2011; Matlaga et al. 2024). In contrast, drought-
tolerant conservative species, with higher wood density and lower SLA, prioritise protecting
their retrieved resources, resulting in lower growth rates but enhanced stress tolerance and
survival (Werden et al. 2018).

General resource-use theory predicts that acquisitive species perform better under resource-
rich, favourable conditions, but that their performance decreases rapidly when resources
become scarce (Méndez-Toribio et al. 2020; Worthy et al. 2020; Mira et al. 2023). Indeed, the
high growth rate of acquisitive species drops rapidly when water (Sterck et al. 2011) or soil
nutrient levels (Baribault et al. 2012; Reich 2014) become limiting. Low soil water availability
or high evaporative demand increases the water potential difference between soil and

atmosphere (Bauman et al. 2022; Méndez-Toribio et al. 2020), causing acquisitive species to

close their stomata to prevent hydraulic failure, resulting in lower growth rates and potentially

even carbon-starvation-induced mortality after prolonged drought (McDowell et al. 2008).

Conservative species are less sensitive to hydraulic failures and can maintain photosynthetic

activity under drier conditions (Sterck et al. 2011; Matlaga et al. 2024). Similarly, acquisitive

species require higher soil fertility, giving them a growth advantage in fertile soils but posing

challenges when soil nutrients become limiting (Worthy et al. 2020; Reich 2014). Moreover, it

has been suggested that acquisitive species are more sensitive to higher temperatures than

conservative species (Sastry & Barua 2017), although this was not confirmed by Slot and

Winter (2017). An additional important characteristic of species in TDFs is the ability to fix

atmospheric nitrogen, which gives nitrogen-fixing species a competitive advantage over non-

fixing species through increased photosynthetic activity and water-use efficiency (Pellegrini et

al. 2016; Menge & Chazdon 2016).

In summary, seedling performance is influenced by both species identity and site-specific

environmental context, and their interactions might play a pivotal role in restoration practices

(Caleño-Ruiz et al. 2023; Zirbel & Brudvig 2020; Mira et al. 2023). These interactions

potentially explain some of the observed variability in restoration outcomes (Dimson &

Gillespie 2020). Most research on drivers of tree performance and their interactions has

focussed on tropical rainforests (e.g. Matlaga et al. 2024; Worthy et al. 2020), often neglecting

the TDF biome (Campo et al. 2023; Schröder et al. 2021). With this study, we aim to improve

our understanding of how functional traits and environmental factors interact to influence

seedling performance in TDF restoration projects. We focus on the Tumbes-Piura dry forest

region of northwestern Peru, thereby contributing to improved tree-planting practices in the

area (Cerrón et al. 2019; Fremout et al. 2022).

Specifically, we evaluate the impacts of interactions between abiotic (i.e. climate and soil)

conditions and adult functional traits (i.e. WD and SLA) of ten tree species on the performance

(i.e. growth and survival) of planted seedlings across nine restoration sites over a gradient of

environmental conditions. We expect acquisitive species (with low WD and high SLA) to (1)

thrive in fertile sites with low atmospheric aridity, showing relatively low mortality and higher

growth rates, but (2) be more sensitive to low resource availability compared to conservative

species (Sterck et al. 2011; Zirbel & Brudvig 2020). In other words, we anticipate little variation

in growth and mortality rates of conservative species across the studied environmental
gradient, while acquisitive species are expected to show relatively high mortality and low
growth rates in unfavourable conditions, but better performance as site conditions improve
(Matlaga et al. 2024).

MATERIALS AND METHODS

Study sites and region

To evaluate the drivers of tree planting performance, we visited nine restoration sites in the
lowland Tumbes-Piura dry forests in northwestern Peru (Figure 1), situated between the
Pacific coast and the Andean mountain range up to altitudes of 500 m.a.s.l. (Linares-Palomino
2006). Average annual precipitation in the Tumbes-Piura region varies between 100 and 500
mm, of which around 85% is centred between January and April (Best & Kessler 1995; Cueva-
Ortiz et al. 2020). Temperature exhibits a comparable seasonal pattern, with mean annual
values varying between 20 and 26°C (Cueva-Ortiz et al. 2020). Every 3 to 7 years, the El Niño
Southern Oscillation (ENSO) brings more intense rains to the region, with precipitation levels
up to ten times higher than the climatic mean, fostering higher tree recruitment and growth
rates (Best & Kessler 1995).

The Tumbes-Piura dry forests are part of the Tumbes-Chocó-Magdalena biodiversity hotspot
(CEPF 2005) and are renowned for their endemism (Best & Kessler 1995; Linares-Palomino et
al. 2010). Nonetheless, substantial portions of these forests have been lost through
deforestation (Escribano-Avila et al. 2017), and much of the remaining patches are strongly
degraded due to overgrazing by livestock and – often illegal – logging (Fremout et al. 2020;
Linares-Palomino 2006). In recent years, the region has witnessed many restoration efforts to
reverse this trend, aiming to restore forest cover and structure, conserve biodiversity, and
raise awareness among the local population (Cerrón et al. 2019). Between August and
September 2022, we visited nine tree-planting restoration sites in the area that varied in size
(0.22 - 12.7 ha), planting year (1988 – 2022), and species composition (Supplement S1). Three
suitable sites were obtained from the project database of Cerrón et al. (2019) and
complemented with six restoration projects suggested by three restoration managers active
in the region. The sites covered a gradient in climatic conditions such as potential

evapotranspiration (1603 - 1968 mm/year), mean annual precipitation (63 – 188 mm) and

mean temperature of the wettest quarter (22.75 – 27.05 °C).

Figure 1: Aridity index map (mean annual precipitation divided by potential evapotranspiration
(MAP/PET) following Zomer et al. (2022)) and site locations in the study area in northwestern
Peru. Map generated using QGIS 3.34.0 (QGIS Development Team 2024).

Abiotic conditions

Both climate and soil conditions were incorporated to assess the importance of environmental

conditions for planted seedling performance (Supplement S1). We extracted climate variables

for every site from the CHELSA 2.1 Climate Database (Karger et al. 2017) for the period 1981

to 2010 in QGIS 3.34.0 (QGIS Development Team 2024). These variables, with a resolution of

30 arcsec (ca. 0.9 km at the equator), included mean annual temperature (MAT), mean daily

air temperatures of the wettest (BIO8) and driest (BIO9) quarter, mean annual precipitation

(MAP) and mean annual potential evapotranspiration (PET), computed by summing up

monthly averages. To account for soil variability, eight subsamples were collected at each site

and combined using the quartering method (EPA 2023) to produce one representative

composite sample (Pennock et al. 2006). The subsamples were taken at depths of 10, 20 and

30 cm from eight separate locations arranged in a zigzag pattern to ensure even distribution.

For two sites exhibiting clear soil heterogeneity, multiple composite samples were acquired

from different zones to assess within-site variance, resulting in four composite samples (i.e. 4

× 8 subsamples) at site ENTL and two at site SAPET. The composite soil samples were then air-

dried and stored in plastic bags until analysis at the *Universidad Nacional Agraria La Molina*

for key factors important for soil fertility (Supplement S1).

Functional traits

To evaluate the effect of functional traits on seedling performance, we included WD, SLA and

nitrogen fixation as explanatory variables (Supplement S2). WD [g/cm³] and SLA [cm²/g] were

measured for five species (i.e. *Neltuma pallida, Beautempsia avicenniifolia, Parkinsonia*

aculeata, Lycium boerhaviifolium and *Morisonia scabrida*), while for *Loxopterygium huasango*

only SLA was obtained in the field. For the SLA and WD values of other species, we used

literature data as we could not identify suitable candidates to sample in the field. Similarly,

the ability to fix atmospheric nitrogen was obtained from literature for all studied species.

SLA was determined by collecting ten undamaged leaves fully exposed to the sun from two to

four isolated mature individuals near the restoration sites (Cornelissen et al. 2003; Pérez-

Harguindeguy et al. 2013). We included the leaf petioles and did not subdivide compound

leaves into individual leaflets. Fresh leaf area was calculated in the software program ImageJ

(Schneider et al. 2012) following the protocol of O'Neal et al. (2002). After oven-drying the

leaves for 48 hours at 80 °C and measuring their mass, the SLA was calculated by dividing the

leaf area by the oven-dry weight. For the wood density measurements, we sampled mature

individuals in the vicinity of the planting sites with a tree increment borer (Chave 2006). The

extracted cores were stored in paper straws, dried at 80°C for 48 hours (Cornelissen et al.

2003), and XCT scanned in the Woodlab at Ghent University in Belgium (Van den Bulcke et al.

2014). The scanned images were converted to mean densities of the whole core using a

reference material of known density and air with a density assumed to be zero.

Tree growth and survival

At the nine visited sites, we measured the height of every tree (Moonlight et al. 2021). For
multi-stemmed trees, commonly found in TDFs, only the tallest stem was considered.
Additionally, we recorded the survival status of each individual, which was evaluated by
scratching the bark and observing whether the cambium displayed any signs of life (Werden
et al. 2020). In the sites with multiple composite soil samples, tree individuals were related to
their soil sampling zone. Absolute height growth rates (hereafter denoted as growth [m/year])
were computed by subtracting the obtained height by 30 cm, reported as the mean planting
height by restoration managers, and divided by the time in years between planting and
monitoring (Visser et al. 2016; Werden et al. 2020):

236
$$HR = \frac{Height - 30\ cm}{2022 - Planting\ year}$$

237   During the survival assessment, we could only evaluate the status of individuals present at the
238   time of monitoring and hence omitted multiple empty spots, which causes an overestimation
239   of survival rates and might potentially induce bias by favouring some species more than
240   others.

Data analysis

Overall, ten species were present in at least two sites and are included in our analysis. For two
sites established in the monitoring year (i.e. SAPET and SICH), we only evaluated survivorship,
as measured heights are expected to represent planting size and not (yet) outcomes of trait-
environment interactions (Visser et al. 2016), which could increase the signal-to-noise ratio
and hamper capturing significant effects (Rodríguez-Alarcón et al. 2024). Across four other
sites, 346 negative and 33 zero height growth rate values were removed (Werden et al. 2020),
as these were likely caused by tree cutting or animal damage. Additionally, we removed all 27
growth rate observations considered to be an outlier within the site following Tukey (1977).
In the end, we obtained a dataset of 5514 tree individuals for the survival assessment, which,
after the removal of dead individuals, non-positive height values and two sites, resulted in
2672 height rate observations (Supplement S3). All explanatory factors, except for the

categorical nitrogen fixation, were standardized, facilitating model fitting and convergence
and allowing for direct comparison of predictor effect sizes (Zuur et al. 2009).

Generalised linear mixed models (GLMMs) were then employed to evaluate the drivers of
growth and survival variability in tropical dry forest restoration, with site identity as a random
intercept to incorporate the ecological assumption that within-site observations are more
closely related (Banin et al. 2023). Since WD and SLA were highly negatively correlated
(Pearson ρ < -0.7), we retained only the former, as it is phylogenetically more conserved,
shows less plastic variability and is not affected by the timing of the measurements (Poorter
et al. 2018; Derroire et al. 2018; Slot & Winter 2017). Additionally, the ability to fix nitrogen
(Nfix) was added to our model. We expanded our functional trait model with potential
evapotranspiration (PET), the mean temperature in the wettest quarter (BIO8) and the
presence of organic matter (OM). Due to the strong correlations between most environmental
factors, we could only include a limited amount of explanatory variables. Since irrigation was
anticipated to be a more significant factor in water availability than mean annual precipitation
(MAP) at our restoration sites, and as MAP differed relatively little between the sites
(Supplement S1), we opted to use PET as a measure of atmospheric aridity (Méndez-Toribio
et al. 2020), rather than the conventional aridity index, which is calculated by dividing annual
precipitation by potential evapotranspiration (Zomer et al. 2022). In addition, we included
BIO8 to account for energy availability in the rainy season (Slot & Winter 2017) and OM as an
indicator of soil fertility and water-holding capacity (Wood et al. 2016). Interaction effects with
WD were included for all abiotic predictor factors except for BIO8, as this increased the
variance inflation factors (VIFs) considerably (Kim 2019). We used the same model structure
for both growth and survival rates:

$$\text{Growth (or survival)} \sim \text{Nfix} + \text{WD*PET} + \text{WD* OM} + \text{BIO8} + (1|\text{Site})$$

Height growth rates were analysed using the *lme4* package (Bates et al. 2015) with gamma
error distribution and Restricted Maximum Likelihood (REML) estimation (Banin et al. 2023;
Luke 2017). Significance of the results was assessed according to Satterwaite's degrees of
freedom method included in the *lmerTest* package (Kuznetsova et al. 2017). Survival rates
were modelled via a binomial error distribution with *cloglog* link function (Banin et al. 2023)
using the *glmmTMB* package (Brooks et al. 2017). We used the *piecewiseSEM* package

(Lefcheck 2016) to compute the marginal (R^2_m) and conditional (R^2_c) coefficients of
determination to assess the goodness-of-fit (Nakagawa & Schielzeth 2013; Werden et al.
2020). The former quantifies the variance explained by the fixed effects, while the latter
evaluates the variance explained by both the fixed and random factors. To test the validity of
our findings, we also ran the same model with SLA instead of WD (Supplement S4).
Furthermore, we built a model to assess the interactive effect of BIO8 and WD, thereby
omitting the interaction between WD and PET due to the high VIFs (Supplement S5). Lastly,
we ran our survival model for a reduced dataset, in which the sites established in the year of
monitoring were excluded (Supplement S6). These additional analyses confirmed our main
results. All analyses were performed in R 4.2.3 (R Core Team 2023).

RESULTS

The performance of planted tree seedlings varied considerably between species and sites
(Supplement S3) and was significantly influenced by trait-environment interactions (Table 1;
Figure 2; Supplement S7). First, the interaction between WD and atmospheric aridity had a
considerable influence on height growth rates. Specifically, the growth of species with a higher
WD was more negatively affected by an increase in PET than that of species with a lower WD
(PET x WD -0.17 $p < 0.001$). Likewise, a significant interaction existed between WD and OM,
as the height growth of species with a low WD remained relatively stable across the OM
gradient, while that of species with high WD decreased rapidly when OM became limiting (OM
x WD 0.20 $p < 0.001$). Besides these significant interaction effects, we observed important
main effects for Nfix (0.35 $p < 0.001$), WD (-0.66 $p < 0.001$), OM (0.14 $p < 0.05$), PET (-0.49
$p < 0.01$) and BIO8 (0.52 $p < 0.01$). Overall, the fixed effects in our model accounted for 56.3%
of the variance in height growth (R^2_m), while adding site as a random factor increased the
explained variance to 58.3% (R^2_c).

Similarly, trait-environment interactions significantly altered survival rates (Table 1 and S5;
Figure 2). In particular, species with a low WD showed rapidly growing survivorship when the
organic matter content increased, while survival rates of species with a high WD were less
affected (WD x OM -0.13 $p < 0.001$). We also observed a significant positive interaction
between WD and PET (WD x PET 0.12 $p < 0.01$). Moreover, the main effect of OM was positive
(OM 0.08 $p < 0.05$), as was the case for WD (0.17 $p < 0.001$) and Nfix (0.45 $p < 0.001$). A higher

temperature in the wet season decreased the survival rate (BIO8 -0.32 $p < 0.05$). In contrast to height growth, our model could only explain limited variation in survival rates through fixed (R^2_m 3.0%) and both fixed and random effects (R^2_c 11.3%).

Figure 2: Significant interaction effects between wood density (WD [g/cm³]) and annual potential evapotranspiration (PET [mm/year]) (a) and WD and organic matter content (OM [%]) (b) on height growth. Effect of the interaction between WD and potential evapotranspiration (c) and between WD and organic matter (d) on survival rate.

Driver	Height growth rate		Survival rate	
	β	CI	β	CI
Nfix	0.35***	[0.28, 0.41]	0.45***	[0.38, 0.51]
WD	-0.66***	[-0.70, -0.62]	0.17***	[0.12, 0.22]
OM	0.14*	[0.08, 0.19]	0.08*	[0.04, 0.12]
PET	-0.49**	[-0.58, -0.40]	-0.08	[-0.23, 0.08]
BIO8	0.52**	[0.44, 0.61]	-0.32*	[-0.48, -0.16]
WD*OM	0.20***	[0.16, 0.24]	-0.13***	[-0.17, -0.10]
WD*PET	-0.17***	[-0.20, -0.14]	0.12**	[0.07, 0.16]
R^2_m (%)	56.3		3.0	
R^2_c (%)	58.3		11.3	

DISCUSSION

Our results demonstrate the importance of incorporating trait-environment interactions
when assessing tree planting performance in restoration projects. While several of the studied
environmental drivers (potential evapotranspiration (PET) and organic matter (OM)) and
functional traits (wood density (WD)) independently influenced growth and survival, the
presence of interactions between these drivers implies that evaluating main effects alone is
insufficient to predict overall TDF restoration success.

Contrasting wood density effects on tree survival and growth rates

Our findings revealed significant but opposing effects of wood density on height growth and
survival, lending empirical support to the *growth-mortality hypothesis* (Wright et al. 2010;
Sterck et al. 2011) and its relevance to TDF restoration (Werden et al. 2023; Rosell et al. 2022).
Specifically, conservative species with higher WD exhibited slower growth but enhanced
survival, which can be attributed to their greater investment in dense leaf and wood tissues
(Markesteijn & Poorter 2009). While this enables better tolerance of stress and disturbances,
increasing survivorship, the strategy comes at the expense of lower resource allocation
towards growth (Chave et al. 2009; Rosell et al. 2022). In addition to wood density, the ability

to fix atmospheric nitrogen enhanced both growth and survival rates, which confirms the
suggested competitive advantage of nitrogen-fixers over non-fixer species (Menge & Chazdon
2016). Nitrogen fixation increases foliar nitrogen contents, enabling higher photosynthetic
activity and water-use efficiency (Pellegrini et al. 2016; Poorter & Markesteijn 2008), and
supports the production of nitrogen-containing defensive compounds, thereby reducing
insect-induced mortality (McDowell et al. 2008; Menge & Chazdon 2016).

The *growth-mortality* hypothesis holds along environmental gradients

General resource-use theory predicts high growth and survival rates for acquisitive species
with a lower WD under favourable conditions, but a rapidly decreasing performance as water
and nutrients become limiting (Sterck et al. 2011; Baribault et al. 2012; Caleño-Ruiz et al.
2023). Conversely, conservative species are better able to tolerate drought or low nutrient
availability, resulting in high survival rates and stable – but rather low – growth along the
resource gradients (Chave et al. 2009; Pinho et al. 2019). Indeed, we observed conservative
species with higher WD to have more stable survival rates along the gradients in atmospheric
aridity and organic matter, here used as a proxy for nutrient content (Wood et al. 2016). This
supports, at least for survivorship, the theory that conservative species are more tolerant of
lower resource availabilities. For growth, however, we observed an opposite impact than
anticipated: the growth of acquisitive species decreased less rapidly along the resource
gradients than the growth of more conservative species. These results suggest that the
growth-mortality hypothesis holds along environmental gradients, as acquisitive species are
even under resource-poor conditions able to retrieve sufficient resources and allocate them
to height growth. Nonetheless, their resource-use strategy results in higher survivorship
vulnerability to dry and low nutrient conditions, indicating that few acquisitive species are able
to survive the harsh initial conditions in TDF restoration sites (Poorter et al. 2019; Schwartz et
al. 2022).

The high and stable growth rates of acquisitive species along the resource gradients (i.e. both
water and organic matter availability) are surprising and indicate that those individuals that
manage to establish achieve high growth rates independent of site resource availability,
suggesting a competitive growth advantage over slow-growing conservative species (Werden
et al. 2023). First, the fast-growing acquisitive species likely invest heavily in belowground

biomass and thereby increase their rooting density (Fagundes et al. 2022; Méndez-Toribio et al. 2020), which gives them an enhanced growth capacity (Werden et al. 2023) and makes them less susceptible to changes in soil fertility or atmospheric aridity (Markesteijn & Poorter 2009). Once established and with a well-developed rooting system, acquisitive species may thus retrieve sufficient nutrients and water to invest in height growth compared to conservative species. However, due to the decoupling between above- and below-ground resource-use strategies (Werden et al. 2023; Worthy et al. 2020), we are not able to confirm this hypothesis due to the lack of root trait data, although it is supported by other studies reporting a relationship between acquisitive resource use and rooting density (Comita & Engelbrecht 2009; Méndez-Toribio et al. 2020). Secondly, the stable growth of acquisitive species along the resource gradient can also be explained by their high growth rates during the short wet season (Mendivelso et al. 2014; Poorter et al. 2018), after which many of them quickly shed their leaves to avoid drought stress (Fagundes et al. 2022; Poorter & Markesteijn 2008), making the effect of atmospheric aridity (high PET) on height growth less relevant (Rodríguez-Alarcón et al. 2024). Conservative species, on the other hand, remain active in (part of) the dry season, extending their growing period but also increasing their susceptibility to higher evaporative demand (McDowell et al. 2008; Wright et al. 2010). This forces them to adapt their growth rates to existing conditions, resulting in a higher sensitivity to potential evapotranspiration.

Acquisitive species benefit more from higher wet season temperatures

Higher temperatures in the wettest quarter decreased survivorship but positively affected growth rates. While the negative effect on survival rates aligns with our expectations, as higher temperatures can exacerbate drought and hydraulic failure (McDowell et al. 2008), the positive effect on growth rates is surprising and contradicts previous studies (Méndez-Toribio et al. 2017; Mendivelso et al. 2014). In TDFs, however, higher temperatures during the growing season can benefit tree growth by enhancing energy availability, thereby increasing photosynthetic activity (Slot & Winter 2017). In addition, we anticipated that acquisitive species would be more sensitive than conservative species to temperature increases, with rapidly decreasing performance under warmer conditions (Sastry & Barua 2017). Although we could not include this interaction in our main model due to high VIFs, testing it separately

revealed that growth rates of acquisitive species actually increased more rapidly along the

temperature gradient than was the case for conservative species. While this contradicts our

initial expectations, it supports the hypothesis by Slot and Winter (2017) that acquisitive

species benefit more from increased energy availability, likely due to their higher electron

transport capacity and larger temperature-response plasticity. It also corroborates our

hypothesis that acquisitive species are less sensitive to high atmospheric aridity by focusing

on achieving high growth rates in the favourable, wet period with sufficient water and energy

availability.

Study limitations and future research avenues

Overall, our results indicate that the *growth-mortality* hypothesis holds along resource

gradients, with acquisitive species able to sustain their high growth rates even when water or

nutrient content becomes limiting, although they exhibit higher vulnerability to mortality

under these conditions compared to conservative species. Despite the importance of this

empirical evidence supporting the significance of trait-environment interactions on seedling

performance in restoration projects, several limitations must be considered. First, survival

rates might be biased as we could only include remaining dead individuals with certain species

identification and had to omit empty planting spots, which may have resulted in an

overrepresentation of conservative species with more durable stems (Chave et al. 2009). This

could also explain the low explanatory power of the survival model. Nonetheless, our results

align with our expectations and are corroborated by previous studies (Wright et al. 2010; Díaz

et al. 2004). Second, the observational nature of our study prevented the assessment of key

management strategies in TDF restoration, such as irrigation (Dimson & Gillespie 2020), which

could have influenced seedling performance. However, our use of potential

evapotranspiration as a proxy for atmospheric aridity provided an indirect measure of drought

stress (Méndez-Toribio et al. 2020), independent of management interactions. Third, we could

only incorporate one functional trait in our models. While incorporating multi-dimensional

trait axes could potentially improve explanatory power (Matlaga et al. 2024; Rodríguez-

Alarcón et al. 2024), our results remained consistent when substituting WD with SLA, probably

due to their strong correlation, further reinforcing our conclusions.

Future research on trait-environment interactions should focus on experimental studies

across resource gradients in the Tumbes-Piura TDFs, using a study design with known initial species numbers and standardised management practices. Furthermore, we encourage future restoration research to incorporate trait plasticity (Derroire et al. 2018; Poorter et al. 2018) and ontogenetic variability (Metz et al. 2023; Poorter & Markesteijn 2008), since they both enable individuals to acclimate to local abiotic conditions and hence might influence trait-environment interactions in restoration projects (Lanuza et al. 2020; Worthy et al. 2020). Additionally, accounting for the high soil and microclimatic heterogeneity of TDFs in future studies can shed additional light on the importance of trait-environment interactions in restoration practices (Rodríguez-Alarcón et al. 2024; Caleño-Ruiz et al. 2023; Schwartz et al. 2022).

Given our observation that seedling performance depends on the interaction between environmental variables and species' functional traits, our findings underscore the importance of context dependence in restoration projects and highlight that species selection should depend on both the site's environmental conditions and overarching restoration objectives (Zirbel & Brudvig 2020; Fremout et al. 2022). This study lays the foundation for future research on these interactions, particularly within the underexplored Tumbes-Piura TDFs.

Acknowledgements

The authors express their gratitude to Michael Perring (UK Centre for Ecology & Hydrology) for his valuable critique of the manuscript and to Nora Grados and Gaston Cruz (UDEP) for their assistance with legal documentation and the analysis of leaf and wood samples. The authors also acknowledge the cooperation of restoration managers who facilitated the research on their sites, as well as the Peruvian Forest Service (SERFOR) for granting research and botanical sampling permissions. The fieldwork was made possible through a VLIR-UOS scholarship (REI-2022-01-01).

Literature cited

Alvarez-Aquino C, Williams-Linera G (2012) Seedling Survival and Growth of Tree Species: Site Condition and Seasonality in Tropical Dry Forest Restoration. Botanical Sciences 90:341–351

Banda KR, Delgado-Salinas A, Dexter KG, Linares-Palomino R, Oliveira-Filho A, Prado D, et al. (2016) Plant diversity patterns in neotropical dry forests and their conservation implications. Science 353

Banin LF, Raine EH, Rowland LM, Chazdon RL, Smith SW, Rahman NEB, et al. (2023) The road to recovery: a synthesis of outcomes from ecosystem restoration in tropical and sub-tropical Asian forests. Philosophical Transactions of the Royal Society B 378

Baribault TW, Kobe RK, Finley AO (2012) Tropical tree growth is correlated with soil phosphorus, potassium, and calcium, though not for legumes. Ecological Monographs 82:189–203

Bates D, Mächler M, Bolker BM, Walker SC (2015) Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software 67:1–48

Bauman D, Fortunel C, Cernusak LA, Bentley LP, McMahon SM, Rifai SW, et al. (2022) Tropical tree growth sensitivity to climate is driven by species intrinsic growth rate and leaf traits. Global Change Biology 28:1414–1432

Best BJ, Kessler M (1995) Biodiversity and Conservation in Tumbesian Ecuador and Peru.

Birch JC, Newton AC, Aquino CA, Cantarello E, Echeverría C, Kitzberger T, et al. (2010) Cost-effectiveness of dryland forest restoration evaluated by spatial analysis of ecosystem services. Proceedings of the National Academy of Sciences of the United States of America 107:21925–21930

Blackie R, Baldauf C, Gautier D, Gumbo D, Kassa H, Parthasarathy N, et al. (2014) Tropical dry forests: The state of global knowledge and recommendations for future

484   research.

485   Brancalion PHS, Holl KD (2020) Guidance for successful tree planting initiatives. Journal
486   of Applied Ecology 57:2349–2361

487   Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et al. (2017)
488   glmmTMB balances speed and flexibility among packages for zero-inflated
489   generalized linear mixed modeling. R Journal 9:378–400

490   Brudvig LA, Catano CP (2021) Prediction and uncertainty in restoration science.
491   Restoration Ecology e13380

492   Van den Bulcke J, Wernersson ELG, Dierick M, Van Loo D, Masschaele B, Brabant L, et al.
493   (2014) 3D tree-ring analysis using helical X-ray tomography. Dendrochronologia
494   32:39–46

495   Caleño-Ruiz BL, Garzón F, López-Camacho R, Pizano C, Salinas V, González-M R (2023) Soil
496   resources and functional trait trade-offs determine species biomass stocks and
497   productivity in a tropical dry forest. Frontiers in Forests and Global Change
498   6:1028359

499   Campo J, Giardina CP, Dirzo R (2023) Tropical Dry Forest Restoration in an Era of Global
500   Change: Ecological and Social Dimensions. Sustainability 2023, Vol. 15, Page 3052
501   15:3052

502   Cárdenas CD, Varón-García D, Suárez-Rodríguez F, Pizano C (2022) Overcoming the
503   regeneration barriers of tropical dry forest: effects of water stress and herbivory on
504   seedling performance and allocation of key tree species for restoration. Journal of
505   Tropical Ecology 38:210–218

506   CEPF (2005) Protecting Nature's Hotspots for People and Prosperity. Annual report 2005.

507   Cerrón J, Fremout T, Thomas E, Cornelius JP, Atkinson R (2019) Experiencias de
508   restauración y fuentes semilleras en el bosque seco tropical del norte del Perú:
509   Estado actual y lecciones aprendidas.

Chave J (2006) Measuring wood density for tropical forest trees. A field manual.

Chave J, Coomes D, Jansen S, Lewis SL, Swenson NG, Zanne AE (2009) Towards a
worldwide wood economics spectrum. Ecology Letters 12:351–366

Comita LS, Engelbrecht BMJ (2009) Seasonal and spatial variation in water availability
drive habitat associations in a tropical forest. Ecology 90:2755–2765

Cornelissen JHC, Lavorel B S, Garnier B E, Díaz C S, Buchmann N, Gurvich DE, et al. (2003)
A handbook of protocols for standardised and easy measurement of plant functional
traits worldwide. Australian Journal of Botany 51:335–380

Cueva-Ortiz J, Espinosa CI, Aguirre-Mendoza Z, Gusmán-Montalván E, Weber M,
Hildebrandt P (2020) Natural Regeneration in the Tumbesian Dry Forest:
Identification of the Drivers Affecting Abundance and Diversity. Scientific Reports
2020 10:1 10:1–13

Derroire G, Powers JS, Hulshof CM, Cárdenas Varela LE, Healey JR (2018) Contrasting
patterns of leaf trait variation among and within species during tropical dry forest
succession in Costa Rica. Scientific reports 8

Díaz S, Hodgson JG, Thompson K, Cabido M, Cornelissen JHC, Jalili A, et al. (2004) The
plant traits that drive ecosystems: Evidence from three continents. Journal of
Vegetation Science 15:295–304

Dimson M, Gillespie TW (2020) Trends in active restoration of tropical dry forest:
Methods, metrics, and outcomes. Forest Ecology and Management 467:118150

EPA (2023) Soil Sampling.

Escribano-Avila G, Cervera L, Ordóñez-Delgado L, Jara-Guerrero A, Amador L, Paladines
B, et al. (2017) Biodiversity patterns and ecological processes in Neotropical dry
forest: the need to connect research and management for long-term conservation.
http://www.tandfonline.com/action/authorSubmission?journalCode=tneo20&pag
e=instructions 3:107–116

Eviner VT, Hawkes C V. (2008) Embracing Variability in the Application of Plant–Soil
Interactions to the Restoration of Communities and Ecosystems. Restoration
Ecology 16:713–729

Fagundes MV, Souza AF, Oliveira RS, Ganade G (2022) Functional traits above and below
ground allow species with distinct ecological strategies to coexist in the largest
seasonally dry tropical forest in the Americas. Frontiers in Forests and Global Change
5:930099

FAO and UNEP (2020) The State of the World's Forests 2020. FAO and UNEP

Fremout T, Thomas E, Gaisberger H, Van Meerbeek K, Muenchow J, Briers S, et al. (2020)
Mapping tree species vulnerability to multiple threats as a guide to restoration and
conservation of tropical dry forests. Global Change Biology 26:3552–3568

Fremout T, Thomas E, Taedoumg H, Briers S, Gutiérrez-Miranda CE, Alcázar-Caicedo C, et
al. (2022) Diversity for Restoration (D4R): Guiding the selection of tree species and
seed sources for climate-resilient restoration of tropical forest landscapes. Journal
of Applied Ecology 59:664–679

Griscom HP, Ashton MS (2011) Restoration of dry tropical forests in Central America: A
review of pattern and process. Forest Ecology and Management 261:1564–1579

HLPE (2017) Sustainable Forestry for Food Security and Nutrition. A report by the High
Level Panel of Experts on Food Security and Nutrition of the Committee on World
Food Security, Rome.

Hoekstra JM, Boucher TM, Ricketts TH, Roberts C (2005) Confronting a biome crisis: global
disparities of habitat loss and protection. Ecology Letters 8:23–29

Janzen DH (1988) Tropical dry forests: The most endangered major tropical ecosystem.
In: Biodiversity. Wilson, EO & Peter, FM, editors. National Academy Press,
Washington, DC pp. 130–137.

Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, et al. (2017)
Climatologies at high resolution for the earth's land surface areas. Scientific Data

2017 4:1 4:1–20

Kim JH (2019) Multicollinearity and misleading statistical results. Korean Journal of
565   Anesthesiology 72:558

566   Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest Package: Tests in Linear
567   Mixed Effects Models. Journal of Statistical Software 82:1–26

568   Lamb D, Erskine PD, Parrotta JA (2005) Restoration of degraded tropical forest
569   landscapes. Science (New York, N.Y.) 310:1628–1632

570   Lanuza OR, Espelta JM, Peñuelas J, Peguero G (2020) Assessing intraspecific trait
571   variability during seedling establishment to improve restoration of tropical dry
572   forests. Ecosphere 11:e03052

573   Lefcheck JS (2016) piecewiseSEM: Piecewise structural equation modelling in r for
574   ecology, evolution, and systematics. Methods in Ecology and Evolution 7:573–579

575   Linares-Palomino R (2006) Phytogeography and Floristics of Seasonally Dry Tropical
576   Forests in Peru. 257–279

577   Linares-Palomino R, Kvist LP, Aguirre-Mendoza Z, Gonzales-Inca C (2010) Diversity and
578   endemism of woody plant species in the Equatorial Pacific seasonally dry forests.
579   Biodiversity and Conservation 19:169–185

580   Luke SG (2017) Evaluating significance in linear mixed-effects models in R. Behavior
581   research methods 49:1494–1502

582   Markesteijn L, Poorter L (2009) Seedling root morphology and biomass allocation of 62
583   tropical tree species in relation to drought- and shade-tolerance. Journal of Ecology
584   97:311–325

585   Matlaga D, Lammerant R, Hogan JA, Uriarte M, Rodriguez-Valle C, Zimmerman JK, et al.
586   (2024) Survival, growth, and functional traits of tropical wet forest tree seedlings
587   across an experimental soil moisture gradient in Puerto Rico. Ecology and Evolution
588   14

McDowell N, Pockman WT, Allen CD, Breshears DD, Cobb N, Kolb T, et al. (2008)
Mechanisms of plant survival and mortality during drought: why do some plants
survive while others succumb to drought? New Phytologist 178:719–739

Méndez-Toribio M, Ibarra-Manríquez G, Navarrete-Segueda A, Paz H (2017) Topographic
position, but not slope aspect, drives the dominance of functional strategies of
tropical dry forest trees. Environmental Research Letters 12:085002

Méndez-Toribio M, Ibarra-Manríquez G, Paz H, Lebrija-Trejos E (2020) Atmospheric and
soil drought risks combined shape community assembly of trees in a tropical dry
forest. Journal of Ecology 108:1347–1357

Mendivelso HA, Camarero JJ, Gutiérrez E, Zuidema PA (2014) Time-dependent effects of
climate and drought on tree growth in a Neotropical dry forest: Short-term
tolerance vs. long-term sensitivity. Agricultural and Forest Meteorology 188:13–23

Menge DNL, Chazdon RL (2016) Higher survival drives the success of nitrogen-fixing trees
through succession in Costa Rican rainforests. New Phytologist 209:965–977

Metz MR, Wright SJ, Zimmerman JK, Hernandéz A, Smith SM, Swenson NG, et al. (2023)
Functional traits of young seedlings predict trade-offs in seedling performance in
three neotropical forests. Journal of Ecology 111:2568–2582

Miles L, Newton AC, DeFries RS, Ravilious C, May I, Blyth S, et al. (2006) A global overview
of the conservation status of tropical dry forests. Journal of Biogeography 33:491–
505

Mira E, Cochard H, Evette A, Dulormne M (2023) Growth, Xylem Vulnerability to
Cavitation and Leaf Cell Response to Dehydration in Tree Seedlings of the Caribbean
Dry Forest. Forests 14:697

Mooney HA, Bullock SH, Medina E (1995) Introduction. In: Seasonally Dry Tropical
Forests. Bullock, SH, Mooney, HA, & Medina, E, editors. Cambridge University Press,
Cambridge pp. 1–8.

Moonlight P, Banda-R K, Phillips OL, Dexter KG, Pennington RT, Baker TR, et al. (2021)

Supplementarty Material: The DryFlor field manual for plot establishment and
remeasurement. Plants, People, Planet 3:295–300

Nakagawa S, Schielzeth H (2013) A general and simple method for obtaining R2 from
generalized linear mixed-effects models. Methods in Ecology and Evolution 4:133–
142

O'Neal ME, Landis DA, Isaacs R (2002) An Inexpensive, Accurate Method for Measuring
Leaf Area and Defoliation Through Digital Image Analysis. Journal of Economic
Entomology 95:1190–1194

Pellegrini AFA, Staver AC, Hedin LO, Charles-Dominique T, Tourgee A (2016) Aridity, not
fire, favors nitrogen-fixing plants across tropical savanna and forest biomes. Ecology
97:2177–2183

Pennock D, Yates T, Braidek J (2006) Soil sampling designs. In: Soil Sampling and Methods
of Analysis: Second Edition. CRC Press pp. 1–14.

Pérez-Harguindeguy NA, Díaz SA, Garnier EB, Lavorel SC, Poorter HD, Jaureguiberry PA,
et al. (2013) New handbook for standardised measurement of plant functional traits
worldwide. Australian Journal of Botany 61:167–234

Pinho BX, Tabarelli M, Engelbrecht BMJ, Sfair J, Melo FPL (2019) Plant functional assembly
is mediated by rainfall and soil conditions in a seasonally dry tropical forest. Basic
and Applied Ecology 40:1–11

Poorter L, Castilho C V., Schietti J, Oliveira RS, Costa FRC (2018) Can traits predict
individual growth performance? A test in a hyperdiverse tropical forest. New
Phytologist 219:109–121

Poorter L, Markesteijn L (2008) Seedling Traits Determine Drought Tolerance of Tropical
Tree Species. Biotropica 40:321–331

Poorter L, Rozendaal DMA, Bongers F, de Almeida-Cortez JS, Almeyda Zambrano AM,
Álvarez FS, et al. (2019) Wet and dry tropical forests show opposite successional
pathways in wood density but converge over time. Nature Ecology & Evolution 2019

3:6 3:928–934

QGIS Development Team (2024) QGIS Geographic Information System.

R Core Team (2023) R: A Language and Environment for Statistical Computing.

Reich PB (2014) The world-wide 'fast–slow' plant economics spectrum: a traits manifesto.
Journal of Ecology 102:275–301

Rodríguez-Alarcón S, González-M R, Carmona CP, Tordoni E (2024) Trait–growth
relationships in Colombian tropical dry forests: Incorporating intraspecific variation
and trait interactions. Journal of Vegetation Science 35:e13233

Rosell JA, Olson ME, Martínez-Garza C, Martínez-Méndez N (2022) Functional Diversity
in Woody Organs of Tropical Dry Forests and Implications for Restoration.
Sustainability (Switzerland) 14:8362

Sastry A, Barua D (2017) Leaf thermotolerance in tropical trees from a seasonally dry
climate varies along the slow-fast resource acquisition spectrum. Scientific Reports
2017 7:1 7:1–11

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image
analysis. Nature Methods 2012 9:7 9:671–675

Schröder JM, Ávila Rodríguez LP, Günter S (2021) Research trends: Tropical dry forests:
The neglected research agenda? Forest Policy and Economics 122:102333

Schwartz NB, Medvigy D, Tijerin J, Pérez-Aviles D, Rivera-Polanco D, Pereira D, et al.
(2022) Intra-annual variation in microclimatic conditions in relation to vegetation
type and structure in two tropical dry forests undergoing secondary succession.
Forest Ecology and Management 511:120132

Sewell A, Van Der Esch S, Löwenhardt H (2020) Goals and Commitments for the
Restoration Decade: A Global Overview of Countries' Restoration Commitments
under the Rio Conventions and Other Pledges. The Hague

Slot M, Winter K (2017) In situ temperature response of photosynthesis of 42 tree and

liana species in the canopy of two Panamanian lowland tropical forests with
contrasting rainfall regimes. New Phytologist 214:1103–1117

Sterck F, Markesteijn L, Schieving F, Poorter L (2011) Functional traits determine trade-
offs and niches in a tropical forest community. Proceedings of the National Academy
of Sciences of the United States of America 108:20627–20632

Tukey J (1977) Exploratory data analysis. Addison-Wesley, Reading, MA

United Nations General Assembly (2019) General Assembly official records, 73rd session:
69th plenary meeting.

Visser MD, Bruijning M, Wright SJ, Muller-Landau HC, Jongejans E, Comita LS, et al. (2016)
Functional traits as predictors of vital rates across the life cycle of tropical trees.
Functional Ecology 30:168–180

Werden LK, Averill C, Crowther TW, Calderón-Morales E, Toro L, Alvarado JP, et al. (2023)
Below-ground traits mediate tree survival in a tropical dry forest restoration.
Philosophical Transactions of the Royal Society B: Biological Sciences 378

Werden LK, Becknell JM, Powers JS (2018) Edaphic factors, successional status and
functional traits drive habitat associations of trees in naturally regenerating tropical
dry forests. Functional Ecology 32:2766–2776

Werden LK, Calderón-Morales E, Alvarado J. P, Gutiérrez L. M, Nedveck DA, Powers JS
(2020) Using large-scale tropical dry forest restoration to test successional theory.
Ecological Applications 30:e02116

Wood SA, Sokol N, Bell CW, Bradford MA, Naeem S, Wallenstein MD, et al. (2016)
Opposing effects of different soil organic matter fractions on crop yields. Ecological
Applications 26:2072–2085

Worthy SJ, Laughlin Daniel C, Zambrano Jenny, N Uma MI, Zhang Caicai, Lin Luxiang, et
al. (2020) Alternative designs and tropical tree seedling growth performance
landscapes.

Wright SJ, Kitajima K, Kraft NJB, Reich PB, Wright IJ, Bunker DE, et al. (2010) Functional
traits and the growth-mortality trade-off in tropical trees. Ecology 91:3664–3674

Zirbel CR, Brudvig LA (2020) Trait–environment interactions affect plant establishment
success during restoration. Ecology 101:e02971

Zomer RJ, Xu J, Trabucco A (2022) Version 3 of the Global Aridity Index and Potential
Evapotranspiration Database. Scientific Data 2022 9:1 9:1–15

Zuur AF, Ieno EN, Walker N, Saveliev AA, Smith GM (2009) Mixed effects models and
extensions in ecology with R. Springer New York, New York, NY