Trait-environment interactions influence tree planting success in Peruvian tropical dry

forests

Running title Trait-environment interactions influence restoration success

Authors and affiliations

Goossens, W.[1*], Fremout, T.[1,2], Muys, B.[1,3], Soto Quispe, J.C. [4,5], Van Meerbeek, K.[1,3], Maes,

S.L.[1]

[1] Department of Earth & Environmental Sciences, Division of Forest, Nature and Landscape,
KU Leuven, Belgium
[2] Alliance Bioversity International - CIAT, Lima, Peru
[3] KU Leuven Plant Institute, KU Leuven, Belgium
[4] Universidad Nacional de Piura, Piura, Peru
[5] Servicio Nacional de Áreas Naturales Protegidas por el Estado (SERNANP), Piura, Peru

Corresponding author detail

Willem Goossens, willem.goossens@kuleuven.be

Department of Earth & Environmental Sciences, Division of Forest, Nature and Landscape, KU
Leuven
Celestijnenlaan 200E, box 2411
B-3001 Leuven, Belgium

Authors

Name	Mail	ORCID-ID
Willem Goossens	willem.goossens@kuleuven.be	0009-0006-4832-0902
Tobias Fremout	t.fremout@cgiar.org	0000-0002-0812-3027
Bart Muys	bart.muys@kuleuven.be	0000-0001-9421-527X
Juan Carlos Soto Quispe	jcarlos030119@gmail.com	0000-0002-6295-5764
Koenraad Van Meerbeek	koenraad.vanmeerbeek@kuleuven.be	0000-0002-9260-3815
Sybryn L. Maes	sybryn.maes@gmail.com	0000-0002-7168-2390

20   **Author contributions**

21   WG, SM, BM conceived and designed the research; WG, JCSQ carried out the fieldwork; WG,

22   SM, KVM analysed the data; WG drafted the manuscript; TF, BM, JCSQ, KVM, SM critically

23   reviewed the manuscript

Keywords

- Forest restoration
- Nitrogen fixation
- Resource-use strategy
- Seedling growth and survival
- Trait-environment interactions
- Tropical dry forests

Abstract

Tropical dry forests (TDFs) are among the most endangered ecosystems globally, despite their critical socio-ecological importance. In recent decades, tree-planting initiatives have been widely implemented to restore these forests, yielding mixed results. To enhance restoration effectiveness, it is imperative to adopt a predictive approach, focusing on the drivers of seedling performance and their interactions. This study explores how functional traits of planted tree species interact with site environmental conditions to influence seedling survival and growth rates across nine restoration sites of varying ages and climatic conditions within the Peruvian Tumbes-Piura TDFs. While the widely accepted resource-use theory predicts that acquisitive species – focusing on the fast retrieval of resources and characterised by higher specific leaf area and lower wood density – achieve high growth rates in resource-rich environments but experience sharp declines under resource-limited conditions, we found that acquisitive species maintained their high growth rates along the environmental gradient. In contrast, conservative species, expected to be less sensitive to resource variability, exhibited marked declines in growth when water or nutrient availability decreased. Additionally, nitrogen-fixing species did not outperform non-fixers in growth or survival, indicating that nitrogen fixation alone does not confer a consistent advantage in early restoration. Overall, our findings highlight the importance of trait-environment interactions in tree planting performance and underscore the need to align species selection and management strategies with site conditions and broader restoration goals to enhance the success of TDF restoration.

Implications for practice

- Our findings highlight the importance of considering trait-environment interactions when evaluating growth rates across TDF restoration sites.

- Planting acquisitive species, which maintained higher growth rates than conservative species even under resource-poor environmental conditions, can accelerate tropical dry forest recovery in resource-poor restoration sites within the Peruvian Tumbes-Piura region.

- While seedling growth is strongly shaped by environmental factors and their interaction with functional traits, survival may depend more on site-specific factors and management interventions, including irrigation and fertilisation.

- Selecting species based solely on nitrogen fixation ability may not enhance restoration success, as nitrogen-fixing species did not exhibit higher growth or survival than non-fixers.

INTRODUCTION

All around the globe, forests face significant threats from deforestation and degradation, leading to biodiversity loss, accelerating climate change and negatively impacting human well-being (FAO & UNEP 2020). In response to these alarming trends, there has been a surge in ambitious commitments to restore degraded forest ecosystems across all continents (Sewell et al. 2020), further amplified by the current 'UN Decade on Ecosystem Restoration' (2021-2030), which underscores the urgency of reviving damaged ecosystems (United Nations General Assembly 2019).

Many restoration initiatives involve the active planting of trees to speed up ecosystem recovery (Brancalion & Holl 2020; Mesa-Sierra et al. 2024), which is particularly important in tropical dry forests (TDFs) that experience a dry season of several months (Mooney et al. 1995). Active planting can considerably accelerate forest establishment in these areas, as natural recovery is often slow and uncertain due to harsh microclimates and severe forest fragmentation (Alvarez-Aquino & Williams-Linera 2012; Dimson & Gillespie 2020).

Due to numerous limitations, TDFs remain among the most threatened terrestrial ecosystems (Miles et al. 2006), despite their socio-ecological importance in housing numerous endemic species and supporting the livelihoods of some of the world's poorest communities (Blackie et al. 2014; Schröder et al. 2021). The urgency to restore these ecosystems has led to various TDF restoration initiatives (Campo et al. 2023; Dimson & Gillespie 2020; Cárdenas et al. 2022). However, tree planting as TDF restoration approach often yields variable and disappointing growth and survival rates (Dimson & Gillespie 2020; Werden et al. 2023), despite substantial resource investments (Toro et al. 2024). Restoration ecology is therefore urged to become more predictive by studying the factors influencing seedling performance regarding growth and survival (Campo et al. 2023; Brudvig & Catano 2021; Lanuza et al. 2020). Both the species' *functional identity* and the site's *environmental conditions* play a crucial role in governing growth and survival rates of planted trees (Rosell et al. 2022; Werden et al. 2020). Hence, understanding how both factors relate to each other is key to tailoring restoration designs successfully to local environmental contexts and improving overall restoration success (Eviner & Hawkes 2008; Zirbel & Brudvig 2020; Werden et al. 2023).

The functioning of tree species can be summarised by placing them along the *acquisitive-*

conservative functional trait spectrum (Díaz et al. 2004; Wright et al. 2010). By grouping

species with similar resource-use strategies, we can identify and predict their response to

environmental gradients (Pinho et al. 2019) and adapt management practices accordingly

(Eviner & Hawkes 2008). In TDFs, interactions between species' functional traits and

environmental conditions have been explored to understand patterns of functional diversity

along edaphic, climatic and topographic gradients (Méndez-Toribio et al. 2017; Comita &

Engelbrecht 2009; Pinho et al. 2019), suggesting that trait-environment interactions play an

important role in shaping TDF tree growth and survival rates (Sterck et al. 2011).

TDF species have developed distinct resource-use strategies to cope with the seasonally

limited water availability, which is widely regarded as the main limiting factor for seedling

growth and survival in these forests (Alvarez-Aquino & Williams-Linera 2012; Méndez-Toribio

et al. 2020; Markesteijn & Poorter 2009). On the one hand, drought-avoidant acquisitive

species are usually deciduous and characterised by a rapid growth response to changes in

water availability (Rosell et al. 2022). They possess relatively large transpirational tissues (i.e.

high specific leaf area (SLA)) and low wood density (WD), facilitating greater hydraulic

conductivity during the wet season (Poorter et al. 2019; Sterck et al. 2011; Matlaga et al.

2024). In contrast, drought-tolerant conservative species, with higher wood density and lower

SLA, prioritise resource retention and structural reinforcement, resulting in slower growth but

enhanced stress tolerance and survival (Werden et al. 2018).

General resource-use theory predicts that acquisitive species perform better under resource-

rich, favourable environmental conditions, but experience sharp performance declines when

resources become scarce (Méndez-Toribio et al. 2020; Worthy et al. 2020; Mira et al. 2023).

Indeed, their high growth rate drops rapidly when water (Sterck et al. 2011) or soil nutrient

(Baribault et al. 2012; Reich 2014) availability decline. Low soil water availability or high

evaporative demand increases the water potential difference between soil and atmosphere

(Bauman et al. 2022; Méndez-Toribio et al. 2020), forcing acquisitive species to close their

stomata to prevent hydraulic failure, resulting in lower growth rates and potentially even

carbon-starvation-induced mortality after prolonged drought (McDowell et al. 2008).

Conservative species are less sensitive to hydraulic failures and can maintain photosynthetic

activity under drier conditions (Sterck et al. 2011; Matlaga et al. 2024). Regarding soil nutrient

limitations, acquisitive species require higher soil fertility, giving them a growth advantage in

fertile soils but posing challenges when nutrients are scarce (Worthy et al. 2020; Reich 2014).

Moreover, while some studies suggest that acquisitive species are more sensitive to rising

temperatures than conservative species (Sastry & Barua 2017), others contradict this (Slot &

Winter 2017).

An additional important trait in TDFs is nitrogen fixation, which can provide a competitive

advantage through increased photosynthetic activity and water-use efficiency (Pellegrini et al.

2016; Menge & Chazdon 2016). As nitrogen-fixing species span the full acquisitive-

conservative spectrum (Gei 2014), both resource-use strategy and nitrogen fixation should be

considered in tropical forest analyses (Batterman et al. 2018; Werden et al. 2023).

Since seedling performance, i.e. their growth and survival rates, is influenced by both species

traits (acquisitive-conservative spectrum) and local environmental contexts (resources and

conditions), their interactions might play a pivotal role in restoration practices (Caleño-Ruiz et

al. 2023; Zirbel & Brudvig 2020; Mira et al. 2023) and explain some of the observed variability

in restoration outcomes (Dimson & Gillespie 2020). However, the trait-performance

relationship is not always consistent, with some studies showing no clear link between

resource-use strategies and tree performance (Martínez-Garza et al. 2013; Poorter et al.

2018). Additionally, while research on trait-environment has expanded, most studies have

focussed on tropical rainforests (Matlaga et al. 2024; Worthy et al. 2020), neglecting the TDF

biome (Campo et al. 2023; Schröder et al. 2021). This study aims to improve understanding of

how functional trait identity and environmental conditions interactively influence seedling

performance across nine TDF restoration projects. We focus on the Tumbes-Piura dry forest

region in northwestern Peru, contributing to improved tree-planting practices in the area

(Cerrón et al. 2019; Fremout et al. 2022).

Specifically, we evaluate the impacts of interactions between abiotic (i.e. climate and soil)

conditions and functional trait identities (i.e. principal component axis of WD and SLA) of ten

adult tree species on the performance (i.e. growth and survival) of planted seedlings across

nine real-world restoration sites covering a gradient of environmental conditions. We expect

that acquisitive species (with lower WD and higher SLA) will (1) experience lower mortality

and higher growth rates in sites with higher fertility and lower atmospheric aridity, but (2) be

more sensitive than conservative species to lower resource availabilities (Sterck et al. 2011;

Zirbel & Brudvig 2020). Thus, we anticipate conservative species to show little variation in
seedling performance across the studied environmental gradient, whereas acquisitive species
are expected to show higher mortality and lower growth rates in unfavourable conditions but
to exhibit increased performance as environmental conditions improve (Matlaga et al. 2024).

MATERIALS AND METHODS

Study sites and region

To evaluate the drivers of tree planting performance, we sampled nine restoration sites in the
lowland Tumbes-Piura dry forests in northwestern Peru (Figure 1), situated between the
Pacific coast and the Andean mountain range up to altitudes of 500 m.a.s.l. (Linares-Palomino
& Alvarez 2005). The forests host 234 species, with 6 to 25 species per hectare, reaching a
basal area of 2.31 to 22.79 m²/ha (Linares-Palomino & Alvarez 2005; Linares-Palomino et al.
2010). Mature forests in the region typically reach a canopy height of 10 to 15 metres. Average
annual precipitation in the Tumbes-Piura region is low, ranging between 100 and 500 mm, of
which around 85% is centred between January and April, coinciding with the warmest period
of the year (Best & Kessler 1995; Cueva-Ortiz et al. 2020). Mean annual temperatures vary
between 20 and 26°C (Cueva-Ortiz et al. 2020). Every 3 to 7 years, the El Niño Southern
Oscillation (ENSO) brings more intense rains to the region, with precipitation levels up to ten
times higher than the climatic mean, fostering higher (natural) tree recruitment and growth
rates (Best & Kessler 1995).

The Tumbes-Piura dry forests belong to the Tumbes-Chocó-Magdalena biodiversity hotspot
(CEPF 2005) and are renowned for their endemism (Best & Kessler 1995; Linares-Palomino et
al. 2010). Nonetheless, substantial portions of these forests have been lost through
deforestation, and much of the remaining patches are strongly degraded due to overgrazing
by livestock and logging (Fremout et al. 2020; Linares-Palomino & Alvarez 2005). In recent
years, the region has witnessed many restoration efforts to reverse this trend, aiming to
restore forest cover and structure, conserve biodiversity, and raise awareness among the local
population (Cerrón et al. 2019). Between August and September 2022, we sampled nine tree-
planting restoration sites in the area, varying in size (0.22 - 12.7 ha), planting year (1988 –
2022), and species composition (Supplement S1). Three suitable sites were obtained from the

project database of Cerrón et al. (2019) and complemented with six projects suggested by

restoration managers in the region. The sites covered a gradient in climatic conditions (Figure

1), including potential evapotranspiration (1603 - 1968 mm/year), mean annual precipitation

(63 – 188 mm) and mean temperature of the wettest quarter (22.75 – 27.05 °C). Although we

obtained some data on restoration design – planting density (200 – 4444 individuals/hectare)

and propagation method (bare-rooted – containerised) across eight sites – and management

practices – irrigation (40 – 80 L/plant/month) and fertilisation (1.2 – 2.5 kg (organic)

fertiliser/plant/year) at four sites – this information was incomplete and hence omitted from

further analyses (Supplement S1).

Figure 1: (a) Map of the aridity index (mean annual precipitation divided by potential
evapotranspiration (MAP/PET) following Zomer et al. (2022)) showing study site locations and
the number of planted species at each site across northwestern Peru. Distribution of functional
traits – specific leaf area [cm²/g] and wood density [g/cm³] (see further) – along gradients of
(b) potential evapotranspiration [mm/year], (c) mean temperature of the wettest quarter [°C],
and (d) organic matter content [%]. Map generated using QGIS 3.34.0 (QGIS Development
Team 2024), graphs created with R 4.2.3 (R Core Team 2023).

Abiotic conditions

Both climatic and soil conditions were assessed for their influence on planted seedling performance (Supplement S1). Since direct quantification of some resources available to seedlings – such as water and energy availability – is challenging (Parkin et al. 2006), we selected established environmental conditions affecting resource availability. We extracted climate variables for every site from the CHELSA 2.1 Climate Database (Karger et al. 2017) for the period 1981 to 2010 in QGIS 3.34.0 (QGIS Development Team 2024). These variables, with a resolution of 30 arcsec (ca. 0.9 km at the equator), included mean annual temperature (MAT), mean daily air temperatures of the wettest (BIO8) and driest (BIO9) quarter, mean annual precipitation (MAP) and mean annual potential evapotranspiration (PET), computed by summing up monthly averages. Rather than directly quantifying resource availability, these variables describe environmental conditions regulating water and energy balance, with MAT, BIO8 and BIO9 reflecting thermal conditions (Denissen et al. 2022), and MAP and PET capturing aspects of water availability (Konapala et al. 2020; Li et al. 2022; Stefanidis & Alexandridis 2021). For example, PET, indicating atmospheric aridity, has previously explained differing performances of conservative and acquisitive species in TDFs (Méndez-Toribio et al. 2020).

To account for soil variability, eight subsamples were collected per site and combined using the quartering method (EPA 2023) to produce one representative composite sample (Pennock et al. 2006). The subsamples were taken at depths of 10, 20 and 30 cm from eight evenly spaced locations in a zigzag pattern. For two sites exhibiting clear soil heterogeneity, multiple composite samples were acquired from different zones to assess within-site variance, resulting in four composite samples (i.e. 4 × 8 subsamples) at site ENTL and two at site SAPET. The composite soil samples were then air-dried and stored in plastic bags until analysis at the *Universidad Nacional Agraria La Molina* for organic matter content, electrical conductivity, pH and concentrations of K, P and $CaCO_3$ (Supplement S1).

Functional traits

To evaluate the effect of functional traits on seedling performance, we included WD, SLA, maximum height (H), seed mass (SM), deciduousness and nitrogen fixation (Supplement S2).

WD [g/cm³] and SLA [cm²/g] were measured from wood cores and leaf samples for five species

(i.e. *Neltuma pallida, Beautempsia avicenniifolia, Parkinsonia aculeata, Lycium*

boerhaviifolium and *Morisonia scabrida*), while for *Loxopterygium huasango* only SLA was

obtained in the field, as we did not receive permission to extract wood cores for the site in

which the species was present as an adult. The SLA and WD values of the other four species

(i.e. *Cordia lutea, Libidibia glabrata, Tara spinosa* and *Vachellia macracantha*) were obtained

from literature, as we could not find healthy, mature trees to sample from. Similarly, the

maximum tree height, seed mass, deciduousness and nitrogen fixation ability of all studied

species were obtained from literature.

SLA was determined by collecting ten undamaged leaves fully exposed to the sun from two to

four isolated mature individuals near the restoration sites (Cornelissen et al. 2003; Pérez-

Harguindeguy et al. 2013). We included the leaf petioles and did not subdivide compound

leaves into individual leaflets. Fresh leaf area was calculated in the software program ImageJ

(Schneider et al. 2012) following the protocol of O'Neal et al. (2002). After oven-drying the

leaves for 48 hours at 80 °C and measuring their mass, the SLA was calculated by dividing the

leaf area by the oven-dry weight. For the wood density measurements, we sampled mature

individuals in the vicinity of the planting sites with a tree increment borer (Chave 2006). The

extracted cores were stored in paper straws, dried at 80°C for 48 hours (Cornelissen et al.

2003), and the density was computed using X-ray Computed Tomography in the Woodlab at

Ghent University in Belgium (Van den Bulcke et al. 2014). The scanned images were converted

to mean densities of the whole core using a reference material of known density and air with

a density assumed to be zero.

After standardising the continuous traits (SM, H, WD and SLA), we conducted a principal

component analysis (PCA) with the *prcomp* function (R Core Team 2023) and retrieved the

first two axes explaining 65% of the trait variation (Jollife & Cadima 2016). Similar to Díaz et

al. (2015), we found that one axis corresponded to the resource-use strategy of species

($PC_{resource-use}$), with higher values corresponding to species with a lower SLA and higher WD

(Supplement S2). The other axis (PC_{size}) was strongly associated with H and SM, increasing

when species showed higher maximum height and lower seed mass.

Tree growth and survival

We measured the height and recorded survival status of every seedling across the nine study sites (Moonlight et al. 2021). For multi-stemmed trees, commonly found in TDFs, only the tallest stem was recorded. Survival status was evaluated by scratching the bark and observing whether the cambium displayed signs of life (Werden et al. 2020). Relative height growth rates (growth rates [ln(m)/year]) were computed by subtracting the natural logarithm of the obtained height by the natural logarithm of 0.30 m, reported as the mean height at planting by five restoration managers, and divided by the time in years between planting and monitoring (Werden et al. 2023; Worthy et al. 2020):

$$RGR = \frac{\ln() - \ln(0.30)}{2022 - Planting\ year}$$

During the survival assessment, we could only evaluate the status of individuals present at the time of monitoring for the seven mixed-species sites, which meant that empty planting spots had to be omitted, leading to an overestimation of survival rates and potentially introducing bias by favouring some species more than others. However, in the two oldest sites, which are monocultures of *Neltuma pallida*, species identity was not a concern, allowing accurate quantification of mortality rates.

Data analysis

Overall, ten species were present in at least two sites and were included in our analysis. For the two sites established in the monitoring year, we only evaluated survivorship, as measured heights likely represent planting height, and not (yet) significant growth rates due to the short growing time period (Rodríguez-Alarcón et al. 2024). We removed all 21 RGR observations considered outliers at the site level following Tukey (1977). Our final dataset consisted of 5514 tree individuals for the survival assessment, which resulted in 3057 growth measurements after removing dead individuals, growth outliers and the two sites established in the monitoring year (Supplement S3). All explanatory factors, except for the categorical nitrogen fixation and two PCA-axes, were standardised to facilitate model fitting and effect size comparison (Zuur et al. 2009).

Generalised linear mixed models (GLMMs) were employed to evaluate the drivers of growth

and survival variability in TDF restoration sites, with site identity as a random intercept to

account for the ecological assumption that within-site observations are more closely related

(Banin et al. 2023), while species identity was also included as a random factor. In sites with

multiple composite soil samples, tree individuals were related to their soil sampling zone. The

model included the first two PCA-axes, depicting the variation in resource-use ($PC_{resource-use}$)

and size (PC_{size}) among the species (Díaz et al. 2015; Liu et al. 2022; Werden et al. 2023), and

ability to fix nitrogen (Nfix). We expanded this three-factor model with the environmental

factors potential evapotranspiration (PET), mean temperature in the wettest quarter (BIO8)

and presence of organic matter (OM). Given the strong correlations between the

environmental factors, only a limited set was used as explanatory variables. We included PET

as a proxy of water availability, considering that it indicates atmospheric aridity and has been

linked previously to differing growth responses of conservative and acquisitive species

(Méndez-Toribio et al. 2020). Similarly, interactive effects are expected between resource-use

strategy and BIO8 – an indicator for energy availability in the rainy season (Slot & Winter 2017;

Denissen et al. 2022) – and OM – representing the soil fertility and water-holding capacity

(Wood et al. 2016). Interaction effects with $PC_{resource-use}$ were included for all abiotic predictors

except BIO8, due to induced multicollinearity and increased variance inflation factors (VIFs)

(Kim 2019). Moreover, we included site age as a fixed variable in the growth model, accounting

for time-related changes in microclimate and growth patterns (Werden et al. 2020; Eviner &

Hawkes 2008). While age is also relevant to model survivorship (Liu et al. 2022), its strong

correlation with the site-level random effect caused model singularity. Since age's effect was

indirectly captured by the site random effect in the survival model, the following two models

were ultimately tested:

RGR ~ Nfix + PC_{size} + $PC_{resource-use}$*PET + $PC_{resource-use}$*OM + BIO8 + AGE + (1|Site) + (1|Species)

Survival ~ Nfix + PC_{size} + $PC_{resource-use}$*PET + $PC_{resource-use}$*OM + BIO8 + (1|Site) + (1|Species)

Height growth rates were analysed using the *lme4* package (Bates et al. 2015) with Gaussian

error distribution and Restricted Maximum Likelihood (REML) estimation (Banin et al. 2023;

Zuur et al. 2009). Significance of the results was assessed via Satterwaite's degrees of freedom

method in the *lmerTest* package (Kuznetsova et al. 2017). Survival was modelled with a

binomial error distribution and *cloglog* link function (Banin et al. 2023; Liu et al. 2022) using

the *glmmTMB* package (Brooks et al. 2017). We used the *piecewiseSEM* package (Lefcheck

2016) to compute the marginal (R^2_m) and conditional (R^2_c) coefficients of determination to

assess the goodness-of-fit (Nakagawa & Schielzeth 2013). The former quantifies the variance

explained by the fixed effects, while the latter evaluates the variance explained by both the

fixed and random factors. Model assumptions were checked using the *DHARMa* package

(Hartig 2024). Two additional models were tested, following the same structure but including

the interaction between resource-use strategy and BIO8 instead of PET. We also ran the

survival model including age as an additional explanatory variable and omitting the random

effect of site (Supplement S4), which confirmed the importance of age in explaining survival

rates. All analyses were performed in R 4.2.3 (R Core Team 2023), using the *tidyverse*

(Wickham et al. 2019), *ggeffects* (Lüdecke 2025) and *ggpubr* (Kassambara 2023) packages.

RESULTS

Tree seedling growth varied considerably across species and sites (Supplement S3) and was

significantly influenced by trait-environment interactions (Table 1; Figure 2a-b). First, the

interaction between $PC_{resource-use}$ and PET interactively influenced height growth. Specifically,

the growth of conservative species – with a higher WD and lower SLA – was more negatively

affected by an increase in PET than that of acquisitive species (PET × $PC_{resource-use}$ -0.09

$p < 0.001$). Likewise, a significant interaction existed between $PC_{resource-use}$ and OM, as the

height growth of acquisitive species remained relatively stable and even slightly increased

with lower values of OM, while that of conservative species decreased when OM decreased

(OM × $PC_{resource-use}$ 0.10 $p < 0.001$). Significant main effects were found for BIO8 (0.48 $p < 0.05$)

and age (0.04 $p < 0.05$), while nitrogen fixation and PC_{size} had no significant impact. Overall, the

fixed effects accounted for 75.3% of the variance in height growth (R^2_m), increasing to 82.3%

when site and species were included as random factors (R^2_c).

In contrast, while seedling survivorship also varied considerably across species and sites

(Supplement S3), the only significant effect in the survival model was a positive interaction

between OM and $PC_{resource-use}$ (OM × $PC_{resource-use}$ 0.06 $p < 0.05$). Specifically, survival rates of

conservative species increased with higher organic matter content, but the opposite trend

was observed for acquisitive species (Table 1; Figure 2c). However, the fixed effects could only

explain 4.6% of the variation (R^2_m) in the survival model, increasing to 24.6% when accounting

for the random intercepts of species and site (R^2_c).

When testing an alternative model replacing the PET interaction with an interaction between

BIO8 and $PC_{resource-use}$, results remained largely consistent. BIO8 and $PC_{resource-use}$ significantly

interacted to influence height growth (Figure 2d), with acquisitive species showing stronger

growth responses to higher temperature in the wettest quarter than conservative species

(BIO8 × $PC_{resource-use}$ 0.16 $p < 0.001$). The OM × $PC_{resource-use}$ interaction remained significant

(OM × $PC_{resource-use}$ 0.09 $p < 0.001$), showing the same pattern as in the PET interaction model.

Similarly, the survival model including the interaction with BIO8 instead of PET yielded

comparable results, with the OM × $PC_{resource-use}$ interaction remaining the only significant effect

(OM × $PC_{resource-use}$ 0.07 $p < 0.05$). Given these consistencies, we present only the BIO8

interaction for growth in Figure 2, with additional statistical details for the growth and survival

models in Supplement S5.

Table 1: Parameter estimates (β) and 95%-confidence intervals (CI) for the generalised linear
mixed models (GLMMs) examining height growth and survival rates focusing on the interaction
between resource-use strategy and water availability. Predictor variables include nitrogen
fixation (Nfix), resource-use strategy ($PC_{resource-use}$), size variation (PC_{size}), potential
evapotranspiration (PET [mm/year]), organic matter content (OM [%]), mean temperature in
the wettest quarter (BIO8 [°C]) and site age. Low values of $PC_{resource-use}$ indicate acquisitive
species, while higher values correspond to more conservative species. Significance of the
parameters is indicated with * p<0.05, ** p<0.01 and *** p<0.001. Goodness-of-fit is given as
both marginal (R^2_m) and conditional (R^2_c) coefficients of determination. Supplement S6 provides
additional details on the degrees of freedom and t-values for the growth model, and z-values
for the survival model.

Driver	Height growth rate		Survival rate	
	β	CI	β	CI
Nfix	-0.17	[-0.36, 0.02]	0.48	[0.06, 0.90]
$PC_{resource-use}$	-0.29**	[-0.35, -0.23]	-0.11	[-0.25, 0.04]
PC_{size}	0.16	[0.07, 0.24]	-0.11	[-0.30, 0.09]
PET	-0.31*	[-0.40, -0.23]	0.05	[-0.11, 0.21]
OM	0.10*	[0.06, 0.14]	-0.04	[-0.08, -0.01]
BIO8	0.48*	[0.38, 0.57]	-0.25	[-0.41, -0.09]
AGE	0.04*	[0.04, 0.05]	NA	NA
$PC_{resource-use}$*PET	-0.09***	[-0.11, -0.08]	-0.03	[-0.06, 0.00]
$PC_{resource-use}$*OM	0.10***	[0.07, 0.12]	0.06*	[0.03, 0.09]
R^2_m (%)	75.3		4.6	
R^2_c (%)	82.3		24.6	

Figure 2: Significant interaction effects from the first model between resource-use strategy (acquisitive versus conservative based on the first principal component axis) and (a) annual potential evapotranspiration (PET [mm/year]) and (b) organic matter content (OM [%]) on height growth. Effect (c) of the interaction between resource-use strategy and organic matter content on survival rates. Results from the additional model, (d) showing the significant interaction effect between the resource-use strategy and mean temperature during the wettest quarter (BIO8 [°C]) on height growth. Lines represent predicted values for three levels of the first principal component axis, corresponding to the most acquisitive, an intermediate and the most conservative species. Points represent individuals, coloured according to their resource-use strategy.

DISCUSSION

Our results demonstrate the importance of incorporating interactions between functional traits and environmental variables when assessing tree planting performance in restoration projects. While most studies assess the individual effects of these factors on seedling growth and survival (Werden et al. 2018; Rosell et al. 2022), the presence of significant interactions suggests that considering them in isolation may not fully capture the complexity of TDF restoration projects.

Acquisitive species are less sensitive to higher evapotranspiration and lower organic matter content

Acquisitive species – characterised by lower WD and higher SLA – maintained relative high growth rates across the environmental gradients in potential evapotranspiration (as a proxy for atmospheric aridity and water availability; Méndez-Toribio et al. 2020) and organic matter content (a proxy for nutrient content; Wood et al. 2016). This suggests that acquisitive species were able to retrieve sufficient resources and allocate them to height growth across a range of resource availability conditions, i.e. even under resource-poorer conditions. This finding opposes the general resource-use theory, which predicts higher growth rates for acquisitive species under favourable conditions but a sharp decline in performance when water and nutrients become limiting, while, conversely, conservative species are expected to be less sensitive to such resource variation (Sterck et al. 2011; Baribault et al. 2012; Caleño-Ruiz et al. 2023).

The growth rates of acquisitive species appeared less sensitive to the environmental variation (i.e. both water and organic matter availability), which indicates that once established, acquisitive seedlings can sustain high growth rates regardless of site resource availability, suggesting a competitive growth advantage over slow-growing conservative species (Werden et al. 2023). Mechanistically, a possible explanation is that fast-growing acquisitive species invest heavily in belowground biomass and thereby increase their rooting density and depth (Fagundes et al. 2022; Méndez-Toribio et al. 2020; Martínez-Garza et al. 2013), which enhances their growth capacity (Werden et al. 2023) and reduces susceptibility to changes in soil fertility and atmospheric aridity (Markesteijn & Poorter 2009). Once well-established,

acquisitive species may thus retrieve nutrients and water more efficiently than conservative

species, allowing them to sustain height growth (Carrasco-Carballido et al. 2019). However,

due to the lack of root trait data, we cannot confirm this hypothesis, although previous studies

support a link between acquisitive resource use and rooting density (Comita & Engelbrecht

2009; Méndez-Toribio et al. 2020).

Another possible explanation for the relatively sustained growth rates of acquisitive species

along the resource gradient is that they prioritise high growth rates during the short wet

season in TDFs (Mendivelso et al. 2014; Poorter et al. 2018), after which many quickly shed

their leaves to avoid drought stress (Fagundes et al. 2022; Poorter & Markesteijn 2008;

Rodríguez-Alarcón et al. 2024). Conservative species, in contrast, remain active during (part

of) the dry season, extending their growing period but also increasing their susceptibility to

the higher evaporative demand (McDowell et al. 2008; Wright et al. 2010). As a result,

conservative species are forced to adjust their growth rates to match available resources,

making them more sensitive to increased potential evapotranspiration.

Our survival analysis further revealed contrasting effects of organic matter availability, with

higher nutrient content increasing survivorship in conservative species but reducing it in

acquisitive species. Although previous studies reported higher survival rates for fast-growing

acquisitive species in early TDF restoration stages (Werden et al. 2023; Martínez-Garza et al.

2013; González-Tokman et al. 2018), the observed decline at greater nutrient availability is

unexpected. However, given the low explanatory power of our survival model, these results

should be interpreted with caution.

Acquisitive species benefit more from higher wet season temperatures

Higher temperatures during the wettest quarter positively affected growth rates, challenging

our initial hypothesis and previous studies (Méndez-Toribio et al. 2017; Mendivelso et al.

2014). One possible explanation is that increased temperatures during the growing season

enhance energy availability, thereby stimulating photosynthetic activity (Slot & Winter 2017).

We also anticipated that acquisitive species would be more sensitive than conservative

species to temperature increases, expecting a more rapid growth decline under warmer

conditions (Sastry & Barua 2017). Although we could not include this interaction in the main

model due to high VIFs, testing it separately revealed that acquisitive species exhibited a
stronger positive response to temperature than conservative species. This supports the
hypothesis by Slot and Winter (2017) that acquisitive species benefit more from increased
energy availability, likely due to their higher electron transport capacity and greater plasticity
in temperature responses. Additionally, these results align with our hypothesis that acquisitive
species are less sensitive to higher atmospheric aridity, as they concentrate growth during the
wet period when water and energy availability are optimal.

Nitrogen fixation and size do not significantly affect seedling performance

Contrary to our hypothesis, nitrogen-fixing species did not exhibit enhanced growth and
survival rates, despite potentially higher photosynthetic activity and water-use efficiency
(Pellegrini et al. 2016; Menge & Chazdon 2016). As in previous TDF studies (Werden et al.
2023; Cárdenas et al. 2022), nitrogen-fixing and non-fixing species showed similar
performance, suggesting that nitrogen fixation alone does not necessarily provide an
advantage in early restoration sites. This reinforces the importance of including mixtures of
fixers and non-fixers in restoration projects (Werden et al. 2023), especially considering that
nitrogen fixing species span the acquisitive-conservative resource-use gradient (Gei 2014).

Similarly, we found no significant effect of the principal component axis representing variation
in species size. This axis was positively associated with maximum height but negatively with
seed mass, whereas Díaz et al. (2015) found that both traits contributed positively to size
variation. This suggests that trait relationships among our study species may differ from those
observed in other systems, possibly explaining why the size axis was not a strong predictor of
seedling performance, in contrast to previous studies on tropical forests that found important,
although contrasting, effects of seed mass and maximum height on seedling growth and
survival (Wright et al. 2010; Martínez-Garza et al. 2013).

Study limitations and future research avenues

Overall, our results indicate that across nine restoration projects in the Tumbes-Piura TDFs,
acquisitive species maintain relative high growth rates even when water or nutrient content
becomes limiting, and respond positively to higher temperatures during the wet season.
Additionally, the relatively small difference between the marginal and conditional R^2 values in

the growth model suggests that the fixed main and interactive effects effectively captured

growth variation. These findings highlight the importance of trait-environment interactions in

seedling performance, providing valuable empirical evidence to inform tree planting

strategies in TDF restoration.

However, we acknowledge the limitations caused by our study's observational nature, which

restricts our ability to directly assess key management strategies, initial planting conditions

and seedling provenance (Dimson & Gillespie 2020; González-Tokman et al. 2018), and

prevented control over factors like the planted number of individuals per species. Additionally,

while temperature, potential evapotranspiration and organic matter content are useful

proxies for resource availability (Stefanidis & Alexandridis 2021; Konapala et al. 2020; Li et al.

2022), they do not fully capture the complexity of water and energy dynamics (Parkin et al.

2006; Bauman et al. 2022) or soil fertility (Pennock et al. 2006). The scarcity of initial planting

data also meant that variations in initial seedling sizes might have influenced the relative

growth rate calculations, although much of this variation is likely captured by the random

effects of species and site. Similarly, we could only include remaining dead individuals with

certain species identification and had to omit empty planting spots in all but two sites, which

may have resulted in an overrepresentation of conservative species with more durable stems

in the survival models (Chave et al. 2009), potentially biasing survival estimates and

contributing to the models' low explanatory power. Notably, the relatively high variance in

survivorship explained by random effects indicates that unmeasured species- and site-specific

factors, including local management interventions (Mesa-Sierra et al. 2024), may have

influenced survival rates.

Nonetheless, our study provides important insights into resource-use dynamics within real-

world tree planting efforts in TDFs, complementing findings from controlled but relatively

small-scale experiments (Werden et al. 2020, 2023) and broader observational studies of

natural forest succession (Poorter et al. 2019; Derroire et al. 2018). While growth was largely

shaped by trait-environment interactions, our findings suggest that survival may depend more

strongly on factors such as management practices (Dimson & Gillespie 2020; Mesa-Sierra et

al. 2024). In contrast, Toro et al. (2024) found limited effects of irrigation and fertilisation on

the growth and survival of most TDF species. However, their findings may not be directly

comparable to ours, as their study was conducted in a region with substantially higher mean

annual precipitation and shorter dry season.

Future research on trait-environment interactions should prioritise controlled experimental
designs with known initial species numbers, management practices and environmental
conditions along a gradient in both water and nutrient availability. We also encourage future
restoration research to incorporate trait plasticity (Derroire et al. 2018; Poorter et al. 2018)
and ontogenetic variability (Metz et al. 2023; Poorter & Markesteijn 2008), since both factors
allow individuals to acclimate to local abiotic conditions and could govern trait-environment
interactions in restoration projects (Lanuza et al. 2020; Worthy et al. 2020). Additionally,
accounting for the high soil and microclimatic heterogeneity of TDFs in future studies can shed
additional light on the importance of trait-environment interactions in restoration practices
(Rodríguez-Alarcón et al. 2024; Caleño-Ruiz et al. 2023).

Given our observation that seedling growth depends on the interaction between
environmental variables and species' resource-use strategy, our findings underscore the
importance of context-dependent management approaches in restoration projects and
highlight that species selection should consider both the site's environmental conditions and
overarching restoration objectives (Zirbel & Brudvig 2020; Fremout et al. 2022), while also
integrating management interventions to enhance seedling survival (Brancalion & Holl 2020).
This study lays the foundation for future research on these interactions, particularly within the
underexplored Tumbes-Piura TDFs.

Acknowledgements

The authors express their gratitude to Michael Perring (UKCEH) for his valuable critique of the
manuscript and to Nora Grados and Gaston Cruz (UDEP) for their assistance with legal
documentation and the analysis of leaf and wood samples. The authors also acknowledge the
cooperation of restoration managers who facilitated the research on their sites, as well as the
Peruvian Forest Service (SERFOR) for granting research and botanical sampling permissions.
The fieldwork was made possible through a VLIR-UOS scholarship (REI-2022-01-01).

Literature cited

Alvarez-Aquino C, Williams-Linera G (2012) Seedling Survival and Growth of Tree Species: Site Condition and Seasonality in Tropical Dry Forest Restoration. Botanical Sciences 90:341–351

Banin LF, Raine EH, Rowland LM, Chazdon RL, Smith SW, Rahman NEB, et al. (2023) The road to recovery: a synthesis of outcomes from ecosystem restoration in tropical and sub-tropical Asian forests. Philosophical Transactions of the Royal Society B 378

Baribault TW, Kobe RK, Finley AO (2012) Tropical tree growth is correlated with soil phosphorus, potassium, and calcium, though not for legumes. Ecological Monographs 82:189–203

Bates D, Mächler M, Bolker BM, Walker SC (2015) Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software 67:1–48

Batterman SA, Hall JS, Turner BL, Hedin LO, LaHaela Walter JK, Sheldon P, et al. (2018) Phosphatase activity and nitrogen fixation reflect species differences, not nutrient trading or nutrient balance, across tropical rainforest trees. Ecology Letters 21:1486–1495

Bauman D, Fortunel C, Cernusak LA, Bentley LP, McMahon SM, Rifai SW, et al. (2022) Tropical tree growth sensitivity to climate is driven by species intrinsic growth rate and leaf traits. Global Change Biology 28:1414–1432

Best BJ, Kessler M (1995) Biodiversity and Conservation in Tumbesian Ecuador and Peru.

Blackie R, Baldauf C, Gautier D, Gumbo D, Kassa H, Parthasarathy N, et al. (2014) Tropical dry forests: The state of global knowledge and recommendations for future research.

Brancalion PHS, Holl KD (2020) Guidance for successful tree planting initiatives. Journal of Applied Ecology 57:2349–2361

Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et al. (2017) glmmTMB balances speed and flexibility among packages for zero-inflated generalized

linear mixed modeling. R Journal 9:378–400

Brudvig LA, Catano CP (2021) Prediction and uncertainty in restoration science. Restoration
Ecology e13380

Van den Bulcke J, Wernersson ELG, Dierick M, Van Loo D, Masschaele B, Brabant L, et al. (2014)
3D tree-ring analysis using helical X-ray tomography. Dendrochronologia 32:39–46

Caleño-Ruiz BL, Garzón F, López-Camacho R, Pizano C, Salinas V, González-M R (2023) Soil
resources and functional trait trade-offs determine species biomass stocks and
productivity in a tropical dry forest. Frontiers in Forests and Global Change 6:1028359

Campo J, Giardina CP, Dirzo R (2023) Tropical Dry Forest Restoration in an Era of Global
Change: Ecological and Social Dimensions. Sustainability 2023, Vol. 15, Page 3052
15:3052

Cárdenas CD, Varón-García D, Suárez-Rodríguez F, Pizano C (2022) Overcoming the
regeneration barriers of tropical dry forest: effects of water stress and herbivory on
seedling performance and allocation of key tree species for restoration. Journal of
Tropical Ecology 38:210–218

CEPF (2005) Protecting Nature's Hotspots for People and Prosperity. Annual report 2005.

Cerrón J, Fremout T, Thomas E, Cornelius JP, Atkinson R (2019) Experiencias de restauración y
fuentes semilleras en el bosque seco tropical del norte del Perú: Estado actual y lecciones
aprendidas.

Chave J (2006) Measuring wood density for tropical forest trees. A field manual.

Chave J, Coomes D, Jansen S, Lewis SL, Swenson NG, Zanne AE (2009) Towards a worldwide
wood economics spectrum. Ecology Letters 12:351–366

Comita LS, Engelbrecht BMJ (2009) Seasonal and spatial variation in water availability drive
habitat associations in a tropical forest. Ecology 90:2755–2765

Cornelissen JHC, Lavorel B S, Garnier B E, Díaz C S, Buchmann N, Gurvich DE, et al. (2003) A
handbook of protocols for standardised and easy measurement of plant functional traits

worldwide. Australian Journal of Botany 51:335–380

Cueva-Ortiz J, Espinosa CI, Aguirre-Mendoza Z, Gusmán-Montalván E, Weber M, Hildebrandt
P (2020) Natural Regeneration in the Tumbesian Dry Forest: Identification of the Drivers
Affecting Abundance and Diversity. Scientific Reports 2020 10:1 10:1–13

Denissen JMC, Teuling AJ, Pitman AJ, Koirala S, Migliavacca M, Li W, et al. (2022) Widespread
shift from ecosystem energy to water limitation with climate change. Nature Climate
Change 2022 12:7 12:677–684

Derroire G, Powers JS, Hulshof CM, Cárdenas Varela LE, Healey JR (2018) Contrasting patterns
of leaf trait variation among and within species during tropical dry forest succession in
Costa Rica. Scientific reports 8

Díaz S, Hodgson JG, Thompson K, Cabido M, Cornelissen JHC, Jalili A, et al. (2004) The plant
traits that drive ecosystems: Evidence from three continents. Journal of Vegetation
Science 15:295–304

Díaz S, Kattge J, Cornelissen JHC, Wright IJ, Lavorel S, Dray S, et al. (2015) The global spectrum
of plant form and function. Nature 2015 529:7585 529:167–171

Dimson M, Gillespie TW (2020) Trends in active restoration of tropical dry forest: Methods,
metrics, and outcomes. Forest Ecology and Management 467:118150

EPA (2023) Soil Sampling.

Eviner VT, Hawkes C V. (2008) Embracing Variability in the Application of Plant–Soil
Interactions to the Restoration of Communities and Ecosystems. Restoration Ecology
16:713–729

Fagundes MV, Souza AF, Oliveira RS, Ganade G (2022) Functional traits above and below
ground allow species with distinct ecological strategies to coexist in the largest seasonally
dry tropical forest in the Americas. Frontiers in Forests and Global Change 5:930099

FAO and UNEP (2020) The State of the World's Forests 2020. FAO and UNEP

Fremout T, Thomas E, Gaisberger H, Van Meerbeek K, Muenchow J, Briers S, et al. (2020)

Mapping tree species vulnerability to multiple threats as a guide to restoration and
conservation of tropical dry forests. Global Change Biology 26:3552–3568

Fremout T, Thomas E, Taedoumg H, Briers S, Gutiérrez-Miranda CE, Alcázar-Caicedo C, et al.
(2022) Diversity for Restoration (D4R): Guiding the selection of tree species and seed
sources for climate-resilient restoration of tropical forest landscapes. Journal of Applied
Ecology 59:664–679

Gei MG (2014) Biological nitrogen fixation in tropical dry forests of costa rica: patterns and
controls. University of Minnesota

González-Tokman DM, Barradas VL, Boege K, Domínguez CA, del-Val E, Saucedo E, et al. (2018)
Performance of 11 tree species under different management treatments in restoration
plantings in a tropical dry forest. Restoration Ecology 26:642–649

Hartig F (2024) Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models
[R package DHARMa version 0.4.7]. CRAN: Contributed Packages

Jollife IT, Cadima J (2016) Principal component analysis: a review and recent developments.
Philosophical transactions. Series A, Mathematical, physical, and engineering sciences
374:20150202

Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, et al. (2017) Climatologies
at high resolution for the earth's land surface areas. Scientific Data 2017 4:1 4:1–20

Kassambara A (2023) 'ggplot2' Based Publication Ready Plots [R package ggpubr version 0.6.0].
CRAN: Contributed Packages

Kim JH (2019) Multicollinearity and misleading statistical results. Korean Journal of
Anesthesiology 72:558

Konapala G, Mishra AK, Wada Y, Mann ME (2020) Climate change will affect global water
availability through compounding changes in seasonal precipitation and evaporation.
Nature Communications 2020 11:1 11:1–10

Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest Package: Tests in Linear Mixed

Effects Models. Journal of Statistical Software 82:1–26

Lanuza OR, Espelta JM, Peñuelas J, Peguero G (2020) Assessing intraspecific trait variability
during seedling establishment to improve restoration of tropical dry forests. Ecosphere
11:e03052

Lefcheck JS (2016) piecewiseSEM: Piecewise structural equation modelling in r for ecology,
evolution, and systematics. Methods in Ecology and Evolution 7:573–579

Li J, Pandey B, Dakhil MA, Khanal M, Pan K (2022) Precipitation and potential
evapotranspiration determine the distribution patterns of threatened plant species in
Sichuan Province, China. Scientific Reports 2022 12:1 12:1–11

Linares-Palomino R, Alvarez SIP (2005) Tree community patterns in seasonally dry tropical
forests in the Cerros de Amotape Cordillera, Tumbes, Peru. Forest Ecology and
Management 209:261–272

Linares-Palomino R, Kvist LP, Aguirre-Mendoza Z, Gonzales-Inca C (2010) Diversity and
endemism of woody plant species in the Equatorial Pacific seasonally dry forests.
Biodiversity and Conservation 19:169–185

Liu X, Huang Y, Chen L, Li S, Bongers FJ, Castro-Izaguirre N, et al. (2022) Species richness,
functional traits and climate interactively affect tree survival in a large forest biodiversity
experiment. Journal of Ecology 110:2522–2531

Lüdecke D (2025) Create Tidy Data Frames of Marginal Effects for 'ggplot' from Model Outputs
[R package ggeffects version 2.2.1]. CRAN: Contributed Packages

Markesteijn L, Poorter L (2009) Seedling root morphology and biomass allocation of 62 tropical
tree species in relation to drought- and shade-tolerance. Journal of Ecology 97:311–325

Martínez-Garza C, Bongers F, Poorter L (2013) Are functional traits good predictors of species
performance in restoration plantings in tropical abandoned pastures? Forest Ecology and
Management 303:35–45

Matlaga D, Lammerant R, Hogan JA, Uriarte M, Rodriguez-Valle C, Zimmerman JK, et al. (2024)

Survival, growth, and functional traits of tropical wet forest tree seedlings across an
experimental soil moisture gradient in Puerto Rico. Ecology and Evolution 14

McDowell N, Pockman WT, Allen CD, Breshears DD, Cobb N, Kolb T, et al. (2008) Mechanisms
of plant survival and mortality during drought: why do some plants survive while others
succumb to drought? New Phytologist 178:719–739

Méndez-Toribio M, Ibarra-Manríquez G, Navarrete-Segueda A, Paz H (2017) Topographic
position, but not slope aspect, drives the dominance of functional strategies of tropical
dry forest trees. Environmental Research Letters 12:085002

Méndez-Toribio M, Ibarra-Manríquez G, Paz H, Lebrija-Trejos E (2020) Atmospheric and soil
drought risks combined shape community assembly of trees in a tropical dry forest.
Journal of Ecology 108:1347–1357

Mendivelso HA, Camarero JJ, Gutiérrez E, Zuidema PA (2014) Time-dependent effects of
climate and drought on tree growth in a Neotropical dry forest: Short-term tolerance vs.
long-term sensitivity. Agricultural and Forest Meteorology 188:13–23

Menge DNL, Chazdon RL (2016) Higher survival drives the success of nitrogen-fixing trees
through succession in Costa Rican rainforests. New Phytologist 209:965–977

Mesa-Sierra N, de la Peña-Domene M, Campo J, Giardina CP (2024) Restoration of tropical dry
forest: an analysis of constraints and successes across a highly threatened biome.
Frontiers in Environmental Science 12:1458613

Metz MR, Wright SJ, Zimmerman JK, Hernandéz A, Smith SM, Swenson NG, et al. (2023)
Functional traits of young seedlings predict trade-offs in seedling performance in three
neotropical forests. Journal of Ecology 111:2568–2582

Miles L, Newton AC, DeFries RS, Ravilious C, May I, Blyth S, et al. (2006) A global overview of
the conservation status of tropical dry forests. Journal of Biogeography 33:491–505

Mira E, Cochard H, Evette A, Dulormne M (2023) Growth, Xylem Vulnerability to Cavitation
and Leaf Cell Response to Dehydration in Tree Seedlings of the Caribbean Dry Forest.
Forests 14:697

Mooney HA, Bullock SH, Medina E (1995) Introduction. In: Seasonally Dry Tropical Forests.
Bullock, SH, Mooney, HA, & Medina, E, editors. Cambridge University Press, Cambridge
pp. 1–8.

Moonlight P, Banda-R K, Phillips OL, Dexter KG, Pennington RT, Baker TR, et al. (2021)
Supplementarty Material: The DryFlor field manual for plot establishment and
remeasurement. Plants, People, Planet 3:295–300

Nakagawa S, Schielzeth H (2013) A general and simple method for obtaining R2 from
generalized linear mixed-effects models. Methods in Ecology and Evolution 4:133–142

O'Neal ME, Landis DA, Isaacs R (2002) An Inexpensive, Accurate Method for Measuring Leaf
Area and Defoliation Through Digital Image Analysis. Journal of Economic Entomology
95:1190–1194

Parkin G, Clarke Topp G, Ferre´ T (2006) Soil Water Content. Soil Sampling and Methods of
Analysis, Second Edition

Pellegrini AFA, Staver AC, Hedin LO, Charles-Dominique T, Tourgee A (2016) Aridity, not fire,
favors nitrogen-fixing plants across tropical savanna and forest biomes. Ecology 97:2177–
2183

Pennock D, Yates T, Braidek J (2006) Soil sampling designs. In: Soil Sampling and Methods of
Analysis: Second Edition. CRC Press pp. 1–14.

Pérez-Harguindeguy NA, Díaz SA, Garnier EB, Lavorel SC, Poorter HD, Jaureguiberry PA, et al.
(2013) New handbook for standardised measurement of plant functional traits
worldwide. Australian Journal of Botany 61:167–234

Pinho BX, Tabarelli M, Engelbrecht BMJ, Sfair J, Melo FPL (2019) Plant functional assembly is
mediated by rainfall and soil conditions in a seasonally dry tropical forest. Basic and
Applied Ecology 40:1–11

Poorter L, Castilho C V., Schietti J, Oliveira RS, Costa FRC (2018) Can traits predict individual
growth performance? A test in a hyperdiverse tropical forest. New Phytologist 219:109–
121

Poorter L, Markesteijn L (2008) Seedling Traits Determine Drought Tolerance of Tropical Tree
Species. Biotropica 40:321–331

Poorter L, Rozendaal DMA, Bongers F, de Almeida-Cortez JS, Almeyda Zambrano AM, Álvarez
FS, et al. (2019) Wet and dry tropical forests show opposite successional pathways in
wood density but converge over time. Nature Ecology & Evolution 2019 3:6 3:928–934

QGIS Development Team (2024) QGIS Geographic Information System.

R Core Team (2023) R: A Language and Environment for Statistical Computing.

Reich PB (2014) The world-wide 'fast–slow' plant economics spectrum: a traits manifesto.
Journal of Ecology 102:275–301

Rodríguez-Alarcón S, González-M R, Carmona CP, Tordoni E (2024) Trait–growth relationships
in Colombian tropical dry forests: Incorporating intraspecific variation and trait
interactions. Journal of Vegetation Science 35:e13233

Rosell JA, Olson ME, Martínez-Garza C, Martínez-Méndez N (2022) Functional Diversity in
Woody Organs of Tropical Dry Forests and Implications for Restoration. Sustainability
(Switzerland) 14:8362

Sastry A, Barua D (2017) Leaf thermotolerance in tropical trees from a seasonally dry climate
varies along the slow-fast resource acquisition spectrum. Scientific Reports 2017 7:1 7:1–
11

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis.
Nature Methods 2012 9:7 9:671–675

Schröder JM, Ávila Rodríguez LP, Günter S (2021) Research trends: Tropical dry forests: The
neglected research agenda? Forest Policy and Economics 122:102333

Sewell A, Van Der Esch S, Löwenhardt H (2020) Goals and Commitments for the Restoration
Decade: A Global Overview of Countries' Restoration Commitments under the Rio
Conventions and Other Pledges. The Hague

Slot M, Winter K (2017) In situ temperature response of photosynthesis of 42 tree and liana

species in the canopy of two Panamanian lowland tropical forests with contrasting
rainfall regimes. New Phytologist 214:1103–1117

Stefanidis S, Alexandridis V (2021) Precipitation and Potential Evapotranspiration Temporal
Variability and Their Relationship in Two Forest Ecosystems in Greece. Hydrology 2021,
Vol. 8, Page 160 8:160

Sterck F, Markesteijn L, Schieving F, Poorter L (2011) Functional traits determine trade-offs
and niches in a tropical forest community. Proceedings of the National Academy of
Sciences of the United States of America 108:20627–20632

Toro L, Torres-Romero F, Salinas SM, Avella-Munoz A, Galatowish S, Secchi S, et al. (2024)
Cost-effectiveness of management strategies in a nucleation experiment in a tropical dry
forest. Restoration Ecology 32:e14094

Tukey J (1977) Exploratory data analysis. Addison-Wesley, Reading, MA

United Nations General Assembly (2019) General Assembly official records, 73rd session: 69th
plenary meeting.

Werden LK, Averill C, Crowther TW, Calderón-Morales E, Toro L, Alvarado JP, et al. (2023)
Below-ground traits mediate tree survival in a tropical dry forest restoration.
Philosophical Transactions of the Royal Society B: Biological Sciences 378

Werden LK, Becknell JM, Powers JS (2018) Edaphic factors, successional status and functional
traits drive habitat associations of trees in naturally regenerating tropical dry forests.
Functional Ecology 32:2766–2776

Werden LK, Calderón-Morales E, Alvarado J. P, Gutiérrez L. M, Nedveck DA, Powers JS (2020)
Using large-scale tropical dry forest restoration to test successional theory. Ecological
Applications 30:e02116

Wickham H, Averick M, Bryan J, Chang W, D' L, Mcgowan A, et al. (2019) Welcome to the
Tidyverse. Journal of Open Source Software 4:1686

Wood SA, Sokol N, Bell CW, Bradford MA, Naeem S, Wallenstein MD, et al. (2016) Opposing

effects of different soil organic matter fractions on crop yields. Ecological Applications

26:2072–2085

Worthy SJ, Laughlin Daniel C, Zambrano Jenny, N Uma MI, Zhang Caicai, Lin Luxiang, et al.

(2020) Alternative designs and tropical tree seedling growth performance landscapes.

Wright SJ, Kitajima K, Kraft NJB, Reich PB, Wright IJ, Bunker DE, et al. (2010) Functional traits

and the growth-mortality trade-off in tropical trees. Ecology 91:3664–3674

Zirbel CR, Brudvig LA (2020) Trait–environment interactions affect plant establishment success

during restoration. Ecology 101:e02971

Zomer RJ, Xu J, Trabucco A (2022) Version 3 of the Global Aridity Index and Potential

Evapotranspiration Database. Scientific Data 2022 9:1 9:1–15

Zuur AF, Ieno EN, Walker N, Saveliev AA, Smith GM (2009) Mixed effects models and

extensions in ecology with R. Springer New York, New York, NY

776