Tools and Resources article

Forecasting protein evolution by integrating birth-death population models with structurally constrained substitution models

David Ferreiro[1,2], Luis Daniel González-Vázquez[1,2], Ana Prado-Comesaña[1] and Miguel Arenas[1,2,*]

[1]CINBIO, Universidade de Vigo, 36310 Vigo, Spain.

[2]Department of Biochemistry, Genetics and Immunology, Universidade de Vigo, 36310 Vigo, Spain.

Email addresses:

    DF: david.ferreiro.garcia@uvigo.es

    LDGV: luisdaniel.gonzalez@uvigo.gal

    APC: ana.maria.prado.comesana@uvigo.gal

    MA: marenas@uvigo.es

* Corresponding author:
Miguel Arenas
Department of Biochemistry, Genetics and Immunology
University of Vigo
36310 Vigo, Spain
Email: marenas@uvigo.es
Phone: +34 986 130047

Running head: Forecasting protein evolution

Keywords: Molecular evolution, Forecasting evolution, Birth-death process, Substitution model, Protein folding stability, Phylogenetics, Molecular adaptation

Abstract

Evolutionary studies in population genetics and ecology were mainly focused on predicting

and understanding past evolutionary events. Recently, however, a growing trend explores the

prediction of evolutionary trajectories toward the future promoted by its wide variety of

applications. In this context, we introduce a forecasting protein evolution method that

integrates birth-death population models with substitution models that consider selection on

protein folding stability. In contrast to traditional population genetics methods that usually

make the unrealistic assumption of simulating molecular evolution separately from the

evolutionary history, the present method combines both processes to simultaneously model

forward-in-time birth-death evolutionary trajectories and protein evolution under structurally

constrained substitution models that outperformed traditional empirical substitution models.

We implemented the method into a freely available computer framework. We evaluated the

accuracy of the predictions with several monitored viral proteins of broad interest. Overall,

the method showed acceptable errors in predicting the folding stability of the forecasted

protein variants but, expectedly, the errors grew up in the prediction of the corresponding

sequences. We conclude that forecasting protein evolution is feasible in certain evolutionary

scenarios and provide suggestions to enhance its accuracy by improving the underlying

models of evolution.

Introduction

Molecular evolution is traditionally investigated through inferences about past evolutionary events, such as phylogenetic tree and ancestral sequence reconstructions, and predictions about the future were considered as inaccessible for a long time because they can be affected by complex processes such as environmental change. Nevertheless, a variety of biological systems display a Darwinian evolutionary process where selection operates toward a limited set of adapted variants. These variants, and in extension the evolutionary trajectories to reach them, would be positively selected and could present a certain degree of predictability (Lässig *et al*, 2017; Wortel *et al*, 2023). The progress made in developing more accurate models of evolution (Arenas, 2015b) and the benefits from predicting the outcome of evolution (i.e., to understand the course of evolution or to prepare for the future (Wortel *et al.*, 2023)) motivated a variety of investigations on forecasting evolution in diverse fields including medicine, agriculture, biotechnology, and conservation biology, among others (e.g., Barton *et al*, 2016; Bull & Molineux, 2008; de Visser *et al*, 2018; Diaz-Uriarte & Vasallo, 2019; Fischer *et al*, 2015; Gerrish & Sniegowski, 2012; Lassig & Luksza, 2014; Lind *et al*, 2019; Luksza & Lassig, 2014; Morris *et al*, 2018; Munck *et al*, 2014; Neher *et al*, 2014; Wortel *et al.*, 2023). Unfortunately, forecasting evolution not always is achievable. Under neutral evolution, all the molecular variants are equally likely to be present in the population, showing lack of repeatability and disallowing accurate prediction of future variants. Thus, forecasting evolution requires a system with measurable selection pressures, and where certain positively selected variants could produce more descendants than other variants and expand in the population (Desai & Fisher, 2007; Goyal *et al*, 2012; Neher & Hallatschek, 2013; Neher *et al.*, 2014). Actually, a rougher fitness landscape resulting from selection can lead to greater accuracy in evolutionary predictions (Papkou *et al*, 2023; Rubin *et al*, 2023; Van Cleve &

Weissman, 2015). Overall, an evolutionary process could be predictable to some extent

(prediction errors are inevitable with any method and in any evolutionary scenario) depending

on the strength of selection driving evolution and the heterogeneity in fitness among different

variants.

Here, we focus on forecasting protein evolution because it involves molecular evolutionary

processes driven by selection pressures and where the fitness of each variant can be

parameterized and predicted (Carneiro & Hartl, 2010; Gilson *et al*, 2017). Traditionally,

evolutionary histories and ancestral sequences of proteins are inferred using probabilistic

methods based on advanced substitution models of protein evolution (e.g., Arenas, 2015b;

Arenas & Bastolla, 2020; Ferreiro *et al*, 2022; Malcom *et al*, 1990; Moreira *et al*, 2023;

Stackhouse *et al*, 1990; Thornton *et al*, 2003; Ugalde *et al*, 2004). The accuracy of these

inferences is affected by the accuracy of the applied substitution model, where substitution

models that better fit with the study data usually produce more accurate phylogenetic trees

and ancestral sequences (Arenas & Bastolla, 2020; Del Amparo & Arenas, 2022a, 2023;

Lemmon & Moriarty, 2004). These findings suggest that accurate substitution models of

evolution are also convenient for forecasting protein evolution. In this regard, a variety of

studies showed that structurally constrained substitution (SCS) models of protein evolution

provide more accurate evolutionary inferences than the traditional empirical substitution

models of protein evolution, in terms of phylogenetic likelihood, distribution of amino acid

frequencies among protein sites, rates of molecular evolution and folding stability of

reconstructed proteins, among other aspects (e.g., Arenas & Bastolla, 2020; Arenas *et al*,

2015; Bastolla *et al*, 2006; Bordner & Mittelmann, 2013; Del Amparo *et al*, 2023; Echave &

Wilke, 2017; Ferreiro *et al*, 2024a; Ferreiro *et al*, 2024b; Fornasari *et al*, 2002; Parisi &

Echave, 2001; Pascual-Garcia *et al*, 2019; Rodrigue *et al*, 2005), although SCS models

usually demand more computational resources than substitution models that only include

information from the protein sequence. Notice that the protein structure provides information

about the location and molecular interactions of amino acids at different protein sites, which

could be far from each other in the sequence but close in the three-dimensional structure and

interact affecting their evolution (Morcos *et al*, 2011; Ruiz-Gonzalez & Fares, 2013). Indeed,

selection from the protein folding stability is relevant in the evolution of multiple proteins,

including those in microbial and viral systems (e.g., Ferreiro *et al.*, 2022; Gong *et al*, 2013;

Jacquier *et al*, 2013; Rodrigues *et al*, 2016; Wylie & Shakhnovich, 2011; Zeldovich *et al*,

2007). Therefore, we believe that it should be taken into account for forecasting protein

evolution in such systems.

Predictions about future evolutionary events can be performed with simulation-based methods

(e.g., Eccleston *et al*, 2023; Neher *et al.*, 2014; Yoshida *et al*, 2023). In order to simulate

molecular evolution, traditional population genetics methods apply two separate steps

(Arenas, 2012; Hoban *et al*, 2012). First, the simulation of the evolutionary history (i.e., a

phylogenetic tree) using approaches such as the coalescent and birth-death population

processes (Gernhard, 2008; Hudson, 1990; Kingman, 1982; Stadler, 2010). Afterward, the

forward-in-time simulation of molecular evolution is performed, from the root node to the tip

nodes, upon the previously simulated evolutionary history (Yang, 2006). This methodology

was implemented into a variety of population genetics frameworks that simulate molecular

evolution (Arenas, 2012; Hoban *et al.*, 2012). However, for technical and computational

simplicity, it assumes that the simulation of the evolutionary history is independent from the

simulation of molecular evolution, which can produce biological incoherences (i.e., the

evolutionary history is usually simulated under neutral evolution while molecular evolution is

usually simulated with substitution models that consider selection). To enhance the realism of

this modeling, here we merged both processes into a single one where evolutionary history

influences molecular evolution and *vice versa*. In particular, we adopted a birth-death

population genetics method to simulate the forward-in-time evolutionary history (already used

for forecasting evolution (Lassig & Luksza, 2014; Neher *et al.*, 2014)), taking into account the

fitness of the molecular variant (through evolutionary constraints from the protein folding

stability (Bastolla & Demetrius, 2005; Gong *et al.*, 2013; Liberles *et al*, 2012; Zeldovich *et*

al., 2007)) at the corresponding node, to determine its subsequent birth or death event, and we

integrated this process with SCS models to model protein evolution along the derived

phylogenetic branches. The method is detailed below, and we implemented it into a new

version of our computer framework *ProteinEvolver* (Arenas *et al*, 2013), which is freely

available from https://github.com/MiguelArenas/proteinevolver. *ProteinEvolver2* includes

detailed documentation and a variety of ready-to-use examples. Next, considering the

potential applications of forecasting evolution to design vaccines and therapies against

pathogens, we evaluated and applied the method to forecasting protein evolution in several

real protein data of viruses monitored over time.

138 Materials and Methods

A method for forecasting protein evolution by combining birth-death population

genetics with structurally constrained substitution models of protein evolution

Following previous methods for forecasting evolution based on simulations (Lassig &

Luksza, 2014; Neher *et al.*, 2014), we developed a method to simulate the forward-in-time

evolutionary history of a protein sample with a birth-death process that considers the fitness

of the protein variant (based on folding stability) at every temporal node. The method derives

the birth and death rates for a protein variant based on the fitness of that variant, where a high

fitness results in a high birth rate and a low death rate, which can lead to a large number of

descendants, and the opposite leading to a few or none (extinction) descendants. Thus, the

fitness of the molecular variant at every node drives its corresponding forward in time birth-

death evolutionary history. The details of this simulation process are outlined below.

First, similarly to common simulators of molecular evolution (Arenas, 2012; Hoban *et al.*,

2012; Yang, 2006), a given protein sequence and structure (hereafter, protein variant) is

assigned to the root node. The fitness (*f*) of the protein variant (*A*) is calculated from its

folding stability (free energy, *ΔG*) following the Boltzmann distribution (Goldstein, 2013) (I,

which takes values from 0 to 1),

$$f(A) = \frac{1}{1 + e^{\Delta G / kT}} \qquad \text{(I)}$$

The user can alternatively choose whether the fitness of the protein variant is based only on its

folding stability or whether it is based on similarity with the stability of a real protein variant

(i.e., a protein structure from the Protein Data Bank, PDB). We believe that the latter can be

convenient since, in nature, a high folding stability does not necessarily indicate a high fitness

but a stability close to real stability may suggest a proper fitness because the real stability is

the result of a selection process. If the fitness is derived from only the folding stability of the

protein variant, the birth rate (*b*) is considered equal to the fitness. Alternatively, if the fitness

is determined based on the similarity in folding stability between the modeled variant and a

real variant, the birth rate is assumed to be 1 minus the root mean square deviation (RMSD) in

folding stability. Notice that the smaller this difference, the higher the birth rate. In both

cases, the death rate (*d*) is considered as *1-b* to allow a constant global (birth-death) rate. The

birth-death process is simulated forward in time deciding whether every next event is a birth

or a death event (Harmon, 2019; Stadler, 2010; Stadler, 2011) according to the fitness of the

corresponding molecular variant, and it ends when a user-specified criterion, such as a

particular sample size (considering or ignoring extinction nodes) or a certain evolutionary

time (t_e), is reached. Starting from an "active" node (i.e., the root node) at current time t_c, the

time to the next event (birth to produce two descendants or death to produce the extinction of

the node) can be calculated (details below). In contrast to standard birth-death processes

where birth and death rates are constant over time and among lineages, the present method

considers heterogeneity where each protein variant at a node has specific birth and death rates

according to its corresponding fitness. The method is described below and summarized in

Figure 1.

1. The process starts at the root node, assigning a user-specified protein sequence and

179   corresponding protein structure (i.e., obtained from the PDB) to that node. In general,

180   for every protein variant assigned to an active node, the corresponding birth and death

181   rates are calculated following the indications presented above.

182   2. Calculation of the time to the next birth or death event. Following common birth-death

183   methods, the time to the next event t_n is calculated through an exponential distribution

184   with rate based on the number of active nodes (s) and the sum of the birth and death

185   rates (Harmon, 2019) (II),

186
$$t_n = e^{(s(b+d))} \qquad \text{(II)}$$

187   As a consequence, since $b+d=1$ at each node, t_n is consistent across all nodes,

188   according to (Harmon, 2019).

189   3. Evaluate whether the simulation concludes before the next event occurs $(t_c + t_n)$.

190   4. If it does conclude (i.e., $t_c + t_n$ higher than t_e), the simulation of the evolutionary

191   history finishes.

192   5. If it does not conclude, a random active node is selected, and its protein variant is

193   analyzed to determine its type of next event (birth or death). The probability of a birth

194   event is $P_b = b/(b + d)$ and the probability of a death event is $P_d = d/(b + d)$. A

195   random sample from those probabilities is taken to determine the type of evolutionary

196   event.

6. 197   If a birth event is selected. Two descendant active nodes from the study node are

198   incorporated with branch lengths t_n, and the study node is then considered inactive.

199   Next, molecular evolution is simulated from the original node to each descendant node

200   based on the specified SCS model of protein evolution (Arenas *et al.*, 2013). Thus, the

201   process results in a protein variant for every descendant node. Finally, the folding

202   stability and subsequent fitness of these descendant protein variants are calculated.

7. 203   If a death event is selected. Then, the study node is considered inactive.

8. 204   Return to step 1 while the user-specified criterion for ending the birth-death process is

not satisfied and at least one active node exists in the evolutionary history. Otherwise,

the simulation ends.

This process simultaneously simulates, forward in time, evolutionary history and protein

evolution, with protein evolution influencing the evolutionary history through selection from

the folding stability. Indeed, selection can vary among protein variants at their corresponding

nodes of the evolutionary history. The process produces a forward in time birth-death

phylogenetic history, that encompasses nodes that reached the ending time, internal nodes and

nodes that were extinct at some time, along with the protein variant associated with each

node.

The method includes several optional capabilities listed in Table S1 (Supplementary Material)

and in the software documentation, with some summarized below.

• The user can fix the birth and death rates along the evolutionary history or specify that

they are based on the fitness of the corresponding protein variant as described before.

For the latter, the fitness can be based on the folding stability of the analyzed protein

variant or on the similarity in folding stability between the analyzed and real protein

variants.

• The birth-death simulation can finish any of the following criteria are met: (a) A

specified sample size, including extinction nodes derived from death events. (b) A

specified sample size, excluding extinction nodes. (c) A specified evolutionary time,

measured from the root to a tip node.

• Extinction nodes can either be preserved or removed from the evolutionary history.

• Evaluation of the simulated birth-death evolutionary history in terms of tree balance

using the Colless index (Colless, 1982; Lemant *et al*, 2022).

• Possible variation of the site-specific substitution rate according to user specifications.

• Customizable substitution model featuring site-specific exchangeability matrices

(relative rates of change among amino acids and their frequencies at the equilibrium).

• Implementation of several SCS models and a variety of empirical substitution models

of protein evolution. The implemented SCS models were evaluated in our previous

work (Arenas *et al.*, 2013) and, the implemented empirical substitution models were

properly identified using simulated data with *ProtTest3* (Darriba *et al*, 2011).

The implemented SCS models consider molecular energy functions based on amino acid

contact matrices and configurational entropies per residue in unfolded and misfolded proteins

(Arenas *et al.*, 2013; Bastolla *et al*, 2007; Mendez *et al*, 2010). Technical details about these

SCS models are presented in our previous study (Arenas *et al.*, 2013). The method also

implements common empirical substitution models of protein evolution (i.e., *Blosum62*,

CpRev, *Dayhoff*, *DayhoffDCMUT*, *FLU*, *HIVb*, *HIVw*, *JTT*, *JonesDCMUT*, *LG*, *Mtart*,

Mtmam, *Mtrev24*, *RtRev*, *VT* and *WAG*; Table S1) and the user can specify any particular

exchangeability matrix for all sites or for each site, allowing for heterogeneity of the

substitution process among sites (details in Table S1 and in the software documentation). In

addition, the framework implements heterogeneous substitution rates among sites by the

traditional Gamma distribution (+G) (Yang, 1996) and proportion of invariable sites (+I)

(Fitch & Margoliash, 1967), and also the user can directly alter the substitution rate at each

site for any empirical or SCS model (Table S1). Regarding the evolutionary history, in

addition to the birth-death process presented before, the user can specify a particular

phylogenetic tree or simulate a coalescent evolutionary history (Hudson, 1983; Kingman,

1982) (Table S1). In this regard, we maintained the capabilities of the previous version,

including the coalescent with recombination (Hudson, 1983) [which can be homogeneous or

heterogeneous along the sequence (according to Wiuf & Posada, 2003)], variable population

size over time (growth rate or demographic periods), several migration models [i.e., island

(Hudson, 1998), stepping-stone (Kimura & Weiss, 1964) and continent-island (Wright, 1931)]

with temporal variation of migration rates and convergence of demes or subpopulations,

simulation of haploid or diploid data, and longitudinal sampling (Navascues *et al*, 2010)

(details in Table S1 and in the software documentation). The framework outputs a simulated

multiple sequence alignment with the protein sequences of the internal and tip nodes, as well

as their folding stabilities, and the evolutionary history, among other information (Table S1

and software documentation).

Study data for evaluating the method for forecasting protein evolution

We evaluated the accuracy of the method for forecasting protein evolution using viral proteins

sampled over time (longitudinal sampling). Specifically, we used protein sequences from

previous experiments of virus evolution monitored over time, which contain consensus

molecular data (avoiding rare variants) that belong to different evolutionary time points.

• The matrix (MA) protein of HIV-1, with data obtained from an *in vitro* cell culture

experiment where samples were collected at different times (Arenas *et al*, 2016). This

data included 21 MA protein consensus sequences collected at times (population

passages, *T*) *T1* (initial time and includes an initial sequence) and *T31* (time after 31

passages and includes 20 sequences), which accumulated 48 amino acid substitutions

(sequence identity 0.973; Figure S1, Supplementary Material).

• The main protease (Mpro) and papain-like protease (PLpro) proteins of SARS-CoV-2.

For each protein, the data includes the first sequenced variant (Wuhan) and a sequence

built with all the substitutions observed in a dataset of 384 genomes of the Omicron

variant of concern collected from the GISAID database. The resulting Mpro and

PLpro sequences presented 10 and 22 substitutions (sequence identity 0.967 and

0.930), respectively.

• The protease (PR) of HIV-1 with data sampled from patients monitored over time,

from 2008 to 2017, available from the Specialized Assistance Services in Sexually

Transmissible Diseases and HIV/AIDS in Brazil (Ferreiro *et al.*, 2022; Santos-Pereira

et al, 2021; Souto *et al*, 2021). These evolutionary scenarios are complex due to the

diverse antiretroviral therapies administered to the patients (Table S2; Supplementary

Material), which could vary during the studied time periods and that could promote the

fixation of specific mutations (i.e., associated with resistance) (Ferreiro *et al.*, 2022;

Santos-Pereira *et al.*, 2021; Souto *et al.*, 2021). For each viral population (patient), the

data included a consensus sequence for each of the four or five samples collected at

different time points. These consensus sequences exhibited between one and 22 amino

acid substitutions respect to the consensus sequence of the corresponding first sample

(Table S2).

We identified a representative protein structure for each data, which was used to predict the

folding stability and to inform the SCS model. In particular, we obtained the protein structures

of the sequences at the initial time (*T1*) by homology modelling. We used *SWISS-MODEL*

(Arnold *et al*, 2006) to identify the best-fitting templates of PDB structures (Table S3,

Supplementary Material). Next, we predicted the protein structures by homology modelling

with *Modeller* (Sali & Blundell, 1993) using the protein sequences and corresponding

structural templates.

Forward in time prediction of viral protein variants

The evaluation was performed with the previously presented real data, which includes a real

protein variant present at the initial time (*T1*) and subsequent protein variants present at a later

time (*Tn*). We applied the method for forecasting protein evolution to predict the most likely

protein variants at *Tn* derived from the real protein variant observed at the initial time (*T1*).

The prediction error was determined by measuring the distance between the real protein

variants and the predicted variants corresponding to time *Tn*.

Thus, we assigned to the initial node (*T1*) the corresponding real protein variant (including its

sequence and structure), and its evolution was simulated forward in time until it reached the

number of substitutions observed in the real data at time *Tn*. Thus, we considered the number

of observed substitutions as a measure of evolutionary time to allow proper comparisons

between real and predicted variants. We simulated 100 alignments of protein sequences, each

containing the same number of sequences as the real data at *Tn*. This included 20 HIV-1 MA

protein sequences, a consensus sequence for the SARS-CoV-2 Mpro, a consensus sequence

for the SARS-CoV-2 PLpro and, a consensus sequence of HIV-1 PR for every viral

population. To avoid rare variants, each sequence of the simulated multiple sequence

alignments was obtained as the consensus of 100 linked simulated sequences.

To investigate the effect of selection on the predictions, we compared the accuracy of

forecasting protein evolution when selection from the protein structure is considered and,

when it is ignored (neutral evolution). If selection is considered, as previously presented, the

probability of birth and death events was based on the fitness of the protein variants, and

protein evolution was modeled using an SCS model (Arenas *et al.*, 2013). For the case of

neutral evolution, all protein variants equally fit and are allowed, leading to only birth events,

and we modeled protein evolution using an exchangeability matrix with the same relative

rates of change for any pair of amino acids.

Accuracy of the predicted protein variants

We assessed the accuracy of the method for forecasting protein evolution by comparing the

predicted and real protein variants present at time *Tn*, both at the protein sequence and

structure levels.

For data containing multiple sequences at time *Tn*, we calculated the Kullback-Leibler (KL)

divergence, which provides a distance between two multiple sequence alignments (the real

and predicted data) based on the distribution of amino acid frequencies along the sequences

(Equation III, where factors *P* and *Q* denote the distribution of amino acid frequencies in the

real and predicted protein sequences at time *Tn*, respectively, *i* refers to protein site)

(Kullback & Leibler, 1951). This distance was only calculated for data with a set of sequences

at time Tn (HIV-1 MA) because a single sequence does not provide site-specific variability.

$$KL\ (P||Q) = \sum_i P_i \times \log(\frac{P_i}{Q_i}) \qquad \text{(III)}$$

We also compared the real and predicted evolutionary trajectories of protein variants using the

Grantham distance, which measures the differences between amino acids based on their

physicochemical properties (Grantham, 1974). In particular, for both real and predicted

protein variants, we calculated the Grantham distance at each protein site that differs between

the two datasets, considering its evolution from $T1$ to the subsequent multiple sequence

alignment at Tn. We examined sites that varied over time, thus the general site-specific

Grantham distance Gi was calculated as the frequency of each amino acid f at site i multiplied

by the specific Grantham distance between amino acid m at time Tn and amino acid n at time

$T1$, normalized with the largest Grantham distance G_{max} to obtain values between 0 and 1

(Equation IV). Next, to compare the real and predicted data, we calculated the site-specific

difference of Grantham distance Gb_i between the real P and predicted Q protein variants

(Equation V),

$$G_i = \sum_{m=1}^{20} f_m \times \frac{G_{(m,n)}}{G_{max}} \qquad \text{(IV)}$$

$$Gb_i = |G_{P,i} - G_{Q,i}| \qquad \text{(V)}$$

In addition, we obtained and compared the protein folding stability (ΔG) of the predicted and

real protein variants observed at time Tn, using their corresponding protein structures, with

DeltaGREM (Arenas *et al.*, 2015; Minning *et al*, 2013).

Results

Implementation of the forecasting protein evolution method

We extended the previous version of our framework *ProteinEvolver* (Arenas *et al.*, 2013),

maintaining its previous capabilities (i.e., simulation of protein evolution upon user-specified

phylogenetic trees and upon phylogenetic trees simulated with the coalescent with or without

recombination, migration, demographics and longitudinal sampling, empirical and SCS

models, among others; Table S1), by adding, among others (Table S1), the forward in time

modeling of protein evolution that combines a birth-death process based on the fitness of

every protein variant (folding stability) at each node to determine its birth and death rates, as

well as SCS models of protein evolution. The framework *ProteinEvolver2* is written in C and

distributed with a detailed documentation and a variety of illustrative practical examples. The

framework is freely available from https://github.com/MiguelArenas/proteinevolver.

Evaluation of predictions of HIV-1 MA evolution

Regarding the evolution of the HIV-1 MA protein, the Grantham distance and the KL

divergence between the real variants at time *Tn* and the corresponding predicted variants were

low (around 5% and 6%, respectively; Table 1), and they did not differ comparing predictions

that consider selection on the folding stability and predictions that ignore it (Table 1). On the

other hand, we found that the folding stability of the protein variants predicted considering

selection on the folding stability was closer to the folding stability of the real protein variants

than that of the protein variants predicted under neutral evolution (Table 1). In particular, the

protein variants predicted ignoring selection were less stable than those predicted considering

selection and also less stable than the real protein variants.

Evaluation of predictions of SARS-CoV-2 Mpro and PLpro evolution

The analyses of the SARS-CoV-2 Mpro and PLpro data the Grantham distance between the

real and predicted sequences was around 25% and 36%, respectively (Figure 2A). Again, this

distance was similar when comparing predictions based on a model that considers selection

on the protein folding stability and a model of neutral evolution. Regarding comparisons

based on the protein folding stability, we found again that the model that considers selection

from the folding stability produces variants closer to the stability of the real protein variants

than the model that ignores selection (Figure 2B). Indeed, the protein sequences derived from

the model that considers selection were more stable than those derived from the model of

neutral evolution.

Evaluation of predictions of HIV-1 PR evolution

In general, the Grantham distance, which compared the evolutionary trajectories of the real

and predicted protein variants from time *T1* to later times, varied among viral populations

(patients) (Figure 3A). However, for the majority of these populations, the distance remained

below 30% and exhibited minor fluctuations over time. One particular population exhibited a

notable trend, with the distance increasing from 10% to nearly 60% over time. Considering

that the length of the evolutionary trajectories of the protein can differ among the studied

populations, we explored whether the accumulated number of amino acid substitutions could

affect the accuracy of the predictions, and we found that the number of substitutions varied

among populations and this variability did not correlate with the Grantham distance between

the real and predicted data (R^2=0.0001, Figure 3B). In general, the folding stability of the

predicted protein variants was similar or slightly less stable than that of the real protein

variants [with a difference ranging from 0 to 9 kcal/mol and a mean of 3.1 ± 0.9 (95% CI)

kcal/mol; Figure 3C]. Indeed, the folding stability exhibited small fluctuations, increasing and

decreasing, over time.

Discussion

While reconstructing evolutionary histories and ancestral sequences, among other inferences

about the past, was popular in the field, predictions about evolution toward the future were

traditionally ignored due to potential high prediction errors. However, in the last decade,

forecasting evolution has gained attention because of its wide variety of applications and

advancements in models of evolution (see the reviews Lässig *et al.*, 2017; Wortel *et al.*,

2023). Several studies showed that forecasting evolution can be feasible in some systems,

including virus evolution (Luksza & Lassig, 2014; Thadani *et al*, 2023). Here, we also

investigated forecasting evolution in viruses but at the molecular level, considering the

relevance of the substitution process to produce molecular variants that affect the viral fitness

(Arenas, 2015a; Arenas *et al.*, 2016; Bloom & Neher, 2023; Poon *et al*, 2007; Watabe &

Kishino, 2010). We believe that forecasting genome evolution remains challenging due to the

complexity of its evolutionary processes [i.e., including epistatic interactions, chromosomal

rearrangements and heterogeneous substitution patterns among genomic regions (Arbiza *et al*,

2011), among others] that complicate the parameterization and prediction of accurate fitness

landscapes. However, we believe that it could be more feasible in structural proteins because

of their relatively simpler evolutionary processes that usually include selection on folding

stability (Bloom *et al*, 2006; Goldstein, 2011).

To make evolutionary predictions over time, either toward the past or toward the future,

considering a substitution model of molecular evolution can be convenient. Actually, a variety

of traditional studies showed that the substitution model influences the phylogenetic

likelihood generated by probabilistic approaches used for evolutionary inference (Lemmon &

Moriarty, 2004; Yang *et al*, 1994; Zhang & Nei, 1997). To study protein evolution, empirical

substitution models of protein evolution are routinely used because they allow rapid

calculations based on the assumption of site-independent evolution and typically apply the

same exchangeability matrix for all the protein sites, which is highly unrealistic (Echave *et al*,

2016). Besides, a small set of empirical substitution models was developed for modeling the

evolution of the diverse range of viral proteins (Dang *et al*, 2010; Del Amparo & Arenas,

2022b; Dimmic *et al*, 2002; Le & Vinh, 2020; Nickle *et al*, 2007). As an alternative, some

substitution models that consider evolutionary constraints on the protein structure

incorporated site-dependent evolution, which allows a more accurate modeling of protein

evolution compared to the empirical substitution models (Arenas *et al.*, 2013; Arenas *et al.*,

2015; Bordner & Mittelmann, 2013; Ferreiro *et al.*, 2024a; Parisi & Echave, 2005).

According to previous methods for forecasting evolution based on computer simulations

(Lassig & Luksza, 2014; Neher *et al.*, 2014), we also adopted the birth-death population

genetics process to allow a forward-in-time evolutionary process where the birth and death

rates can differ among nodes (Stadler, 2010; Stadler, 2011). Traditional population genetics

methods to simulate the evolution of molecular data implement a two-step simulation process,

where first the evolutionary history is simulated [i.e., with the coalescent theory or a forward-

in-time simulation approach (Arenas, 2012; Hoban *et al.*, 2012), often assuming neutral

evolution] and, in a subsequent step, molecular evolution is simulated upon the previously

obtained evolutionary history (Yang, 2006). However, this methodology is unrealistic because

the fitness of the data can affect its evolutionary history (i.e., a variant with high fitness is

likely to have more descendants than a variant with low fitness). Thus, we designed and

implemented a method for forecasting protein evolution that integrates a birth-death

population genetics process with a SCS model of protein evolution, where the folding stability

of each protein variant is evaluated to predict its future trajectory in terms of both

evolutionary history and molecular evolution. We implemented this forward-in-time

simulation of protein evolution in a new version of our previous framework *ProteinEvolver*

(Arenas *et al.*, 2013), while maintaining its previous capabilities and extending some of them

(i.e., incorporation of site-specific exchangeability matrices and additional substitution

models of protein evolution, among others; see Table S1 and software documentation).

Next, we evaluated the forward-in-time evolutionary predictions with real data of HIV-1 and

SARS-CoV-2 proteins. We determined the prediction errors between the real and the

predicted protein variants by examining dissimilarity in evolutionary trajectory (Grantham

distance based on physicochemical properties among the differing amino acids during the

evolutionary trajectories), sequence divergence (distribution of amino acid frequencies among

sites using the KL divergence), and protein structure (protein folding stability). Additionally,

we analyzed the influence of accounting for selection, based on the protein folding stability,

on the predictions.

In general, the sequence and evolutionary trajectory dissimilarities between the real and

predicted protein variants were relatively small. For the HIV-1 MA protein, the SARS-CoV-2

Mpro and the SARS-CoV-2 PLpro, the sequence dissimilarity was below 10%, 26% and 36%,

respectively. The low prediction errors for the HIV-1 MA protein were expected because the

data was derived from an *in vitro* cell culture experiment that is not influenced by a variety of

external factors that could affect the predictions in *in vivo* experiments. As a rough reference,

in traditional ancestral sequence reconstruction (ASR) of protein data with high sequence

identity the error was approximately 2% (Arenas & Posada, 2010). Notice that ASR methods

can exhibit lower prediction errors due to their statistical evaluation based on a set of original

sequences, rather than relying on a single original sequence used in the case of forecasting

evolution. The contrasting scenario regarding evolutionary complexity was the HIV-1 PR

data, which was sampled from populations (patients) undergoing various antiretroviral

treatments (Ferreiro *et al.*, 2022). There, the sequence dissimilarity varied among viral

populations, although most were below 30%. A viral population exhibited higher

dissimilarities (near 60%) that we believe could be caused by molecular adaptations promoted

by the therapies, although this needs formal investigation.

An unexpected result was that the model of neutral evolution produced sequence

dissimilarities between the real and predicted protein variants that were similar to those

obtained with the SCS model (Table 1 and Figure 2). A few studies indicated that the

substitution model has negligible effects on the reconstructed phylogenetic trees (Abadi *et al*,

2019; Spielman, 2020). Subsequent studies found that the influence of the substitution model

on phylogenetic reconstructions is dependent on the diversity of the data, where data with

high diversity is more sensible to the applied substitution model due to containing more

evolutionary information to be modeled (Del Amparo & Arenas, 2022a, 2023). Considering

that the data studied here has overall low diversity (i.e., sequence identity of 0.973, 0.967 and

0.930 for the HIV MA data, SARS-CoV-2 Mpro data and SARS-CoV-2 PLpro data,

respectively; it is important to note that, in general, longitudinal data derived from monitored

evolutionary processes usually show a low diversity because involve relatively short

evolutionary histories, among other factors), we believe that the influence of the applied

substitution models on the prediction of the sequences was small because of the small number

of modeled substitution events, as found in phylogenetic reconstructions (Del Amparo &

Arenas, 2022a, 2023). In all, the prediction error related to sequence dissimilarity varied

among the studied evolutionary scenarios; as expected, it was higher for the more complex

scenarios.

We also evaluated the prediction error between the real and predicted protein variants

regarding their folding stability, again comparing the predictions made under a model that

considers structural constraints and a model of neutral evolution. In general, the protein

variants predicted under the SCS model presented a folding stability close to the folding

stability of the respective real protein variants, with differences below 1, 2 and 7 kcal/mol for

the HIV MA data, SARS-CoV-2 Mpro data and SARS-CoV-2 PLpro data, respectively. The

higher differences (around 9 kcal/mol) were again observed for the HIV-1 PR data. In contrast

to the prediction error based on sequence dissimilarity, the prediction error based on folding

stability varies between predictions obtained under the SCS model and those obtained under

the neutral model. In the studied evolutionary scenarios, the protein variants predicted under

the neutral model were less stable and farther from the stability of the real protein variants

compared to those predicted under the SCS model. These results were expected because,

under SCS models, protein stability can be modeled with greater accuracy than sequence

similarity due to selection for maintaining stability in the protein structure despite amino acid

changes (Arenas & Bastolla, 2020; Illergard *et al*, 2009; Pascual-Garcia *et al*, 2010). Indeed,

previous studies showed that models that ignore structural constraints often produce proteins

with unrealistic folding instability (Arenas & Bastolla, 2020; Del Amparo *et al.*, 2023), which

suggests that accounting for protein folding stability in the modeling of protein evolution is

recommended for predicting protein variants with appropriate structural properties.

We present a method to simulate forward-in-time protein evolution accounting for

evolutionary constraints from the protein structure and a birth-death population process, and

where the evolutionary history is influenced by the protein evolution and *vice versa*. The

524 method is implemented in the computer framework *ProteinEvolver2*, which is freely

525 distributed with several practical examples and detailed documentation. We believe that

526 implementing methods into freely available phylogenetic frameworks is important to facilitate

527 practical applications, as well as future improvements and evaluations. We applied the

528 method to forecast protein evolution in some viral proteins. We found that the method

529 provides acceptable approximations to the real evolution, especially in terms of protein

530 folding stability, suggesting that combining structural constraints with birth-death population

531 processes in the modeling of protein evolution is convenient. Still, to advance in methods for

532 forecasting protein evolution, we believe that further efforts should be made in the field to

533 improve the modeling of protein evolution, such as the incorporation of site-dependent

534 evolutionary constraints from the protein activity.

536 **Supplementary Material**

537 The supplementary material is available at the journal online.

539 **Data Availability**

540 The computer framework *ProteinEvolver2* is freely available from

541 https://github.com/MiguelArenas/proteinevolver. The SARS-CoV-2 data is available from

542 GISAID database with DOI https://doi.org/10.55876/gis8.250206gt. The real and predicted

543 protein variants are available from Zenodo repository at the URL

544 https://doi.org/10.5281/zenodo.14767475.

546 **Acknowledgments**

548   This work was supported by the Project PID2023-151032NB-C22 funded by

549   MCIU/AEI/10.13039/501100011033 and by FEDER, UE. D.F. was funded by a fellowship

550   from Xunta de Galicia [ED481A-2020/192]. We thank the "Centro de Supercomputación de

551   Galicia (CESGA)" for the computer resources. The funders had no role in study design, data

552   collection and analysis, decision to publish, or preparation of the manuscript.

Declaration of Interests

555   The authors declare no competing interests.

References

Abadi S, Azouri D, Pupko T, Mayrose I (2019) Model selection may not be a mandatory step
for phylogeny reconstruction. *Nat Commun* 10: 934

Arbiza L, Patricio M, Dopazo H, Posada D (2011) Genome-wide heterogeneity of nucleotide
substitution model fit. *Genome Biol Evol* 3: 896-908

Arenas M (2012) Simulation of Molecular Data under Diverse Evolutionary Scenarios. *PLoS*
Comput Biol 8: e1002495

Arenas M (2015a) Genetic Consequences of Antiviral Therapy on HIV-1. *Comput Math*
Methods Med 2015: 9

Arenas M (2015b) Trends in substitution models of molecular evolution. *Front Genet* 6: 319

Arenas M, Bastolla U (2020) ProtASR2: Ancestral reconstruction of protein sequences
accounting for folding stability. *Methods Ecol Evol* 11: 248-257

Arenas M, Dos Santos HG, Posada D, Bastolla U (2013) Protein evolution along phylogenetic
histories under structurally constrained substitution models. *Bioinformatics* 29: 3020-
3028

Arenas M, Lorenzo-Redondo R, Lopez-Galindez C (2016) Influence of mutation and
recombination on HIV-1 in vitro fitness recovery. *Mol Phylogenet Evol* 94: 264-270

Arenas M, Posada D (2010) The effect of recombination on the reconstruction of ancestral
sequences. *Genetics* 184: 1133-1139

Arenas M, Sanchez-Cobos A, Bastolla U (2015) Maximum likelihood phylogenetic inference
with selection on protein folding stability. *Mol Biol Evol* 32: 2195-2207

Arnold K, Bordoli L, Kopp J, Schwede T (2006) The SWISS-MODEL workspace: a web-
based environment for protein structure homology modelling. *Bioinformatics* 22: 195-201

Barton JP, Goonetilleke N, Butler TC, Walker BD, McMichael AJ, Chakraborty AK (2016)
Relative rate and location of intra-host HIV evolution to evade cellular immunity are
predictable. *Nature Communications* 7: 11660

Bastolla U, Demetrius L (2005) Stability constraints and protein evolution: the role of chain
length, composition and disulfide bonds. *Protein Eng Des Sel* 18: 405-415

Bastolla U, Porto M, Roman HE, Vendruscolo M (2006) A protein evolution model with
independent sites that reproduces site-specific amino acid distributions from the Protein
Data Bank. *BMC Evol Biol* 6: 43

Bastolla U, Porto M, Roman HE, Vendruscolo M (2007) *Structural approaches to sequence
evolution*. Springer, Berlin, Heidelberg

Bloom JD, Labthavikul ST, Otey CR, Arnold FH (2006) Protein stability promotes
evolvability. *Proc Natl Acad Sci U S A* 103: 5869-5874

Bloom JD, Neher RA (2023) Fitness effects of mutations to SARS-CoV-2 proteins. *Virus
Evol* 9: vead055

Bordner AJ, Mittelmann HD (2013) A new formulation of protein evolutionary models that
account for structural constraints. *Mol Biol Evol* 31: 736-749

Bull JJ, Molineux IJ (2008) Predicting evolution from genomics: experimental evolution of
bacteriophage T7. *Heredity* 100: 453-463

Carneiro M, Hartl DL (2010) Colloquium papers: Adaptive landscapes and protein evolution.
Proc Natl Acad Sci U S A 107 Suppl 1: 1747-1751

Colless DH (1982) Review of Phylogenetics: The Theory and Practice of Phylogenetic
Systematics. *Systematic Zoology* 31: 100-104

Dang CC, Le QS, Gascuel O, Le VS (2010) FLU, an amino acid substitution model for
influenza proteins. *BMC Evol Biol* 10: 99

Darriba D, Taboada GL, Doallo R, Posada D (2011) ProtTest 3: fast selection of best-fit
models of protein evolution. *Bioinformatics* 27: 1164-1165

de Visser J, Elena SF, Fragata I, Matuszewski S (2018) The utility of fitness landscapes and
big data for predicting evolution. *Heredity (Edinb)* 121: 401-405

Del Amparo R, Arenas M (2022a) Consequences of Substitution Model Selection on Protein
Ancestral Sequence Reconstruction. *Molecular Biology and Evolution* 39: msac144

Del Amparo R, Arenas M (2022b) HIV Protease and Integrase Empirical Substitution Models
of Evolution: Protein-Specific Models Outperform Generalist Models. *Genes* 13

Del Amparo R, Arenas M (2023) Influence of substitution model selection on protein
phylogenetic tree reconstruction. *Gene* 865: 147336

Del Amparo R, González-Vázquez LD, Rodríguez-Moure L, Bastolla U, Arenas M (2023)
Consequences of Genetic Recombination on Protein Folding Stability. *Journal of*
Molecular Evolution 91: 33-45

Desai MM, Fisher DS (2007) Beneficial Mutation–Selection Balance and the Effect of
Linkage on Positive Selection. *Genetics* 176: 1759-1798

Diaz-Uriarte R, Vasallo C (2019) Every which way? On predicting tumor evolution using
cancer progression models. *PLoS Comput Biol* 15: e1007246

Dimmic MW, Rest JS, Mindell DP, Goldstein RA (2002) rtREV: an amino acid substitution
matrix for inference of retrovirus and reverse transcriptase phylogeny. *J Mol Evol* 55: 65-
73

Eccleston RC, Manko E, Campino S, Clark TG, Furnham N (2023) A computational method
for predicting the most likely evolutionary trajectories in the stepwise accumulation of
resistance mutations. *eLife* 12: e84756

Echave J, Spielman SJ, Wilke CO (2016) Causes of evolutionary rate variation among protein
sites. *Nat Rev Genet* 17: 109-121

Echave J, Wilke CO (2017) Biophysical Models of Protein Evolution: Understanding the
Patterns of Evolutionary Sequence Divergence. *Annu Rev Biophys* 46: 85-103

Ferreiro D, Branco C, Arenas M (2024a) Selection among site-dependent structurally
constrained substitution models of protein evolution by approximate Bayesian
computation. *Bioinformatics* 40: btae096

Ferreiro D, Khalil R, Gallego MJ, Osorio NS, Arenas M (2022) The evolution of the HIV-1
protease folding stability. *Virus Evol* 8: veac115

Ferreiro D, Khalil R, Sousa SF, Arenas M (2024b) Substitution Models of Protein Evolution
with Selection on Enzymatic Activity. *Molecular Biology and Evolution* 41: msae026

Fischer A, Vazquez-Garcia I, Mustonen V (2015) The value of monitoring to control evolving
populations. *Proc Natl Acad Sci U S A* 112: 1007-1012

Fitch WM, Margoliash E (1967) A method for estimating the number of invariant amino acid
coding positions in a gene using cytochrome c as a model case. *Biochem Genet* 1: 65-71
Fornasari MS, Parisi G, Echave J (2002) Site-specific amino acid replacement matrices from
structurally constrained protein evolution simulations. *Mol Biol Evol* 19: 352-356
Gernhard T (2008) The conditioned reconstructed process. *J Theor Biol* 253: 769-778
Gerrish PJ, Sniegowski PD (2012) Real time forecasting of near-future evolution. *J R Soc*
Interface 9: 2268-2278
Gilson AI, Marshall-Christensen A, Choi JM, Shakhnovich EI (2017) The Role of
Evolutionary Selection in the Dynamics of Protein Structure Evolution. *Biophys J* 112:
1350-1365
Goldstein RA (2011) The evolution and evolutionary consequences of marginal
thermostability in proteins. *Proteins* 79: 1396-1407
Goldstein RA (2013) Population Size Dependence of Fitness Effect Distribution and
Substitution Rate Probed by Biophysical Model of Protein Thermostability. *Genome Biol*
Evol 5: 1584-1593
Gong LI, Suchard MA, Bloom JD (2013) Stability-mediated epistasis constrains the evolution
of an influenza protein. *eLife* 2: e00631
Goyal S, Balick DJ, Jerison ER, Neher RA, Shraiman BI, Desai MM (2012) Dynamic
Mutation–Selection Balance as an Evolutionary Attractor. *Genetics* 191: 1309-1319
Grantham R (1974) Amino acid difference formula to help explain protein evolution. *Science*
185: 862-864
Harmon LJ (2019) Introduction to birth-death models. In: *Phylogenetic Comparative*
Methods, p. https://lukejharmon.github.io/pcm/chapter10_birthdeath/.
Hoban S, Bertorelle G, Gaggiotti OE (2012) Computer simulations: tools for population and
evolutionary genetics. *Nat Rev Genet* 13: 110-122
Hudson RR (1983) Properties of a neutral allele model with intragenic recombination.
Theoretical Population Biology 23: 183-201
Hudson RR (1990) Gene genealogies and the coalescent process. *Oxford Surveys in*
Evolutionary Biology 7: 1-44
Hudson RR (1998) Island models and the coalescent process. *Mol Ecol* 7: 413-418
Illergard K, Ardell DH, Elofsson A (2009) Structure is three to ten times more conserved than
sequence--a study of structural response in protein cores. *Proteins* 77: 499-508

Jacquier H, Birgy A, Le Nagard H, Mechulam Y, Schmitt E, Glodt J, Bercot B, Petit E,

Poulain J, Barnaud G *et al* (2013) Capturing the mutational landscape of the beta-

lactamase TEM-1. *Proceedings of the National Academy of Sciences* 110: 13067-13072

Kimura M, Weiss GH (1964) The Stepping Stone Model of Population Structure and the

Decrease of Genetic Correlation with Distance. *Genetics* 49: 561-576

Kingman JFC (1982) The coalescent. *Stochastic Processes and their Applications* 13: 235-

248

Kullback S, Leibler RA (1951) On information and sufficiency. *Ann Math Stat* 22: 79-86

Lassig M, Luksza M (2014) Can we read the future from a tree? *Elife* 3

Lässig M, Mustonen V, Walczak AM (2017) Predicting evolution. *Nature Ecology &*

Evolution 1: 0077

Le TK, Vinh LS (2020) FLAVI: An amino acid substitution model for flaviviruses. *J Mol*

Evol In press

Lemant J, Le Sueur C, Manojlović V, Noble R (2022) Robust, Universal Tree Balance

Indices. *Systematic Biology* 71: 1210-1224

Lemmon AR, Moriarty EC (2004) The importance of proper model assumption in bayesian

phylogenetics. *Syst Biol* 53: 265-277

Liberles DA, Teichmann SA, Bahar I, Bastolla U, Bloom J, Bornberg-Bauer E, Colwell LJ,

de Koning AP, Dokholyan NV, Echave J *et al* (2012) The interface of protein structure,

protein biophysics, and molecular evolution. *Protein Sci* 21: 769-785

Lind PA, Libby E, Herzog J, Rainey PB (2019) Predicting mutational routes to new adaptive

phenotypes. *Elife* 8

Luksza M, Lassig M (2014) A predictive fitness model for influenza. *Nature* 507: 57-61

Malcom BA, Wilson KP, Matthews BW, Kirsch JF, Wilson AC (1990) Ancestral lysozymes

reconstructed, neutrality tested, and thermostability linked to hydrocarbon packing.

Nature 345: 86-89

Mendez R, Fritsche M, Porto M, Bastolla U (2010) Mutation bias favors protein folding

stability in the evolution of small populations. *PLoS Comput Biol* 6: e1000767

Minning J, Porto M, Bastolla U (2013) Detecting selection for negative design in proteins

through an improved model of the misfolded state. *Proteins* 81: 1102-1112

Morcos F, Pagnani A, Lunt B, Bertolino A, Marks DS, Sander C, Zecchina R, Onuchic JN,
Hwa T, Weigt M (2011) Direct-coupling analysis of residue coevolution captures native
contacts across many protein families. *Proc Natl Acad Sci U S A* 108: E1293-1301
Moreira F, Arenas M, Videira A, Pereira F (2023) Evolution of TOP1 and TOP1MT
Topoisomerases in Chordata. *Journal of Molecular Evolution* 91: 192-203
Morris DH, Gostic KM, Pompei S, Bedford T, Łuksza M, Neher RA, Grenfell BT, Lässig M,
McCauley JW (2018) Predictive Modeling of Influenza Shows the Promise of Applied
Evolutionary Biology. *Trends Microbiol* 26: 102-118
Munck C, Gumpert HK, Wallin AI, Wang HH, Sommer MO (2014) Prediction of resistance
development against drug combinations by collateral responses to component drugs.
Science translational medicine 6: 262ra156
Navascues M, Depaulis F, Emerson BC (2010) Combining contemporary and ancient DNA in
population genetic and phylogeographical studies. *Mol Ecol Resour* 10: 760-772
Neher RA, Hallatschek O (2013) Genealogies of rapidly adapting populations. *Proceedings of*
the National Academy of Sciences 110: 437-442
Neher RA, Russell CA, Shraiman BI (2014) Predicting evolution from the shape of
genealogical trees. *Elife* 3
Nickle DC, Heath L, Jensen MA, Gilbert PB, Mullins JI, Kosakovsky Pond SL (2007) HIV-
specific probabilistic models of protein evolution. *PLoS One* 2: e503
Papkou A, Garcia-Pastor L, Escudero JA, Wagner A (2023) A rugged yet easily navigable
fitness landscape. *Science* 382: eadh3860
Parisi G, Echave J (2001) Structural constraints and emergence of sequence patterns in
protein evolution. *Mol Biol Evol* 18: 750-756
Parisi G, Echave J (2005) Generality of the structurally constrained protein evolution model:
assessment on representatives of the four main fold classes. *Gene* 345: 45-53
Pascual-Garcia A, Abia D, Mendez R, Nido GS, Bastolla U (2010) Quantifying the
evolutionary divergence of protein structures: the role of function change and function
conservation. *Proteins* 78: 181-196
Pascual-Garcia A, Arenas M, Bastolla U (2019) The Molecular Clock in the Evolution of
Protein Structures. *Syst Biol* 68: 987-1002

Poon AF, Kosakovsky Pond SL, Richman DD, Frost SD (2007) Mapping protease inhibitor

resistance to human immunodeficiency virus type 1 sequence polymorphisms within

patients. *J Virol* 81: 13598-13607

Rodrigue N, Lartillot N, Bryant D, Philippe H (2005) Site interdependence attributed to

tertiary structure in amino acid sequence evolution. *Gene* 347: 207-217

Rodrigues JV, Bershtein S, Li A, Lozovsky ER, Hartl DL, Shakhnovich EI (2016)

Biophysical principles predict fitness landscapes of drug resistance. *Proceedings of the*

National Academy of Sciences 113: E1470-E1478

Rubin IN, Ispolatov Y, Doebeli M (2023) Adaptive diversification and niche packing on

rugged fitness landscapes. *J Theor Biol* 562: 111421

Ruiz-Gonzalez MX, Fares MA (2013) Coevolution analyses illuminate the dependencies

between amino acid sites in the chaperonin system GroES-L. *BMC Evol Biol* 13: 156

Sali A, Blundell TL (1993) Comparative protein modelling by satisfaction of spatial

restraints. *J Mol Biol* 234: 779-815

Santos-Pereira A, Triunfante V, Araujo PMM, Martins J, Soares H, Poveda E, Souto B,

Osorio NS (2021) Nationwide Study of Drug Resistance Mutations in HIV-1 Infected

Individuals under Antiretroviral Therapy in Brazil. *Int J Mol Sci* 22

Souto B, Triunfante V, Santos-Pereira A, Martins J, Araujo PMM, Osorio NS (2021)

Evolutionary dynamics of HIV-1 subtype C in Brazil. *Sci Rep* 11: 23060

Spielman SJ (2020) Relative Model Fit Does Not Predict Topological Accuracy in Single-

Gene Protein Phylogenetics. *Molecular Biology and Evolution* 37: 2110-2123

Stackhouse J, Presnell SR, McGeehan GM, Nambiar KP, Benner SA (1990) The ribonuclease

from an extinct bovid ruminant. *FEBS Lett* 262: 104-106

Stadler T (2010) Sampling-through-time in birth–death trees. *Journal of Theoretical Biology*

267: 396-404

Stadler T (2011) Simulating trees with a fixed number of extant species. *Syst Biol* 60: 676-684

Thadani NN, Gurev S, Notin P, Youssef N, Rollins NJ, Ritter D, Sander C, Gal Y, Marks DS

(2023) Learning from prepandemic data to forecast viral escape. *Nature* 622: 818-825

Thornton JW, Need E, Crews D (2003) Resurrecting the ancestral steroid receptor: ancient

origin of estrogen signaling. *Science* 301: 1714-1717

Ugalde JA, Chang BSW, Matz MV (2004) Evolution of Coral Pigments Recreated. *Science*

305: 1433-1433

Van Cleve J, Weissman DB (2015) Measuring ruggedness in fitness landscapes. *Proceedings*
of the National Academy of Sciences 112: 7345-7346

Watabe T, Kishino H (2010) Structural considerations in the fitness landscape of a virus. *Mol*
Biol Evol 27: 1782-1791

Wiuf C, Posada D (2003) A coalescent model of recombination hotspots. *Genetics* 164: 407-
417

Wortel MT, Agashe D, Bailey SF, Bank C, Bisschop K, Blankers T, Cairns J, Colizzi ES,
Cusseddu D, Desai MM *et al* (2023) Towards evolutionary predictions: Current promises
and challenges. *Evolutionary Applications* 16: 3-21

Wright S (1931) Evolution in Mendelian populations. *Genetics* 16: 97-159

Wylie CS, Shakhnovich EI (2011) A biophysical protein folding model accounts for most
mutational fitness effects in viruses. *Proceedings of the National Academy of Sciences*
108: 9916-9921

Yang Z (1996) Among-site rate variation and its impact on phylogenetic analysis. *Trends*
Ecol Evol 11: 367-372

Yang Z (2006) *Computational Molecular Evolution*. Oxford University Press, Oxford,
England.

Yang Z, Goldman N, Friday A (1994) Comparison of models for nucleotide substitution used
in maximum-likelihood phylogenetic estimation. *Molecular Biology and Evolution* 11:
316-324

Yoshida K, Hata K, Kawakami K, Hiradate S, Osawa T, Kachi N (2023) Predicting
ecosystem changes by a new model of ecosystem evolution. *Scientific Reports* 13: 15353

Zeldovich KB, Chen P, Shakhnovich EI (2007) Protein stability imposes limits on organism
complexity and speed of molecular evolution. *Proceedings of the National Academy of*
Sciences 104: 16152-16157

Zhang J, Nei M (1997) Accuracies of ancestral amino acid sequences inferred by the
parsimony, likelihood, and distance methods. *J Mol Evol* 44 Suppl 1: S139-146

Tables

Table 1. Comparison of real and predicted sequences of the HIV-1 MA protein considering predictions based on the SCS and

neutral models. For the data simulated under the SCS and neutral models, the table shows the Grantham distance between the amino acids

that changed during the real and predicted evolutionary trajectories and, the Kullback-Leibler (KL) divergence between the real and

predicted multiple sequence alignments. Next, it shows the folding stability (ΔG) of the real protein variants at times *T1* and *T31*, and the

folding stability of the predicted protein variants at time *T31*. The error corresponds to the 95% confidence interval from the mean of

predictions of folding stability.

	Grantham distance	KL divergence	ΔG of the real variant at *T1* (kcal/mol)	ΔG of the real variants at *T31* (kcal/mol)	ΔG of the predicted variants at *T31* (kcal/mol)	$\Delta\Delta G$ (kcal/mol) at *T31* (predicted – real variants)
SCS model	5%	6%	-9.72	-10.34 ± 0.14	-9.96 ± 0.02	0.38
Neutral model	5%	6%	-9.72	-10.34 ± 0.14	-9.21 ± 0.07	1.14

Figure captions

Figure 1. Illustrative example of forward in time simulation of protein evolution

integrating a birth-death population evolutionary process with fitness from the protein

folding stability and the modeling of protein evolution with a structurally constrained

substitution model. Given a protein variant assigned to a node at time t (blue node), its

fitness is calculated considering its protein folding stability. Then, the fitness is used to

determine the birth and death rates for that variant, which provide the time to the next birth or

death event (horizontal dashed line) that corresponds to the forward-in-time branch length.

Next, the variant is evolved forward in time toward each descendant, upon the previously

determined branch length, under a SCS model of protein evolution. The process is repeated,

forward in time, starting at each new variant. If a death event occurs, the variant of the extinct

node (pink node) is obtained but it does have descendants. The process finishes when a

particular sample size or simulation time is reached (i.e., $t+n$).

Figure 2. Prediction error of SARS-CoV-2 Mpro and PLpro evolution regarding

physicochemical properties of the amino acid changes accumulated during the

evolutionary trajectories and protein folding stability. (A) Grantham distance calculated

from the amino acid changes that occurred during the real and predicted evolutionary

trajectories based on SCS and neutral models of protein evolution. (B) Variation of protein

folding stability ($\Delta\Delta G$) between real and predicted protein variants based on SCS and neutral

models of protein evolution. Notice that positive $\Delta\Delta G$ indicates that the real protein variants

are more stable than the predicted protein variants and *vice versa*. Error bars correspond to the

95% confidence interval of the mean of prediction error from 100 multiple sequence

alignments simulated for the corresponding population and time.

Figure 3. Prediction error of HIV-1 PR evolution at diverse populations regarding physicochemical properties of the amino acid changes accumulated during the evolutionary trajectories and protein folding stability. (A) For each viral population (patient, represented with a particular color) and time, Grantham distance calculated from the amino acid changes that occurred during the real and predicted evolutionary trajectories. For each population, the mean of distances obtained over time is shown on the right. (B) Relationship between Grantham distances and accumulated number of substitution events (R^2 = 0.0001, which indicates a lack of correlation between these parameters). (C) Variation of protein folding stability ($\Delta\Delta G$) between real and predicted protein variants at each viral population and time. For each population, the mean of distances obtained over time is shown on the right. Notice that positive $\Delta\Delta G$ indicates that the real protein variants are more stable than the predicted protein variants and *vice versa*. Error bars correspond to the 95% confidence interval of the mean of prediction error from 100 multiple sequence alignments simulated for the corresponding viral population and time.

842   *Figure 1.*

844   *Figure 2.*

846   *Figure 3.*