Forecasting protein evolution by integrating birth-death population models with structurally constrained substitution models

David Ferreiro[1,2], Luis Daniel González-Vázquez[1,2], Ana Prado-Comesaña[1] and Miguel Arenas[1,2,*]

[1]CINBIO, Universidade de Vigo, 36310 Vigo, Spain.

[2]Department of Biochemistry, Genetics and Immunology, Universidade de Vigo, 36310 Vigo, Spain.

Email addresses:

    DF: david.ferreiro.garcia@uvigo.es

    LDGV: luisdaniel.gonzalez@uvigo.gal

    APC: ana.maria.prado.comesana@uvigo.gal

    MA: marenas@uvigo.es

* Corresponding author:
Miguel Arenas
Department of Biochemistry, Genetics and Immunology
University of Vigo
36310 Vigo, Spain
Email: marenas@uvigo.es
Phone: +34 986 130047

Running head: Forecasting protein evolution

Keywords: Molecular evolution, Forecasting evolution, Birth-death process, Substitution model, Protein folding stability, Phylogenetics, Molecular adaptation

Abstract

Evolutionary studies in population genetics and ecology were mainly focused on predicting

and understanding past evolutionary events. Recently, however, a growing trend explores the

prediction of evolutionary trajectories toward the future promoted by its wide variety of

applications. In this context, we introduce a forecasting protein evolution method that

integrates birth-death population models with substitution models that consider selection on

protein folding stability. In contrast to traditional population genetics methods that usually

make the unrealistic assumption of simulating molecular evolution separately from the

evolutionary history, the present method combines both processes to simultaneously model

forward-in-time birth-death evolutionary trajectories and protein evolution under structurally

constrained substitution models that outperformed traditional empirical substitution models.

We implemented the method into a freely available computer framework. We evaluated the

accuracy of the predictions with several monitored viral proteins of broad interest. Overall,

the method showed acceptable errors in predicting the folding stability of the forecasted

protein variants but, expectedly, the errors were larger in the prediction of the corresponding

sequences. We conclude that forecasting protein evolution is feasible in certain evolutionary

scenarios and provide suggestions to enhance its accuracy by improving the underlying

models of evolution.

Introduction

Molecular evolution is traditionally investigated through inferences about past evolutionary events, such as phylogenetic tree and ancestral sequence reconstructions, and predictions about the future were considered as inaccessible for a long time because they can be affected by complex processes such as environmental change. Nevertheless, a variety of biological systems display a Darwinian evolutionary process where selection operates toward a limited set of adapted variants. These variants, and in extension the evolutionary trajectories to reach them, would be positively selected and could present a certain degree of predictability (Lässig *et al*, 2017; Wortel *et al*, 2023). The progress made in developing more accurate models of evolution (Arenas, 2015b) and the benefits from predicting the outcome of evolution (i.e., to understand the course of evolution or to prepare for the future (Wortel *et al.*, 2023)) motivated a variety of investigations on forecasting evolution in diverse fields including medicine, agriculture, biotechnology, and conservation biology, among others (e.g., Barton *et al*, 2016; Bull & Molineux, 2008; de Visser *et al*, 2018; Diaz-Uriarte & Vasallo, 2019; Fischer *et al*, 2015; Gerrish & Sniegowski, 2012; Lassig & Luksza, 2014; Lind *et al*, 2019; Luksza & Lassig, 2014; Morris *et al*, 2018; Munck *et al*, 2014; Neher *et al*, 2014; Wortel *et al.*, 2023). Unfortunately, forecasting evolution not always is achievable. Under neutral evolution, all the molecular variants are equally likely to be present in the population, showing lack of repeatability and disallowing accurate prediction of future variants. Thus, forecasting evolution requires a system with measurable selection pressures, and where certain positively selected variants could produce more descendants than other variants and expand in the population (Desai & Fisher, 2007; Goyal *et al*, 2012; Neher & Hallatschek, 2013; Neher *et al.*, 2014). Actually, a rougher fitness landscape resulting from selection can lead to greater accuracy in evolutionary predictions (Papkou *et al*, 2023; Rubin *et al*, 2023; Van Cleve &

Weissman, 2015). Overall, an evolutionary process could be predictable to some extent

(prediction errors are inevitable with any method and in any evolutionary scenario) depending

on the strength of selection driving evolution and the heterogeneity in fitness among different

variants.

Here, we focus on forecasting protein evolution because it involves molecular evolutionary

processes driven by selection pressures and where the fitness of each variant can be

parameterized and predicted (Carneiro & Hartl, 2010; Gilson *et al*, 2017). Traditionally,

evolutionary histories and ancestral sequences of proteins are inferred using probabilistic

methods based on advanced substitution models of protein evolution (e.g., Arenas, 2015b;

Arenas & Bastolla, 2020; Ferreiro *et al*, 2022; Malcom *et al*, 1990; Moreira *et al*, 2023;

Stackhouse *et al*, 1990; Thornton *et al*, 2003; Ugalde *et al*, 2004). The accuracy of these

inferences is affected by the accuracy of the applied substitution model, where substitution

models that better fit with the study data usually produce more accurate phylogenetic trees

and ancestral sequences (Arenas & Bastolla, 2020; Del Amparo & Arenas, 2022a, 2023;

Lemmon & Moriarty, 2004). These findings suggest that accurate substitution models of

evolution are also convenient for forecasting protein evolution. In this regard, a variety of

studies showed that structurally constrained substitution (SCS) models of protein evolution

provide more accurate evolutionary inferences than the traditional empirical substitution

models of protein evolution, in terms of phylogenetic likelihood, distribution of amino acid

frequencies among protein sites, rates of molecular evolution and folding stability of

reconstructed proteins, among other aspects (e.g., Arenas & Bastolla, 2020; Arenas *et al*,

2015; Bastolla *et al*, 2006; Bordner & Mittelmann, 2013; Del Amparo *et al*, 2023; Echave &

Wilke, 2017; Ferreiro *et al*, 2024a; Ferreiro *et al*, 2024b; Fornasari *et al*, 2002; Parisi &

Echave, 2001; Pascual-Garcia *et al*, 2019; Rodrigue *et al*, 2005), although SCS models

usually demand more computational resources than substitution models that only include

information from the protein sequence. Notice that the protein structure provides information

about the location and molecular interactions of amino acids at different protein sites, which

could be far from each other in the sequence but close in the three-dimensional structure and

interact affecting their evolution (Morcos *et al*, 2011; Ruiz-Gonzalez & Fares, 2013). Indeed,

selection from the protein folding stability is relevant in the evolution of multiple proteins,

including those in microbial and viral systems (e.g., Ferreiro *et al.*, 2022; Gong *et al*, 2013;

Jacquier *et al*, 2013; Rodrigues *et al*, 2016; Wylie & Shakhnovich, 2011; Zeldovich *et al*,

2007). Therefore, we believe that it should be taken into account for forecasting protein

evolution in such systems.

Predictions about future evolutionary events can be performed with simulation-based methods

(e.g., Eccleston *et al*, 2023; Neher *et al.*, 2014; Yoshida *et al*, 2023). In order to simulate

molecular evolution, traditional population genetics methods apply two separate steps

(Arenas, 2012; Hoban *et al*, 2012). First, the simulation of the evolutionary history (i.e., a

phylogenetic tree) using approaches such as the coalescent and birth-death population

processes (Gernhard, 2008; Hudson, 1990; Kingman, 1982; Stadler, 2010). Afterward, the

forward-in-time simulation of molecular evolution is performed, from the root node to the tip

nodes, upon the previously simulated evolutionary history (Yang, 2006). This methodology

was implemented into a variety of population genetics frameworks that simulate molecular

evolution (Arenas, 2012; Hoban *et al.*, 2012). However, for technical and computational

simplicity, it assumes that the simulation of the evolutionary history is independent from the

simulation of molecular evolution, which can produce biological incoherences (i.e., the

evolutionary history is usually simulated under neutral evolution while molecular evolution is

usually simulated with substitution models that consider selection). To enhance the realism of

this modeling, here we merged both processes into a single one where evolutionary history

influences molecular evolution and *vice versa*. In particular, we adopted a birth-death

population genetics method to simulate the forward-in-time evolutionary history (already used

for forecasting evolution (Lassig & Luksza, 2014; Neher *et al.*, 2014)), taking into account the

fitness of the molecular variant (through evolutionary constraints from the protein folding

stability (Bastolla & Demetrius, 2005; Gong *et al.*, 2013; Liberles *et al*, 2012; Zeldovich *et*

al., 2007)) at the corresponding node, to determine its subsequent birth or death event, and we

integrated this process with SCS models to model protein evolution along the derived

phylogenetic branches. The method is detailed below, and we implemented it into a new

version of our computer framework *ProteinEvolver* (Arenas *et al*, 2013), which is freely

available from https://github.com/MiguelArenas/proteinevolver. *ProteinEvolver2* includes

detailed documentation and a variety of ready-to-use examples. Next, considering the

potential applications of forecasting evolution to design vaccines and therapies against

pathogens, we evaluated and applied the method to forecasting protein evolution in several

real protein data of viruses monitored over time.

Materials and Methods

A method for forecasting protein evolution by combining birth-death population

genetics with structurally constrained substitution models of protein evolution

Following previous methods for forecasting evolution based on simulations (Lassig &

Luksza, 2014; Neher *et al.*, 2014), we developed a method to simulate the forward-in-time

evolutionary history of a protein sample with a birth-death process that considers the fitness

of the protein variant (based on folding stability) at every temporal node. The method derives

the birth and death rates for a protein variant based on the fitness of that variant, where a high

fitness results in a high birth rate and a low death rate, which can lead to a large number of

descendants, and the opposite leading to a few or none (extinction) descendants. Thus, the

fitness of the molecular variant at every node drives its corresponding forward in time birth-

death evolutionary history. The details of this simulation process are outlined below.

First, similarly to common simulators of molecular evolution (Arenas, 2012; Hoban *et al.*,

2012; Yang, 2006), a given protein sequence and structure (hereafter, protein variant) is

assigned to the root node. The fitness (*f*) of the protein variant (*A*) is calculated from its

folding stability (free energy, *ΔG*) following the Boltzmann distribution (Goldstein, 2013) (I,

which takes values from 0 to 1),

$$f(A) = \frac{1}{1 + e^{\Delta G / kT}} \qquad \text{(I)}$$

Protein folding stability constrains protein evolution and is commonly used to obtain protein

fitness (Bastolla *et al*, 2007; Goldstein, 2013; Liberles *et al.*, 2012; Lobkovsky *et al*, 2010;

Mendez *et al*, 2010; Sella & Hirsh, 2005; Zeldovich *et al.*, 2007). The user can alternatively

choose whether the fitness of the modeled protein variant is determined solely by its folding

stability or by its similarity to the stability of a real protein variant (i.e., a protein structure

from the Protein Data Bank, PDB). We believe the latter can be more realistic, as in nature,

high folding stability does not necessarily indicate high fitness, but a stability that closely

resembles that of a real protein may suggest high fitness since the real stability is the result of

a selection process (which also incorporates negative design). If the fitness is derived from

only the folding stability of the protein variant, the birth rate (*b*) is considered equal to the

fitness. Alternatively, if the fitness is determined based on the similarity in folding stability

between the modeled variant and a real variant, the birth rate is assumed to be 1 minus the

root mean square deviation (RMSD, which offers advantages such as minimizing the

influence of small deviations while amplifying larger differences, thereby enhancing the

detection of remarkable molecular changes) in folding stability. Notice that the smaller this

difference, the higher the birth rate. In both cases, the death rate (d) is considered as *1-b* to

allow a constant global (birth-death) rate. In this model, the fitness influences reproductive

success, where protein variants with higher fitness have higher birth rates leading to more

birth events, while those with lower fitness have higher death rates leading to more extinction

events. This parameterization is meaningful in the context of protein evolution because the

fitness of a protein variant can affect its survival (birth or extinction) without necessarily

altering its rate of evolution. Although a higher growth rate can sometimes correlate with

higher fitness, a variant with high fitness does not necessarily accumulate substitutions more

rapidly.

Additionally, we incorporated another birth-death model that follows the proposal by Neher et

al (2014), in which the death rate is fixed at 1 and the birth rate is modeled as *1+fitness*. In

this model, fitness not only affects reproductive success but also influences the global birth-

death rate, which can vary among lineages.

The birth-death process is simulated forward in time deciding whether every next event is a

birth or a death event (Harmon, 2019; Stadler, 2010; Stadler, 2011) according to the fitness of

the corresponding molecular variant, and it ends when a user-specified criterion, such as a

particular sample size (considering or ignoring extinction nodes) or a certain evolutionary

time (t_e), is reached. Starting from an "active" node (i.e., the root node) at current time t_c, the

time to the next event (birth to produce two descendants or death to produce the extinction of

the node) can be calculated (details below). In contrast to standard birth-death processes

where birth and death rates are constant over time and among lineages, the present method

considers heterogeneity where each protein variant at a node has specific birth and death rates

according to its corresponding fitness. The method is described below and summarized in

Figure 1.

1. The process starts at the root node, assigning a user-specified protein sequence and

corresponding protein structure (i.e., obtained from the PDB) to that node. In general,

for every protein variant assigned to an active node, the corresponding birth and death

rates are calculated following the indications presented above.

2. Calculation of the time to the next birth or death event. Following common birth-death

methods, the time to the next event t_n is calculated through an exponential distribution

with rate based on the number of active nodes (s) and the sum of the birth and death

rates (Harmon, 2019) (II),

203
$$t_n = e^{(s(b+d))} \qquad \text{(II)}$$

One of the birth-death models that we implemented considers that $b+d=1$ at each

node, allowing variation of the reproductive success among nodes while keeping t_n

consistent among them, according to (Harmon, 2019). In contrast, in the other birth-

death model we implemented, $b+d$ can vary among nodes (Neher *et al.*, 2014),

thereby allowing variation of both reproductive success and t_n among nodes.

3. Evaluate whether the simulation concludes before the next event occurs $(t_c + t_n)$.

4. If it does conclude (i.e., $t_c + t_n$ higher than t_e), the simulation of the evolutionary

history finishes.

5. If it does not conclude, a random active node is selected, and its protein variant is

analyzed to determine its type of next event (birth or death). The probability of a birth

event is $P_b = b/(b + d)$ and the probability of a death event is $P_d = d/(b + d)$. A

random sample from those probabilities is taken to determine the type of evolutionary

event.

6. If a birth event is selected. Two descendant active nodes from the study node are

incorporated with branch lengths t_n, and the study node is then considered inactive.

Next, molecular evolution is simulated from the original node to each descendant node

based on the specified SCS model of protein evolution (Arenas *et al.*, 2013). The

integration of SCS models to evolve protein variants along given branch lengths

followed standards approaches of molecular evolution in population genetics, in which

the branch length and the substitution model inform the number and type of

substitution events, respectively (Arenas, 2012; Carvajal-Rodriguez, 2010; Hoban *et*

al., 2012; Yang, 2006). Thus, the process results in a protein variant for every

descendant node. Finally, the folding stability and subsequent fitness of these

descendant protein variants are calculated.

7. If a death event is selected. Then, the study node is considered inactive.

8. Return to step 1 while the user-specified criterion for ending the birth-death process is

not satisfied and at least one active node exists in the evolutionary history. Otherwise,

the simulation ends.

This process simultaneously simulates, forward in time, evolutionary history and protein

evolution, with protein evolution influencing the evolutionary history through selection from

the folding stability. Indeed, selection can vary among protein variants at their corresponding

nodes of the evolutionary history. The process produces a forward in time birth-death

phylogenetic history, that encompasses nodes that reached the ending time, internal nodes and

nodes that were extinct at some time, along with the protein variant associated with each

node.

The method includes several optional capabilities listed in Table S1 (Supplementary Material)

and in the software documentation, with some summarized below.

• The user can fix the birth and death rates along the evolutionary history or specify that

they are based on the fitness of the corresponding protein variant as described before.

For the latter, the fitness can be based on the folding stability of the analyzed protein

variant or on the similarity in folding stability between the analyzed and real protein

variants. In addition, the global birth-death rate can be constant or vary among

lineages depending on the specified birth-death model.

• The birth-death simulation can finish when any of the following criteria is met: (a) A

specified sample size, including extinction nodes derived from death events. (b) A

specified sample size, excluding extinction nodes. (c) A specified evolutionary time,

measured from the root to a tip node.

• Extinction nodes can either be preserved or removed from the evolutionary history.

• Evaluation of the simulated birth-death evolutionary history in terms of tree balance

using the Colless index (Colless, 1982; Lemant *et al*, 2022).

• Possible variation of the site-specific substitution rate according to user specifications.

• Customizable substitution model featuring site-specific exchangeability matrices

(relative rates of change among amino acids and their frequencies at the equilibrium).

• Implementation of several SCS models and a variety of empirical substitution models

of protein evolution. The implemented SCS models were evaluated in our previous

work (Arenas *et al.*, 2013) and, the implemented empirical substitution models were

properly identified using simulated data with *ProtTest3* (Darriba *et al*, 2011).

The implemented SCS models consider molecular energy functions based on amino acid

contact matrices and configurational entropies per residue in unfolded and misfolded proteins

(Arenas *et al.*, 2013; Bastolla *et al.*, 2007; Mendez *et al.*, 2010). These models incorporate

both positive and negative design strategies. In particular, the evaluation of the target protein

structure while taking into account a database of residue contacts from alternative protein

structures in the PDB, thus considering background genetic information that helps reduce

prediction biases (Minning *et al*, 2013). Technical details about these SCS models are

presented in our previous study (Arenas *et al.*, 2013). Next, SCS models outperformed models

that ignore structural evolutionary constraints in terms of phylogenetic likelihood, among

other properties (Arenas *et al.*, 2013; Arenas *et al.*, 2015; Bordner & Mittelmann, 2013). The

method also implements common empirical substitution models of protein evolution (i.e.,

Blosum62, *CpRev*, *Dayhoff*, *DayhoffDCMUT*, *FLU*, *HIVb*, *HIVw*, *JTT*, *JonesDCMUT*, *LG*,

Mtart, *Mtmam*, *Mtrev24*, *RtRev*, *VT* and *WAG*; Table S1) and the user can specify any

particular exchangeability matrix for all sites or for each site, allowing for heterogeneity of

the substitution process among sites (details in Table S1 and in the software documentation).

In addition, the framework implements heterogeneous substitution rates among sites by the

traditional Gamma distribution (+G) (Yang, 1996) and proportion of invariable sites (+I)

(Fitch & Margoliash, 1967), and also the user can directly alter the substitution rate at each

site for any empirical or SCS model (Table S1). Regarding the evolutionary history, in

addition to the birth-death process presented before, the user can specify a particular

phylogenetic tree or simulate a coalescent evolutionary history (Hudson, 1983; Kingman,

1982) (Table S1). In this regard, we maintained the capabilities of the previous version,

including the coalescent with recombination (Hudson, 1983) [which can be homogeneous or

heterogeneous along the sequence (according to Wiuf & Posada, 2003)], variable population

size over time (growth rate or demographic periods), several migration models [i.e., island

(Hudson, 1998), stepping-stone (Kimura & Weiss, 1964) and continent-island (Wright, 1931)]

with temporal variation of migration rates and convergence of demes or subpopulations,

simulation of haploid or diploid data, and longitudinal sampling (Navascues *et al*, 2010)

(details in Table S1 and in the software documentation). The framework outputs a simulated

multiple sequence alignment with the protein sequences of the internal and tip nodes, as well

as their folding stabilities, and the evolutionary history, among other information (Table S1

and software documentation).

Study data for evaluating the method for forecasting protein evolution

We evaluated the accuracy of the method for forecasting protein evolution using viral proteins

sampled over time (longitudinal sampling). Specifically, we used protein sequences from

previous experiments of virus evolution monitored over time, which contain consensus

molecular data (avoiding rare variants) that belong to different evolutionary time points.

- The matrix (MA) protein of HIV-1, with data obtained from an *in vitro* cell culture
  experiment where samples were collected at different times (Arenas *et al*, 2016). This
  data included 21 MA protein consensus sequences collected at times (population
  passages, *T*) *T1* (initial time and includes an initial sequence) and *T31* (time after 31
  passages and includes 20 sequences), which accumulated 48 amino acid substitutions
  (sequence identity 0.973; Figure S1, Supplementary Material).

- The main protease (Mpro) and papain-like protease (PLpro) proteins of SARS-CoV-2.
  For each protein, the data includes the first sequenced variant (Wuhan) and a sequence
  built with all the substitutions observed in a dataset of 384 genomes of the Omicron
  variant of concern collected from the GISAID database. The resulting Mpro and
  PLpro sequences presented 10 and 22 substitutions (sequence identity 0.967 and
  0.930), respectively.

• The non-structural protein 1 (NS1) of the influenza virus. We retrieved the NS1

sequences from the years 2005, 2015 and 2020 from the Influenza Virus Resource

(Bao *et al*, 2008). Next, we obtained the consensus sequence of the 2005 dataset as

initial time point (*T1*), and the consensus sequences from the 2015 (*T2*) and 2020 (*T3*)

datasets as subsequent time points. The resulting consensus sequences for 2015 and

2020 showed 40 and 37 substitutions (sequence identity 0.802 and 0.817),

respectively.

• The protease (PR) of HIV-1 with data sampled from patients monitored over time,

from 2008 to 2017, available from the Specialized Assistance Services in Sexually

Transmissible Diseases and HIV/AIDS in Brazil (Ferreiro *et al.*, 2022; Santos-Pereira

et al, 2021; Souto *et al*, 2021). These evolutionary scenarios are complex due to the

diverse antiretroviral therapies administrated to the patients (Table S2; Supplementary

Material), which could vary during the studied time periods and that could promote the

fixation of specific mutations (i.e., associated with resistance) (Ferreiro *et al.*, 2022;

Santos-Pereira *et al.*, 2021; Souto *et al.*, 2021). For each viral population (patient), the

data included a consensus sequence for each of the four or five samples collected at

different time points. These consensus sequences exhibited between one and 22 amino

acid substitutions respect to the consensus sequence of the corresponding first sample

(Table S2).

We identified a representative protein structure for each dataset, which was used to predict the

folding stability and to inform the SCS model. In particular, we obtained the protein structures

of the sequences at the initial time (*T1*) from the PDB (Table S3, Supplementary Material).

For the case of the HIV-1 protease, we obtained the protein structure through homology

modelling. We used *SWISS-MODEL* (Arnold *et al*, 2006) to identify the best-fitting templates

of PDB structures (Table S3). Next, we predicted the protein structures by homology

modelling with *Modeller* (Sali & Blundell, 1993) using the protein sequences and

corresponding best-fitting structural templates.

Forward in time prediction of viral protein variants

The evaluation was performed with the previously presented real data, which includes a real

protein variant present at the initial time ($T1$) and subsequent protein variants present at a later

time (Tn). We applied the method for forecasting protein evolution to predict the most likely

protein variants at Tn derived from the real protein variant observed at the initial time ($T1$).

The prediction error was determined by measuring the distance between the real protein

variants and the predicted variants corresponding to time Tn.

Thus, we assigned to the initial node ($T1$) the corresponding real protein variant (including its

sequence and structure), and its evolution was simulated forward in time until it reached the

number of substitutions observed in the real data at time Tn. Thus, we considered the number

of observed substitutions as a measure of evolutionary time to allow proper comparisons

between real and predicted variants. We simulated 100 alignments of protein sequences, each

containing the same number of sequences as the real data at Tn. This included 20 HIV-1 MA

protein sequences, a consensus sequence for the SARS-CoV-2 Mpro, a consensus sequence

for the SARS-CoV-2 PLpro, a consensus sequence of the influenza NS1 protein for each time

point and, a consensus sequence of HIV-1 PR for each viral population. To avoid rare

variants, each sequence of the simulated multiple sequence alignments was obtained as the

consensus of 100 linked simulated sequences.

To investigate the effect of selection on the predictions, we compared the accuracy of

forecasting protein evolution when selection from the protein structure is considered and,

when it is ignored (neutral evolution). If selection is considered, as previously presented, the

probability of birth and death events was based on the fitness of the protein variants, and

protein evolution was modeled using an SCS model (Arenas *et al.*, 2013). In the case of

neutral evolution, all protein variants equally fit and are allowed. Since variants are observed,

we allowed birth events. However, it assumed the absence of death events as no information

independent of fitness is available to support their inclusion, thereby avoiding the imposition

of arbitrary death events based on an arbitrary death rate. Also, to model neutral evolution, we

used an exchangeability matrix with the same relative rates of change to all amino acid pairs.

Accuracy of the predicted protein variants

We assessed the accuracy of the method for forecasting protein evolution by comparing the

predicted and real protein variants present at time *Tn*, both at the protein sequence and

structure levels.

For data containing multiple sequences at time *Tn* (i.e., HIV-1 MA dataset), we calculated the

Kullback-Leibler (KL) divergence, which provides a distance between two multiple sequence

alignments (the real and predicted data) based on the distribution of amino acid frequencies

along the sequences (Equation III, where factors *P* and *Q* denote the distribution of amino

acid frequencies in the real and predicted protein sequences at time *Tn*, respectively, *i* refers to

protein site) (Kullback & Leibler, 1951). This distance was only calculated for data with a set

of sequences at time *Tn* (HIV-1 MA) because a single sequence does not provide site-specific

variability.

$$KL\ (P||Q) = \sum_i P_i \times \log(\frac{P_i}{Q_i}) \qquad \text{(III)}$$

We also compared the real and predicted evolutionary trajectories of protein variants using the

Grantham distance, which measures the differences between amino acids based on their

physicochemical properties (Grantham, 1974). In particular, for both real and predicted

protein variants, we calculated the Grantham distance at each protein site that differs between

the two datasets, considering its evolution from *T1* to the subsequent multiple sequence

alignment at *Tn*. We examined sites that varied over time, thus the general site-specific

Grantham distance *Gi* was calculated as the frequency of each amino acid *f* at site *i* multiplied

by the specific Grantham distance between amino acid *m* at time *Tn* and amino acid *n* at time

T1, normalized with the largest Grantham distance G_{max} to obtain values between 0 and 1

(Equation IV). Next, to compare the real and predicted data, we calculated the site-specific

difference of Grantham distance Gb_i between the real *P* and predicted *Q* protein variants

(Equation V),

$$G_i = \sum_{m=1}^{20} f_m \times \frac{G_{(m,n)}}{G_{max}} \qquad (IV)$$

$$Gb_i = |G_{P,i} - G_{Q,i}| \qquad (V)$$

In addition, we obtained and compared the protein folding stability (ΔG) of the predicted and

real protein variants observed at time *Tn*, using their corresponding protein structures, with

DeltaGREM (Arenas *et al.*, 2015; Minning *et al.*, 2013).

Results

Implementation of the forecasting protein evolution method

We extended the previous version of our framework *ProteinEvolver* (Arenas *et al.*, 2013),

maintaining its previous capabilities (i.e., simulation of protein evolution upon user-specified

phylogenetic trees and upon phylogenetic trees simulated with the coalescent with or without

recombination, migration, demographics and longitudinal sampling, empirical and SCS

models, among others; Table S1), by adding, among others (Table S1), the forward in time

modeling of protein evolution that combines a birth-death process based on the fitness of

every protein variant (folding stability) at each node to determine its birth and death rates, as

well as SCS models of protein evolution. The framework *ProteinEvolver2* is written in C and

distributed with a detailed documentation and a variety of illustrative practical examples. The

framework is freely available from https://github.com/MiguelArenas/proteinevolver.

Evaluation of predictions of HIV-1 MA evolution

Regarding the evolution of the HIV-1 MA protein, the Grantham distance and the KL

divergence between the real variants at time *Tn* and the corresponding predicted variants were

low (around 5% and 6%, respectively; Table 1), and they did not differ comparing predictions

that consider selection on the folding stability (including birth-death models with constant and

variable global birth-death rate among lineages) and predictions that ignore it (Table 1). On

the other hand, we found that the folding stability of the protein variants predicted considering

selection on the folding stability (again, including birth-death models with constant or

variable global birth-death rate among lineages) was closer to the folding stability of the real

protein variants than that of the protein variants predicted under neutral evolution (Table 1).

In particular, the protein variants predicted ignoring selection were less stable than those

predicted considering selection and also less stable than the real protein variants.

Evaluation of predictions of SARS-CoV-2 Mpro and PLpro evolution

The analyses of the SARS-CoV-2 Mpro and PLpro data showed Grantham distances between

the real and predicted sequences around 25% and 36%, respectively (Figure 2A). Again, this

distance was similar when comparing predictions based on models that consider selection on

the protein folding stability (including birth-death models with constant or variable global

birth-death rate among lineages) and a model of neutral evolution. Regarding comparisons

based on the protein folding stability, we found again that the models that consider selection

from the folding stability produce variants closer to the stability of the real protein variants

than the model that ignores selection (Figure 2B). Indeed, protein variants derived from the

models that consider selection were more stable than those derived from the model of neutral

evolution. Next, we did not find statistically significant differences in sequence similarity or

folding stability between variants predicted under birth-death models with constant or

variable global birth-death rate among lineages (Figure 2).

Evaluation of predictions of influenza NS1 protein evolution

The evolutionary predictions for the influenza NS1 protein varied depending on the model

used. Specifically, at the sequence level and for the two prediction time points studied,

Grantham distances between the real and predicted protein sequences were around 23.5% for

the models that incorporated structural evolutionary constraints, compared to about 25.5% for

the neutral model (Figure 3A). These differences became more pronounced when examining

predictions based on protein folding stability. For both time points, models that included

selection consistently generated protein variants with stability more similar to that of the

observed variants than those predicted by the neutral model (Figure 3B). Indeed, sequences

predicted by the model that accounts for selection were generally more stable than those

predicted under neutral evolution. Again, we found no statistically significant differences in

sequence similarity or folding stability between variants predicted under birth-death models

with constant or variable global birth-death rate among lineages (Figure 3).

Evaluation of predictions of HIV-1 PR evolution

In general, the Grantham distance, which compared the evolutionary trajectories of the real

and predicted protein variants from time $T1$ to later times, varied among viral populations

(patients) (Figure 4A). However, for the majority of these populations, the distance remained

below 30% and exhibited minor fluctuations over time. One particular population exhibited a

notable trend, with the distance increasing from 10% to nearly 60% over time. Considering

that the length of the evolutionary trajectories of the protein can differ among the studied

populations, we explored whether the accumulated number of amino acid substitutions could

affect the accuracy of the predictions, and we found that the number of substitutions varied

among populations and this variability did not correlate with the Grantham distance between

the real and predicted data (R^2=0.0001, Figure 4B). In general, the folding stability of the

predicted protein variants was similar or slightly less stable than that of the real protein

variants [with a difference ranging from 0 to 9 kcal/mol and a mean of 3.1 ± 0.9 (95% CI)

kcal/mol; Figure 4C]. Indeed, the folding stability exhibited small fluctuations, increasing and

decreasing, over time.

Discussion

While reconstructing evolutionary histories and ancestral sequences, among other inferences

about the past, was popular in the field, predictions about evolution toward the future were

traditionally ignored due to potential high prediction errors. However, in the last decade,

forecasting evolution has gained attention because of its variety of applications and

advancements in models of evolution (see the reviews Lässig *et al.*, 2017; Wortel *et al.*,

2023). Several studies showed that forecasting evolution can be feasible in some systems,

including virus evolution (Luksza & Lassig, 2014; Thadani *et al*, 2023). Here, we also

investigated forecasting evolution in viruses but at the molecular level, considering the

relevance of the substitution process to produce molecular variants that affect the viral fitness

(Arenas, 2015a; Arenas *et al.*, 2016; Bloom & Neher, 2023; Poon *et al*, 2007; Watabe &

Kishino, 2010). We believe that forecasting genome evolution remains challenging due to the

complexity of its evolutionary processes [i.e., including epistatic interactions, chromosomal

rearrangements and heterogeneous substitution patterns among genomic regions (Arbiza *et al*,

2011), among others] that complicate the parameterization and prediction of accurate fitness

landscapes. However, we believe that it could be more feasible in structural proteins because

of their relatively simpler evolutionary processes that usually include selection on folding

stability (Bloom *et al*, 2006; Goldstein, 2011).

To make evolutionary predictions over time, either toward the past or toward the future,

considering a substitution model of molecular evolution can be convenient. Actually, a variety

of traditional studies showed that the substitution model influences the phylogenetic

likelihood generated by probabilistic approaches used for evolutionary inference (Lemmon &

Moriarty, 2004; Yang *et al*, 1994; Zhang & Nei, 1997). To study protein evolution, empirical

substitution models of protein evolution are routinely used because they allow rapid

calculations based on the assumption of site-independent evolution and typically apply the

same exchangeability matrix for all the protein sites, which is highly unrealistic (Echave *et al*,

2016). Besides, a small set of empirical substitution models was developed for modeling the

evolution of the diverse range of viral proteins (Dang *et al*, 2010; Del Amparo & Arenas,

2022b; Dimmic *et al*, 2002; Le & Vinh, 2020; Nickle *et al*, 2007). As an alternative, some

substitution models that consider evolutionary constraints on the protein structure

incorporated site-dependent evolution, which allows a more accurate modeling of protein

evolution compared to the empirical substitution models (Arenas *et al.*, 2013; Arenas *et al.*,

2015; Bordner & Mittelmann, 2013; Ferreiro *et al.*, 2024a; Parisi & Echave, 2005).

According to previous methods for forecasting evolution based on computer simulations

(Lassig & Luksza, 2014; Neher *et al.*, 2014), we also adopted the birth-death population

genetics process to allow a forward-in-time evolutionary process where the birth and death

rates can differ among nodes (Stadler, 2010; Stadler, 2011). Traditional population genetics

methods to simulate the evolution of molecular data implement a two-step simulation process,

where, first, the evolutionary history is simulated [i.e., with the coalescent theory or a

forward-in-time simulation approach (Arenas, 2012; Hoban *et al.*, 2012), often assuming

neutral evolution] and, in a subsequent step, molecular evolution is simulated upon the

previously obtained evolutionary history (Yang, 2006). However, this methodology is

unrealistic because the fitness of the data can affect its evolutionary history (i.e., a variant

with high fitness is likely to have more descendants than a variant with low fitness). Thus, we

designed and implemented a method for forecasting protein evolution that integrates a birth-

death population genetics process (including the modeling of constant and variable global

birth-death rate among lineages) with a SCS model of protein evolution, where the folding

stability of each protein variant is evaluated to predict its future trajectory in terms of both

evolutionary history and molecular evolution. We implemented this forward-in-time

simulation of protein evolution in a new version of our previous framework *ProteinEvolver*

(Arenas *et al.*, 2013), while maintaining its previous capabilities and extending some of them

(i.e., incorporation of site-specific exchangeability matrices and additional substitution

models of protein evolution, among others; see Table S1 and software documentation). As

any other method for forecasting evolution, the present method ignores possible

environmental shifts that are inherently unpredictable and that could affect the accuracy of the

predictions. Next, we evaluated the forward-in-time evolutionary predictions with real data of

HIV-1, SARS-CoV-2 and influenza virus proteins. We determined the prediction errors

between the real and the predicted protein variants by examining dissimilarity in evolutionary

trajectory (Grantham distance based on physicochemical properties among the differing

amino acids during the evolutionary trajectories), sequence divergence (distribution of amino

acid frequencies among sites using the KL divergence), and protein structure (protein folding

stability). Additionally, we analyzed the influence of accounting for selection, based on the

protein folding stability, on the predictions. We also evaluated birth-death models

incorporating either constant or variable global birth-death rates among lineages. Notably, in

the variable-rate model, fitness can influence both reproductive success and the rate of

molecular evolution.

In general, the sequence and evolutionary trajectory dissimilarities between the real and

predicted protein variants were relatively small, with some variations among the study

proteins. For the HIV-1 MA protein, the SARS-CoV-2 Mpro and PLpro, and influenza NS1

protein, the sequence dissimilarity was below 10%, 26%, 36% and 26%, respectively. The

low prediction errors for the HIV-1 MA protein were expected because this dataset was

derived from an *in vitro* cell culture experiment that is not influenced by a variety of external

factors that could affect the predictions in *in vivo* experiments. As a rough reference, in

traditional ancestral sequence reconstruction (ASR) of protein data with high sequence

identity the error was approximately 2% (Arenas & Posada, 2010). Notice that ASR methods

can exhibit low prediction errors due to their statistical evaluation based on a set of original

sequences, rather than relying on a single original sequence used in the case of forecasting

evolution. The contrasting scenario regarding evolutionary complexity was the HIV-1 PR

data, which was sampled from populations (patients) undergoing various antiretroviral

treatments (Ferreiro *et al.*, 2022). There, the sequence dissimilarity varied among viral

populations, although most were below 30%. A viral population exhibited higher

dissimilarities (near 60%) that we believe could be caused by molecular adaptations promoted

by the therapies, although this needs formal investigation.

An unexpected result was that the model of neutral evolution produced sequence

dissimilarities between the real and predicted protein variants that were quite similar to those

obtained with the SCS model (Table 1 and Figures 2 and 3). A few studies indicated that the

substitution model has negligible effects on the reconstructed phylogenetic trees (Abadi *et al*,

2019; Spielman, 2020). Subsequent studies found that the influence of the substitution model

on phylogenetic reconstructions is dependent on the diversity of the data, where data with

high diversity is more sensible to the applied substitution model due to containing more

evolutionary information to be modeled (Del Amparo & Arenas, 2022a, 2023). Considering

that the studied data present overall low diversity [i.e., sequence identity of 0.973, 0.967,

0.930, 0.802 and 0.817 for the HIV MA data, SARS-CoV-2 Mpro data, SARS-CoV-2 PLpro

data, and influenza NS1 protein data (for each prediction time point, T2 and T3), respectively;

it is important to note that, in general, longitudinal data derived from monitored evolutionary

processes usually show a low diversity because involve relatively short evolutionary histories,

among other factors], we believe that the influence of the applied substitution models on the

prediction of the sequences was small because of the small number of modeled substitution

events, as found in phylogenetic reconstructions (Del Amparo & Arenas, 2022a, 2023).

Actually, in the case of the influenza NS1 protein dataset, which had the highest sequence

diversity among the study datasets, the sequences predicted under the SCS models were more

similar to the real sequences than those derived from predictions under neutral evolution.

Overall, the prediction accuracy varied among the studied evolutionary scenarios; as

expected, it was lower in the more complex scenarios. Indeed, datasets with higher sequence

diversity contain more evolutionary signals, which can improve prediction quality.

We also evaluated the prediction error between the real and predicted protein variants

regarding their folding stability, again comparing the predictions made under a model that

considers structural constraints and a model of neutral evolution. In general, the protein

variants predicted under the SCS model presented a folding stability close to the folding

stability of the respective real protein variants, with differences below 1, 2, 7 and 1 kcal/mol

for the HIV MA data, SARS-CoV-2 Mpro data, SARS-CoV-2 PLpro data, and influenza NS1

protein data, respectively. The higher differences (around 9 kcal/mol) were again observed for

the HIV-1 PR data. In contrast to the prediction error based on sequence dissimilarity, the

prediction error based on folding stability varies between predictions obtained under the SCS

model and those obtained under the neutral model. In the studied evolutionary scenarios, the

protein variants predicted under the neutral model were less stable and farther from the

stability of the real protein variants compared to those predicted under the SCS model. These

results were expected because, under SCS models, protein stability can be modeled with

greater accuracy than sequence similarity due to selection for maintaining stability in the

protein structure despite amino acid changes (Arenas & Bastolla, 2020; Illergard *et al*, 2009;

Pascual-Garcia *et al*, 2010). Indeed, previous studies showed that models that ignore

structural constraints often produce proteins with unrealistic folding instability (Arenas &

Bastolla, 2020; Del Amparo *et al.*, 2023), which suggests that accounting for protein folding

stability in the modeling of protein evolution is recommended for predicting protein variants

with appropriate structural properties.

Therefore, we found a good accuracy in predicting the real folding stability of forecasted

protein variants, while predicting the exact sequences was more challenging, which was not

surprising considering previous studies. In particular, inferring specific sequences is

considerably challenging even for ancestral molecular reconstruction (Arenas & Bastolla,

2020; Arenas *et al*, 2017). Indeed, observed sequence diversity is much greater than observed

structural diversity (Illergard *et al.*, 2009; Pascual-Garcia *et al.*, 2010), and substitutions

between amino acids with similar physicochemical properties can yield modeled protein

variants with more accurate folding stability, even when the exact amino acid sequences

differ. Further work is demanded in the field of substitution models of molecular evolution.

We also found that datasets with relatively high sequence diversity can improve the accuracy

of the predictions due to containing more evolutionary information. Apart from that,

forecasting the folding stability of future real proteins is an important advance in forecasting

protein evolution, given the essential role of folding stability in protein function (Bloom *et al.*,

2006; Scheiblhofer *et al*, 2017) and its variety of applications.

Variation in the global birth-death rate among lineages showed minor effects on prediction

accuracy, suggesting a limited role in protein evolution modeling. Molecular evolution

parameters, particularly the substitution model, appear to be more critical in this regard.

In the context of protein evolution, substitution models are a critical factor (Arenas *et al.*,

2017; Bordner & Mittelmann, 2013; Echave *et al.*, 2016; Echave & Wilke, 2017; Liberles *et*

al., 2012; Wilke, 2012), and the presented combination with a birth-death model constitutes a

first approximation upon which next studies can build to better understand this evolutionary

system. Next, the present method assumes that the protein structure is maintained over the

studied evolutionary time, which can be generally reasonable for short timescales where the

structure is conserved (Illergard *et al.*, 2009; Pascual-Garcia *et al.*, 2010). Over longer

evolutionary timescales, structural changes may occur and, in such cases, modeling the

evolution of the protein structure would be necessary. To our knowledge, modeling the

evolution of the protein structure remains a challenging task that requires substantial

methodological developments. Recent advances in artificial intelligence, particularly in

protein structure prediction from sequence (Abramson *et al*, 2024; Jumper *et al*, 2021), may

offer promising tools for addressing this challenge. However, we believe that evaluating such

approaches in the context of structural evolution would be difficult, especially given the

limited availability of real data with known evolutionary trajectories involving structural

change. In any case, this is probably an important direction for future research.

We present a method to simulate forward-in-time protein evolution accounting for

evolutionary constraints from the protein structure and a birth-death population process, and

where the evolutionary history is influenced by the protein evolution and *vice versa*. The

method is implemented in the computer framework *ProteinEvolver2*, which is freely

distributed with several practical examples and detailed documentation. We believe that

implementing methods into freely available phylogenetic frameworks is important to facilitate

practical applications, as well as future improvements and evaluations. We applied the

method to forecast protein evolution in some viral proteins. We found that the method

provides acceptable approximations to the real evolution, especially in terms of protein

folding stability, suggesting that combining structural constraints with birth-death population

processes in the modeling of protein evolution is convenient. Still, to advance in methods for

forecasting protein evolution, we believe that further efforts should be made in the field to

improve the modeling of protein evolution, such as the incorporation of site-dependent

evolutionary constraints from the protein activity.

Supplementary Material

The supplementary material is available at the journal online.

Data Availability

The computer framework *ProteinEvolver2* is freely available from

https://github.com/MiguelArenas/proteinevolver. The SARS-CoV-2 data is available from

GISAID database with DOI https://doi.org/10.55876/gis8.250206gt. The real and predicted

protein variants are available from Zenodo repository at the URL

https://doi.org/10.5281/zenodo.15548146.

Acknowledgments

This work was supported by the Project PID2023-151032NB-C22 funded by

MCIU/AEI/10.13039/501100011033 and by FEDER, UE. D.F. was funded by a fellowship

from Xunta de Galicia [ED481A-2020/192]. We thank the "Centro de Supercomputación de

Galicia (CESGA)" for the computer resources. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

Declaration of Interests

The authors declare no competing interests.

References

Abadi S, Azouri D, Pupko T, Mayrose I (2019) Model selection may not be a mandatory step
    for phylogeny reconstruction. *Nat Commun* 10: 934

Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L,
    Ballard AJ, Bambrick J *et al* (2024) Accurate structure prediction of biomolecular
    interactions with AlphaFold 3. *Nature* 630: 493-500

Arbiza L, Patricio M, Dopazo H, Posada D (2011) Genome-wide heterogeneity of nucleotide
substitution model fit. *Genome Biol Evol* 3: 896-908

Arenas M (2012) Simulation of Molecular Data under Diverse Evolutionary Scenarios. *PLoS*
Comput Biol 8: e1002495

Arenas M (2015a) Genetic Consequences of Antiviral Therapy on HIV-1. *Comput Math*
Methods Med 2015: 9

Arenas M (2015b) Trends in substitution models of molecular evolution. *Front Genet* 6: 319

Arenas M, Bastolla U (2020) ProtASR2: Ancestral reconstruction of protein sequences
accounting for folding stability. *Methods Ecol Evol* 11: 248-257

Arenas M, Dos Santos HG, Posada D, Bastolla U (2013) Protein evolution along phylogenetic
histories under structurally constrained substitution models. *Bioinformatics* 29: 3020-
3028

Arenas M, Lorenzo-Redondo R, Lopez-Galindez C (2016) Influence of mutation and
recombination on HIV-1 in vitro fitness recovery. *Mol Phylogenet Evol* 94: 264-270

Arenas M, Posada D (2010) The effect of recombination on the reconstruction of ancestral
sequences. *Genetics* 184: 1133-1139

Arenas M, Sanchez-Cobos A, Bastolla U (2015) Maximum likelihood phylogenetic inference
with selection on protein folding stability. *Mol Biol Evol* 32: 2195-2207

Arenas M, Weber CC, Liberles DA, Bastolla U (2017) ProtASR: An Evolutionary
Framework for Ancestral Protein Reconstruction with Selection on Folding Stability.
Syst Biol 66: 1054-1064

Arnold K, Bordoli L, Kopp J, Schwede T (2006) The SWISS-MODEL workspace: a web-
based environment for protein structure homology modelling. *Bioinformatics* 22: 195-
201

Bao Y, Bolotov P, Dernovoy D, Kiryutin B, Zaslavsky L, Tatusova T, Ostell J, Lipman D
(2008) The influenza virus resource at the National Center for Biotechnology
Information. *J Virol* 82: 596-601

Barton JP, Goonetilleke N, Butler TC, Walker BD, McMichael AJ, Chakraborty AK (2016)
Relative rate and location of intra-host HIV evolution to evade cellular immunity are
predictable. *Nature Communications* 7: 11660

Bastolla U, Demetrius L (2005) Stability constraints and protein evolution: the role of chain
length, composition and disulfide bonds. *Protein Eng Des Sel* 18: 405-415

703 Bastolla U, Porto M, Roman HE, Vendruscolo M (2006) A protein evolution model with

704   independent sites that reproduces site-specific amino acid distributions from the Protein

705   Data Bank. *BMC Evol Biol* 6: 43

706 Bastolla U, Porto M, Roman HE, Vendruscolo M (2007) *Structural approaches to sequence*

707   *evolution*. Springer, Berlin, Heidelberg

708 Bloom JD, Labthavikul ST, Otey CR, Arnold FH (2006) Protein stability promotes

709   evolvability. *Proc Natl Acad Sci U S A* 103: 5869-5874

710 Bloom JD, Neher RA (2023) Fitness effects of mutations to SARS-CoV-2 proteins. *Virus*

711   *Evol* 9: vead055

712 Bordner AJ, Mittelmann HD (2013) A new formulation of protein evolutionary models that

713   account for structural constraints. *Mol Biol Evol* 31: 736-749

714 Bull JJ, Molineux IJ (2008) Predicting evolution from genomics: experimental evolution of

715   bacteriophage T7. *Heredity* 100: 453-463

716 Carneiro M, Hartl DL (2010) Colloquium papers: Adaptive landscapes and protein evolution.

717   *Proc Natl Acad Sci U S A* 107 Suppl 1: 1747-1751

718 Carvajal-Rodriguez A (2010) Simulation of genes and genomes forward in time. *Curr*

719   *Genomics* 11: 58-61

720 Colless DH (1982) Review of Phylogenetics: The Theory and Practice of Phylogenetic

721   Systematics. *Systematic Zoology* 31: 100-104

722 Dang CC, Le QS, Gascuel O, Le VS (2010) FLU, an amino acid substitution model for

723   influenza proteins. *BMC Evol Biol* 10: 99

724 Darriba D, Taboada GL, Doallo R, Posada D (2011) ProtTest 3: fast selection of best-fit

725   models of protein evolution. *Bioinformatics* 27: 1164-1165

726 de Visser J, Elena SF, Fragata I, Matuszewski S (2018) The utility of fitness landscapes and

727   big data for predicting evolution. *Heredity (Edinb)* 121: 401-405

728 Del Amparo R, Arenas M (2022a) Consequences of Substitution Model Selection on Protein

729   Ancestral Sequence Reconstruction. *Mol Biol Evol* 39: msac144

730 Del Amparo R, Arenas M (2022b) HIV Protease and Integrase Empirical Substitution Models

731   of Evolution: Protein-Specific Models Outperform Generalist Models. *Genes* 13

732 Del Amparo R, Arenas M (2023) Influence of substitution model selection on protein

733   phylogenetic tree reconstruction. *Gene* 865: 147336

Del Amparo R, González-Vázquez LD, Rodríguez-Moure L, Bastolla U, Arenas M (2023)
Consequences of Genetic Recombination on Protein Folding Stability. *Journal of*
Molecular Evolution 91: 33-45

Desai MM, Fisher DS (2007) Beneficial Mutation–Selection Balance and the Effect of
Linkage on Positive Selection. *Genetics* 176: 1759-1798

Diaz-Uriarte R, Vasallo C (2019) Every which way? On predicting tumor evolution using
cancer progression models. *PLoS Comput Biol* 15: e1007246

Dimmic MW, Rest JS, Mindell DP, Goldstein RA (2002) rtREV: an amino acid substitution
matrix for inference of retrovirus and reverse transcriptase phylogeny. *J Mol Evol* 55:
65-73

Eccleston RC, Manko E, Campino S, Clark TG, Furnham N (2023) A computational method
for predicting the most likely evolutionary trajectories in the stepwise accumulation of
resistance mutations. *eLife* 12: e84756

Echave J, Spielman SJ, Wilke CO (2016) Causes of evolutionary rate variation among protein
sites. *Nat Rev Genet* 17: 109-121

Echave J, Wilke CO (2017) Biophysical Models of Protein Evolution: Understanding the
Patterns of Evolutionary Sequence Divergence. *Annu Rev Biophys* 46: 85-103

Ferreiro D, Branco C, Arenas M (2024a) Selection among site-dependent structurally
constrained substitution models of protein evolution by approximate Bayesian
computation. *Bioinformatics* 40: btae096

Ferreiro D, Khalil R, Gallego MJ, Osorio NS, Arenas M (2022) The evolution of the HIV-1
protease folding stability. *Virus Evol* 8: veac115

Ferreiro D, Khalil R, Sousa SF, Arenas M (2024b) Substitution Models of Protein Evolution
with Selection on Enzymatic Activity. *Mol Biol Evol* 41: msae026

Fischer A, Vazquez-Garcia I, Mustonen V (2015) The value of monitoring to control evolving
populations. *Proc Natl Acad Sci U S A* 112: 1007-1012

Fitch WM, Margoliash E (1967) A method for estimating the number of invariant amino acid
coding positions in a gene using cytochrome c as a model case. *Biochem Genet* 1: 65-71

Fornasari MS, Parisi G, Echave J (2002) Site-specific amino acid replacement matrices from
structurally constrained protein evolution simulations. *Mol Biol Evol* 19: 352-356

Gernhard T (2008) The conditioned reconstructed process. *J Theor Biol* 253: 769-778

Gerrish PJ, Sniegowski PD (2012) Real time forecasting of near-future evolution. *J R Soc*
Interface 9: 2268-2278

Gilson AI, Marshall-Christensen A, Choi JM, Shakhnovich EI (2017) The Role of
Evolutionary Selection in the Dynamics of Protein Structure Evolution. *Biophys J* 112:
1350-1365

Goldstein RA (2011) The evolution and evolutionary consequences of marginal
thermostability in proteins. *Proteins* 79: 1396-1407

Goldstein RA (2013) Population Size Dependence of Fitness Effect Distribution and
Substitution Rate Probed by Biophysical Model of Protein Thermostability. *Genome*
Biol Evol 5: 1584-1593

Gong LI, Suchard MA, Bloom JD (2013) Stability-mediated epistasis constrains the evolution
of an influenza protein. *eLife* 2: e00631

Goyal S, Balick DJ, Jerison ER, Neher RA, Shraiman BI, Desai MM (2012) Dynamic
Mutation–Selection Balance as an Evolutionary Attractor. *Genetics* 191: 1309-1319

Grantham R (1974) Amino acid difference formula to help explain protein evolution. *Science*
185: 862-864

Harmon LJ (2019) Introduction to birth-death models. In: *Phylogenetic Comparative*
Methods, p. https://lukejharmon.github.io/pcm/chapter10_birthdeath/.

Hoban S, Bertorelle G, Gaggiotti OE (2012) Computer simulations: tools for population and
evolutionary genetics. *Nat Rev Genet* 13: 110-122

Hudson RR (1983) Properties of a neutral allele model with intragenic recombination.
Theoretical Population Biology 23: 183-201

Hudson RR (1990) Gene genealogies and the coalescent process. *Oxford Surveys in*
Evolutionary Biology 7: 1-44

Hudson RR (1998) Island models and the coalescent process. *Mol Ecol* 7: 413-418

Illergard K, Ardell DH, Elofsson A (2009) Structure is three to ten times more conserved than
sequence--a study of structural response in protein cores. *Proteins* 77: 499-508

Jacquier H, Birgy A, Le Nagard H, Mechulam Y, Schmitt E, Glodt J, Bercot B, Petit E,
Poulain J, Barnaud G *et al* (2013) Capturing the mutational landscape of the beta-
lactamase TEM-1. *Proceedings of the National Academy of Sciences* 110: 13067-13072

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K,
Bates R, Žídek A, Potapenko A *et al* (2021) Highly accurate protein structure prediction
with AlphaFold. *Nature* 596: 583-589

Kimura M, Weiss GH (1964) The Stepping Stone Model of Population Structure and the
Decrease of Genetic Correlation with Distance. *Genetics* 49: 561-576

Kingman JFC (1982) The coalescent. *Stochastic Processes and their Applications* 13: 235-
248

Kullback S, Leibler RA (1951) On information and sufficiency. *Ann Math Stat* 22: 79-86

Lassig M, Luksza M (2014) Can we read the future from a tree? *Elife* 3

Lässig M, Mustonen V, Walczak AM (2017) Predicting evolution. *Nature Ecology &*
Evolution 1: 0077

Le TK, Vinh LS (2020) FLAVI: An amino acid substitution model for flaviviruses. *J Mol*
Evol In press

Lemant J, Le Sueur C, Manojlović V, Noble R (2022) Robust, Universal Tree Balance
Indices. *Systematic Biology* 71: 1210-1224

Lemmon AR, Moriarty EC (2004) The importance of proper model assumption in bayesian
phylogenetics. *Syst Biol* 53: 265-277

Liberles DA, Teichmann SA, Bahar I, Bastolla U, Bloom J, Bornberg-Bauer E, Colwell LJ,
de Koning AP, Dokholyan NV, Echave J *et al* (2012) The interface of protein structure,
protein biophysics, and molecular evolution. *Protein Sci* 21: 769-785

Lind PA, Libby E, Herzog J, Rainey PB (2019) Predicting mutational routes to new adaptive
phenotypes. *Elife* 8

Lobkovsky AE, Wolf YI, Koonin EV (2010) Universal distribution of protein evolution rates
as a consequence of protein folding physics. *Proc Natl Acad Sci U S A* 107: 2983-2988

Luksza M, Lassig M (2014) A predictive fitness model for influenza. *Nature* 507: 57-61

Malcom BA, Wilson KP, Matthews BW, Kirsch JF, Wilson AC (1990) Ancestral lysozymes
reconstructed, neutrality tested, and thermostability linked to hydrocarbon packing.
Nature 345: 86-89

Mendez R, Fritsche M, Porto M, Bastolla U (2010) Mutation bias favors protein folding
stability in the evolution of small populations. *PLoS Comput Biol* 6: e1000767

Minning J, Porto M, Bastolla U (2013) Detecting selection for negative design in proteins
through an improved model of the misfolded state. *Proteins* 81: 1102-1112

Morcos F, Pagnani A, Lunt B, Bertolino A, Marks DS, Sander C, Zecchina R, Onuchic JN,
Hwa T, Weigt M (2011) Direct-coupling analysis of residue coevolution captures native
contacts across many protein families. *Proc Natl Acad Sci U S A* 108: E1293-1301

Moreira F, Arenas M, Videira A, Pereira F (2023) Evolution of TOP1 and TOP1MT
Topoisomerases in Chordata. *Journal of Molecular Evolution* 91: 192-203

Morris DH, Gostic KM, Pompei S, Bedford T, Łuksza M, Neher RA, Grenfell BT, Lässig M,
McCauley JW (2018) Predictive Modeling of Influenza Shows the Promise of Applied
Evolutionary Biology. *Trends Microbiol* 26: 102-118

Munck C, Gumpert HK, Wallin AI, Wang HH, Sommer MO (2014) Prediction of resistance
development against drug combinations by collateral responses to component drugs.
Science translational medicine 6: 262ra156

Navascues M, Depaulis F, Emerson BC (2010) Combining contemporary and ancient DNA in
population genetic and phylogeographical studies. *Mol Ecol Resour* 10: 760-772

Neher RA, Hallatschek O (2013) Genealogies of rapidly adapting populations. *Proceedings of
the National Academy of Sciences* 110: 437-442

Neher RA, Russell CA, Shraiman BI (2014) Predicting evolution from the shape of
genealogical trees. *Elife* 3

Nickle DC, Heath L, Jensen MA, Gilbert PB, Mullins JI, Kosakovsky Pond SL (2007) HIV-
specific probabilistic models of protein evolution. *PLoS One* 2: e503

Papkou A, Garcia-Pastor L, Escudero JA, Wagner A (2023) A rugged yet easily navigable
fitness landscape. *Science* 382: eadh3860

Parisi G, Echave J (2001) Structural constraints and emergence of sequence patterns in
protein evolution. *Mol Biol Evol* 18: 750-756

Parisi G, Echave J (2005) Generality of the structurally constrained protein evolution model:
assessment on representatives of the four main fold classes. *Gene* 345: 45-53

Pascual-Garcia A, Abia D, Mendez R, Nido GS, Bastolla U (2010) Quantifying the
evolutionary divergence of protein structures: the role of function change and function
conservation. *Proteins* 78: 181-196

Pascual-Garcia A, Arenas M, Bastolla U (2019) The Molecular Clock in the Evolution of
Protein Structures. *Syst Biol* 68: 987-1002

Poon AF, Kosakovsky Pond SL, Richman DD, Frost SD (2007) Mapping protease inhibitor
resistance to human immunodeficiency virus type 1 sequence polymorphisms within
patients. *J Virol* 81: 13598-13607

Rodrigue N, Lartillot N, Bryant D, Philippe H (2005) Site interdependence attributed to
tertiary structure in amino acid sequence evolution. *Gene* 347: 207-217

Rodrigues JV, Bershtein S, Li A, Lozovsky ER, Hartl DL, Shakhnovich EI (2016)
Biophysical principles predict fitness landscapes of drug resistance. *Proceedings of the*
National Academy of Sciences 113: E1470-E1478

Rubin IN, Ispolatov Y, Doebeli M (2023) Adaptive diversification and niche packing on
rugged fitness landscapes. *J Theor Biol* 562: 111421

Ruiz-Gonzalez MX, Fares MA (2013) Coevolution analyses illuminate the dependencies
between amino acid sites in the chaperonin system GroES-L. *BMC Evol Biol* 13: 156

Sali A, Blundell TL (1993) Comparative protein modelling by satisfaction of spatial
restraints. *J Mol Biol* 234: 779-815

Santos-Pereira A, Triunfante V, Araujo PMM, Martins J, Soares H, Poveda E, Souto B,
Osorio NS (2021) Nationwide Study of Drug Resistance Mutations in HIV-1 Infected
Individuals under Antiretroviral Therapy in Brazil. *Int J Mol Sci* 22

Scheiblhofer S, Laimer J, Machado Y, Weiss R, Thalhamer J (2017) Influence of protein fold
stability on immunogenicity and its implications for vaccine design. *Expert Rev*
Vaccines 16: 479-489

Sella G, Hirsh AE (2005) The application of statistical physics to evolutionary biology. *Proc*
Natl Acad Sci USA 102: 9541-9546

Souto B, Triunfante V, Santos-Pereira A, Martins J, Araujo PMM, Osorio NS (2021)
Evolutionary dynamics of HIV-1 subtype C in Brazil. *Sci Rep* 11: 23060

Spielman SJ (2020) Relative Model Fit Does Not Predict Topological Accuracy in Single-
Gene Protein Phylogenetics. *Molecular Biology and Evolution* 37: 2110-2123

Stackhouse J, Presnell SR, McGeehan GM, Nambiar KP, Benner SA (1990) The ribonuclease
from an extinct bovid ruminant. *FEBS Lett* 262: 104-106

Stadler T (2010) Sampling-through-time in birth–death trees. *Journal of Theoretical Biology*
267: 396-404

Stadler T (2011) Simulating trees with a fixed number of extant species. *Syst Biol* 60: 676-684

Thadani NN, Gurev S, Notin P, Youssef N, Rollins NJ, Ritter D, Sander C, Gal Y, Marks DS
(2023) Learning from prepandemic data to forecast viral escape. *Nature* 622: 818-825

Thornton JW, Need E, Crews D (2003) Resurrecting the ancestral steroid receptor: ancient
origin of estrogen signaling. *Science* 301: 1714-1717

Ugalde JA, Chang BSW, Matz MV (2004) Evolution of Coral Pigments Recreated. *Science*
305: 1433-1433

Van Cleve J, Weissman DB (2015) Measuring ruggedness in fitness landscapes. *Proceedings*
of the National Academy of Sciences 112: 7345-7346

Watabe T, Kishino H (2010) Structural considerations in the fitness landscape of a virus. *Mol*
Biol Evol 27: 1782-1791

Wilke CO (2012) Bringing molecules back into molecular evolution. *PLoS Comput Biol* 8:
e1002572

Wiuf C, Posada D (2003) A coalescent model of recombination hotspots. *Genetics* 164: 407-
417

Wortel MT, Agashe D, Bailey SF, Bank C, Bisschop K, Blankers T, Cairns J, Colizzi ES,
Cusseddu D, Desai MM *et al* (2023) Towards evolutionary predictions: Current
promises and challenges. *Evolutionary Applications* 16: 3-21

Wright S (1931) Evolution in Mendelian populations. *Genetics* 16: 97-159

Wylie CS, Shakhnovich EI (2011) A biophysical protein folding model accounts for most
mutational fitness effects in viruses. *Proceedings of the National Academy of Sciences*
108: 9916-9921

Yang Z (1996) Among-site rate variation and its impact on phylogenetic analysis. *Trends*
Ecol Evol 11: 367-372

Yang Z (2006) *Computational Molecular Evolution*. Oxford University Press, Oxford,
England.

Yang Z, Goldman N, Friday A (1994) Comparison of models for nucleotide substitution used
in maximum-likelihood phylogenetic estimation. *Molecular Biology and Evolution* 11:
316-324

Yoshida K, Hata K, Kawakami K, Hiradate S, Osawa T, Kachi N (2023) Predicting
ecosystem changes by a new model of ecosystem evolution. *Scientific Reports* 13:
15353

Zeldovich KB, Chen P, Shakhnovich EI (2007) Protein stability imposes limits on organism

complexity and speed of molecular evolution. *Proceedings of the National Academy of*

Sciences 104: 16152-16157

Zhang J, Nei M (1997) Accuracies of ancestral amino acid sequences inferred by the

parsimony, likelihood, and distance methods. *J Mol Evol* 44 Suppl 1: S139-146

Tables

Table 1. Comparison of real and predicted sequences of the HIV-1 MA protein considering predictions based on the SCS and

neutral models. For the data simulated under the SCS [including birth-death models with constant (SCS) and variable global birth-death

rate among lineages (GlobalBDvar)] and neutral models, the table shows the Grantham distance between the amino acids that changed

during the real and predicted evolutionary trajectories and, the Kullback-Leibler (KL) divergence between the real and predicted multiple

sequence alignments. Next, it shows the folding stability (ΔG) of the real protein variants at times $T1$ and $T31$, and the folding stability of

the predicted protein variants at time $T31$. The error corresponds to the 95% confidence interval from the mean of predictions of folding

stability.

	Grantham distance	KL divergence	ΔG of the real variant at $T1$ (kcal/mol)	ΔG of the real variants at $T31$ (kcal/mol)	ΔG of the predicted variants at $T31$ (kcal/mol)	$\Delta\Delta G$ (kcal/mol) at $T31$ (predicted – real variants)
SCS model	5%	6%	-9.72	-10.34 ± 0.14	-9.96 ± 0.02	0.38
SCS GlobalBDvar model	5%	6%	-9.72	-10.34 ± 0.14	-10.03 ± 0.03	0.31
Neutral model	5%	6%	-9.72	-10.34 ± 0.14	-9.21 ± 0.07	1.14

Figure captions

Figure 1. Illustrative example of forward in time simulation of protein evolution

integrating a birth-death population evolutionary process with fitness from the protein

folding stability and the modeling of protein evolution with a structurally constrained

substitution model. Given a protein variant assigned to a node at time t (blue node), its

fitness is calculated considering its protein folding stability. Then, the fitness is used to

determine the birth and death rates for that variant, which provide the time to the next birth or

death event (horizontal dashed line) that corresponds to the forward-in-time branch length.

Next, the variant is evolved forward in time toward each descendant, upon the previously

determined branch length, under a SCS model of protein evolution. The process is repeated,

forward in time, starting at each new variant. If a death event occurs, the variant of the extinct

node (pink node) is obtained but it does have descendants. The process finishes when a

particular sample size or simulation time is reached (i.e., $t+n$).

Figure 2. Prediction error of SARS-CoV-2 Mpro and PLpro evolution under SCS and

neutral models regarding physicochemical properties of the amino acid changes

accumulated during the evolutionary trajectories and protein folding stability.

Predictions based on data simulated under the SCS [including birth-death models with

constant (SCS) and variable global birth-death rate among lineages (GlobalBDvar)] and neutral

models. (A) Grantham distance calculated from the amino acid changes that occurred during

the real and predicted evolutionary trajectories based on SCS and neutral models of protein

evolution. (B) Variation of protein folding stability ($\Delta\Delta G$) between real and predicted protein

variants based on SCS and neutral models of protein evolution. Notice that positive $\Delta\Delta G$

indicates that the real protein variants are more stable than the predicted protein variants and

vice versa. Error bars correspond to the 95% confidence interval of the mean of prediction

error from 100 multiple sequence alignments simulated for the corresponding population and

time.

Figure 3. Prediction error of influenza NS1 protein evolution under SCS and neutral

models regarding physicochemical properties of the amino acid changes accumulated

during the evolutionary trajectories and protein folding stability. Predictions were

performed for two time points (longitudinal samples T2 and T3). Predictions based on data

simulated under the SCS [including birth-death models with constant (SCS) and variable

global birth-death rate among lineages (GlobalBDvar)] and neutral models. (A) Grantham

distance calculated from the amino acid changes that occurred during the real and predicted

evolutionary trajectories based on SCS and neutral models of protein evolution. (B) Variation

of protein folding stability ($\Delta\Delta G$) between real and predicted protein variants based on SCS

and neutral models of protein evolution. Notice that positive $\Delta\Delta G$ indicates that the real

protein variants are more stable than the predicted protein variants and *vice versa*. Error bars

correspond to the 95% confidence interval of the mean of prediction error from 100 multiple

sequence alignments simulated for the corresponding population and time.

Figure 4. Prediction error of HIV-1 PR evolution at diverse populations regarding

physicochemical properties of the amino acid changes accumulated during the

evolutionary trajectories and protein folding stability. (A) For each viral population

(patient, represented with a particular color) and time, Grantham distance calculated from the

amino acid changes that occurred during the real and predicted evolutionary trajectories. For

each population, the mean of distances obtained over time is shown on the right. (B)

Relationship between Grantham distances and accumulated number of substitution events (R^2

= 0.0001, which indicates a lack of correlation between these parameters). (C) Variation of

protein folding stability ($\Delta\Delta G$) between real and predicted protein variants at each viral

population and time. For each population, the mean of distances obtained over time is shown

on the right. Notice that positive $\Delta\Delta G$ indicates that the real protein variants are more stable

than the predicted protein variants and *vice versa*. Error bars correspond to the 95%

confidence interval of the mean of prediction error from 100 multiple sequence alignments

simulated for the corresponding viral population and time.

Figure 1.

Figure 2.

Figure 3.

Figure 4.