Saccadic Suppression Enhances Saliency of Prey-Like Stimuli in the Optic Tectum

Giulia Soto[1], Carina Thomas[1], Tim C. Hladnik[1,2], Florian A. Dehmelt[1], Ziad M. Hafed[1,3], Aristides B.

Arrenberg[1,*]

[1]Werner Reichardt Centre for Integrative Neuroscience, University of Tübingen, Tübingen, Germany

[2]Graduate Training Centre of Neuroscience, University of Tübingen, Tübingen, Germany

[3]Hertie Institute for Clinical Brain Research, University of Tübingen, Tübingen, Germany

*e-mail: aristides.arrenberg@uni-tuebingen.de

Abstract (200 words)

Saccadic suppression, a reduction in visual sensitivity around the time of a rapid eye movement, is a

robust perceptual phenomenon, extensively studied in primate psychophysics. However, the

mechanisms and purpose of saccadic suppression are still debated in the field. Here, we demonstrate

saccadic suppression in zebrafish, and trace its origins not only in retinal circuits, but also in

downstream visual and motor-related neurons of the optic tectum. Using electrophysiology, we first

established that retinal ganglion cells jumpstart saccadic suppression, in a spatial frequency dependent

manner. Calcium imaging of the optic tectum, combined with 360° visual stimulation and behavioral

tracking, revealed that motor signals enhance peri-saccadic suppression strength up to 0.5 s after the

saccade. Notably, vision-mediated saccadic suppression lasts more than 3 s. Suppression strength

depends on saccade size. Interestingly, we found much weaker saccadic suppression in the optic tectum

for stimuli related to hunting behavior than for behaviorally less relevant global flashes, suggesting a

selectivity of saccadic suppression aimed at increasing the salience of particular types of visual stimuli

after saccades. These findings demonstrate that saccadic suppression in zebrafish integrates visual and

motor signals to optimize sensory processing within neural constraints, providing insights into

evolutionarily conserved visual strategies.

Introduction

Saccadic eye movements are an efficient way of rapidly sampling the visual environment, employed by many species across the animal kingdom. However, by moving the eyes at several hundred degrees per second from one direction of gaze to another, visual flow is greatly disrupted, a disruption that we are blissfully unaware of in our daily perception. From a perceptual standpoint, this can be explained through a phenomenon known as saccadic suppression. Saccadic suppression describes a brief peri-saccadic reduction in perceptual sensitivity and it has been investigated in primates for decades[1–5]. Physiological studies furthermore showed that individual neuronal responses in visual areas are concomitantly reduced peri-saccadically[5], and that responses to stimuli of lower spatial frequency are suppressed more strongly[3,6]. While the benefit of suppressing visual processing during saccades intuitively makes sense, its computation in the brain as well as the reasons underlying variable suppression strengths under different image conditions are still unclear.

Zebrafish (*Danio rerio*) offer the opportunity to investigate neural correlates of saccadic suppression across the visual system in a vertebrate "blueprint" brain without a visual cortex. Larval zebrafish start exhibiting spontaneous saccadic eye movement prior to 4 days post-fertilization (dpf); by 5 dpf, saccades are well coordinated and comparable to adults[7]. Moreover, brainstem control of saccadic eye movements is highly conserved across evolutionarily distant species like zebrafish and primates[8]. Finally, the optic tectum (OT) of zebrafish is homologous to the primate superior colliculus (SC), and both of them are highly sensitive to visual stimulation[9]. Thus, zebrafish are well suited to answer the long-standing debate in the field on the origins and purposes of saccadic suppression.

Considering the frequent occurrence of saccadic eye movements, it might be naïve to think that saccadic suppression simply serves the purpose of suppressing a visual disturbance. Suppression of visual information for fractions of a second with every executed saccade seems simply like too large of a loss

of sensory information, particularly in a system strapped for neural resources, such as the small larval

zebrafish brain. A potential purpose of saccadic suppression could, instead, be as a precise tool for

efficient task-specific encoding. The observation that visual stimulus properties affect suppression

dynamics would be in line with this possibility. For example, saccades across low spatial frequency

surroundings will lead to stronger and longer lasting suppression, compared to high spatial frequency

surroundings[6,10–12]. Similar effects can be observed with luminance contrast dependency[6,13] and

chromatic contrast[14]. Thus, saccadic suppression can be highly specific and depends on image content.

As we show here, the stimulus-dependence of saccadic suppression could even be beneficial to optimize

sensory encoding in behavioral contexts, such as prey capture, where saccadic eye movements play a

vital role in the behavioral sequence[15].

The preponderance of evidence for visual dependencies of saccadic suppression both illuminates and

complicates the search for the circuit origins of saccadic suppression. For example, is saccadic

suppression purely visual? Masking, or the reduced visibility of one brief visual stimulus due to the

presence of a second brief visual stimulus[16], plays a vital role in generating saccadic suppression. In the

1970's, there was already behavioral evidence that visual masking leads to the reduced perception of a

"blur", a visual disturbance caused by the saccadic eye movements itself[17,18]. More recently Idrees and

colleagues showed in isolated mammalian retinae, that visual masking leads to reduced responses to a

probe flash in retinal ganglion cells, following a saccade-like texture displacement[6,19]. However, saccadic

motor signals still likely play a role. For example, it is known from psychophysics studies that the peri-

saccadic window of suppression was significantly shortened[6] when participants performed real saccades

instead of watching a simulated corresponding visual stimulus (a texture displacement). Corollary

discharge signals originating from motor commands can act as inhibitory signals for visual neurons

during rapid eye movements[4,20]. Recent work by Ali and colleagues has shown that for locomotion, a

corollary discharge signal can in fact be found in the zebrafish OT, the sensory structure responsible for

processing most visual inputs in the zebrafish brain[21]. In primates, behavioral and neuronal evidence

suggest the presence of a saccade-related corollary discharge signal in parts of the visual system[22–25].

It is evident that both visual and motor-related signals shape saccadic suppression properties. However,

how the two act together on the individual neuronal level to modulate visual responses is poorly

understood. Here, we show in larval zebrafish how visual responses of tectal neurons are modulated by

sensory and motor-related signals during peri-saccadic suppression, by imaging large populations of

OT neurons through calcium imaging, while simultaneously using 360° whole-field visual stimulation

and behavioral tracking in tethered animals. We find that both visual and motor related processes shape

saccadic suppression differentially in the OT. The optic tectum receives purely visually suppressed

signals from the retina and further modulates them. Motor-related signals in the OT enhance saccadic

suppression. Furthermore, stimuli related to hunting behavior appear more salient, because they are

less suppressed than behaviorally irrelevant global flashes, indicating optimized signal processing.

Results

Purely visual saccadic suppression exists in the zebrafish retina

To enable our investigation of peri-saccadic stimulus encoding, we first need to demonstrate that

saccadic suppression does exist in zebrafish. We started with recording in the retina, because this is the

location where saccadic suppression first occurs in the mammalian visual system[6]. Thus, we isolated

adult zebrafish retinae and performed multi-electrode array (MEA) recordings of retinal ganglion cells

(RGCs) during visual stimulation (Fig. 1a). In order to simulate the visual transient of a saccade, retinae

Fig. 1. Visual saccadic suppression in zebrafish retinae. a. RGC activity of adult zebrafish retinal explants was recorded on low-density multi-electrode array (LD MEA) with simultaneous visual stimulation with DLP projector from below. **b.** Textured stimulus was rapidly displaced in a saccade-like manner (blue arrow). After a variable delay period (d), a whole-field dark or bright flash was shown. **c.** Average spike rate (black line) to texture displacement of example RGC across multiple trials (blue lines). Blue bar indicates time span of texture displacement. **d.** Spike count for individual trials (upper panel) and average spike rate across trials (middle panel) for the same example RGC as in **c,** when texture displacement was paired with a whole-field flash after various delay periods. Isolated flash responses (lower panel) were acquired by subtracting "saccade" activity (blue dashed line) from flash activity (solid green lines). Green bars indicate time points of flash presentation; gray bar indicates time span of texture displacement. **e.** Percent suppression (mean ± SEM) of visual responses to whole-field flash around the time of a saccade-like texture displacement (gray bar) across all RGCs recorded. RGC activity was significantly suppressed during and up to 250 ms after texture displacement onset (# p < 0.0001, two-tailed Wilcoxon signed-rank test). Suppression was significantly stronger for coarse textures, than fine (solid vs. dashed line), where indicated by asterisk (* p < 0.05, two-tailed Wilcoxon signed-rank test, Bonferroni corrected for n = 18 conditions (6 delays vs. baseline for coarse texture, 6 delays vs. baseline for fine texture, 6 coarse vs. fine texture at each delay time)). *N = 87 RGCs from 10 animals.*

were shown a coarse or fine texture which was rapidly displaced in a saccade-like manner (Fig 1b, left).

Texture displacements generated a visual response in RGCs as can be seen in Fig. 1c. To test if saccade-

like texture displacements could suppress responses of a secondary visual stimulus, at various time

points in close temporal proximity to the texture displacement, a 17 ms long dark or bright whole-field

flash was shown (Fig. 1b, right, Fig. 1d). Flash responses, when shown in temporal isolation ("Baseline")

were robust and in most cases much larger than texture displacement responses. In order to isolate

flash-associated responses around the time of the texture displacement, texture displacement-associated

responses were subtracted from flash responses as shown in Fig. 1d (second and third panel). After

removal of this "saccade"-associated response it can be seen that flash responses are significantly

suppressed around the time of the texture displacement (Fig, 1d, third panel). When averaging all RGCs

recorded (n=87), it became apparent that suppression is strongest at 17 and 50 ms after texture

displacement onset and that responses return to baseline levels after ~300 ms (Fig. 1e). This is a similar

time course to that described in mouse and pig retinae[6]. Overall suppression strength is, however,

weaker in zebrafish, with a maximal suppression of 15.8 ± 3.0% (mean ± SEM), whereas mammalian

retinal saccadic suppression can be as strong as ~30% (see Ref. 19, Fig. 3). Furthermore, when

comparing the suppression induced by a coarse texture displacement versus a fine texture displacement,

coarse will generally lead to stronger and longer lasting suppression, than fine, reproducing spatial

frequency dependencies of saccadic suppression in primates[3,6,10].

To ensure that stimulus parameters were well matched with the RGC receptive field properties of

zebrafish, we additionally characterized receptive field sizes, response latencies and response polarities

of individual RGCs (Fig. 2). Adult zebrafish RGCs had receptive field sizes of 21.1 ± 6.6° (mean ± SD, n

= 87) (Fig. 2a). The coarse and fine textures used in this experiment had blob sizes of approximately

14.3° and 3.6° respectively (Methods), falling within the range resolvable by RGCs. With whole-field

Fig. 2. **Receptive field properties of zebrafish RGCs. a.** Receptive field sizes and response latencies were characterized using a binary checkerboard flicker with a checker size of 55 x 55 μm. Orange ellipse indicates receptive filed size and orange line indicates response latency of an example RGCs, both determined as described in Methods. Group data (right panels) shows receptive filed sizes and response latencies of all RGCs recorded, error bars indicating mean ± SD. *N = 87 RGCs*. **b.** Whole-field contrast steps, each lasting 2 seconds were used to determine preferred polarity of RGCs. A majority of RGCs preferred light-to-dark (OFF) contrast steps over dark-to-light (ON) contrast steps. **c.** Spike count for individual trials and mean firing rate across trials during whole-field contrast steps for one example OFF RGC (upper panels) and one example ON RGC (lower panels).

contrast steps, preferred polarity of RGC responses was determined (Fig. 2b, c). Most RCGs were

characterized as OFF cells and preferred a bright-to-dark contrast step (72.4%, 63/87 RGCs). Fewer

RGCs were characterized as ON cells (27.6%, 24/87 RGCs). For the texture displacement experiment,

RGCs were only analyzed for their preferred flash polarity; ON cells for bright flashes and OFF cells for

dark flashes.

151 In summary, we showed with visual stimuli adapted to the visual acuity of zebrafish, that zebrafish

152 retinae modulate visual responses around the time of a saccade-like texture displacements with similar

153 temporal dynamics as mammalian retinae. While suppression strength is reduced, compared to

154 mammals, spatial frequency dependencies remain intact.

156 Tectal neurons respond to dark flash stimuli in a 360° surround arena

The optic tectum is the major retinorecipient area in the zebrafish brain[9]. To investigate how the above-

described retina-modulated responses are further processed in higher visual processing areas, we used

two-photon calcium imaging to image large populations of tectal neurons in larval zebrafish. Due to the

extremely large visual field of zebrafish[26,27], it is vital to provide true whole-field visual stimulation, when

studying saccadic suppression. Our recently established visual stimulation setup allows for high-

resolution 360° whole-field visual stimulation during calcium imaging (Fig. 3a). This enables us to

investigate the consequences of saccade-like texture displacements, analog to the retinal experiments at

the next step of visual processing. Tethered animals were again presented with global coarse and fine

textures. Textures were displaced horizontally in a saccade-like manner and at various delay times a

probe flash was shown. Probe flash parameters were slightly adjusted, to optimize for calcium imaging

in the OT (Fig. 3b). The duration of the flash was extended to 500 ms to allow for the relatively slow

calcium signal to integrate and only dark flashes were presented, since our own observations showed

that most OT neurons preferred dark stimuli over bright. Calcium recordings were registered and

segmented into individual ROIs, each representing one neuron (Methods). For each ROI, $\Delta F/F$ was

computed and regression-based analysis revealed many ROIs (14.9%, 12,397/83,104 ROIs) showing

significant responses to the flash stimulus (Fig. 3c). These responses are truly stimulus-associated, as

comparison with time-permuted traces showed (Fig. 3d), thus further allowing us to investigate how

these visual responses are modulated by a texture displacement.

Fig. 3. Novel visual stimulation paradigm to study saccadic suppression in zebrafish optic tectum. a. Visual
stimulation setup allows for whole-field visual stimulation onto a spherical projection surface, which encloses a
live and behaving zebrafish larva (see left inset for photo of visual stimulus projected onto fish bowl). During visual
stimulation eye position can be tracked using an infrared camera and activity of tectal neurons can be measured
using 2P-imaging (right inset). Stimulus light was split by pyramidal mirror into four optical paths projecting onto
the fish bowl screen surface **b.** Coarse and fine textures were displaced in azimuth on the spherical projection
screen in a saccade-like manner. After a variable delay period (d), a dark probe flash was shown by reducing
texture luminance. **c.** Regression-based analysis of fluorescence time traces of individual ROIs (each representing
one tectal neuron) revealed many neurons responding significantly to the dark probe flash, while some neurons
did not respond to the dark flash. The left panel shows ΔF/F traces of example neurons from the tectal slice shown
in (a), across the entire recording; the right panel shows *z*-scores of responses to a flash shown in isolation for the
same ROIs. Black lines indicate the stimulus regressor, blue bars indicate the time points of texture displacements
and pink and gray traces indicate neuronal activity. Pink ROIs correspond to neurons with strong correlation
coefficient, R to the stimulus regressor, gray ROIs represent neurons without strongly correlated responses to the
visual stimulus. **d.** Average flash responses of all real ROIs recorded are larger compared to simulated ROIs with
responses that were circularly permuted in time (p < 0.0001, two-tailed Wilcoxon rank-sum test, *n = 83,104 ROIs*).

Texture displacement suppresses visual responses to global dark flash in the optic tectum

To investigate how texture displacements modulate visual responses in the OT, we normalized flash responses around the time of saccade to each ROI's response to a flash shown in temporal isolation. Analogous to our analysis in zebrafish retinae, any responses associated to the texture displacement were removed and remaining flash responses were quantified. When looking at all flash-responding ROIs, we can see that more than 70% of neurons showed overall responses that were modulated downward, thus suppressed, by the presence of a texture displacement (Fig. 4a, Supplementary Fig. 1a). When comparing the 20% most suppressed and most enhanced cells to the same dataset with shuffled delay identities, we can see that, not only are responses more suppressed in the real data (more green in Fig. 4b, left), but enhancement is much weaker in the real data than expected by chance (less orange in Fig. 4b, left).

We next tested whether the temporal profile of saccadic suppression is simply inherited from the retina or does change in the tectum. From Fig. 4b it is apparent that most of the suppressed neurons have their strongest suppression at 1000 ms or later after texture displacement onset (Fig. 4b, red box). This was quite surprising to us as retinal suppression usually ends before 1000 ms (~250 ms in our dataset). Texture displacements seem to modulate visual responses in the OT in a less straightforward fashion than in the retina. This can not only be seen by this very late suppression in Fig. 4b, but also when looking at the whole population of visual neurons in the OT. While there are, as previously stated, much more suppressed than enhanced neurons, a spectrum of modulation exists, with many neurons also being enhanced or not modulated at all (Supplementary Fig. 1a). Further, modulated neurons can be found in the tectum in a more posterior-dorsal location, compared to all neurons registered in the recording window (p<0.0001). This location corresponds to upper temporal visual space. The recording

Fig. 4. Prolonged suppression in the OT after texture displacement. a. Cumulative distribution of flash response suppression across all flash-responsive neurons. Neurons were suppressed (or enhanced) by displacement of the coarse texture in a saccade-like manner (all delay times pooled). The majority of neurons have suppressed flash responses (curve for real data is shifted to the left relative to the curve for data with shuffled delay identities, p < 0.0001, two-tailed Wilcoxon rank-sum test, *n = 3,758 ROIs*). **b.** Modulation of flash responses at various flash delays of the 20% most suppressed and 20% most enhanced ROIs. Real data (left panel) shows stronger suppression and much weaker enhancement, compared to shuffled data (right panel). Red dashed box indicates that most of the suppressed neurons have their strongest suppression for delays ≥ 1000 ms. On the y-axis cells are sorted according strength of suppression/enhancement, peak delay time, and response magnitude at peak delay (for details see suppl. Fig. 1 and methods description). **c.** Center positions of all flash-responsive ROIs. Modulated ROIs can be found more posteriorly, compared to all registered ROIs (p < 0.0001 for enhanced vs. all ROIs and suppressed vs. all ROIs on AP-axis, two-tailed Wilcoxon rank-sum test, Bonferroni corrected, n = 866 enhanced ROIs, 2,330 suppressed ROIs, 31,944 total registered ROIs). There was no difference in anatomical distribution between enhanced and suppressed ROIs (p = 0.18 for enhanced vs. suppressed on AP-axis, p = 0.78 for enhanced vs. suppressed on ML-axis, two-tailed Wilcoxon rank-sum test, Bonferroni corrected). **d.** Dependence of percent suppression (mean ± SEM) on delay time for all suppressed neurons. Flash responses were significantly suppressed where indicated by a hash symbol (# p < 0.0001, two-tailed Wilcoxon signed-rank test, Bonferroni corrected for n = 24 conditions). Suppression was significantly stronger for coarse textures, than fine (solid vs. dashed line) for all delay times (* p < 0.001, two-tailed Wilcoxon signed-rank test). *N = 869 ROIs from 6 animals.*

window expanded to parts of the dorsal pretectum, where the proportion of modulated neurons was

smaller, compared to the tectum (Supplementary Fig. 2). However, there was no difference in

anatomical distribution between enhanced and suppressed neurons (Fig. 4c, Supplementary Fig. 2). To

further investigate the temporal dynamics of saccadic modulation in the OT of not all, but only

suppressed neurons, we used a permutation test to identify truly suppressed neurons independent of

their actual temporal dynamics (Methods, Supplementary Fig. 3). Of the flash-responsive neurons 23.1

% (869/3758) were significantly suppressed and these neurons do in fact appear to inherit retinal

suppression for delays up to 500 ms: Peak suppression for coarse and fine texture before 500 ms is nearly

identical to retinal suppression (Fig. 4d, compare to Fig. 1e), with 13.8 ± 0.7% suppression for coarse

and 6.8 ± 0.7% suppression for fine textures in the OT, compared to 15.8 ± 3.0% and 8.6 ± 3.4% in retina

(mean ± SEM for all). The spatial frequency dependency observed in the retina remains intact as well.

However, at delays ≥500 ms from texture displacement onset, a secondary, much stronger suppression

occurs, with its peak at 2000 ms and 21.0 ± 0.7% (mean ± SEM) suppression of visual responses for

coarse texture displacements.

We were startled by this finding, since saccadic suppression lasting >1 second in an ecological sense,

would be quite a devastating loss of visual information. One possible explanation could be that our

experiment did not truly assess saccadic suppression, but rather texture displacement suppression. In

this paradigm the tectum only received visual information about an apparent saccade. It is, however,

known in primates, that saccadic suppression is shorter with real saccades compared to visually

simulated saccades[6,20]. This is why we investigated in our next steps, if the presence of a motor signal

could shorten tectal suppression to values similar to other species. After all, in ecological settings, both

visual and motor signals are present to shape saccadic suppression.

Motor signals strengthen early saccadic suppression, but do not abolish late suppression

Surprised by the exceptionally strong and long lasting extra-retinal suppression we observed in flash responses of tectal neurons following a visually simulated saccade, we tested if the presence of a motor signal could shorten saccadic suppression to an ecologically reasonable length. For this purpose, we had fish perform spontaneous horizontal saccades, either in front of a textured or uniform background. Saccades were detected online and again at various delay times after the saccade onset a global dark flash was presented (Fig. 5a, b). With the same regression-based analysis described above, visually responsive ROIs were identified and analyzed for their flash responses (Fig. 5b).

Once again, visual responses are diversely modulated up or down (Supplementary Fig. 1b). When looking at the 20% most suppressed neurons, it can be seen that the temporal suppression profile is much more structured than would be expected by chance (compare Fig. 5c and d, left with Fig. 5d, right), with and without visual feedback. Consistent with our findings in simulated saccades, more neurons are suppressed than enhanced (Supplementary Fig. 1b). With visual feedback, similar to the texture displacement experiment, more than 70% of neurons are suppressed across all delay times. Without visual feedback ~60% of neurons are still modulated downwards by a saccade (Fig. 5e), albeit less than with visual feedback. Therefore, suppression in the OT can also be mediated by a real saccade.

The visually induced saccadic suppression, which can be seen in texture displacement experiments in retina and OT is further altered in the presence of a motor signal. Peak suppression for early delays <500 ms is nearly doubled with a motor signal (13.8 ± 0.7% suppression for texture displacement vs. 26.4 ± 1.1% suppression saccade (mean ± SEM for both, coarse stimulus condition)). Further, we had anticipated that through the presence of a motor signal, long-lasting suppression after >500 ms would be weakened or even abolished. To our surprise, when looking at suppressed neurons, we could still see strong suppression multiple seconds after saccade onset (Fig. 5f). This time course of saccadic

282   suppression was also not altered by visual feedback. When saccades were made across a uniform

283   background, we saw the same suppression strength and length as with a textured background.

284   Therefore, saccadic suppression in the OT can be accomplished by motor signals alone in absence of

285   visual feedback. This suppression is stronger than visual suppression alone, showing the strong

286   influence of a corollary discharge signal in saccadic suppression.

Fig. 5. Motor signals induce strong and long-lasting tectal saccadic suppression. a. Schematic of experimental paradigm. Fish performed spontaneous saccades either in front of a textured or uniform background. Saccades were detected online (blue trace) and after a variable delay period (d) a dark probe flash was shown, by reducing luminance of the texture or uniform background (pink line). **b.** Sample recording of closed-loop stimulus design. Eye position is tracked via IR camera and saccades are detected online (upper panel). Stimulus luminance is reduced after delay period (middle panel). An example neuron responds with a weaker $\Delta F/F$ signal for shorter delay periods, than for longer delay periods (lower panel). **c. & d.** Saccade-induced modulation of flash responses at various flash delays of the 20% most suppressed ROIs, without visual feedback (uniform background, panel c) and with visual feedback (textured background, panel d left). Temporal suppression profiles are more structured than expected by chance (shuffled data, panel d right). **e.** Cumulative distribution of saccade-induced modulation of flash responses in visual neurons across all flash delay times. A majority of visual neurons showed suppressed flash responses after a saccade across a coarse texture or across a uniform background; more than expected by chance ($p < 0.0001$, two-tailed Wilcoxon rank-sum test, n = 3,242 ROIs for textured background and shuffled data, 5,397 ROIs for uniform background). **f.** Temporal profiles of percent suppression of response (mean ± SEM) for all neurons suppressed by a saccade across coarse, fine or uniform background. Flash responses were significantly suppressed for all delay times and all backgrounds (# $p < 0.0001$, two-tailed Wilcoxon signed-rank test). Suppression was significantly stronger for coarse than fine textures (solid vs. dashed red line) for all delay times (* $p < 0.001$, two-tailed Wilcoxon signed-rank test). *N = 664 ROIs from 6 animals for textured background, n = 540 ROIs from 6 animals for uniform background.* **g.** Normalized flash response for delay times smaller than 500 ms and saccade size are negatively correlated, where larger saccades lead to more strongly suppressed flash responses ($p < 0.001$, Spearman rank correlation, n = 657 saccades).

If the purpose of such a corollary discharge signal is to act as a suppressive command for visual neurons during rapid eye movements, larger saccades, that lead to larger visual transient, should induce stronger suppression. Therefore, we tested if motor driven saccadic suppression in the OT might also correlate with saccade size. Particularly for early delay times <500 ms, where motoric suppression has its greatest effect, saccade size is indeed negatively correlated with normalized flash response, thus for greater saccade sizes, responses are down-modulated more strongly or suppressed (Fig. 5g), not however for late delay times ≥500 ms (Supplementary Fig. 4). This is in congruence with recent findings of a corollary discharge signal of saccade-like swim bouts in the zebrafish OT, which in size is positively correlated with swim vigor[21].

In summary, visual responses to global flashes are further modulated in the OT by the presence of a motor signal. Early saccadic suppression is stronger in the presence of a motor signal. In fact, motor signals alone, in the absence of visual feedback can elicit this suppression. Suppression strength is

correlated with saccade size, indicating that a corollary discharge signal is graded to the expected

sensory signal that is to be suppressed. And lastly, both simulated and real saccades induce long lasting

(>500 ms) suppression for global flash stimuli.

Saliency filtering can explain long lasting suppression in OT

After having found that motor signals can enhance early saccadic suppression but cannot abolish long-

lasting suppression to a global flash stimulus, we considered the behavioral context of saccades in larval

zebrafish. Larval zebrafish will perform non-reflexive saccades primarily to reorient their gaze and

examine different parts of the visual field or in the context of prey capture to locate prey objects on the

area centralis[15,28]. In both cases, it is beneficial for the animal to focus visual attention to local stimuli,

such as an approaching predator or a small prey object. Now, if the animal was visually inhibited, due

to saccadic suppression lasting several seconds past an actual saccade, behaviorally highly salient stimuli

like predator and prey, could not be detected. Therefore, we suspected that late suppression, lasting

>1000 ms could act like a visual saliency filter, ensuring efficient sensory processing, rather than

saccadic suppression per se. While behaviorally relevant local stimuli could remain fully visible, less

relevant global stimuli are suppressed.

To test this hypothesis, we combined a texture displacement with a local prey-like moving dot, rather

than a global flash, as in previous experiments (Fig. 6a). The local dot stimulus elicited visual responses

in fewer and more anterior-medially localized neurons, as is expected, due to the tectum's retinotopy[29]

(Fig. 6b, corresponding to stimulus positions in the upper-nasal visual field). The responding neurons,

however, had comparable response levels as those responsive to global flash stimuli in the previous

experiments (Supplementary Fig. 5). As expected, visual responses were barely modulated across all

Fig. 6. Responses to local stimuli persist after simulated saccades. a. Texture displacement was paired with a local prey-like moving dot stimulus as a probe stimulus. Coarse texture was displaced in a saccade-like manner (blue line) and after a variable delay period (d) a dark dot with a diameter of 20° was shown at 10° elevation, moving in azimuth at 60°/s across the left frontal visual field (pink line, Methods). b. Center positions of ROIs that showed responses to the moving dot stimulus. Moving dot-responsive ROIs (pink) were located more medially and more anteriorly, compared to all ROIs (grey) that could be segmented within the recording window (p < 0.0001 for ML-axis, p < 0.0001 for AP-axis, two-tailed Wilcoxon rank-sum test, Bonferroni corrected, n = 2,048 dot-responsive ROIs, 18,497 total registered ROIs) c. Cumulative distribution of texture displacement-induced modulation of moving dot responses in visual neurons across all flash delay times. Unlike in experiments with a global flash probe stimulus (dashed blue line), for the local dot stimulus (solid blue line), visual neurons showed equal amounts of suppressed and enhanced responses following texture displacement as expected by chance (p < 0.0001 for local dot vs. global flash, p > 0.05 for local dot vs. shuffled data, two-tailed Wilcoxon rank-sum test, n = 2,259 ROIs) d. Temporal profile of response suppression for local dots (mean ± SEM) for all neurons suppressed by a displacement of a coarse or fine texture. Local dot responses are only suppressed for delay times ≤1000ms for both coarse and fine textures (# p < 0.0001, two-tailed Wilcoxon signed-rank test). There was no difference between the two textures (p > 0.05, two-tailed Wilcoxon signed-rank test). *N = 504 ROIs from 3 animals.*

delay times, and we did not see more suppressed than enhanced neurons, like we did in previous global flash experiments (Fig. 6c). When looking at truly suppressed neurons only, we can see that there is only very weak suppression for early delay times; late suppression is completely abolished (Fig. 6d). This early suppression is likely inherited directly from the retina, as it follows a roughly similar time course and suppression strength (compare two black lines in Fig. 7a).

Overall, we show that late saccadic suppression lasting >1000 ms is heavily dependent on the probe stimulus used. While global stimuli, such as a global flash, remain suppressed for several seconds after a saccade, behaviorally relevant local stimuli, such as a prey-like dot, become quickly perceivable again. This task-specific processing of visual information in the tectum could function as a saliency filter, ensuring efficient sensory encoding for zebrafish that have limited neural capacities.

Saccadic suppression is shaped by the type of visual stimuli and motor signals in zebrafish OT

When combining the observations from all experiments described above, we can identify both visual and motor-related mechanisms that shape saccadic suppression dynamics across the zebrafish visual system (Fig. 7a, Fig. 8). In isolated adult retinae, RGC responses to global flash stimuli are weakly suppressed for up to 250 ms; the suppressed signal is relayed to the OT. Here, visual responses of larval tectal neurons are even more strongly suppressed by a real saccade, than a simulated saccade, particularly in the first 500 ms after saccade onset. At later time points (≥1000 ms) after saccade onset, global and local visual stimuli are differentially processed by tectal neurons. While responses to global flashes are strongly suppressed for several seconds, responses to a local prey-like dot, after weak early suppression in the first 1000 ms after saccade onset, quickly return to baseline response level.

Fig. 7. Visual and motor signals act on non-overlapping neural populations in the OT. a. Temporal suppression profiles in retina and OT of all visual and motor conditions tested in this study. Visual Only refers to experiments with simulated saccades on a textured background, Visual & Motor Combined refers to experiments with real saccades across a textured background, and Motor Only refers to experiments with real saccades across a uniform background. **b. i.** Proportions of neurons suppressed exclusively in the Visual & Motor Combined condition (red) or exclusively in the Visual Only condition (blue) versus neurons suppressed in both conditions (purple) when tested within the same animal, compared to the proportions in data with shuffled condition identities. There are more Visual & Motor, exclusive suppressed neurons than expected by chance ($p < 0.05$, χ^2-test for proportions) and less Visual Only, exclusive suppressed neurons than expected by chance ($p < 0.05$, χ^2-test for proportions). **ii.** Proportions of neurons suppressed exclusively in the Visual & Motor Combined condition (red) or exclusively in the Motor Only condition (yellow) versus neurons suppressed in both conditions (green) when tested within the same animal, compared to the proportions in data with shuffled condition identities. Only few neurons are suppressed in both conditions, no more than expected by chance ($p = 0.27$, χ^2-test for proportions), suggesting that they are separate populations.

Lastly, when investigating motor and visual effects in neurons of the same animal, we found non-overlapping populations of neurons. While some neurons were exclusively suppressed through the visual feedback of a texture displacement (50/94 ROIs, 53.2%), a large proportion of those suppressed through texture displacement could also be suppressed through a saccade across texture (44/94 ROIs, 46.8%), more than would be expected by chance (Fig. 7b, i.). However, most of the neurons, that were suppressed through a real saccade, were not suppressed in the simulated condition (172/216 ROIs, 79.6%), indicating that those neurons required both motor and visual feedback; visual feedback alone did not suffice. When comparing saccades with visual feedback to saccades without visual feedback in

neurons of the same animal, we found that almost all neurons are exclusively suppressed in either one

of the two cases, saccades with visual feedback (137/149 ROIs, 91.9%) or saccades without visual

feedback (149/161 ROIs, 92.5%). Very few neurons are suppressed for both, no more than expected by

chance (Fig. 7b., ii.), indicating that this could be two separate populations of neurons.

Discussion

We identified distinct visual and motor-related processes, shaping saccadic suppression on a neuronal

level across the zebrafish visual system (Fig. 8). Purely visual effects, such as masking, already take place

in the zebrafish retina prior to entering the OT. Following a saccade-like texture displacement, RGCs

show suppressed responses to peri-saccadic visual stimuli. The length of the peri-saccadic suppression

window in zebrafish retina is comparable to that found in mammals, but suppression strength is much

weaker[6]. Spatial frequency dependencies of suppression are similar to those observed in mammals in

that low spatial frequency content is more strongly suppressed than high spatial frequency content.

Zebrafish have lower visual acuity though, so the stimuli were coarser in this study than in research on

mammals.

This modulated retinal signal gets further relayed to the OT, the major retinorecipient area in the

zebrafish brain. There, saccadic suppression is further shaped by both visual and motor signals. Saccade-

like texture displacements elicit saccadic suppression of visual responses in the OT just as in the retina.

In fact, suppression in the early peri-saccadic window (delays <500 ms) is likely directly inherited from

the retina, as it follows the same time course and has nearly identical suppression strength. When

comparing purely visually induced "saccadic" suppression to saccadic suppression following an actual

spontaneous saccade, we found that the execution of an actual motor command significantly increases

Fig. 8. Visual and motor mechanisms shape saccadic suppression throughout the zebrafish visual system. Both visual and motor-related signals shape the temporal profile of saccadic suppression in the zebrafish. Purely visual mechanisms (i.e. visual masking) lead to early, weak saccadic suppression in the retina (blue box). Retinal signals are sent to the OT and further modulated. Motor signals, likely originating from the hindbrain oculomotor circuits, strengthen early saccadic suppression in the OT (red box). Visual filtering mechanism in the OT can alter late saccadic suppression to last more than 1000 ms (gray box). Behaviorally salient stimuli, such as a local dot, are not further suppressed, whereas behaviorally irrelevant stimuli, such as a global flash are suppressed for long time scales (gray box). Motor and visual signals act together on different types of tectal neurons. While some neurons are only suppressed by either a visual or a motor signal, most neurons integrate suppression from both signals (yellow box).

suppression strength in the early peri-saccadic window. This result suggests that corollary discharge signals play an important role in strengthening saccadic suppression. The observed saccade-size dependency (Fig. 5g) supports this idea. Corollary discharge signals are known to play a vital role in optimizing sensory processing during self-motion across the animal kingdom[30–32] and recently, it has been shown that the zebrafish OT receives corollary discharge signals during saccade-like swim bouts[21].

It is likely that eye movement signals could elicit a similar corollary discharge signal in zebrafish, and our observed "motor only" saccadic suppression with a homogeneous visual background suggests that these corollary discharge signals exist. Future studies are needed to measure the saccadic corollary discharge signal more directly in the zebrafish OT and demonstrate its mechanistic involvement in saccadic suppression.

On a behavioral level, in primates, motor signals are known to shorten the peri-saccadic suppression window[33], potentially due to post-saccadic excitability enhancement in cortical areas[34–36]. The lack of a cortex is thus a potential proximal explanation for the lack of a motor-associated temporal restriction of saccadic suppression in zebrafish.

Another striking difference between saccadic suppression in the zebrafish OT and saccadic suppression in primates is a strong secondary suppression in zebrafish, particularly of global flash responses for several seconds past the saccade onset. This was true for simulated saccades and real saccades with and without visual feedback. We were struck by this finding, as from an ecological perspective, it seems like a massive loss of visual information, to discard sensory signals for up to 4 seconds after a saccade. Using an imaging technique with very slow temporal resolution of ~2 Hz, attempting to resolve an effect that ought to play in the tens and hundreds of millisecond time scale, we were concerned that the slow suppression dynamics were an artifact of our stimulus design and poor temporal resolution. Thus, in a small subset of recordings, we tested a short 20 ms flash stimulus, comparable to the brief probe flash used in the high temporal resolution MEA recordings in the retina. In order to capture small and rapidly changing neural responses to this brief visual stimulus, we chose to image a much smaller brain region, but at 50 fps temporal resolution. We found that even with this brief visual stimulus, and with recordings at high temporal resolution, suppression lasting >1000 ms remained (Supplementary Fig. 6).

We propose that the persistent suppression is a task-specific efficient encoding strategy for the larval

zebrafish, which has to deal with limited neuronal resources. Since spontaneous saccades are often

followed by the detection of a local visual stimulus, either in a prey capture setting or when detecting a

predator[15,28], it would be sensible for the fish to focus its neural resources to accurately detecting local

visual stimuli and discarding information about global stimuli. This is exactly what we found when

testing a local prey-like dot stimulus for saccadic suppression in the OT. While the visual suppression

inherited from the retina remains intact, there is no further modulation of the visual signal after the

initial peri-saccadic window. Such attentional filtering for behaviorally salient stimuli following a

saccade has already been predicted in computational models of cortical population activity; in a

competitive interaction between two stimuli, modeling studies predict suppression of unattended

stimuli to enhance sensitivity to relevant stimuli[37]. Further, studies in archerfish have shown that the

teleost optic tectum is capable of using visually driven long-range inhibition, i.e. inhibition modulated

by signals from outside the receptive field, to aid in selecting behaviorally salient stimuli, such as prey

over behaviorally irrelevant stimuli[38]. In the archerfish retina, saccades might serve to encode visual

stimuli with higher fidelity[39]. Overall, future studies are needed to reveal the differential suppression of

various ecologically relevant visual stimuli in the context of saccadic suppression. In the cortical area

V1 of primates, preliminary data suggests that dark contrasts are not suppressed at all, unlike in superior

colliculus and primate perception[40,41], which could represent an ecological adaptation as well, and this

phenomenon needs to be addressed in comparative studies. Furthermore, a better understanding of the

receptive field properties and functional roles of tectal neurons will be required to fully understand why

and how different visual signals are suppressed in the OT.

Lastly, we investigated how visual and motor signals can act on tectal neurons within the same animal.

We have found that while there are many neurons that can be suppressed purely by a visual signal of a

simulated or real saccade across a texture, a larger proportion of neurons requires a simultaneous motor

signal, arising from an actual saccade in order to be suppressed. Further, when comparing which

neurons are suppressed through the execution of a saccade in the absence of any visual feedback to those

suppressed through a saccade with visual feedback, we found that there is almost no overlap between

these two cell populations. It is unclear why motor suppression and visually driven suppression are

segregated in such a manner. It seems that saccadic suppression is used as a precise tool for various

different contexts. This suppression layout across the neuronal population could be the results of

balancing suppression against information loss, since only small populations are suppressed, and only

under certain conditions.

In summary, we identified that saccadic suppression exists in the small-brained larval zebrafish, and

that visual and motor signals play differential roles. Compared to mammals, zebrafish saccadic

suppression is weaker but longer-lasting. Further studies are needed to reveal conserved mechanisms

that govern saccadic suppression and potentially explain the observed differential expression of

suppression dynamics across species.

511 Methods

512 Zebrafish

For retinal recordings we used retinae of *nacre +/-* adult zebrafish (*Danio rerio*), which are functionally

wild type. Ten retinae from 6 male and 4 female fish (age 8-23 months) were used. Adult zebrafish were

housed at our local animal facility at 28°C water temperature at a 14/10h light/dark cycle. Animals were

brought to the laboratory space and dark-adapted for several hours prior to retinal tissue preparation.

For *in-vivo* calcium recordings, larval zebrafish, age 5-7 dpf, with pan-neuronal GCaMP6f expression

and homozygous for the *mitfa* mutation were used (*Tg(HuC:H2B-GCaMP6f)jf7; nacre -/-*)[42]. Zebrafish

were reared in E3 solution at 28°C on a 14/10 light/dark cycle. At 4 dpf, animals with the strongest GFP

expression were selected for subsequent recordings at 5-7 dpf. Animal experiments and all experimental

protocols were approved by the responsible ethics committee of the Regierungspräsidium Tübingen in

accordance with German federal law and Baden-Württemberg state law. All animal procedures also

conformed to the institutional guidelines of the University of Tübingen.

Retinal electrophysiology

Tissue preparation (MEA). Fish were dark adapted for a minimum of 2 hours, before they were killed

under dim red light either with anesthetic overdose (250 mg/L Tricaine, Pharmaq, in fish water) or

through rapid chilling (~2-4°C). Afterwards, animals were decapitated and the eyes were removed and

placed into chilled Ringer solution (in mM: 110 NaCl, 2.5 KCl, 1.0 CaCl2, 1.6 MgCl2, 10 D-Glucose,

and 22 NaHCO3) bubbled with 5% CO_2 and 95% O_2. Retina was isolated inside a Ringer bath, placed

into 5% Hyaluronidase solution for 3 min to remove remaining vitreous and then placed into warmed

Ringer solution (~30°C) for one hour to recover tissue. Following recovery, retina was mounted onto a

nitrocellulose filter (Millipore) with a central 1.5 x 1.5 mm hole and placed RGC side down onto a

perforated low-density multi-electrode array (60pMEA200/30iR-Ti-gr, Multichannel Systems,

Reutlingen, Germany) with 60 electrodes in a square-grid arrangement and 200 μm inter-electrode

distance. Good electrode contact was ensured through negative pressure through the perforation over

at least 30 minutes, prior to recording begin. Tissue was superfused with bubbled Ringer solution at

~30-32°C. Data were acquired at 25 kHz using MC Rack version 4.6.2 (Multichannel Systems).

Visual stimulation (MEA). Visual stimuli were adapted from Idrees and colleagues[6]. A spectral

intensity profile (in µW cm^{-2} nm^{-1}) of the visual stimulus was measured with a calibrated USB2000+

spectrometer (OceanOptics, Orlando, FL, USA) and converted into biological units (R* rod^{-1} s^{-1})[43]

normalized zebrafish rod absorbance[44] (2.4 µm^2 for zebrafish rods[45]). On the DLP projector scale (0-

255) a stimulus intensity of "60" with ND5 filter (which was used in almost all retinal recordings)

equaled 3442 R* rod^{-1} s^{-1}.

Two binary Gaussian blur textures with differing spatial scales were used in the experiment. The fine

texture had image blobs with a size of 75 µm, the coarse texture had blobs of 300 µm. In adult zebrafish,

this equates to approximately 3.5° and 14.3° of retinal surface for fine and coarse texture respectively[46].

The pixel intensity of the textures ranged from 30-90 with a mean intensity of 60 on the 8-bit DLP

projector scale (0-255).

In order to simulate saccades, the texture was rapidly displaced for 100 ms at 100 to 300°/s at a straight

trajectory in random direction. At various time points around the texture displacement (-150, 17, 50,

100, 250, 500, and 2000 ms after texture displacement onset) a probe flash was shown. The probe flash

was a full-screen positive or negative luminance step ("bright" or "dark" flash), that lasted for one frame

(16.7 ms). Local contrast within the texture remained constant between the probe flash and the

background texture. In total a single trial consisted of 15 conditions (7 delays with dark flash, 7 delays

with bright flash, 1 control displacement without flash) shown in random order; trials were repeated at

least four times for each texture type. Additionally, the following stimuli were shown to characterize

RGCs types and receptive field properties: 1. Full field contrast steps. To test ON/OFF properties of

RCGs, uniform background luminance steps from 30 to 60 and from 60 to 90 pixel intensity steps in

positive (ON) and negative (OFF) direction were shown. 2. Binary checkerboard flicker. For spatial

receptive field mapping a binary checkerboard pattern, with checkers of 55 x 55 μm size, randomly

switching between 10 or 50 pixel intensity independent of other checkers was shown. The stimulus was

shown for 10 minutes. Linear filters, representing spatial and temporal receptive fields were calculated

through reverse correlation.

Data analysis (MEA). MEA recordings were high-pass filtered at a 500 Hz cutoff frequency using a

tenth-order Butterworth filter. Spikes were sorted semi-manually using Matlab-based custom software,

and spike waveforms and times were extracted according to Idrees and colleagues[6]. RGC types were

characterized based on their responses to full-field contrast steps. For each cell an ON/OFF index was

calculated by dividing differences between ON and OFF step peak response by their sum. The index

ranged from -1 to +1, where -1 indicates a full OFF cell and +1 indicates a full ON cell. Cells with an

ON/OFF index > 0 where only analyzed for bright probe flashes and cells with an ON/OFF index < 0

were only analyzed for dark probe flashes. Further, we characterized the spatial receptive field of each

cell, by examining the linear filters for each region (checker) defined by the binary checkerboard

stimulus. A 500 ms window prior to each spike was summed and the modulation strength of each linear

filter, measured as the standard deviation along the 500 ms kernel, determined how strongly that

checker drives the RGC. The resulting 2D map of standard deviation values was fitted with a 2D

Gaussian, and receptive field size was determined as an ellipse with 1.5σ. Receptive field sizes were

converted from x μm to α degrees approximating the focal length (f) of the adult zebrafish eye as

1.2mm[46], such that $\alpha = 2\tan^{-1}\left(\frac{x}{2f}\right)$. Response latency was determined as the time from stimulus onset

to 90% peak response. All spike times were converted into firing rates (spikes/s) by convolving spike

times with a Gaussian of σ = 10 ms standard deviation and amplitude $0.25\frac{\sqrt{e}}{\sigma}$.

To determine retinal saccadic suppression for each recorded RGC, first baseline responses were defined

as peak firing rates in response to probe flashes at 2000 ms delay from texture displacement onset,

averaged across trials. Then, average peak firing rates of the same RGC to all other delay times were

normalized to this baseline response. Any texture displacement related responses were identified in

control trials, without probe flashes and averaged across trials for coarse and fine textures

independently. Flash responses were isolated from texture displacement responses by subtracting

average texture displacement responses from the total responses, leaving only the flash-related response.

In-vivo **calcium imaging**

Visual stimulation and recording setup. All visual stimuli used in this study were displayed on a

custom-built spherical stimulation arena, with simultaneous eye position recordings. In the stimulation

arena, fish were placed inside a custom printed Ø 80 mm resin bowl (Accura ClearVue, SLA resin, 3D

Systems GmbH, Meorfelden-Walldorf, Germany), evenly coated in white acrylic spray paint and sealed

with clear varnish. The resin fish bowl served as a projection surface for a DMD video projector

(LightCrafter E4500 MKII, EKB Technologies Ltd., Israel), powered by a multispectral high power LED

light source (CHROLIS, 365/420/470/625 nm, Thorlabs GmbH, Bergkirchen, Germany). In order to

cover the entire projection surface of the bowl, the image of the video projector was split into four paths

by a pyramidal mirror and refocused onto the fish bowl from all four sides (Fig. 3a) to cover -90° to

+45° in elevation and 360° in azimuth. All stimuli were presented at 60 Hz refresh rate at 625 nm (red)

to avoid light interference with calcium imaging. Irradiance and corresponding Michelson contrast

inside the fishbowl were measured using an optical power meter (Newport 1918-R optical power meter

with 918D-SL-OD3R detector, Newport Corporation, Irvine, CA, USA) and a Ø 5 mm liquid core light

guide (Thorlabs GmbH, Bergkirchen, Germany) positioned on the inside wall of one half bowl at

various elevations. At 625 nm stimulation wavelength the following measurements for high irradiance,

low irradiance and Michelson contrast were made at 22.5°, -22.5°, and -77.5° elevation: 22.5°: 8.56

mW/m^2 high, 0.51 mW/m^2 low, 0.89 contrast; -22.5°: 17.83 mW/m^2 high, 0.61 mW/m^2 low, 0.93

contrast; -77.5°: 5.79 mW/m^2 high, 0.61 mW/m^2 low, 0.81 contrast.

A custom-built 850 nm IR-LED ring provided illumination for eye position tracking through the

excitation path. A small window in the spherical stimulation arena below the fish (~5 mm diameter)

allowed for recording of eye potion with an infrared-filtered camera at 100 Hz (Basler ace 2 R, a2A1920-

160umBAS, Basler AG, Ahrensburg, Germany).

VxPy (Version 0.1.4) was used to generate and display visual stimuli, as well as record eye position traces

and trigger visual stimuli in a closed-loop manner.

Visual stimulation protocol (Calcium Imaging). Visual stimuli were designed in analogy to retinal

stimulation, but adjusted to the geometry of our stimulation arena. Instead of using a rectangular array

of black and white pixels and convolving these with Gaussians, we designed a stimulus in spherical

coordinates by first placing equidistant nodes on the surface of a unit sphere. Half of the nodes were

randomly selected as "bright" nodes, and identical von-Mises-Fisher distributions with concentration,

κ= 5.6 were placed centered on each of these. Nothing was placed on the remaining, "dark" nodes. After

summing up all distributions, intensities were rescaled to impose a desired minimum and maximum

across the sphere surface to ensure roughly uniform local luminance and contrast across the visual field.

The number of nodes is the parameter that determines pairwise distances between them. Its values were

chosen to produce stimuli with spatial frequencies (Supplementary Fig. 7) that reliably elicit behavioral

responses. For the coarse texture 2,000 nodes (1,000 bright) were used, for the fine texture 4,000 nodes

(2,000 bright) were used.

In "texture displacement" experiments, saccades were simulated through a rapid rotation of the texture

on the vertical axis by 30° at 300°/s either to the left or right. In "real saccade" experiments, saccades

were detected with a velocity threshold of at least 200°/s. Saccade threshold was individually adjusted

for each fish, to ensure best saccade detection, without noise interference.

At various delay times around texture displacement onset or saccade detection a probe stimulus was

presented. Delay times were usually -500, 20, 100, 250, 500, 1000, 2000, and 4000 ms, unless stated

otherwise. In "real saccade" experiments, no pre-saccadic stimulus could be shown. Probe stimuli were

shown twice without texture displacement or 8 s after real saccade onset and used as baseline stimuli.

Additionally, texture displacement/saccade occurred without probe stimulus. Therefore, one trial

consisted of 9-11 conditions (6-8 delays, 2 baseline probes, 1 control displacement/saccade), shown in

random order; trials were repeated at least 4 times.

In flash experiments the secondary visual stimulus consisted of a whole-field luminance decrease with

a cosine shape, lasting 500 ms at 0.5 Hz and an amplitude of 50% luminance change. For high frequency

recordings, a luminance step, lasting 20 ms with an amplitude of 50% luminance decrease was used.

In moving dot experiments, a dark dot, moving across the static texture was used as a secondary visual

stimulus. The moving dot had a 20° diameter and was located in the upper left visual field at 10°

elevation. The dot moved at 60°/s in azimuth from -30° to -60° and back to 0° over a time span of 1.5 s.

To avoid movement overlap between the texture displacement (lasting 100 ms) and the moving dot

stimulus, only delay times ≥ 100 ms were used in the moving dot experiment.

Two-photon calcium imaging. For in-vivo calcium imaging, fish were embedded in 1.6% low-melting

agarose and placed onto a custom-built stage, which could be attached to a holder on the inside of the

spherical stimulation arena. For the stage, the tip of a plastic Pasteur pipette was cut and shaped into an

arrowhead shape. Then, the pipette tip was placed into a custom 3D-printed mold and 2.5% low-melting

agarose was poured into the mold resulting in a transparent stage that allowed the fish to be placed

central inside the spherical stimulation arena. For "real saccade" experiments, agarose was removed

around the eyes of the fish, to allow for free eye movements. Once the fish was placed in the center of

the spherical stimulation arena dorsal side up, it was filled with clear E3 medium. The position of the

fish could be adjusted, to correct for potential roll.

Calcium activity of tectal neurons was recorded with a movable objective two-photon setup (Sutter

Instruments, Novato, California, USA), previously described[47]. The calcium indicator was excited at 920

nm and emitted light was collected with a 20x/1.0 Zeiss objective. In most recordings image acquisition

rate was 2.18 fps, with 512 x 512 pixels imaged. Only in high-speed recordings a smaller window (40 x

40 pixels) was imaged at 50 fps acquisition rate. Magnification ranged from 1-2x, where 1x results in a

resolution of 0.88 µm x 0.88 µm per pixel.

Data analysis (Calcium imaging). Calcium recordings with 2.18 Hz acquisition rate were registered

and segmented with suite2p[48] into individual ROIs, each representing one tectal neuron. A time

constant of 1.61 s was used and for recordings at 1.6x magnification a cell diameter of 10 pixels (5.5 µm)

was used. For different magnifications, the cell size was adjusted accordingly. The suite2p built-in

selection algorithm *is-cell* was ignored. High-speed recordings with 50 fps acquisition rate were

segmented manually in Fiji (ImageJ) into elliptical ROIs, based on standard deviation projections.

Center positions of segmented ROIs were registered to the Max Planck Zebrafish Brain (mapzebrain)

atlas[49] using ANTsPy[50].

Further data analysis was performed using custom-written Matlab (The MathWorks Inc., Natick, USA)

and Python (PyCharm, JetBrains, Prague, Czech Republic) scripts. Fluorescence traces for each ROI

were normalized into $\Delta F/F_0$, with a sliding window median of 120 seconds for F_0. Through a regression-

based analysis, ROIs that responded to the secondary visual stimulus used in the experiment (flash or

moving dot), when shown in temporal isolation ("baseline", i.e. without saccade or texture

displacement), were identified and only those ROIs were considered for further analysis

(19,784/119,220 ROIs across all flash experiments, 16.6%, 2,210/20,789 ROIs for moving dot

experiment, 10.6%).

Saccade- and texture displacement-associated activity in ROIs was identified, by averaging calcium

traces during the saccade/texture displacement from control conditions (without secondary visual

stimulus) and long delay conditions ($\geq$2000 ms) up to the point of secondary visual stimulus

presentation. Average saccade-/texture displacement-associated activity was then subtracted from the

calcium trace for the time points during and shortly after the saccade/texture displacement.

Peak $\Delta F/F$ responses to the secondary visual stimulus at each delay time were quantified and averaged

across trials for each ROI. For "real saccade" experiments, trials in which fish performed undesired

saccades between the initial saccade and presentation of the secondary visual stimulus were discarded.

Trials, in which the fish struggled, thus rapidly moved its head, rather than performing an actual saccade

were also discarded. Average peak $\Delta F/F$ responses were normalized to average baseline responses

without saccade/texture displacement. Individual modulation curves for each ROI were generated as in

Supplementary Fig. 2b.

To identify suppressed ROIs, amongst all ROIs sensitive to the visual stimulus, modulation curves of

individual ROIs were first correlated against 6-8 model suppression curves (Supplementary Fig. 2c),

depending on the number of delays used in the experiment. Model suppression curves had different

delay times of peak suppression and each a unimodal modulation. The average modulation across all

delays had to be <1, thus suppressed. Out of the model suppression curves, the curve with the best

correlation coefficient was then confirmed through a permutation test with a Bonferroni-corrected (n

= 6-8) significance cut-off of $p < 0.05$. ROIs that passed that permutation test were considered truly

suppressed (Supplementary Fig. 2).

Statistics

Graphical presentation and statistical analysis was done in Matlab. For retinal electrophysiology

recordings, all recorded RGCs were considered in group data. For *In-vivo* calcium imaging, only truly

suppressed ROIs (as described above) were considered for group data. Group data were tested for

normality using a Lilliefors test. Since almost all data did not pass this test we chose non-parametric

tests for further statistical analysis. Wilcoxon signed-rank tests were used for paired observations;

Wilcoxon rank-sum tests were used for independent sample testing. For regression-based analysis and

to identify truly suppressed ROIs in *in-vivo* calcium recordings, Pearson linear correlation was used.

For correlation between saccade size and normalized flash response (Fig. 5g), Spearman rank

correlation was used. A χ^2-test for proportions was used to compare proportions of suppressed ROIs

across experiments (Fig. 7b). For all tests p-value cutoffs were Bonferroni-corrected for multiple testing,

when necessary and $p < 0.05$ was considered as significance cut-off.

Data Availability

The data that support the findings of this study are available from the corresponding author upon

request.

References

1. Zuber, B. L. & Stark, L. Saccadic suppression: Elevation of visual threshold associated with
saccadic eye movements. *Exp Neurol* **16**, 65–79 (1966).

2. Beeler, G. W. Visual threshold changes resulting from spontaneous saccadic eye movements.
Vision Res **7**, 769–775 (1967).

3. Burr, D. C., Morrone, M. C. & Ross, J. Selective suppression of the magnocellular visual
pathway during saccadic eye movements. *Nature 1994 371:6497* **371**, 511–513 (1994).

4. Bremmer, F., Kubischik, M., Hoffmann, K. P. & Krekelberg, B. Neural dynamics of saccadic
suppression. *Journal of Neuroscience* **29**, 12374–12383 (2009).

5. Hafed, Z. M. & Krauzlis, R. J. Microsaccadic suppression of visual bursts in the primate
superior colliculus. *Journal of Neuroscience* **30**, 9542–9547 (2010).

6. Idrees, S., Baumann, M. P., Franke, F., Münch, T. A. & Hafed, Z. M. Perceptual saccadic
suppression starts in the retina. *Nat Commun* **11**, 1–19 (2020).

7. Easter, S. S. & Gregory Nicola, J. N. The Development of Eye Movements in the Zebrafish
(Danio rerio). *Dev Psychobiol* **31**, 267–276 (1997).

8. Schoonheim, P. J., Arrenberg, A. B., Del Bene, F. & Baier, H. Optogenetic localization and
genetic perturbation of saccade-generating neurons in Zebrafish. *Journal of Neuroscience* **30**,
7111–7120 (2010).

9. Bollmann, J. H. The Zebrafish Visual System: From Circuits to Behavior.
https://doi.org/10.1146/annurev-vision-091718-014723 **5**, 269–293 (2019).

10. Baumann, M. P., Idrees, S., Münch, T. & Hafed, Z. Selective peri-saccadic suppression of low
spatial frequencies is a visual phenomenon. *J Vis* **19**, 253–253 (2019).

11. Ridder, W. H. & Tomlinson, A. A comparison of saccadic and blink suppression in normal
observers. *Vision Res* **37**, 3171–3179 (1997).

12. Chen, C. Y. & Hafed, Z. M. A neural locus for spatial-frequency specific saccadic suppression
in visual-motor neurons of the primate superior colliculus. *J Neurophysiol* **117**, 1657–1673
(2017).

13. Michels, L. & Lappe, M. Contrast dependency of saccadic compression and suppression. *Vision
Res* **44**, 2327–2336 (2004).

14. Knöll, J., Binda, P., Concetta Morrone, M. & Bremmer, F. Spatiotemporal profile of peri-
saccadic contrast sensitivity. *J Vis* **11**, 15–15 (2011).

15. Bianco, I. H., Kampff, A. R. & Engert, F. Prey capture behavior evoked by simple visual stimuli
in larval zebrafish. *Front Syst Neurosci* **5**, 18496 (2011).

16. Sperling, G. Temporal and Spatial Visual Masking. I. Masking by Impulse Flashes. *JOSA, Vol.*
55, Issue 5, pp. 541-559 **55**, 541–559 (1965).

17. Matin, E., Clymer, A. B. & Matin, L. Metacontrast and Saccadic Suppression. *Science (1979)*
178, 179–182 (1972).

18. Campbell, F. W. & Wurtz, R. H. Saccadic omission: Why we do not see a grey-out during a
saccadic eye movement. *Vision Res* **18**, 1297–1303 (1978).

19. Idrees, S. *et al.* Suppression without inhibition: how retinal computation contributes to
saccadic suppression. *Communications Biology 2022 5:1* **5**, 1–23 (2022).

20. Diamond, M. R., Ross, J. & Morrone, M. C. Extraretinal control of saccadic suppression. *J*
Neurosci **20**, 3449–55 (2000).

21. Ali, M. A., Lischka, K., Preuss, S. J., Trivedi, C. A. & Bollmann, J. H. A synaptic corollary
discharge signal suppresses midbrain visual processing during saccade-like locomotion. *Nature*
Communications 2023 14:1 **14**, 1–18 (2023).

22. Kennard, D. W., Hartmann, R. W., Kraft, D. P. & Boshes, B. Perceptual suppression of
afterimages. *Vision Res* **10**, 575–585 (1970).

23. Volkmann, F. C. Human visual suppression. *Vision Res* **26**, 1401–1416 (1986).

24. Kayama, Y., Riso, R. R., Bartlett, J. R. & Doty, R. W. Luxotonic responses of units in macaque
striate cortex. *https://doi.org/10.1152/jn.1979.42.6.1495* **42**, 1495–1517 (1979).

25. Ibbotson, M. R., Price, N. S. C., Crowder, N. A., Ono, S. & Mustari, M. J. Enhanced Motion
Sensitivity Follows Saccadic Suppression in the Superior Temporal Sulcus of the Macaque
Cortex. *Cerebral Cortex* **17**, 1129–1138 (2007).

26. Easter, S. S. & Nicola, G. N. The development of vision in the zebrafish (Danio rerio). *Dev Biol*
180, 646–663 (1996).

27. Dehmelt, F. A. *et al.* Spherical arena reveals optokinetic response tuning to stimulus location,
size, and frequency across entire visual field of larval zebrafish. *Elife* **10**, (2021).

28. Dowell, C. K., Lau, J. Y. N. & Bianco, I. H. The saccadic repertoire of larval zebrafish reveals
kinematically distinct saccades that are used in specific behavioural contexts. *bioRxiv*
2023.11.07.565345 (2023) doi:10.1101/2023.11.07.565345.

29. Wang, K., Hinz, J., Zhang, Y., Thiele, T. R. & Arrenberg, A. B. Parallel Channels for Motion
Feature Extraction in the Pretectum and Tectum of Larval Zebrafish. *Cell Rep* **30**, 442-453.e6
(2020).

30. Crapse, T. B. & Sommer, M. A. Corollary discharge across the animal kingdom. *Nat Rev*
Neurosci **9**, 587 (2008).

31. Fukutomi, M. & Carlson, B. A. A History of Corollary Discharge: Contributions of Mormyrid
Weakly Electric Fish. *Front Integr Neurosci* **14**, 561912 (2020).

32. Cavanaugh, J., Berman, R. A., Joiner, W. M. & Wurtz, R. H. Saccadic Corollary Discharge
Underlies Stable Visual Perception. *Journal of Neuroscience* **36**, 31–42 (2016).

33. Idrees, S., Baumann, M. P., Franke, F., Münch, T. A. & Hafed, Z. M. Perceptual saccadic
suppression starts in the retina. *Nat Commun* **11**, 1–19 (2020).

34. Rajkai, C. *et al.* Transient Cortical Excitation at the Onset of Visual Fixation. *Cerebral Cortex*
18, 200–209 (2008).

35. Ibbotson, M. R., Crowder, N. A., Cloherty, S. L., Price, N. S. C. & Mustari, M. J. Saccadic
Modulation of Neural Responses: Possible Roles in Saccadic Suppression, Enhancement, and
Time Compression. *The Journal of Neuroscience* **28**, 10952 (2008).

36. Cloherty, S. L., Mustari, M. J., Rosa, M. G. P. & Ibbotson, M. R. Effects of saccades on visual
processing in primate MSTd. *Vision Res* **50**, 2683–2691 (2010).

37. Hamker, F. H., Zirnsak, M., Calow, D. & Lappe, M. The Peri-Saccadic Perception of Objects
and Space. doi:10.1371/journal.pcbi.0040031.

38. Volotsky, S., Vinepinsky, E., Donchin, O. & Segev, R. Long-range neural inhibition and
stimulus competition in the archerfish optic tectum. *J Comp Physiol A Neuroethol Sens Neural*
Behav Physiol **205**, 537–552 (2019).

39. Segev, R., Schneidman, E., Goodhouse, J. & Ii, M. J. B. Role of Eye Movements in the Retinal
Code for a Size Discrimination Task. *J Neurophysiol* **98**, 1380–1391 (2007).

40. Malevich, T., Zhang, T., Baumann, M. P., Bogadhi, A. R. & Hafed, Z. M. Faster Detection of
"Darks" than "Brights" by Monkey Superior Colliculus Neurons. *Journal of Neuroscience* **42**,
9356–9371 (2022).

41. Wu, W. & Hafed, Z. M. Stronger premicrosaccadic sensitivity enhancement for dark contrasts
in the primate superior colliculus. *Scientific Reports 2025 15:1* **15**, 1–17 (2025).

42. Dunn, T. W. *et al.* Brain-wide mapping of neural activity controlling zebrafish exploratory
locomotion. *Elife* **5**, (2016).

43. Tikidji-Hamburyan, A. *et al.* Retinal output changes qualitatively with every change in ambient
illuminance. **6**, (2014).

44. Cameron, D. A. Mapping absorbance spectra, cone fractions, and neuronal mechanisms to
photopic spectral sensitivity in the zebrafish. *Vis Neurosci* **19**, 365–372 (2002).

45. Vogalis, F. *et al.* Ectopic expression of cone-specific G-protein-coupled receptor kinase GRK7
in zebrafish rods leads to lower photosensitivity and altered responses. *Journal of Physiology*
589, 2321–2348 (2011).

46. Collery, R. F., Veth, K. N., Dubis, A. M., Carroll, J. & Link, B. A. Rapid, Accurate, and Non-
Invasive Measurement of Zebrafish Axial Length and Other Eye Dimensions Using SD-OCT
Allows Longitudinal Analysis of Myopia and Emmetropization. *PLoS One* **9**, 110699 (2014).

47.   Zhang, Y., Huang, R., Nörenberg, W. & Arrenberg, A. B. A robust receptive field code for optic
823   flow detection and decomposition during self-motion. *Current Biology* **32**, 2505-2516.e8
824   (2022).

825   48.   Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon microscopy.
826   *bioRxiv* 061507 (2017) doi:10.1101/061507.

827   49.   Shainer, I. *et al.* A single-cell resolution gene expression atlas of the larval zebrafish brain.
828   (2023).

829   50.   Avants, B. B. *et al.* A reproducible evaluation of ANTs similarity metric performance in brain
830   image registration. *Neuroimage* **54**, 2033–2044 (2011).

Acknowledgements

This work was funded by the Deutsche Forschungsgemeinschaft (DFG; German Research Foundation) through the Special Priority Programme: "SPP 2205 Evolutionary optimization of neural processing" (430158665; AR 1076/1-2, HA 6749/3-2, AR 1076/1-1), by the Major Research Instrumentation programme of the DFG (INST 37/1254-1 FUGG), and by the Human Frontier Science Program (HFSP) Young Investigator grant RGY0079. We want to thank all members of the Arrenberg group for fruitful discussions regarding this work, Saad Idrees for his incredible support and expertise regarding retinal electrophysiology and we want to thank Lidia García-Pradas for her excellent fish care.

Author Contributions

G.S. performed experiments in the retina and optic tectum, C.T. performed experiments in the optic tectum. G.S. and C.T. analyzed data and generated figures. T.C.H. developed the visual stimulation hardware and software. F.A.D. helped design visual stimuli and generated supplementary materials. G.S., Z.M.H. and A.B.A. designed the overall study. G.S. and A.B.A. conceived the experiments and associated analyses and wrote the manuscript. Z.M.H and A.B.A acquired funding for the project.

Competing Interests

The authors declare no competing interest.