N-terminal region of PetD is essential for cytochrome b_6f function and controls
STT7 kinase activity via STT7-dependent feedback loop phosphorylation

Afifa Zaeem[1], Yuval Milrad[1], Simon Bütfering[1], Carolyne Stoffel[2], Adrien Burlacot[2,3], Martin
Scholz[1], Felix Buchert[1*], Michael Hippler[1,4*]

[1]Institute of Plant Biology and Biotechnology, University of Münster, 48143 Münster, Germany
[2]Biosphere Sciences and Engineering Division, Carnegie Science, Stanford, CA 94305
[3]Department of Biology, Stanford University, Stanford, CA 94305
[4]Institute of Plant Science and Resources, Okayama University, Kurashiki, Japan
11
[*]Correspondence to: felix.buchert@uni-muenster.de; mhippler@uni-muenster.de

Abstract

The cytochrome b_6f complex (b_6f) links photosystem I (PSI) and photosystem II (PSII) in the
photosynthetic electron transfer chain and is distributed across appressed and non-appressed
thylakoid membranes. The b_6f also activates the state-transition 7 kinase (STT7), which
phosphorylates light-harvesting complex proteins, triggering their migration to enable energy
redistribution between PSII and PSI. STT7-dependent phosphorylation has also been observed at
T4 in the N-terminal domain of the b_6f subunit-IV (PetD), though its functional significance has
remained unclear. To investigate its role, we generated several chloroplast mutants. The
phosphomimic mutation PetD T4E inhibited STT7 kinase activity, as indicated by the absence of
STT7-dependent phosphorylation and the strain being locked in State 1. This reveals a novel
feedback mechanism regulating phosphorylation facilitated by STT7. The deletion of five N-
terminal amino acids resulted in a comparable inhibition of STT7 activity and, additionally, in the
disruption of electron transfer which underscores a crucial role of the PetD N-terminus in b_6f
function.

Main

Photosynthesis captures light through photosystem (PS) I and PSII along with light-harvesting complexes (LHCI and LHCII). Light energy drives water splitting at PSII and ultimately reduces ferredoxin (FDX) through PSI. In linear electron transfer (LEF), reduced FDX transfers electrons to FDX-NADP(H) oxidoreductase (FNR), producing NADPH for CO_2 fixation and biomass production. In addition, electron transport deposits H^+ in the lumen and separates charges across the thylakoid membrane, thereby generating both a proton gradient (ΔpH) and membrane potential ($\Delta\Psi$), which collectively drive ATP synthesis. Electron flow between the photosystems is mediated by the cytochrome b_6f complex (b_6f; reviewed in [1]), which shuttles electrons from plastoquinol (PQH_2) in the plastoquinone (PQ) pool to plastocyanin. Besides small peripheral subunits, the b_6f core comprises cytochrome-f (PetA), cytochrome-b_6 (PetB), Rieske iron-sulfur protein (ISP, PetC), and subunit-IV (PetD). High-potential redox centers contrast with low-potential cofactors, including heme c_i at the Qi-site. The latter is the stromal PQH_2 binding pocket opposing the lumenal Qo-site. By cycling protons between sites Qo and Qi, the ΔpH associated with b_6f activity substantially contributes to ATP production for CO_2 fixation. Moreover, the c_i heme within the Qi-site is relevant for cyclic electron flow (CEF) and the b_6f complex may function as a FDX-PQ reductase (FQR) in CEF [2,3].

The lumen acidification stemming from photosynthetic electron flows is important for efficient photoprotection under conditions of excess excitation energy: it establishes "photosynthetic control" at the Qo-site where PQH_2 oxidation (and therefore electron flow) ceases at acidic pH [4,5], and it reduces photochemistry by stimulating controlled heat dissipation via non-photochemical quenching (NPQ) [6-8]. Moreover, the b_6f is linked to state transitions (ST) – another photoprotective mechanism to cope with overexcitation [9]. During this process, LHCII migrates between PSII and PSI upon its phosphorylation by the thylakoid-bound STT7 kinase [10,11]. STT7 is activated by a reduced PQ pool while specific phosphatases act as STT7 antagonists [12]. The activity of the STT7 kinase is regulated by its transmembrane domain, with a disulfide bond between two lumenal cysteine residues being essential for its function [13]. Active discussions and research on the kinase redox regulation are ongoing [14-16]. Recently it was shown that ST involve PetD residues N122, Y124, and R125 in the stromal "fg Loop", linking helices F and G of this b_6f subunit [17]. Moreover, the C-terminus of the b_6f subunit PetB interacts with the fg Loop in PetD, thereby facilitating STT7 activation [18]. Mechanistically, binding of STT7 to b_6f enhances its autophosphorylation and R125 is directly involved in this process. In this report we focus on another stromal element of b_6f in the model green microalgae *Chlamydomonas reinhardtii* (see Extended data Fig. 1): The N-terminus in PetD where T4 undergoes phosphorylation in an STT7-dependent manner [19] (see also Extended data Fig. 2).

To investigate the functional significance of STT7-dependent phosphorylation at PetD T4, we

created PetD site-directed mutants T4A and T4E to mimic constitutive absence and presence of

phosphorylation, respectively. Moreover, the function of the five N-terminal residues was

addressed in a truncation mutant (ΔN) whereas the wild-type PetD (WT) was used as control. To

achieve this, chloroplast transformation targeted recombination within the *petD* locus,

incorporating an *aadA* cassette to confer spectinomycin resistance for selection. The recipient

strain used carried a six-histidine tag at the C-terminus of PetA [20] for optional b_6f purification.

Spot test on TP plates, requiring photoautotrophic growth, revealed that mutant strains T4A and

T4E performed similar to WT under normal (40 µmol photons $m^{-2} s^{-1}$) and moderate high light

(200 µmol photons $m^{-2} s^{-1}$; Extended data Fig. 3a). In contrast, the ΔN strain showed a growth

deficit at both light intensities compared to the other strains. Proteomic analysis showed no

significant differences in the levels of PetD and other core b_6f subunits between WT, T4A, and

T4E (Extended data Fig. 3b). Compared to WT, only PETO was significantly downregulated in

ΔN, while LHCSR1 was markedly upregulated in T4E relative to T4A (Extended data Fig. 4).

Given the observed growth phenotypes and the changes in the amounts of the CEF effector

protein PETO [21] and LHCSR1 – required for pH-dependent NPQ [22] – key photoprotection

pathways associated with b_6f function might be altered in the mutants. To this end, we first

present data investigating ST as they require the physical presence of b_6f [23], and then performed

electron transfer measurements to directly assess b_6f activity in the mutants.

To assess if ST are affected in the mutants, the antenna redistribution between PSI and PSII was

measured using 77 K fluorescence emission spectroscopy [24] after cells were shifted to either State

1 or State 2 inducing conditions (Fig. 1). Thereby, an increase in antennae size under State 2

versus State 1 conditions was observed for WT and T4A (Figs. 1a, 1b). In contrast, T4E and ΔN

strains displayed no differences in their 77 K fluorescence emission spectra, indicating they were

unable to perform ST (Figs. 1c, 1d).

The 77 K data suggest impaired STT7-dependent phosphorylation in T4E and ΔN strains, which

was supported by an additional phospho-proteomic analysis of PetD variants. Here, absence of

STT7 phosphorylation levels of S569 and T571 (Extended data Fig. 5) coincided with lack of ST.

To look for STT7 targets in our phospho-proteomic search we also included a CRISPR/Cas9 *stt7*

KO mutant and its respective WT in the analysis as a control (Extended data Fig. 6). Differential

expression analysis revealed STT7-dependent phosphorylation sites with significant upregulation

or downregulation in samples from State 2 conditions (absolute log2-fold change ≥ 1, $p < 0.05$,

Extended data Tab. 1, Extended data Fig. 2). The chloroplast protein PTAC16 showed the highest

number of differentially regulated phosphorylation sites. Generally, decreased or absent

phosphorylation of PTAC16 was observed in the *stt7* strain (Extended data Fig. 2). T4E and ΔN

showed similar PTAC16 patterns but retained some phosphorylation at WT sites (Fig. 2). The

T4A strain showed no significant differences. Phosphorylation differences varied between strains and may depend on genetic background. PTAC16 was recognized as a protein of the plastid transcriptionally active chromosome complex [25], and has also been identified as a substrate of STN7 in Arabidopsis [26]. WT cells showed substantially higher levels of dual T7/T11 phosphorylation on LHCB4 compared to *stt7*, with a 6.8-Log2fold change. The *stt7* mutation had widespread effects, eliminating phosphorylation at numerous key sites. These included single phosphorylation sites on proteins like PetD (T4), PETO (T161/S164), and ANR1 (T171), as well as several critical threonine clusters - notably the T33/T35/T37 region of LHCB4, T23/T36 of LHCB5, and the N-terminal phospho-site cluster in LHCSR3 (S26/S28/T32/T33/T39) [27]. LHCB5 retained only minimal phosphorylation at T20 in the *stt7* background (Extended data Fig. 2). The ΔN and T4E strains also showed distinct phosphorylation patterns. Some sites undetectable in *stt7* were similarly absent in these strains – such as ANR1 (T171) in ΔN, and LHCBM1 (T27) in both ΔN and T4E. PETO showed an interesting pattern where T185 phosphorylation occurred only in conjunction with S187 phosphorylation in *stt7* and ΔN, though at lower levels than in wild type. The ΔN and T4E strains both exhibited reduced phosphorylation across several thylakoid proteins, including LHCBM5 (double phosphorylation of S28+T33), PSB34 (T39), PSAG (T71), and PSBR (T37, S43, and T37+S43), while also showing decreased STT7 autophosphorylation at S569 and T571 (Extended data Fig. 5). ΔN specifically showed decreased phosphorylation of LHCBM3 and LHCBM6 but was the only strain with augmented phosphorylation at T72 of PSAG. The T4A strain behaved distinctly, with most of the aforementioned phosphorylation sites remaining unchanged, though it uniquely showed substantially increased phosphorylation of PSBR (T39, S43, and T37+S43) compared to the other strains. Most of the STT7-dependent phosphorylation sites listed above had also been previously identified [19,28].

While most phosphorylations of thylakoid proteins decreased in the ST-deficient mutants, PSBD (T2) showed increased phosphorylation in *stt7* (5.5-Log2fold change), confirming previous results obtained from a different *stt7* mutant (Extended data Fig. 2, and [19]). PSBD phosphorylation also increased in T4E (3.4-fold log2), while ΔN and T4A strains showed no significant differences. Previous studies have also documented STT7-dependent phosphorylation sites on threonine residues in LHCBM1, LHCBM4, 6, 8, 9, LHCBM3, and LHCBM5 [19,28]. The STT7-dependent phosphorylation of LHCBM1 and LHCBM5 have been structurally resolved in the PSI-LHCI-LHCII ST complex [29]. Thus, we conclude that a single amino acid substitution mimicking constitutive phosphorylation in PetD (T4 to E4) markedly altered phosphorylation patterns of STT7-dependent substrates.

To assess functional consequences of the alterations in the engineered strains, we measured photosynthetic electron transfer rates (ETR) using electrochromic shift (ECS) signals. In

photoautotrophic cells, neither T4A nor T4E were different from WT which suggests that T4

does not impact ETR directly (Figs. 3a, 3b). However, ETR in ΔN slowed significantly 30–50 ms

after light exposure in cells grown under oxic conditions (Fig. 3a). Under anoxic conditions,

which favour CEF, ETR increased initially but reached a lower steady state, with the initial boost

notably reduced in ΔN (Fig. 3b). This suggests that the PetD N-terminus ensures ETR via b_6f.

Redox kinetics of P700 were also monitored (Figs. 3c, 3d). In ΔN, P700 oxidized rapidly at light

onset, followed by slow re-reduction in the dark, indicating donor-side limitation under both oxic

and anoxic conditions. Steady-state ETR (Figs. 3a, 3b) were established in the presence of a

substantial P700 acceptor-side limitation, except in ΔN (Figs. 3c, 3d). This donor-side limitation

in ΔN persisted even with the H^+/K^+ exchanger nigericin (Extended Data Figs. 7a, 7b), ruling out

a pH-dependent photosynthetic control mechanism. Addition of DCMU and nigericin resulted in

rapid P700 oxidation across all strains, slightly faster in ΔN (Extended Data Figs. 7c, 7d),

suggesting impaired b_6f function in LEF and CEF.

To confirm whether these ETR and P700 redox kinetics differences arose from b_6f dysfunction,

we assessed b_6f electrogenicity and redox kinetics (Extended Data Fig. 8, Fig. 4). ECS rise from

Q-cycle activity was impaired only in ΔN, despite the presence of heme c_i (Extended Data Figs. 8

and 9), the terminal electron acceptor in the low-potential chain [30]. Under oxic conditions, WT b-

hemes showed a transient reduction (~10 ms) before net oxidation (~16 ms half-life), while

cytochrome-f reduction (~6 ms half-life) began ~1 ms post-flash (Fig. 4a). In ΔN, the apparent b-

heme reduction was enhanced because their oxidation slowed ~20-fold (Fig. 4b). This coincided

with a ~10-fold slowdown in cytochrome-f reduction (Fig. 4b), indicating Qi-site impairment

affecting the Qo-site as has been shown in b_6f [3,31] and the related cytochrome-bc_1 complex [32,33].

Under anoxic conditions, WT displayed flat b-heme signals in the first 10 ms, followed by

accelerated oxidation (Fig. 4c), consistent with prior findings [3,31]. In ΔN, a redox-inactive low-

potential chain caused a ~25-fold slowdown in the high-potential chain, as reflected in

cytochrome-f reduction (cf. Figs. 4c, 4d).

This slowdown of electron transfer within b_6f very likely explains the significantly diminished

ETR and enhanced P700 donor side limitation (Fig. 3) as well as the strong impact on

electrogenicity of the complex when the N-terminus in PetD is truncated (Extended Data Fig. 8).

The reason remains to be determined but recent cryo-EM structures of spinach b_6f at 1.9 Å and

2.2 Å resolution have revealed that PQ does not make direct contact with the heme c_i at the Qi-

site [34]. Instead, the Qi-site is connected to water channels that link to the aqueous exterior of the

enzyme, suggesting that binding of PQ at the Qi-site displaces water through these channels. It

has been proposed that, upon entering the Qi-site, PQ pushes water molecules out of the cavity in

a piston-like fashion. The only remaining water molecule, coined wat1 [34], is crucial for allowing

the quinone ring to adopt an orientation that facilitates both electron and proton transfer. The

binding of wat1 is guided by PetD D35 which is believed to donate H^+ to wat1 since it is

surrounded by several polar residues.

The PetD N-terminus we studied is exposed to the aqueous phase, contains two lysine residues,

and is positioned near PetD D35. Our findings suggest that the ΔN truncation could influence the

entry and exit function of the water channel and/or affect the functional properties of PetD D35

itself. This could, in turn, impair electron injection and ejection from the low-potential chain,

potentially slowing down electron transfer through the b_6f as observed in the kinetic data (Fig. 4,

Extended data Fig. 8). It may also explain why STT7 kinase is not activated in the ΔN mutant

(Fig. 2b) as the turnover of the Qo-site was significantly slowed. In line, it has been previously

described that a strong impairment in PQH_2 oxidation at the Qo-site of b_6f prevents activation of

STT7 [35].

Intriguingly, in the ΔN and T4E strains, STT7 autophosphorylation at S569 and T571 was

significantly reduced. The failure to modulate these novel phospho-sites may be mechanistically

linked to the loss of STT7 kinase activity. Thus, our data highlight the importance of STT7-

dependent phosphorylation, which is negatively affected by the phosphomimic T4E mutation,

while the negative charge at PetD position 4 does not influence b_6f function. Notably, the failure

of STT7 to operate in ΔN, did not lead to the phosphorylation of PSBD T2, as seen in *stt7*

(Extended data Fig. 2 and [19]) and PetD T4E (Fig. 2). In some instances, such as PSBR, the T4A

showed increased phosphorylation, implying that PetD T4 phosphorylation may function as a

feedback regulator during STT7 activation. The dependence of STT7 function on PetD T4

phosphorylation reveals a novel feedback regulation mechanism, complementing previously

described STT7 autophosphorylation pathways involving other b_6f structural elements in stromal

proximity [17,18].

In conclusion, we found that the N-terminal region of PetD is essential for cytochrome b_6f

function. We further conclude that STT7-dependent phosphorylation of PetD T4 regulates STT7

kinase activity, establishing a feedback loop that adds a new layer to the intricate regulation of b_6f

and STT7 kinase. This feedback loop transiently reduces STT7-dependent phosphorylation via

the phosphorylation at PetD T4, which coincides with increasing PSBD T2 PSII core

phosphorylation, as previously described for vascular plants [36] and algae [37]. As a result, PetD T4

phosphorylation mediates a balance between STT7-dependent ST and other processes, as

indicated by PSBD T2 PSII core phosphorylation.

Methods

WT Plasmid construction

The *petD/trnR1* region (481 bp upstream and 1032 bp downstream of *petD*) was amplified from WT chloroplast DNA using restriction site-containing primer 1 (*SalI*) and 2 (*EcoRI*; see Extended Data Tab. 2). The *SalI/EcoRI* digested PCR product was ligated into a double-digested pUC18 cloning vector and propagated as pUC18-3'-*petD*-5' in NEB 5-alpha *E. coli* cells using ampicillin selection. In the next step, the aminoglycoside adenyl transferase (aadA) spectinomycin resistance cassette [38] was amplified from a pUC18-aadA vector using primers 3 and 4, each containing a *SpeI* restriction site. The cloned *petD/trnR1* region contains a single *SpeI* site into which the aadA cassette was cloned. To do so, intermediate vectors were created. First (pUC18-int1), the *SalI/SpeI* fragment from pUC18-3'-*petD*-5' (1.46 kb) was ligated with the *SpeI* digested aadA PCR fragment, followed by PCR (primers 1 and 5) and blunt ligation of the ~3 kb product into *SmaI* digested empty pUC18 vector. Second (pUC18-int2), the *EcoRI/SpeI* fragment from pUC18-3'-*petD*-5' (0.51 kb) was ligated with the *SpeI* digested aadA PCR fragment, followed by PCR (primers 2 and 6), *XmaI/EcoRI* digest of the ~2 kb product, and ligation into an empty pUC18 vector. The 2.4 kb fragment of *BamHI/BlpI* digested pUC18-int1 was ligated with the 3.7 kb fragment of *BamHI/BlpI* digested pUC18-int2 to yield pUC18-3'-*petD*-aadA-5'

Point Mutation and Truncation Plasmid Construction

To generate T4E and T4A mutation as well as the *petD* truncation ΔN (lacking underlined N-terminal amino acids M<u>SVTKK</u>PD…), primers 8-11 and 13 were phosphorylated. Using pUC18-3'-*petD*-aadA-5' as a template, five independent PCRs were done with the following primers: 1 and 8 (I), 1 and 9 (II), 1 and 10 (III), 11 and 12 (IV) and 12 and 13 (V). Ligated PCR fragments I and IV served as templates for the T4E PCR (primers 1 and 12) which was further digested (*PacI/SalI*) for ~1.36 kb insert swapping and mutant introduction into pUC18-3'-*petD*-aadA-5' (ligation with ~4.78 kb backbone). In a similar fashion, for T4A PCR (primers 1 and 12) PCR fragments II and IV served as templates. For ΔN PCR (primers 1 and 12), fragments III and V served as templates.

Biolistic Transformation of WT Strain

Mutant strains were generated through chloroplast transformation using plasmids containing the modified *petD* gene version along with an *aadA* cassette to confer spectinomycin resistance (150 µg/ml). The recipient wild-type (WT) strain carries a six-histidine tag at the C-terminus of PetA [20]. Spectinomycin resistant clones were selected and replated several times to obtain homoplastic strains (for sequencing primers, see Extended Data Tab. X).

Growth conditions and treatments

Cells were maintained at 50 µmol photons m^{-2} s^{-1} at 23°C on agar plates of Tris-acetate-

phosphate (TAP) medium or Tris-minimal medium (TP) in the absence of acetate [39] by bubbling

sterile air under a 16-h light/8-h dark regime. For state transition treatment, TAP-grown

synchronized cultures were freshly diluted one day before the experiment and, upon harvesting

(4000 g, 5 min, 23°C), adjusted to 5 µg chlorophyll ml^{-1}. Aliquots were either kept for 45 min in

aerobic conditions by shaking in 40 µmol photons m^{-2} s^{-1} light upon adding 10 µM 3-(3,4-

Dichlorophenyl)-1,1-dimethylurea (DCMU; State 1), or were given anoxic treatment in the dark

using a glucose oxidase/catalase cocktail (State 2) [40]. For proteomic analyses of $stt7$ experiments,

TAP grown cells were shifted to State 2 conditions for 30 min and samples for phospho-

proteomic analyses were collected. The strains were checked for photoautotrophic growth for one

week by spotting 20 µl of 1×10^6 cells/ml on TP agar plates at 25°C kept under 40 µmol photons

m^{-2} s^{-1} (16 h light/8 h dark) and continuous 500 µmol photons m^{-2} s^{-1}.

Liquid TP cultures were harvested for optical spectroscopy and Ficol (20% w/v) was added to the

TP resuspension medium.

Biochemical methods

Liquid cultures of WT, T4A, T4E, and ΔN were grown in TAP medium and freshly diluted one

day before harvesting. The cells were pelleted and mixed in a buffer containing 0.2 M dichloro-

diphenyl-trichloroethane, 0.2 M Na$_2$CO$_3$, 10 mM NaF, and protease inhibitors (0.2 mM

phenylmethylsulfonyl fluoride, 1 mM benzamidine, 20 mM amino caproic acid) to be analyzed

via SDS-PAGE [41]. For heme staining the b_6f was isolated using the standard protocol [42] to

analyze their purity via SDS-PAGE. The high-spin hemes were stained in gel using 3,3',5,5'-

tetramethylbenzidine [43].

Generation of $stt7$ mutants with CRISPR-CAS9

Chlamydomonas strain CC-125 ($nit1$; $nit2$; $mt+$) was grown in tris-acetate-phosphate (TAP)

media under continuous illumination of 50 µmol photons m^{-2} s^{-1} at 22 °C. Single guide RNAs

(sgRNAs) were designed by CHOPCHOP using version 5.6 of the $Chlamydomonas$

$reinhardtii$ genome, and 5'-CTCCTGTAGACCGCTCAATGCGG was selected as the targeting

sequence, and sgRNAs were bought will all accessory parts to make it functional (IDT, USA).

Ribonucleic protein (RNP) complexes were prepared by duplexing the sgRNA (23% volume per

volume, v/v; IDT) and Cas-9 (19% v/v; IDT) within a duplex buffer (58% v/v; IDT) and

incubated at 37°C for 25 to 30 min. After treatment with autolysin for 3 hours without shaking,

cells were washed with TAP media supplemented with 40 mM of sucrose, resuspended with the

same supplemented media to approximately 2 x 10^8 cells mL^{-1}, and electroporated in the presence

of a hygromycin resistance DNA cassette (2 µg mL^{-1}) and the 37°C-incubated RNP mixture.

After 10 min at room temperature, cells were resuspended in 10 mL of TAP media supplemented

with 40 mM of sucrose and left without shaking overnight in dim light (10-20 µmol photons m^{-2} s^{-1})

^{2} s^{-1}) at 22°C. Hygromycin resistant transformants were selected on solid TAP media (2% Agar

w/v) containing hygromycin (20 µg mL^{-1}). sgRNA targeted regions were amplified via

polymerase chain reaction (PCR) to check for full or partial insertion of the hygromycin

resistance cassette at the target site using the following primers: STT7 F/R (STT7F: 5'-

GCACGAACCAAGACACACATAG-3'; STT7 R: 5'-GTAGACGATGTCACCGCACTT3')

with an annealing temperature of 62.6°C (Extended Data Fig. 6). Impaired accumulation of the

STT7 protein was then verified by immunodetection (Agrisera, AS15 3080).

Sample preparation for mass spectrometry

Protein isolation from cell pellets was carried out as described [37]. Reduction/alkylation of

cysteines and tryptic digestion of proteins (100 µg per sample) was performed following the SP4

protocol [44] . After digestion, peptides were desalted using PurePep H50 SPE Spin columns

(Affinisep). The peptide samples were divided into two aliquots: 5 µg for whole proteome

analysis and 95 µg for phosphopeptide enrichment. Both aliquots were dried using vacuum

centrifugation before further processing.

To enrich phosphopeptides, titanium dioxide (TiO2) beads with 5 µm particle size (Titansphere,

GL Sciences) in a metal oxide affinity chromatography (MOAC) procedure were used. TiO2 (1

mg per sample) was activated once with acetonitrile and equilibrated three times with loading

buffer (LB; 80% (v/v) acetonitrile/5% (v/v) TFA/1 M glycolic acid). Loading buffer was added to

create a 10% (w/v) TiO2 suspension. Peptide samples were dissolved in 50 µl LB, combined with

10 µl TiO2 suspension, and incubated for 30 minutes at 25°C in an Eppendorf Thermomixer at

1,200 rpm). The mixture was transferred to SDB-XC STAGE tips prepared in-house [45]. Unless

stated otherwise, all subsequent washing and elution steps were performed using 60 µl buffer

volumes with in-between centrifugation at 2000 × g at room temperature. The buffer was

removed by centrifugation and the TiO2 beads were washed twice with LB and once with 1%

(v/v) acetonitrile/0.1% (v/v) TFA (W1). Non-phosphorylated peptides that were removed from

the beads and subsequently bound to the SDB-XC membrane plug were removed by one wash

with 80% (v/v) acetonitrile/0.1% (v/v) TFA (W2). The phosphopeptides were eluted from the

TiO2 beads onto the SDB-XC material using 160 µl of pH 11 buffer (0.4 M sodium hydrogen

phosphate/NaOH with 1% acetonitrile). After washing once with 1% acetonitrile/0.1% TFA, the

phosphopeptides were sequentially eluted in three fractions using increasing concentrations of

acetonitrile (7.5%, 20%, and 60%) in 100 mM ammonium formate (pH10). For samples from the

STT7 experiment, only one fraction (60% acetonitrile in 100 mM ammonium formate (pH10))

was collected. The strongly bound phosphopeptides were eluted in two steps: first from the TiO2

beads using 5% ammonia, then from the SDB-CX membrane using 60% acetonitrile in 100 mM

ammonium formate at pH 10. All fractions were dried by vacuum centrifugation and stored

at -80°C until MS analysis.

Mass spectrometry

Whole proteome analysis

Dried peptide samples were reconstituted in 5 μl of 2% (v/v) acetonitrile/0.05% (v/v) TFA in

LC/MS grade water. Samples were analyzed on an LC-MS/MS system consisting of an Ultimate

3000 NanoLC (Thermo Fisher Scientific) coupled via a Nanospray Flex ion source (Thermo

Fisher Scientific) to a Q Exactive Plus mass spectrometer (Thermo Fisher Scientific). Peptides

were concentrated on a trap column (Acclaim Pepmap C18, 5 × 0.3 mm, 3 μm particle size,

Thermo Scientific) for 3 min using 4% (v/v) acetonitrile/0.05% (v/v) in LC/MS grade water at a

flow rate of 10 μl/min. The trap column was operated in back-flush mode, allowing the transfer

of peptides on a reversed-phase column (PepSep Fifty, 500 × 0.075 mm, 1.9 μm particle size,

Bruker) for chromatographic separation. The eluents used were 0.1% (v/v) formic acid in LC/MS

grade water (A) and 0.1% (v/v) formic acid/80% (v/v) acetonitrile in LC/MS grade water (B).

The gradient was programmed as follows: 2.5% to 5% B over 5 min, 5% to 17.5% B over 47

min, 17.5% to 40% B over 105 min, 40% to 99% B over 10 min, and 99% B for 20 min. Flow

rate was 250 nl/min. The mass spectrometer was operated in data-dependent acquisition (DDA)

mod, alternating between one MS1 full scan and up to 12 MS2 scans. Full scans were acquired

with the following settings: AGC target 3e6, MS1 resolution 70000, maximum injection time 50

ms, scan range: 350–1400 m/z. Settings for MS were: AGC target 5e4, resolution 17500, MS2

maximum injection time 80 ms, intensity threshold 1e4, normalized collision energy 27 (HCD)).

Dynamic exclusion was set to 'auto', assuming a chromatographic peak width (FWHM) of 60 s.

Phosphoproteomic analysis

Prior to analysis, dried peptide samples were reconstituted in 5 μl of 2% (v/v) acetonitrile/0.05%

(v/v) TFA in LC/MS grade water. LC-MS/MS system, flow rate and eluent compositions were

the same as described above. The gradient for peptide separation was as follows: 2.5% to 5% B

over 5 min, 5% to 40% B over 92 min, 40% to 99% B over 10 min, and 99% B for 20 min.

The mass spectrometer used the same DDA mode settings as the whole proteome analysis, with

two changes: MS2 maximum injection time was increased to 120 ms, and dynamic exclusion was

set to 45 s.

MS data analysis

For quantitative phosphoproteome analysis, mass spectrometry raw data from nonenriched and enriched samples were searched using Fragpipe [46,47] against a concatenated, non-redundant database containing *Chlamydomonas reinhardtii* protein sequences from v5.6 and v6.1 gene models (Phytozome 13, phytozome-next.jgi.doe.gov). In addition, the polypeptide sequences of mutated/tagged proteins were added to the database. Default 'LFQ-MBR' workflow settings were applied to data from whole proteome samples, while the 'LFQ-phospho' workflow was used for phosphopeptide samples. A false discovery rate of 1% was applied at both peptide and protein levels.

Fragpipe output files containing peptide/protein identifications and quantitative data ('combined_modified_peptide.tsv' from enriched samples and 'combined_protein.tsv' from whole proteome samples) were reformatted using a custom Python script to ensure compatibility with Phospho-Analyst (v. 1.0.2), which was used for differential expression analysis [48]: Missing values were imputed using the 'MinProb' function, and variance stabilizing transformation was applied to normalize phosphosite intensities. Since the whole proteome LFQ data was already normalized in Fragpipe, no additional normalization steps were needed. Phospho-Analyst automatically corrected phosphosite abundance changes for underlying protein levels. A phosphorylation site localization probability threshold of 0.33 was applied, and p-values were adjusted using the Benjamini-Hochberg method. Predicted protein localizations are based on recently published data (PB-Chlamy)[49].

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via the PRIDE partner repository [50] with the dataset identifier PXD060640.

Optical spectroscopy

The flash-induced b_6f measurements were carried out by using a Joliot-type spectrophotometer (JTS-150, SpectroLogiX) equipped with the interference filters and deconvolution procedures as reported previously [3]. Cells were adapted to alternating 7.5 s darkness and 5 s actinic light (AL; 550 µmol photons m^{-2} s^{-1} peaking at 630 nm). Sub-saturating 6-ns flashes (Q-switched Nd:YAG, Continuum) were fired at 3 s darkness, hitting ~40% of PSI. Three averaged baseline spectra (in the absence of a flash) were subtracted from each flash measurement at the b_6f-specific wavelengths. For multiple turnover conditions, previously described conditions and protocols used cells which were adapted to alternating 2.5 s AL and dark. Briefly, ETR were obtained in the presence of LEF but referred to PSI charge separation activity. This stems from separate ECS calibration measurements that allowed, via saturating single-turnover flashes, for the conversion

of optical changes (obtained as $\Delta I/I$ at 520 nm - 546nm) into the ECS signals corresponding to 1

PSI charge separation [51]. The ETR protocol was based on a Dark Interval Relaxation Kinetics

approach [52,53]. P700 measurements during 2.5 s AL and dark relied on the Klughammer &

Schreiber method [54].

References

1   Malone, L. A., Proctor, M. S., Hitchcock, A., Hunter, C. N. & Johnson, M. P.
Cytochrome b(6)f - Orchestrator of photosynthetic electron transfer. *Biochim Biophys*
Acta Bioenerg **1862**, 148380 (2021). https://doi.org:10.1016/j.bbabio.2021.148380

2   Mitchell, P. The protonmotive Q cycle: a general formulation. *FEBS Lett* **59**, 137-139
(1975). https://doi.org:10.1016/0014-5793(75)80359-0

3   Buchert, F., Mosebach, L., Gabelein, P. & Hippler, M. PGR5 is required for efficient Q
cycle in the cytochrome b6f complex during cyclic electron flow. *Biochem J* **477**, 1631-
1650 (2020). https://doi.org:10.1042/BCJ20190914

4   Rumberg, B. & Siggel, U. Ph Changes in Inner Phase of Thylakoids during
Photosynthesis. *Naturwissenschaften* **56**, 130-+ (1969). https://doi.org:Doi
10.1007/Bf00601025

5   Nishio, J. N. & Whitmarsh, J. Dissipation of the Proton Electrochemical Potential in
Intact Chloroplasts (II. The pH Gradient Monitored by Cytochrome f Reduction Kinetics).
Plant Physiol **101**, 89-96 (1993).

6   Li, X. P. *et al.* A pigment-binding protein essential for regulation of photosynthetic light
harvesting. *Nature* **403**, 391-395 (2000).

7   Peers, G. *et al.* An ancient light-harvesting protein is critical for the regulation of algal
photosynthesis. *Nature* **462**, 518-521 (2009). https://doi.org:10.1038/nature08587

8   Girolomoni, L. *et al.* LHCSR3 is a nonphotochemical quencher of both photosystems in.
P Natl Acad Sci USA **116**, 4212-4217 (2019). https://doi.org:10.1073/pnas.1809812116

9   Allorent, G. *et al.* A dual strategy to cope with high light in Chlamydomonas reinhardtii.
Plant Cell **25**, 545-557 (2013). https://doi.org:10.1105/tpc.112.108274

10   Depege, N., Bellafiore, S. & Rochaix, J. D. Role of chloroplast protein kinase Stt7 in
LHCII phosphorylation and state transition in Chlamydomonas. *Science* **299**, 1572-1575
(2003). https://doi.org:10.1126/science.1081397

11   Rochaix, J. D. Regulation and dynamics of the light-harvesting system. *Annu Rev Plant*
Biol **65**, 287-309 (2014). https://doi.org:10.1146/annurev-arplant-050213-040226

12   Cariti, F. *et al.* Regulation of Light Harvesting in Chlamydomonas reinhardtii Two
Protein Phosphatases Are Involved in State Transitions. *Plant Physiol* **183**, 1749-1764
(2020). https://doi.org:10.1104/pp.20.00384

13   Lemeille, S. *et al.* Analysis of the chloroplast protein kinase Stt7 during state transitions.
PLoS Biol **7**, e45 (2009). https://doi.org:10.1371/journal.pbio.1000045

14   Shapiguzov, A. *et al.* Activation of the Stt7/STN7 Kinase through Dynamic Interactions
with the Cytochrome b_6f Complex. *Plant Physiol* **171**, 82-92 (2016).
https://doi.org:10.1104/pp.15.01893

15   Singh, S. K. *et al.* Trans-membrane signaling in photosynthetic state transitions: Redox-
and structure-dependent interaction *in vitro* between STT7 kinase and the cytochrome b_6f
complex. *J Biol Chem* **291**, 21740-21750 (2016).
https://doi.org:10.1074/jbc.M116.732545

16   Wu, J. *et al.* Functional redox links between Lumen Thiol Oxidoreductase1 and
Serine/Threonine-protein kinase STN7. *Plant Physiol* (2021).
https://doi.org:10.1093/plphys/kiab091
17   Dumas, L. *et al.* A stromal region of cytochrome b_6f subunit IV is involved in the
activation of the Stt7 kinase in *Chlamydomonas*. *Proc Natl Acad Sci U S A* **114**, 12063-
12068 (2017). https://doi.org:10.1073/pnas.1713343114
18   Riche, A. *et al.* The stromal side of the cytochrome b6f complex regulates state
transitions. *Plant Cell* **36**, 4234-4244 (2024). https://doi.org:10.1093/plcell/koae190
19   Bergner, S. V. *et al.* STATE TRANSITION7-dependent phosphorylation is modulated by
changing environmental conditions, and its absence triggers remodeling of photosynthetic
protein complexes. *Plant Physiol* **168**, 615-634 (2015).
https://doi.org:10.1104/pp.15.00072
20   Choquet, Y., Zito, F., Wostrikoff, K. & Wollman, F. A. Cytochrome *f* translation in
Chlamydomonas chloroplast is autoregulated by its carboxyl-terminal domain. *Plant Cell*
15, 1443-1454 (2003).
21   Takahashi, H. *et al.* PETO Interacts with Other Effectors of Cyclic Electron Flow in
Chlamydomonas. *Mol Plant* **9**, 558-568 (2016).
https://doi.org:10.1016/j.molp.2015.12.017
22   Dinc, E. *et al.* LHCSR1 induces a fast and reversible pH-dependent fluorescence
quenching in LHCII in Chlamydomonas reinhardtii cells. *Proc Natl Acad Sci U S A* **113**,
7673-7678 (2016). https://doi.org:10.1073/pnas.1605380113
23   Wollman, F.-A. & Lemaire, C. Studies on kinase-controlled state transitions in
Photosystem II and b_6f mutants from *Chlamydomonas reinhardtii* which lack quinone-
binding proteins. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **933**, 85-94 (1988).
https://doi.org:https://doi.org/10.1016/0005-2728(88)90058-8
24   Wollman, F.-A. & Bennoun, P. A new chlorophyll-protein complex related to
Photosystem I in *Chlamydomonas reinhardtii*. *Biochim Biophys Acta* **680**, 352-360
(1982).
25   Pfalz, J., Liere, K., Kandlbinder, A., Dietz, K. J. & Oelmuller, R. pTAC2, -6, and -12 are
components of the transcriptionally active plastid chromosome that are required for
plastid gene expression. *Plant Cell* **18**, 176-197 (2006).
https://doi.org:10.1105/tpc.105.036392
26   Ingelsson, B. & Vener, A. V. Phosphoproteomics of Arabidopsis chloroplasts reveals
involvement of the STN7 kinase in phosphorylation of nucleoid protein pTAC16. *FEBS*
Lett **586**, 1265-1271 (2012). https://doi.org:10.1016/j.febslet.2012.03.061
27   Scholz, M. *et al.* Light-dependent N-terminal phosphorylation of LHCSR3 and LHCB4
are interlinked in *Chlamydomonas reinhardtii*. *Plant J* **99**, 877-894 (2019).
https://doi.org:10.1111/tpj.14368
28   Lemeille, S., Turkina, M. V., Vener, A. V. & Rochaix, J. D. Stt7-dependent
phosphorylation during state transitions in the green alga Chlamydomonas reinhardtii.
Molecular & cellular proteomics : MCP **9**, 1281-1295 (2010).
https://doi.org:10.1074/mcp.M000020-MCP201
29   Pan, X. *et al.* Structural basis of LhcbM5-mediated state transitions in green algae. *Nat*
Plants **7**, 1119-1131 (2021). https://doi.org:10.1038/s41477-021-00960-8
30   Alric, J., Pierre, Y., Picot, D., Lavergne, J. & Rappaport, F. Spectral and redox
characterization of the heme c_i of the cytochrome b_6f complex. *Proc Natl Acad Sci U S A*
102, 15860-15865 (2005). https://doi.org:10.1073/pnas.0508102102

31   Buchert, F., Scholz, M. & Hippler, M. Electron transfer via cytochrome b_6f complex
displays sensitivity to antimycin A upon STT7 kinase activation. *Biochem J* **479**, 111-127
(2022). https://doi.org:10.1042/BCJ20210802

32   Valkova-Valchanova, M., Darrouzet, E., Moomaw, C. R., Slaughter, C. A. & Daldal, F.
Proteolytic Cleavage of the Fe−S Subunit Hinge Region of Rhodobacter capsulatus bc1
Complex: Effects of Inhibitors and Mutations. *Biochemistry* **39**, 15484-15492 (2000).
https://doi.org:10.1021/bi000751d

33   Cooley, J. W., Lee, D.-W. & Daldal, F. Across Membrane Communication between the
Qo and Qi Active Sites of Cytochrome bc1. *Biochemistry* **48**, 1888-1899 (2009).
https://doi.org:10.1021/bi802216h

34   Pintscher, S. *et al.* Molecular basis of plastoquinone reduction in plant cytochrome b(6)f.
Nat Plants (2024). https://doi.org:10.1038/s41477-024-01804-x

35   Zito, F. *et al.* The Qo site of cytochrome b_6f complexes controls the activation of the
LHCII kinase. *EMBO J* **18**, 2961-2969 (1999). https://doi.org:10.1093/emboj/18.11.2961

36   Mekala, N. R., Suorsa, M., Rantala, M., Aro, E. M. & Tikkanen, M. Plants Actively
Avoid State Transitions upon Changes in Light Intensity: Role of Light-Harvesting
Complex II Protein Dephosphorylation in High Light. *Plant physiology* **168**, 721-734
(2015). https://doi.org:10.1104/pp.15.00488

37   Younas, M., Scholz, M., Marchetti, G. M. & Hippler, M. Remodeling of algal
photosystem I through phosphorylation. *Biosci Rep* **43** (2023).
https://doi.org:10.1042/BSR20220369

38   Fu, H. Y. *et al.* Redesigning the Q(A) binding site of Photosystem II allows reduction of
exogenous quinones. *Nat Commun* **8**, 15274 (2017).
https://doi.org:10.1038/ncomms15274

39   Gorman, D. S. & Levine, R. P. Cytochrome *f* and plastocyanin: their sequence in the
photosynthetic electron transport chain of *Chlamydomonas reinhardtii. Proc Natl Acad*
Sci U S A **54**, 1665-1669 (1965).

40   Buchert, F., Mosebach, L., Gäbelein, P. & Hippler, M. PGR5 is required for efficient Q
cycle in the cytochrome b_6f complex during cyclic electron flow. *Biochemical Journal*
477, 1631-1650 (2020). https://doi.org:10.1042/bcj20190914

41   Schägger, H. & Von Jagow, G. Tricine-sodium dodecyl sulfate-polyacrylamide gel
electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Analytical*
biochemistry **166**, 368-379 (1987).

42   Stroebel, D., Choquet, Y., Popot, J. L. & Picot, D. An atypical haem in the cytochrome b_6f
complex. *Nature* **426**, 413-418 (2003). https://doi.org:10.1038/nature02155

43   Thomas, P. E., Ryan, D. & Levin, W. An improved staining procedure for the detection of
the peroxidase activity of cytochrome P-450 on sodium dodecyl sulfate polyacrylamide
gels. *Anal Biochem* **75**, 168-176 (1976). https://doi.org:10.1016/0003-2697(76)90067-1

44   Johnston, H. E. *et al.* Solvent Precipitation SP3 (SP4) Enhances Recovery for Proteomics
Sample Preparation without Magnetic Beads. *Anal Chem* **94**, 10320-10328 (2022).
https://doi.org:10.1021/acs.analchem.1c04200

45   Rappsilber, J., Mann, M. & Ishihama, Y. Protocol for micro-purification, enrichment, pre-
fractionation and storage of peptides for proteomics using StageTips. *Nature Protocols* **2**,
1896-1906 (2007). https://doi.org:10.1038/nprot.2007.261

46   Kong, A. T., Leprevost, F. V., Avtonomov, D. M., Mellacheruvu, D. & Nesvizhskii, A. I.
MSFragger: ultrafast and comprehensive peptide identification in mass spectrometry–

based proteomics. *Nature Methods* **14**, 513-520 (2017). https://doi.org/10.1038/nmeth.4256

Yu, F. *et al.* Analysis of DIA proteomics data using MSFragger-DIA and FragPipe computational platform. *Nat Commun* **14**, 4154 (2023). https://doi.org/10.1038/s41467-023-39869-5

Zhang, H. *et al.* Phospho-Analyst: An Interactive, Easy-to-Use Web Platform To Analyze Quantitative Phosphoproteomics Data. *J Proteome Res* **22**, 2890-2899 (2023). https://doi.org/10.1021/acs.jproteome.3c00186

Wang, L. *et al.* A chloroplast protein atlas reveals punctate structures and spatial organization of biosynthetic pathways. *Cell* **186**, 3499-3518.e3414 (2023). https://doi.org:https://doi.org/10.1016/j.cell.2023.06.008

Perez-Riverol, Y. *et al.* The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic acids research* **47**, D442-D450 (2019).

Bailleul, B., Cardol, P., Breyton, C. & Finazzi, G. Electrochromism: a useful probe to study algal photosynthesis. *Photosynthesis Research* **106**, 179-189 (2010). https://doi.org/10.1007/s11120-010-9579-z

Sacksteder, C. A. & Kramer, D. M. Dark-interval relaxation kinetics (DIRK) of absorbance changes as a quantitative probe of steady-state electron transfer. *Photosynth Res* **66**, 145 (2000). https://doi.org/10.1023/a:1010785912271

Joliot, P. & Joliot, A. Cyclic electron transfer in plant leaf. *Proc Natl Acad Sci U S A* **99**, 10209-10214 (2002).

Klughammer, C. & Schreiber, U. An improved method, using saturating light-pulses, for the determination of photosystem-I quantum yield via P700$^+$-absorbance changes at 830nm. *Planta* **192**, 261-268 (1994).

Acknowledgements

M.H. (507704013) and F. B. (507704013) acknowledge funding by the Deutsche Forschungsgemeinschaft (DFG FOR 5573/1). M.H. acknowledges the RECTOR program in association with the University of Okayama, Japan. F. B. acknowledges Deutsche Forschungsgemeinschaft 461765884. Y.M. acknowledges the Alexander von Humboldt Foundation for a Research Fellowship for postdocs (1219125). A.B. acknowledges the support of Carnegie Science.

Author contributions

M.H. and F.B. designed the project. A.Z. and F.B. did genetic engineering and kinetic measurements and analyses. S.B. contributed to mutant analyses. C.S. and A.B. produced the *stt7* KO strains. M.S. did the mass spectrometric measurements and data evaluation. M.S. and M.H. analysed the mass spectrometric data. M.H wrote the manuscript with contributions from F.B., A.Z., A.B. and M.S. All authors contributed to the analysis and the final version of the manuscript.

Competing interests

566   The authors declare no competing interests.

Fig. 1. The relative antenna cross section is shown via chlorophyll fluorescence emission spectra at 77 K. These spectra were averaged ($N = 3 \pm$ SD) and normalized to photosystem II-attached antenna emission signals at ~687 nm. Panels a – d highlight the differences in antenna attached to photosystem I under State 1 (oxic) and State 2 (anoxic), contributing to ~712 nm emissions.

574
Fig. 2. Volcano plots visualizing effect of PetD mutations on protein phosphorylation.
Phosphorylation sites with FDR-adjusted p-value < 0.05 and |log₂ fold change| > 2 are highlighted
in blue, with threshold boundaries shown as dashed lines. Data points are shown in orange instead
of blue when all values for one strain in the fold-change calculation are imputed due to completely
missing data. Green halos indicate chloroplast-encoded proteins and nuclear-encoded proteins with
confirmed or predicted (PB-Chlamy) chloroplast localization. Phosphorylation log₂ fold changes
are normalized to corresponding protein abundance levels to account for changes in protein
expression. Numbers at he end of data labels indicate peptide multiplicity (total phosphorylations
per peptide, e.g. PRPS10_S42_2 = PRPS10 with phosphorylated S42 and one additional
phosphorylation on the same peptide). A) WT vs. ΔN. B) WT vs. T4A. C) WT vs. T4E. D) T4A vs.
T4E.
586

Fig. 3. Photosynthetic activities of the cytochrome _b₆f_ variants are shown. Averaged kinetics are depicted ($N = 3 \pm$ SD). **a**, Electron transfer rates (ETR) developed over the 2.5-s illumination period under oxic conditions. The letters at the punctual rates indicate statistical significances (One-Way ANOVA/Tukey-HSD, $P < 0.05$) **b**, ETR developments are shown under anoxic conditions. **c**, Quantification of oxidized primary PSI donor ([P700⁺]) is plotted under oxic conditions. Red, yellow, and black bar represent actinic light intensities of 550, 3000, and 0 µmol photons m⁻² s⁻¹. **d**, The P700 kinetics are shown as in panel c but under anoxic conditions.

Fig. 4. Single turnover redox changes of the cytochrome b_6f hemes are shown. Averaged kinetics are depicted ($N = 3 \pm$ SD). WT (**a**, **c**) as well as the ΔN mutant (**b**, **d**) were measured under oxic (**a**, **b**) and anoxic conditions (**c**, **d**). The averaged half-times of cytochrome-f reduction and b-hemes oxidation are given in the panels' legends (n. d.; not determined).

602
603
604 **Extended Data Fig. 1. Graphical abstract of the cytochrome b_6f complex interplay with**
605 **STT7 kinase regulation.** The N-terminus of PetD (subunit-IV) has been investigated and the
606 focus was on T4. Previously known processes are included for reference. The drawing was
607 created with Biorender.

Extended Data Fig. 2. Volcano plots visualizing differential phosphorylation (A) and protein expression (B) between WT and *stt7* mutant. Proteins with FDR-adjusted p-value < 0.01 and |log₂ fold change| > 1 are highlighted in blue, with threshold boundaries shown as dashed lines. Data points are shown in orange instead of blue when all values for one strain in the fold-change calculation are imputed due to completely missing data. Green halos indicate chloroplast-encoded proteins and nuclear-encoded proteins with confirmed or predicted (PB-Chlamy) chloroplast localization.

609
610 **Extended Data Fig. 3. Growth analysis in varying light conditions and protein expression in**
611 **the mutants**. **a**. Spot tests on minimal media compare the growth under two light conditions (40
612 and 200 µmol photons m^{-2} s^{-1}). **b**. Protein expression of b_6f subunits in the engineered PetD strains.
613 The box plots visualize the log$_2$ protein LFQ intensities.

Extended Data Fig. 4. Volcano plots visualizing differential protein expression between WT and PetD mutants. Proteins with FDR-adjusted p-value < 0.01 and |log₂ fold change| > 1 are highlighted in blue, with threshold boundaries shown as dashed lines. Data points are shown in orange instead of blue when all values for one strain in the fold-change calculation are imputed due to completely missing data. Green halos indicate chloroplast-encoded proteins and nuclear-encoded proteins with confirmed or predicted (PB-Chlamy) chloroplast localization. A) WT vs. ΔN. B) WT vs. T4A. C) WT vs. T4E. D) T4A vs. T4E.

Extended Data Fig. 5. STT7 protein expression and selected autophosphorylation amounts in the engineered PetD strains. The box plots visualize $\log_2$ protein LFQ (A) and phosphosite intensities (B+C).

620
621
Extended Data Fig. 6. Generation of the *stt7* CRISPR/Cas mutant. a. The gene map of the
STT7 gene showing 5' UTR (green), numbered exons (blue) and 3'UTR (yellow) with predicted
cut site (vertical dotted line) and primers used to verify the genotype of the insertion (blue
arrows). **b.** Genetic characterization of the *stt7* mutant and its control (CC125) using PCR on
genomic DNA. **c.** Immunodetection of STT7 in the *stt7* mutant and its control (CC125).

Extended Data Fig. 7. Photosynthetic activities of the cytochrome *b₆f* variants are shown for P700. Averaged kinetics are depicted ($N = 3 \pm SD$) and the reference measurements are shown in main text Figs. 3c and 3d. The redox changes of primary PSI donor ([P700$^+$]) is plotted (**a**) under oxic and (**b**) anoxic conditions in the presence of 10 µM nigericin, acting as H$^+$/K$^+$ exchanger to attenuate ΔpH in favor of an electric field. Red, yellow, and black bar represent actinic light intensities of 550, 3000, and 0 µmol photons m^{-2} s^{-1}. **c**, The conditions were as in panel a but photosystem II was inhibited by hydroxyl amine (HA; 2 mM) and 3-(3,4-Dichlorophenyl)-1,1-dimethylurea (DCMU; 20 µM). **d**, The conditions were as in panel c except for the anoxic pre-treatment.

Extended Data Fig. 8. Laser flash-induced electrochromic shift (ECS) shows multiphasic kinetics that rely on activities of both photosystems, the cytochrome b_6f complex and ATP synthase. Averaged measurements are shown ($N=3 \pm SD$).

heme c (cyt *f*)

heme c_i (cyt b_6)

WT ΔN

40 kDa

35 kDa
25 kDa

15 kDa

671 **Extended Data Fig. 9. Heme staining of isolated *b6f* using the TMBZ method.** Heme c_i was
672 present in the ΔN mutant strain.

673 **Extended Data Tab. 2. List of primers used for plasmid construction and biolistic**
674 **transformation of the chloroplast genome.**
675

Oligo number in text	5'-3' oligo sequence
1	tatagtcgactgccactgacgtcccgtcagg
2	ttaagaattcctgccactgagctcccgtaagg
3	ttaatactagtggatcgcactctaccgattgagttac
4	ccaggactagtcttttacgaatacacatatgg
5	ccaggtctagacttttacgaatacacatatgg
6	ttaatcccggggggatcgcactctaccgattgagttac
7	gtaactcaatcggtagagtgcgatcc
8	caggttttttctcaactgacatatttatttatc
9	caggtttttttgcaactgacatatttatttatc
10	catatttatttatccgttaattttcaataaaac
11	atttaagcgatccagtttttaaaagc
12	ctcctaattaattaattcccattttatactg
13	cctgatttaagcgatccag