DateBack, a digital open access repository of archaeobotanical data for reconstructing date palm history

Besseiche, Margot[1], Chambraud, Elora[2], Dabrowski, Vladimir[2,3], Brandstätt, Elisa[2], Sabot, François[1], Bouchaud, Charlène[2+], Gros-Balthazard, Muriel*[1]

1. DIADE, Univ Montpellier, CIRAD, IRD, Montpellier, France.

2. BioArch Laboratory - UMR 7209, National Center of Scientific Research (CNRS), National Museum of Natural History (MNHN), Paris, France.

3.Orient et Méditerranée - UMR 8167, Université Paris 1 Panthéon-Sorbonne, Sorbonne Université, CNRS, EPHE, Collège de France, Paris, France.

 * Correspondence: charlene.bouchaud@mnhn.fr and muriel.gros-balthazard@ird.fr.

ORCID ID

Margot Besseiche: https://orcid.org/0009-0007-2353-1612

Elisa Brandstätt: https://orcid.org/0009-0002-3351-9223

Vladimir Dabrowski: https://orcid.org/0000-0002-7428-5702

Elora Chambraud: https://orcid.org/0009-0001-7363-1159

Francois Sabot: https://orcid.org/0000-0002-8522-7583

Charlène Bouchaud: https://orcid.org/0000-0002-1318-027X

Muriel Gros-Balthazard: https://orcid.org/0000-0002-2587-3946

Abstract

- The date palm (*Phoenix dactylifera* L.) has been a cornerstone of oasis agrosystems in West Asia and North Africa for millenia, yet the timeline and processes of its domestication and spread remain poorly documented. Archaeobotanical remains provide critical insights into its cultivation history, but no comprehensive review or dedicated platform currently exists to synthesize and analyze these data.
- To address this gap, we conducted an extensive literature review and developed *DateBack* (https://cloudapps.france-bioinformatique.fr/dateback), a digital open access repository of archaeobotanical remains of *Phoenix* spp. In its first version, *DateBack* records macroremains (e.g., seeds, fruits, stems, petioles) from prehistoric to Late Antiquity contexts in Southwest and South Asia.
- We assembled 140 entries from 99 archaeological sites across 110 references, along with a separate table of 55 radiocarbon-dated entries from 17 sites, refining chronological assessments. Most entries comprise charred seeds (55.7%), with a smaller proportion of charred vegetative parts or desiccated remains. Our findings highlight regional disparities in remain distribution, concentrated in the Levant and Gulf, while Saudi Arabia and southern Iran are underrepresented, and in data reliability, particularly dating resolution, complicating the reconstruction of date cultivation history. Our evidence shows that the earliest securely dated remains, indicating date consumption, appear in the Gulf around 5000 BCE, while cultivation emerges later, first in southern Mesopotamia and the northern Gulf in the 4th millennium BCE, then in eastern Arabia and the Indus Valley in the 3rd millennium BCE. The Levant presents challenges, with numerous presumed early finds but significant uncertainties, with secure evidence of cultivation only appearing by the late 2nd millennium BCE, as in northwestern Arabia, while in the southern Arabian Peninsula, it is not attested until the 1st millennium BCE.
- By compiling and standardizing archaeobotanical data, *DateBack* facilitates advanced spatiotemporal analyses and supports additional archaeobotany-based research including ancient DNA studies. Moreover, the platform is dynamic, scalable, and collaborative, enhancing data integration and refinement, with future expansions planned to include additional periods (Islamic era and beyond), geographic regions (North Africa), and new types of evidence, such as plant microremains and iconography.

Keywords

Agrobiodiversity; Archaeobotany; Date palm (*Phoenix dactylifera* L.); Domestication; Open access database; Radiocarbon dating; Spatiotemporal analysis.

Introduction

The date palm *(Phoenix dactylifera* L.) is a keystone species in oasis agrosystems of North Africa and West Asia, where it plays a crucial role in agricultural activities, human livelihoods, and cultural traditions. Cultivated in more than 30 countries today, the date palm yields over 10 million tons of sugar-rich dates annually, a figure that continues to grow (FAOSTAT, 2023). Beyond its role as a dietary staple, all parts of the plant have long been used for diverse purposes: leaves and fibers for making baskets and ropes, as well as the stem and petiole for building (as beam, roof) and fuel use, especially in wood-scarce environments (e.g., Mouton et al., 2021; Tengberg, 2012). Its canopy and surroundings further supports the cultivation of other crops, such as cereals, pulses, herbs, legumes, and fruit trees, in arid environments (Tengberg, 2012). The cultural and traditional significance of the date palm is recognized internationally, as reflected in the 2022 UNESCO inscription of its associated knowledge, skills, and practices on the Representative List of the Intangible Cultural Heritage of Humanity.

While the date palm's significance in agriculture and culture is well-established, critical aspects of its early history remain unclear, including its origins, routes and timing of spread, and its role in shaping oasis agriculture (reviewed in Gros-Balthazard & Flowers, 2021). *Phoenix dactylifera* belongs to the genus *Phoenix*, which comprises 13 other non-cultivated species (Barrow, 1998; Gros-Balthazard et al., 2021). Although genetic studies confirm that domestication stemmed from *P. dactylifera* itself (Pintaud et al., 2010; Gros-Balthazard et al., 2017), the prehistoric distribution of its ancestral populations remains poorly understood, hindering efforts to fully reconstruct its early interactions with human societies (Gros-Balthazard & Flowers, 2021).

Archaeological evidence points to date consumption as early as ~5000 BCE in the Gulf region, based on date seeds recovered from ancient deposits (Beech & Shepherd, 2001; Parker et al., 2011). By the late 4th millennium BCE, archaeobotanical and iconographic evidence indicate widespread date palm cultivation in Mesopotamia, leading some researchers to propose it as a domestication center (Tengberg, 2012; Zohary & Spiegel-Roy, 1975). Written records, appearing in the 3rd millennium BCE, further attest to its economic and cultural significance in the region. Alternatively, others suggest that fruit tree horticulture, including date palm, may have first developed in the Chalcolithic Levant (Zohary et al., 2012; Langgut & Sasi, 2023; Langgut, 2024).

In North Africa, date palm cultivation is attested later, with evidence from the 2nd millennium BCE in Egypt (Tengberg & Newton, 2016), early 1st mill. BCE in the central Sahara (Van der Veen, 1992; Kaczmarek et al., 2024), and only as late as medieval times in Morocco (Ros et al., 2024). While this pattern suggests a westward diffusion of date palm cultivation from West Asia across its historical range, spanning Morocco to Pakistan (Tengberg, 2012), genetic data challenge this straightforward narrative. Cultivated date palms are divided into two primary genepools — North African and West Asian (Zehdi et al., 2015; Hazzouri et al., 2015) — which some researchers interpret as evidence of two distinct domestication centers (Zehdi et al., 2015). North African date palms exhibit higher genetic diversity than would be expected from a simple diffusion model (Hazzouri et al., 2015; Gros-Balthazard et al., 2017). This unexpected diversity may partly be explained by inter-specific hybridization with *Phoenix theophrasti*, a wild relative currently restricted to Crete and Turkey, though the

timing and mechanisms of such hybridization events remain unclear (Flowers et al., 2019; Gros-Balthazard et al., 2020; 2021; Pérez-Escobar et al., 2021). Together, these findings highlight a complex history of spread and diversification, requiring further investigation to untangle the interactions between migration, hybridization, and local adaptation.

Archaeobotanical disciplines bring major insights to understand the origins, domestication, diffusion and diversification of crops, including fruits, cereals, pulses, and oil or fiber crops. Plant macro- (wood, seeds, fruits, and other non-woody remains) and microremains (pollen, phytoliths, starch) recovered from archaeological sites provide insights into several aspects of plants-human interactions of past societies such as plant domestications, subsistence strategies, agropastoral practices, crop-processing, or trade and craft activities (Cappers & Neef, 2012; Marston et al., 2014; Van der Veen, 2018). Macroremains are particularly valuable for reconstructing crop domestication and diffusion patterns due to their wide availability on archaeological sites, their diverse modes of conservation (e.g., as charred, mineralized, waterlogged, desiccated items), which increase their likelihood of being preserved, and their higher accuracy in crop identification compared to other botanical proxies, such as phytoliths. For example, identifying phytoliths at a low taxonomic level is challenging, as the morphological similarities within the Arecaceae family make it difficult to differentiate *Phoenix* from other taxa (e.g., Bretzke et al., 2022). Some anatomical parts (e.g., seeds or fruits) exhibit traits associated with domestication syndrome (Cappers & Neef, 2021; Fuller, 2018). In particular, seed morphometrics can be used to differentiate between wild and cultivated accessions and thus trace the emergence of domestication and cultivation (e.g., in grape, Terral et al., 2010; date palm, Gros-Balthazard et al., 2016). However, depending on the context the presence of macroremains (e.g., seeds) can suggest consumption rather than cultivation (Van der Veen, 2011). While fruits and seeds can travel with humans and may not necessarily be of local origin, the presence of vegetative parts, such as leaves or stems, suggests a higher likelihood of local cultivation. Additionally, desiccated macroremains recovered from arid environments are common (Celant et al., 2015; Van der Veen, 2007; Malleson et al., 2024) and provide a suitable material for ancient DNA analysis to better understand patterns of genetic diversity and evolutionary change (e.g., on date palm, Pérez-Escobar et al., 2021). However, this state of preservation may raise issues when it comes to precise dating since there is a possibility of modern contamination (e.g., Kaczmarek et al., 2024).

Despite the potential of archaeobotanical data to reconstruct crop history, significant challenges persist. A significant amount is scattered across grey literature, excavation reports, and peer-reviewed publications, with inconsistencies in reporting, lack of standardization (e.g., Bates & Jiménez-Arteaga, 2024), and barriers posed by diverse formats and languages. Efforts to compile archaeobotanical information, such as ArchbotLit (Kirleis et al., 2021), provide valuable lists of publications mentioning specific taxa, including the date palm. However, these tools do not integrate datasets or visualization capabilities, limiting their usefulness for broader analyses. While other platforms aim to create structured databases for archaeobotanical data - such as ADEMNES (https://www.ademnes.de/), ArboDat+ (http://arbodat.info/), I2AF (https://i2af.mnhn.fr/), they often face similar issues, such as incomplete coverage or a lack of analytical tools. Yet, meta-analyses of archaeobotanical remains have proven invaluable for uncovering the history and processes of crop

domestication and dispersal (e.g., Fuller, 2018; Silva et al., 2015; Terral et al., 2010), highlighting the significant opportunities that such integrative resources can unlock.

Among crops, these challenges are especially marked for the date palm, where existing reviews, while valuable, remain narrow in scope. Focused on specific geographic areas (e.g., the Jordan Valley, Langgut, 2024; or the Persian Gulf, Tengberg, 2012) or particular types of remains (e.g., seeds, Rivera et al., 2014), these studies often lack the standardization and comprehensive datasets needed for broader syntheses. As a result, gaining insights from archaeobotanical data into the history, domestication, and diffusion of the date palm remains an overwhelming task. Additionally, previously published evidence must be reassessed in light of recent advances, including the identification of domestication syndrome and improved taxonomic resolution (e.g., Thomas, 2013; Gros-Balthazard et al., 2016; 2017).

In this study, we aim to address these gaps by developing a centralized and evolving web platform that compiles archaeobotanical evidence for the date palm and its wild relatives (*Phoenix* spp.). The first version of *DateBack*, presented here, focuses on macroremains from pre-Islamic periods, spanning prehistory to Late Antiquity, in West and South Asia, creating a dedicated repository based on an extensive literature review. The platform consolidates data with a strong emphasis on methodological rigor, verification of the oldest occurrences, standardization, contextual details, and integration of radiocarbon information.

By providing tools for visualization and spatial analysis, *DateBack* facilitates archaeobotanical syntheses, allowing researchers to identify patterns and trends in the data. Furthermore, it lays the foundation for future applications, including ancient DNA studies and seed morphometric analyses. As a dynamic resource, it fosters interdisciplinary collaboration and provides new insights into the domestication, cultivation, and diffusion of the date palm. Ultimately, it contributes to a broader understanding of early agriculture in arid environments, making archaeobotanical data more accessible to both researchers and the wider public, supporting academic research and knowledge dissemination alike.

Material and Methods

Scope of the literature review

The review focuses on archaeobotanical remains attributed to the genus *Phoenix*, encompassing all species mentioned in the original sources. This inclusive approach acknowledges the challenges of species-level identification due to morphological and anatomical similarities within the genus (e.g., Thomas, 2013; Gros-Balthazard et al., 2016), as well as the potential contributions of wild relatives to the evolutionary history and diversity of cultivated date palms (e.g., Flowers et al., 2019).

Given the vast number of studies mentioning *Phoenix* remains across its entire distribution range and throughout prehistoric and historical periods, we restricted the scope to work with a manageable subset of data. The first iteration of the *DateBack* platform was thus designed to focus on the emergence and early diffusion of date palm cultivation, which was achieved by applying restrictions based on material type, geographical focus, and chronology:

- The review focused on macroremains, including seeds and fruits, inflorescence (flower, perianth, and pedicel), and vegetative parts (stem, petiole, leaf/leaflet, fibre, root). Microremains, including phytoliths, were excluded from the review due to the difficulties in confidently attributing them to *Phoenix*, as they are often indistinguishable from those of other members of the Arecaceae family (e.g., Bretzke et al., 2013). Alongside with pollen, they are usually sampled on off-site, non-anthropogenic locations (e.g., palaeolakes, stalagmites) for the purpose of palaeoenvironmental reconstructions, which fall outside the study (e.g., Parker et al. 2004). Additionally, examples of pollen analysis from archaeological contexts are particularly limited (e.g., Bellini et al. 2011).
- Geographically, the review concentrated on Southwest Asia and part of South Asia, covering the Levantine region, the Arabian Peninsula, and extending eastward to Iran and Pakistan. This focus was motivated by two main reasons: first Southwest Asia is widely considered as the region where date palm cultivation emerged (Tengberg, 2012; Langgut & Sasi, 2023), whereas in North Africa, it developed later (Tengberg & Newton, 2016). Second, limiting the scope to Southwest Asia in this first version allowed for a more focused review, given the extensive literature available for North Africa, particularly Egypt.
- Chronologically, the review focused on remains dating up to 630 CE, marking the early emergence of Islam. This cut-off was chosen to manage the large volume of literature and to prioritize the early emergence, diffusion and development of phoeniciculture. By this time, the date palm was cultivated across Southwest Asia and part of South Asia at a broad regional scale (Tengberg, 2012; Ros et al., 2024), although new cultivation areas may have continued to emerge.

Methodology of the literature review

The literature review included a wide range of sources documenting *Phoenix* macroremains, encompassing peer-reviewed journal articles, books, and grey literature such as PhD theses and archaeological excavation reports. For this first version, only references published before November 2024 were considered.

The review began by compiling documents mentioning *Phoenix* remains, using existing literature reviews (e.g., Tengberg, 2012; Flowers et al., 2019: Dataset S1) and other studies referencing such remains (e.g., Weiss, 2015; Langgut & Sasi, 2023). Rather than relying solely on these secondary mentions, we actively sought the original papers documenting the remains. When original papers could not be located, or when they were written in languages not mastered by any of the co-authors, these references were excluded. Particular attention was given to studies documenting early *Phoenix* remains and to key studies from underrepresented regions.

The search was then expanded in two ways. First, we leveraged existing knowledge of excavated sites within the surveyed region, drawing from archaeological databases (e.g., ArchbotLit, Kirleis et al., 2021), excavation reports, insights shared by researchers and archaeobotanists, and bibliographies from articles identified in the initial phase. Second, systematic searches were conducted on Google Scholar, Academia, and the Web of Science using the query "*Phoenix*" to identify documents potentially mentioning archaeobotanical

remains. Studies were thoroughly examined and included in the dataset only if they fell within the review's defined scope.

The objective of this compilation was to be as exhaustive as possible within the defined scope. Particular emphasis was placed on rigorously verifying and critically assessing the occurrences of date palm remains to ensure the reliability of the dataset.

Data table structure and organization

Based on the review and existing archaeobotanical remains databases, and in line with the objectives of our study, we identified the information to recover (fields), structured the data accordingly, and considered how to standardize this information. The details of this process are described in the results section ("Structure and content of the database"). The data tables are stored as CSV files.

Web application and visualization tools

The web application and visualization tools were developed using R (v. 4.4.0, R Core Team, 2022) and the shiny package v.1.9.1 (Chang et al., 2024) to facilitate user interaction with the data. These tools were designed to allow users to explore the data table, visualize trends and distributions, and generate interactive maps. The following R packages were used: leaflet v.2.2.2 (Cheng et al., 2024), plotly v.4.10.4 (Sievert, 2020), tidyverse v.2.0.0 (Wickham et al., 2019), viridis v.0.6.5 (Garnier et al., 2024), bslib v.0.8.0 (Sievert et al., 2024), leaflegend v.1.2.1 (Roh, 2024), markdown v.1.13 (Xie et al., 2024). These codes are in open access under GNU GPL3 on a git repository (https://forge.ird.fr/diade/besseiche/dateback).

The web application is hosted by the Institut Français de Bioinformatique (IFB), at https://cloudapps.france-bioinformatique.fr/dateback.

Results

Structure and content of the database

The database was built from a systematic review of literature mentioning *Phoenix* archaeobotanical remains, consolidating previously scattered data into a structured format. It was designed to compile and organize key information necessary for both assessing data reliability and enabling visualization and interpretation. It includes details on archaeological sites, chronological contexts, remains data (e.g., taxon, anatomical parts, preservation state), and radiocarbon dating records performed on date palm remains.

The gathered information was organized into two complementary data tables. The structure and detailed descriptions of the main table are provided in Appendix 1. This main table (Appendix 2) serves as the primary repository, capturing detailed information about archaeobotanical records of *Phoenix*. The second table focuses exclusively on radiocarbon-dated date palm remains (Appendix 3). Depending on the level of detail available for each field, some were standardized to finite lists (e.g., "type of archaeological site"), while others were left as free text to capture nuanced or unique information (e.g., "archaeological context") (Appendix 1).

Briefly, the main data table encompasses the following categories of information:

- Archaeological site details: Name of the site, modern country, type of site (e.g., burial area, settlement), geographic coordinates, and excavation years.
- Chronological context: Description and dating of the context in which the remains were found. Depending on the site, on the number of distinct periods represented, and on the precision of the descriptions in the articles, the context may refer to one stratigraphic unit, locus, area, or several. As a result, multiple "chronological context" sections may exist for the same "archaeological site" section. Information on the nature of the context, including its identification number (if available) or other relevant details, are filled out. The start and end dates represent the earliest and latest possible dates for the context. Broad cultural period and dating quality of the context (whether relative, absolute on material other than date palm, or absolute directly on date palm remain) are also provided when available. For cases of absolute dating on date palm remains, detailed information is provided in the dedicated radiocarbon data table.
- Remains: Type(s) of remains (e.g., seed, stem, petiole), preservation state (e.g., charred, desiccated), taxon as reported in the original paper, and, if necessary, our correction of the taxon with an explanation (see Appendix 1 for details). The number of remains is included as free text to accommodate entries such as "10 seeds," ">10 seeds," or "1 seed fragment + unquantified fruits," reflecting a qualitative approach to quantification, as the complexities and debates surrounding standardization in archaeobotanical studies are beyond the scope of this paper (e.g., Bates & Jiménez-Arteaga, 2024).
- References: The original reference(s) reporting the remain(s) in question, as well as additional sources where other critical information was gathered (e.g., follow-up radiocarbon dating).

Each entry in the main table represents a unique combination of an archaeological site, contexts with similar start and end dates, and a taxon. To clarify:

- Chronological grouping: Contexts at a given site with similar start and end dates were grouped into a single entry, with identification numbers or other details recorded in the field "chronological context". For example, at Dalma Island DA11, two contexts (contexts 4 and 15) dated to the same period both yielded date palm remains and were therefore combined into a single entry (Beech & Shepherd, 2001). Conversely, when contexts corresponded to distinct periods, they were recorded as separate entries. For instance, for the site of Qal'at al-Bahrain, four entries were created for four distinct periods: the early, middle, and late Dilmun periods (2200–1750 BCE, 1450–1350 BCE, and 1000–500 BCE, respectively) and the Tylos period (300 BCE–300 CE) (Tengberg & Lombard, 2001). In some cases, overlapping contexts were kept separate when one offered more specific dating within a broader timeframe, as seen at Mleiha (Mouton et al., 2012; Dabrowski, 2019; Tengberg, 1999a; Pena-Chocarro & Barron Lopez 1999). Additionally, some contexts were retained as distinct entries despite close date ranges when they corresponded to different archaeological areas. For instance, at Khirbat al-Jariya, two entries have very similar dating (-1100 to -900 BCE and -1150 to -900 BCE), which were kept separate since they correspond to a stone

structure and a slag mound, respectively (Liss et al., 2020; Stroth et al., 2023; Ben-Yosef et al., 2010).

- Taxon-specific entries: Entries were also categorized based on the taxon reported in the study. When different studies documented distinct taxa at the same site, separate entries were created to reflect these variations. For example, at Jericho, one entry records *Phoenix dactylifera* seeds (Hopf, 1969; Hopf, 1983), while another records charcoal identified as either *Phoenix* or *Hyphaene* (Western, 1971), which was thus classified under Arecaceae.

In addition to the main data table, a complementary table was created to compile radiocarbon data on date palm remains. This so-called C14 date palm table includes fields identical to those in the main table to link each radiocarbon analysis to its corresponding site and context. It also contains specific fields such as laboratory codes, anatomical parts dated and their preservation states, raw ^{14}C ages (with associated uncertainty), and calibrated age ranges. Multiple radiocarbon analyses performed on the same site or context are recorded as separate entries to ensure comprehensive documentation.

Development of the web application and visualization tools

The visualization application consists of a website with two visualization tabs: a *Data Tables* tab which allows the navigation in and the download of the data tables and an *Interactive Map* tab (Figure 1). Both can be interactively explored thanks to various filters. In particular, we provide a filter on the chronological range to explore a given time period. We also allow users to select on the present-day countries they wish to explore, the anatomical parts, their preservation states, the dating qualities of the context and the taxa.

In addition, four informative tabs (*Versioning*, *Bibliography*, *License* and *Contact*) are accessible. The Contact tab was added to encourage community engagement, thanks to a dedicated email address (dateback@ird.fr), for contributions or questions about *DateBack*.

Figure 1. Overview of the *DateBack* platform Homepage.

Screenshot of the homepage showcasing the main features of the *DateBack* web platform, including options for data exploration, visualization, and user interaction.

Results of the literature review

Our literature review encompassed 110 accessible documents that report *Phoenix* spp. remains from prehistoric to pre-Islamic Southwest Asia and South Asia, all of which have been included in the first iteration of the *DateBack* database. While we systematically relied on original sources to ensure accuracy, one exception was made: for the numerous studies authored by Liphschitz in Hebrew, we instead referenced a comprehensive English-language review by the same author, which consolidated detailed information on the sites, contexts, and archaeobotanical findings (Liphschitz, 1996).

The main data table includes 140 entries from 99 sites across 14 present-day countries (Appendix 2). The C14 date palm datatable contains 55 entries from 17 sites across 6 countries (Appendix 3).

In the original sources, the majority of remains were identified as date palm (*Phoenix* (cf.) *dactylifera*), accounting for 129 entries (92.1%), with only a single report of other species, namely *Phoenix theophrasti*. When original sources lacked precise identification, remains were classified more broadly in the database as *Phoenix* sp. (n=8, 5.7%) or as Arecaceae when identified as either *Phoenix* or another palm genus (n=3, 2.1%). In addition to this compilation of the taxa as reported in the source, we also provided taxonomic corrections in cases of uncertainty. Specifically, for three entries from the Levant, at 'Atlit (Atlit Yam) and Ohalo II, where both *P. theophrasti* and *P. dactylifera* had been reported, we reclassified them as *Phoenix* sp. to reflect the uncertainty (Galili et al., 1993; Liphschitz & Nadel, 1997; Kislev et al., 2004).

Of the 140 recorded entries, 56 (40.0%) reported multiple anatomical parts of *Phoenix*. The majority of entries comprise seeds (n=98, 70.0%; Figure 2A), a pattern influenced by taphonomic, analytical, and disciplinary biases. Taphonomic factors favor seed preservation due to their durability and resilience, with *Phoenix* seeds in particular demonstrating exceptional resistance to charring (Ivorra et al., 2024). Analytical biases arise because seeds are more easily identifiable at the species or genus level compared to other anatomical parts, owing to their distinct morphological features and the availability of extensive reference collections and seed atlases for taxonomic identification (Nesbitt et al., 2003; Cappers & Neef, 2012). Disciplinary biases further shape the types of remains identified. For example, stems are frequently reported from Levantine sites, reflecting the research emphasis of Nili Liphshitz, whose numerous studies in the region account for half of these entries, illustrating how regional expertise and research specializations influence the types of remains reported.

Most entries comprise charred remains (n=112, 80.0%, Figure 2B), which aligns with expectations given the prevalence of burning practices and the preservation conditions typical of many archaeological contexts (Van der Veen, 2007). However, the discovery of some desiccated remains (n=15, 10.7%) is particularly noteworthy, as their preservation provides unique opportunities for ancient DNA studies (e.g., Pérez-Escobar et al., 2021). Only two entries report waterlogged remains, both from the submerged site of 'Atlit Yam in the Mediterranean (Galili et al., 1993; Kislev et al., 2004). As expected, such remains are rare in arid West Asia, unlike in Europe, where waterlogged remains, such as grape, are more commonly found (Colledge & Connolly, 2014; Bouby et al., 2023).

Figure 2. Overview of *Phoenix* sp. remain characteristics documented in the review (n=140).

A. Distribution by type of remains. B. Distribution by preservation state. Since one entry may include several types of remains and multiple preservation states, the total count exceeds the number of entries in the dataset.

The earliest recorded *Phoenix* remains are from Ohalo II in the Levant (21,000 BCE to 17,000 BCE, Liphshitz & Nadel, 1997), followed by a long gap until approximately 10,000 years ago, with remains only becoming more frequent after 5,000 BCE, steadily increasing through the 4th millennium BCE to the pre-Islamic period (up to 630 CE) (Figures 3-4). Half of the entries come from contexts with radiocarbon dating, either from layers that are radiocarbon-dated but not necessarily the date palm remains themselves (n=46) or from directly radiocarbon-dated date palm remains (n=23), while the rest are either relatively dated or lack chronological information.

Figure 3. Chronological distribution of *Phoenix* spp. remains documented in the review.

Number of entries in the main data table categorized by millennium. The pie chart on the top left represents the total proportions of dating qualities.

Figure 4. Chronological distribution of archaeological sites with *Phoenix* sp..

Geographically, date palm remains are concentrated in certain regions, while other areas are notably lacking (Figures 4-5). Although this partly reflects actual distribution patterns, the disparity may also result from excavation biases. For example, the Levant has a long history of excavations that is reflected in the results with 64 entries (45.7%). Conversely, although Saudi Arabia covers a third of the surface of the studied region, solely nine entries (6.4%) are from this country, with solely five archaeological sites with published recorded archaeobotanical remains; this will change as ongoing archaeological projects in Saudi Arabia continue to expand (Dabrowski, Bouchaud et al., 2024). Additionally, biases in the retrieval of archaeobotanical remains can arise from early excavation priorities, such as in Iraq, where the focus was predominantly on monumental architecture and artifacts, leaving plant remains largely uncollected or unstudied. This trend was common in the early 20[th] century, when archaeobotany was not yet a well-established field (Zohary et al., 2012), resulting in significant gaps in the botanical record from such excavations.

Prior to 4000 BCE:

Between 4000 and 2000 BCE:

After 2000 BCE:

Dating quality: ▪ relative or unknown ▲ absolute ● absolute (on date palm)

Figure 5. Geographical distribution of archaeological sites with *Phoenix* remains across different time periods.

The dating quality of the archaeological context is represented by different symbols.

Discussion

In this study, we reviewed the literature and compiled archaeobotanical macroremains of *Phoenix* sp. in West and South Asia from prehistory to Late Antiquity. The findings were integrated into the web platform *DateBack*, which centralizes the data in structured tables and visualization tools to support the study of spatial and temporal patterns of *Phoenix* remains. These records, in turn, provide insights into the early history of date use and cultivation in the region.

Synthesizing archaeobotanical evidence of *Phoenix*: Presenting *DateBack*, a platform for data compilation and visualization

Our literature review drew from 110 accessible sources. Despite our efforts toward comprehensive coverage, some references remained inaccessible due to language barriers, lack of digitization, or restricted availability. These challenges reflect broader issues in archaeobotanical research (Lodwick, 2019), and emphasize the importance of open science practices in ensuring data accessibility across disciplines (Wilkinson et al., 2016). By consolidating scattered records into an open-access platform, *DateBack* helps mitigate these challenges, providing a centralized and structured resource for future studies. Additionally, researchers can contribute new data or corrections via the provided contact email, ensuring the platform remains dynamic and continuously refined.

Our reliance on original sources may have led to the omission of some data, yet this approach ensured the accuracy and reliability of the recorded dataset by avoiding errors often introduced through secondary citations. For instance, we found that *Phoenix* remains were reported from Ain Ghazal (present-day Jordan), yet no such evidence appears in the original publication (Rollefson et al., 1985). By systematically verifying each original reference, we corrected such misattributions, ensuring that misreported data were not perpetuated in our dataset.

With this verified dataset, we integrated 140 entries from 99 sites across 14 countries into *DateBack*'s main dataset, alongside 55 radiocarbon-dated entries in a separate table. Compared to previous reviews, our dataset offers more standardized and detailed records. For instance, we systematically assigned geographic coordinates to each site, facilitating spatial analyses and visualizations, and recorded excavation years, which may serve as an indirect measure of data reliability, as identification methods and counting methods may differ between older and more recent excavations (Marston et al. 2014; Cappers and Neef 2012). Although previous reviews have recorded dating quality (e.g., Flowers et al., 2019), this study is the first to not only distinguish between absolute and relative dating but also systematically compile all radiocarbon dates performed on *Phoenix* remains within our review's scope, enhancing chronological reliability.

Future enhancements are expected to improve the usability, transparency, accessibility, and analytical depth of the platform, reinforcing *DateBack*'s role as a key resource for archaeobotanical research For instance, the present dataset does not yet include an assessment of data confidence, as suggested in previous studies (e.g., Fuller & Weber, 2005, discussed in Flowers et al., 2019). Implementing such a grading system in future versions, with the potential for community contributions, could allow users to distinguish between

well-documented remains and those with ambiguous identification or uncertain dating. Additional metadata fields are planned to track re-examinations of remains, such as seed morphometric analyses or ancient DNA studies, with links to relevant publications, a functionality that is already under development. Documenting storage locations will facilitate further research and validation, while integrating photographs of remains, when available, will support identification and comparative analyses.

Reassessing the earliest archaeobotanical evidence of *Phoenix* before 4000 BCE

The earliest archaeobotanical evidence of date palm (*Phoenix* sp.) prior to 4000 BCE, as recorded in the database, consists of eight entries from seven archaeological sites across the Persian Gulf, the Levant, and Pakistan (Figure 5). However, as detailed below, the nature, preservation state, and contextual or chronological reliability of these remains vary significantly. Their interpretation is further complicated by unknown past distributions of *Phoenix* species, including wild date palms (Barrow, 1998; Gros-Balthazard & Flowers, 2021) and difficulty to differentiate them morphologically (e.g., Gros-Balthazard et al., 2016).

The earliest reported *Phoenix* evidence comes from Ohalo II (21000-17000 BCE) in the Levant, where a single "wood" fragment was recovered from a surface layer of the excavated campsite (Liphschitz & Nadel, 1997). Given the scarcity of material, uncertain depositional context, and lack of direct radiocarbon dating, this evidence remains highly uncertain and cannot be confidently attributed to this early period.

Additional finds from the Levant also present challenges in terms of dating and interpretation. At 'Atlit (8100-7400 BCE), a waterlogged seed and a "wood" fragment were recovered, but their precise archaeological context is unclear, as sources report their presence in either a well or hearths (Galili et al., 1993; Kislev et al., 2004). Furthermore, the taxonomic identification is inconsistent: while the "wood" was identified as *P. dactylifera* (Galili et al., 1993), the seeds were later attributed to *P. theophrasti* (Kislev et al., 2004). Another site, the Cave of Treasure, has yielded numerous date seeds, interpreted as domesticated based on their size (Fuller, 2018). Presumably from the Chalcolithic layer (4250–4050 BCE), their attribution remains uncertain due to the absence of direct radiocarbon dating and the site's later Roman-period occupation, as noted by the authors (Zaitschek, 1961; Bar-Adon, 1980).

On the eastern side of the date palm's range, early evidence from present-day Iran and Pakistan presents similar dating uncertainties. At Mehrgarh, two date seeds were recovered from 6[th] millennium BCE contexts (Costantini, 1983, 1985), but their silicified state prevents direct radiocarbon dating. Another site, Tepe Gaz Tavila (5400–4800 BCE), has also yielded date seeds (Costantini & Biasini, 1985, cited in Tengberg, 2012), but as we could not access the original publication, they were not recorded in *DateBack*, and their precise context, quantity, and preservation state remain unknown. For both sites, beyond dating issues, the limited quantity of seeds, lack of associated charcoal evidence, and the suggested presence of wild date palms in the region (Fuller & Stevens, 2019) make it unclear whether the seeds were locally produced or introduced, and whether they came from wild or cultivated palms.

In contrast to the uncertainties surrounding early *Phoenix* remains in the Levant and Pakistan, archaeobotanical evidence from the Persian Gulf appears more chronologically and contextually reliable, with several sites yielding remains from the late 6[th] to early 5[th] millennium BCE. The least-documented case is Tell al-Oueli in present-day Iraq, where stem or petiole fragments were recovered from Late Ubaid contexts (5200-4350 BCE; Neef, 1991). While the archaeological context is well-documented, the remains are unquantified, their state of preservation unknown, and the absence of radiocarbon dating prevents confirmation of their exact age, altogether making their interpretation difficult.

Stronger evidence comes from further south with the site of Aş Şabīyah (5300-4900 BCE) in present-day Kuwait, where four mineralized date seeds were found (Beech, 2003). Although their mineralized state precludes direct radiocarbon dating, indirect dating based on associated materials supports their chronological placement (Parker, 2011), and morphometric analyses suggest they correspond to dates gathered from wild populations (Gros-Balthazard et al., 2017), aligning with this ancient timeframe.

The more securely dated remains come from Dalma 11, on Dalma Island in the southern Persian Gulf, where two charred date seeds and seed imprints were discovered (Beech & Shepherd, 2001). The seeds, directly radiocarbon-dated to 5290-4940 cal. BCE and 4810-4540 cal. BCE, are the oldest yet reported for the species. The presence of Ubaid pottery, a material culture tradition originating in southern Mesopotamia, suggests that these date fruits may have been imported from the northern Gulf, although their exact origin remains uncertain (Tengberg, 2012). Regrettably, their poor conditions prevent morphometric analysis from determining whether they were wild or cultivated.

Overall, these findings illustrate both the insights and limitations that archaeobotanical evidence provides in reconstructing early date palm use. In the Persian Gulf, date consumption is securely attested by 5000 BCE, whereas elsewhere, the evidence remains ambiguous. Among these finds, no clear evidence of cultivation has emerged, as the remains that could be analyzed align with wild populations, while others are too poorly preserved for assessment.

The transition to date palm cultivation during the 4[th]–3[rd] millennia BCE

During the 4[th] and 3[rd] millennia BCE, our review revealed that the number of archaeobotanical records of *Phoenix* increases significantly with 33 recorded entries from 29 sites spanning the Levant, Mesopotamia, the Gulf and Pakistan. While earlier evidence remains ambiguous, finds from this period suggest a growing reliance on date palms, with indications of early cultivation emerging, particularly in southern Mesopotamia and the Persian Gulf.

Archaeobotanical remains from southern Mesopotamia and the northern Gulf contrast with earlier findings, as they appear in significantly larger quantities and more consistently include a diverse range of anatomical parts, such as fibers and leaf fragments, alongside seeds. At Eridu (4000-3000 BCE), abundant charred seeds were excavated and considered domesticated by the author (Gillett, 1981) and others (e.g., Fuller & Stevens, 2019), though the absence of radiocarbon dating limits chronological certainty. In contrast, at Al-Khidr (3200–1200 BCE), dozens of seeds and leaf imprints on bitumen were recovered, with radiocarbon dating of the occupation providing a more secure chronological framework

(Hajnalova et al., 2009). This substantial and varied assemblage suggests that date palms were actively managed and cultivated rather than merely exploited from wild wild populations.

This limited number of finds, largely due to geopolitical constraints that have restricted archaeological investigations in the region, likely leaves additional evidence undocumented. However, other sources complement the archaeobotanical record. Iconographic depictions, such as the Warka vase from Late Uruk (4th millennium BCE), illustrate date palm gardens, while textual sources from the 3rd millennium BCE attest to the presence of date palm groves where date palms were cultivated alongside other crops (Tengberg, 2012; Miller, 2016; Michel-Dansac & Caubet, 2013; Paszke, 2019). Together, these records strengthen the hypothesis of early cultivation in southern Mesopotamia and the northern Gulf.

Archaeobotanical evidence from southern Iran and the Indo-Iranian borderland is limited, resulting in a fragmented and uneven picture of date palm management during the 4th and 3rd millenia BCE. Yet, at Konār Şandal (3rd mill. BCE), the discovery of both seeds and vegetative parts alongside cereals and grape remains provides compelling evidence for date palm cultivation (Tengberg, 2012; Mashkour et al., 2013). This interpretation is further supported by depictions of date palms chlorite vases from the same site (Perrot & Madjidzadeh, 2005). Conversely, in southwestern Pakistan, date seeds from Miri Kalat and Shahi Tump (Tengberg, 1998, 1999b) were directly radiocarbon dated to the early 3rd millennium BCE and identified as wild through geometric morphometric analyses, though their classification as *P. dactylifera* or *P. sylvestris* remains uncertain (Ivorra et al., 2024). The contrast with Konār Şandal underscores how, within the same region, human populations engaged with date palms in diverse ways, ranging from reliance on wild stands to active cultivation. Aside from these discoveries, available data remain scarce (Costantini, 1985; Miller, 1982), largely due to limited excavations and geopolitical constraints. Notably, no archaeobotanical evidence has been recovered from the arid lowland of southwestern and south-central Iran, where conditions would have been suitable for date palm cultivation, prompting Tengberg (2012) to suggest that date palms may have been grown there as early as in Mesopotamia.

In eastern Arabia, apart from the two previously mentioned 5th millennium BCE seeds from Dalma islands, which may have been imported from Mesopotamia (Beech & Shepherd, 2001; Tengberg, 2012), no date palm remains are found before the end of the 4th millennium. At Hili 8, one of the earliest finds in the region (3000-2700 BCE), abundant seeds and carbonised vegetative parts were recovered alongside cereals, with the occupation layer securely dated by radiocarbon (Cleuziou, 1982; 1997; Tengberg, 1998). Through the 3rd millennium BCE, charred date palm seeds and vegetative fragments alongside cereals appear at several sites, including Bāt (Tengberg, 1998; Deckers et al., 2019) and Qal'at al-Bahrain (Tengberg & Lombard, 2001). Although not all finds are radiocarbon-dated, their abundance and association with other cultivated species suggest that date palm cultivation was established in eastern Arabia by the 3rd millennium BCE.

Further east, in the Indus Valley, within the context of the Harappan culture (second half of the 3rd millennium BCE), date palm remains include numerous palm branches and stem fragments at Mehrgarh, likely used for construction (Jarrige et al., 1995), and abundant seeds at Nausharo (Costantini, 1990a). These findings, associated with the broader diffusion of

crops such as cereals, pulses, and cotton (*Gossypium* sp.) into the region, have been interpreted as evidence of date palm cultivation by Tengberg (2012).

In the Levant, numerous sites have yielded date palm remains from the 4th and 3rd millennia BCE. However, as with the presumed earlier finds discussed above, these are limited to either seeds or charcoal and appear in small quantities. Only Qumrân, cave 24, has two secured direct datings providing the earliest confirmed evidence of date fruit consumption in the region (2920-2657 cal. BCE and 2897-2618 cal. BCE; Patrich, 1994; Liphschitz & Bonani, 2001). Based on seed size, these were interpreted as wild (Liphschitz & Bonani, 2001), and other scholars suggest the presence of wild date palm in the region (Zohary & Spiegel-Roy, 1975), supported by *Phoenix* pollen evidence through the Holocene (e.g., Litt et al., 2012). However, taxonomic uncertainties persist, and re-examining charcoal assemblages analyzed decades ago using updated anatomical criteria (Bouchaud et al., 2012; Thomas, 2013) could help clarify identifications, particularly distinguishing between stipe and petiole fragments or refining genus-level classifications (e.g., Bronze Age charcoal fragments from Jericho; Western, 1971). While the recurrent finds reviewed here strongly suggest the presence and use of date palms during the Chalcolithic and Bronze Age, the current evidence remains inconclusive regarding local cultivation in the Levant, as proposed in previous studies (Langgut & Sasi, 2023; Fuller, 2018). Additional well-contextualized and larger botanical assemblages would help further evaluate this hypothesis.

Our findings, in light of the literature, suggest that date palm cultivation emerged in southern Mesopotamia and the northern Gulf by the 4th millennium BCE, while in eastern Arabia and the Indus Valley, the first evidence of phoeniciculture appears in the 3rd millennium BCE. This pattern aligns with broader agricultural developments in the region. Cereal agriculture is known to have been introduced to eastern Arabia (Cleuziou, 2004), and trade networks linked Mesopotamia, Iran, and the Indus Valley, with the Gulf serving as a hub for exchange and agricultural transmission (Beech & Shepherd, 2001; Tengberg, 2012). Given that the earliest secure evidence for date palm cultivation comes from southern Mesopotamia and the northern Gulf, it is plausible that, like cereals, date palm cultivation in eastern Arabia—and perhaps also in the Indus Valley—was introduced rather than developing independently. In contrast, while consumption evidence exists in the Indo-Iranian borderland and the Levant, cultivation remains uncertain during this period.

The late expansion of date palm cultivation throughout Southwest and South Asia in the 2nd and 1st millennia BCE

Although date palm cultivation may have begun earlier, the first secure evidence in the Levant comes from the Iron Age, with well-documented finds at two sites in Jordan: Khirbet An Nahhas (1199-844 BCE, Baierle et al., 1989; Engel, 1993) and Khirbat al-Jariya (1100-900 BCE, Liss et al., 2020; Stroth et al., 2023). Their dating relies on radiocarbon analysis, including on date palm remains at Khirbat al-Jariya, and both seeds and charcoal were found in substantial quantities at both sites.

Unlike the Levant, where older remains exist but are often difficult to interpret due to uncertain dating and taxonomic challenges, in northwestern Arabia, the earliest securely dated remains also represent the oldest known evidence of date palm cultivation in the region. At Tall al-Sālimīyyah in the al-'Ulā region, seeds, stem, and petiole fragments have been

radiocarbon-dated to 1215–1015 cal. BCE, supporting the hypothesis of local cultivation (Rohmer et al., 2022). At the same period, charcoal and seeds, some of them radiocarbon-dated, are attested at the neighboring site of Tayma (Dinies et al., 2016; Tourtet et al., 2021).

In southern Arabia, the dataset compiled in *DateBack* is limited due to both geopolitical constraints limiting archaeological excavations and restricted access to original reports. While a potential early mention of date palm use is reported at ar-Raqlah 1, a site occupied during the 3rd millennium BCE (Costantini, 1990b, cited in Schiettecatte, 2013), direct access to the documentation was unavailable for this review. As a result, our dataset is constrained to three sites. At Şabir, remains described generically as "palm" have been reported from the 2nd millennium BCE (Vogt & Sedov, 1998; Görsdorf & Vogt, 2001), but without clear taxonomic identification their attribution to *Phoenix dactylifera* remains uncertain, and direct radiocarbon dating is lacking. In contrast, securely identified desiccated date seeds from Raybūn (Levkovskaya & Filatenko, 1992) and charred seeds and imprints from Barāqish (de Maigret et al., 1986) provide more reliable evidence for date consumption by the 1st millennium BCE. Since early to mid-1st millennium BCE inscriptions explicitly mention palm groves (reviewed in Schiettecatte, 2013), it is highly likely that date palms were already under cultivation during this period.

In central Arabia, data are limited to the unquantified remains from Qaryat al-Fāw, a site occupied approximately 2000 years ago (Muṣṭafā Ḥasan et al., 2019). More material from the contemporaneous site of al-Yamāma (Schiettecatte & al-Ghazzi, 2015) remains unpublished (Bouchaud, unpublished; Chambraud et al., ongoing). The currently available data thus suggest a late emergence of date palm gardens in central Arabia compared to neighboring regions, aligning with the relatively late development of urban oases in the area (Schiettecatte & al-Ghazzi, 2016). However, the lack of archaeobotanical studies makes it difficult to determine whether earlier cultivation may have occurred but remains undocumented.

In the continental part of northeast Arabia, archaeobotanical data are notably lacking, probably due to the limited number of large-scale archaeological projects. However, archaeobotanical analysis on the Classical city of Thaj (present-day Saudi Arabia), dated to the 7th century BCE-7th century CE, is currently underway and will be published soon (Dabrowski, ongoing).

Conclusion & Prospects

In this study, we have developed the *DateBack* platform, a centralized resource designed to facilitate research on the history, domestication, and diffusion of the date palm (*Phoenix dactylifera* L.). By rigorously compiling and standardizing archaeological and archaeobotanical data, the platform enables a refined assessment of the spatial and temporal patterns of date palm use and management.

Our findings confirm that the earliest secure evidence for date consumption is concentrated in the Persian Gulf, while evidence from other regions, particularly the Levant, remains ambiguous due to uncertain contexts, taxonomic issues, and a lack of direct dating. Date palm cultivation appears established in southern Mesopotamia and the northern Gulf by the 4th–3rd

millennium BCE and may have developed as early in southern Iran, though the scarcity of archaeobotanical studies prevents confirmation. In the Indus Valley and eastern Arabia, cultivation is attested by the 3rd millennium BCE, and connections with the Gulf may indicate a diffusion process rather than independent emergence.

In the Levant and northwestern Arabia, secure evidence for cultivation only appears at the end of the 2nd millennium BCE, while in southern Arabia, it is not attested until the 1st millennium BCE. The late emergence of date palm cultivation in various regions of the Arabian Peninsula seems to result from a combination of regional factors. In northwestern Arabia, available data suggest a distinct regional agricultural pattern (Dinies et al., 2016; Rohmer et al., 2022; Chambraud, unpublished), whereas in other areas, limitations stem primarily from the scarcity of archaeobotanical studies or challenges in interpreting existing evidence. Additionally, the lack of Bronze Age contexts narrows the chronological perspective, and restricted access to reports and publications continues to hinder data collection.

While this study has reviewed date palm remains from earliest evidence through Late Antiquity, their interpretation has primarily focused on early date consumption and emergence of cultivation in South and Southwest Asia. Further research is needed to explore additional questions, including continuity—whether date palm cultivation was consistently maintained in certain regions or periodically abandoned—and diffusion routes. In addition, morphometric analyses of well-preserved seeds could help assess their wild or domesticated status and refine the understanding of date palm consumption and cultivation in different regions. Expanding archaeological research, particularly in data-poor areas, would also be essential to filling critical gaps in current knowledge.

The *DateBack* platform is designed as a dynamic and scalable resource, with future developments aiming to extend its scope to later periods (e.g., Islamic and beyond) and additional regions such as North Africa. Beyond macroremains, forthcoming updates may incorporate data on microremains (e.g., pollen, phytoliths), with initial work on this expansion already underway but not yet included in the current public version. Further iterations could also integrate impressions, objects, and textual or iconographic evidence, enhancing the interpretative framework. By fostering interdisciplinary collaboration among archaeobotanists, historians, and geneticists, *DateBack* will continue to refine our understanding of the cultural and agricultural significance of the date palm, adapting as new discoveries and methodologies emerge.

Appendices

Appendix 1. Structure and details of the Main Data Table fields. (https://doi.org/10.23708/4ZHYZU)

Appendix 2. Main Data Table summarizing the results of the literature review. (https://doi.org/10.23708/4ZHYZU)

Details and formats of the fields are provided in Appendix 1. For entries with radiocarbon dating performed on date palm remains, corresponding results are available in Appendix 3.

Appendix 3. Radiocarbon data table for date palm remains. (https://doi.org/10.23708/C5N9CM)

This table documents radiocarbon analyses of date palm remains, linking each analysis to its site and context. It includes specific fields such as laboratory codes, anatomical parts, preservation states, raw ^{14}C ages (with uncertainty), and calibrated age ranges.

Acknowledgments

We would like to acknowledge Naomi Miller, Margareta Tengberg, Guy Bar-Oz, Daniel Fuks and Laurent Bouby for their valuable contributions to the archaeobotanical review. Special thanks to Yann Bourgois and Vincent Battesti for their insightful discussions and comments on the manuscript. We also appreciate Stéphanie Thiébault, Michael Purugganan, and Jonathan Flowers for their assistance in accessing some key documents. Additionally, we would like to thank the IRD documentation team in Montpellier, particularly Aure Franceschi and Hanka Hensen, for their help in locating references, as well as the librarians at NYU for their support. The *DateBack* logo has been designed by Emeline Pobelle (https://www.malt.fr/profile/emelinepobelle). AI tools, specifically ChatGPT and DeepL, were used to assist in polishing and refining the text throughout the manuscript, and in writing the code of the web application.

Data, scripts, code, and supplementary information availability

The structure of the data tables is licensed under CC-BY, and the contents of the data tables are licensed according to the original datasource or CC-BY.

The supplementary information is provided as an appendix, and both data tables (version 1) are accessible online via DataSuds. The general repository can be found at https://dataverse.ird.fr/dataverse/dateback, and include the data tables (main: https://doi.org/10.23708/4ZHYZU and radiocarbon dating: https://doi.org/10.23708/C5N9CM). Scripts and code are available online in a git repository: https://forge.ird.fr/diade/besseiche/dateback (https://doi.org/10.23708/KNQIN1 for v.1.0). All codes are released under the license GNU GPL v3.

The web application with the currently described version of the data table is hosted by the French "Institut Français de Bioinformatique" (IFB) at https://cloudapps.france-bioinformatique.fr/dateback.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Funding

This work was funded by the regular budgets of IRD, MNHN and CNRS, with additional support from the "al-'Ulā Date Palm Agrobiodiversity" project (directed by Muriel Gros-Balthazard and Vincent Battesti) and "ECO-Seed" project (directed by Charlène Bouchaud), part of the Oasis program of AFALULA (Agence Française pour le Développement d'AlUla).

Contribution of each author

The design of the data table was primarily carried out by MB, EC, VD, CB, MGB, with additional contributions from EB. The literature review and data entry were mainly performed by EB, EC, VD, CB, MGB, with contributions from MB. The curation of the data table was led by MB and MGB. The design and development of the web application was primarily handled by MB, with input from FS and MGB. The manuscript was primarily written by MB and MGB, with revisions and input from all other co-authors.

References

Barrow, S. C. (1998). A Monograph of *Phoenix* L. (Palmae: Coryphoideae). Kew Bulletin, 53(3), 513-575. https://doi.org/10.2307/4110478

Baierle, H., Frey, W., Jagiella, C., & Kürschner, H. (1989). Die Brennstoffressourcen im Raum Fenan (Wadi Araba, Jordanien) und die bei der Kupferverhüttung verwendeten Brennstoffe. In A. Hauptmann, E. Pernicka, & G. Wagner (Eds.), Archäometallurgie der alten Welt. Der Anschnitt, Beiheft 7 (pp. 213–222). Deutsches Bergbau-Museum.

Bates, J., & Jiménez-Arteaga, C. (2024). Comparable quantification methodologies in archaeobotany – a work-in-progress and debate. Vegetation History and Archaeobotany, 33(5), 671–686. https://doi.org/10.1007/s00334-023-00982-6

Beech, M. (2003). Archaeobotanical evidence for early date consumption in the Arabian Gulf. In The Emirates Center for Strategic Studies & Research (Ed.), The Date Palm: From Traditional Resource to Green Wealth (pp. 11–31). The Emirates Center for Strategic Studies & Research.

Beech, M., & Shepherd, E. (2001). Archaeobotanical evidence for early date consumption on Dalma Island, United Arab Emirates. Antiquity, 75(287), 83–89. https://doi.org/10.1017/S0003598X00052765

Bellini, C., Pavan, A., Pignotti, L., Gonnelli, T., & Lippi, M. M. (2020). Food plants in pollen records from ancient Southern Arabia: The evidences from Sumhuram (Southern Oman). Journal of Arid Environments, 177, 104131.

https://doi.org/10.1016/j.jaridenv.2020.104131

Ben-Yosef, E., Levy, T. E., Higham, T., Najjar, M., & Tauxe, L. (2010). The beginning of Iron Age copper production in the southern Levant: New evidence from Khirbat al-Jariya, Faynan, Jordan. Antiquity, 84(325), 724–746. https://doi.org/10.1017/S0003598X00100195

Bouby, L., Bonhomme, V., Ivorra, S., & Pastor, T. (2023). Experimental waterlogging of grape seeds, impact on seed shape and geometrical reversing for morphometric inference. Journal of Archaeological Science: Reports, 51, 104204. https://doi.org/10.1016/j.jasrep.2023.104204

Bretzke, K., Armitage, S. J., Parker, A. G., Walkington, H., & Uerpmann, H.-P. (2013). The environmental context of Paleolithic settlement at Jebel Faya, Emirate Sharjah, UAE. Quaternary International, 300, 83–93. https://doi.org/10.1016/j.quaint.2013.01.028

Bretzke, K. (2022). Archaeological records indicate a complex history of Pleistocene hunter-gatherer societies in Arabia. L'Anthropologie, 126(3), 103047. https://doi.org/10.1016/j.anthro.2022.103047

Cappers, R.T.J., Neef, R., 2012. Handbook of plant palaeoecology, Groningen archaeological studies. Barkhuis, Groningen.

Cappers, R. T. J., & Neef, R. (2021). Handbook of Plant Palaeoecology (2nd ed., Vol. 19). Barkhuis; JSTOR. https://doi.org/10.2307/j.ctv23wf387

Celant, A., Magri, D., Stasolla, F. (2015). Collection of Plant Remains from Archaeological Contexts. In: Yeung, E., Stasolla, C., Sumner, M., Huang, B. (eds) Plant Microtechniques and Protocols. Springer, Cham. https://doi.org/10.1007/978-3-319-19944-3_25

Chang, W., Cheng, J., Allaire, J., Sievert, C., Schloerke, B., Xie, Y., Allen, J., McPherson, J., Dipert, A., & Borges, B. (2024). shiny: Web application framework for R [Manual]. https://CRAN.R-project.org/package=shiny

Cheng, J., Schloerke, B., Karambelkar, B., & Xie, Y. (2024). leaflet: Create interactive web maps with the JavaScript "leaflet" library [Manual]. https://CRAN.R-project.org/package=leaflet

Costantini, L. (1983). The Beginning of Agriculture in the Kachi Plain: The Evidence of Mehrgarh. In B. Allchin, F. R. Allchin, & M. Sidell (Eds.), South Asian Archaeology 1981: Proceedings of the Sixth International Conference of the Association of South Asian Archaeologists in Western Europe (pp. 29–33). Cambridge University Press.

Costantini, L. (1985). Considerazioni su alcuni reperti di palma da dattero e sul centro di origine e l'area di coltivazione della Phoenix dactylifera L.. Rome. In G. Gnoli & S. Lanciotti (Eds.), Orientalia Josephi Tucci Memoriae Dicata (Vol. 1, pp. 209–218). Istituto Italiano per il Medio ed Estremo Oriente.

Costantini, L. (1990a). Harappan agriculture in Pakistan: The evidence of Nausharo. In M. Taddei (Ed.), South Asian archaeology 1987 (pp. 321–332). Istituto italiano per il Medio ed Estremo Oriente.

Costantini, L. (1990b). Ecology and Farming of the Protohistoric Communities in the Central Yemeni Highlands. In The Bronze Age Culture of Hawlān at-Tiyāl and al-Hadā, A First General Report. (IsMEO, pp. 187–204).

Costantini, L., & Biasini, L. C. (1985). Le piante di Yelkhi. Ln Ferra Tra i Due Fiurrii. Venti Anni Di Archeologia Italiana in Medio Oriente. La Mesopotamia Dei Tesori. Torino: Il Quadrante Edizioni, 57.

Cleuziou, S. (1982). Hili and the beginning of oasis life in eastern Arabia. Proceedings of the Seminar for Arabian Studies, 12, 15–22. http://www.jstor.org/stable/41219277

Cleuziou, S. (1997). Construire et protéger son terroir: Les oasis d'Oman à l'Âge du Bronze. In J. Burnouf, J.-P. Bravard, & G. Chouquer (Eds.), Dynamique des paysages

protohistoriques antiques, médiévaux et modernes (pp. 389–412). APDCA.

Cleuziou, S. (2004). Pourquoi si tard? Nous avons pris un autre chemin. L'Arabie des chasseurs-cueilleurs de l'Holocène au début de l'Âge du Bronze. In J. Guilaine (Ed.), Aux marges des grands foyers du Néolithique: Périphéries débitrices ou créatrices? (pp. 123–148). Éditions Errance.

Colledge, S., & Conolly, J. (2014). Wild plant use in European Neolithic subsistence economies: A formal assessment of preservation bias in archaeobotanical assemblages and the implications for understanding changes in plant diet breadth. Quaternary Science Reviews, 101, 193–206. https://doi.org/10.1016/j.quascirev.2014.07.013

Dabrowski, V. (2019). Systèmes d'approvisionnement et gestion des ressources végétales en Arabie orientale aux périodes antique et islamique (IVème s. Av. J.-C. - XVIème s. Ap. J.-C.): Approches archéobotanique et archéoentomologique [PhD thesis, Muséum national d'Histoire naturelle]. https://theses.fr/2019MNHN0003

Dabrowski, V., Bouchaud, C., Desormeau, X., Herveux, L., Chambraud, E., Ryan, S. E., & Tengberg, M. (2024). A tale of new crops in the arid Arabian Peninsula oasis from antiquity to the early Islamic period. Vegetation History and Archaeobotany, 1–14. https://doi.org/10.1007/s00334-023-00976-4

de Maigret, A., Antonini, S., & Gnoli, G. (1986). YEMEN: Archaeological Activities in the Yemen Arab Republic. East and West, 36(4), 376–470.

Deckers, K., Döpper, S., & Schmidt, C. (2019). Vegetation, land, and wood use at the sites of Bat and Al‑Khashbah in Oman (fourth–third millennium BC). Arabian Archaeology and Epigraphy, 30(1), 1–14. https://doi.org/10.1111/aae.12123

Dinies, M., Neef, R., Plessen, B., & Kürschner, H. (2016). Holocene Vegetation, Climate, Land Use and Plant Cultivation in the Tayma Region, Northwestern Arabia. Proceedings of the International Congress Held at the University of Vienna, 5-8 December, 2013, 4, 57–78.

Engel, T. (1993). Charcoal remains from an Iron Age copper smelting slag heap at Feinan, Wadi Arabah (Jordan). Vegetation History and Archaeobotany, 2(4), 205–211. https://doi.org/10.1007/BF00198162

Flowers, J. M., Hazzouri, K. M., Gros-Balthazard, M., Mo, Z., Koutroumpa, K., Perrakis, A., Ferrand, S., Khierallah, H. S. M., Fuller, D. Q., Aberlenc, F., Fournaraki, C., & Purugganan, M. D. (2019). Cross-species hybridization and the origin of North African date palms. Proceedings of the National Academy of Sciences, 116(5), 1651–1658. https://doi.org/10.1073/pnas.1817453116

Fuller, D. Q. (2018). Long and attenuated: Comparative trends in the domestication of tree fruits. Vegetation History and Archaeobotany, 27(1), 165–176. https://doi.org/10.1007/s00334-017-0659-2

Fuller, D. Q., & Stevens, C. J. (2019). Between domestication and civilization: The role of agriculture and arboriculture in the emergence of the first urban societies. Vegetation History and Archaeobotany, 28(3), 263–282. https://doi.org/10.1007/s00334-019-00727-4

Fuller, Dorian Q and Weber, Steven A. (2005) Formation Processes and Paleoethnobotanical Interpretation in South Asia. Journal of Interdisciplinary Studies in History and Archaeology 2(1): 91-114.

Fuks, D., Schmidt, F., García‑Collado, M. I., Besseiche, M., Payne, N., Bosi, G., Bouchaud, C., Castiglioni, E., Dabrowski, V., Frumin, S., Fuller, D. Q., Hovsepyan, R., Muthukumaran, S., Peña-Chocarro, L., Jordá, G. P., Ros, J., Rottoli, M., Ryan, P., Spengler, R., … Gros-Balthazard, M. (2024). Orphan crops of archaeology‑based crop history research. PLANTS, PEOPLE, PLANET, ppp3.10468. https://doi.org/10.1002/ppp3.10468

Galili, E., Weinstein-Evron, M., Hershkovitz, I., Gopher, A., Kislev, M., Lernau, O., Kolska-Horwitz, L., & Lernau, H. (1993). Atlit-Yam: A Prehistoric Site on the Sea Floor off the Israeli Coast. Journal of Field Archaeology, 20(2), 133–157. https://doi.org/10.2307/529950

Garnier, Simon, Ross, Noam, Rudis, Robert, Camargo, Pedro, A., Sciaini, Marco, Scherer, & Cédric. (2024). viridis(Lite)—Colorblind-friendly color maps for R [Manual]. https://doi.org/10.5281/zenodo.4679423

Gillet, J. B. (1981). Botanical samples. In F. Safar, M. A. Muṣṭafá, & S. Lloyd (Eds.), Eridu (pp. 317–318). Republic of Iraq, Ministry of Culture and Information, State Organization of Antiquites and Heritage.

Görsdorf, J., & Vogt, B. (2001). Excavations at Ma'layba and Sabir, Republic of Yemen: Radiocarbon Datings in the Period 1900 to 800 cal BC. Proceedings of the 17th International Radiocarbon Conference (Part 3 of 3), 43, 1353–1361. https://doi.org/10.1017/S0033822200038595

Gros‑Balthazard, M., Battesti, V., Ivorra, S., Paradis, L., Aberlenc, F., Zango, O., Zehdi‑Azouzi, S., Moussouni, S., Naqvi, S. A., Newton, C., & Terral, J. (2020). On the necessity of combining ethnobotany and genetics to assess agrobiodiversity and its evolution in crops: A case study on date palms (*Phoenix dactylifera* L.) in Siwa Oasis, Egypt. Evolutionary Applications, 13(8), 1818–1840. https://doi.org/10.1111/eva.12930

Gros-Balthazard, M., & Flowers, J. M. (2021). A Brief History of the Origin of Domesticated Date Palms. In J. M. Al-Khayri, S. M. Jain, & D. V. Johnson (Eds.), The Date Palm Genome, Vol. 1: Phylogeny, Biodiversity and Mapping (pp. 55–74). Springer International Publishing. https://doi.org/10.1007/978-3-030-73746-7_3

Gros-Balthazard, M., Galimberti, M., Kousathanas, A., Newton, C., Ivorra, S., Paradis, L., Vigouroux, Y., Carter, R., Tengberg, M., Battesti, V., Santoni, S., Falquet, L., Pintaud, J.-C., Terral, J.-F., & Wegmann, D. (2017). The Discovery of Wild Date Palms in Oman Reveals a Complex Domestication History Involving Centers in the Middle East and Africa. Current Biology, 27(14), 2211-2218.e8. https://doi.org/10.1016/j.cub.2017.06.045

Gros-Balthazard, M., Newton, C., Ivorra, S., Pierre, M.-H., Pintaud, J.-C., & Terral, J.-F. (2016). The Domestication Syndrome in *Phoenix dactylifera* Seeds: Toward the Identification of Wild Date Palm Populations. PLoS ONE, 11(3), e0152394. https://doi.org/10.1371/journal.pone.0152394

Hajnalová, M., Miklíková, Z., & Belanová-Štolcová, T. (2009). Environmental research at al-Khiḍr, Failakah Island, Kuwait. Proceedings of the Seminar for Arabian Studies, 39, 197–202. https://www.jstor.org/stable/41223980

Hazzouri, K. M., Flowers, J. M., Visser, H. J., Khierallah, H. S. M., Rosas, U., Pham, G. M., Meyer, R. S., Johansen, C. K., Fresquez, Z. A., Masmoudi, K., Haider, N., El Kadri, N., Idaghdour, Y., Malek, J. A., Thirkhill, D., Markhand, G. S., Krueger, R. R., Zaid, A., & Purugganan, M. D. (2015). Whole genome re-sequencing of date palms yields insights into diversification of a fruit tree crop. Nature Communications, 6(1), 8824. https://doi.org/10.1038/ncomms9824

Hopf, M. (1969). Plant remains and early farming in Jericho. In P. J. Ucko & G. W. Dimbleby (Eds.), The domestication and exploitation of plants and animals (pp. 355–359). Gerald Duckworth & Co Ltd.

Hopf, M. (1983). Appendix B. Jericho Plant remains. In K. M. Kenyon & T. A. Holland (Eds.), Excavations at Jericho: Vol. Volume 5: The pottery phases of the tell and other finds (pp. 576–624). The British School of Archaeology in Jerusalem.

Ivorra, S., Tengberg, M., Bonhomme, V., Kaczmarek, T., Pastor, T., Terral, J.-F., &

Gros-Balthazard, M. (2024). Leveraging the potential of charred archaeological seeds for reconstructing the history of date palm. Journal of Archaeological Science, 170, 106052. https://doi.org/10.1016/j.jas.2024.106052

Jarrige, C., Jarrige, J.-F., Meadow, R., & Quivron, G. (Eds.). (1995). Mergharh. Mehrgarh. Department of Culture and Tourism, Government of Sindh. https://hal.science/halshs-03868706v1

Kaczmarek, T., Van Der Veen, M., Ivorra, S., Mattingly, D., Terral, J.-F., & Gros-Balthazard, M. (2024). Origins and evolution of oasis agriculture in the Sahara: Evidence from morphometric analyses of archaeological date palm seeds. The Holocene, 34(3), 353–365. https://doi.org/10.1177/09596836231211879

Kirleis, W., Kroll, H., Reiser, T., Schmid, C., & Schmütz, K. (2021). ArchbotLit—the archaeobotanical literature database: An update of the search engine for literature on archaeological remains of cultivated plants since 1981. Vegetation History and Archaeobotany, 30(1), 171–174. https://doi.org/10.1007/s00334-020-00794-y

Kislev, M. E., Hartmann, A., & Galili, E. (2004). Archaeobotanical and archaeoentomological evidence from a well at Atlit-Yam indicates colder, more humid climate on the Israeli coast during the PPNC period. Journal of Archaeological Science, 31(9), 1301–1310. https://doi.org/10.1016/j.jas.2004.02.010

Langgut, D. (2024). The core area of fruit-tree cultivation: Central Jordan Valley (Levant), ca. 7,000 BP. Palynology, 2347905. https://doi.org/10.1080/01916122.2024.2347905

Langgut, D., & Sasi, A. (2023). The Emergence of Fruit Tree Horticulture in Chalcolithic Southern Levant. In E. Ben-Yosef & I. W. N. Jones (Eds.), "And in Length of Days Understanding" (Job 12:12) (pp. 39–58). Springer International Publishing. https://link.springer.com/10.1007/978-3-031-27330-8_3

Levkovskaya, G. M., & Filatenko, A. A. (1992). Palaeobotanical and palynological studies in South Arabia. Review of Palaeobotany and Palynology, 73(1–4), 241–257. https://doi.org/10.1016/0034-6667(92)90061-K

Liphschitz, N. (1996). The vegetational landscape of the Negev during antiquity as evident from archaeological wood remains. Israel Journal of Plant Sciences, 44(2–3), 161–179. https://doi.org/10.1080/07929978.1996.10676645

Liphschitz, N., & Bonani, G. (2001). Wild and Cultivated Date Palm (*Phoenix Dactylifera*) from Qumran Cave 24. Tel Aviv, 28(2), 305–309. https://doi.org/10.1179/tav.2001.2001.2.305

Liphschitz, N., & Nadel, D. (1997). Charred Wood Remains from Ohalo II (19,000 B.P.), Sea of Galilee,. Journal of the Israel Prehistoric Society, 27, 5–18.

Liss, B., Howland, M. D., Lorentzen, B., Smitheram, C., Najjar, M., & Levy, T. E. (2020). Up the Wadi: Development of an Iron Age Industrial Landscape in Faynan, Jordan. Journal of Field Archaeology, 45(6), 413–427. https://doi.org/10.1080/00934690.2020.1747792

Litt, T., Ohlwein, C., Neumann, F. H., Hense, A., & Stein, M. (2012). Holocene climate variability in the Levant from the Dead Sea pollen record. Quaternary Science Reviews, 49, 95–105. https://doi.org/10.1016/j.quascirev.2012.06.012

Lodwick, L. (2019). Sowing the Seeds of Future Research: Data Sharing, Citation and Reuse in Archaeobotany. Open Quaternary, 5, 7. https://doi.org/10.5334/oq.62

Malleson, C., & Srour, F. J. (2024). "It's all just barley and figs!" Identifying patterns of plant waste accumulation in House 169, Elephantine Island, Egypt (1750–1650 bc) using machine learning. Vegetation History and Archaeobotany. https://doi.org/10.1007/s00334-024-01010-x

Marston, J.M., Guedes, J.D., Warinner, C. (Eds.), 2014. Method and Theory in Paleoethnobotany. University Press of Colorado.

Mashkour, M., Tengberg, M., Shirazi, Z., & Madjidzadeh, Y. (2013). Bio-archaeological studies at Konar Sandal, Halil Rud basin, southeastern Iran. Environmental Archaeology, 18(3), 222–246. https://doi.org/10.1179/1749631413Y.0000000006

Michel-Dansac, F., & Caubet, A. (2013). L'iconographie et le symbolisme du palmier dattier dans l'Antiquité (Proche-Orient, Égypte, Méditerranée orientale). Revue d'ethnoécologie, 4. https://doi.org/10.4000/ethnoecologie.1275

Miller, N. (1982). Economy and Environment of Malyan, A Third Millennium B.C. Urban Center in Southern Iran [PhD thesis, The University of Michigan]. https://hdl.handle.net/2027.42/102023

Miller, N., Jones, P., & Pittman, H. (2016). Sign and Image: Representations of Plants on the Warka Vase of Early Mesopotamia. Origini, 39, 53–73.

Mouton, M., Tengberg, M., Bernard, V., Maguer, S. L., Reddy, A., & Soulié, D. (2021). Building H at Mleiha: New evidence of the late pre-Islamic period D phase (PIR.D) in the Oman peninsula (second to mid-third century AD). Proceedings of the Seminar for Arabian Studies, 2012, Vol. 42, Papers from the forty-fifth meeting of the Seminar for Arabian Studies held at the British Museum, London, 28 to 30 July 2011 (2012), pp. 205-221. Published by Archaeopress. https://www.jstor.org/stable/41623638

Muṣṭafā Ḥasan, Ḥasan, al-Ḥasan, A. A. al-Qāsim, & Khaybar, ʿAbd al-Raḥīm Muḥammad. (2019). Plant remains of Qaryat al-Fāw, Pre-Islamic: Comestible plants and fruits. In A. R. T. al-Ansary (Ed.), Qaryat (al-Fāw) (Saudi Commission ofr Tourism and National Heritage, pp. 319–344). King Saud University.

Neef, R. (1991). Plant remains of archaeological sites in lowland Iraq: Tell el'Oueili. In J.-L. Huot (Ed.), 'Oueili. Travaux de 1985 (pp. 322–329). Editions recherche sur les civilisations.

Nesbitt, M., Colledge, S., & Murray, M. A. (2003). Organisation and Management of Seed Reference Collections. Environmental Archaeology, 8(1), 77–84. https://doi.org/10.1179/env.2003.8.1.77

Parker, A. G., Eckersley, L., Smith, M. M., Goudie, A. S., Stokes, S., Ward, S., White, K., & Hodson, M. J. (2004). Holocene vegetation dynamics in the northeastern Rub' al-Khali desert, Arabian Peninsula: A phytolith, pollen and carbon isotope study. Journal of Quaternary Science, 19(7), 665–676. https://doi.org/10.1002/jqs.880

Parker, A.G. (2010). Palaeoenvironmental evidence from H3, Kuwait. In Maritime Interactions in the Arabian Neolithic: Evidence from H3, AsSabiyah, an Ubaid-Related Site in Kuwait, R.A. Carter, and H.E.W. Crawford, eds. (Brill), pp. 189–201.

Paszke, M. Z. (2019). DATE PALM AND DATE PALM INFLORESCENCES IN THE LATE URUK PERIOD (C. 3300 B.C.): BOTANY AND ARCHAIC SCRIPT. Iraq, 81, 221–239. https://doi.org/10.1017/irq.2019.15

Patrich, J. (1994). Khirbet Qumran in Light of New Archaeological Explorations in the Qumran Caves. Annals of the New York Academy of Sciences, 722(1), 73–95. https://doi.org/10.1111/j.1749-6632.1994.tb30464.x

Pena-Chocarro, L., & Barron Lopez, E. (1999). Plant remains from the site of Mleiha. In M. Mouton (Ed.), Mleiha I. Environnement, stratégies de subsistance et artisanat (pp. 63–69). Maison de l'Orient et de la Méditerranée. https://www.persee.fr/doc/mom_1274-6525_1999_rpm_29_1_1887

Perrot, J., & Madjidzadeh, Y. (2005). L'iconographie des vases et objets en chlorite de Jiroft (Iran). Paléorient, 31(2), 123–152. https://doi.org/10.3406/paleo.2005.5129

Pérez-Escobar, O. A., Bellot, S., Przelomska, N. A. S., Flowers, J. M., Nesbitt, M., Ryan, P., Gutaker, R. M., Gros-Balthazard, M., Wells, T., Kuhnhäuser, B. G., Schley, R., Bogarín, D., Dodsworth, S., Diaz, R., Lehmann, M., Petoe, P., Eiserhardt, W. L.,

Preick, M., Hofreiter, M., … Baker, W. J. (2021). Molecular clocks and archeogenomics of a late period Egyptian date palm leaf reveal introgression from wild relatives and add timestamps on the domestication. Molecular Biology and Evolution, 38(10), 4475–4492. https://doi.org/10.1093/molbev/msab188

Pintaud, J.-C., Zehdi, S., Couvreur, T., Barrow, S., Henderson, S., Aberlenc-Bertossi, F., Tregear, J., & Billotte, N. (2010). Species Delimitation in the Genus *Phoenix* (Arecaceae) Based on SSR Markers, with Emphasis on the Identity of the Date Palm (*Phoenix dactylifera*). 20.

Rollefson, G. O., Simmons, A. H., Donaldson, M. L., Gillespie, W., Kafafi, Z., Köhler-Rollefson, I., McAdam, E., Rolston, S. L., & Tubb, K. M. (1985). Excavation at the Pre-Pottery Neolithic B (PPNB) village of ʿAin Ghazal (Jordan), 1983. Mitteilungen Der Deutschen Orient-Gesellschaft, 117, 69–116.

Rivera, D., Obón, C., García-Arteaga, J., Egea, T., Alcaraz, F., Laguna, E., Carreño, E., Johnson, D., Krueger, R., Delgadillo, J., & Ríos, S. (2014). Carpological analysis of *Phoenix* (Arecaceae): Contributions to the taxonomy and evolutionary history of the genus. Botanical Journal of the Linnean Society, 175(1), 74–122. https://doi.org/10.1111/boj.12164

Roh, T. (2024). leaflegend: Create custom legends for leaflet [Manual]. https://leaflegend.roh.engineering

Rohmer, J., Lesguer, F., Bouchaud, C., Purdue, L., Alsuhaibani, A., Tourtet, F., Monchot, H., Dabrowski, V., Decaix, A., Desormeau, X., Alkhatib Alkontar, R., & Reiller, H. (2022). New clues to the development of the oasis of Dadan. Results from a test excavation at Tall al-Sālimīyyah (al-ʿUlā, Saudi Arabia). In R. Foote, M. Guagnin, I. Périssé, & S. Karacic (Eds.), Revealing Cultural Landscapes in Northwest Arabia. (pp. 157–190). Archaeopress. https://cnrs.hal.science/hal-03930118v1

Ros, J., Badri, F.-E., & Pelling, R. (2024). Agrobiodiversity and crop diffusion in Morocco from Antiquity to the early Modern period (8th century bce–17th century ce): An archaeobotanical review. Vegetation History and Archaeobotany. https://doi.org/10.1007/s00334-024-01013-8

Schiettecatte, J. (2013). Le palmier dattier (*Phoenix dactylifera* L.) dans l'Arabie méridionale préislamique. Revue d'ethnoécologie, 4. https://doi.org/10.4000/ethnoecologie.1356

Schiettecatte, J., Al-Ghazzi, A., Bouchaud, C., Chabrol, A., Crassard, R., Cuny, J., Dinies, M., Hilbert, Y., Monchot, H., Sagory, T., & Siméon, P. (2015). Preliminary Report. Fourth season of the Saudi-French mission in al-Kharj, Province of Riyadh, 23 January—27 February 2015 [Research Report]. CNRS, UMR8167 Orient & Méditerranée. https://shs.hal.science/halshs-01186755

Schiettecatte J. & A. al-Ghazzi (dir.), (2016). Al-Kharj I. Report on two excavation seasons in the oasis of al-Kharj (2011–2012). Saudi Arabia (Series of Archaeological Refereed studies, No. 40), Riyad, Saudi Commission for Tourism and National Heritage, 378 pp. [Pdf: https://halshs.archives-ouvertes.fr/halshs-01678594].

Sievert, C. (2020). Interactive web-based data visualization with R, plotly, and shiny. Chapman and Hall/CRC. https://plotly-r.com

Sievert, C., Cheng, J., & Aden-Buie, G. (2024). bslib: Custom "bootstrap" "sass" themes for "shiny" and "rmarkdown" [Manual]. https://CRAN.R-project.org/package=bslib

Silva, F., Stevens, C. J., Weisskopf, A., Castillo, C., Qin, L., Bevan, A., & Fuller, D. Q. (2015). Modelling the Geographical Origin of Rice Cultivation in Asia Using the Rice Archaeological Database. PLoS ONE, 10(9), e0137024. https://doi.org/10.1371/journal.pone.0137024

Stroth, L., Garvin Suero, A., Liss, B., Howland, M. D., & D'Alpoim Guedes, J. (2023). Archaeological Evidence of Casual Snacking and Resource Provisioning at Khirbat

al-Jariya (ca. Eleventh to Tenth Centuries BCE), an Iron-Age Copper Production Site. In E. Ben-Yosef & I. W. N. Jones (Eds.), "And in Length of Days Understanding" (Job 12:12): Essays on Archaeology in the Eastern Mediterranean and Beyond in Honor of Thomas E. Levy (pp. 1133–1153). Springer. https://doi.org/10.1007/978-3-031-27330-8_48

Tengberg, M. (1998). Paléoenvironnements et économie végétale en milieu aride et semiaride: Recherches archéobotaniques dans la région du Golfe arabo-persique et dans le Makran pakistanais (4ème millénaire av. Notre ère e 1er millénaire de notre ère) [PhD thesis, Université Montpellier 2]. https://www.theses.fr/1998MON20266

Tengberg, M. (1999a). L'exploitation des ligneux à Mleiha. Étude anthracologique. In M. Mouton (Ed.), Mleiha I. Environnement, stratégies de subsistance et artisanat (pp. 71–82). Maison de l'Orient et de la Méditerranée. https://www.persee.fr/doc/mom_1274-6525_1999_rpm_29_1_1888

Tengberg, M. (1999b). Crop husbandry at Miri Qalat Makran, SW Pakistan (4000–2000 B.C.). Vegetation History and Archaeobotany, 8(1–2), 3–12. https://doi.org/10.1007/BF02042836

Tengberg, M. (2012). Beginnings and early history of date palm garden cultivation in the Middle East. Journal of Arid Environments, 86, 139–147. https://doi.org/10.1016/j.jaridenv.2011.11.022

Tengberg, M., & Lombard, P. (2001). Environnement et économie végétale à Qal'at al-Bahreïn aux périodes Dilmoun et Tylos. Premiers éléments d'archéobotanique. Paléorient, 27(1), 167–181. https://doi.org/10.3406/paleo.2001.4727

Tengberg, M., & Newton, C. (2016). Origine et évolution de la phéniciculture au Moyen-Orient et en Égypte. In M.-P. Ruas (Ed.), Des fruits d'ici et d'ailleurs: Regards sur l'histoire de quelques fruits consommés en Europe. Mouans-Sartoux, Éditions Omniscience.

Terral, J.-F., Tabard, E., Bouby, L., Ivorra, S., Pastor, T., Figueiral, I., Picq, S., Chevance, J.-B., Jung, C., Fabre, L., Tardy, C., Compan, M., Bacilieri, R., Lacombe, T., & This, P. (2010). Evolution and history of grapevine (Vitis vinifera) under domestication: New morphometric perspectives to understand seed domestication syndrome and reveal origins of ancient European cultivars. Annals of Botany, 105(3), 443–455. https://doi.org/10.1093/aob/mcp298

Terral, J.-F., Tabard, E., Bouby, L., Ivorra, S., Pastor, T., Figueiral, I., Picq, S., Chevance, J.-B., Jung, C., Fabre, L., Tardy, C., Compan, M., Bacilieri, R., Lacombe, T., & This, P. (2010). Evolution and history of grapevine (Vitis vinifera) under domestication: New morphometric perspectives to understand seed domestication syndrome and reveal origins of ancient European cultivars. Annals of Botany, 105(3), 443–455. https://doi.org/10.1093/aob/mcp298

Thomas, R. (2013). Anatomy of the endemic palms of the Near and Middle East: Archaeobotanical perspectives. Revue d'ethnoécologie, 4. https://doi.org/10.4000/ethnoecologie.1366

Tourtet, F., Daszkiewicz, M., Hausleiter, A. (2021). Pottery from Tayma: Chronostratigraphy, Archaeometric Studies, Cultural Interaction, in: Luciani, M. (Ed.), The Archaeology of the Arabian Peninsula 2. Connecting the Evidence. Proceedings of the Workshop Held at the 10th ICAANE in Vienna, April 2016, Oriental and Europaean Archaeology. Austrian Academy of Science Press, Vienna, pp. 43–88.

Van der Veen, M. (1992). Botanical evidence for Garamantian agriculture in Fezzan, southern Libya. Review of Palaeobotany and Palynology, 73(1–4), 315–327. https://doi.org/10.1016/0034-6667(92)90066-P

Van der Veen, M. (2007). Formation processes of desiccated and carbonized plant remains –

the identification of routine practice. Journal of Archaeological Science, 34(6), 968–990. https://doi.org/10.1016/j.jas.2006.09.007

Van Der Veen, M. (2011). Consumption, Trade and Innovation (Vol. 6). Africa Magna Verlag. ISBN 978-3-937248-23-3.

Van der Veen, M., 2018. Archaeobotany: the archaeology of human-plant interactions, in: Scheidel, W. (Ed.), The Science of Roman History. Biology, Climate, and the Future of the Past. Princeton University Press, Princeton, pp. 53–95.

Vogt, B., & Sedov, A. (1998). The Sabir culture and coastal Yemen during the second millennium BC - the present state of discussion. Proceedings of the Seminar for Arabian Studies, 28, 261–270. https://www.jstor.org/stable/41223633

Weiss, E. (2015). Beginnings of Fruit Growing in the Old World two generations later. Israel Journal of Plant Sciences, 62(1–2), 75–85. https://doi.org/10.1080/07929978.2015.1007718

Western, C. A. (1971). The Ecological Interpretation of Ancient Charcoals from Jericho. Levant, 3(1), 31–40. https://doi.org/10.1179/lev.1971.3.1.31

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019). Welcome to the tidyverse. Journal of Open Source Software, 4(43), 1686. https://doi.org/10.21105/joss.01686

Wilkinson, M. D., Dumontier, M., Aalbersberg, Ij. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., Da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., … Mons, B. (2016). The FAIR Guiding Principles for scientific data management and stewardship. Scientific Data, 3(1), 160018. https://doi.org/10.1038/sdata.2016.18

Xie, Y., Allaire, J., & Horner, J. (2024). markdown: Render markdown with "commonmark" [Manual]. https://CRAN.R-project.org/package=markdown

Zaitschek, D. V. (1961). Remains of Cultivated Plants from the Caves of Naḥal Mishmar: Preliminary Note. Israel Exploration Journal, 11(1/2), 70–72.

Zehdi-Azouzi, S., Cherif, E., Moussouni, S., Gros-Balthazard, M., Abbas Naqvi, S., Ludeña, B., Castillo, K., Chabrillange, N., Bouguedoura, N., Bennaceur, M., Si-Dehbi, F., Abdoulkader, S., Daher, A., Terral, J.-F., Santoni, S., Ballardini, M., Mercuri, A., Ben Salah, M., Kadri, K., … Aberlenc-Bertossi, F. (2015). Genetic structure of the date palm (*Phoenix dactylifera*) in the Old World reveals a strong differentiation between eastern and western populations. Annals of Botany, 116(1), 101–112. https://doi.org/10.1093/aob/mcv068

Zohary, D., Hopf, M., & Weiss, E. (2012). Domestication of Plants in the Old World: The origin and spread of domesticated plants in Southwest Asia, Europe, and the Mediterranean Basin. Oxford: Oxford University Press. https://doi.org/10.1093/acprof:osobl/9780199549061.001.0001

Zohary, D., & Spiegel-Roy, P. (1975). Beginnings of Fruit Growing in the Old World: Olive, grape, date, and fig emerge as important Bronze Age additions to grain agriculture in the Near East. Science, 187(4174), 319–327. https://doi.org/10.1126/science.187.4174.319