Co-expression of Bovine leukemia virus and Bovine foamy virus-derived miRNAs in naturally infected cattle

Petersen MI[1,2], Suarez Archilla G[3], Perez CV[1,2], Gonzalez DD[1,2], Trono KG[1,2], Mongini C[1,2], Jaworski JP[1,2], Carignano HA[1,2*].

Affiliations

[1] Instituto de Virología e Innovaciones Tecnológicas, Instituto Nacional de Tecnología Agropecuaria - Consejo Nacional de Investigaciones Científicas y Técnicas, B1686 Hurlingham, Argentina.

[2] Consejo Nacional de Investigaciones Científicas y Técnicas, C1033AAJ Ciudad Autónoma de Buenos Aires, Argentina.

[3] Instituto de Investigación de la Cadena Láctea, Estación Experimental Agropecuaria Rafaela, Instituto Nacional de Tecnología Agropecuaria - Consejo Nacional de Investigaciones Científicas y Técnicas, S2300 Rafaela, Argentina.

[*]Address correspondence to Hugo A. Carignano, carignano.hugo@inta.gob.ar

Abstract

MicroRNAs (miRNAs) derived from viruses play a central role in modulating the expression of both viral and host gene, facilitating immune evasion, viral persistence, and pathogenesis. Among the viruses that encode miRNAs and infect cattle such as Bovine leukemia virus (BLV), Bovine foamy virus (BFV), Bovine herpesvirus type 1 (BoHV-1), and Bovine herpesvirus type 5 (BoHV-5), BLV has gained increasing attention due to the critical role that BLV-miRNAs may play in inducing leukemia and lymphoma in infected animals. BLV is highly prevalent in the Americas and has a negatively affects dairy herds, primarily due to restrictions on the commercialization of dairy products from infected animals and a decrease in milk production.

Mixed infections involving BLV and BFV appear to be common in cattle. Considering the ability of foamy viruses to cross species barriers, preventing their presence within the food chain is essential. In this study, through small RNA sequencing, we identified the co-expression of seven BLV-derived miRNAs (*blv-miR-B1-3p*, *blv-miR-B2-5p*, *blv-miR-B2-3p*, *blv-miR-B3-5p*, *blv-miR-B3-3p*, *blv-miR-B4-3p,* and *blv-miR-B5-5p*) and three BFV-derived (*bfv-miR-BF1-5p*, *bfv-miR-BF1-3p*, and *bfv-miR-BF2-5p*) in naturally BLV-infected cattle, while no BoHV-1/5 miRNAs were detected. A differential expression analysis revealed seven bovine miRNAs (*bta-miR-375*, *bta-miR-133a*, *bta-miR-677*, bta-miR-1, *bta-miR-3613a*, *bta-miR-9-5p*, and *bta-miR-95*) with altered expression levels (fold change > |1.5| and q-value < 0.05) between cows with a high BLV proviral load (PVL) and uninfected counterparts. A comprehensive functional analysis of protein-protein interaction networks for genes targeted by both viral and host-derived miRNAs highlighted key pathways implicated in tumorigenesis and the regulation of the immune response. Notably, *bta-miR-375*, previously proposed as an early biomarker for the progression of enzootic bovine leukosis progression, exhibited differential expression in cows with elevated PVL, reinforcing its diagnostic potential. The functional interaction among miRNAs derived from BLV and BFV raises important concerns regarding their potential influence on disease severity and the increased dissemination of BFV. The findings derived from this study offer new perspectives for creating diagnostic and treatment approaches to manage viral persistence and tumor growth in cattle.

Introduction

The study of virus-host interactions has traditionally focused on how viral proteins modulate host proteins to hijack cellular processes in favor of viral replication and pathogenesis (Ruivinho and Gama-Carvalho, 2023). However, like their host cells, many viruses encode regulatory non-coding RNAs, including small RNAs such as microRNAs (miRNAs), which provide new capacities for regulating host gene expression to ensure successful viral infection (Sullivan and Ganem, 2005; Samols et al., 2005; Grundhoff and Sullivan, 2011; Guo and Steitz, 2014). miRNAs are RNA fragments, typically 19–24 nucleotides (nt) long, that fine-tune gene expression at the post-transcriptional level (Bartel, 2004; Griffiths-Jones et al., 2006). The canonical biogenesis of a mature miRNA begins with the transcription of a precursor in the cell

nucleus mediated by RNA polymerase II. This precursor, known as pri-miRNA, is processed by the RNase III enzyme Drosha into a characteristic stem-loop structure called pre-miRNA (approximately 60 nt). Once exported to the cytoplasm by Exportin-5, the loop of the pre-miRNA is cleaved by the RNase III-like enzyme Dicer, yielding a mature miRNA duplex of approximately 20 nucleotides. The regulatory function of miRNAs is primarily mediated by the RNA-induced silencing complex (RISC), which facilitates base-pairing between the mature miRNA´s 2–7 nucleotide seed region and the 3' untranslated region (3'UTR) of the target messenger RNA (mRNA) (Kim, 2005; Bartel, 2009). The miRNA-mRNA interaction induces mRNA degradation by cellular endonucleases or translational repression (He and Hannon, 2004; Bartel et al., 2009). It is estimated that miRNAs may regulate at least 60% of the human coding genome (Friedman et al., 2009; Bartel et al., 2018).

Host-derived microRNAs (miRNAs) regulate viral infections by exhibiting a dual nature that can benefit either the virus or the host. For example, host miRNAs that directly target viral RNA often enhance the viral life cycle. Conversely, host miRNAs can exert an indirect negative effect on the virus by targeting mRNAs that encode essential host factors necessary for the viral cycle or for establishing immune responses and defense mechanisms (Guo and Steitz, 2014; Girardi et al., 2018). Given their high versatility in fine-tuning gene expression and their small coding size, it is not surprising that viruses have evolved the ability to express their own miRNAs (Skalsky and Cullen, 2010). Since the first report of miRNAs encoded by the Epstein-Barr virus (EBV) (Pfeffer et al., 2004), miRNAs encoded by DNA viruses have proven to be the most abundant, particularly those expressed by the Herpesviridae family. In contrast, RNA viruses that express miRNAs are far less common and are typically restricted to those with a DNA intermediate. Additionally, several miRNA-like molecules have been identified in both positive and negative-sense RNA viruses (Pfeffer et al., 2005; Sullivan et al., 2005; Seo et al., 2008; Ghosh et al., 2009; Hussain et al., 2012, 2014; Liang et al., 2014; Kinkaid et al., 2012, 2014; Whisnant et al., 2014; Yao et al., 2014; Teng et al., 2015; Cao et al., 2018; Hashimoto-Gotoh et al., 2020; Aso et al 2021).

Usually, viral miRNAs follow the same biogenesis and effector pathways as host miRNAs. However, certain retroviruses, such as members of the Spumaretrovirinae subfamily (e.g., Simian foamy virus (SFV) and Bovine foamy virus (BFV)) and the Orthoretrovirinae subfamily (e.g., Bovine leukemia virus (BLV)), decouples miRNA transcription from genomic transcription by employing RNA polymerase III instead of RNA polymerase II (Kinkaid et al., 2012, 2014; Whisnant et al., 2014). This mechanism likely prevents the auto-degradation of viral genomes mediated by endonucleases during miRNA biogenesis. (Hartwig, 2014; Aguado and tenOever, 2018).

Generally, the functional role of viral miRNAs involves the up-regulation or down-regulation of viral and host protein expression to regulate the viral life cycle (e.g., promoting or inhibiting cell proliferation and apoptosis), evade the host immune response, and establish latent or persistent infections (Seo et al., 2008; Cullen, 2009; Grundhoff and Sullivan, 2011; Cullen, 2013; Tycowski et al., 2015). This is particularly evident in retroviruses and herpesviruses, where viral protein expression is strongly restricted

during latency. The production of a low-antigenicity regulatory molecule provides a significant advantage for viral persistence while evading host immune detection (Skaslky and Cullen, 2010; Pluta et al 2020).

Viruses that encode miRNAs and infect cattle include BLV (Kinkaid et al., 2012), BFV (Whisnant et al., 2014), Bovine herpesvirus 1 (BoHV-1) (Glazov et al., 2010), and Bovine herpesvirus 5 (BoHV-5) (Tang et al., 2014). BoHV-1 encodes at least ten miRNAs, two of which have been implicated in maintaining latency both *in vivo and in vitro* (Jaber et al., 2010; da Silva and Jones, 2012). In contrast, the functional roles and target genes of the three BFV-derived miRNAs remain poorly understood (Cao et al., 2020). Among BLV-encoded miRNAs, blv-miR-B4-3p is the most extensevely studied. It shares a seed sequence with miR-29a, an oncomiR that targets tumor suppressors such as peroxidasin homolog (*PXDN*) and HMG-box transcription factor 1 (*HBP1*), as reported in murine tumor-induction studies (Han et al., 2010; Santanam et al., 2010). *In vitro* reporter assays have demonstrated that *blv-miR-B4-3p* exerts a repressive effect on the expression of *HBP1* and *PXDN*. However, a negative correlation between *blv-miR-B4-3p* and *HBP1* expression has been documented, while levels of *PXDN* remained unchanged in ovine primary tumor B cells (Kinkaid et al., 2012; Rosewick et al., 2013). Additionally, recent findings indicate that *PXDN* expression is significantly downregulated in naturally BLV-infected cattle expressing *blv-miR-B4* when compared to their non-infected counterparts. Notably, *miR-29a* expression remains unaffected in both groups (Petersen et al., 2023).

The BLV prevalence reaches up to 85% in North American at farms-level and over 80% in lactating cows of Argentina's main dairy region (VanLeeuwen et al., 2005; 2006; NAHMS, 2007; Gutierrez et al., 2012). International trade restrictions of livestock products from affected herds impact negatively the economy (OIE, 2012), but greater losses stem from reduced milk production and earlier culling of asymptomatic BLV carriers compared to BLV-free herds (Erskine et al., 2012; Bartlett et al., 2013).

Mixed infections involving BLV and BFV appear to be common in cattle (Materniak et al., 2019). Given the ability of foamy viruses (FVs) to cross species barriers and cause zoonotic infections in humans, their presence in the food chain may be a potential risk to both human and animal health. Furthermore, cattle infected with BLV are more susceptible to co-infections with BoHV-1 (de Brun et al., 2023; Mendez et al., 2023; Oviedo-Pastrana et al., 2024).

This study identified the expression of three BFV-derived miRNAs and seven BLV-derived miRNAs in naturally BLV-infected cattle through small RNA analysis sequencing. No miRNAs from BoHV-1/5 were found. Seven bovine miRNAs showed differential expression between cattle infected with BLV/BFV and those uninfected with BLV. A functional analysis of the protein-protein interaction network for genes potentially targeted by both viral and host-derived miRNAs revealed key metabolic pathways associated with tumorigenesis and immune response regulation.

Methods

Selection of Animals

Briefly, adult cows (over 3 years old, >1 lactation) were sampled from the central region of Argentina, an area known for its high dairy farming activity, where the average individual prevalence of BLV exceeds 80% (Gutiérrez et al., 2012). Cows were screened over a 10-month period (T-10, T-5, T-3, T0 months before final sampling) to evaluate both antibody levels and proviral load using anti-BLV ELISA and quantitative PCR (qPCR) assays, respectively (details provided later). All sample collection and handling procedures adhere to the recommendations and guidelines of the Institutional Committee for the Care and Use of Experimental Animals at the Instituto Nacional de Tecnología Agropecuaria.

Isolation of Peripheral Blood Mononuclear Cells (PBMCs)

Fresh blood samples from animals were collected via jugular venipuncture and supplemented with EDTA (225 μM). Peripheral blood mononuclear cells (PBMCs) were isolated on the same day of collection using Ficoll-Paque Plus (GE Healthcare, Uppsala, Sweden) density gradient centrifugation, following the manufacturer's protocol. The plasma fraction was kept for anti-BLV ELISA serology. After isolation, PBMCs were preserved in RNAlater solution (Ambion, Austin, TX) and stored at −80°C until further use.

Serology

The anti-BLV ELISA assay, as described by Trono et al. (2001), was employed to identify BLV-infected animals. The whole BLV viral particle served as the antigen for detecting plasma anti-BLV antibodies in each sample. Briefly, a sample-to-positive (S/P) ratio, referred to as the percentage of reactivity (PR), was calculated using the following formula: $S/P = [(OD_{Sample} - OD_{WS-}) / (OD_{WS+1/7} - OD_{WS-})] \times 100$, where OD refers to optical density, WS- indicates the negative serum, and WS+1/7 is the international standard weak positive control serum (diluted 1:7 in negative serum). Samples with a PR greater than 25% were considered positive (BLV(+)), while those below were regarded as negative (BLV(-)).

BLV proviral load (PVL) Quantification

Genomic DNA was extracted from PBMCs using the High Pure PCR Template Preparation Kit (Roche, Penzberg, Germany) following the manufacturer's instructions. The quality and concentration of genomic DNA from whole blood samples, extracted with the Blood Genomic DNA AxyPrep™ kit (Axygen Biosciences, Union City, USA), were assessed using a microvolume spectrophotometer (NanoDrop™ Technologies, Inc., Wilmington, USA).

A BLV POL gene-based PVL qPCR assay based on the SYBR Green dye detection system was conducted as described by Petersen et al. (2018). Each 25 μL qPCR reaction contained Fast Start Universal SYBR Green Master Mix (2×; Roche), forward and reverse primers (800 nM; BLVpol_5f:5′-

CCTCAATTCCCTTTAAACTA-3′ and BLVpol_3r:5′-GTACCGGGAAGACTGGATTA-3′; Thermo Fisher

Scientific), and 200 ng of genomic DNA template. Amplification and detection were carried out using a

Step One Plus system (Applied Biosystems, Foster City, CA). The specificity of each BLV-positive

reaction was confirmed by melting temperature dissociation curve (Tm) analysis. Based on the

assumption of a low natural infection rate (1% of BLV-infected cells in the peripheral blood), PVL values

below 1,500 copies/μg of total DNA were classified as low, while values above this threshold were

classified as high (Hopkins and DiGiacomo, 1997).

Small RNA Sequencing

Total RNA was extracted from PBMCs using the miRNeasy kit (Qiagen, Hilden, Germany) according to

the manufacturer's protocol. RNA quality (integrity) and concentration were assessed via digital

electrophoresis on an Agilent 2200 TapeStation system (Agilent Technologies, Santa Clara, US).

The QIAseq miRNA Library Kit (Qiagen, Hilden, Germany) was used for miRNA sequencing library

construction, which integrates Unique Molecular Indices (UMIs) tags into the adapters during the cDNA

synthesis step. These UMI tags help to mitigate PCR amplification biases and sequencing artifacts by

allowing identification and collapsing reads with the same amplification origin. A specific 8-nt barcode

was assigned to each sample for multiplexing. The multiplexed libraries were pooled to an equal molar

concentration and processed in a NovaSeq run (Illumina, San Diego, US) with a 75x50 bp configuration

(300–400 million reads pairs per lane).

Small RNA sequencing reads processing and deduplication

The quality control of sequencing reads was performed sequentially to ensure reliable reads for

unbiased miRNome identification and quantification. The 12-nt UMI pattern was extracted from each

read using the umi_tools v1.1.6 software (Smith et al., 2017). Forward (Fw) and reverse (Rv) reads were

processed independently before collapsing. Reads with low quality (quality cutoff < 30), a minimum

length < 16 nt, the presence of ambiguous bases (N) or adapter sequences were removed using

Cutadapt v5.0 (Martin, 2011).

Singleton reads resulting were discarded, and only paired Fw and Rv reads were considered. Forward

read sequences were aligned to the Rv-complement and merged into a single consensus read using

BBMap v25.85 (Bushnell, 2014) with a minimum insert size of 16 nt and a minimum overlap of 18 nt.

After merging, Qiagen-specific adapter sequences were removed using Cutadapt v5.0 (Martin, 2011).

A stringent mapping strategy was employed for read deduplication using Bowtie2 v2.5.4 (Langmead and

Salzberg, 2012). Reads were aligned to the bovine reference genome ARS-UCD1.2 (acc.

GCF_002263795.1) and to viral genomes encoding miRNAs that infect cattle, including BLV (acc.

NC_001414.1), BFV (acc. NC_001831.1), BoHV-1 (acc. NC_063268.1) and BoHV-5 (acc.

NC_005261.3). No mismatches were allowed in the first 18 bases of the read (left end). After alignment

to the reference sequences, reads were deduplicated based on their mapping coordinates and the

previously extracted UMI tags using UMICollapse v1.0.0 (Liu, 2019).

miRNAs identification and quantification

Deduplicated reads were mapped to the reference precursor miRNA sequences for bovine, BLV, BFV,

BoHV-1 and BoHV-5 obtained from miRBase v22.1 (Kozomara et al., 2019). Mature miRNA expression

was quantified using miRDeep2 v0.1.3 (Friedländer et al., 2012) with default parameters, generating a

count matrix of miRNA expression for each sample.

Differential miRNA Expression analysis

The raw count matrix for all samples was analyzed using DESeq2 v1.44.0 (Love et al., 2014) to assess

differential miRNA expression between BLV-infected and non-infected animals. Based on miRNA

expression profiles, an exploratory analysis of variation patterns between infected and non-infected

animals was performed using principal component analysis (PCA). To ensure homoskedasticity of the

data, raw miRNA counts were transformed using the Variance Stabilizing Transformation (VST) method

implemented in DESeq2 v1.44.0. Pairwise distances between samples were calculated based on the

root-mean-square deviation of miRNA expression levels.

To test for differential miRNA expression, the DESeq2 model adjusts miRNA counts by accounting for

sample-specific differences, such as library size, sequencing composition, and miRNA-specific biases,

using negative binomial models. Under the null hypothesis, the model assumes that a miRNA has the

same mean expression across the two conditions (BLV(-) and BLV(+) groups). The statistical

significance of the Log_2 fold change (Log_2FC) between conditions was assessed using the Wald test.

Correction for multiple comparisons was performed using the Benjamini-Hochberg false discovery rate

(BH-FDR) method (Benjamini and Hochberg, 1995). The miRNAs with BH-FDR-adjusted p-values (q) <

0.05 and a $Log_2FC > |1.5|$ were considered statistically significant. The PCA and Volcano plots were

performed under the R environment (R Core Team, 2024).

miRNAs gene target prediction

Significant differentially expressed bovine and viral-derived miRNAs between BLV(+) and non-infected

cows were used to predict potential bovine gene targets *in silico* (miRNA:mRNA interactions) using

miRanda v3.3 with default parameters (Enright et al., 2003). The analysis utilized 3′-UTR sequences of

21,400 bovine genes from the ARS-UCD1.2 annotation (Rosen et al., 2020). The miRanda algorithm

evaluates potential gene targets based on seed-site alignment and the thermodynamic stability of RNA-

RNA duplexes. Predicted targets were filtered using a score threshold of ≥140 and a free energy ≤−20

kcal/mol.

Gene ontology and graph-based pathway analysis of miRNA target genes

The functional annotation of the potential target genes was carried out using STRINGdb (Szklarczyk et al., 2021) with Gene Ontology (GO) terms and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways (Kanehisa and Gotoh, 2000). Next, a protein-protein interaction (PPI) network of the identified target genes was constructed using STRINGdb. This platform integrates curated and predicted interactions from multiple sources, including experimental evidence, computational predictions, scientific literature, co-expression data and primary public databases. Highly interconnected protein clusters within the network were identified using the Markov Cluster Algorithm (MCL) with default parameters (inflation parameter = 2.5). Only clusters with more than 15 nodes (genes) were considered (Dongen et al., 2008). GO terms and KEGG pathways overrepresentation analysis was performed on the identified clusters using Fisher's exact test based on the hypergeometric distribution, using a term similarity threshold > 0.7 (Tavazoie et al., 1999). Wordclouds and Treemaps plots of significant GO terms grouped by semantic similarity were obtained using the rrvgo package with default parameters (Sayols, 2023).

Results

In this study, four animals consistently exhibited both high antibody reactivity across all sampling times (5671 = 178.6 ± 28.4; 5841 = 191.3 ± 32.8; 6021 = 169.8 ± 42.7 and 6097 = 183.2 ± 25.4) and high proviral loads at T-3 and T0 (average proviral load: 5671 = 63,148.0; 5841 = 44,234.3; 6021 = 52,902.4 and 6097 = 91,255.4). These cows were classified as BLV(+). In contrast, three animals that consistently tested negative for both anti-BLV ELISA and qPCR were classified as BLV(-)

The quality of extracted total RNA was assessed using the RNA Integrity Number (RIN), which ranged 7.8 to 9.6 across all samples. Small RNA sequencing libraries were then constructed by incorporating adapters with UMIs tags. A multi-step processing pipeline was performed to obtain high-quality, non-redundant reads for unbiased downstream analyses. A descriptive characterization of each sample and the number of sequencing reads before and after quality control is summarized in Table 1.

Demultiplexed reads were mapped to the reference miRNA sequences of bovine, BLV, BFV, BoHV-1 and BoHV-5 (n = 1060) to identify and quantify the miRNAs present in each sample. On average, 351 miRNAs (with mapped reads ≥ 5) derived from both bovine and viral sources were identified and quantified across all samples (Supp. Table S1). A multivariate analysis of sample distances based on the normalized miRNA expression matrix revealed that BLV(-) samples clustered together, forming a distinct group separate from BLV(+) samples. PC1 explains 62% of the variance in the expression matrix, while PC2 accounts for 17% (Figure 1).

In the BLV(+) group, a total of 10 viral-derived miRNAs were consistently expressed, including 7 from BLV: *blv-miR-B1-3p* (avgexp = 27,681.1), *blv-miR-B2-5p* (avgexp = 59,058.0), *blv-miR-B2-3p* (avgexp = 112.5), *blv-miR-B3-5p* (avgexp = 309.6), *blv-miR-B3-3p* (avgexp = 49,312.2), *blv-miR-B4-3p* (avgexp = 65.1), and *blv-miR-B5-5p* (avgexp = 951.7). Additionally, 3 BFV-derived miRNAs were identified: *bfv-miR-BF1-5p* (avgexp = 6,844.7), *bfv-miR-BF1-3p* (avgexp = 8,257.3), and *bfv-miR-BF2-5p* (avgexp =

14,920.6). In contrast, viral-derived miRNAs expression was low (avgexp < 30.0) or undetectable in the

BLV(-) group, except for *blv-miR-B2-5p* (avgexp = 67.4) and *blv-miR-B3-3p* (avgexp = 49.8) (Table 2).

Differential expression analysis of host miRNAs, identified a total of seven bovine miRNAs (bta) with a

fold change (FC) > |1.5| and a q-value < 0.05 were considered differentially expressed (BTA-miRNAs-

DE) between the BLV(+) and BLV(-) groups: *bta-miR-375* (FC = 8.87, q-value = 0.00003), *bta-miR-133a*

(FC = 4.14, q-value = 0.0007), *bta-miR-677* (FC = 8.22, q-value = 0.01), bta-miR-1 (FC = 7.31, q-value =

0.01), *bta-miR-3613a* (FC = 3.83, q-value = 0.01), *bta-miR-9-5p* (FC = 3.83, q-value = 0.04) and *bta-*

miR-95 (FC = 5.09, q-value = 0.04) (Figure 2; Supp. Table S2).

To evaluate the potential functional impact of these seven differentially expressed bta-miRNAs and the

ten viral-derived miRNAs, a target gene prediction study was conducted, identifying a total of 1,911

potentially regulated genes. Of these, 379, 1,046 and 315 were exclusively targeted by BTA-miRNAs-

DE, blv-miRNAs, and bfv-miRNAs, respectively (Supp. Table S3). A low level of overlap was observed

among the potential target genes, with only seven genes shared across all three miRNA groups:

Synaptotagmin (*SYT12*), N-alpha-acetyltransferase (*NAA60*), G-protein-coupled receptor kinase 5

(*GRK5*), Potassium inwardly rectifying channel subfamily J member 9 (*KCNJ9*), Zinc finger and BTB

domain containing 38 (*ZBTB38*), Methyl-CpG binding protein 2 (*MECP2*), and Paternally expressed

gene 10 (*PEG10*). Meanwhile, 69, 20, and 75 genes were shared between BTA-DE – BLV, BTA – BFV,

and BLV – BFV, respectively (Figure 3 and Supp. Table S3).

Functional annotation of the 1,911 potential target genes was conducted using Gene Ontology (GO)

terms and metabolic pathways, revealing 94 significantly overrepresented GO biological process (BP)

terms, 18 GO molecular function (MF) terms and 37 GO cellular component (CC) terms. The word cloud

visualizations for the enriched GO terms in the GO BP and GO MF categories are shown in Figure 4, the

most frequent occurring words for the overrepresented processes and functions included "Regulation,"

"Process," "Response," "Binding," "Activity," and "Kinase," among others. The overrepresentation

analysis of KEGG and Reactome metabolic pathways revealed five significant terms: "*Ras signaling*

pathway" (bta04014, q = 0.04), "*Pathways in cancer*" (bta05200, q = 0.02), "*Oxytocin signaling pathway*"

(bta04921, q = 0.024), "*Morphine addiction*" (bta05032, q = 0.02), and "*Cushing syndrome*" (bta04934, q

= 0.04). Additionally, the terms "*Signal Transduction*" (BTA-162582, q = 0.0002) and "*Immune System*"

(BTA-168256, q = 0.01) were significant in Reactome pathways analysis (Supp. Table S4).

A protein-protein interaction (PPI) network was developed, comprising 1,911 nodes (genes) and 9,198

edges (interactions) (Supp. Figure S1). From this network, five highly interconnected protein clusters

(C1, C2, C3, C4, and C5) were identified (PPI p-value < $1.0e^{-16}$), which potentially represent hubs of

specialized biological function (Figure 5). Functional overrepresentation analysis for these clusters

revealed key cellular processes involved in immune response modulation, cell signaling, infection

persistence, and mechanisms related to cancer development (Supp. Table S4).

For example, Figure 6 illustrates the grouping based on the semantic similarity of significantly

overrepresented GO BP terms for C1 (q < 0.05), whit representative terms including "*Cell surface*

receptor signaling pathway" "*Extrinsic apoptotic signaling pathway in absence of ligand*" "*Regulation of leukocyte proliferation*" "*Anatomical structure morphogenesis*", "*Cell activation*", "*Regulation of cell migration*", "*Cellular developmental process*" and "*Positive regulation of immune system process*". For C2, the enriched GO BP terms were primarily link to immune system processes, including "*Cytokine-mediated signaling pathway*", "*Regulation of T cell activation*", "*Regulation of interferon-gamma production*", "*Positive and negative regulation of cytokine production*" and "*Regulation of leukocyte differentiation*" among others. In C3, mostly associated with transcriptional regulation through chromatin modification, including "*Histone deacetylation*", "*Histone modification*", "*Positive regulation of transcription, DNA-templated*" and "*Chromatin remodeling*". Finally, the enriched biological processes in C4 and C5 were releated to signaling pathways, such as the "*Regulation of Wnt signaling pathway*" and the "*G protein-coupled receptor signaling pathway*" for C4 and C5, respectively.

Significantly enriched KEGG pathways for C1 included: "*Pathways in cancer*" (bta05200, q-value = $3.2e^{-09}$), "*Apoptosis*" (bta04210, q-value = $6.4e^{-06}$), "*MicroRNAs in cancer*" (bta05206, q-value = 0.0006), "*Human T-cell leukemia virus 1 infection*" (bta05166, q-value = 0.0004) and "*Regulation of actin cytoskeleton*" (bta04810, q-value = 0.001). For C2, significant pathways included "*Viral protein interaction with cytokine and cytokine receptor*" (bta04061, q-value = 8.99e-13), "*JAK-STAT signaling pathway*" (bta04630, q-value = 6.58e-09) and "*Chemokine signaling pathway*" (bta04062, q-value = 1.44e-05). Additionally, for C4 and C5, the enriched pathways were "*Wnt signaling pathway*" (bta04310, q-value = 5.64e-21) and "*Apelin signaling pathway*" (bta04371, q-value = 1.58e-10), respectively.

Discussion

This study, conducted a comprehensive characterization of circulating miRNAs in the peripheral blood of naturally BLV-infected and non-infected cattle was performed. Noteworthy, all BLV-infected cows had a high PVL, which has been associated with progression of the BLV pathogenesis (Mirsky et al., 1996; Jimba et al., 2010), and with an increase in viral dissemination within herds (Mammerickx et al., 1987). Following the identification of BLV-miRNAs in persistently infected cell lines (Kinkaid et al., 2012), their expression was assessed using next-generation sequencing for complete miRNA profiling and RT-qPCR to test the expression of candidate miRNAs. BLV-miRNAs were identified in primary leukemic B-cells and B-cell lymphomas isolated from BLV-infected ovine/bovine tumors (Rosewick et al., 2013), experimentally infected cattle (Gillet et al., 2016), and naturally BLV-infected cattle (Frie et al., 2018; Casas et al., 2020; Petersen et al., 2023).

On the other hand, BFV-miRNAs have been identified in both persistently and recently infected MDBK cells, as well as in BFV experimentally challenged cattle (Whisnant et al., 2014). However, to our knowledge, this is the first report of the co-expression of miRNAs derived from two distinct viruses (BLV

and BFV) in naturally infected cattle. We identified and quantified seven BLV-derived and three BFV-derived miRNAs. No miRNAs from BoHV-1 and -5 were detected.

Coinfections of BLV and BFV are prevalent among cattle (Materniak et al., 2019), akin to the occurrences in cats infected with Feline foamy virus (FFV) and Feline leukemia virus (FeLV) (Cavalcante et al., 2018), as well as in baboons with Simian foamy virus (SFV) and Simian T-cell leukemia virus (STLV) (Alais et al., 2018). While BFV infections in cattle are generally regarded as mild and asymptomatic (Whisnant et al., 2014; Materniak-Kornas et al., 2019), there is evidence to suggest that co-infections with BFV and BLV could amplify pathogenicity and impair the bovine immune system, thereby aiding in the transmission and spread of BFV (Amborski et al., 1989; Jacobs et al., 1992; Materniak et al., 2013; 2019). Infectious BFV has been extracted from raw milk, and considering that foamy viruses can cross species barriers, there is rising concern over the zoonotic potential of BFV in humans (Romen et al., 2007; Pinto-Santini et al., 2017). Moreover, increased pathogenic effects have been noted in mixed infections of macaques with SFV and Simian immunodeficiency virus (SIV) (Choudhary et al., 2013), as well as in cats coinfected with Feline immunodeficiency Virus (FIV) and FFV or FeLV and FFV (Cavalcante et al., 2018; Powers et al., 2018).

The seven miRNAs derived from BLV comprised no more than 8% of the total miRNA sequencing reads in all analyzed samples (data not provided), consistent with the expression levels reported by Casas et al. (2020) for natural BLV infections. In contrast, Rosewick et al. (2013) found that all ten mature BLV-derived miRNAs accounted for about 40% of total miRNAs present in B-cell lymphomas from sheep infected with BLV. Similarly, Ochiai et al. (2021) noted that BLV-derived miRNAs made up 38% of total miRNAs in Japanese black cattle diagnosed with enzootic bovine leukosis (B-cell lymphoma). Additionally, Kincaid et al. (2012) reported that BLV miRNAs were found in greater quantities than cellular miRNAs in the BL3.1 cell line persistently infected with BLV. Although the relative expression levels of individual BLV miRNAs differ across studies, *blv-miR-B4-5p* and *blv-miR-B5-3p* appear to be either weakly expressed or undetectable. The dominant BLV-miRNA may change depending on the disease stage or phase of infection analyzed. Overall, these findings suggest that BLV miRNAs are highly expressed in *ex vivo* cell cultures, persistently infected cell lines, primary B-cell tumors, and cattle affected by enzootic bovine leukosis. In cattle with subclinically BLV-infections, their expression levels might vary based on several factors, including the infection route (natural or experimental) and the animal's age (whether calf or adult cow), among other factors (Kincaid et al., 2012; Rosewick et al., 2013; Gillet et al., 2016; Frie et al., 2018; Casas et al., 2020; Ochiai et al., 2021). Conversely, the other BLV transcripts show little to no expression *in vivo*, highlighting the essential function of BLV miRNAs in viral persistence (Lagarias and Radke, 1989; Radke et al., 1992).

Likewise, three BFV-derived miRNAs (*bfv-miR-BF1-5p*, *bfv-miR-BF1-3p*, and *bfv-miR-BF2-5p*) have been identified in MDBK cells that are persistently and recently infected with BFV, showing higher expression levels during persistent infections. Furthermore, two BFV-miRNAs were found in the peripheral blood leukocytes of cattle experimentally infected with BFV, utilizing a qPCR assay (Whisnant

et al., 2014). This research is the first to document the expression of these three BFV-derived miRNAs in
natural setting infections.

A previous study compared BLV seropositive and seronegative cattle but without testing for BLV
genomic material (Casas et al., 2020). The detection of BLV-miRNAs in seronegative cows suggests a
previous exposure to the virus, likely resulting in an inadequate immune response that did not generate a
positive result in the anti-BLV ELISA test. However, a longitudinal analysis of seroconversion was not
performed. This study found that all ten virus-derived miRNAs were consistently and highly expressed in
the BLV (+) group, whereas only a few showed weak expression in the BLV(-) group. Although
undetected infections cannot be entirely excluded, it is improbable that adult animals over two years old,
which tested negative via qPCR and ELISA at two intervals (three months apart), were unrecognized
cases. The potential for BLV- or BFV-miRNAs to function as xeno-miRs and be transmitted from infected
to uninfected animals remains a question for future research.

Notably, the expression profiles of BLV(+) / BFV-miRNAs(+) and BLV(-) samples distinctly differentiate
the two groups of cows groups, as shown in the PCA plot. This indicates a shift in the host miRNA
expression profile between them.

Among the significant differentially expressed bovine miRNAs, *bta-miR-375* was identified, which has
been previously reported to be associated with BLV infection in cattle (Casas et al., 2020; Ochiai et al.,
2021; Ma et al., 2022), and suggested it as an early biomarker for diagnosing enzootic bovine leukosis
(EBL). Moreover, its levels effectively distinguished EBL-affected cattle from asymptomatic cattle with
high sensitivity and specificity (Murakami et al., 2024).

In a similar manner, *bta-miR-133a* was found to be differentially expressed in the serum of BLV-
seropositive cows compared to seronegative cows (Ma et al., 2022). The other significantly differentially
expressed miRNAs (*bta-miR-677*, *bta-miR-95*, *bta-miR-9-5p*, *bta-miR-3613a*, and *bta-miR-1*) have not
been previously reported in relation to BLV and/or BFV infections.

To date, efforts to assign biological functions to BLV- and BFV-derived miRNAs have focused on
predicting potential target genes for specific BLV- and BFV-miRNAs, in addition to selecting candidate
target genes for functional assays (Kinkaid et al., 2012; Rosewick et al., 2013; Gillet et al., 2016; Frie et
al., 2018; Casas et al., 2020; Cao et al., 2020; Petersen et al., 2023). In this study, we evaluated the
functional interaction (protein-protein interaction network) of potential target genes of BLV-miRNAs, while
also considering target genes of BFV-miRNAs and differentially expressed bovine-miRNAs.

It has been demonstrated that BLV fails to induce leukemia/lymphoma in sheeps (oncogenicity
suppressed) challenged with an isogenic BLV provirus lacking the miRNA genomic region. Additionally,
BLV-miRNAs are crucial in promoting efficient viral propagation and disease progression in cattle (Gillet
et al., 2016; Safari et al., 2020).

The analysis of overrepresented metabolic pathways for the complete potential target gene set identified
"*Pathway in cancer*" as one of the significant KEGG pathways. This pathway involves a series of cellular
signaling pathways that activate crucial hallmarks of tumorigenesis, including tissue invasion and

metastasis, evading apoptosis, genomic instability, cell proliferation, genomic damage, and insensitivity to anti-growth signals (Weinberg, 2007). Supporting this, the grouping by semantic similarity of the overrepresented GO BP terms for the most interconnected protein cluster (biological function hub) C1 included terms associated with cellular processes that, when disrupted, could be linked to tumor development events. These terms include "*Regulation of cell migration*," "*Extrinsic apoptotic signaling pathway in the absence of ligand*", "*Chromatin remodeling*", "*Regulation of leukocyte proliferation*", and "*Cell surface receptor signaling*". Interestingly, other overrepresented KEGG terms in C1 included the "*Ras signaling pathway*", "*Rap1 signaling pathway*", "*PI3K-Akt signaling pathway*", "*MAPK signaling pathway*", "*Regulation of actin cytoskeleton*" and Reactome terms such as "*RAF/MAP kinase cascade*" and "*Signaling by Receptor Tyrosine Kinases*". This signaling pathways are amplified and propagated intracellularly by various kinases, ultimately affecting how transcription factors and histone-modifying complexes control downstream gene expression (Suganuma and Workman, 2013). In this context, a crucial signaling cascade most altered in human cancers is the mitogen-activated protein kinase (MAPK) pathway, which includes the RAS–RAF–MAPK kinase (MEK)–extracellular signal-regulated kinase (ERK) pathway (Gimple and Wang, 2019; Bahar et al., 2023). Using the ovine BLV pathogenesis model, sheeps were experimentally challenged with an isogenic BLV-miRNA deletion mutant, and global transcriptome analysis revealed that BLV-miRNAs primarily promote the proliferation of BLV-infected B-cells by inhibiting immune response and cell signaling pathways (Safari et al., 2020). In turn, in a reporter assay, the FBJ murine osteosarcoma viral oncogene homolog (*FOS*) was identified as a direct target of BLV-miRNAs. The *FOS* gene is a component of the activator protein-1 (AP-1) complex, which is involved in the primary response to B-cell receptor signaling and is frequently downregulated in many cancers, including breast carcinomas (Huo and Rothstein, 1995; Gillet et al., 2016).

The functional annotation of the proteins within hub C2 revealed that immune system-related terms, such as "Cytokine-mediated signaling pathway", "*Regulation of T cell activation*" and "*Regulation of interferon-gamma production*" were significant overrepresented, along with the Reactome term "*Cytokine Signaling in immune system*". Additionally, enriched GO BP terms across C3 included "*Histone modification*", "*Positive regulation of cell population proliferation*", and "*Histone deacetylation*" among others; as well as the Reactome terms "*Regulation of TP53 Activity through Acetylation*" and "*Chromatin modifying enzymes*". The transcription factor *TP53*, a tumor suppressor activated by DNA damage, plays a critical role in maintaining genomic integrity (Aubrey et al., 2016). The expression of two transcription factors, B-lymphocyte-induced maturation protein 1 (*BLIMP1*) and B-cell lymphoma 6 (*BCL6*), is negatively correlated with BLV-miRNA expression in BLV(+) cows. These transcription factors play a pivotal role regulating B-cell differentiation, antibody affinity and T-cells immune function. Particularly, an important function of *BCL6* is to enable GC B-cells to proliferate in response to T-cell antigens by specifically repressing gene expression related to DNA damage sensing. This repression allows for the tolerance of DNA breaks induced during immunoglobulin class rearrangement and somatic hypermutation (Nutt et al., 2007; Swaminathan et al., 2014; Frie et al., 2018).

The functional role of BFV-miRNAs remains poorly investigated. Predicted gene targets for *bfv-miR-BF2-5p* were Ankyrin repeat domain-containing protein (*ANKRD17*) and Bax-interacting factor 1 (*BIF1*) (Cao et al., 2020). *BIF1* plays a key role in activating the pro-apoptotic Bax protein in the intrinsic apoptosis pathway as well as in autophagy and autophagosome formation, thus acting as a tumor suppressor (Coppola et al., 2008; Xu et al., 2016). *ANKRD17,* on the other hand, is involved in DNA replication and cell cycle progression. It also interacts with genes that are responsible for sensing viral RNAs and trigger immune responses (Deng et al., 2009; Wang et al., 2012).

Among the genes targeted by BLV-miRNAs, BFV-miRNAs, and BTA-miRNAs-DE, we identified G-protein-coupled-receptor-kinase-5 (*GRK5*), which is a member of the G protein-coupled receptor kinases (GPCRKs). The GRK proteins use ATP to phosphorylate serine and threonine residues in specific receptor locations, initiating their desensitization and down-regulation (Black et al., 2016). Additionally, two genes associated with DNA methylation binding, methyl-CpG binding protein 2 (*MECP2*) and Zinc-finger-and-BTB-domain-containing-38 (*ZBTB38*), act as epigenetic readers of DNA, by binding to methylated DNA and regulating transcription. Cytosine methylation is crucial for silencing exogenous retroviruses, and *MECP2* seems to mediate this transcriptional silencing (Challita and Kohn, 1994; Lorincz et al., 2001; Vuu et al., 2023). Remarkably, another identified target gene is *PEG10*, an oncogene involved in cell proliferation, apoptosis, and metastasis in B-cell lymphocytic leukemia (Xie et al., 2018).

Regarding the differentially expressed BTA-miRNAs, *miR-1*, *miR-133a*, and *miR-375* are recognized as tumor suppressors and are downregulated in various types of cancer, exerting their effects through distinct mechanisms (Yoshino et al., 2011; Hua et al., 2021). For instance, *miR-1* interacts with the proto-oncogene B-cell lymphoma 2 (*BCL-2*), an anti-apoptotic gene. Overexpression of *miR-1* inhibits cell proliferation, migration, and invasion while promoting apoptosis in breast cancer (Peng et al., 2020). Similarly, *miR-133a* modulates biological processes such as proliferation, apoptosis, and autophagy (Hua et al., 2021). On the other hand, *miR-375* is significantly downregulated in several cancers and considered a biomarker of poor prognosis (Mathé et al., 2009; Avissar et al., 2009; Yan et al., 2014; He et al., 2017), but it is overexpressed in breast cancer (de Souza Rocha Simonini et al., 2010). Similarly, *miR-95-3p* and *miR-9-3p* act as prognostic markers, promoting the progression of cervical and breast cancers, respectively (Li et al., 2020; Feng et al., 2022).

Conversely, the differentially expressed bovine miRNA *bta-miR-677* enhances the production of type I interferons (IFN-I) and interferon-stimulated genes (ISGs) (Zhong et al., 2022). Similarly, inhibiting *miR-3613-3p* decreases the expression of IFN-α and IFN-β, thereby affecting the anti-hepatitis B activity of interferons (Zhao et al., 2019). Moreover, *miR-3613-3p* has been recognized as a tumor suppressor, with its deletion associated with poor prognosis in estrogen receptor-positive breast cancer patients (Chen et al., 2021).

Conclusion

In this study, we report for the first time the co-expression of seven miRNAs derived from BLV and three from BFV in cattle naturally infected with BLV. Several bovine BTA-miRNAs were found to be differentially expressed (DE) between cows with BLV high proviral load (PVL) and non-infected cows. Among them, *bta-miR-375* was previously identified as a potential early biomarker for the progression of enzootic bovine leukosis, and this study further emphasizes its potential utility in identifying high PVL cows alongside other candidate miRNAs.

The functional analysis of protein-protein interaction networks involving BLV-, BFV-, and BTA-DE miRNAs-targeted genes identified key metabolic pathways that potentially underlie tumorigenesis and immune response modulation, including cellular signaling pathways, regulation of T cell activation, apoptosis, chromatin remodeling, and cytoskeleton regulation. These findings emphasize the interplay between viral miRNAs and host cellular processes in promoting disease progression. The co-expression of miRNAs derived from BLV and BFV raises concerns about whether their interaction may exacerbate disease pathogenesis or facilitate the dissemination of BFV in light of the zoonotic potential that BFV presents. Ultimately, the identified miRNAs and associated metabolic pathways offer promising opportunities for developing diagnostic tools and therapeutic strategies aimed at controlling viral persistence and tumorigenesis in cattle.

Declarations

Competing interests

The authors declare that they have no competing interests.

Funding

The present study was partially supported by the Instituto Nacional de Tecnología Agropecuaria (INTA, Argentina) projects 2023-2017-PD-I113 and 2023-2027-PD-I114, the Fondo para la Investigación Científica y Tecnológica (FONCyT) PICT-2017-0262, and by the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, Argentina) PIP 2021- 2023-11220200101106CO01

Authors' contributions

Conceptualization: DDG, CM, JPJ, HAC.

Data curation: HAC.

Formal analysis: MIP, HAC.

Funding acquisition: CM, JPJ, HAC

Investigation: MIP, GSA, CP, DDG, CM, JPJ, HAC.

Methodology: HAC.

Visualization: HAC.

Writing – original draft: HAC.

Writing – review & editing: MIP, GSA, CP, DDG, CM, JPJ, HAC.

Roles as defined by: CRediT (contributor role taxonomy)

Availability of data

553   Raw small sequencing reads are available in the NCBI Sequence Read Archive (SRA) under BioProject

554   accession number PRJNA1219522.

Acknowledgments

558   We would like to thank Dr. Andrea Puebla and Pablo Vera from the Genomic Unit (IABIMO, INTA-

559   CONICET) for their technical assistance with the small RNA next-generation sequencing assay.

References

.- Aguado LC, tenOever B. 2018. RNA virus building blocks-miRNAs not included. PLoS Pathog. 14(5):e1006963.
doi: 10.1371/journal.ppat.1006963.

.- Alais S, Pasquier A, Jegado B, Journo C, Rua R, Gessain A, Tobaly-Tapiero J, Lacoste R, Turpin J, Mahieux R.
2018. STLV-1 co-infection is correlated with an increased SFV proviral load in the peripheral blood of SFV/STLV-1
naturally infected non-human primates. PLoS Negl Trop Dis. 12(10):e0006812. doi: 10.1371/journal.pntd.0006812.

.- Aso S, Kitao K, Hashimoto-Gotoh A, Sakaguchi S, Miyazawa T. 2021. Identification of Feline Foamy Virus-
derived MicroRNAs. Microbes Environ. 36(4):ME21055. doi: 10.1264/jsme2.ME21055.

.- Aubrey BJ, Strasser A, Kelly GL. 2016. Tumor-Suppressor Functions of the TP53 Pathway. Cold Spring Harb
Perspect Med. 6(5):a026062. doi: 10.1101/cshperspect.a026062.

.- Avissar M, Christensen BC, Kelsey KT, Marsit CJ. 2009. MicroRNA expression ratio is predictive of head and
neck squamous cell carcinoma. Clin Cancer Res. 15(8):2850-5. doi: 10.1158/1078-0432.CCR-08-3131.

.- Bahar ME, Kim HJ, Kim DR. 2023 Targeting the RAS/RAF/MAPK pathway for cancer therapy: from mechanism
to clinical studies. Signal Transduct Target Ther. 8(1):455. doi: 10.1038/s41392-023-01705-z.

.- Bartlett PC, Norby B, Byrem TM, Parmelee A, Ledergerber JT, Erskine RJ. 2013. Bovine leukemia virus and cow
longevity in Michigan dairy herds. J Dairy Sci. 96:1591–1597.

.- Bartel DP. 2004 MicroRNAs: genomics, biogenesis, mechanism, and function. Cell. 116(2):281-97. doi:
10.1016/s0092-8674(04)00045-5.

.- Bartel DP. 2009. MicroRNAs: target recognition and regulatory functions. Cell. 136(2):215-33. doi:
10.1016/j.cell.2009.01.002.

.- Bartel DP. 2018. Metazoan MicroRNAs. Cell. 173(1):20-51. doi: 10.1016/j.cell.2018.03.006.

.- Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: A practical and powerful approach to
multiple testing. J Roy Statist Soc Ser B. 57:289–300.

.- Black JB, Premont RT, Daaka Y. 2016. Feedback regulation of G protein-coupled receptor signaling by GRKs and arrestins. Semin Cell Dev Biol. 50:95–104. doi: 10.1016/j.semcdb.2015.12.015.

.- Bushnell, B. 2014. BBMap: A Fast, Accurate, Splice-Aware Aligner. Berkeley: Lawrence Berkeley National Lab.(LBNL).

.- Cao W, Heit A, Hotz-Wagenblatt A, Löchelt M. 2018. Functional characterization of the bovine foamy virus miRNA expression cassette and its dumbbell-shaped pri-miRNA. Virus Genes. 54(4):550-560. doi: 10.1007/s11262-018-1574-z.

.- Cao W, Stricker E, Hotz-Wagenblatt A, Heit-Mondrzyk A, Pougialis G, Hugo A, Kuźmak J, Materniak-Kornas M, Löchelt M. 2020. Functional Analyses of Bovine Foamy Virus-Encoded miRNAs Reveal the Importance of a Defined miRNA for Virus Replication and Host-Virus Interaction. Viruses. 12(11):1250. doi: 10.3390/v12111250.

.- Casas E, Ma H, Lippolis JD. 2020. Expression of Viral microRNAs in Serum and White Blood Cells of Cows Exposed to Bovine Leukemia Virus. Front Vet Sci. 7:536390. doi: 10.3389/fvets.2020.536390.

.- Cavalcante LTF, Muniz CP, Jia H, Augusto AM, Troccoli F, Medeiros SO, Dias CGA, Switzer WM, Soares MA, Santos AF. 2018. Clinical and Molecular Features of Feline Foamy Virus and Feline Leukemia Virus Co-Infection in Naturally-Infected Cats. Viruses. 10(12):702. doi: 10.3390/v10120702.

.- Challita PM, Kohn DB. 1994. Lack of expression from a retroviral vector after transduction of murine hematopoietic stem cells is associated with methylation in vivo. Proc Natl Acad Sci U S A. 91(7):2567-71. doi: 10.1073/pnas.91.7.2567.

.- Chen C, Pan Y, Bai L, Chen H, Duan Z, Si Q, Zhu R, Chuang TH, Luo Y. 2021. MicroRNA-3613-3p functions as a tumor suppressor and represents a novel therapeutic target in breast cancer. Breast Cancer Res. 23(1):12. doi: 10.1186/s13058-021-01389-9.

.- Coppola D, Khalil F, Eschrich SA, Boulware D, Yeatman T, Wang HG. 2008. Down-regulation of Bax-interacting factor-1 in colorectal adenocarcinoma. Cancer. 113(10):2665-70. doi: 10.1002/cncr.23892.

.- Cullen BR. 2009. Viral and cellular messenger RNA targets of viral microRNAs. Nature. 22;457(7228):421-5. doi: 10.1038/nature07757.

.- Cullen BR. 2013. MicroRNAs as mediators of viral evasion of the immune system. Nat Immunol. 14(3):205-10. doi: 10.1038/ni.2537.

.- da Silva LF, Jones C. 2012. Two microRNAs encoded within the bovine herpesvirus 1 latency-related gene promote cell survival by interacting with RIG-I and stimulating NF-κB-dependent transcription and beta interferon signaling pathways. J Virol. 86(3):1670-82. doi: 10.1128/JVI.06550-11.

.- De Brun L, Mionetto M, Rodríguez F, et al. 2023. Natural Association between Bovine Leukemia Virus and Reproductive Infectious Diseases. Acta Scientiae Veterinariae. 51:1922.

.- de Souza Rocha Simonini P, Breiling A, Gupta N, Malekpour M, Youns M, Omranipour R, Malekpour F, Volinia S, Croce CM, Najmabadi H, Diederichs S, Sahin O, Mayer D, Lyko F, Hoheisel JD, Riazalhosseini Y. 2010. Epigenetically deregulated microRNA-375 is involved in a positive feedback loop with estrogen receptor alpha in breast cancer cells. Cancer Res. 70(22):9175-84. doi: 10.1158/0008-5472.CAN-10-1318.

.- Deng M, Li F, Ballif BA, Li S, Chen X, Guo L, Ye X. 2009. Identification and functional analysis of a novel cyclin e/cdk2 substrate ankrd17. J Biol Chem. 284(12):7875-88. doi: 10.1074/jbc.M807827200.

.- Friedländer M, Mackowiak S, Li N, Chen W, Rajewsky N. 2012. miRDeep2 identifica con precisión genes de microARN conocidos y cientos de nuevos en siete clados animales, Nucleic Acids Research 40(1):37–52.

.- Enright AJ, John B, Gaul U, Tuschl T, Sander C, Marks DS. 2003. MicroRNA targets in Drosophila. Genome Biol. 5(1):R1. doi: 10.1186/gb-2003-5-1-r1.

.- Erskine RJ, Bartlett PC, Byrem TM, Render CL, Febvay C, Houseman JT. 2012. Using a herd profile to determine age-specific prevalence of bovine leukemia virus in Michigan dairy herds. Vet Med Int.

.- Feng S, Lu Y, Sun L, Hao S, Liu Z, Yang F, Zhang L, Wang T, Jiang L, Zhang J, Liu S, Pang H, Wang Z, Wang H. 2022. MiR-95-3p acts as a prognostic marker and promotes cervical cancer progression by targeting VCAM1. Ann Transl Med. 10(21):1171. doi: 10.21037/atm-22-5184.

.- Frie MC, Droscha CJ, Greenlick AE, Coussens PM. 2018. MicroRNAs Encoded by Bovine Leukemia Virus (BLV) Are Associated with Reduced Expression of B Cell Transcriptional Regulators in Dairy Cattle Naturally Infected with BLV. Front Vet Sci. 4:245. doi: 10.3389/fvets.2017.00245.

.- Friedman RC, Farh KK, Burge CB, Bartel DP. 2009. Most mammalian mRNAs are conserved targets of microRNAs. Genome Res. 19(1):92-105. doi: 10.1101/gr.082701.108.

.- Ghosh Z, Mallick B, Chakrabarti J. 2009. Cellular versus viral microRNAs in host-virus interaction. Nucleic Acids Res. 37(4):1035-48. doi: 10.1093/nar/gkn1004.

.- Gillet N, Florins A, Boxus M, Burteau C, Nigro A, Vandermeers F, Balon H, Bouzar AB, Defoiche J, Burny A,
Reichert M, Kettmann R, Willems L. 2007. Mechanisms of leukemogenesis induced by bovine leukemia virus:
prospects for novel anti-retroviral therapies in human. Retrovirology. 4:18. doi: 10.1186/1742-4690-4-18.

.- Gillet NA, Hamaidia M, de Brogniez A, Gutiérrez G, Renotte N, Reichert M, Trono K, Willems L. 2016. Bovine
Leukemia Virus Small Noncoding RNAs Are Functional Elements That Regulate Replication and Contribute to
Oncogenesis In Vivo. PLoS Pathog. 12(4):e1005588. doi: 10.1371/journal.ppat.1005588.

.- Girardi E, López P, Pfeffer S. 2018 On the Importance of Host MicroRNAs During Viral Infection. Front Genet.
9:439. doi: 10.3389/fgene.2018.00439.

.- Glazov EA, Horwood PF, Assavalapsakul W, Kongsuwan K, Mitchell RW, Mitter N, Mahony TJ. 2010.
Characterization of microRNAs encoded by the bovine herpesvirus 1 genome. J Gen Virol. 91:32-41. doi:
10.1099/vir.0.014290-0.

.- Gimple RC, Wang X. 2019. RAS: Striking at the Core of the Oncogenic Circuitry. Front Oncol. 9:965. doi:
10.3389/fonc.2019.00965.

.- Griffiths-Jones S. 2006 miRBase: the microRNA sequence database. Methods Mol Biol. 342:129-38. doi:
10.1385/1-59745-123-1:129.

.- Grundhoff A, Sullivan CS. 2011. Virus-encoded microRNAs. Virology. 15;411(2):325-43. doi:
10.1016/j.virol.2011.01.002.

.- Guo YE, Steitz JA. 2014. Virus meets host microRNA: the destroyer, the booster, the hijacker. Mol Cell Biol.
34(20):3780-7. doi: 10.1128/MCB.00871-14.

.- Gutiérrez G, Carignano H, Alvarez I, Martínez C, Porta N, Politzki R, Gammella M, Lomonaco M, Fondevila N,
Poli M, Trono K. 2012. Bovine leukemia virus p24 antibodies reflect blood proviral load. BMC Vet Res. 8:187. doi:
10.1186/1746-6148-8-187.

.- Han YC, Park CY, Bhagat G, Zhang J, Wang Y, Fan JB, et al. 2010. microRNA-29a induces aberrant self-
renewal capacity in hematopoietic progenitors, biased myeloid development, and acute myeloid leukemia. J Exp
Med. 207: 475–489. doi: 10.1084/jem.20090831.

.- Harwig A, Das AT, Berkhout B. 2014. Retroviral microRNAs. Curr Opin Virol. 7:47-54. doi:
10.1016/j.coviro.2014.03.013.

.- Hashimoto-Gotoh A, Kitao K, Miyazawa T. 2020. Persistent Infection of Simian Foamy Virus Derived from the
Japanese Macaque Leads to the High-Level Expression of microRNA that Resembles the miR-1 microRNA
Precursor Family. Microbes Environ. 35(1):ME19130. doi: 10.1264/jsme2.ME19130.

.- He L, Hannon GJ. 2004. MicroRNAs: small RNAs with a big role in gene regulation. Nat Rev Genet. 5(7):522-31.
doi: 10.1038/nrg1379.

.- He Y, Jin J, Wang L, Hu Y, Liang D, Yang H, Liu Y, Shan B. 2017. Evaluation of miR-21 and miR-375 as
prognostic biomarkers in oesophageal cancer in high-risk areas in China. Clin Exp Metastasis. 34(1):73-84. doi:
10.1007/s10585-016-9828-4.

.- Hopkins SG, DiGiacomo RF. 1997. Natural transmission of bovine leukemia virus in dairy and beef cattle. Vet
Clin North Am Food Anim Pract. 13(1):107-28. doi: 10.1016/s0749-0720(15)30367-4.

.- Hua YT, Xu WX, Li H, Xia M. 2021. Emerging roles of MiR-133a in human cancers. J Cancer. 12(1):198-206. doi:
10.7150/jca.48769.

.- Huo L, Rothstein TL. 1995. Receptor-specific induction of individual AP-1 components in B lymphocytes. J
Immunol. 154(7):3300-9.

.- Hussain M, Torres S, Schnettler E, Funk A, Grundhoff A, Pijlman GP, Khromykh AA, Asgari S. 2012. West Nile
virus encodes a microRNA-like small RNA in the 3' untranslated region which up-regulates GATA4 mRNA and
facilitates virus replication in mosquito cells. Nucleic Acids Res. 40(5):2210-23. doi: 10.1093/nar/gkr848.

.- Hussain M, Asgari S. 2014 MicroRNA-like viral small RNA from Dengue virus 2 autoregulates its replication in
mosquito cells. Proc Natl Acad Sci US 111(7):2746-51. doi: 10.1073/pnas.1320123111.

.- Jaber T, Workman A, Jones C. 2010. Small noncoding RNAs encoded within the bovine herpesvirus 1 latency-
related gene can reduce steady-state levels of infected cell protein 0 (bICP0). J Virol. 84(13):6297-307. doi:
10.1128/JVI.02639-09.

.- Jimba M, Takeshima S, Matoba K, Endoh D, Aida Y. 2010. BLV-CoCoMo-qPCR: quantitation of bovine leukemia
virus proviral load using the CoCoMo algorithm. Retrovirology. 7:91. doi: 10.1186/1742-4690-7-91.

.- Kanehisa M, Goto S. 2000. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids Res. 28(1):27-30.
doi: 10.1093/nar/28.1.27.

.- Kim VN. 2005. MicroRNA biogenesis: coordinated cropping and dicing. Nat Rev Mol Cell Biol. 6(5):376-85. doi:
10.1038/nrm1644.

.- Kincaid RP, Burke JM, Sullivan CS. 2012. RNA virus microRNA that mimics a B-cell oncomiR. Proc Natl Acad
Sci US 109(8):3077-82. doi: 10.1073/pnas.1116107109.

.- Kincaid RP, Chen Y, Cox JE, Rethwilm A, Sullivan CS. 2014. Noncanonical microRNA (miRNA) biogenesis gives
rise to retroviral mimics of lymphoproliferative and immunosuppressive host miRNAs. mBio 5(2):e00074. doi:
10.1128/mBio.00074-14.

.- Kozomara A, Birgaoanu M, Griffiths-Jones S. 2019. miRBase: from microRNA sequences to function. Nucleic
Acids Res. 47(D1):D155-D162. doi: 10.1093/nar/gky1141.

.- Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods. 9(4):357-9. doi:
10.1038/nmeth.1923.

.- Li X, Zeng Z, Wang J, Wu Y, Chen W, Zheng L, Xi T, Wang A, Lu Y. 2020. MicroRNA-9 and breast cancer.
Biomed Pharmacother. 122:109687. doi: 10.1016/j.biopha.2019.109687.

.- Liang H, Zhou Z, Zhang S, Zen K, Chen X, Zhang C. 2014. Identification of Ebola virus microRNAs and their
putative pathological function. Sci China Life Sci. 57(10):973-81. doi: 10.1007/s11427-014-4759-2.

.- Liu D. 2019. Algorithms for efficiently collapsing reads with Unique Molecular Identifiers. PeerJ. 7:e8275. doi:
10.7717/peerj.8275.

.- Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. Genome Biol. 15(12):550. doi: 10.1186/s13059-014-0550-8.

.- Lorincz MC, Schübeler D, Groudine M. 2001. Methylation-mediated proviral silencing is associated with MeCP2
recruitment and localized histone H3 deacetylation. Mol Cell Biol. 21(23):7913-22. doi: 10.1128/MCB.21.23.7913-
7922.2001.

.- Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal.
17(1):pp10-12. doi:https://doi.org/10.14806/ej.17.1.200.

.- Materniak-Kornas M, Tan J, Heit-Mondrzyk A, Hotz-Wagenblatt A, Löchelt M. 2019. Bovine Foamy Virus: Shared
and Unique Molecular Features In Vitro and In Vivo. Viruses. 11(12):1084. doi: 10.3390/v11121084.

.- Ma H, Lippolis JD, Casas E. 2022. Expression Profiles and Interaction of MicroRNA and Transcripts in Response
to Bovine Leukemia Virus Exposure. Front Vet Sci. 9:887560. doi: 10.3389/fvets.2022.887560.

.- Mammerickx M, Portetelle D, de Clercq K, Burny A. 1987. Experimental transmission of enzootic bovine leukosis
to cattle, sheep and goats: infectious doses of blood and incubation period of the disease. Leuk Res.11:353–358.
doi: 10.1016/0145-2126(87)90180-9.

.- Mathé EA, Nguyen GH, Bowman ED, Zhao Y, Budhu A, Schetter AJ, Braun R, Reimers M, Kumamoto K,
Hughes D, Altorki NK, Casson AG, Liu CG, Wang XW, Yanaihara N, Hagiwara N, Dannenberg AJ, Miyashita M,
Croce CM, Harris CC. 2009. MicroRNA expression in squamous cell carcinoma and adenocarcinoma of the
esophagus: associations with survival. Clin Cancer Res. 15(19):6192-200. doi: 10.1158/1078-0432.CCR-09-1467.

.- Mirsky ML, Olmstead CA, Da Y, Lewin HA. 1996. The prevalence of proviral bovine leukemia virus in peripheral
blood mononuclear cells at two subclinical stages of infection. J Virol. 70:2178–2183. doi: 10.1128/jvi.70.4.2178-
2183.1996.

.- Méndez A, Cerón-Téllez F, Sarmiento R, Tórtora-Pérez J, Rojas-Anaya E, Álvarez H. 2023. Presence of co-
infection between bovine leukemia virus and bovine herpesvirus 1 in herds vaccinated against bovine respiratory
complex. Canadian Journal of Veterinary Research. 87(2):105-109(5).

.- Murakami K, Matsunaga T, Matsuzaki T, Naruke Y, Miyauchi S, Kobayashi S, Yoneyama S, Sakai Y, Ichijo T,
Hirata TI, Kimura A, Chiba Y, Matsuda KI, Yamada S, Hikono H. 2024. Serum bta-miRNA-375 as a potential
biomarker for the early diagnosis of enzootic bovine leukosis. PLoS One. 19(5):e0302868. doi:
10.1371/journal.pone.0302868.

.- NAHMS,2007.https://www.aphis.usda.gov/animal_health/nahms/dairy/downloads/dairy07/Dairy07_is_BLV.pdf.

.- Nutt SL, Fairfax KA, Kallies A. 2007. BLIMP1 guides the fate of effector B and T cells. Nat Rev Immunol.
7(12):923-7. doi: 10.1038/nri2204.

.-Ochiai C, Miyauchi S, Kudo Y, Naruke Y, Yoneyama S, Tomita K, Dongze L, Chiba Y, Hirata TI, Ichijo T, Nagai K,
Kobayashi S, Yamada S, Hikono H, Murakami K. 2021 Characterization of microRNA expression in B cells derived
from Japanese black cattle naturally infected with bovine leukemia virus by deep sequencing. PLoS One.
16(9):e0256588. doi: 10.1371/journal.pone.0256588.

.- OIE. Manual of Diagnostic Tests and Vaccines for Terrestrial Animals. 2012. 7th Ed.

.- Oviedo-Pastrana M, Doria-Ramos M, Mattar S, Oviedo-Socarras T, Vallejo-Timarán D. 2024. Seroprevalence of
bovine leukemia virus and association with bovine infectious abortion in Creole breeds from tropical grazing herds
in the Colombian Caribbean. Vet World. 17(8):1715-1721. doi: 10.14202/vetworld.2024.1715-1721.

.- Peng J, Yuan C, Wu Z, Wang Y, Yin W, Lin Y, Zhou L, Lu J. 2020. Upregulation of microRNA-1 inhibits proliferation and metastasis of breast cancer. Mol Med Rep. 22(1):454-464. doi: 10.3892/mmr.2020.11111.

.- Petersen MI, Alvarez I, Trono KG, Jaworski JP. 2018. Quantification of bovine leukemia virus proviral DNA using a low-cost real-time polymerase chain reaction. J Dairy Sci. 101(7):6366-6374. doi: 10.3168/jds.2017-14253.

.- Petersen MI, Carignano HA, Suarez Archilla G, Caffaro ME, Alvarez I, Miretti MM, Trono K. 2021. Expression-based analysis of genes related to single nucleotide polymorphism hits associated with bovine leukemia virus proviral load in Argentinean dairy cattle. J Dairy Sci. 104(2):1993-2007. doi: 10.3168/jds.2020-18924.

.- Petersen MI, Carignano HA, Mongini C, Gonzalez DD, Jaworski JP. 2023. Bovine leukemia virus encoded blv-miR-b4-3p microRNA is associated with reduced expression of anti-oncogenic gene in vivo. PLoS One 18(2):e0281317. doi: 10.1371/journal.pone.0281317.

.- Pfeffer S, Zavolan M, Grässer FA, Chien M, Russo JJ, Ju J, John B, Enright AJ, Marks D, Sander C, Tuschl T. 2004. Identification of virus-encoded microRNAs. Science. 304(5671):734-6. doi: 10.1126/science.1096781.

.- Pfeffer S, Sewer A, Lagos-Quintana M, Sheridan R, Sander C, Grässer FA, van Dyk LF, Ho CK, Shuman S, Chien M, Russo JJ, Ju J, Randall G, Lindenbach BD, Rice CM, Simon V, Ho DD, Zavolan M, Tuschl T. 2005. Identification of microRNAs of the herpesvirus family. Nat Methods. 2(4):269-76. doi: 10.1038/nmeth746.

.- Pluta A, Jaworski JP, Douville RN. 2020. Regulation of Expression and Latency in BLV and HTLV. Viruses. 12(10):1079. doi: 10.3390/v12101079.

.- R Core Team, 2024. R: A Language and Environment for Statistical Computing. Vienna, Austria.

.- Rosen BD, Bickhart DM, Schnabel RD, Koren S, Elsik CG, Tseng E, Rowan TN, Low WY, Zimin A, Couldrey C, Hall R, Li W, Rhie A, Ghurye J, McKay SD, Thibaud-Nissen F, Hoffman J, Murdoch BM, Snelling WM, McDaneld TG, Hammond JA, Schwartz JC, Nandolo W, Hagen DE, Dreischer C, Schultheiss SJ, Schroeder SG, Phillippy AM, Cole JB, Van Tassell CP, Liu G, Smith TPL, Medrano JF. 2020. De novo assembly of the cattle reference genome with single-molecule sequencing. Gigascience. 1;9(3):giaa021. doi: 10.1093/gigascience/giaa021.

.- Rosewick N, Momont M, Durkin K, Takeda H, Caiment F, Cleuter Y, Vernin C, Mortreux F, Wattel E, Burny A, Georges M, Van den Broeke A. 2013. Deep sequencing reveals abundant noncanonical retroviral microRNAs in B-cell leukemia/lymphoma. Proc Natl Acad Sci U S A. 110(6):2306-11. doi: 10.1073/pnas.1213842110.

.- Ruivinho C, Gama-Carvalho M. 2023. Small non-coding RNAs encoded by RNA viruses: old controversies and new lessons from the COVID-19 pandemic. Front Genet. 21;14:1216890.

.- Safari R, Jacques JR, Brostaux Y, Willems L. 2020. Ablation of non-coding RNAs affects bovine leukemia virus B lymphocyte proliferation and abrogates oncogenesis. PLoS Pathog. 16(5):e1008502. doi: 10.1371/journal.ppat.1008502.

.- Samols MA, Hu J, Skalsky RL, Renne R. 2005 Cloning and identification of a microRNA cluster within the latency-associated region of Kaposi's sarcoma-associated herpesvirus. J Virol. 79(14):9301-5. doi: 10.1128/JVI.79.14.9301-9305.2005.

.- Santanam U, Zanesi N, Efanov A, Costinean S, Palamarchuk A, Hagan JP, et al. 2010 Chronic lymphocytic leukemia modeled in mouse by targeted miR-29 expression. Proc Natl Acad Sci U S A. 107: 12210–12215. doi: 10.1073/pnas.1007186107.

.- Sayols S. 2023. rrvgo: a Bioconductor package for interpreting lists of Gene Ontology terms. MicroPubl Biol. 10.17912/micropub.biology.000811. doi: 10.17912/micropub.biology.000811.

.- Seo GJ, Fink LH, O'Hara B, Atwood WJ, Sullivan CS. 2008. Evolutionarily conserved function of a viral microRNA. J Virol. 82(20):9823-8. doi: 10.1128/JVI.01144-08.

.- Smith T, Heger A, Sudbery I. 2017. UMI-tools: modeling sequencing errors in Unique Molecular Identifiers to improve quantification accuracy. Genome Res. 27(3):491-499. doi: 10.1101/gr.209601.116.

.- Skalsky RL, Cullen BR. 2010. Viruses, microRNAs, and host interactions. Annu Rev Microbiol. 64:123-41. doi: 10.1146/annurev.micro.112408.134243.

.- Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, Doncheva NT, Legeay M, Fang T, Bork P, Jensen LJ, von Mering C. 2021. The STRING database in 2021: customizable protein-protein networks, and functional characterization of user-uploaded gene/measurement sets. Nucleic Acids Res. 49(D1):D605-D612. doi: 10.1093/nar/gkaa1074. Erratum in: Nucleic Acids Res. 2021 Oct 11;49(18):10800. doi: 10.1093/nar/gkab835.

.- Suganuma T, Workman JL. 2013. Chromatin and signaling. Curr Opin Cell Biol. 25(3):322-6. doi: 10.1016/j.ceb.2013.02.016.

.- Sullivan CS, Ganem D. 2005 MicroRNAs and viral infection. Mol Cell. 7;20(1):3-7. doi: 10.1016/j.molcel.2005.09.012.

.- Tang Q, Wu YQ, Chen DS, Zhou Q, Chen HC, Liu ZF. 2014. Bovine herpesvirus 5 encodes a unique pattern of microRNAs compared with bovine herpesvirus 1. J Gen Virol. 95:671-678. doi: 10.1099/vir.0.061093-0.

.- Swaminathan S, Duy C, Müschen M. 2014. BACH2-BCL6 balance regulates selection at the pre-B cell receptor checkpoint. Trends Immunol. 35(3):131-7. doi: 10.1016/j.it.2013.11.002.

.- Tavazoie S, Hughes JD, Campbell MJ, Cho RJ, Church GM. 1999. Systematic determination of genetic network architecture. Nat Genet. 22(3):281-5.

.- Teng Y, Wang Y, Zhang X, Liu W, Fan H, Yao H, Lin B, Zhu P, Yuan W, Tong Y, Cao W. 2015. Systematic Genome-wide Screening and Prediction of microRNAs in EBOV During the 2014 Ebolavirus Outbreak. Sci Rep. 5:9912. doi: 10.1038/srep09912.

.- Trono KG, Pérez-Filgueira DM, Duffy S, Borca MV, Carrillo C. 2001. Seroprevalence of bovine leukemia virus in dairy cattle in Argentina: comparison of sensitivity and specificity of different detection methods. Vet Microbiol. 83(3):235-48. doi: 10.1016/s0378-1135(01)00420.

.- Tycowski KT, Guo YE, Lee N, Moss WN, Vallery TK, Xie M, Steitz JA. 2015. Viral noncoding RNAs: more surprises. Genes Dev. 29(6):567-84. doi: 10.1101/gad.259077.115.

.- VanLeeuwen JA, Forsythe L, Tiwari A, Chartier R. 2005. Seroprevalence of antibodies against bovine leukemia virus, bovine viral diarrhea virus, Mycobacterium avium subspecies paratuberculosis, and Neospora caninum in dairy cattle in Saskatchewan. Can Vet J. 46:56–58.

.- VanLeeuwen JA, Tiwari A, Plaizier JC, Whiting TL. Seroprevalences of antibodies against bovine leukemia virus, bovine viral diarrhea virus, Mycobacterium avium subspecies paratuberculosis, and Neospora caninum in beef and dairy cattle in Manitoba. Can Vet J. 47:783–786.

.- Vuu YM, Roberts CT, Rastegar M. 2023. MeCP2 Is an Epigenetic Factor That Links DNA Methylation with Brain Metabolism. Int J Mol Sci. 24(4):4218. doi: 10.3390/ijms24044218.

.- Wang Y, Tong X, Li G, Li J, Deng M, Ye X. 2012. Ankrd17 positively regulates RIG-I-like receptor (RLR)-mediated immune signaling. Eur J Immunol. 42(5):1304-15. doi: 10.1002/eji.201142125.

.- Weinberg, R.A. 2007. The Biology of Cancer Garland Science. Taylor & Francis Group, LLC, New York, NY.

.- Whisnant AW, Kehl T, Bao Q, Materniak M, Kuzmak J, Löchelt M, Cullen BR. 2014. Identification of novel, highly expressed retroviral microRNAs in cells infected by bovine foamy virus. J Virol. 88(9):4679-86. doi: 10.1128/JVI.03587-13.

.- Xie T, Pan S, Zheng H, Luo Z, Tembo KM, Jamal M, Yu Z, Yu Y, Xia J, Yin Q, Wang M, Yuan W, Zhang Q, Xiong J. 2018. PEG10 as an oncogene: expression regulatory mechanisms and role in tumor progression. Cancer Cell Int. 18:112. doi: 10.1186/s12935-018-0610-3.

.- Xu L, Wang Z, He SY, Zhang SF, Luo HJ, Zhou K, Li XF, Qiu SP, Cao KY. 2016. Bax-interacting factor-1 inhibits cell proliferation and promotes apoptosis in prostate cancer cells. Oncol Rep. 36(6):3513-3521. doi: 10.3892/or.2016.5172.

.- Yan J, She Q, Shen X, Zhang Y, Liu B, Zhang G. 2017. Potential Role of MicroRNA-375 as Biomarker in Human Cancers Detection: A Meta-Analysis. Biomed Res Int. 2017:1875843. doi: 10.1155/2017/1875843.

.- Yao Y, Smith LP, Nair V, Watson M. 2014. An avian retrovirus uses canonical expression and processing mechanisms to generate viral microRNA. J Virol. 88(1):2-9. doi: 10.1128/JVI.02921-13.

.- Yoshino H, Chiyomaru T, Enokida H, Kawakami K, Tatarano S, Nishiyama K, Nohata N, Seki N, Nakagawa M. 2011. The tumour-suppressive function of miR-1 and miR-133a targeting TAGLN2 in bladder cancer. Br J Cancer. 104(5):808-18. doi: 10.1038/bjc.2011.23.

.- Zhao Y, Yu Y, Ye L. 2019. MiR-3613-3p impairs IFN-induced immune response by targeting CMPK1 in chronic hepatitis B. Infect Genet Evol. 74:103919. doi: 10.1016/j.meegid.2019.103919.

.- Zhong C, Liao Z, Zhang B, Xiao L, Li J, Zhu X. 2022. Bta-miR-677 contribute to interferon pathway affecting the proliferation of caprine parainfluenza virus type 3. Microb Pathog. 169:105642. doi: 10.1016/j.micpath.2022.1056.

Table 1. Sample characterization and small RNA sequencing output

ID	PR (SD)[1]	PVL	RIN	Raw reads (n)	Reads after QC (n)
5671	178.6(28.4)	63,148.0	9.6	8,090,674	4,243,257
5841	191.3(32.8)	44,234.3	9.6	11,869,092	4,038,887
6021	169.8(42.8)	52,902.4	7.8	15,487,902	6,148,370
6097	183.2(25.4)	91,255.4	9.3	12,511,860	4,744,543
5830	<25	-	9.0	11,903,095	5,115,654
6493	<25	-	9.3	10,871,552	4,312,484
6962	<25	-	9.0	11,545,292	4,950,884

[1] Anti-BLV enzyme-linked immunosorbent assay % of reactivity; % of reactivity values < 25.0 indicated negative samples

Abbreviatures: ID = sample identification; PR = % of reactivity; SD = standard deviation; RIN = RNA integrity number; PVL = T-3 and T0 BLV average proviral load; QC = quality control.

Figure 1. Principal component analysis of miRNAs expression matrix. Blue triangles and Red circles represent BLV(+) and BLV(-) samples, respectively. PC = principal component. PC1 and PC2, represent 62% and 17% of variance contained in the expression matrix data, respectively.

Table 2. Viral derived-miRNAs expression

Condition	BLV(+)					BLV(-)			
miRNA/ID	5671	5841	6021	6097	avgexp(SD)	5830	6493	6962	avgexp(SD)
bfv-miR-BF1-5p	8,973.2	719.9	14,813.3	2,872.6	6,844.7(6,359.3)	15.6	5.8	15.4	12.3(5.6)
bfv-miR-BF1-3p	12,947.2	1,329.6	15,826.4	2,925.9	8,257.3(7,204.2)	7.1	11.6	8.6	9.1(2.3)
bfv-miR-BF2-5p	17,694.3	1,735.7	33,383.7	6,868.5	14,920.6(13,991.2)	0.0	15.4	17.1	10.8(9.4)
bfv-miR-BF2-3p	2.8	0.0	6.3	6.2	3.8(3.0)	0.0	0.0	0.0	0.0(0.0)
blv-miR-B1-5p	0.0	0.0	0.0	0.0	0(0.0)	0.0	0.0	0.0	0.0(0.0)
blv-miR-B1-3p	7,556.4	13,726.1	46,243.5	43,198.3	27,681.1(19,875.4)	28.3	24.1	24.0	25.5(2.5)
blv-miR-B2-5p	30,941.9	35,856.1	88,113.4	81,320.4	59,057.9(29,825.4)	60.9	69.4	71.9	67.4(5.7)
blv-miR-B2-3p	34.2	47.4	171.3	197.0	112.5(83.6)	0.0	0.0	0.0	0.0(0.0)
blv-miR-B3-5p	262.0	371.8	259.8	344.7	309.6(57.3)	0.0	0.0	0.0	0.0(0.0)
blv-miR-B3-3p	37,074.5	33,573.4	61,540.1	65,060.6	49,312.2(16,278.9)	39.7	55.9	53.9	49.8(8.8)
blv-miR-B4-5p	1.4	0.0	0.0	1.0	0.6(0.7)	0.0	0.0	0.0	0.0(0.0)
blv-miR-B4-3p	39.9	69.9	79.8	70.8	65.1(17.4)	0.0	0.0	0.0	0.0(0.0)
blv-miR-B5-5p	864.3	972.1	1,218.4	752.0	951.7(199.2)	0.7	1.0	0.9	0.8(0.1)
blv-miR-B5-3p	0.0	1.2	0.8	2.1	1.0(0.9)	0.0	0.0	0.0	0.0(0.0)

Abbreviatures: ID = Sample ID; avgexp = normalized mean expression; SD = Standard deviation

Figure 2. Volcano plot. Differential expression analysis of bovine miRNAs between BLV(+) and BLV(-) groups. Each circle represents a bovine miRNA. Red circles indicate miRNAs with FC < |1.5| and non-significant, while blue circles represent miRNAs with FC > |1.5| and q < 0.05 (-log(q-value) > 1.3). Viral-derived miRNAs, primarily quantified in BLV(+) samples, were excluded from the plot.

Figure 3. Venn diagram of number of predicted gene target for bta-miRNAs-DE, blv-miRNAs and bfv-miRNAs.

A) GO Biological process   B) GO Molecular function

Figure 4. Word cloud for GO term overrepresented in target genes for biological process A) and molecular fuction B) categories.

Figure 5. Protein-protein interaction (PPI) cluster analysis. Five cluster (> 15 nodes) were identified A) Cluster 1 (C1): Nº nodes: 53 - Nº edges: 282, avg. node degree: 10.6; B) Cluster 2 (C2): Nº nodes: 30 – Nº edges: 130, avg. node degree: 8.67; C) Cluster 3 (C3): Nº nodes: 19 – Nº edges: 45, avg. node degree: 4.74; D) Nº nodes: 16 – Nº edges: 42, avg. node degree: 5.25; E) Nº nodes: 16 – Nº edges: 53, avg node degree: 6.62. Nodes represent proteins encoded by a predicted miRNA gene target. Edges represent protein-protein interactions as determined by STRINGdb evidence, including: known interactions (curated databases and/or experimentally validated), predicted interactions (gene fusions, gene co-occurrence), text mining, co-expression, and protein homology.

Figure 6. Treemaps plot for GO BP enriched in C1 cluster. The terms were grouped (colored) based on semantic similarity, where the term with the lowest q-value is considered representative of the group.