A Critical Role for the Fascin Family of Actin Bundling Proteins in Axon Development, Brain Wiring and Function

[1] Katherine R. Hardin, [1]Arjolyn B. Penas, [1]Shuristeen Joubert, [1] Changtian Ye, [1]Kenneth R. Myers, and [1,2,3] James Q. Zheng

1. Department of Cell Biology, Emory University School of Medicine, Atlanta, GA 30322

2. Center for Neurodegenerative Diseases, Emory University School of Medicine, Atlanta, GA 30322

3. Department of Neurology, Emory University School of Medicine, Atlanta, GA 30322

Keywords: Actin cytoskeleton, growth cones, brain wiring, *Drosophila*, mushroom body, sensorimotor function

Abstract: 200, *Introduction:* 545, *Discussion:* 1415, *Figures:* 9, *Tables:*1

Correspondence: James Zheng, PhD, Department of Cell Biology, 615 Michael Street, Atlanta, Georgia 30322, USA. Email: james.zheng@emory.edu

Acknowledgements: We thank Drs. Kenneth Moberg and Dorothy Lerit for their help and guidance with the study. We sincerely thank Hengrui Zhang for his help with primary hippocampal neuronal culture. This research is supported in part by a NRSA F31 fellowship from NIH-NINDS to KRH (NS127574), a R21 research grant from NIH-NINDS to JQZ (NS132393), and a Whitehall Foundation research grant to KRM.

Author contributions: KRH led the effort in conducting most of the experiments and data analysis. ABP performed *Drosophila* brain dissections, immunostaining, imaging, and data analysis. SJ performed RNAi western blots and CAD cell imaging. CY helped with brain dissection and imaging. KRM helped with the experiments concerning cultured mammalian neurons and fascin manipulation. JQZ helped conceptualize the experiments, provided guidance and constructive input to the study. KRH wrote the manuscript draft and worked with JQZ and KRM to finalize the manuscript.

Abstract

Actin-based cell motility drives many neurodevelopmental events including guided axonal growth. Fascin is a major family of F-actin bundling proteins, but its role in axon development and brain wiring is unknown. Here, we report that fascin is required for axon development, brain wiring and function. We show that fascin is enriched in the motile filopodia of axonal growth cones and its inhibition impairs axonal extension and branching of hippocampal neurons in culture. We next provide evidence that fascin is essential for axon development and brain wiring using *Drosophila melanogaster* as an *in vivo* model. *Drosophila* express a single ortholog of mammalian fascin called Singed (SN), which is highly expressed in the mushroom body (MB) of the central nervous system. We observe that loss of SN results in drastic MB disruption, highlighted by α- and β-lobe defects that are consistent with altered axonal guidance. SN-null flies also exhibit defective sensorimotor behaviors as assessed by the negative geotaxis assay. MB-specific expression of SN in SN-null flies rescues MB structure and sensorimotor deficits, indicating that SN functions autonomously in MB neurons. Together, our data from primary neuronal culture and *in vivo* models highlight a critical role for fascin in brain development and function.

Fascin in neuronal development

Significance statement:

- Fascin is a major family of actin-binding proteins (ABPs) that crosslink and bundle actin filaments to underline membrane protrusions essential for cell motility and neuronal development.

- Inhibition of fascin in cultured cells causes it to dissipate from F-actin-rich membrane protrusions and become cytosolic, reduces filopodia formation, and decreases axon outgrowth and branching.

- In fruit flies, loss of the fascin orthologue, Singed, causes defects in brain development, specifically in mushroom body (MB) lobes, and leads to behavioral impairments which can be rescued upon MB-specific expression of Singed.

Introduction

The actin cytoskeleton powers a wide range of cellular activities ranging from cell migration to membrane trafficking[1,2]. The structure and dynamics of the actin cytoskeleton are regulated by over 100 actin binding proteins (ABPs)[3]. Several families of ABPs crosslink actin filaments (F-actin) to generate F-actin bundles of different densities that underlie distinct membrane protrusions such as microvilli and filopodia[4,5]. Filopodia are finger-like protrusions found on motile cells that play an important role in cell migration and cancer cell metastasis[6]. In developing neurons, they serve as a hallmark feature of axonal growth cones, contributing significantly to axon guidance and branching[7-9]. These dynamic structures detect and respond to external axon guidance signals, stabilizing when encountering an attractive molecule or retracting in response to a repellent one[10]. Research has demonstrated that while eliminating growth cone filopodia does not hinder axonal elongation, it disrupts the proper response to guidance cues[11,12]. Additionally, filopodia are essential for collateral branching, a process vital for both neurodevelopment and injury recovery[9]. While the importance of filopodia in axon guidance and branching is recognized, the molecular mechanisms that underlie axon filopodia regulation and function remain to be fully elucidated.

While several families of ABPs can crosslink actin filaments to form different F-actin bundles, fascin is known to generate tight parallel F-actin bundles constituting the core F-actin structure in motile filopodia of nerve growth cones and motile cells[13,14]. There are three mammalian isoforms of fascin, with fascin1 being the most widely expressed and studied[15]. Increased fascin1 expression in cancer cells is associated with increased metastasis and poor clinical prognosis, while knockdown of fascin1 leads to a decrease

in cell migration[16-19]. Interestingly, fascin1 exhibits an intriguing expression profile in mammalian brains: it is highly expressed during development, but its expression is substantially decreased in the adult brain[20,21], suggesting an important role for fascin in early neurodevelopment.

Despite its high level of expression in the developing brain and its importance in the formation of filopodia, a role for fascin1 in neural development and function, especially guided axonal development, has not been investigated. An early study of gross brain morphology in a fascin1 knockout (KO) mouse model revealed an increase in the size of the lateral ventricles and the loss of the posterior region of the anterior commissure, suggesting a potential role for fascin1 in axon development[22]. In addition, KO neurons exhibited smaller growth cones with fewer and shorter filopodia than wild-type control neurons [22]. These findings, along with the high perinatal lethality exhibited by fascin1 KO mice (~50%), suggest a crucial role in mouse development[22]. However, no detailed analysis of brain structure, neuronal connections, synaptic function, or behavior in fascin1 KO mice was performed[22]. In support of these findings, a separate fascin1 knockout mouse developed by the University of California, Davis Knock Out Mouse Project (KOMP) showed homozygous fascin1 KO animals exhibited a high degree of lethality[23]. Interestingly, the heterozygous mice were reported to exhibit neurobehavioral abnormality, further supporting a role for fascin in brain development and function[23].

In this study, we utilized a combination of mammalian cell culture and *Drosophila melanogaster* as an *in vivo* model to investigate the role of fascin in brain development and function. Our findings establish a critically important role for the fascin family of actin bundling proteins in brain development and function.

Methods and Materials

<u>Cell culture</u>

Dissociated primary hippocampal neuron cultures were prepared as previously described[24]. Briefly, 18.5-day old embryos were collected from timed-pregnant Sprague Dawley rats (Charles River Laboratories), and then hippocampi were dissected in ice-cold Hank's Balanced Salt Solution (HBSS). Hippocampi were pooled together, trypsinized for 12 minutes (mins.), briefly incubated in 20% fetal bovine serum (Atlanta Biologicals, #S11150) and then plated on 25 mm coverslips coated with 100 µg/ml poly-L-lysine (P2636; Sigma-Aldrich) at two different densities, ~150,000 or 25,000 cells per 35 mm dish for two different treatment conditions (see below). Hippocampal neurons were cultured in Neurobasal medium (21103049; Gibco) supplemented with 1% B-27 (17504044; Gibco), 1% Penicillin/Streptomycin (30002CI; ThermoFisher), and 1% GlutaMax (35050061; Gibco). The culture medium was replaced with fresh medium 24 hours (hrs.) after plating. Animal care and use was conducted following National Institutes of Health guidelines, and procedures were approved by the Institutional Animal Care and Use Committee at Emory University.

Cath.-a-differentiated (CAD) cells[25] were used for analyses of fascin localization in lamellipodia and filopodia. CAD cells were cultured in DMEM/F12 (10-092-CV; Corning) media supplemented with 8% fetal bovine serum (Atlanta Biologicals, #S11150) and 1% Penicillin/Streptomycin (Invitrogen, #15140122). For imaging, we resuspended CAD cells by mechanical force and plated them on laminin-coated No. 1.5 glass coverslips for one hr. (laminin: CAS#: 114956-81-9, Millipore Sigma; 20 µg/mL for 1 hr. at 37 °C) before fixation and immunocytochemistry.

Fascin in neuronal development

Immunohistochemistry of cultured cells

For immunofluorescent labeling of fascin and its cellular localization, both CAD cells and hippocampal neurons were fixed by cold (-20°C) anhydrate methanol (Sigma CAS# 67-56-1) for 10 min, followed by rehydration in 1x phosphate-buffered saline (PBS). Cells were blocked in PBS containing 2% normal Donkey serum (D9663, Millipore-Sigma) for one hr. at room temperature. The cells were then incubated with primary antibodies at 4°C overnight and labeled with fluorescent secondary antibodies at room temperature for 1 hr. The following antibodies were used: fascin (1:1000 mouse anti-fascin, Sigma MAB3582, clone 55K2) and rabbit anti-β-actin (1:400, ab8227, Abcam). Alexa Fluor-conjugated secondary antibodies (Invitrogen) were diluted at a concentration of 1:500 in PBS-containing 2% donkey serum. All labeled coverslips were mounted on glass slides using Fluoromount-G (Southern Biotech #0100-01) for imaging.

NP-G2-044 treatment

The fascin inhibitor NP-G2-044 (HY-125506, MedChemExpress, Monmouth Junction, NJ) was stored at a concentration of 10 mM in DMSO. CAD cells were plated on 20 µg/ml laminin-coated 25 mm coverslips and allowed to attach and spread for one hr. CAD cells were then treated with either 15 µM of NP-G2-044 or 0.15% DMSO as a control. Coverslips were fixed at either 30 mins or 240 mins post-treatment application and subsequently labeled for fascin and β-actin. Images were acquired using widefield fluorescence microscopy. For this study, we selected fully flattened CAD cells with a "pancake" morphology to keep analysis consistent. For analysis, the segmented line tool in FIJI was used measure the average intensity of both the actin and fascin fluorescence in filopodia and the polygon tool was used to measure the cytosolic intensity of actin and

fascin. The ratios of filopodial to cytosolic actin and fascin fluorescence intensities were calculated and compared using One-way ANOVA with multiple comparisons.

To examine growth cone morphology, cultured hippocampal neurons (150,000/coverslip) were grown to days in vitro (DIV) 3. The cells were then treated with 15 μM NP-G2-004 or 0.15% DMSO for 24 hrs., followed by methanol fixation and immunolabeling for fascin and β-actin on DIV4. Fluorescence microscopy and quantitative analyses were performed using the line tool in FIJI to trace the filopodia in each growth cone to determine length and number.

To investigate neurite extension, we cultured hippocampal neurons at a low density (~25,000 cells/coverslip). On DIV1 the coverslips were inverted over a feeder layer of high-density rat cortical neurons (300,000/well) in 6 well plates with paraffin wax dots to separate the coverslips from the feeder layer. Starting DIV1, the neurons were treated with either 15 μM NP-G2-044, or DMSO vehicle control for 4 days, with fresh DMSO or NP-G2-044 added every day until fixation. On DIV5, the coverslips were removed, washed with PBS, fixed in 4% paraformaldehyde in PBS with 4% sucrose for 15 mins at room temperature, and then permeabilized in PBS containing 0.1% Triton X-100 for ten mins. The cells were blocked for one hr. at room temperature in 2% donkey serum (D9663, Millipore-Sigma) in PBS, and then incubated overnight at 4°C with mouse anti-tau (Sigma MAB3420, clone PC1C6) and rabbit anti-tubulin (Covance, PRB-435P) antibodies at 1:1000 in blocking buffer. Coverslips were next incubated in Alexa Fluor-conjugated secondary antibodies (Invitrogen A10037 and A11036) diluted in PBS 1:500 for an hr. at room temperature. Coverslips were then mounted in Fluoromount-G containing Dapi (Southern Biotech, Cat. No.: 0100-20). Axon length and the number of

axon branch points were measured using the Simple Neurite Tracer (SNT) plugin for FIJI[26].

Unless otherwise noted, all data were collected from three independent biological replicates. Images of mammalian cells were taken using a NIKON Ti2-E inverted microscope equipped with a Hamamatsu FLASH4 eCOMS camera, and a Nikon C2+ confocal system. Slides were blinded for imaging and quantitative analyses. Statistical analysis was performed using GraphPad Prism Version 10.4.1 for macOS.

Fly stocks

Fly stocks were obtained from Bloomington Drosophila Stock Center or were generously provided by Dr. Tina Tootle (University of Iowa), Dr, Kenneth Moberg (Emory University), and Dr. Jennifer Zanet (University of Toulouse, France). All crosses and stocks were maintained in standard conditions.

Name	BL # (if applicable)	Source
w[1118]	3605	Bloomington
Oregon R	2376	Bloomington
sn[28]/fm7Kr-Gal4 UAS GFP		Tina Tootle
sn[x2] (In(1)Mud,Mudd[1]/In(1)dl-49 sn[X2]	1035	Bloomington
w*; if/CyO;TM2/TM6B		Ken Moberg
UAS-snRNAI (III) y[1]sc[*]v[1]sev[21];P{y[+t7.7]v[+t1.8]=TRiP.HMS 02450}dattP2	42615	Bloomington

Fascin in neuronal development

UAS-Dicer2		Ken Moberg
OK107 GAL4 w[*];P{w[+mW.hs]=GawB}OK107ey[OK107]/In(4) ci[D,ci[D] pan[ciD] sv[spa-pol]	854	Ken Moberg
UAS myristoylated GFP		Ken Moberg
Split Mb GAL 4 w[1118];P{y[+t7.7]w[+mC]=R13F02-p65.AD}attP40; P{y[+t7.7] w[+mC]=R58F02-GAL4.DBD}attP2	68255	Bloomington
UAS-GFP fascin (Singed)		Jennifer Zanet
238Y-GAL4 w[1118]; P{w[+mW.hs]=GawB}238Y	81009	Bloomington
Nsyb-GAL4		Ken Moberg
Act5-GAL4/CyO		Ken Moberg

A balanced sn^{28} line was created by crossing sn28/fm7Kr-Gal4 UAS GFP flies with w*; if/CyO;TM2/TM6B flies. The resulting line was: sn28/fm7Kr-Gal4 UAS GFP; if/CyO;TM2/TM6B. This line was used to generate the Split-GAL4 and SN-rescue lines that are in the sn^{28} background.

The following genotypes were made through crosses and used experimentally:

1. Nsyb-GAL4; UAS myristoylated GFP

2. 238Y-GAL4; UAS myristoylated GFP

3. Split MB-GAL 4/ UAS myristoylated GFP; Split MB-GAL 4

4. sn^{28}; Split MB-GAL 4/ UAS myristoylated GFP; Split MB-GAL 4

5. UAS-Dicer2/Act5-GAL4; UAS-snRNAi

6. UAS-Dicer2; UAS-snRNAi; OK107-GAL4

7. sn^{28}; UAS-GFP Singed/238Y-GAL4

8. sn^{28}; UAS myristoylated GFP/238Y-GAL4

<u>Fly brain dissection and staining</u>

For anti-Fasciclin II (Fas II) and/or anti-GFP staining, dissections were performed as previously described in Behnke, et al.[27]. Briefly, whole flies were fixed for 3 hr. in 4% paraformaldehyde (PFA) in 0.01% Triton X100-containing PBS (PBST). The flies were rinsed with PBST 4X for 15 min each, and brains were dissected in PBST. Brains were permeabilized overnight in 0.5% PBST at 4°C with nutation, then blocked in 5% normal Donkey serum (D9663, Millipore-Sigma) in PBST for 90 min at room temperature with nutation. Brains were incubated in mouse anti-Fas II (DSHB, AB528235; 1:100) antibody and/or rabbit anti-GFP (Invitrogen, A11122; 1:1000) in blocking buffer overnight at 4°C with nutation. Brains were washed 3X with PBST and then incubated in secondary antibodies (Invitrogen A-11001 and A11032, 1:500) overnight at 4°C with nutation. The brains were then washed 3X with PBST and mounted in *SlowFade* Gold (ThermoFisher S36936) mounting media (ThermoFisher S36936)

Brains stained for SN were prepared using a protocol adapted from the FlyLight Double Label IHC for Adult Drosophila CNS[28]. Briefly, flies were anesthetized with CO_2 and then put in PBS on ice. Brains were dissected in PBST and then stored on ice prior to fixation. The brains were fixed in 2% PFA in 0.5% PBST for 55 mins at room temperature with nutation. Brains were then washed 4x in 0.5% PBST for 10 mins and

then blocked in 2% donkey serum (D9663, Millipore-Sigma) in 0.5% PBST for 90 mins at room temperature with nutation. Brains were incubated with anti-SN mouse antibody (DSHB, AB528239; undiluted) for 4 hrs. at room temperature with nutation and then incubated at 4°C with nutation for an additional 2 days. Brains were washed 4x for 15 min. in 0.5% PBST and then incubated in secondary antibody (Invitrogen A-11001 or A11032, 1:500) for 4 hrs. at room temperature followed by incubation at 4°C with nutation for 2 days. The brains were again washed 4x for 15 mins in 0.5% PBST and then mounted in *SlowFade* Gold mounting media.

<u>*Drosophila* Image Acquisition and Data Analysis</u>

Unless otherwise noted, all data were collected from three independent biological replicates. *Drosophila* brains were imaged using an Olympus Fluoview FV1000-MPE two-photon confocal system with a Spectra-Physics MaiTi Deepsee Ti:sapphire femtosecond laser, using a 25X/1.05NA water immersion objective. Mushroom body defects were quantified by comparing blinded images to reference control images, as described previously in Kelly, *et al.*[29]. Representative control images were selected for male and female brains from either w^{1118} or Oregon-R flies based on previously reported characteristics of MB morphology. Images of brains from SN null and control flies were then blinded, compared to representative images, and scored for defects. The α- and β-lobes were scored in 4 categories: normal, thin, split (appearance of two axon distinct tracts), or missing. Midline defects were defined as present (either β-lobes crossing the midline or β-lobes failing to reach the midline) or absent (both β-lobes project to the midline and stop). GraphPad Prism Version 10.4.1 for macOS was used to calculate the N-1 Chi-squared test for comparison of two proportions for MB characterization, with each

null line compared to its respective control line. A minimum of 25 brains were analyzed per genotype and 3 independent replicates were collected, except for the Split-GAL4 experiments. For comparisons between multiple proportions, Marascuilo's Procedure was used post-hoc[30].

For split-Gal4, a minimum of ten brains were analyzed per condition. N-1 Chi-squared test was used for statistical analysis for comparison of two proportions for MB characterization, with each null line compared to its respective control line.

Modified Negative Geotaxis Assay (mNGA)

Data were collected as previously described in Ye, *et al.*[31]. Briefly, flies were placed in vials containing a layer of 2% agar at the bottom. Two to four vials were assayed at the same time using a custom 3-D Polylactic Acid (PLA) printed rig. In each trial, one vial contained control flies (either w^{1118} or Oregon-R) while the other contained SN null flies (either sn^{28} or sn^{x2}) or SN null-rescue flies. Flies were separated and tested by sex. For each trial, the rig was lightly tamped three times, and fly movement was recorded using a Panasonic HC-V800 digital video recorder at 60 frames per second. Individual vials from each video were cropped, and the first 10 seconds were trimmed in Image J. Results were quantified by dividing the vial into 10 equal-size bins and counting the number of flies in each bin every second after climbing begins. The climbing index was calculated based the number of flies in each bin [see Ye, *et al.*[31] for details]. All testing took place between ZT 3 and ZT 8 (ZT, Zeitgeber time; lights are turned on at ZT 0 and turned off at ZT 12) and testing occurred between 20 and 23°C under laboratory lighting conditions. The climbing index of each group of flies were plotted and statistically analyzed using paired T-tests for control versus null flies and two-way ANOVA with multiple comparisons

for control, null, and rescue comparisons. Tracing images were obtained by creating Z-stacks of the video frames of the standard deviation of the fly positions five seconds into climbing.

Drosophila western blots

10 fly heads per genotype were collected and homogenized in RIPA buffer (150mM sodium chloride, 50 mM Tris-HCL, 1% Nonidet P-40, 0.5% sodium deoxycholate, 0.1% sodium dodecyl sulfate). The samples were spun down for 10 mins at 14,000 rpm, after which supernatant was collected. Samples were mixed 1:1 with 2x Laemmli buffer and run on a 4–20% Mini-PROTEAN® TGX™ Precast Protein Gel (BioRad, #4561094). After transfer to nitrocellulose, the membrane was blocked in 5% donkey serum in PBS + 0.2% Tween-20 for 1 hr. at room temperature. Blots were then incubated with the following primary antibodies in blocking buffer overnight at 4°: anti-SN mouse antibody (DSHB, AB528239; 1:50) and anti- α/β tubulin sheep antibody (Cytoskeleton, Inc. ATN02; 1:250). Blots were then incubated with the following secondary antibodies in blocking buffer for 1 hr. at room temperature: IRDye® 800CW Goat anti mouse IgG (Li-Cor, 926-32210; 1:1000) and Alexa Fluo3 647 Donkey anti-Sheep IgG (Invitrogen, AB_2535865; 1:1000). Densitometry was performed using FIJI.

Results

Fascin in axonal development in culture

Fascin is the primary actin crosslinker in several membrane protrusions, including filopodia and microspikes, where it arranges up to 15-20 individual parallel actin filaments into tight bundles[32,33]. Using a mouse antibody against fascin (clone 55k2), we first performed immunofluorescent staining on cultured CAD cells. CAD cells form large actin-rich membrane protrusions when plated on laminin-coated surface, which enables high resolution imaging of detailed F-actin structures, including lamellipodia and filopodia[34]. Fascin is concentrated in filopodia and microspikes with thick F-actin bundles, as revealed by actin staining (Supplemental Figure S1A, blue arrows). Interestingly, we also observed fascin signals that appear to be associated with branched F-actin networks in lamellipodia, especially at the leading edge (Supplemental Figure S1B, red arrowheads). While fascin is best known for its role in forming tight F-actin bundles mostly seen in filopodia[35], recent studies suggest that fascin may function in branched F-actin networks to regulate the cofilin-mediated actin turnover underlying lamellipodia dynamics[36,37], further supporting a key role for fascin in actin-based cell motility.

The elaboration and extension of axonal projections during neural development depends on actin-based cell motility[38]. In particular, the motile tip of developing axons features a club-shaped actin-based motile structure called the growth cone that is responsible for guided axonal elongation[8]. We next used cultured primary hippocampal neurons to examine the spatiotemporal pattern of fascin, and its function in axon development. Over the first five days in culture, hippocampal neurons undergo stereotypical neurodevelopmental stages that result in a highly polarized morphology

consisting of a single axonal projection and multiple minor dendritic processes[39-41]. Immunofluorescent imaging of fascin in hippocampal neurons on DIV4 shows that fascin is highly enriched in axonal growth cones, especially in the filopodia of growth cones where β-actin is concentrated (Figure 1A&B, arrows). Consistently, fascin signals were also observed in the lamellipodial region of the growth cone (Figure 1A&B, arrowheads), suggesting a potential role for fascin in the regulation of branched F-actin networks underlying growth cone lamellipodial protrusion. Furthermore, filopodia protruding from the axonal shaft, referred to as collateral filopodia, are also enriched in fascin (Figure 1A &B, asterisks). Previous studies have shown that the formation of collateral filopodia along the axon is essential for branching[42]. Therefore, fascin may also function in axonal branching. It is worth noting that many of the minor processes, which are believed to be the precursors of dendrites[40], also contain a relatively high concentration of fascin (Figure 1A). While a previous study suggested that dendritic filopodia lack fascin[2], it is plausible that fascin could be present and function during the early stages of dendritic development.

Exogenously expressed GFP-tagged fascin concentrates in neuronal growth cones and is mostly associated with F-actin bundles in filopodia, although GFP-fascin signals are also observed in growth cone lamellipodia (Figure 1C, see also Supplemental movie S1). Consistent with previous reports[43,44], activation of protein kinase C (PKC) via treatment with 12-O-tetradecanoylphorbol-13-acetate (TPA) causes the dissociation of fascin from F-actin bundles, leading to growth cone collapse (Figure 1C, see also Supplemental movie S1). This observation supports the notion that the cellular localization of fascin in the growth cones is associated with its F-actin bundling activity.

Therefore, fascin likely plays an important role in actin-based axon development leading to the wiring of the brain.

To determine the role of fascin in growth cone motility and axon development, we utilized a potent small molecule inhibitor of fascin, NP-G2-044 (hereafter referred to as G2 for simplicity), which binds to fascin and prevents bundling activity[45,46]. We first tested the effects of this inhibitor in CAD cells by treating them with either 15 μM G2 or DMSO as a vehicle control for 30 mins or 4 hrs., followed by methanol fixation and immunofluorescent staining to determine the effect of G2 treatment on the spatial distribution of fascin and β-actin. We observed that G2 treatment resulted in a gradual shift in fascin localization away from the actin-rich lamellipodia and filopodia at the cell edge and towards the central region of the cell (Figure 2A-B). We quantified the peripheral to central shift of fascin by measuring the ratio of the mean fascin fluorescence intensity at the cell edge over the mean fascin intensity in the center of the cell. At both the 30-minute and 4 hr. timepoints the mean fluorescence intensity of fascin in filopodia, relative to the cell center, significantly decreases in response to treatment with G2 (Figure 2A-B). This suggests that 15 μM G2 is sufficient to inhibit fascin-binding to actin in CAD cells. As a control, we also determined the ratio of β-actin intensity in filopodia and the cytosol. We did not detect a shift in β-actin localization to the cytosol after treatment with G2, showing that the inhibitor specifically affects fascin1 localization to filopodia (Figure 2 A-B).

We next investigated the effect of G2 treatment on filopodia morphology in neuronal growth cones by treating cultured hippocampal neurons on DIV3 with either DMSO or 15 μM G2 for 24 hrs., followed by methanol fixation and immunofluorescent labeling of fascin and β-actin on DIV4. After 24 hr. G2 exposure, axonal growth cones

had significantly fewer filopodia (Figure 2 C-D). In addition, the remaining filopodia were shorter in length (Figure 2 C-D). These findings are consistent with neurons cultured from fascin1 KO mice[22], further suggesting that G2 successfully inhibits fascin activity in cultured neurons.

We next examined whether fascin inhibition impacts axon outgrowth and branching during early neuronal development. We applied G2 to DIV1 hippocampal neurons and examined their morphology on DIV5. Here, neurons were treated with either 15 µM of G2 or DMSO as a vehicle control. In mice, G2 was shown to have a half-life ($T_{1/2}$) of ~ 4 hrs.[47]. Therefore, to ensure that fascin was constantly inhibited for 4 days, fresh G2 or DMSO was applied to neurons every day. We fixed the neurons on DIV5 with PFA and stained for tau to visualize axons (Figure 3A), followed by tracing and quantification of axon length and branching for each condition (Figure 3B). We observe that neurons treated with 15 µM G2 exhibited reduced axon length and branching when compared to DMSO control neurons (Figure 3A-D). These data indicate that fascin-mediated actin bundling promotes axonal growth and branching.

Drosophila melanogaster as an _in vivo_ model to study axon development and brain wiring

Neurons in culture allow high-resolution imaging, easy manipulation, and quantitative assessment of their axonal growth and branching. However, they lack the complex and dynamic molecular and cellular components found _in vivo_ that guide their axonal elongation to specific targets for previse brain wiring. To investigate how fascin regulates axonal development and brain wiring _in vivo_, we used _Drosophila melanogaster_ as a model system. Unlike mammals, _Drosophila_ expresses a single ortholog of fascin called Singed (SN), thus eliminating the possibility of compensation by other fascin

isoforms. SN null flies exhibit defects in actin-based bristle morphology and female sterility due to defective actin structures in nurse cells[48], demonstrating that SN functions in F-actin bundling. However, little is known about the expression pattern of SN in *Drosophila* brains and its potential functions in central nervous system development[48]. Previous work showed that mushroom body (MB) γ-neurons isolated from SN null fly brains exhibit a filagree phenotype when cultured *in vitro*, but SN expression in the fly brain and its *in vivo* function were not examined[49]. We thus first examined the expression of SN in *Drosophila* brains by immunofluorescent labeling and whole brain two-photon imaging.

<u>SN is expressed in mushroom body neurons</u>

Using an antibody against *Drosophila* SN, we detect SN signal in many neuronal cell bodies in the cortex and optic lobes (Figure 4A). Strikingly, SN highlights the mushroom body (MB), the part of the *Drosophila* central nervous system critical for olfactory learning, olfactory memory, and locomotion (Figure 4A)[50-52]. SN signals are enriched in both the α- and β- lobes, which are the axonal projections of Kenyon neurons (Figure 4A). Interestingly, the SN signal appears to be restricted within a narrow subset of MB axons (Figure 4A). To confirm this, we used a MB-GAL4 (238-Y)[50] line to drive expression of UAS-Myristoylated-GFP throughout the entire MB and then co-stained with anti-SN antibody. These data revealed that SN is only expressed in axons found within the centers of the α- and β- lobes (Figure 4B). Together, our findings confirm the expression of SN in the adult *Drosophila* brain and indicate the selective expression of SN in distinct populations of neurons not only within the central nervous system (CNS), but also within the MB.

To investigate the function of fascin in *Drosophila* brain development, we obtained a null SN line (sn[28]) from Dr. Tina Tootle[53]. We first performed a western blot of female sn[28] heterozygous and homozygous null fly brains to confirm SN knockout (Supp. Figure 2A). We also confirmed via immunocytochemistry that SN is not expressed in sn[28] brains (Supp. Figure 2B). As a negative control we first looked at pigment dispersing factor (PDF) neurons, which do not appear to express SN (Figure 4A), to see if there were any effects on their morphology. This subgroup of ventral lateral neurons belongs to the circadian clock neuron family and project axons from the optic lobe of the brain to the central brain[54]. As expected, we did not observe any differences between PDF neuron morphology in control and sn[28] brains (Supp. Figure 2C).

Loss of SN results in axonal defects *in vivo*

The MB α- and β- lobes are formed by axons extended from Kenyon cells that undergo bifurcation and then project dorsally and ventrally (Figure 5A). The formation of MB lobes is dependent on both axon bifurcation and guidance[55-59], giving us a system to observe the effects of fascin depletion on both processes. Brains were dissected from control w[1118 flies] flies and compared to sn[28] flies. The brains were stained for Fas II, a marker for the MB α/β lobes[60]. We observed significant defects in the morphology of MB α- and β- lobes in SN null flies (Figure 5B). In control brains, the projecting axons bifurcate to form α and β lobes, with the α lobes projecting vertically and the β lobes projecting medially and stopping at the midline of the brain. The control α and β lobes also appear to have tightly bundled axon tracks. In sn[28] flies, we observe a range of defects, including missing, thin, or split (i.e. loosely bundled) α and β lobes (Figure 5C&D). We also observe a highly penetrative midline defect, with the β lobes either crossing over the midline or in

some cases failing to reach the midline (Figure 5E&F). The full statistical comparison of defects in control versus null brains is shown in Table 1. We conclude that global loss of SN results in significant MB α- and β- lobe defects.

To better understand the MB defects in SN null fly brains, we employed a Split-GAL4 line to drive the expression of myristoylated GFP in a subset of MB axons[61]. In wildtype flies (Split-GAL4/UAS-myristoylated GFP; Split-GAL4/+), MB axons form stereotypical α- and β- lobes (Figure 6A), see Supplemental Movie 2 for 3-dimensional visualization. In sn[28] flies (sn[28]; Split-GAL4/UAS-myristoylated GFP; Split-GAL4/+), we observed similar MB defects as described above (Figure 5), including MB lobes displaying thin, split, and missing phenotypes, as well as midline crossing defects (Figure 6B&C). Several of the guidance defects are highlighted by 3D rendering of the MB lobes in sn[28] brains, including the midline crossing defects and split β-lobe (Figure 6B&C, orange and yellow arrows, see also Supplemental Movie S3). We also observed several instances of β lobe axons appearing to initially project medially, only to turn back towards the branch point (Figure 6C, white arrow, see also Supplemental Movie S4) or project vertically (Figure 6C, orange arrow, see Supplemental Movie S4). The MB lobe and midline defects observed in the sn[28] flies expressing the split GAL4 driver are similar to those observed in the sn[28] flies stained for Fas II (Figure 6 D&E). These data indicate that SN is required for proper axon guidance.

Loss of SN results in sensorimotor defects

Previous studies have implicated the MB in locomotion, with most MB output neurons having direct connections to the central complex, which controls locomotion and navigation in flies[51,61-63]. Consequently, locomotor defects are commonly associated with

MB lobe defects[64,65]. We next used the modified negative geotaxis assay[31] to assess if SN loss affects startle-induced climbing behavior. We found that there is a significant delay in climbing response for both SN null lines when compared to controls, with SN null flies taking longer to reach the top of the vial (Figure 7A&C, Supplemental Movie 5). Negative geotaxis represents a sensorimotor response that requires flies to first detect their position in respect to gravity and then move against the pull of gravity. Our data thus show that SN null flies exhibit significant sensorimotor defects as compared to control flies (Figure 7B&D).

To confirm the robust phenotypes observed in SN[28] flies, we obtained a second SN null line (sn[x2])[48] to examine MB lobe morphology and the effects on sensorimotor behavior. We used Oregon-R flies as the control line and found that sn[x2] flies displayed similar defects in MB α- and β- lobe formation, midline crossing, and sensorimotor function (Supp. Figure 3). Together, these data confirm the importance of SN for MB development and brain function.

While SN appears to be strongly expressed in MB neurons and SN null lines display MB lobe defects, we wanted to determine if malformed MB lobes and deficits in startle-induced climbing were due to the loss of SN specifically in MB neurons. To answer this question, we knocked down SN expression using the GAL4-UAS system to express an SN shRNA construct (snRNAi)[66]. For the initial characterization of SN knockdown, the Actin5-GAL4[67] driver was used to globally express UAS-snRNAi and UAS-Dicer2[68]. We found that expression of SN snRNAi in combination with Dicer2, resulted in an ~60-70% reduction in SN protein levels as compared to control flies (Figure 8A). Flies that expressed both the UAS-snRNAi and UAS-Dicer2 constructs also displayed the gnarled

bristle phenotype seen in the SN null flies as previously reported[48] (Figure 8B). Next, we used OK107-GAL4[69] to drive the expression of snRNAi specifically in MB neurons and their axons. We observed marked MB defects in UAS-Dicer2;UAS-snRNAi;OK107-GAL4 brains compared to control brains (Figure 8 C&D), although they appear to be less severe than SN null brains. This is likely due to the incomplete KD of SN which is consistent with the lack of midline crossing defects in these brains. our findings suggest that SN functions autonomously in MB to regulate its development and patterning.

<u>MB-specific expression of SN is able to rescue defects</u>

To further determine the role of SN in MB development, we used a MB-specific GAL4 driver (238Y)[50] to drive the expression of either UAS-myristoylated-GFP (UAS myrs-GFP) as a negative control or wildtype UAS-GFP-Singed (UAS SN-GFP)[70] constructs in the sn^{28} background. sn^{28} flies exhibiting the SN-null bristle defect[48] and expressing both 238Y-GAL4 and either of the UAS-GFP constructs were collected and compared to control w^{1118} and sn^{28} flies in the mNGA and staining for MB morphology. We first examined startle-induced climbing in sn^{28} female flies expressing either UAS-myrs-GFP or wildtype UAS SN-GFP specifically in their MB neurons. Significantly, we found that MB-specific expression of SN-GFP in the sn^{28} background was able to restore startle-induced climbing to closely resemble control w1118 flies (Figure 9A). On the other hand, MB-specific expression of myrs-GFP in the sn^{28} background did not affect the impaired climbing behavior, and they exhibited the same deficits as sn^{28} flies (Figure 9A). Next, we examined whether MB-specific GFP-SN expression could rescue the structural MB defects in SN null flies. We found that the expression of UAS SN-GFP via the 238Y-Gal4 driver was sufficient to fully rescue MB α- and β- lobe defects to the levels seen in

control w[1118] flies (Figure 9B&C). Likewise, MB-specific SN-GFP expression successfully rescued the sn[28] midline defects (Figure 9D). Together with our snRNAi results, these data indicate that SN functions in MB neurons to regulate the formation and patterning of α- and β- lobes during development.

It should be noted that MB-specific SN-GFP expression in male sn[28] flies was also able to rescue both structural and behavioral deficits, but with some differences from female flies (Supplemental figure 4A-D). Whether there exists a sex difference in terms of SN-GFP expression or MB development remains to be investigated. Nonetheless, our *in vivo* data, together with results from cell cultures, not only demonstrate the importance of the fascin family of actin bundling proteins in actin based neuronal motility but also provide the first evidence for Fascin/SN in brain development and function.

Discussion

      As the major F-actin bundling protein, fascin is concentrated in protrusive cellular structures such as invadopodia and filopodia that play critical roles in directed cell motility and invasion of the cellular matrix[10,13,71-75]. Fascin is well known for its concentration in the filopodia of nerve growth cones[14], the sensory motile apparatus that is responsible for sensing the environmental cues and subsequently steering the axons to their specific targets. Few studies have been conducted so far to investigate the role of fascin in axon development and brain wiring in culture or *in vivo*. Here, we take advantage of mammalian primary neuronal cultures and a *Drosophila* model, combined with pharmacological, genetic and imaging approaches, to establish, for the first time that fascin plays a critical role in axonal development and brain wiring. Here, primary neurons in culture have enabled us to not only reveal the distinct cellular localization of fascin in developing neurons but also understand how fascin-mediated actin-bundling regulates axon outgrowth and branching. Additionally, the use of *Drosophila* as an *in vivo* model reveals the critical importance of fascin in axon development, brain wiring and function. Through a combination of genetic manipulation, whole brain imaging, and sensorimotor behavioral assays, we provide direct evidence that fascin is required for the proper development of the mushroom body (MB), which is analogous to the mammalian hippocampus[76]. Loss of fascin in *Drosophila* leads to defects in both α- and β- lobes, which represent guided axonal projections from Kenyon cells[60]. We additionally demonstrate that loss of fascin leads to deficits in startle-induced climbing. Importantly, MB-specific expression of fascin is sufficient to rescue observed structural and behavioral deficits. Together, our study has established a critically important role for fascin-mediated actin bundling in brain development and function.

Fascin in neuronal development

<u>Fascin in developing neurons</u>

Primary rat hippocampal neurons in culture represent a well-established model to investigate axon formation as they undergo distinct stages leading to the establishment of a single axon within a one-week period[39-41,77]. Importantly, hippocampal neurons in culture allow high-resolution imaging that not only helps to determine the cellular localization of fascin at various stages of axon development but also enables the tracking and quantification of axonal branching and the growth of individual neurons. In cultured hippocampal neurons, we found that fascin is highly enriched in growth cones, especially growth cone filopodia. Importantly, fascin signals were also detected in growth cone lamellipodia, suggesting a potential role in the regulation of branched F-actin networks underlying lamellipodial protrusions[78,79]. Furthermore, fascin signals were also found enriched in collateral filopodial protrusions, suggesting a potential role for fascin-mediated actin dynamics in axonal branching. Axonal branching is an important developmental event that enables a neuron to establish synaptic connections with multiple targets[80], Axon branching occurs through two mechanisms. The first is when growth cones split to form two separate axon paths, also known as bifurcation, and the second is collateral branch formation, which occurs when new branch points are established on the axon shaft[81]. For either mechanism, the presence of filopodia is important to initiate branch formation[82]. Indeed, inhibition of fascin by the small molecule inhibitor NP-G2-044 prominently decreased axon branch points and the outgrown of secondary and tertiary axon branches, supporting a role for fascin in axon branching and growth. NP-G2-044 is a well-established and highly-specific fascin inhibitor[83]. Importantly, crystal structure and cryo-electron microscopy studies have gained the mechanistic understanding of how and

where this small compound binds to fascin to interfere with its bundling activities[84,85]. As a result, the rapid action of fascin inhibition by NP-G2-044 has enabled us to gain important insights into the immediate, short-term, and long-term effects of fascin inhibition on axonal development in culture. The subsequent use of a *Drosophila* model combined with genetic manipulation has substantiated our work on cultured neurons and allowed us to elucidate the *in vivo* function of fascin.

<u>Expression and roles of fascin in the mushroom body</u>

Unlike vertebrates, *Drosophila* expresses only one orthologue of fascin, called Singed (SN). While SN has been shown to be required for actin bundles in bristles[86] and nurse cells[48] in female flies, the expression pattern of SN in *Drosophila* brains was unknown. Previous work by Kraft, *et al.*[49] found that SN is required for MB γ-neuron morphology in culture but it did not evaluate SN expression in the MB nor its function in the MB. We have shown for the first time that SN is expressed in a narrow subset of adult MB α- and β- lobe axons. We speculate that this subset of SN-expressing axons come from late-born neurons, as they appear similar to the subset of neurons labeled by the NP7175 GAL4 driver which marks MB neurons born later in CNS development that go on to form the central region of the α- and β- lobes[87]. We speculate that SN is likely more widely expressed in the brain including MB neurons during early development, like that observed in rodent and human[20,21]. It is possible that the reason we see SN staining in a narrow subset of adult MB axons because these later-born neurons require SN for axon pathfinding. Axons that have already projected to their targets likely no longer require SN and thus are not positive for SN. Future work could further examine the expression profile of SN in MB neurons in flies of different stages.

Fascin in neuronal development

The observed defects in MB lobes most closely resemble guidance errors. MB axons are guided by multiple guidance factors and signaling pathways[55-57,88-90]. One of the most well-studied and conserved pathways is the Slit/Robo family which governs midline crossing in *Drosophila*[90]. Previous work in mammalian cultured neurons has shown that Slit-mediated repulsion is dependent on filopodia dynamics[91]. Here, we show that loss of fascin in MB axons results in a highly penetrative midline defect. This further supports a role for fascin's regulation of filopodia in axon guidance. Our *Drosophila* Split-GAL4 data show that the axon paths extending from Kenyon cell bodies appear to successfully bifurcate and then fail to follow proper guidance pathways, hence the prominence of the "split lobe" phenotype that we observe in SN null brains. We speculate that this phenotype is due to α-lobe axons following β-lobe guidance pathways and vice versa. The determination of which axons become α-lobe and which become β-lobe is dependent on tightly regulated expression and localization of the Eph/Ephrin family of guidance molecules[58]. The interaction of fascin and this family has not been studied in neurons, however a study in prostate cancer cells showed that the addition of an ephrin ligand resulted in an increase in fascin-positive filopodia[92], suggesting that the split lobe phenotype we observed could be due to disruption in response to Eph/Ephrin signaling. At this moment, we are unable to definitively determine if MB axon bifurcation is affected by SN loss. Future experiments at the single cell level, such as mosaic analysis with a repressible cell marker (MARCM)[93], will likely provide the answer.

Fascin's function in axon development and brain wiring

Developing axons are guided by a myriad of attractive and repulsive cues that are surface-bound or diffusible in nature[90,94-102]. It is believed that the motile growth cones at

the tip of axons sense the environment via dynamic filopodia and steer the axons through actin-based motility[103-105]. Given that fascin is mostly enriched in growth cone filopodia, it may serve as an important cytoskeletal component to support the formation and extension of filopodia. On the other hand, fascin can be phosphorylated at serine 39 by protein kinase C (PKC), and S39 phosphorylation diminishes its bundling activity[43,44]. An additional phosphorylation site of fascin (S274)[106] has also been identified along with two lysines (K247, K250)[107] that can be monoubiquitinated. Therefore, fascin-mediated F-actin bundles underlying filopodial protrusions can be dynamically regulated in response to signaling cascades during axon pathfinding. Furthermore, fascin has been shown to interact with Dishevelled-associated activator of morphogenesis (DAAM1), a member of the formin family[108]. DAAM functions as a link between the Wnt5/planar cell polarity (PCP) guidance system and the actin cytokeleton, which guides MB lobe projection[57] in *Drosophila* neurons. DAAM1 is also enriched in the newly formed axons of MB neurons, where it is required for proper lobe projection[109]. Finally, both DAAM1 and fascin1 have been shown to bind to microtubules and actin,[110,111] raising the possibility that fascin may coordinate these two cytoskeletal systems for guided axon elongation during brain wiring. Therefore, it is reasonable to postulate that fascin, and fascin-mediated actin bundling activities may function beyond just making F-actin bundles for filopodia but be actively involved in the intricate signal transduction that underlies axon guidance and brain wiring.

Supplemental methods

PDF neuron staining

Whole flies were fixed for 3 h in 4% paraformaldehyde (PFA) in phosphate-buffered saline containing 0.01% Triton-X (PBST). The flies were rinsed 4 times for 15 min each with PBST and brains were subsequently dissected in PBST. Brains were permeabilized overnight in 0.5% PBST at 4°C with nutation then blocked in 5% normal goat serum in PBST for 90 min at room temperature with nutation. After blocking, the brains were incubated in mouse anti-PDF (DSHB, PDF C7, AB_760350 AB_2315084) antibody diluted 1:100 in blocking buffer overnight at 4°C with nutation. Brains were washed 3 times with PBST and then incubated in secondary antibody (Invitrogen A-11001) overnight at 4°C with nutation. The brains were then washed 3 times in PBS and mounted in *SlowFade* Gold mounting media (ThermoFisher S36936)

Works Cited:

1. Dominguez, R., and Holmes, K.C. (2011). Actin structure and function. Annu Rev Biophys *40*, 169-186. 10.1146/annurev-biophys-042910-155359.
2. Svitkina, T. (2018). The Actin Cytoskeleton and Actin-Based Motility. Cold Spring Harb Perspect Biol *10*. 10.1101/cshperspect.a018267.
3. Pollard, T.D. (2016). Actin and Actin-Binding Proteins. Cold Spring Harb Perspect Biol *8*. 10.1101/cshperspect.a018226.
4. Tseng, Y., Kole, T., Lee, J., Fedorov, E., Almo, S., Schafer, B., and Wirtz, D. (2005). How actin crosslinking and bundling proteins cooperate to generate an enhanced cell mechanical response. Biochemical and biophysical research communications *334*, 183-192. 10.1016/j.bbrc.2005.05.205.
5. Rajan, S., Kudryashov, D.S., and Reisler, E. (2023). Actin Bundles Dynamics and Architecture. Biomolecules *13*. 10.3390/biom13030450.
6. Stevenson, R.P., Veltman, D., and Machesky, L.M. (2012). Actin-bundling proteins in cancer progression at a glance. Journal of Cell Science *125*, 1073-1079. 10.1242/jcs.093799.
7. Cajal, S.R. (1890). Textura de las circunvolutiones cerebrales de los mamíferos inferiores: nota preventiva.
8. Letourneau, P.C. (1983). Axonal growth and guidance. Trends in Neurosciences *6*, 451-455. 10.1016/0166-2236(83)90217-5.
9. Armijo-Weingart, L., and Gallo, G. (2017). It takes a village to raise a branch: Cellular mechanisms of the initiation of axon collateral branches. Mol Cell Neurosci *84*, 36-47. 10.1016/j.mcn.2017.03.007.
10. Dent, E.W., Gupton, S.L., and Gertler, F.B. (2011). The growth cone cytoskeleton in axon outgrowth and guidance. Cold Spring Harb Perspect Biol *3*, a001800. 10.1101/cshperspect.a001800.
11. Zheng, J., Wan, J., and Poo, M. (1996). Essential role of filopodia in chemotropic turning of nerve growth cone induced by a glutamate gradient. The Journal of Neuroscience *16*, 1140-1149. 10.1523/jneurosci.16-03-01140.1996.
12. Chien, C.B., Rosenthal, D.E., Harris, W.A., and Holt, C.E. (1993). Navigational errors made by growth cones without filopodia in the embryonic Xenopus brain. Neuron *11*, 237-251.
13. Vignjevic, D., Kojima, S., Aratyn, Y., Danciu, O., Svitkina, T., and Borisy, G.G. (2006). Role of fascin in filopodial protrusion. J Cell Biol *174*, 863-875. 10.1083/jcb.200603013.
14. Cohan, C.S., Welnhofer, E.A., Zhao, L., Matsumura, F., and Yamashiro, S. (2001). Role of the actin bundling protein fascin in growth cone morphogenesis: Localization in filopodia and lamellipodia. Cell Motility *48*, 109-120. https://doi.org/10.1002/1097-0169(200102)48:2<109::AID-CM1002>3.0.CO;2-G.
15. Jayo, A., and Parsons, M. (2010). Fascin: A key regulator of cytoskeletal dynamics. The International Journal of Biochemistry & Cell Biology *42*, 1614-1617. 10.1016/j.biocel.2010.06.019.
16. Chen, Y., Tian, T., Li, Z.-Y., Wang, C.-Y., Deng, R., Deng, W.-Y., Yang, A.-k., Chen, Y.-F., and Li, H. (2019). FSCN1 is an effective marker of poor prognosis

and a potential therapeutic target in human tongue squamous cell carcinoma. Cell Death & Disease *10*, 356.

17. Wu, D., Chen, L., Liao, W., Ding, Y., Zhang, Q., Li, Z., and Liu, L. (2010). Fascin1 expression predicts poor prognosis in patients with nasopharyngeal carcinoma and correlates with tumor invasion. Annals of oncology *21*, 589-596.

18. Fu, H., Wen, J.-F., Hu, Z.-L., Luo, G.-Q., and Ren, H.-Z. (2009). Knockdown of fascin1 expression suppresses the proliferation and metastasis of gastric cancer cells. Pathology *41*, 655-660. https://doi.org/10.3109/00313020903273100.

19. Hwang, J.H., Smith, C.A., Salhia, B., and Rutka, J.T. (2008). The Role of Fascin in the Migration and Invasiveness of Malignant Glioma Cells. Neoplasia *10*, 149-159. https://doi.org/10.1593/neo.07909.

20. De Arcangelis, A., Georges-Labouesse, E., and Adams, J.C. (2004). Expression of fascin-1, the gene encoding the actin-bundling protein fascin-1, during mouse embryogenesis. Gene Expr Patterns *4*, 637-643. 10.1016/j.modgep.2004.04.012.

21. Zhang, F.R., Tao, L.H., Shen, Z.Y., Lv, Z., Xu, L.Y., and Li, E.M. (2008). Fascin expression in human embryonic, fetal, and normal adult tissue. The journal of histochemistry and cytochemistry : official journal of the Histochemistry Society *56*, 193-199. 10.1369/jhc.7A7353.2007.

22. Yamakita, Y., Matsumura, F., and Yamashiro, S. (2009). Fascin1 is dispensable for mouse development but is favorable for neonatal survival. Cell motility and the cytoskeleton *66*, 524-534. 10.1002/cm.20356.

23. Valenzuela, D.M., Murphy, A.J., Frendewey, D., Gale, N.W., Economides, A.N., Auerbach, W., Poueymirou, W.T., Adams, N.C., Rojas, J., Yasenchak, J., et al. (2003). High-throughput engineering of the mouse genome coupled with high-resolution expression analysis. Nat Biotechnol *21*, 652-659. 10.1038/nbt822.

24. Gu, J., Firestein, B.L., and Zheng, J.Q. (2008). Microtubules in dendritic spine development. J Neurosci *28*, 12120-12124.

25. Qi, Y., Wang, J.K., McMillian, M., and Chikaraishi, D.M. (1997). Characterization of a CNS cell line, CAD, in which morphological differentiation is initiated by serum deprivation. J Neurosci *17*, 1217-1225. 10.1523/jneurosci.17-04-01217.1997.

26. Arshadi, C., Günther, U., Eddison, M., Harrington, K.I.S., and Ferreira, T.A. (2021). SNT: a unifying toolbox for quantification of neuronal anatomy. Nature Methods *18*, 374-377. 10.1038/s41592-021-01105-7.

27. Behnke, J.A., Ye, C., Setty, A., Moberg, K.H., and Zheng, J.Q. (2021). Repetitive mild head trauma induces activity mediated lifelong brain deficits in a novel Drosophila model. Scientific Reports *11*, 9738. 10.1038/s41598-021-89121-7.

28. FlyLight Double label IHC for adult Drosophila CNS. (2015). Janelia.

29. Kelly, S.M., Bienkowski, R., Banerjee, A., Melicharek, D.J., Brewer, Z.A., Marenda, D.R., Corbett, A.H., and Moberg, K.H. (2016). The Drosophila ortholog of the Zc3h14 RNA binding protein acts within neurons to pattern axon projection in the developing brain. Developmental Neurobiology *76*, 93-106. https://doi.org/10.1002/dneu.22301.

30. Marascuilo, L.A. (1966). Large-sample multiple comparisons. Psychological Bulletin *65*, 280-290. 10.1037/h0023189.

31.   Ye, C., Ho, R., Moberg, K.H., and Zheng, J.Q. (2024). Adverse impact of female reproductive signaling on age-dependent neurodegeneration after mild head trauma in Drosophila. Elife *13*. 10.7554/eLife.97908.

32.   Otto, J.J., Kane, R.E., and Bryan, J. (1979). Formation of filopodia in coelomocytes: Localization of Fascin, a 58,000 Dalton actin cross-linking protein. Cell *17*, 285-293. https://doi.org/10.1016/0092-8674(79)90154-5.

33.   Aramaki, S., Mayanagi, K., Jin, M., Aoyama, K., and Yasunaga, T. (2016). Filopodia formation by crosslinking of F-actin with fascin in two different binding manners. Cytoskeleton *73*, 365-374. https://doi.org/10.1002/cm.21309.

34.   Vitriol, E.A., McMillen, L.M., Kapustina, M., Gomez, S.M., Vavylonis, D., and Zheng, J.Q. (2015). Two functionally distinct sources of actin monomers supply the leading edge of lamellipodia. Cell Rep *11*, 433-445. 10.1016/j.celrep.2015.03.033.

35.   Vignjevic , D., Kojima , S.-i., Aratyn , Y., Danciu , O., Svitkina , T., and Borisy , G.G. (2006). Role of fascin in filopodial protrusion. Journal of Cell Biology *174*, 863-875. 10.1083/jcb.200603013.

36.   Nemethova, M., Auinger, S., and Small, J.V. (2008). Building the actin cytoskeleton: filopodia contribute to the construction of contractile bundles in the lamella. J Cell Biol *180*, 1233-1244. 10.1083/jcb.200709134.

37.   Chikireddy, J., Lengagne, L., Le Borgne, R., Durieu, C., Wioland, H., Romet-Lemonne, G., and Jégou, A. (2024). Fascin-induced bundling protects actin filaments from disassembly by cofilin. J Cell Biol *223*. 10.1083/jcb.202312106.

38.   Omotade, O.F., Pollitt, S.L., and Zheng, J.Q. (2017). Actin-based growth cone motility and guidance. Mol Cell Neurosci *84*, 4-10. 10.1016/j.mcn.2017.03.001.

39.   Banker, G. (2018). The Development of Neuronal Polarity: A Retrospective View. The Journal of Neuroscience *38*, 1867-1873. 10.1523/jneurosci.1372-16.2018.

40.   Dotti, C.G., Sullivan, C.A., and Banker, G.A. (1988). The establishment of polarity by hippocampal neurons in culture. J Neurosci *8*, 1454-1468. 10.1523/jneurosci.08-04-01454.1988.

41.   Kaech, S., and Banker, G. (2006). Culturing hippocampal neurons. Nature Protocols *1*, 2406-2415. 10.1038/nprot.2006.356.

42.   Gallo, G. (2011). The cytoskeletal and signaling mechanisms of axon collateral branching. Dev Neurobiol *71*, 201-220. 10.1002/dneu.20852.

43.   Ono, S., Yamakita, Y., Yamashiro, S., Matsudaira, P.T., Gnarra, J.R., Obinata, T., and Matsumura, F. (1997). Identification of an actin binding region and a protein kinase C phosphorylation site on human fascin. Journal of Biological Chemistry *272*, 2527-2533.

44.   Yamakita, Y., Ono, S., Matsumura, F., and Yamashiro, S. (1996). Phosphorylation of Human Fascin Inhibits Its Actin Binding and Bundling Activities (∗). Journal of Biological Chemistry *271*, 12632-12638. 10.1074/jbc.271.21.12632.

45.   Chung, V., Tsai, F., Chen, W., Von Hoff, D.D., Garmey, E.G., Zhang, J.J., and Huang, X.Y. (2022). 96 (PB086) - NP-G2-044, a First-in-Class Fascin Inhibitor, Inhibits Growth and Metastasis of Gynecologic Cancers. European Journal of Cancer *174*, S34. https://doi.org/10.1016/S0959-8049(22)00891-7.

46.   Chung, V., Tsai, F., Jhaveri, K., Wang, Y., Hoff, D.D.V., Garmey, E.G., Zhang, J.J., and Huang, X.-Y. (2019). Abstract C053: NP-G2-044, a novel fascin inhibitor,

blocks tumor metastasis and increases antitumor immune response. Molecular Cancer Therapeutics *18*, C053-C053. 10.1158/1535-7163.Targ-19-c053.

47. Wang, Y., Zhang, J.J., and Huang, X.Y. (2020). Anti-Metastasis Fascin Inhibitors Decrease the Growth of Specific Subtypes of Cancers. Cancers (Basel) *12*. 10.3390/cancers12082287.

48. Cant, K., Knowles, B.A., Mooseker, M.S., and Cooley, L. (1994). Drosophila singed, a fascin homolog, is required for actin bundle formation during oogenesis and bristle extension. J Cell Biol *125*, 369-380. 10.1083/jcb.125.2.369.

49. Kraft, R., Escobar, M.M., Narro, M.L., Kurtis, J.L., Efrat, A., Barnard, K., and Restifo, L.L. (2006). Phenotypes of Drosophila Brain Neurons in Primary Culture Reveal a Role for Fascin in Neurite Shape and Trajectory. The Journal of Neuroscience *26*, 8734-8747. 10.1523/jneurosci.2106-06.2006.

50. Tettamanti, M., Armstrong, J.D., Endo, K., Yang, M.Y., Furukubo-Tokunaga, K., Kaiser, K., and Reichert, H. (1997). Early development of the Drosophila mushroom bodies, brain centres for associative learning and memory. Development Genes and Evolution *207*, 242-252. 10.1007/s004270050112.

51. Martin, J.R., Ernst, R., and Heisenberg, M. (1998). Mushroom bodies suppress locomotor activity in Drosophila melanogaster. Learn Mem *5*, 179-191.

52. McGuire, S.E., Le, P.T., and Davis, R.L. (2001). The role of Drosophila mushroom body signaling in olfactory memory. Science *293*, 1330-1333. 10.1126/science.1062622.

53. Lamb, M.C., Kaluarachchi, C.P., Lansakara, T.I., Mellentine, S.Q., Lan, Y., Tivanski, A.V., and Tootle, T.L. (2021). Fascin limits Myosin activity within Drosophila border cells to control substrate stiffness and promote migration. eLife *10*, e69836. 10.7554/eLife.69836.

54. Reinhard, N., Schubert, F.K., Bertolini, E., Hagedorn, N., Manoli, G., Sekiguchi, M., Yoshii, T., Rieger, D., and Helfrich-Förster, C. (2022). The Neuronal Circuit of the Dorsal Circadian Clock Neurons in Drosophila melanogaster. Frontiers in Physiology *13*. 10.3389/fphys.2022.886432.

55. Oliva, C., Soldano, A., Mora, N., De Geest, N., Claeys, A., Erfurth, M.L., Sierralta, J., Ramaekers, A., Dascenco, D., Ejsmont, R.K., et al. (2016). Regulation of Drosophila Brain Wiring by Neuropil Interactions via a Slit-Robo-RPTP Signaling Complex. Dev Cell *39*, 267-278. 10.1016/j.devcel.2016.09.028.

56. Reynaud, E., Lahaye, L.L., Boulanger, A., Petrova, I.M., Marquilly, C., Flandre, A., Martianez, T., Privat, M., Noordermeer, J.N., Fradkin, L.G., and Dura, J.M. (2015). Guidance of Drosophila Mushroom Body Axons Depends upon DRL-Wnt Receptor Cleavage in the Brain Dorsomedial Lineage Precursors. Cell Rep *11*, 1293-1304. 10.1016/j.celrep.2015.04.035.

57. Shimizu, K., Sato, M., and Tabata, T. (2011). The Wnt5/Planar Cell Polarity Pathway Regulates Axonal Development of the Drosophila Mushroom Body Neuron. Journal of Neuroscience *31*, 4944-4954. 10.1523/jneurosci.0154-11.2011.

58. Boyle, M., Nighorn, A., and Thomas, J.B. (2006). Drosophila Eph receptor guides specific axon branches of mushroom body neurons. Development *133*, 1845-1854. 10.1242/dev.02353.

59. Lee, T., Lee, A., and Luo, L. (1999). Development of the Drosophila mushroom bodies: sequential generation of three distinct types of neurons from a neuroblast. Development *126*, 4065-4076. 10.1242/dev.126.18.4065.
60. Kurusu, M., Awasaki, T., Masuda-Nakagawa, L.M., Kawauchi, H., Ito, K., and Furukubo-Tokunaga, K. (2002). Embryonic and larval development of the Drosophila mushroom bodies: concentric layer subdivisions and the role of fasciclin II. Development *129*, 409-419. 10.1242/dev.129.2.409.
61. Aso, Y., Hattori, D., Yu, Y., Johnston, R.M., Iyer, N.A., Ngo, T.-T.B., Dionne, H., Abbott, L.F., Axel, R., Tanimoto, H., and Rubin, G.M. (2014). The neuronal architecture of the mushroom body provides a logic for associative learning. eLife *3*, e04577. 10.7554/eLife.04577.
62. Li, F., Lindsey, J.W., Marin, E.C., Otto, N., Dreher, M., Dempsey, G., Stark, I., Bates, A.S., Pleijzier, M.W., Schlegel, P., et al. (2020). The connectome of the adult Drosophila mushroom body provides insights into function. Elife *9*. 10.7554/eLife.62576.
63. Hulse, B.K., Haberkern, H., Franconville, R., Turner-Evans, D., Takemura, S.-y., Wolff, T., Noorman, M., Dreher, M., Dan, C., Parekh, R., et al. (2021). A connectome of the Drosophila central complex reveals network motifs suitable for flexible navigation and context-dependent action selection. eLife *10*, e66039. 10.7554/eLife.66039.
64. Bienkowski, R.S., Banerjee, A., Rounds, J.C., Rha, J., Omotade, O.F., Gross, C., Morris, K.J., Leung, S.W., Pak, C., Jones, S.K., et al. (2017). The Conserved, Disease-Associated RNA Binding Protein dNab2 Interacts with the Fragile X Protein Ortholog in Drosophila Neurons. Cell Rep *20*, 1372-1384. 10.1016/j.celrep.2017.07.038.
65. Lancaster, C.L., Yalamanchili, P.S., Goldy, J.N., Leung, S.W., Corbett, A.H., and Moberg, K.H. (2024). The RNA-binding protein Nab2 regulates levels of the RhoGEF Trio to govern axon and dendrite morphology. Molecular Biology of the Cell *35*, ar109. 10.1091/mbc.E24-04-0150.
66. Perkins, L.A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K., Yang-Zhou, D., Flockhart, I., Binari, R., Shim, H.S., et al. (2015). The Transgenic RNAi Project at Harvard Medical School: Resources and Validation. Genetics *201*, 843-852. 10.1534/genetics.115.180208.
67. Xu, R., Deng, K., Zhu, Y., Wu, Y., Ren, J., Wan, M., Zhao, S., Wu, X., Han, M., Zhuang, Y., and Xu, T. (2008). A large-scale functional approach to uncover human genes and pathways in Drosophila. Cell Research *18*, 1114-1127. 10.1038/cr.2008.295.
68. Lee, Y.S., Nakahara, K., Pham, J.W., Kim, K., He, Z., Sontheimer, E.J., and Carthew, R.W. (2004). Distinct Roles for Drosophila Dicer-1 and Dicer-2 in the siRNA/miRNA Silencing Pathways. Cell *117*, 69-81. 10.1016/S0092-8674(04)00261-2.
69. Aso, Y., Grübel, K., Busch, S., Friedrich, A.B., Siwanowicz, I., and Tanimoto, H. (2009). The Mushroom Body of Adult Drosophila Characterized by GAL4 Drivers. Journal of Neurogenetics *23*, 156-172. 10.1080/01677060802471718.

70. Zanet, J., Stramer, B., Millard, T., Martin, P., Payre, F.o., and Plaza, S. (2009). Fascin is required for blood cell migration during Drosophilaembryogenesis. Development *136*, 2557-2565. 10.1242/dev.036517.

71. Adams, J.C. (2004). Roles of fascin in cell adhesion and motility. Current opinion in cell biology *16*, 590-596. 10.1016/j.ceb.2004.07.009.

72. Jayo, A., and Parsons, M. (2010). Fascin: a key regulator of cytoskeletal dynamics. The international journal of biochemistry & cell biology *42*, 1614-1617. 10.1016/j.biocel.2010.06.019.

73. Kureishy, N., Sapountzi, V., Prag, S., Anilkumar, N., and Adams, J.C. (2002). Fascins, and their roles in cell structure and function. Bioessays *24*, 350-361. 10.1002/bies.10070.

74. Li, A., Dawson, J.C., Forero-Vargas, M., Spence, H.J., Yu, X., König, I., Anderson, K., and Machesky, L.M. (2010). The actin-bundling protein fascin stabilizes actin in invadopodia and potentiates protrusive invasion. Current biology : CB *20*, 339-345. 10.1016/j.cub.2009.12.035.

75. Cohan, C.S., Welnhofer, E.A., Zhao, L., Matsumura, F., and Yamashiro, S. (2001). Role of the actin bundling protein fascin in growth cone morphogenesis: localization in filopodia and lamellipodia. Cell motility and the cytoskeleton *48*, 109-120. 10.1002/1097-0169(200102)48:2<109::Aid-cm1002>3.0.Co;2-g.

76. Hourcade, B., Muenz, T.S., Sandoz, J.C., Rössler, W., and Devaud, J.M. (2010). Long-term memory leads to synaptic reorganization in the mushroom bodies: a memory trace in the insect brain? J Neurosci *30*, 6461-6465. 10.1523/jneurosci.0841-10.2010.

77. Bradke, F., and Dotti, C.G. (2000). Establishment of neuronal polarity: lessons from cultured hippocampal neurons. Current Opinion in Neurobiology *10*, 574-581. https://doi.org/10.1016/S0959-4388(00)00124-0.

78. He, Y., Ren, Y., Wu, B., Decourt, B., Lee, A.C., Taylor, A., and Suter, D.M. (2015). Src and cortactin promote lamellipodia protrusion and filopodia formation and stability in growth cones. Mol Biol Cell *26*, 3229-3244. 10.1091/mbc.E15-03-0142.

79. Hou, X., Nozumi, M., Nakamura, H., Igarashi, M., and Sugiyama, S. (2021). Coactosin Promotes F-Actin Protrusion in Growth Cones Under Cofilin-Related Signaling Pathway. Frontiers in Cell and Developmental Biology *9*. 10.3389/fcell.2021.660349.

80. Gibson, D.A., and Ma, L. (2011). Developmental regulation of axon branching in the vertebrate nervous system. Development *138*, 183-195. 10.1242/dev.046441.

81. Bodakuntla, S., Nedozralova, H., Basnet, N., and Mizuno, N. (2021). Cytoskeleton and Membrane Organization at Axon Branches. Front Cell Dev Biol *9*, 707486. 10.3389/fcell.2021.707486.

82. Wit, C.B., and Hiesinger, P.R. (2023). Neuronal filopodia: From stochastic dynamics to robustness of brain morphogenesis. Seminars in Cell & Developmental Biology *133*, 10-19. https://doi.org/10.1016/j.semcdb.2022.03.038.

83. Chung, V., Tsai, F., Chen, W., Von Hoff, D.D., Garmey, E.G., Zhang, J.J., and Huang, X.Y. (2022). NP-G2-044, a First-in-Class Fascin Inhibitor, Inhibits Growth and Metastasis of Gynecologic Cancers. European Journal of Cancer *174*, S34. 10.1016/S0959-8049(22)00891-7.

84.   Huang, J., Dey, R., Wang, Y., Jakoncic, J., Kurinov, I., and Huang, X.-Y. (2018). Structural Insights into the Induced-fit Inhibition of Fascin by a Small-Molecule Inhibitor. Journal of Molecular Biology *430*, 1324-1335. https://doi.org/10.1016/j.jmb.2018.03.009.

85.   Pan, J., Wang, L., Xu, L.-Y., Li, E.-M., and Dong, G. (2024). Prediction of The Binding Mode of NP-G2-044 Targeting Fascin and Its Drug Resistance Mutations. Advanced Theory and Simulations *7*, 2300427. https://doi.org/10.1002/adts.202300427.

86.   Mohr, O.L. (1922). Cases of mimic mutations and secondary mutations in the X-chromosome of Drosophila melanogaster. Zeitschrift für Induktive Abstammungs- und Vererbungslehre *28*, 1-22.

87.   Murthy, M., Fiete, I., and Laurent, G. (2008). Testing odor response stereotypy in the Drosophila mushroom body. Neuron *59*, 1009-1023. 10.1016/j.neuron.2008.07.040.

88.   Zwarts, L., Goossens, T., Clements, J., Kang, Y.Y., and Callaerts, P. (2016). Axon Branch-Specific Semaphorin-1a Signaling in Drosophila Mushroom Body Development. Front Cell Neurosci *10*, 210. 10.3389/fncel.2016.00210.

89.   Bashaw, G.J., Kidd, T., Murray, D., Pawson, T., and Goodman, C.S. (2000). Repulsive axon guidance: Abelson and Enabled play opposing roles downstream of the roundabout receptor. Cell *101*, 703-715. 10.1016/s0092-8674(00)80883-1.

90.   Kidd, T., Bland, K.S., and Goodman, C.S. (1999). Slit Is the Midline Repellent for the Robo Receptor in Drosophila. Cell *96*, 785-794. https://doi.org/10.1016/S0092-8674(00)80589-9.

91.   McConnell, R.E., Edward van Veen, J., Vidaki, M., Kwiatkowski, A.V., Meyer, A.S., and Gertler, F.B. (2016). A requirement for filopodia extension toward Slit during Robo-mediated axon repulsion. Journal of Cell Biology *213*, 261-274. 10.1083/jcb.201509062.

92.   Astin, J.W., Batson, J., Kadir, S., Charlet, J., Persad, R.A., Gillatt, D., Oxley, J.D., and Nobes, C.D. (2010). Competition amongst Eph receptors regulates contact inhibition of locomotion and invasiveness in prostate cancer cells. Nature Cell Biology *12*, 1194-1204. 10.1038/ncb2122.

93.   Wu, J.S., and Luo, L. (2006). A protocol for mosaic analysis with a repressible cell marker (MARCM) in Drosophila. Nature Protocols *1*, 2583-2589. 10.1038/nprot.2006.320.

94.   Drescher, U., Kremoser, C., Handwerker, C., Löschinger, J., Noda, M., and Bonhoeffer, F. (1995). In vitro guidance of retinal ganglion cell axons by RAGS, a 25 kDa tectal protein related to ligands for Eph receptor tyrosine kinases. Cell *82*, 359-370.

95.   Cheng, H.-J., Nakamoto, M., Bergemann, A.D., and Flanagan, J.G. (1995). Complementary gradients in expression and binding of ELF-1 and Mek4 in development of the topographic retinotectal projection map. Cell *82*, 371-381.

96.   Tamagnone, L., Artigiani, S., Chen, H., He, Z., Ming, G.-l., Song, H.-j., Chedotal, A., Winberg, M.L., Goodman, C.S., and Poo, M.-m. (1999). Plexins are a large family of receptors for transmembrane, secreted, and GPI-anchored semaphorins in vertebrates. Cell *99*, 71-80.

97.   Kolodkin, A.L., Matthes, D.J., O'Connor, T.P., Patel, N.H., Admon, A., Bentley, D., and Goodman, C.S. (1992). Fasciclin IV: sequence, expression, and function during growth cone guidance in the grasshopper embryo. Neuron *9*, 831-845.

98.   Battye, R., Stevens, A., and Jacobs, J.R. (1999). Axon repulsion from the midline of the Drosophila CNS requires slit function. Development *126*, 2475-2481.

99.   Kidd, T., Brose, K., Mitchell, K.J., Fetter, R.D., Tessier-Lavigne, M., Goodman, C.S., and Tear, G. (1998). Roundabout controls axon crossing of the CNS midline and defines a novel subfamily of evolutionarily conserved guidance receptors. Cell *92*, 205-215.

100.   Seeger, M., Tear, G., Ferres-Marco, D., and Goodman, C.S. (1993). Mutations affecting growth cone guidance in Drosophila: genes necessary for guidance toward or away from the midline. Neuron *10*, 409-426.

101.   Hedgecock, E.M., Culotti, J.G., and Hall, D.H. (1990). The unc-5, unc-6, and unc-40 genes guide circumferential migrations of pioneer axons and mesodermal cells on the epidermis in C. elegans. Neuron *4*, 61-85.

102.   Culotti, J.G., and Merz, D.C. (1998). DCC and netrins. Current opinion in cell biology *10*, 609-613.

103.   Zheng, J.Q., Wan, J., and Poo, M. (1996). Essential role of filopodia in chemotropic turning of nerve growth cone induced by a glutamate gradient. Journal of Neuroscience *16*, 1140-1149.

104.   Bentley, D., and Toroian-Raymond, A. (1986). Disoriented pathfinding by pioneer neurone growth cones deprived of filopodia by cytochalasin treatment. Nature *323*, 712-715.

105.   Chien, C.-B., Rosenthal, D.E., Harris, W.A., and Holt, C.E. (1993). Navigational errors made by growth cones without filopodia in the embryonic Xenopus brain. Neuron *11*, 237-251.

106.   Zanet, J., Jayo, A., Plaza, S., Millard, T., Parsons, M., and Stramer, B. (2012). Fascin promotes filopodia formation independent of its role in actin bundling. Journal of Cell Biology *197*, 477-486.

107.   Lin, S., Lu, S., Mulaj, M., Fang, B., Keeley, T., Wan, L., Hao, J., Muschol, M., Sun, J., and Yang, S. (2016). Monoubiquitination inhibits the actin bundling activity of fascin. Journal of Biological Chemistry *291*, 27323-27333.

108.   Liu, W., Sato, A., Khadka, D., Bharti, R., Diaz, H., Runnels, L.W., and Habas, R. (2008). Mechanism of activation of the Formin protein Daam1. Proceedings of the National Academy of Sciences *105*, 210-215. doi:10.1073/pnas.0707277105.

109.   Gombos, R., Migh, E., Antal, O., Mukherjee, A., Jenny, A., and Mihály, J. (2015). The Formin DAAM Functions as Molecular Effector of the Planar Cell Polarity Pathway during Axonal Development in Drosophila. J Neurosci *35*, 10154-10167. 10.1523/jneurosci.3708-14.2015.

110.   Szikora, S., Földi, I., Tóth, K., Migh, E., Vig, A., Bugyi, B., Maléth, J., Hegyi, P., Kaltenecker, P., Sanchez-Soriano, N., and Mihály, J. (2017). The formin DAAM is required for coordination of the actin and microtubule cytoskeleton in axonal growth cones. J Cell Sci *130*, 2506-2519. 10.1242/jcs.203455.

111.   Villari, G., Jayo, A., Zanet, J., Fitch, B., Serrels, B., Frame, M., Stramer, B.M., Goult, B.T., and Parsons, M. (2015). A direct interaction between fascin and

microtubules contributes to adhesion dynamics and cell migration. J Cell Sci *128*, 4601-4614. 10.1242/jcs.175760.

	Female sn^{28}	Female sn^{x2}	Male sn^{28}	Male sn^{x2}
Thin α-lobe	p < 0.0001****	p = 0.0054**	p = 0.0333*	p = 0.0012**
Split α-lobe	p = 0.0939	p < 0.0001****	p = 0.0056**	p < 0.0001****
Multiple α-lobe defects	p = 0.1479	p < 0.0001****	p = 0.4392	p = 0.0012**
Missing α-lobe	p = 0.1171	p < 0.0001****	p = 0.0003***	p < 0.0001****
Thin β-lobe	p = 0.0195*	p = 0.0011**	p = 0.6071	p < 0.0001****
Split β-lobe	p = 0.0022**	p < 0.0001****	p < 0.0001****	p < 0.0001****
Multiple β-lobe defects	p = 0.4144	p = 0.0489*	p = 0.4392	p = 0.0005***
Missing β-lobe	p = 0.2469	p = 0.0227*	p = 0.0175*	p = 0.0886

Table 1: Statistical analysis of α- and β- lobe defects in singed null flies. Chi square analysis was used to calculate p-values. sn^{28} flies were compared to control w^{1118} flies and sn^{x2} flies were compared to Oregon-R flies. Female w^{1118} n=32, female sn^{28} n=43, male w^{1118} n=32, and male sn^{28} n=42. Oregon-R n=54, female sn^{x2} n=43, male Oregon-R n=49, and male sn^{x2} n=34.

Hardin et al "Fascin in neuronal development"

Figures (1-9)

Figure 1: Fascin is concentrated in growth cone filopodia and collateral filopodia. (A) A DIV4 primary rat cortical neuron was methanol fixed and stained for fascin and β-actin. The black arrows indicate fascin1 enrichment at growth cone filopodia while the asterisks indicate fascin1 enrichment at collateral filopodial protrusions. Color panels on the right show the enlarged areas with fascin in green and β-actin in red. (B) enlarged images of a growth cone from panel A show the presence of fascin in lamellipodia in addition to filopodia. (C) GFP-fascin localizes to growth cones and is mostly associated with filopodia. 5 minutes after treatment with 12-O-tetradecanoylphorbol-13-acetate (TPA), a protein kinase C (PKC) activator, GFP-fascin is seen coming off filopodia, followed by filopodia retraction and resulting in growth cone collapse.

Figure 2. Inhibition of fascin1 actin bundling results in shift in localization and reduction in growth cone filopodia. (A) CAD cells were treated for 240 minutes with either NP-G2-044 or DMSO and methanol-fixed. A decrease in fascin signal at the edge of the NP-G2 treated cells and an increase in cytosolic signal is observed while β-actin signal does not undergo this shift in localization, as shown by ratio comparison. (B) Quantification of fascin and β-actin cell edge and center intensities shows a decrease in fascin localization to filopodia after application of NP-G2-044 for both 30 minutes (n=58, p= 0.0368) and 240 minutes (n=70, p<0.0001) when compared to cells treated with DMSO (n=53 and n=69). β-actin signal does not undergo the same cytosolic shift. Analysis conducted via One-way ANOVA. (C) Representative images of growth cone filopodia for control and NP-G2-044 treated neurons. Fascin is shown in cyan and β-actin is shown in magenta. A decrease in number of filopodia as well as length of filopodia is seen. (D) Quantification of filopodia defects observed in fascin1-inhibited growth cones (n= 345, p<0.0001) and DMSO control growth cones (n=328, p<0.0001). Statistical significance calculated via unpaired t-test.

Figure 3. Pharmacological inhibition of fascin1 results in axon growth and branching defects.
(A) Representative images of low-density primary rat hippocampal neurons PFA-fixed on DIV 5 and
stained with anti-tubulin antibody. Neurons treated with 15 µM of NP-G2-044 for 4 days show
truncated axon growth and decreased branch points compared with DMSO-treated cells. (B)
Tracings of representative DMSO control and NP-G2-044 treated neurons, obtained via Simple
Neurite Tracer for FIJI. Cell bodies are in green, primary axons are shown in blue, secondary in
magenta, and tertiary in black. Neurons treated with 15 µM NP-G2-044 show less axonal outgrowth
and elaboration. (C) There are significant decreases in primary (p=0.0338), secondary (p<0.001),
and tertiary (p=0.0024) axon length for the 15 µM NP-G2-044-treated group (n=160) compared to
DMSO control neurons (n=134). Total secondary and tertiary axon lengths are shown. (D)
Quantification of the number of secondary and tertiary branch points shows a significant decrease
for neurons treated with 15 µM of NP-G2-044 (p<0.0001) Statistical analyses done via one-way
ANOVA with Šídák's multiple comparisons test.

Figure 4. SN is expressed in a subset of MB α- and β-lobes in adult _Drosophila_ brains. (A) Brains from flies expressing pan-neuronal UAS-myristoylated GFP via the _nSyb-GAL4_ driver were dissected and stained with anti-GFP and anti-Singed antibodies. We observe that SN staining appears to be enriched in the MB α- and β-lobe axonal tracts, indicated by magenta arrows. (B) Expression of UAS-myristoylated GFP via the 238Y GAL4 driver labels neurons in the MB born post-pupae formation and shows overlap with SN signal. GFP is labelled as cyan, and Singed is labelled as magenta. Axonal tracts labelled by SN staining appear to be centered in the MB axon bundles, as shown in the bottom cross sections. Yellow lines indicate location of cross section along the x-axis.

Figure 5. Loss of fascin1 results in axonal mushroom body defects. (A) Schematic of the Drosophila brain and the structure of the lobes making up the mushroom body (MB). The MB lobes are formed by projection of axons (dotted lines) from Kenyon cell bodies (spheres) which undergo bifurcation as well as guidance responses to form the anterior α- and medial β-lobes. (B) Brains were dissected from control (w^{1118}) and SN null (sn^{28}) flies and stained for Fasciculin II, a marker of MB axons. The SN null flies display significant defects in both α- and β lobes, as well as midline crossing errors (brain midline indicated by dashed black lines). Female w^{1118} brain, displaying normal α- and β-lobes and no midline crossing of the β-lobes and female sn^{28} brain showing a missing α-lobe, thin α-lobe (black asterisk) and split β-lobe (black star). Male ^{w1118} and male sn^{28} brains, showing similar defects to the female brains. (C) Quantification of α- and β- lobe defects. Chi square analysis was used to calculate p-values, shown in these graphs is the analysis of total defects (**** indicates a p-value of <0.0001). For a more detailed comparison of defects, see chart. (D) Schematics of α- and β- lobe defects that we observe in SN null flies. E) Schematics of the midline defects observed in SN null flies. F) Quantification of midline crossing defects in female w^{1118} (n=32), female sn^{28} (n=43), male w^{1118} (n=32), and male sn^{28} (n=42).Chi square analysis was used for statistical testing (p-value <0.0001).

Figure 6. Fluorescent labeling of a subset of mushroom body axons reveals axon guidance deficiencies. Wildtype and sn[28] flies expressing a MB-specific split GAL4 driver were crossed with wildtype or sn[28] flies expressing UAS-Myristoylated GFP. Brains were dissected and stained for GFP and Fas II. (A) Wildtype MB axons project from Kenyon cells and undergo bifurcation to form two separate tracts (α- and β-) which then undergo axon guidance to form the vertical α- and medial β-lobes. (B) sn[28] MB axons undergo bifurcation but then fail to follow midline guidance signals, as shown by β-lobes crossing the midline (dotted blue line). To the right, enlargement of the region outlined by the orange dotted box. We see the β-lobes crossing over the midline (orange arrows) and the split appearance of the β-lobe (yellow arrows). (C) Example of sn[28] MB axons failing to properly follow guidance cues. White arrow shows a β-lobe turning back on itself, shown enlarged in panel on the right. The orange arrow points to a β-lobe that is starting to cross the midline (dashed blue line) and is also turning upwards at the midline, shown enlarged to the right. A schematic of the β-lobe trajectories is also shown. (D) Quantification of α- and β- defects (female control n= 23, female sn[28] n=12, male control n=25, male sn[28]=11), analyzed via Chi square analysis (p= 0.0004, p= 0.0007, p= 0.015, p=0.0007). E) Quantification of midline defects, analyzed via Chi square (p= 0.0015, p=0.0048).

Figure 7. Singed null flies have a delayed climbing response. (A) Climbing behavior was evaluated using the negative geotaxis assay, flies are tamped to the bottom of a vial and results were quantified by dividing the vial into 10 bins (shown on the side of the image) and counting the number of flies in each bin 10 seconds after climbing begins Representative tracing of female w^{1118} and sn^{28} flies 5 seconds after beginning to climb. (B) Comparison of the climbing indices of w^{1118} (n=52) and sn^{28} (n=30) flies shows a climbing deficit for the sn^{28} flies, with the singed null flies taking longer to reach the top of the vial (analyzed using two-way ANOVA, p<0.0001). (C-D) w^{1118} male (n=48) and male sn^{28} (n= 33) flies show the same deficit seen in the female flies (analyzed using two-way ANOVA, p<0.0001).

Figure 8. Mushroom body-specific knockdown of Fascin results in defects. (A) Quantification of Singed knockdown efficiency via western blot shows a decrease in SN expression but does not completely eliminate the presence of SN in heads collected from flies expressing a UAS-snRNAi construct as well as a UAS-Dicer2 construct ubiquitously expressed by an Actin GAL4 driver (n=3). (B) Flies expressing Act5-GAL4, UAS-Dicer2, and UAS-snRNAi display the gnarled bristle defect that is associated with loss of SN. (C) Representative images of control (UAS-Dicer2/CyO;UAS-snRNA/TM2) and snRNAi (UAS-Dicer2/+;UAS-snRNAi/+;OK107-GAL4 /+) brains stained for fascicullin II. (D) Quantification of α- and β- lobe defects in female control (n=29), female MB KD (n=41), male control (n=25), and male MB KD (n=32). Midline (indicated by dashed black line) defects were not observed in the snRNAi brains. Chi square analysis was performed for comparisons, p values were as follows: p=0.0001,p =0.0261, p= 0.001, p= 0.003.

Fig 9. MB-specific expression of fascin1 is sufficient to rescue both climbing and MB lobe defects in females. (A) Female flies from 4 genotypes, w[1118] (Control), sn[28] (SN28), sn[28];238Y-GAL4/UAS-Myristoylated GFP (SN28 + M-GFP), and sn[28];238Y-GAL4/UAS-Singed GFP (SN + SN-GFP) were collected and assessed using the NGA protocol. Flies expressing UAS-GFP-Singed driven by 238Y-GAL4 are able to restore climbing ability to control w[1118] levels (analyzed using two-way ANOVA, p<0.0001). (B) Representative images of MB α- and β- lobes in the four genotypes examined. (C) Quantification of MB α- and β- lobe defects shows that there is a complete rescue of MB lobe defects observed in sn[28] flies upon MB-specific expression of WT GFP-Fascin1. Statistical analysis was performed using Chi-square proportional analysis with post-hoc Marascuilo's Procedure used, all experimental groups were compared to w[1118] control. Female w[1118] n=37, Female sn[28] n=40 (β-lobe p-value <0.0001, α-lobe p-value <0.0001), Female sn[28]; 238Y-GAL4/UAS Myristoylated-GFP n=36 (β-lobe p-value = 0.0465 , α-lobe p-value = 0.0004), female sn[28]; 238Y-GAL4/UAS Singed-GFP n=31 (β-lobe p-value = 0.9763 , α-lobe p-value = 0.6298). (D) MB-specific expression of UAS Singed-GFP is sufficient to restore midline axon projection morphology. Statistical analysis was performed using Chi-square proportional analysis with post-hoc Marascuilo's Procedure used. Female w[1118] n=37, female sn[28] n=40 (p-value <0.0001), female sn[28]; 238Y-GAL4/UAS Myristoylated-GFP n=36 (p-value <0.0001), female sn[28]; 238Y-GAL4/UAS Singed-GFP n=31 (p-value = 0.61).

Supplemental figures (1-4)

Figure S1: Fascin is enriched in actin filament bundles in CAD cells. (A) fascin signal is concentrated in thick F-actin bundles such as filopodia and microspikes. (B) Fascin signal is also enriched in lamellipodia. Bue arrows: thick F-actin bundles in filopodia and microspikes. Red arrowheads: Lamellipodia/branched F-actin network.

Supplemental figure S2: sn²⁸ brains do not show any differences in PDF neuron projection compared to control w¹¹¹⁸ brains. (A) Confirmation of singed knockout in sn²⁸ line via western blot (n=1). Quantification of western blot via densitometry shows a reduction of SN in the heterozygous line and only background levels in the null line. (B) Brains dissected from sn²⁸ flies and stained with anti-Singed antibody do not show any signal, further confirming that these flies are not producing SN protein. (C) Brains from female and male sn²⁸ and w¹¹¹⁸ flies were dissected and stained with an anti-PDF antibody. When imaged, we do not observe any differences in PDF axon projection.

Supplemental figure S3: MB defects and NGA climbing deficits are also observed in another SN null line. (A) Representative images of Fas II staining in Control female Oregon R and female sn^{x2} brains showing split α-lobe, thin α-lobe, split β-lobe, and a thin β-lobe (black arrowhead). Male Oregon R and male sn^{x2} brains also show similar defects to the female brains. female Oregon-R (n=54), female sn^{x2} (n=43), male Oregon-R (n=49), and male sn^{x2} (n=34). (B) Representative tracings of female Oregon-R and sn^{x2} flies 5 seconds after beginning to climb. Comparison of the climbing indices of female Oregon-R(n=54) and female sn^{x2} (n=40) flies shows a climbing deficit for the sn^{x2} flies, with the SN null flies taking longer to reach the top of the vial (analyzed using two-way ANOVA, p<0.0001). The same delay in reaching the top bin position is observed in male sn^{x2} (n=33) flies when compared to male Oregon-R (n=55), p<0.0001. (C) Quantification of α- and β- lobe defects. Chi square analysis was used to calculate p-values, shown in these graphs is the analysis of total defects (**** indicates a p-value of <0.0001). (D) Quantification of midline defects observed in sn^{x2} brains (analyzed using Chi-square analysis, p<0.0001).

Supplemental figure S4. MB-specific expression of fascin1 is sufficient to rescue climbing and midline defects but not MB β-lobe and α-lobe defects in males. (A) Flies from 4 genotypes were collected and assessed using the NGA protocol. Flies expressing UAS-SN-GFP driven by 238Y-GAL4 are able to restore climbing ability to control w^{1118} levels (analyzed using two-way ANOVA, p=0.02. (B) MB-specific expression of UAS SN-GFP is sufficient to restore midline axon projection morphology in male flies. Statistical analysis was performed using Chi-square proportional analysis with post-hoc Marascuilo's Procedure used. Male w^{1118} n=32, male sn^{28} n=37 (p-value =0.0012), male sn^{28}; 238Y-GAL4/UAS Myrs-GFP n=27 (p-value <0.0001), male sn^{28}; 238Y-GAL4/UAS SN-GFP n=36 (p-value = 0.3255). (C) Expression of SN-GFP in the MB is unable restore MB lobe morphology. Brains were dissected and stained for Fas II. Quantification of MB α- and β- lobe defects shows that there is not a rescue of MB lobe defects observed in sn^{28} flies upon MB-specific expression of WT GFP-Fascin1. Statistical analysis was performed using Chi-square proportional analysis with post-hoc Marascuilo's Procedure used. Male w^{1118} n=32, male sn^{28} n=37 (β-lobe p-value <0.0001, α-lobe p-value <0.0001), male sn^{28}; 238Y-GAL4/UAS Myrs-GFP n=27 (β-lobe p-value = 0.0099, α-lobe p-value <0.0001), male sn^{28}; 238Y-GAL4/UAS SN-GFP n=36 (β-lobe p-value = 0.0009, α-lobe p-value <0.0001). (D) Representative images of MB α- and β- lobes in the four genotypes examined.

Supplemental movies (1-5)

Movie #S1. Time-lapse video showing GFP-Fascin in the growth cone of a rat primary hippocampal neuron in culture. Playback speed: X150.

Movie #S2. **Wildtype mushroom body axons bifurcate and project vertically and medially**. Brains from control flies (Split-GAL4/UAS-myristoylated GFP; Split-GAL4/+) were dissected, PFA-fixed, and stained with anti-GFP and anti-Singed antibodies. 2-photon imaging generates a 3-D image of the axon projection from Kenyon cells and shows axons undergoing bifurcation into α and β lobe tracts which then follow guidance cues to form the α and β lobes.

Movie #S3. **sn[28] mushroom body axons bifurcate but appear to have guidance defects.** Brains from SN null flies expressing Split-GAL4/UAS-myristoylated GFP; Split-GAL4/+ were dissected, PFA-fixed, and stained with anti-GFP and anti-Singed antibodies. 2-photon imaging reveals that the axon tract extending from the Kenyon cells bifurcates but the α and β lobe tracts then do not follow proper guidance pathways. We observe midline crossing of the β lobes and additionally see the appearance of a split β lobe.

Movie #S4. **sn[28] mushroom body axons bifurcate but appear to have midline guidance defects.** As previously described, brains from SN null flies expressing Split-GAL4/UAS-myristoylated GFP; Split-GAL4/+ were dissected, PFA-fixed, and stained with anti-GFP and anti-Singed antibodies. When imaged, we see in this case a β lobe that appears to initially project medially but does not reach the midline and instead turns around. We also observe a β lobe that does project to the midline, but some axons cross over the midline while the rest begin to turn vertically.

Movie #S5. **sn[28] flies show a locomotor defect.** Female w[1118] flies are shown on the left and sn[28] flies are on the right. When the flies are tamped to the bottom of the vial, the SN null flies show a delay in reaching the top when compared to control flies.