Hyperreflexia after corticospinal tract lesion reflects 1A afferent circuit changes not increased KCC2 hyperexcitability

Thelma Bethea[1,2], Temitope Adegbenro[1], and John H. Martin[1,2]

[1]Department of Molecular, Cellular, and Biomedical Sciences, Center for Discovery and Innovation, City University of New York School of Medicine, New York, NY, USA; [2]Neuroscience Program, Graduate Center of the City University of New York, New York, NY, USA

Abbreviated title: Hyperreflexia after unilateral corticospinal tract lesion

Main text figures: 6
Supplemental figures: 5

Correspondence should be addressed to:
John H. Martin, Ph.D.
Department of Molecular, Cellular, and Biomedical Sciences
City University of New York School of Medicine & Center for
Discovery and Innovation
160 Convent Avenue
New York, NY 10031
Tel: 212.650.5956
E-Mail: jmartin@med.cuny.edu

Funding: This work was supported by the NIH (JHM: 2R01NS064004); New York State Spinal Cord Injury Board (JHM: C37716GG; JHM, TB: C39060GG) .

Acknowledgements: We thank XiuLi Wu for histology.

The authors have no competing interests to declare.

Abstract

Hyperreflexia, spasticity, and a loss of skilled motor function are consequences of spinal cord injury (SCI). Multiple potential mechanisms underpinning hyperreflexia have been reported, including: aberrant proprioceptive afferent (PA) sprouting, which could enhance reflex signaling; reduced GABAergic inhibitory presynaptic regulation of 1A terminals (GABApre); and increased excitability produced by reduced motor neuron membrane-bound potassium-chloride co-transporter #2 (KCC2), which has only studied after SCI and bilateral CST lesion. Here we examine how selective CST injury allows for specific investigation of the different mechanisms to determine their contributions to hyperreflexia. Hoffmann (H)-reflex rate-dependent depression (RDD) of the forelimb and hindlimb 5th-digit abductor muscles was used to assess hyperreflexia. We compared RDD in naïve and unilateral-PTX animals at 7-dpi and 42-dpi, supplemented with additional timed tests to examine the time-course of hyperreflexia development. Immunohistochemistry was used to identify PA synapses (VGlut1) and GABApre (GABApre), motor neurons (ChAT), and KCC2. Following unilateral PTX, which eliminates the CST from one hemisphere, we observed significant hyperreflexia in the contralesional forelimb only. Membrane-bound KCC2 was unchanged in contralesional cervical motor neurons. Whereas both cervical and lumbar motor neurons showed increased PA sprouting contralesionally, there was a concomitant increase in GABApre terminals for the lumbar not cervical cord. Our findings show that KCC2 is disassociated from hyperreflexia in the uniPTX model. Instead, forelimb hyperreflexia can be explained by cervical motor neuron PA sprouting and an uncompensated GABApre regulation.

Introduction

Hyperreflexia after motor cortex stroke or spinal cord injury (SCI) contributes to significant disease burden in patients suffering a reduction, or loss of skilled volitional movement, and spasticity(Hiersemenzel, Curt, & Dietz, 2000),(Thompson, Reier, Lucas, & Parmer, 1992), (Hultborn, 2003). Although much is known about the mechanistic underpinnings of hyperreflexia—with animal studies stressing a reduction in the potassium-chloride co-transporter 2 (KCC2), sprouting of 1A afferent fiber terminals on motor neurons, and reduced

GABApre inhibition of 1A presynaptic terminals on motor neurons—key questions about the differential contributions of these, and additional, mechanisms have not been addressed.

KCC2 is a membrane-bound protein that maintains low intracellular [Cl$^-$] with extrusion of K$^+$ (Lee et al., 2007) . In the developing spinal cord, increased KCC2 expression is associated with GABA-A and Glycine receptor mediated hyperpolarization (Boulenguez et al., 2010), (Hübner et al., 2001). SCI produces a reduction in the membrane-bound KCC2, thereby driving up motor neuron excitability (Chamma, Chevy, Poncer, & Levi, 2012), contributing to hyperreflexia (Boulenguez et al., 2010); (Cˆote, 2021);(Beverungen, 2020). SCI damages many sensory and motor projection systems as well as segmental afferent connections and intrinsic interneurons and circuits (Oudega, 2012). Earlier we showed that unilateral selective loss of the corticospinal tract (CST) from one hemisphere, by pyramidal tract lesion (PTX) in the caudal medulla, is sufficient to produce contralesional hyperreflexia. We further showed that this led to 1A afferent fiber sprouting on the affected side of the cervical spinal cord. This increase in 1A synapses on motor neurons, which are VGlut1-positive, could contribute to hyperreflexia through enhanced 1A signaling. Subsequently, we showed that a CST lesion or motor cortex inactivation increased 1A afferent projections to the affected (contralateral) motor neurons, creating an imbalance in activity-dependent completion exerted by CST terminals on target neurons, leading to 1A fiber sprouting (Jiang, Zaaimi, & Martin, 2016). Bilateral PTX also produces 1A afferent sprouting and, like SCI, a reduction in membrane-bound KCC2 (Kathe, Hutson, McMahon, & Moon, 2016). That study also showed that there was a reduction in GABApre contacts on 1A motor neuron terminals. Taken together, these findings suggest three potential contributing mechanisms underlying hyperreflexia.

The present study aimed to distinguish the potentially complementary contributions to hyperreflexia of KCC2-mediated hyperexcitability and 1A circuit mechanisms (1A sprouting and associated GABApre inhibition). We used a unilateral PTX to cut all corticospinal projections from one motor cortex. These eliminates most CST axons in the contralateral spinal cord, sparing only 5% of the ipsilateral projections. Hyperreflexia was assessed using rate-dependent depression of the H-reflex, (RDD). We characterized RDD changes for the forelimb and hind limb during an early subacute period (7 days post-injury) and the chronic period (42 days post-injury). We compared RDD changes for the forelimbs and hindlimbs to explore possible distinctions. In the rodent, the CST has fewer ventral projections in the lumbar than cervical

enlargement (Moreno-Lopez, Bichara, Delbecq, Isope, & Cordero-Erausquin, 2021)) and this difference may be associated with important functional distinctions(Moreno-López, Olivares-Moreno, Cordero-Erausquin, & Rojas-Piloni, 2016). Clinically, lower limb spasticity is less prominent (Sunnerhagen, Opheim, & Alt Murphy, 2019) and there are reports of an adaptive association for the lower (Urban et al., 2010), but not so for the upper extremity (Lundström, Smits, Terént, & Borg, 2010). We assayed the 1A connections on motor neurons and their presynaptic inhibition (GABApre) at the two timepoints, using highly sensitive motor neuron 3-D reconstruction, and segmental KCC2.

After unilateral PTX, our findings show contralateral forelimb but not hindlimb hyperreflexia. Although sprouting of 1A afferent terminals on motor neurons occurs in both the contralateral cervical and lumbar enlargements, GABApre sprouting commensurate with the 1A sprouting only occurs at the lumbar level, where hyperreflexia did not occur. Surprisingly, membrane-bound KCC2 did not change in either the cervical or lumbar enlargements, suggesting a 1A circuit-based, not intrinsic motor neuronal excitability, mechanism for hyperreflexia. Thus, using a model brain lesion that axotomizes all CST axons from one motor cortex, we distinguish the contributions of three potential contributing mechanisms to hyperreflexia, showing that only loss of GABApre regulation explains hyperreflexia.

METHODS

Animals

Experiments were performed in accordance with the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals. All animal protocols were approved by the Advanced Science Research Center Institutional Animal Care and Use Committee. Adult male Sprague Dawley rats (250 –275 g) were used for this study. Animals were housed under a 12 h light/dark cycle in a pathogen-free area with water and food provided ad libitum.
Animals comprised three principal groups: naïve (uninjured), early post-injury (3-days-post-injury, dpi), chronic post-injury (42-dpi).

H-Reflex Recording

Animals were anesthetized with an induction dose of ketamine and xylazine (70mg/kg Ketamine; 5 mg/kg Xylazine, i.p.) and maintained on ketamine alone during the course of

physiological experiments (Hosoidoa, 2009); (Ho, 2002). Animals were monitored continuously throughout all procedures and supplemental doses of ketamine (25mg/kg) were provided as needed. To determine the rate-dependent depression (RDD) of the H-reflex, paired-pulse stimulation with control and test pulses were used at increasing inter-pulse intervals (IPIs) (20-2000 ms). We used percutaneous muscle stimulation and recorded the evoked the direct M (muscle) and H-reflex responses through the same electrodes. For the forelimb, we studied the abductor digiti quinti muscle (ADQ) and, for the hindlimb, the abductor digiti minimi muscle (ADM). We chose the percutaneous method because we wished to examine the time-course of development of hyperreflexia repeatedly in the same animal (see Supp Fig1). In a prior study involving the extensor carpi radialis (ECR) muscle, we found that this percutaneous method did not reliably distinguish the M-and H-waves. In initial experiments, we found that ADQ and ADM H-reflex recordings using this method produces reliable separation of the M- and H-waves, resulting in consistent RDD. Thus, we were able to use percutaneous stimulation/recording repeatedly and consistently with this method.

Retrograde labeling of motor neurons.

Assessment of KCC2 was made in extensor carpi radialis (ECR) and tibialis anterior (TA) motor neurons of the cervical and lumbar segments. We confirmed the identities and locations of the motor neurons by intramuscular injection of cholera toxin subunit B (CTB) (FITC conjugated, List Biological Laboratories) into the muscles, to retrogradely label the motor neurons. CTB injections were made 7 days prior to euthanizing the animals (3-dpi early time point group; 42-dpi later timepoint group), which is sufficient time for retrograde labeling of motor neurons and anterograde transport transganglionic labeling of 1A afferent terminals (Jiang et al., 2016). CTB (10 μm for each muscle, 1% solution) was injected into the target muscles bilaterally using a Hamilton syringe (Tan, Chakrabarty, Kimura, & Martin, 2012). A total of 40 μl of 1% CTB was injected into each rat.

Unilateral Pyramidotomy.

To perform the unilateral pyramidal tract injury, animals (n=12) were anesthetized with Ketamine-Xylazine (70mg/kg Ketamine; 5 mg/kg Xylazine, i.p.). Supplemental doses of Ketamine (25 mg/kg) were administered as needed. Similar to our earlier studies (Jiang et al., 2016) ; (Tan et al., 2012), an incision was made through the skin of the ventral neck. The tissues

were bluntly dissected, and the trachea was displaced laterally, to expose the ventral surface of the occipital bone. Using a rongeur, a small craniotomy was made to expose the ventral medulla. The dura was cut, using an iridectomy scissor (Fine Science Tools), to expose the pyramids. The scissor was used to cut the right pyramid, to a depth of 1.2 mm below the ventral medullary surface to completely transect the pyramid. Care was taken to not damage the basilar artery. Completeness of the lesion was confirmed postmortem with histological assessment of the loss of PKC-γ immunoreactivity in the dorsal CST, contralateral to the injury, at C3.

Tissue Processing and Immunohistochemistry

At the termination of experiments, rats were deeply anesthetized with a mixture of ketamine and xylazine (70/5 mg/kg, i.p.) and transcardially perfused with 250 ml of 0.1M phosphate buffer (PB) followed by 500ml of 4% paraformaldehyde solution. The brain and spinal cord were removed and postfixed in paraformaldehyde solution for 3hrs before being transferred to a 30% sucrose solution. Tissue was sectioned transversely at a thickness of 40 μms. To identify 1A terminals on motor neurons and their presynaptic contacts, sections were labelled with: 1) the vesicular glutamate transporter 1(rabbit anti-Vglut1)(Millipore 1:5000 conc), a marker for PA terminals (Alvarez, Villalba, Zerda, & Schneider, 2004), 2) glutamic acid decarboxylase 65 (chicken anti-GAD65)(Millipore, 1:1000 conc.), a marker of GABAergic presynaptic inhibition, and 3) CTB (List Biologicals goat anti-CTB 1:2000 conc.), to label motor neurons. Secondary antibodies used included: guinea pig-Cy5 for Vglut1(Millipore 1:5000 conc.), rabbit-Cy3 for GAD65 (Jackson Labs 1:1000 conc.), and goat-FITC for CTB (Invitrogen 1:2000). A separate set of tissue sections were stained for KCC2 (Millipore, rabbit-KCC2; 1:100 conc.), and co-stained with choline acetyltransferase ChAt (Millipore Goat-ChAT 1:100 conc.) that is used as a marker for motor neurons. Secondary antibodies used were: guinea pig-Cy5 for KCC2 (Millipore 1:5000 conc.) and goat-FITC against ChAT (Invitrogen 1:100).

1A afferent terminals Image Analysis

Sections within the cervical (C6-C8) and lumbar (L1-L4) spinal cord were visualized and imaged digitally using a Zeiss LSM 880 confocal microscope. Individual motor neurons as well as their proximal dendrites ipsilateral and contralateral to the injury were imaged as a z-stack at 63x magnification with a 2.23μm optical step size. Motor neurons (soma and dendrites) were 3D-rendered along with accompanying 1A proprioceptive afferent (PA) terminals and GABApre bouton numbers using Imaris Software (Oxford Instruments ver. 9.7.1). Quantitative analyses were

also performed using Imaris. We previously used ImageJ to count presynaptic 1A afferent synaptic puncta on motoneurons (Tan et al., 2012); (Sharif, Alexander, Azam, & Martin, 2021) . However, the counting of contacts in a single focal plane revealed the limitation of the 2-Danalysis. By comparison, Imaris allowed us to visualize all potential interactions by rendering all components from a z-stack confocal file into 3D rendered objects. This method allows us to accurately and consistently assess the extent to which presynaptic excitatory, as well as inhibitory inputs influence motoneuron excitability. We used a series of thresholding steps for each component starting with CTB-positive motor neurons, then Vglut1-positive synapses, then GAD65-positive synapses. Interactions between rendered motor neurons and Vglut1-positive synapses were defined by a separation distance of zero micrometers. Similarly, interactions between rendered Vglut1-positive synapses that contact the motor neuron, and rendered GAD65 synapses were defined by a separation distance of zero micrometers. Motor neurons were imaged and analyzed individually in the cervical (ipsi-lesional: n= 37 motoneurons; contralesional: n= 33 motoneurons) and lumbar (ipsilesional: n=20 motoneurons; contralesional: n= 20 motoneurons) segments for the early timepoint, and the cervical (ipsilesional: n= 53 motoneurons; contralesional: n= 50 motoneurons) and lumbar segments (ipsilesional: n= 39 motoneurons; contralesional: n= 46 motoneurons) for the late timepoint.

KCC2/ChAT Image Analysis

KCC2 in motor neurons (ChAT-positive) within cervical (C6-C8), and lumbar(L1-L4) segments were analyzed (ImageJ analysis software, ver. 1.53a).   Three lines were drawn cross-sectionally through each motor neuron to measure KCC2 labeling grayscale pixel intensity at the two membrane intersection points and in the cytosol. Supplemental Figure 1 shows a labeled motor neuron (A) and one representative line for constructing the average histogram (B). Histograms were generated for each line. Area under the curve for the membrane, and cytosolic regions were measured as integrated pixel intensities. The intensity for the membrane and cytosolic regions are averaged (Kathe et al., 2016), and a ratio of membrane bound over cytosolic bound KCC2 is determined (Boulenguez et al., 2010). The mean intensity of membrane bound and cytosolic bound KCC2 were separately determined by normalizing gray matter KCC2 labelled motor neurons, to white matter KCC2 labelling (Boulenguez et al., 2010). The following motor neurons were analyzed: cervical (naïve right=19, left=20; early ipsilesional: n=25; early contralesional: n=26; chronic ipsilesional: n=17; chronic contralesional n=19), and lumbar (naïve

right=24, left=22; early ipsilesional: n=21; early contralesional: n=23; chronic ipsilesional: n=19; chronic contralesional n=19).

Western blots

To validate immunohistochemical KCC2 measurements, we used western blotting. Animals were euthanized at 7 days post unilateral CST injury. The cervical (C4-C6) and lumbar (L3-L5) enlargements of the spinal cord (Naïve N=5;PTX N=3) were dissected, split along the midline to separate ipsilesional and contralesional segments, and placed into ice cold Hank's balanced salt solution (HBBS; Invitrogen). The segments were placed in 4 microtubes and homogenized promptly in N-PER buffer (Invitrogen) supplemented with a protease-inhibiting cocktail (Invitrogen). Approximately 50 μg of protein was subjected to SDS-acrylamide-bisacrylamide gel electrophoreses along with a protein ladder (Crystalgen) for size comparison. Following electrophoresis, the protein bands were transferred to PVDF membrane (Millipore) and probed against the primary antibodies. An ECL chemiluminescent system (Cell Signaling Technology) was used to develop the Western blots. The following primary antibodies were used: rabbit anti-KCC2 (Millipore) at 1:500; and mouse anti-GAPDH at1:5000 (Sigma Aldrich). Secondary antibodies used were: guinea pig for KCC2 (1:10 conc.; Millipore) and goat for GAPDH (1:1000 conc.;Vector Laboratories).

Statistics

Statistical analysis was performed using Graphpad (Prism Software, ver. 9.1.2). Unpaired Student's t-test and one-way ANOVA followed by post hoc corrections for multiple comparisons were used as needed. Results are displayed as mean ± SEM, and P-value. Analyses were performed by laboratory personnel blinded to the experimental protocol.

Results

Experimental design

We examined the physiological and morphological effects of a complete unilateral CST injury at the pyramid in the caudal medulla (PTX) at early (7-days post injury, dpi) and chronic (42-dpi) timepoints (Figure 1A). Unilateral PTX was verified to be complete in all animals by the absence of Pkc-γ staining at the C7 level (Fig 1B). H-reflex assessments (n=6 rats) were made prior to PTX (baseline) and at the early and chronic timepoints (Figure 1A). Additional timepoints were assessed (n=5 rats) to show the progression of hyperreflexia. H-reflex changes were assessed using the method of pair-pulse stimulation to measure rate-dependent depression (RDD) of the reflex (Fig 1C). Stimulation evokes the M-wave and the H-wave (Fig 1C). In a naïve animal (Fig 1C), the amplitude of the H-wave evoked by the second test pulse becomes reduced with short interstimulus intervals (ISI; Fig 1C, 2000 ms versus 20 ms). This is rate-dependent depression (RDD) of the H-reflex (Ho, 2002). Seven-days prior to euthanizing the animals, they were subjected to an intramuscular CTB injection into ECR and TA (see Methods). These muscles were chosen because their motor pools are large and extend throughout most of the cervical enlargement and lumbar segments (Romanes, 1964); (Hardman & Brown, 1985), making them good candidates for identifying a generalized motor neuronal response to CST loss. Additionally, ECR has been shown to display RDD after unilateral PTX (Tan et al., 2012). These animals were euthanized at 42-dpi, (chronic time point) and spinal tissue was collected for performing analyses to assess KCC2 and 1A terminals on motor neurons.

Figure 1. Experimental design. A. Time line of experiments. Early. Experiments were designed for collecting tissue samples only (CTB -labeled motor neurons, immunohistochemistry for KCC2, VGlut1, PKCγ and Gad65 (n=5 rats PTX; n=5controls/ shams). Chronic. Experiments designed for repetitive H-reflex testing (42-dpi) and collection of tissue samples for immunohistochemistry, as for the early cohort of rats (n=4). B. Unilateral PTX with PKCγ anatomical validation of lesion completeness. C. Rate dependent depression of the H-reflex. Top part of figure shows experimental setup with single electrode pair in the muscle to electrically activate afferent fibers and record the H- and M-responses.

As indicated in the timeline, a separate set of animals we examined at 7-dpi (early time point; n=5) to assess morphology (KCC2 and 1A terminals).

Unilateral PTX produces forelimb, not hind limb, hyperreflexia

We first determined if there were differential effects of a complete unilateral CST injury on RDD of the H-reflex of the fore-and hind limbs. RDD plots (Figure 2A, B) show that the test H-wave changes as a function of the preceding conditioning pulse for the different ISIs. The plots for both ipsilesional limbs show the characteristic H-wave reduction with progressively shorter ISIs and robust negative slopes. By contrast, the contralesional forelimb shows that the H-wave amplitude at 100 ms, 50 ms, and 20 ms at both 1 week and 6 weeks is maintained; this indicates a reduction in RDD. Surprisingly, the contralesional hindlimb did not show any H-wave reduction, and displays the characteristic negative slope. H-wave reduction for the different conditions is compared at the 20 ms ISI (Fig 2C). Only the contralesional forelimb bar graphs show a significate increase in the H-wave response percentage compared to baseline at 7 dpi (one-way ANOVA; F=5.77; p=0.01) and 42 dpi (one-way ANOVA; F=5.77; p=0.02); all others are not significantly different (see Fig 2 legend for statistics). This indicates that

Figure 2. Rate-dependent depression. A, B. Rate-dependent depression plots for forelimb (A) and hindlimb (B). Insets show linear regression slopes for pre-PTX, post-PTX at 7-days, and post-PTX at 42-days. Color coding of groups is: pre-PTX, black; post-PTX at 7-days, light blue, post-PTX at 42 days, red (code also shown in part C). Inset linear regression plots for (A) Forelimb-Contralesional (slopes; Pre=-0.03435, 7 dpi= -0.01770, 42 dpi=-0.003551; F=7.065; p=0.009). (A) Forelimb-Ipsilesional (slopes; Pre= -0.03548; 7 dpi= -0.03673; 42 dpi=-0.04104; F=0.158; p=0.855). (B) Hindlimb-Contralesional (slopes; Pre=-0.03640; 7 dpi=-0.02919; 42 dpi=-0.03382, F=0.2397, p=0.7905). (B) Hindlimb-Ipsilesional (slopes; Pre=-0.03264; 7 dpi=-0.03326; 42 dpi=-0.03346; F=0.00419, p=0.9958). C, D. Rate-dependent depression measured at 20 ms for the forelimb (C, contralateral, left; ipsilateral, right; D, contralateral, left; ipsilateral, right). Contra-lesional hindlimb (Figure 2C) shows no significant difference from baseline at 7 DPI (one-way ANOVA; F=1.41; p=0.21) and 42 dpi (one-way ANOVA; F=1.41.77; p=0.48), which indicates that RDD does not extend to the hindlimb.

hyperreflexia is present only for the contralateral forelimb. Importantly, that RDD is not different at the two timepoints indicates that it also is persistent. We combined data from the H-reflex animals with that of an additional cohort (n=5), where we added two additional time points (2-dpi and 14-dpi), to determine the time course of development of hyperreflexia (Supplemental Figure 1). We observed a progressive reduction in RDD (one-way ANOVA; F= 6.548), with a numerical reduction at 2-dpi, significance from baseline by 7 dpi (p=0.005 Tukey post hoc), and maintenance of this difference at 42 dpi (p=0.0005). This suggests gradual development of hyperreflexia, as early as 2-dpi and maintenance from 6-dpi. Our findings show that a complete unilateral CST lesion results in only a segmentally specific forelimb hyperreflexia; is presents only in the contralesional forelimb and not the contralesional hindlimb.

Figure 3. Motor neuron KCC2. A. KCC2 immunostaining in the cervical and lumbar levels for the early and chronic cohorts, and naïve rats. B) Ratio of membrane/cytosolic KCC2 for cervical cord showed no significant difference between early, chronic, and naïve groups(one-way ANOVA; F=0.507; p=0.77).C) Same as B but for lumbar cord (one-way ANOVA; F=0.756; p=0.58).

Hyperreflexia is not due to KCC2-mediated intrinsic motor neuronal hyperexcitability

Reduction in membrane-bound KCC2, by internalization and subsequent cytosolic degradation (Côté, Murray, & Lemay, 2017), after injury is a well-established mechanism supporting a role for motor neuron hyperexcitability in hyperreflexia (Boulenguez et al., 2010); (Kathe et al., 2016); (Cˆote, 2021). We next determined if unilateral PTX also produces a KCC2 reduction. High-resolution confocal micrographs (Supplemental Fig 2A; Figure 3A) show characteristic dense KCC2 membrane staining in motor neurons in the naïve animals. Surprisingly, after PTX, we did not observe a change in KCC2 staining for the contralesional cervical of lumbar segments (Figure 3 A). Recall that animals showed significant hyperreflexia only for the contralesional forelimb. We quantified the ratio of membrane-bound to cytosolic KCC2 (Supplemental Fig 2B, C). For the cervical enlargement ECR motor pool, comparison of ipsilesional and contralesional labeling in both early and chronic timepoints show no difference in KCC2 ratios. Importantly, these ratios are not significantly different from the naïve, ruling out a bilateral reduction (see Fig 3 legend for statistics). Surprisingly, despite a reduction in

contralesional forelimb RDD, there is no associated change in contralesional cervical KCC2. Also, there was no change in lumbar KCC2 (see Fig 3C legend for statistics), which is expected since there was no RDD reduction in the contralesional hindlimb. To validate KCC2 immunohistochemistry, we performed western blot analysis and calculated relative protein levels of KCC2 from extracted cervical and lumbar tissue. Analysis of ipsilesional, contralesional, and naive KCC2 from both cervical and lumbar levels collected at the early timepoint show no significant differences (Supp Fig 3 legend). Further, the absence of a KCC2 effect is not due to normalizing gray matter levels to white matter KCC2 (Boulenguez et al., 2010). Comparisons of ipsilesional, and contralesional sides, at early and chronic timepoints, and with the two sides in naïve rats, show no significant differences in the cervical membrane KCC2 (Supplemental Fig 4A; statistics described in legend). The same is true for comparisons done with labeling in the cervical cytosol, lumbar membrane, and lumbar cytosol (Supplemental Fig 4B, C, D; see legends for statistic values). Taken together, our findings show that KCC2 is not a contributing mechanism driving motor neuronal hyperreflexia after PTX.

Figure 4. 3-dimensional reconstruction of motor neurons. A. Representative motor neuron from a naïve rat (whole-cell reconstruction, left; 3-D cluster of synapses, top right; single optical slice, bottom right). B. representative motor neurons after unilateral PTX. (Calibration: 20μm.

VGlugt1 1A terminals increase on cervical and lumbar motor neurons but GABApre terminals increase only on lumbar motor neurons

Previously we showed that unilateral PTX produces afferent fiber sprouting (Tan et al., 2012) both increased axon density in the dorsal horn and intermediate zone, as well as increased 1a terminals on motor neurons. With hyperreflexia occurring only in the contralesional forelimb after PTX, and no reduction in membrane KCC2, we next examined 1a afferent circuit changes at cervical and lumbar levels. We used high-resolution 3-D

reconstruction (Imaris; Figure 4A) to monitor all contacts between VGlut1+ afferent terminals on motor neurons and associated GABA presynaptic inhibitory synapses on single motor neurons. In a representative control ECR motor neuron, we see VGlut1-labeled presynaptic terminals (red) scattered over the somatic and dendritic membrane. GAD-positive GABApre terminals are located on many of the VGlut1 terminals (Betley et al., 2009). An enlargement of the boxed area (A, top right) and associated single optical slice (A, bottom right) reveal a cluster of synapses. After PTX, we observed in the cervical cord, more 1a terminals on the contralesional compared with ipsilesional cervical motor neurons (4B, red arrows), showing axon terminal sprouting. By contrast, the number of GABApre (green) on the two sides were similar. The contralesional lumbar motor neuron (4B, bottom) revealed both increased 1A presynaptic terminals (red arrowheads) and a commensurate increase in GABApre presynaptic terminals (green) on the 1A terminals compared with the ipsilesional side.

Figure 5. Quantification of 1A and GABApre synapses after unilateral PTX; data from the early and chronic cohorts are shown. A. Cervical 1A synapses. B. Cervical GABApre (early-unpaired t-test; 1.663 ± 2.291 ; p<0.47), (chronic-unpaired t-test; 3.160 ± 2.550 ; p<0.21). C. Lumbar 1A synapses (unpaired t-test 0.0002288 ± 0.0002961; p=0.44) D. Lumbar GABApre (chronic-unpaired t-test; 3.971 ± 3.118 ; p<0.206).

Quantification of these morphological changes for the cervical segment (Figure 5A) revealed significant increases in 1a terminals, normalized to motor neuron surface area, for both the 7 dpi and 42 dpi time points (7 dpi: unpaired t-test $0.001. \pm 0.0003$; p<0.0005; 42 dpi: unpaired t-test test 0.001 ± 0.0003; p<0.0008; Fig5A). We then quantified the number of GABApre at early and chronic timepoints and found no change between contralesional, and ipsilesional (Fig 5B).

In the lumbar cord, we observed an increase in normalized 1a terminals at the onto TA motor neurons. Quantification showed a significant increase contralesional to the injury compared to ipsilesional at the early

timepoint (unpaired t-test 0.001 ± 0.0004;p<0.0083)(Fig5C) with no significant difference at the chronic timepoint. Importantly, unlike in the cervical cord, we observed a significant increase in GABApre synaptic inputs (unpaired t-test 3.88 ± 2.94; p<0.014) (Fig5D) suggesting appropriate presynaptic inhibitory regulation of the actions of afferent sprouting. These findings show sprouting of 1A presynaptic terminals for both the cervical and lumbar motor neurons, but a balanced increase in GABApre terminals only in the lumbar cord. Interestingly, for the chronic timepoint, there is neither a significant increase in 1A terminals, nor an increase in GABApre Fig5D, suggesting that the lumbar circuits have pruned the sprouted 1A terminals.

Discussion

In addition to upper limb muscle weakness and loss of voluntary control after injury to the motor systems, hyperreflexia is a key gain-of-function impairment that disrupts control of

	Level	Hyperreflexia	KCC2	1A sprouting	GABAPre
7dpi	Forelimb/Cervical	↑	↔	↑	↔
	Hindlimb/Lumbar	↔	↔	↑	↑
42dpi	Forelimb/Cervical	↑	↔	↑	↔
	Hindlimb/Lumbar	↔	↔	↔	↔

Figure 6. Summary of contralateral forelimb and hindlimb changes in reflex strength, measured as rate-dependent depression, and corresponding changes in KCC2, 1A afferent density on motor neurons, and GABApre at the cervical and lumbar levels, at 7-dpi and 42-dpi. Note, there were no changes ipsilesionally.

arm and hand movements. Hyperreflexia includes stronger reflexes with aberrant patterns of muscle contraction and spasticity (Thompson et al., 1992). In this study, we found intriguing dissociations between unilateral CST loss, 1A circuit plasticity, and motor neuron membrane bound KCC2 that provide new insights into the mechanisms of hyperreflexia (Figure 6, which summarizes data for the contralesional side at early and chronic timepoints). Only the contralateral forelimb developed hyperreflexia, as measured by a reduction in RDD. Neither the contralesional hindlimb nor both ipsilateral limbs showed hyperreflexia. This leads to the expectation that an injury-dependent change associated with, and possibly causal to, hyperreflexia in this model will be present in the contralateral cervical cord only. KCC2 reduction, key to hyperreflexia after SCI and bilateral CST lesion (Kathe et al., 2016), (Chen, 2018),(Beverungen, 2020), does not associate with cervical hyperreflexia in the unilateral PTX model, showing that hyperreflexia can occur without enhanced motor neuronal excitability due to downregulation of KCC2. Whereas 1A sprouting is similarly present at both the cervical and lumbar levels at 7dpi, likely due to loss of synaptic

competition after CST lesion (Jiang et al., 2016), hyperreflexia is only present in the forelimb. Thus, 1A sprouting alone does not associate with hyperreflexia when the cervical and lumbar segments are compared. Finally, the lack of a change in GABApre in the cervical cord suggests that 1A afferent sprouting leads to enhanced reflex signaling because of less presynaptic inhibitory regulation. By contrast, the increase in GABApre terminals in the lumbar spinal cord parallels the increase in 1A terminals. This is consistent with effective presynaptic inhibitory regulation for hindlimb 1A afferent signaling and, in consequence, no hyperreflexia.

The forelimb is more vulnerable than the hindlimb to developing hyperflexia

Clinical studies show that after unilateral cortical stroke, spasticity is more prevalent and severe in the upper limb than for the lower limb (Katoozian, Tahan, Zoghi, & Bakhshayesh, 2018) (Sunnerhagen et al., 2019). Paralleling this findings, we found that unilateral PTX produces forelimb not hindlimb hyperreflexia. As described above, we have used this finding in support of potential causality of a lack of GABApre sprouting failing to check 1A afferent sprouting, and contributing to hyperreflexia in the unilateral PTX model. This is a novel finding, as animal studies examine the forelimb or hindlimb, but rarely both. Animal studies further suggest that the CST may have differential functional roles for movement control of the forelimb and hindlimb. The CST in mice has a more dorsal projection to the lumbar spinal cord (Bareyre, Kerschensteiner, Misgeld, & Sanes, 2005), compared with more extensive projections into the intermediate zone and even ventral projections for the cervical projection (Asante & Martin, 2013), (Steward et al., 2021), (Bennett, Das, & Emmady, 2025). Although the lumbar CST projection was shown to be involved in sensory gating—via deep dorsal horn interneurons (Moreno-Lopez et al., 2021), with motor actions relayed via brain stem and rostral spinal neurons (Moreno-López et al., 2016)—this has not been similarly studied for the cervical spinal cord. The lack of hyperreflexia after CST loss, in the absence of other spinal pathology, may reflect a diminished role for the lumbar-projecting CST in feed-forward control of hindlimb/lower limb motor neurons and muscle contractions. Hyperreflexia may thus reflect a 'dose effect,' with a higher density of CST terminations on ventral motor circuits before injury, and hyperreflexia occurring with its loss after injury. Whereas the CST has different roles in control of the upper and lower limbs in humans, owing to the differential movement capabilities, it is not known if there are CST density differences.

Hyperreflexia is not a progressive condition

Our assessment of hyperreflexia showed RDD was numerically reduced at day 2, the earliest time point tested, and was significant at 1-week post-injury. This early presentation suggests a pathophysiological process triggered by the sudden loss of connections, or loss of their activity. The significant RDD reduction at 1-week post-injury was not different from 6-weeks, showing both persistence of hyperreflexia and that it did not worsen. Taken together, hyperreflexia as measured by RDD is not progressive. This early presence and maintenance over several weeks, lead us to consider the process of hyperreflexia development. The RDD reduction at the earliest timepoint, although not significant, suggests an immediate circuit level reorganization: the sudden loss of CST connections led to competitive primary afferent sprouting after injury (Jiang et al., 2016). The rapid onset and lack of progressive worsening over the 6 weeks of testing parallels observations in humans after stroke. Upper limb spasticity occurs 3 days after first presentation of a stroke in 25% of patients examined (Opheim, Danielsson, Alt Murphy, Persson, & Sunnerhagen, 2014), (Opheim, Danielsson, Alt Murphy, Persson, & Sunnerhagen, 2015). Using logistic regression analysis, the best predictor for persistent spasticity at 12 months was spasticity at 10 days post injury (Sunnerhagen et al., 2019).

We identified sprouting of glutamatergic terminals on motor neurons at 1-week post injury, showing that a morphological change can occur remarkably early. Additionally, in the lumbar cord, GABApre terminals also showed an increase at this early time point. Similarities between unilateral CST injury and stroke, as they relate to time course of hyperreflexia in animals and spasticity onset in humans, further establish the unilateral PTX model as a tool to help determine optimal therapeutic strategies aimed at reducing spasticity after a brain injury.

Do spared ipsilateral CST axons protect from increased excitability after unilateral PTX?

We found no evidence for a reduction in membrane-bound KCC2 contralesionally. This shows that hyperreflexia after CST loss is not due to KCC2 mediated motor neuron hyperexcitability. SCI and bilateral PTX, which completely lesion all CST fibers have identified KCC2 downregulation, as an important contributor to motor neuron excitability and, in turn, hyperreflexia (Boulenguez et al., 2010), (Beverungen, 2020). Therapeutic interventions targeting restoration of membrane bound KCC2, or that result in restoring KCC2, are associated with a significant reduction of hyperreflexia, and spasticity (Kathe et al., 2016), (Cˆote, 2021). Though

the association between these injury models and KCC2 downregulation is clear, the unilateral PTX model used in this study shows evidence of a disassociation between CST injury, hyperreflexia, and motor neuron hyperexcitability. The question remains why with unilateral PTX, KCC2 is not involved when it is a contributing factor after bilateral CST lesion and SCI. We suspect the presence of ipsilateral CST is sufficient to protect against hyperreflexia by providing necessary segmental excitatory input; bilateral CST lesion eliminates that protection. Moreover, there is evidence of injury-induced plasticity of spared ipsilateral CST axons after a unilateral injury; whereby, with the loss of the contralateral CST, ipsilateral CST fibers sprout (Brus-Ramer, Carmel, Chakrabarty, & Martin, 2007). There can be a doubling of this injury-dependent ipsilateral CST sprouting (Brus-Ramer et al., 2007), which could provide compensatory connections to help support segmental excitability. Utilizing these spared CST projections, could be further strengthened with electrical stimulation(Brus-Ramer et al., 2007). It is possible the ipsilateral CST projects sufficient excitatory inputs to mitigate motor neuron hyperexcitability that results from KCC2 reduction, but not enough to prevent hyperreflexia in the affected forelimb.

Constitutive activation of the serotonin receptor 5-HT$_{2c}$ is another well studied mechanism contributing to motor neuron hyperexcitability and spasticity in humans and rodents (Gorassini, Knash, Harvey, Bennett, & Yang, 2004), (Murray et al., 2010). The switch to constitutive 5-HT$_{2c}$ activation following spinal cord injury it is thought to be provoked by the loss of descending monoaminergic inputs to the spinal cord. Although the shift to constitutively active 5-HT$_{2c}$ was not examined in this study, it is not expected to contribute to hyperreflexia, at least early on. A study on the time course of 5-HT$_{2c}$ receptor expression in rats after SCI estimates 1.3x fold increase in 5-HT$_{2c}$ receptor activity when compared to the sham group as early as 14 days after injury (Ren et al., 2013). This time course indicates that the shift to the constitutively active form cannot explain our finding that hyperreflexia begins within days after PTX. Nor does the presence of constitutive 5-HT$_{2c}$ activity explain maintenance without further loss of RDD. It may, however, contribute to development of spasms.

Conclusion: Multiple pathophysiological mechanisms are able to trigger hyperreflexia after injury

Our findings show that hyperreflexia after unilateral PTX is explained by a change in the

1A afferent fiber circuit: sprouting of 1A VGlut1 synapses on motor neurons is not accompanied by commensurate GABApre sprouting in the cervical segments. The unilateral PTX model emphasizes that hyperreflexia can be mediated by multiple factors. Among the three factors that we have examined—1A sprouting, reduced GABApre regulation, and reduced membrane-bound KCC2—each could have, in principle, led to a stronger reflex. But only the lack of cervical GABApre sprouting explained restricted hyperreflexia in the contralateral forelimb.

Considering our findings in the broader context of motor system injuries, our findings show that hyperreflexia can present as a circuit disorder without need for KCC2 motor neuron hyperexcitability. The physiological actions of sprouting of 1A afferent glutamatergic terminals on motor neurons are not checked by concomitant GABApre sprouting; there is a higher proportion of VGlut1 terminals on motor neurons without GABApre regulation.

Supplemental Figure 1. KCC2 immunohistochemistry measurement and analyses. KCC2 Normalized membrane/cytosol averages. A. Methods. B. Cervical membrane comparison (one-way ANOVA; F=2.030; p=0.079). C. Cervical cytosol comparison (one-way ANOVA; F=0.519; p=0.760). D. Lumbar membrane comparison (one-way ANOVA; F=1.872; p=0.141). E) Lumbar cytosol comparison (one-way ANOVA; F=0.234; p=0.872).

Supplemental Figure 2. Hyperreflexia Development. Multiple comparison analysis averaged from main, and supplemental animal cohorts(n=11) shows the only significant increase in RDD (hyperreflexia) at PTX 6/7-DPI (one-way ANOVA; F=6.548; p=0.005), and at PTX 42-DPI(one-way ANOVA; F=6.548; p=0.0005).

Supplemental Figure 3. Western blotting for KCC2. A. Representative western blots from cervical (top) and lumbar (bottom) spinal tissue. B. Cervical protein level comparisons (one-way ANOVA; F=0.4697; p=0.63). C. Lumbar protein level comparisons (one-way ANOVA; F=1.789; p=0.19).

A

B

Supplemental Figure 4. Motor neuron surface area averages. Quantification of average ipsilateral, and contralateral motor neuron size by surface area (μm^2). A. Cervical, early and chronic comparisons (one-way ANOVA; F=1.71; p=0.16). B. Lumbar, early & chronic comparisons (one-way ANOVA; F=1.68; p=0.17).

Supplemental Figure 5. Cervical level assessment of normalized VGlut1 and GABApre-positive contacts on ADQ motor neurons. A. Normalized VGlut1 (unpaired t-test 0.0007970 ± 0.0003057; p=0.011). B) GABApre (unpaired t-test 0.6022 ± 2.694; p=0.823).

References

Alvarez, F. J., Villalba, R. M., Zerda, R., & Schneider, S. P. (2004). Vesicular glutamate transporters in the spinal cord, with special reference to sensory primary afferent synapses. *J Comp Neurol, 472*(3), 257-280. doi:10.1002/cne.20012

Asante, C. O., & Martin, J. H. (2013). Differential joint-specific corticospinal tract projections within the cervical enlargement. *PLOS ONE, 8*(9), e74454. doi:10.1371/journal.pone.0074454

Bareyre, F. M., Kerschensteiner, M., Misgeld, T., & Sanes, J. R. (2005). Transgenic labeling of the corticospinal tract for monitoring axonal responses to spinal cord injury. *Nat Med, 11*(12), 1355-1360. doi:10.1038/nm1331

Bennett, J., Das, J. M., & Emmady, P. D. (2025). Spinal Cord Injuries. In *StatPearls*. Treasure Island (FL) ineligible companies. Disclosure: Joe Das declares no relevant financial relationships with ineligible companies. Disclosure: Prabhu Emmady declares no relevant financial relationships with ineligible companies.: StatPearls Publishing

Copyright © 2025, StatPearls Publishing LLC.

Betley, J. N., Wright, C. V., Kawaguchi, Y., Erdelyi, F., Szabo, G., Jessell, T. M., & Kaltschmidt, J. A. (2009). Stringent specificity in the construction of a GABAergic presynaptic inhibitory circuit. *Cell, 139*(1), 161-174. doi:10.1016/j.cell.2009.08.027

Beverungen, H. C. t., MP. (2020). Rehabilitation Decreases Spasticity by Restoring Chloride Homeostasis through the Brain-Derived Neurotrophic Factor–KCC2 Pathway after Spinal Cord Injury. *JOURNAL OF NEUROTRAUMA, 37*.

Boulenguez, P., Liabeuf, S., Bos, R., Bras, H., Jean-Xavier, C., Brocard, C., . . . Vinay, L. (2010). Down-regulation of the potassium-chloride cotransporter KCC2 contributes to spasticity after spinal cord injury. *Nat Med, 16*(3), 302-307. doi:10.1038/nm.2107

Brus-Ramer, M., Carmel, J. B., Chakrabarty, S., & Martin, J. H. (2007). Electrical stimulation of spared corticospinal axons augments connections with ipsilateral spinal motor circuits after injury. *J Neurosci, 27*(50), 13793-13801. doi:10.1523/jneurosci.3489-07.2007

Cˆote, J. N. B. M.-P. (2021). Enhancing KCC2 activity decreases hyperreflexia and spasticity after chronic spinal cord injury. *Experimental Neurology*.

Chamma, I., Chevy, Q., Poncer, J. C., & Levi, S. (2012). Role of the neuronal K-Cl co-transporter KCC2 in inhibitory and excitatory neurotransmission. *Frontiers in CELLULAR NEUROSCIENCE, 6*. Retrieved from https://www.frontiersin.org/articles/10.3389/fncel.2012.00005

Chen, B. (2018). Reactivation of Dormant Relay Pathways in Injured Spinal Cord by KCC2 Manipulations. *Cell*.

Côté, M. P., Murray, M., & Lemay, M. A. (2017). Rehabilitation Strategies after Spinal Cord Injury: Inquiry into the Mechanisms of Success and Failure. *J Neurotrauma, 34*(10), 1841-1857. doi:10.1089/neu.2016.4577

Gorassini, M. A., Knash, M. E., Harvey, P. J., Bennett, D. J., & Yang, J. F. (2004). Role of motoneurons in the generation of muscle spasms after spinal cord injury. *Brain, 127*(Pt 10), 2247-2258. doi:10.1093/brain/awh243

Hardman, V. J., & Brown, M. C. (1985). Spatial organization within rat motoneuron pools. *Neurosci Lett, 60*(3), 325-329. doi:10.1016/0304-3940(85)90598-1

Hiersemenzel, L. P., Curt, A., & Dietz, V. (2000). From spinal shock to spasticity: neuronal adaptations to a spinal cord injury. *Neurology, 54*(8), 1574-1582. doi:10.1212/wnl.54.8.1574

Ho, S. W., Waite, P.M. (2002). Effects of Different Anesthetics on the Paired-Pulse Depression of the H Reflex in Adult Rat. *Experimental Neurology, 177,* 494-502.

Hosoidoa, T. (2009). Characteristics of H- and M-waves recorded from rat forelimbs. *Neuroscience Letters, 450.*

Hübner, C. A., Stein, V., Hermans-Borgmeyer, I., Meyer, T., Ballanyi, K., & Jentsch, T. J. (2001). Disruption of KCC2 Reveals an Essential Role of K-Cl Cotransport Already in Early Synaptic Inhibition. *Neuron, 30*(2), 515-524. doi:https://doi.org/10.1016/S0896-6273(01)00297-5

Hultborn, H. (2003). Changes in neuronal properties and spinal reflexes during development of spasticity following spinal cord lesions and stroke: studies in animal models and patients. *J Rehabil Med*(41 Suppl), 46-55. doi:10.1080/16501960310010142

Jiang, Y. Q., Zaaimi, B., & Martin, J. H. (2016). Competition with Primary Sensory Afferents Drives Remodeling of Corticospinal Axons in Mature Spinal Motor Circuits. *J Neurosci, 36*(1), 193-203. doi:10.1523/JNEUROSCI.3441-15.2016

Kathe, C., Hutson, T. H., McMahon, S. B., & Moon, L. D. (2016). Intramuscular Neurotrophin-3 normalizes low threshold spinal reflexes, reduces spasms and improves mobility after bilateral corticospinal tract injury in rats. *eLife, 5.* doi:10.7554/eLife.18146

Katoozian, L., Tahan, N., Zoghi, M., & Bakhshayesh, B. (2018). The Onset and Frequency of Spasticity After First Ever Stroke. *J Natl Med Assoc, 110*(6), 547-552. doi:10.1016/j.jnma.2018.01.008

Lee, H. H. C., Walker, J. A., Williams, J. R., Goodier, R. J., Payne, J. A., & Moss, S. J. (2007). Direct Protein Kinase C-dependent Phosphorylation Regulates the Cell Surface Stability and Activity of the Potassium Chloride Cotransporter KCC2 *. *Journal of Biological Chemistry, 282*(41), 29777-29784. doi:10.1074/jbc.M705053200

Lundström, E., Smits, A., Terént, A., & Borg, J. (2010). Time-course and determinants of spasticity during the first six months following first-ever stroke. *J Rehabil Med, 42*(4), 296-301. doi:10.2340/16501977-0509

Moreno-Lopez, Y., Bichara, C., Delbecq, G., Isope, P., & Cordero-Erausquin, M. (2021). The corticospinal tract primarily modulates sensory inputs in the mouse lumbar cord. *eLife, 10.* doi:10.7554/eLife.65304

Moreno-López, Y., Olivares-Moreno, R., Cordero-Erausquin, M., & Rojas-Piloni, G. (2016). Sensorimotor Integration by Corticospinal System. *Front Neuroanat, 10*, 24. doi:10.3389/fnana.2016.00024

Murray, K. C., Nakae, A., Stephens, M. J., Rank, M., D'Amico, J., Harvey, P. J., . . . Fouad, K. (2010). Recovery of motoneuron and locomotor function after spinal cord injury depends on constitutive activity in 5-HT2C receptors. *Nat Med, 16*(6), 694-700. doi:10.1038/nm.2160

Opheim, A., Danielsson, A., Alt Murphy, M., Persson, H. C., & Sunnerhagen, K. S. (2014). Upper-limb spasticity during the first year after stroke: stroke arm longitudinal study at the University of Gothenburg. *Am J Phys Med Rehabil, 93*(10), 884-896. doi:10.1097/phm.0000000000000157

Opheim, A., Danielsson, A., Alt Murphy, M., Persson, H. C., & Sunnerhagen, K. S. (2015). Early prediction of long-term upper limb spasticity after stroke: part of the SALGOT study. *Neurology, 85*(10), 873-880. doi:10.1212/wnl.0000000000001908

Oudega, M. (2012). Corticospinal reorganization after spinal cord injury. *The Journal of Physiology*.

Ren, L. Q., Wienecke, J., Chen, M., Møller, M., Hultborn, H., & Zhang, M. (2013). The time course of serotonin 2C receptor expression after spinal transection of rats: an immunohistochemical study. *Neuroscience, 236*, 31-46. doi:10.1016/j.neuroscience.2012.12.063

Romanes, G. J. (1964). The Motor Pools of the Spinal Cord. In J. C. Eccles & J. P. Schadé (Eds.), *Progress in Brain Research* (Vol. 11, pp. 93-119): Elsevier.

Sharif, H., Alexander, H., Azam, A., & Martin, J. H. (2021). Dual motor cortex and spinal cord neuromodulation improves rehabilitation efficacy and restores skilled locomotor function in a rat cervical contusion injury model. *Exp Neurol, 341*, 113715. doi:10.1016/j.expneurol.2021.113715

Steward, O., Yee, K. M., Metcalfe, M., Willenberg, R., Luo, J., Azevedo, R., . . . Gandhi, S. P. (2021). Rostro-Caudal Specificity of Corticospinal Tract Projections in Mice. *Cereb Cortex, 31*(5), 2322-2344. doi:10.1093/cercor/bhaa338

Sunnerhagen, K. S., Opheim, A., & Alt Murphy, M. (2019). Onset, time course and prediction of spasticity after stroke or traumatic brain injury. *Ann Phys Rehabil Med, 62*(6), 431-434. doi:10.1016/j.rehab.2018.04.004

Tan, A. M., Chakrabarty, S., Kimura, H., & Martin, J. H. (2012). Selective corticospinal tract injury in the rat induces primary afferent fiber sprouting in the spinal cord and hyperreflexia. *J Neurosci, 32*(37), 12896-12908. doi:10.1523/JNEUROSCI.6451-11.2012

Thompson, F. J., Reier, P. J., Lucas, C. C., & Parmer, R. (1992). Altered patterns of reflex excitability subsequent to contusion injury of the rat spinal cord. *J Neurophysiol, 68*(5), 1473-1486. doi:10.1152/jn.1992.68.5.1473

Urban, P. P., Wolf, T., Uebele, M., Marx, J. J., Vogt, T., Stoeter, P., . . . Wissel, J. (2010). Occurence and clinical predictors of spasticity after ischemic stroke. *Stroke, 41*(9), 2016-2020. doi:10.1161/strokeaha.110.581991