Title: The economic strategies of superorganisms

Authors: Lily Leahy[1*], Hannah L. Riskas[1], Ben Halliwell[2,3], Ian J. Wright[4,5], Amelia G. Carlesso[1,6], Nathan J. Sanders[7], Tom R. Bishop[8,9], Catherine L. Parr[9,10], Steven L. Chown[11], Jonathan Z. Shik[12], Alan N. Andersen[13], Heloise Gibb[6, 14]

Affiliations:

[1] Department of Ecological, Plant, and Animal Sciences, La Trobe University; Bundoora, VIC, Australia

[2] Biological Sciences, School of Natural Sciences, University of Tasmania; Hobart, TAS, Australia

[3] ARC Centre of Excellence for Plant Success in Nature and Agriculture, University of Tasmania; Hobart, TAS, Australia

[4] Hawkesbury Institute for the Environment, Western Sydney University; Penrith, NSW, Australia

[5] School of Natural Sciences, Macquarie University; North Ryde, NSW Australia

[6] School of Life and Environmental Sciences, Deakin University; Burwood, VIC, Australia

[7] Department of Ecology and Evolutionary Biology, University of Michigan; Ann Arbor, MI, USA

[8] School of Biosciences, Cardiff University; Cardiff, UK

[9] Department of Zoology and Entomology, University of Pretoria; Pretoria, South Africa

[10] School of Environmental Sciences, University of Liverpool; Liverpool, UK

[11] School of Biological Sciences, Monash University; Monash Victoria, Australia

[12] Section for Ecology and Evolution, Department of Biology, University of Copenhagen, Copenhagen, Denmark

[13] Research Institute for the Environment and Livelihoods, Charles Darwin University, Darwin, NT, Australia

[14] School of Biological, Earth and Environmental Sciences (BEES), University of New South Wales; Kensington NSW, Australia

*Corresponding author: Lily Leahy. Email: lily.leahy4@gmail.com

Abstract

The leaf economics spectrum links strategies of plant investment in resource-acquiring leaves to
overall fitness. We test whether an economic spectrum can also explain variation in ecological
strategies of ant species across environmental gradients, where colony investment in workers is
analogous to plant investment in leaves. A fast return of resource investment was associated with large
colonies of smaller, less robust, short-lived workers with low nitrogen:phosphorus ratios. Slow
resource payback was associated with small colonies of densely built, energetically conservative and
longer-lived workers with high nitrogen:phosphorus ratios. Species representing the entire economic
continuum co-occurred in all communities. Phylogenetic analyses suggest genus level conservation of
core investment templates. These results unify studies of plants and ants, suggesting common
economic principles apply across the tree of life.

Main Text:

One of the basic principles common to all life is that organisms are sustained by the storage and turnover of energy for fueling metabolism and chemical elements for constructing biomass. Pace-of-life theory posits that organisms trade-off allocation of these resources to assimilation, growth, and reproduction, with species falling on a slow-fast life history continuum (*1-4*). Modular organisms such as plants and social insects, however, organize the allocation of these tasks in a unique way. In modular organisms, resource acquisition and assimilation are specifically tasked to resource-acquiring modules that are non-reproductive (*5*). The evolution and optimisation of these modules can be described using economic concepts of investment and return of resources and nutrients (*6*). Such economic concepts have been widely developed in plants. For example, the Leaf Economics Spectrum (LES) scheme (*7*) underpins current understanding of plant investment strategies: plant species vary from investing in cheaply constructed 'fast' leaves with high nutrient concentrations of nitrogen and phosphorus, high metabolic rates, rapid assimilation of resources and short lifespans, to species with expensive but 'slow' leaves that are long-lived, with slow assimilation and metabolic rates.

It has been proposed that the leaf economic spectrum may also be applicable to other modular organisms such as social insects (*5*), but this hypothesis awaits testing. Ants represent a particularly complex form of social organisation where genetically related members of the colony act collectively as "superorganisms" to complete the functional tasks of a unitary organism (*8*). Under this social organisation, billions of individual ant workers gather and recycle the planet's resources every day, performing essential ecosystem processes, from seed dispersal to nutrient cycling and predation (*9-12*). As for modular plants, this has been a highly successful ecological strategy with both ants and plants playing dominant roles in most terrestrial ecosystems (*13, 14*). Gibb et al. (*5*) hypothesised that ants and plants have converged on the same suite of approaches to acquire and distribute resources and that this could be indexed by a set of measurable traits. This leading hypothesis could unify the study of ant colonies with that of plants under the same economic spectrum umbrella, but it has not been empirically tested.

Here, we advance generality in economic theories of life by testing whether ant species fall along a fast-slow economic spectrum akin to the leaf economics spectrum. Sampling across extensive climatic and soil phosphorus gradients (fig. S1, table S1), we acquired trait data from individual workers collected from 123 species from seven subfamilies, including the five most diverse globally. Previous work to describe the resource economics of ant colonies has been fragmented and has failed to formally quantify the process of colony investment of nutrients and biomass in ant workers with a methodology that is scalable to large datasets (*15-19*). Here, we use eight traits to quantify the economic spectrum (*5*). Six traits directly parallel those used in the leaf economics spectrum: (1) Mass-specific standard metabolic rate, representing the *energy maintenance costs* of workers, as it does for plant leaves (*20*); (2) Mass density measured as mass:volume ratio and indicative of anatomical structure (e.g., cuticle thickness (*21*)), representing an *investment cost* equivalent to leaf-mass per area; (3) Nitrogen and (4) phosphorus concentrations of body tissue representing chemical *investment costs* (*22*); (5) Mass-specific assimilation rate (intake of proteins and carbohydrates per unit time per unit mass measured in laboratory feeding trials), representing *potential revenue* for the colony and equivalent to photosynthetic rate in plant leaves (*5*); (6) Relative worker

lifespan (measured as days of survival in laboratory trials), equivalent to leaf lifespan,
representing the *duration of the revenue stream* (*7*) that should be *optimised* to balance
lifetime net resource costs and benefits (*23, 24*). Finally, we also include two ant-specific
traits, (7) worker body mass, and a relative measure of (8) worker number (here referred to as
colony size), that can be altered to change overall colony mass thereby influencing the total
colony budget (*25*).

Chemical elements of nitrogen and phosphorus play slightly different roles in plant leaves
and metazoan bodies, and we would expect a divergence in the co-variation of these two
traits for plant and ant economic spectra. In leaves, concentrations of nitrogen and
phosphorus are positively correlated, with high values of both indicating a fast economic
strategy (*7*). In small-bodied animals like ants, however, nitrogen and phosphorus should
trade-off (*26*). Nitrogen creates biological structure, such as costly cuticle, spines and dense
muscles, while phosphorus is a critical component of the protein synthesis machinery of
ribosomes (r-RNA) that sets the pace of worker growth (*27*). Broadly, greater allocation to
nitrogen-rich structure is hypothesised to slow growth, while greater allocation to phosphorus
rich r-RNA results in fast growth; hence these two elements should be negatively correlated
in ants creating a fast-slow stoichiometric spectrum (*26*).

We present our predictions of how these eight traits interrelate to form a fast-slow economic
spectrum and propose four key hypotheses that underlie these predictions (Fig. 1). These four
hypotheses integrate across a portfolio of underlying ecological theory to support an
economic spectrum in ants from first principles (*28*). Firstly, metabolic theories relate mass
and energy use to allometric scaling laws that apply to all organisms (*3, 20*). Secondly,
theories of ageing support the hypothesis that initial investment from the colony, resource
acquisition rates, and extrinsic foraging risks are correlated to shape intrinsic lifespans of ant
workers (*23, 29, 30*). Thirdly, pace of life theory posits a trade-off between offspring quality
and quantity (*21, 31*), while social evolutionary theory suggests more complex colony
organization reduces offspring reproductive potential (*32*): both theories link worker number
(quantity) to worker structural biomass (size and quality). Fourthly, as outlined above,
ecological stoichiometry theory suggests a trade-off in nitrogen and phosphorus allocation
that should shape the pace of economic growth of the colony (*26*). Finally, we predict that the
relative positions of ant species along the economic spectrum will be both phylogenetically
conserved and modulated by the environment (Fig.1) and explore this by testing for
evolutionary and biogeographic signatures.

The ant economic spectrum

We quantified relationships among the eight economic traits and decomposed the portion of
trait correlation associated with phylogeny (conserved trait correlation *sensu* (*33*)) and that
independent of phylogeny (non-phylogenetic) using multi response phylogenetic mixed
models (*33, 34*). Our analyses revealed sets of phylogenetically conserved, co-varying traits
that represent a clear trade-off in investment and resource return, forming an overall spectrum
of fast-slow economic strategy (Fig. 2a, b). Species that invest more biomass and nitrogen in
workers have small colonies of larger, longer-lived workers with low %P, lower maintenance
costs (low mass-specific metabolic rate; H1, Fig. 1) and slow rates of resource return (low
mass-specific assimilation rate) – an overall slow economic strategy (Fig. 2a, b). Species that
invest less biomass and nitrogen into their workers have higher metabolic costs, high %P

(representing fast growth rate; H4, Fig. 1) and quick rates of return on revenue invested (*20*).
High metabolic rates likely increase the foraging tempo of ant workers, increasing the rate of
resource uptake and thereby offsetting these high energy maintenance costs (*35*). These
species with a fast economic strategy achieve larger colony sizes made up of smaller workers
with less structural investment, shorter lives, and faster worker turnover (as predicted by H2,
H3: Fig. 1) (Fig. 2a, b).

Trait-trait phylogenetic correlations were strong (Fig. 2a, table S2) (majority >90% credible
interval; Data S1). Two traits with less data coverage, %P and mass-specific assimilation rate
showed weaker phylogenetic trait-trait correlations than other traits (Fig. 2a, table S2).
Phylogeny-independent trait-trait correlations were much weaker overall for all trait pairs
(table S2, fig. S2, S3, Data S1). Ant economic traits also had strong phylogenetic signal
indicating conserved trait correlation (average posterior mean λ across traits: 0.62, range:
0.36 – 0.84; fig. S2, table S3). Lifespan (mean $\lambda = 0.36$) and nitrogen (mean $\lambda = 0.41$),
however, were less phylogenetically conserved than other traits. Model validation confirmed
that the fitted model generated plausible data for all response variables including for traits
that had lower data coverage (posterior predictive checks: fig. S4). Leave-one-out cross
validation indicated that model predictions were robust for predicting new data and were not
overly influenced by any one species (LOO-CV: fig. S5).

Selection has therefore favoured workers that gather resources at a rate that reflects their
investment costs and the duration of their revenue stream (*28, 29*). This parallels optimal
lifespan theory for leaves (*23*), where average net resource carbon gain is maximized over the
leaf life cycle (*36*). Consequently, some trait combinations will be rare or never observed
(*37*). For example, tough leaves, or robust ant workers with short lifespans, will not have
enough photosynthetic/foraging time to pay off high resource investment.

Biophysical laws shape the economic spectrum

The independent component of trait-trait relationships (not associated with phylogenetic
history) revealed three notable correlations that are likely to be driven by fundamental
biophysical principles dictating energy-mass balance (*3, 20*). Mass-specific metabolic and
assimilation rates showed an allometric negative relationship with body mass for both
phylogenetic (Fig. 2a, b) and independent trait correlations (table S2, fig. S3). From cells to
organisms to superorganisms (*38*), metabolic rate (*B*) scales allometrically with mass,
according to a power function $B = a\mathrm{M}^\beta$ where, β is typically less than 1 (>-1 and <0 when
mass-specific). Here, the mean slope of the independent component of these relationships
was -0.55 (mass-metabolic) and -0.58 (mass-assimilation) respectively, indicating that
smaller workers use more energy and consume food resources at a faster rate per unit body
mass than do larger workers (as predicted in H1: Fig. 1, table S2).

Several theories have proposed that physio-chemical factors place fundamental mass-based
limits on the metabolic and assimilation rates of organisms (*39*). These allometric
relationships are hypothesised to relate to either nutrient supply networks (the basis of
metabolic theory of ecology (*20*)) or to surface-area scaling dynamics of energy reserve pools
and volume of structural biomass (dynamic energy budget theory (*3*)). The non-phylogenetic
proportion of these correlations is therefore likely due to biophysical laws that determine the
boundaries of economic-strategy space.

Clades partition economic-strategy space

When visualized in multidimensional trait space, the ant economic spectrum was observed to occur over two axes of variation. The first axis represents the ant worker economic spectrum (AWES), explaining 61% of variance. This axis was characterised by the correlation between biomass investment and lifespan trading off against mass-specific metabolic rate, mass-specific assimilation rate, and colony size (Fig. 3). The second axis of variation represents a stoichiometry spectrum (SS), represented by the trade-off of N and P, explaining 13% of variance (as predicted by H4: Fig. 1). This shows a distinct departure from the leaf economics spectrum in which N and P in plant leaves are positively correlated and co-vary with other LES traits over one axis of variation (7). The N-P trade-off in ants aligns with ecological stoichiometry theory applicable to small metazoans (27). Although we did not explicitly test worker growth rates here, the negative association between N and P is consistent with our hypothesis that that biomass investment in nitrogen-rich structures and chemicals (e.g., thick cuticle (21), proteinaceous venoms (40)) is likely to be traded-off with worker growth rate represented by %P (21, 31) (Fig. 2b, Fig. 3).

The different subfamilies occupy distinct regions of multidimensional economic strategy space (MANOVA – Axis 1: $F_{(6, 116)}$ = 26.86, p < 0.0001, Axis 2: $F_{(6, 116)}$ = 16.02, p < 0.0001, Fig. 3, 4). For example, two of the most globally diverse subfamilies, Dolichoderinae and Myrmicinae, generally occupy the fast end of the ant worker economic spectrum. Within that, myrmicines are placed on the slower end of the stoichiometry spectrum with higher N content compared with dolichoderines (Fig. 3, fig. S6). Many dolichoderine genera had a relatively fast strategy over all traits including the behaviourally dominant *Iridomyrmex* and *Anonychomyrma* (Fig. 4). Larger-bodied species of Formicinae, another globally diverse clade, fall on the slower end of the ant worker economic spectrum, but on the fast end of stoichiometry spectrum, with high phosphorus and low nitrogen concentrations, likely due to lower investment in cuticle thickness and a lack of proteinaceous venoms (21, 40) (Fig. 3, Fig. 4). Meanwhile, the more densely built Ectatomminae and Myrmeciinae (41) exhibit a slow strategy across both axes (Fig. 3, 4, fig. S6). This partitioning of economic spectrum space across the two axes indicates that there are different templates for being a "slow" or "fast" ant (Fig. 4).

Full economic spectrum persists in all environments

Other studies have shown that some ant traits obey universal physiological constraints (38, 42) and are strongly phylogenetically conserved (43), while others vary across both geographic (25) and regional environmental gradients (44, 45). Here, however, we found that the full suite of economic strategies was present in all environmental conditions tested (fig. S8, table S4). Local-scale climate or soil phosphorus, for example, did not shape key traits linked to economic strategies (MR-PMM trait:fixed effects, average p-value = 0.7, Data S2). This is consistent with other studies showing that local communities can contain the entire regional pool of ant trait variation (46). Similar global patterns are found in plants: some leaf traits have relatively strong relationships with climate (e.g., leaf mass per area with aridity (47)) but species positions on the slow-fast economic spectrum appear largely independent of climate (7).

We further note that economic strategies can vary within-species at local scales through phenotypic plasticity. For instance, common garden experiments have revealed that exposure

to predators can promote colony defensive allocation (e.g., to major castes (*48*)) and thus
likely caste-specific stoichiometric allocation demands (*49*). Further, resource manipulation
studies have shown that overall ant activity can increase with phosphorus supplementation
(*50*). Thus, core features of the economic spectrum are presumably subject to environmental
modulation at the population level. Here, it appears that species with both slow and fast
strategies persist under a broad range of environmental conditions. Future studies could
investigate whether the relative abundances of species differing in economic spectrum
positions changes with the environment (e.g. through community weighted means), and
particularly with respect to disturbance regimes, which have previously been linked with the
key trait of body size at a global scale (*51, 52*).

Conclusion

We have described a phylogenetically conserved economic spectrum for a globally dominant
group of superorganisms. Based on species presence, the spectrum did not vary
systematically with climate and environmental gradients. Our results suggest that early
lineages of ants likely diverged in their economic strategies and subsequently diversified
(*53*), thereby locking in these fundamental templates of investment and resource return. We
have presented a foundational test of this hypothesis using Australian ant species. We call for
more global sampling, for example, comparing economic spectrum traits in ants across
broader biogeographic gradients, such as temperate versus tropical, and also in other social or
modular organisms. Given the strong role of evolutionary history, compared with climate and
environment, in the economic spectrum, we predict that more globally extensive sampling of
ant communities would align with our key results.

The economic spectrum framework has clear potential for modelling the resource economics
of other social insects who likewise invest in their non-reproductive workers to bring in
resources that promote colony fitness. Workers of social insect colonies gather resources
across vast scales and have major impacts on habitats across our planet. Globally, ants can
increase crop yields through their positive role on soil health (*12*), managed bee colonies
pollinate one third of our crops (*54*) and termites decompose more than half of the deadwood
in tropical forests (*9*). In plants, aggregated economic spectrum trait values of communities,
together with biotic species interactions, correlate with ecosystem productivity, directly
linking the scaling of resource economics from leaves to ecosystems (*55*). Mapping social
insects along the fast-slow worker economic spectrum could be similarly used to understand
the link between rates of resource use and the allocation of energy and nutrients from colony
to community scales. Economic trait values of local assemblages could further be combined
with data on ecological roles to estimate the rates of ecosystem services such as
decomposition or pollination (*51*).

A major goal of ecology is to identify broadly applicable principles that provide a
mechanistic understanding of how living things behave and interact with their environment.
Our findings show that evolutionary optimisation and the biophysical laws of energy and
mass balance have jointly shaped the economics of superorganisms such as ants in similar
ways to plants indicating common solutions to resource allocation problems across the tree of
life.

References

1. K. J. Mathot, W. E. Frankenhuis, Models of pace-of-life syndromes (POLS): a systematic review. *Behavioral Ecology and Sociobiology* **72**, 41 (2018).
2. J. R. Burger, C. Hou, C. A. Hall, J. H. Brown, Universal rules of life: metabolic rates, biological times and the equal fitness paradigm. *Ecology Letters* **24**, 1262-1281 (2021).
3. S. A. L. M. Kooijman, *Dynamic energy budget theory for metabolic organisation*. (Cambridge university press, 2010).
4. C. Zhang, I. J. Wright, U. N. Nielsen, S. Geisen, M. Liu, Linking nematodes and ecosystem function: a trait-based framework. *Trends in Ecology & Evolution*, (2024).
5. H. Gibb *et al.*, Ecological strategies of (pl)ants: Towards a world-wide worker economic spectrum for ants. *Functional Ecology* **37**, 13-25 (2023).
6. R. R. Junker *et al.*, Towards an animal economics spectrum for ecosystem research. *Functional Ecology* **37**, 57-72 (2022).
7. I. J. Wright *et al.*, The worldwide leaf economics spectrum. *Nature* **428**, 821-827 (2004).
8. J. J. Boomsma, R. Gawne, Superorganismality and caste differentiation as points of no return: how the major evolutionary transitions were lost in translation. *Biological Reviews* **93**, 28-54 (2018).
9. H. M. Griffiths, L. A. Ashton, T. A. Evans, C. L. Parr, P. Eggleton, Termites can decompose more than half of deadwood in tropical rainforest. *Current Biology* **29**, 118-119 (2019).
10. H. M. Griffiths *et al.*, Ants are the major agents of resource removal from tropical rainforests. *Journal of Animal Ecology* **87**, 293-300 (2018).
11. A.-M. Klein *et al.*, Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences* **274**, 303-313 (2007).
12. D. Wu, E. Du, N. Eisenhauer, J. Mathieu, C. Chu, Global engineering effects of soil invertebrates on ecosystem functions. *Nature*, (2025).
13. J. M. Kass *et al.*, The global distribution of known and undiscovered ant biodiversity. *Science Advances* **8**, eabp9908 (2022).
14. S. Díaz *et al.*, The global spectrum of plant form and function. *Nature* **529**, 167-171 (2016).
15. W. S. Creighton, The ants of North America. *Bulletin of the Museum of Comparative Zoology* **104**, 1-585 (1950).
16. W. M. Wheeler, *Ants: their structure, development and behavior*. (Columbia University Press, New York, 1910), pp. 663.
17. G. Oster, E. Wilson, *Caste and ecology in the social insects*. (Princteon University Press, Princeton, NJ, 1978), pp. 352.
18. P. Calabi, S. D. Porter, Worker longevity in the fire ant Solenopsis invicta: ergonomic considerations of correlations between temperature, size and metabolic rates. *Journal of Insect Physiology* **35**, 643-649 (1989).
19. M. Kaspari, M. M. Byrne, Caste allocation in litter Pheidole: lessons from plant defense theory. *Behavioral Ecology and Sociobiology* **37**, 255-263 (1995).
20. J. H. Brown, J. F. Gillooly, A. P. Allen, V. M. Savage, G. B. West, Toward a metabolic theory of ecology. *Ecology* **85**, 1771-1789 (2004).
21. C. Peeters, M. Molet, C.-C. Lin, J. Billen, Evolution of cheaper workers in ants: a comparative study of exoskeleton thickness. *Biological Journal of the Linnean Society* **121**, 556-563 (2017).
22. J. J. Elser *et al.*, Biological stoichiometry from genes to ecosystems. *Ecology Letters* **3**, 540-550 (2000).
23. K. Kikuzawa, A cost-benefit analysis of leaf habit and leaf longevity of trees and their geographical pattern. *The American Naturalist* **138**, 1250-1263 (1991).
24. B. H. Kramer, G. S. van Doorn, B. M. S. Arani, I. D. Pen, Eusociality and the Evolution of Aging in Superorganisms. *American Naturalist* **200**, 63-80 (2022).
25. M. Kaspari, Global energy gradients and size in colonial organisms: worker mass and worker number in ant colonies. *Proceedings of the National Academy of Sciences* **102**, 5079-5083 (2005).
26. J. J. Elser *et al.*, Growth rate–stoichiometry couplings in diverse biota. *Ecology Letters* **6**, 936-943 (2003).
27. J. F. Gillooly *et al.*, The metabolic basis of whole-organism RNA and phosphorus content. *Proceedings of the National Academy of Sciences* **102**, 11923-11927 (2005).
28. P. A. Marquet *et al.*, Scaling and power-laws in ecological systems. *Journal of Experimental Biology* **208**, 1749-1769 (2005).
29. L. Keller, M. Genoud, Extraordinary lifespans in ants: a test of evolutionary theories of ageing. *Nature* **389**, 958-960 (1997).

30. N. J. Lemanski, N. H. Fefferman, How life history shapes optimal patterns of senescence: implications
from individuals to societies. *The American Naturalist* **191**, 756-766 (2018).
31. E. R. Pianka, On r-and K-selection. *The american naturalist* **104**, 592-597 (1970).
32. Bourke, Colony size, social complexity and reproductive conflict in social insects. *Journal of*
Evolutionary Biology **12**, 245-257 (1999).
33. M. Westoby, L. Yates, B. Holland, B. Halliwell, Phylogenetically conservative trait correlation:
Quantification and interpretation. *Journal of Ecology* **111**, 2105-2117 (2023).
34. B. Halliwell, B. R. Holland, L. A. Yates, Multi-response phylogenetic mixed models: concepts and
application. *Biological Reviews* **100**, 1294-1316 (2025).
35. K. S. Mason, C. L. Kwapich, W. R. Tschinkel, Respiration, worker body size, tempo and activity in
whole colonies of ants. *Physiological Entomology* **40**, 149-165 (2015).
36. H. Wang *et al.*, Leaf economics fundamentals explained by optimality principles. *Science Advances* **9**,
eadd5667 (2023).
37. M. Castorena, M. E. Olson, B. J. Enquist, A. Fajardo, Toward a general theory of plant carbon
economics. *Trends in Ecology & Evolution* **37**, 829-837 (2022).
38. J. Z. Shik, C. Hou, A. Kay, M. Kaspari, J. F. Gillooly, Towards a general life-history model of the
superorganism: predicting the survival, growth and reproduction of ant societies. *Biology Letters* **8**,
1059-1062 (2012).
39. J. L. Maino, M. R. Kearney, R. M. Nisbet, S. A. Kooijman, Reconciling theories for metabolic scaling.
Journal of Animal Ecology **83**, 20-29 (2014).
40. B. D. Blanchard, C. S. Moreau, Defensive traits exhibit an evolutionary trade-off and drive
diversification in ants. *Evolution* **71**, 315-328 (2017).
41. J. T. Buxton, K. A. Robert, A. T. Marshall, T. L. Dutka, H. Gibb, A cross-species test of the function of
cuticular traits in ants (Hymenoptera: Formicidae). *Myrmecological News* **31**, 31-46 (2021).
42. C. Hou, M. Kaspari, H. B. Vander Zanden, J. F. Gillooly, Energetic basis of colonial living in social
insects. *Proceedings of the National Academy of Sciences* **107**, 3634-3638 (2010).
43. J. Vizueta *et al.*, Adaptive radiation and social evolution of the ants. *Cell*, (2025).
44. J. Z. Shik, X. Arnan, C. S. Oms, X. Cerdá, R. Boulay, Evidence for locally adaptive metabolic rates
among ant populations along an elevational gradient. *Journal of Animal Ecology* **88**, 1240-1249 (2019).
45. L. Leahy, B. R. Scheffers, S. E. Williams, A. N. Andersen, Arboreality drives heat tolerance while
elevation drives cold tolerance in tropical rainforest ants. *Ecology* **103**, e03549 (2022).
46. M. Kaspari, N. A. Clay, J. Lucas, S. P. Yanoviak, A. Kay, Thermal adaptation generates a diversity of
thermal limits in a rainforest ant community. *Global Change Biology* **21**, 1092-1102 (2015).
47. Ü. Niinemets, Global-scale climatic controls of leaf dry mass per area, density, and thickness in trees
and shrubs. *Ecology* **82**, 453-469 (2001).
48. A. S. Yang, C. H. Martin, H. F. Nijhout, Geographic variation of caste structure among ant
populations. *Current Biology* **14**, 514-519 (2004).
49. A. D. Kay, S. Rostampour, R. W. Sterner, Ant Stoichiometry: Elemental Homeostasis in Stage-
Structured Colonies. *Functional Ecology* **20**, 1037-1044 (2006).
50. J. Bujan, S. J. Wright, M. Kaspari, Biogeochemical drivers of Neotropical ant activity and diversity.
Ecosphere **7**, e01597 (2016).
51. F. de Bello *et al.*, Raunkiæran shortfalls: Challenges and perspectives in trait-based ecology.
Ecological monographs **95**, e70018 (2025).
52. H. Gibb *et al.*, Habitat disturbance selects against both small and large species across varying climates.
Ecography **41**, 1184-1193 (2018).
53. C. S. Moreau, C. D. Bell, R. Vila, S. B. Archibald, N. E. Pierce, Phylogeny of the Ants: Diversification
in the Age of Angiosperms. *Science* **312**, 101-104 (2006).
54. FAOSTAT, ProdSTAT Database. 2022 (https://www.fao.org/faostat/en/#data/QCL Accessed 28-09-
2022).
55. P. B. Reich, The world-wide 'fast–slow' plant economics spectrum: a traits manifesto. *Journal of*
ecology **102**, 275-301 (2014).
385

Acknowledgements: We thank and acknowledge the Wurundjeri people of the Kulin Nation, the
Wotjobaluk, the Ngiyampaa, the Durramurragal, the Gubbi Gubbi, the Boonwurung, Bunurong, and
Gunaikurnai people on whose lands this field work was conducted. We thank Emily House for access
to Glen Echo. Field work was carried out under Permit SL102675 NSW Department of Planning,
Industry and Environment, Permit AA-0000328 Parks Victoria, and under animal ethics committee
permit AEC22001. We thank Luke A. Yates for advice and support with statistical modelling. We

thank Ian J. Hammer for technical assistance with metabolic rate assays. We thank Wednesday Elgar,
Stephanie Nicholls, Andrew Chalmers, Hannah Elmes, Nicholas Gale, Lia Herrera Grau and
volunteers for helping with field and lab work.

Funding:
Australian Research Council Discovery Project DP210101630 (HG, IJW, NJS, CLP, TRB)

Australian Government RTPS PhD Funding (HLR)

Australian Research Council Discovery Project DP190100341 (SLC)

Human Frontiers of Science Program grant RGEC26/2024 (TRB)

The Royal Society RGS\R1\221207 (TRB)

Fig. 1: Conceptual framework of the hypothesised relationships between eight ant worker traits that together place species along a fast-slow economic spectrum. Predicted positive trait correlations shown by blue, predicted negative trait correlations shown in red. Four key trait trade-offs and their hypothesised relationships are highlighted and explained in text boxes H1, H2, H3, H4 with reference to underlying bodies of theory (see reference list: (*3, 20-23, 26, 29-32*)). High trait values of body mass, lifespan, mass density, and nitrogen concentration and low trait values for the other four traits are predicted to indicate a slow strategy. High trait values of mass-specific metabolic rate, mass-specific assimilation (resource intake) rate, colony size (worker number), and phosphorus concentration and low trait values for the other four traits are predicted to indicate a fast strategy. Species positions along the superorganism economic spectrum indicate the pace of investment and return of resources into ant workers and back to the colony. Species positions may be modulated by the abiotic and biotic environment, for example, ambient temperature could shift communities towards fast ecological strategies.

Fig. 2: Two sets of correlated traits trade-off to create a fast-slow economic spectrum. (A)
Posterior summaries of phylogenetic correlation coefficients from MR-PMM analysis, showing the
posterior mean (point), 50% (heavy wick) and 95% (light wick) credible intervals. (B) Network
diagram showing phylogenetic trait correlations significant at 95% (bold) and 70-90% (faded)
credible interval. Blue lines = positive relationships, red lines = negative relationships. Line width
indicates strength of correlation. Phylogenetic correlations are estimated with greater uncertainty than
standard between-species correlations (*34*), meaning 95% CI are likely to be conservative. Despite
some long-tailed posterior distributions, the absolute posterior mean estimates for each correlation
coefficients were all >0.4, identifying an integrated trait network that strongly supports a fast-slow
economic spectrum. A_{mass} = mass-specific resource assimilation, MR_{mass} = mass-specific metabolic
rate.

Fig. 3: Mapping the superorganism economic spectrum onto multidimensional space. Showing
varimax-rotated PCA from gap-filled species-averaged trait data from 305 colonies, with the 123
species as points and coloured as subfamily groups. Percentage variation explained in brackets for
axis 1 (ant worker economic spectrum - AWES) and axis 2 (stoichiometry spectrum - SS). Size of
points represents dry body mass scaled between 1-10. Ant icons are representative genera from each
subfamily, anticlockwise from top left: Formicinae = *Camponotus*, Ponerinae = *Anochetus*,
Myrmeciinae = *Myrmecia*, Ectatomminae = *Rhytidoponera*, Dorylinae = *Lioponera*, Myrmicinae =
Meranoplus, Dolichoderinae = *Iridomyrmex*.

Fig. 4: Mapping the superorganism economic spectrum onto the ant phylogenetic tree at the genus level. Showing PCA scores from Fig. 3. For axis 1: Ant Worker Economic Spectrum (AWES) and for axis 2: Stoichiometry Spectrum (SS). Slow values (darker colours) for AWES signify large body mass, long lifespan, high mass density, low mass-specific metabolic and assimilation rates and small colony sizes. Fast values (lighter colours) represent the opposite values for each trait. Slow values (darker colours) for stoichiometry spectrum (SS) represent high nitrogen concentration and low phosphorus concentration, with the opposite ratio for fast ecological strategies (lighter colours). This demonstrates that niche partitioning is occurring along two axes of the ant economic spectrum (see PCA Fig. 3 to see these patterns mapped onto multidimensional space, see fig. S6 for subfamily tree). Note, genus *Arnoldius* scored very high on the SS spectrum and is removed in this figure to improve visualization of colour gradient, see fig. S7 for figure that includes *Arnoldius*.

456