Revealing year-round activity of cave-dwelling insectivorous bats with a sonotype classifier in data-deficient areas.

Morgane Labadie*[1,2], Serge Morand[3,4,5], Alexandre Caron[1,2,6], Helene Marie De Nys[1,2,7], Fabien Roch Niama[8], Franel Nguilili[9], N'Kaya Tobi[9], Mathieu Bourgarel[1,2,6], Yves Bas[10,11], Charlotte Roemer[10,11]

[1] CIRAD, UMR ASTRE, F-34398 Montpellier, France,

[2] ASTRE, CIRAD, INRAE, Univ Montpellier, Montpellier, France

[3] IRL HealthDEEP CNRS – Kasetsart University – Mahidol University, Bangkok, Thailand

[4] Faculty of Veterinary Technology, Kasetsart University, Bangkok, Thailand

[5] Department of Social and Environmental medicine, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand

[6] Faculdade de Veterinaria, Universidade Eduardo Mondlane, Maputo, Mozambique

[7] CIRAD, UMR ASTRE, Harare, Zimbabwe

[8] Laboratoire National de Santé Publique, Brazzaville, République du Congo

[9] Direction Générale de l'Élevage (Service vétérinaire), Ministère de l'Agriculture, de l'élevage et de la pêche, Brazzaville, République du Congo

[10] Centre d'Ecologie et des Sciences de la Conservation (CESCO), Muséum National d'Histoire Naturelle, Centre National de la Recherche Scientifique, Sorbonne Université, Paris, France

[11] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

*Corresponding author
Correspondence: morgane.labadie@gmail.com

ABSTRACT

Context: Bats are the only mammals who have conquered the skies and the leading mammals
in terms of total biomass with the exception of humans and domestic mammals. However, the
ecology of many bat species is still unknown, especially in poorly sampled regions such as
Central Africa.

Aims: We present the first application of an acoustic method for studying the nocturnal
activity patterns and annual phenology of bat activity of cave bat communities in regions
where no bat acoustic classifier exists.

Methods: In two caves (Mont Belo and Boundou) in the Republic of Congo, which are home
to several species of insectivorous bats throughout the year, we set up a passive acoustic
monitoring (PAM) over a period of 19 months. To identify the massive recordings collected,
we used a sonotype classifier. To assign a species name or species group to the different
sonotypes identified by the classifier, we collected local reference calls by capturing bats at
the cave entrances. This enabled us to identify two bat groups and three species despite the
absence of local call libraries. The capture sessions also allowed us to collect information
about the biological status of the individuals, which we put in perspective with the annual
phenology of activity. To quantify bat activity, we defined the activity of each acoustic group
as the number of bat passes, where a bat pass corresponds to the detection of one or several
echolocation calls within a five-second interval. This standardized measure allowed us to
quantify nightly and seasonal variations in activity.

Key results: The nightly activity displayed strong peaks in the evening and the morning,
except during the long rainy season, when a tri- or quadri-modal activity was detected. This
more evenly distributed activity throughout the night is related to the period of juvenile
rearing (gestation, parturition and lactation). For all acoustic groups, we also observed a
higher activity during the rainy seasons and the short dry season, compared to the long dry
season. This high activity is linked to the lactation period, the independence of young bats
and potentially the mating period. Finally, the reproductive activity of insectivorous bat
species is synchronized with the period of high resource availability.

Conclusion : This study represents the first long-term acoustic monitoring of cave bats in the
Republic of Congo. This was allowed by the use of a classifier built without any pre-existing
regional call data, highlighting a practical pathway for studying bat activity in other data-
deficient regions.

Implications : Information about the biological status of individuals captured at the monitored
caves in association with the cave annual activity phenology produced with this new method
are key to progress in the understanding of the ecology and the conservation issues of cave
bat communities, which are particularly threatened globally.

Keywords *:* PAM, Tadarida, activity pattern, abundance, bats phenology, Congo

Introduction

Human activities are currently triggering a massive loss of biodiversity, with rates of species
decline or extinction unprecedented for several millennia (Barnosky et al., 2011; Ceballos et al.,
2015; IPBES, 2019). However, the scale of this crisis is probably still concealed by the lack of
data on the populations of many species around the world (Hughes, 2017). Chiropterans are one
of these taxa, as information on their populations is still poor despite the many threats they face
(Furey & Racey, 2016; Mickleburgh et al., 2002).

Of the 1,500 recognised bat species worldwide, 38% are cave-dwelling species (IUCN, 2023;
Simmons & Cirranello, 2025, Tanalgo et al., 2022), living in caves or natural or man-made
underground cavities (e.g. mines or tunnels). These habitats are crucial for bats, which use them
to reproduce, raise their young, rest on a daily or temporary basis and hibernate (Barros et al.,
2020; Keeley & Tuttle, 1999; Kunz, 1982; Kunz et al., 2003; Lewis, 1995; Mering & Chambers,
2014; Ormsbee et al., 2007; Russo & Ancillotto, 2015; Struebig et al., 2009; The Protection of
Bat Roost Guidelines Subcommittee et al., 1992). In tropical environments, most cave-dwelling
species use caves as permanent roosts and several species can cohabit together by hundreds or
thousands (Kunz, 1982; Monadjem et al., 2020). However, despite their ecological importance,
these fragile environments are threatened and still poorly protected on a global scale (Balogh et
al., 2020; Medellin et al., 2017; Neuhuber et al., 2020). They are exposed to resources extraction
(minerals, guano, etc.) or recreational activities (caving, tourism, etc.) (Okonkwo et al., 2017;
Simons, 1998). In return, these case dwelling bats create through the resource they produce (e.g.,
live and dead-bats, guano) niches for cave-dwelling communities (Labadie et al. 2025a).

Deleva et al. (2023) highlight the need to identify and protect key underground roosts. Broader
global syntheses emphasize that effective conservation of bats roosting underground requires
coordinated protection, monitoring, and policy measures (Meierhofer et al., 2024). Recent tools
such as the bat cave vulnerability index provide integrative frameworks for assessing and ranking
cave vulnerability at multiple scales (Tanalgo & Hughes, 2024). Unfortunately, due to largescale
underfunding for bat research and conservation, the ecological information available on cave-
dwelling bats is often too limited to inform these tools. In particular, data on bat populations and
annual phenology of the occupation of caves is lacking, especially in African countries
(Moutaouakil et al., 2024; Pretorius et al., 2021). Understanding when and how bats use caves
throughout the year is therefore fundamental to assigning appropriate conservation value to each
site. For example, a cave that hosts colonies during key reproductive periods (e.g. parturition or
lactation) contributes critically to population persistence and should thus receive a higher
conservation rank than a site used only for transient or non-reproductive activities.

The annual phenology of species at a given site is the result of the adaptation of its activity
patterns to meet three primary needs (feeding, reproduction or rest) (Alcock, 2013; Appel et al.,
2019; Jones & Rydell, 1994). The temporal phenology of bat activity in tropical caves reflects
reproductive cycles adapted to local environmental conditions. In African regions, the timing of
parturition in cave-dwelling insectivorous bats is closely linked to rainfall and subsequent insect
availability (Cumming & Bernard, 1997). Insect abundance peaks about a month after peak
rainfall, while births occur earlier to ensure weaning before maximum prey availability
(Cumming & Bernard, 1997; Happold & Happold, 1990). This synchrony, though variable
among species, highlights strong selective pressures imposed by seasonal food dynamics. Such
cave-associated activity patterns are especially critical during maternity, when energetic demands

are highest, and their alteration may signal adaptation to changing microclimatic or trophic
conditions (Nkoana, 2020; Wolkovich et al., 2014).

Historically, bat communities have been studied using both invasive and non-invasive methods.
Invasive methods such as bat capture remain often necessary to make an exhaustive species
inventory, verify individual reproductive status and collect biometric data or samples (Appel et
al., 2021; Barnhart & Gillam, 2014; Flaquer et al., 2007). Direct observation of individuals in
their roost (e.g., cave, tree) or when they emerge to forage have also provided information on
their behavior, abundance and movements (Thomas & West, 1989). Visual counts are sometimes
difficult because of the large number of bats or the difficulty accessing the roosting sites. In
addition, human observations are time-consuming and bring observer bias (Forsyth et al., 2006;
Koger et al., 2023; Westcott & McKeown, 2004).

The development of new technologies have made it possible to set up precise counts of
individuals in caves using digital image analysis, thermal cameras (Azmy et al., 2012; Sabol &
Hudson, 1995) or bioacoustics (Sugai et al., 2020; Whiting et al., 2021). Passive acoustic
monitoring (PAM) has a number of advantages: it is non-invasive, applicable in all environments
(even the most remote ones), can collect a large amount of information such as abundance
proxies, and the analysis can be automated (Hill et al., 2019). PAM exploits the use of
echolocation to move and forage by more than 70% of the world's bats (Browning et al., 2017;
Gibb et al., 2019), Pteropodidae excluded. Recording these often species-specific sounds makes it
possible to make a quantitative monitoring of bat species and/or communities. In recent years,
several protocols, including that of Revilla-Martin et al (2020), have demonstrated the reliability
of this technique compared to other methods for counting bat populations as they leave caves.
Recent work by Appel et al. (2025) demonstrated that combining active and passive monitoring
and optimally placing recorders at cave entrances maximizes species detection and comparability
across sites and seasons.

However, PAM also faces limitations, such as the difficulty to distinguish between overlapping
calls. This issue is compounded by the lack of reference call libraries in many regions, especially
in Central Africa, and thus the impossibility to identify species in massive recordings. Sonotype
classifiers were recently developed to counter this issue (Roemer et al., 2019) and opened the
possibility to classify bat sounds in shapes and frequency groups where no species classifier is
available. Labadie et al. (2025b) proposed a method to apply a sonotype classifier in association
with local reference calls collected at cave entrances to obtain sound identification at the species
level, or acoustic group, in massive recordings in the Republic of Congo. Yet, no application of
this method was published so far.

The aim of this study was to demonstrate the possibility to conduct quantitative studies of cave-
dwelling bat communities using acoustics in data-deficient areas. The Republic of Congo has
very limited information on the acoustic repertoire of bats (Bates et al., 2013). Over a period of
19 months, we set up a PAM to describe the annual phenology of bat activity in two caves. We
used the Tadarida sonotype classifier to classify bat vocalizations. To associate each sonotype
with a species, we performed captures to obtain a species list and record reference calls,
following the approach of Labadie et al. (2025b, c). To progress in the understanding of the
biology and the ecology of bats in Central Africa, we compared seasonal variation in activity with
the reproductive status of bats assessed from capture data. We expected species to show higher
activity during periods of breeding and juvenile rearing (Catto et al., 1995; Dietz & Kalko, 2006).

Methods

Study area

Our study took place in the Republic of Congo, in a highly karstic region favouring the presence
of numerous caves and/or cavities of various sizes in a rolling habitat of herbaceous savannah
with more or less important patches of tropical forest.

Boundou and Mont Belo caves, are located in the Bouenza and Niari departments, about 50 km
apart. Both caves are close to Dolisie, one of the three most important towns in the Republic of
Congo on the southbound Brazzaville - Pointe Noire route. In this region, the seasons are divided
into four: the short dry season (January to February), the short rainy season (March to May), the
long dry season (June to mid-September) and the long rainy season (mid-September to
December).

The Mont Belo Cave has several chambers (a main chamber and two secondary chambers). The
Boundou Cave, on the other hand, is a smaller cavity integrated into a rocky complex, with a
main chamber and an annex chamber that is very difficult for humans to access, and houses a
small internal lake.

Details of the configuration of the two caves in this study are presented in Fig. 1.

Figure 1 - General diagram of the configuration of the Boundou and Mont Belo Cave, the location of the acoustic recorder and the bat roosting sites.

Ethic statements

The protocols presented below have all been validated by the Ethics Committee of the Ministry of
Scientific Research and Technological Innovation and by the Ministry of Forest Economy of the
Republic of Congo (N°212/MRSIT/IRSSA/CERSSA and N°687/MEF/CAB/DGEF-DFAP). This
work was also carried out with the agreement and participation of the authorities in the Niari and
Bouenza departments, as well as the communities and landowners close to our study sites.

Capturing bats

During the 19 months of data collection, eleven capture sessions using a harp trap were set up at
the main entrance of each cave, in order to identify the different bat species. During these capture
sessions, a wide range of information was collected, including the time of capture, the family and
species identified in the field, sex, age and reproductive status. The reproductive status of
individuals was determined by assessing females for pregnancy (abdominal palpation) and
lactation (mammary gland size, absence of fur, presence of milk at the nipple). Males were
assessed for sexual activity based on testes size and epididymal coloration. At the beginning of
the study, in order to confirm the genera and species of bats present in the caves, we collected
genetic samples, acoustic recordings and specimens for reference purposes.

Passive acoustic monitoring (PAM)

At each main entrance of the two caves, at a maximum distance of five meters, we placed an
SM4BAT acoustic recorder with a U1 microphone (Wildlife acoustics, Maynard, MA, USA).
Each microphone was placed on a pole or tree and oriented to best record the acoustic calls of
bats emerging and re-entering through the main entrance. These recorders ran continuously
several weeks per month for a period of 19 months (September 2021 to first day of March 2023).
The SM4BATs were set up according to the stationary points protocol of Vigie-chiro, a citizen
science programme for the acoustic monitoring of bat populations in France (French Museum of
Natural History, see (Mariton et al., 2023), starting 30 minutes before sunset and 30 minutes after
sunrise. Figure 1 show the location of the acoustic recorder in relation to the configuration of the
study site.

Processing acoustic data

Recordings were cut in durations of five seconds maximum. In order to process all the acoustic
files collected during this study in the two caves, we used the automatic sonotype classifier
published in Roemer et al., (2021) and modified in Labadie et al. (2025b) to identify the presence
of different sonotypes inside each file. The classifier returns a table in which each line
corresponds to a group of calls with similar call shape and frequency, associated with information
about their acoustic parameters (e.g., frequency, duration). Because of an overlap in the shape and
frequency of certain acoustic calls from bat species of the same family, we decided to group the
species at each study site using a posteriori classification (detail for the different acoustic groups
in Labadie et al., 2025b). The Boundou cave has four distinct groups: (1) the *Miniopterus* group
(*Miniopterus cf minor/ Miniopterus cf. inflatus*), (2) *Hipposideros caffer* (*Hipposideros cf.
caffer*), (3) *Rhinolophus denti* (*Rhinolophus cf. denti*) and (4) the *Rhinolophus* - Macro - Tria
group (*R. cf. landeri-alcyone /Macronycteris gigas/Triaenops cf. afer*) (Fig. 2). The Mont Belo
cave also has four distinct groups: (1) the *Miniopterus* group (*M. cf minor/ M. cf. inflatus*), (2)
Hipposideros caffer (*Hipposideros cf. caffer*), (3) *Rhinolophus denti* (*Rhinolophus cf. denti)* and

(4) *Rhinolophus landeri-alcyone (Rhinolophus cf. landeri-alcyone)* (Fig. 2). For more details on
the automatic classification and on a posteriori sorting of acoustic groups, please respectively
refer to Roemer et al. (2021) and Labadie et al. (2025b).

Statistical analysis

Post-sorting of the Tadarida output which represents the raw results of the acoustic classifier after
analyzing all recorded files. All the statistical analyses presented below were carried out using R
software v. 4.3.1 (R Core Team, 2023). The graphics were produced using ggplot2 (Wickham,
2016) or ggradar (Bion, 2023).

Bat counting techniques and acoustics analyses

Bats may emerge from and enter a cave at any time during the night; nonetheless, most of them
emerge after sunset and enter before sunrise (Erkert, 1982). In order to assess the activity of the
five groups of bats during emergence and their entry at each cave during the 19 months, we
decided to standardise our methodology based on the protocol of Mariton et al. (2023). As in
other acoustic studies (Millon et al., 2015), we defined the activity of the different groups of bats
as the number of bat passes. A bat pass is the occurrence of a single or several calls detected
during a five seconds interval.

As our study took place in the tropics and in two caves that are geographically close, the
characterisation of the diel activity of groups of bats can be based on a time scale (on sunrise and
sunset). To begin with, we defined the time of sunset and sunrise based on the geographical
coordinates and using the getSunlightTimes function in the suncalc package (Thieurmel et al.,
2022). We then quantified the number of bat passes, over two specific time periods: (a) the period
of emergence, from 30 minutes before sunset to two hours after sunset, and (b) the period of entry
into the cave, from two hours before sunrise to 30 minutes after sunrise. This calculation was
carried out for each species, each site and each day on which a complete night was recorded by
the SM4BATs.

Distribution of activity during the night

To calculate the distribution of the activity of acoustic groups of bats throughout the night, we
estimated the activity density using the density function in the R package. This calculation was
done for the twelve months of the year (some months were sampled repetitively during different
years) for each species and each study site.

Monthly pattern of the emergence and the entry activity at the roost

We used Generalised Linear Mixed Models (GLMM) to model the number of bat passes (proxy
of activity) during their emergence and entry as a function of the month of the year in each cave
and for each acoustic group.

We added month and year as a random effect (1|month_year). In total, we have eight different
models (see Appendix 1, p23). We used the glmmTMB package (Brooks et al., 2017) to build our
models. We used a lognormal distribution and checked the conditions and diagnostics using the
DHARMa package (QQ plot residuals, Residuals vs predicted, dispersion, zero-inflation) (Hartig

& Hartig, 2022). We used a Bonferroni correction to limit the type I error rate (false positive)
when interpreting generalised linear models (Cabin & Mitchell, 2000).

Results

Passive acoustic monitoring
In total, during nineteen months, we collected 2,375,956 acoustic files (five seconds duration) with 48.7% coming from Boundou and 51.3% from Mont Belo. The acoustic groups identified at each cave are shown in Figure 2 (Appendix 2, raw data for each group).

Nocturnal activity patterns
In both caves, for the five acoustic groups (Hipposideros caffer, Miniopterus group, Rhinolophus denti, Rhinolophus landeri-alcyone, Rhinolophus Macro Tria group), , nocturnal activity patterns—specifically emergence and return activity— were generally bi-modal. More precisely, the first peak of activity was at the beginning of the night around 6pm (after sunset) and another peak between 3am and 4am, depending on the species (Fig. 3 and Fig. 4).

In December, bat activity was more evenly distributed throughout the night than during the other months for all five acoustic groups and in both caves (Fig. 3 and Fig. 4). In addition, the Boundou cave activity was more spread-out during January and February (Fig. 3). In the Mont Belo cave, activity was also more spread out throughout the night in November and July (Fig. 4).

Annual phenology of bat activity and species-specific patterns predictions
Because analyses using emergence and entry activity produced similar annual phenology of bat activity, we decided to comment our results for the emergence activity only (Fig. 5) (Appendix 2, 4, 5 and 6 for entry activity prediction). The annual phenology of activity varied according to the acoustic group and the month (Appendix 2 and 3). For *Rhinolophus denti*, their low activity during emergence in Mont Belo cave meant that the model had no results (the model had too little data to run) (Appendix 5 and 6).

For the *Miniopterus* group, in both caves, predicted activity (including emergence and entry) showed an increase in activity during the short dry season (Jan, Feb; of which January is a copulation period) and the short rainy season (March, April, May; with March a period for rearing young). For this group, we observed a decline in activity during the long dry season (June, July, August; June was a period for copulation and rearing young, depending on the individual) and the long rainy season (September, October, November, December; period for rearing young) (Figure 5 and Appendix 7).

For *Hipposideros caffer*, in both caves we observed an increase in activity during the short dry season (January, February) and the short rainy season (March, April, May). In this group, these two periods are linked to the mating activity by certain individuals and the rearing of young. Secondly, *Hipposideros caffer* showed a sharp drop-in predicted activity during the long dry season, which stabilized during the long rainy season (end of the year and period of juvenile rearing) (Figure 5 and Appendix 7).

For Mont Belo cave, the predicted activity for *Rhinolophus landeri-alcyone* showed an increase in activity during the short dry season and the short rainy season. Then, during the long dry season, we observed a sharp drop-in activity, which rose again during the long rainy season

(the period when females are pregnant, give birth and nurse their young) (Figure 5 and Appendix
7).

For Boundou cave, the *Rhinolophus denti* showed a strong increase in predicted activity
during the long rainy season and the start of the short dry season (end of the year and start of the
new one; period of rearing young), and low activity during two seasons: the short rainy season
and the long dry season (no individuals captured with individuals rearing young or mating)
(Figure 5 and Appendix 7).

Finally, in Boundou cave, we observed a predictive activity for the *Rhinolophus* Macro Tria
group that increased during the short dry and rainy seasons (Jan, Feb, Mar, Apr) and decreased
during the long dry and rainy seasons (Figure 5 and Appendix 7).

Figure 2 - Acoustic characteristics for the different bat species captured at the two study sites and categorization into different acoustic groups or species. Detail of sonotype, sonogram and acoustic frequency recorded or listed in reference works for each bat genus and species captured in the two caves (species with * are present only at Boundou). In the Boundou Cave, there are four acoustic groups (*Rhinolophus* – Macro – Tria group; *Rhinolophus denti*; *Hipposideros caffer* and *Miniopterus* group) and in the Mont Belo Cave there are four acoustic groups (*Rhinolophus landeri-alcyone*; *Rhinolophus denti*; *Hipposideros caffer* and *Miniopterus* group).

Figure 3 - Number of bat passes for each acoustic group according to month at Boundou Cave. The reproductive status of the species or acoustic group is
indicated by two pictograms with the color of the acoustics groups: 🛒 young rearing (lactation, pregnant) and 💜 : mating period/male activity. The months
of May, July and August, which have an *, are month when no captures were made, hence the absence of data on the reproductive status of the acoustic group.

Figure 4 - Number of bat passes for each acoustic group according to month at Mont Belo Cave. The reproductive status of the species or acoustic group is indicated by two pictograms with the color of the acoustics groups: 🍼 young rearing (lactation, pregnant) and 💜 : mating period/male activity. The months of May, July and August, which have an *, are month when no captures were made, hence the absence of data on the reproductive status of the acoustic group.

Figure 5 - Prediction of acoustic group emergence activity with 95% confidence interval as a function of months and study sites (Boundou and Mont Belo Cave). The *Rhinolophus denti* group at Mont Belo has not been represented here due to lack of data. The reproductive status of the species or acoustic group is indicated by two pictograms with the color of the acoustics groups:

 young rearing (lactation, pregnant) and : mating period/male activity. The months of May, July and August, which have an *, are month when no captures were made, hence the absence of data on the reproductive status of the acoustic group.

Discussion

We used passive acoustic monitoring to characterize bat communities and their phenologies at the entrances of caves in the Republic of Congo, addressing key challenges in cave-dwelling bat research in data_deficient countries. A global sonotype classifier augmented with locally recorded reference calls allowed to overcome the main challenges of bat research in these contexts: the absence of pre-existing reference acoustic libraries; the difficulty of monitoring nocturnal and cryptic species in remote places; and the absence of standardized frameworks for quantifying bat activity.

In two caves, we quantified the activity of five acoustic groups of bats and described their nightly and annual phenologies. This methodological adaptation combining sonotypes with on-site reference recordings offers a framework that can be replicated in other regions lacking acoustic data, potentially across different biomes and bat communities.

Nocturnal activity pattern

Five acoustic groups were detected in both caves: *Hipposideros caffer*, the *Miniopterus* group, *Rhinolophus denti*, *Rhinolophus landeri-alcyone*, and the *Rhinolophus Macro Tria* group. Overall, their emergence and entry patterns showed similar dynamics, with two bi-modal nocturnal activity patterns (a period of emergence and a period of entry into the cave) for most months in both caves. In several months, however, nocturnal activity patterns was more evenly distributed throughout the night—particularly in December for all groups, in January–February at Boundou, and in November and July at Mont Belo. When outside the cave, the large number of individuals observed are hypothesized to go hunting before returning to the cave. Some individuals may move to another resting place or return later.

Nightly observations revealed patterns consistent with known ecological behaviors. With a few exceptions, bats emerged from their daytime roosts at dusk—around 6 p.m.—to fly to feeding sites, as expected (Bullock et al., 1987). The time of this nocturnal emergence was correlated with the time of sunset (Erkert, 1982). A second peak of nocturnal activity generally occurred during the early morning hours, around 3–4 a.m., corresponding to their return to the caves. Central African countries such as Congo have little variation in sunrise and sunset throughout the year (no more than half an hour) (WorldData, 2024), leading to little variation in emergence time across the year. In our study, we also observed "burst activity" patterns during emergence, where several bats exited the cave in rapid succession followed by short periods of reduced activity, a behavior previously described in other bat colonies (Swift, 1980). However, during some months (October – January), nightly activity patterns was more evenly distributed across the night, reflecting changes in foraging and maternal behavior. These nocturnal activity patterns provide the detailed temporal resolution necessary to interpret the annual phenology of bat activity, which is strongly shaped by reproductive cycles and resource availability.

Annual phenology of bat activity and reproductive period

Building on nocturnal activity patterns, we examined the annual phenology of bat activity across the year to understand seasonal trends. In our study, activity peaks during the main rainy season (October–December) coinciding with maternity colony formation (gestation, birth, and beginning of lactation). In December and January, emergence nocturnal activity was spread throughout the

night likely reflecting females alternating between foraging and suckling their young, as reported
in previous studies (Catto et al., 1995 ; Dietz & Kalko, 2006 ; Maier, 1992 ; Swift, 1980, 1997).
Field observations and the ecological knownledge of these species support this interpretation. We
therefore hypothesise that the activity spread throughout the night corresponds to females making
repeated trips between the cave and foraging sites to alternately nurse their offspring and feed.
This pattern is consistent with Henry et al. (2002), who showed that *Myotis lucifugus* reduced
their foraging range and returned to the roost one to two times per night. Such repeated returns
created secondary and/or tertiary nocturnal activity patterns several hours after the initial
emergence observed in our data.

Our results also show peaks during the short rainy season (March–May), likely reflecting the
independence of juveniles. The period when young reach their independence can vary according
to bat species. For example, young *Hipposideros caffer* become independent between three and
three and a half months after birth (Wright, 2009). However, the young are not carried by their
mother throughout this period, and may begin to fly one month after birth, staying in the roost
with other individuals of the same generation when their mothers forage out of the cave (Wright,
2009).

Finally, based on capture observations, we hypothesise that the annual phenology of bat activity
peaks observed during the dry season (January–February) correspond to the initiation of
reproductive cycles for certain species, such as *Hipposideros caffer* and the *Miniopterus group*.

Some acoustic groups in our study showed two reproductive periods within the year. For
example, the *Miniopterus* group exhibited peaks of reproductive activity in March–April and
again from September to December, while *Rhino Macro Tria group* showed similar patterns
between September–January and March–June. These patterns likely reflect the presence of
multiple species within each acoustic group. Each species may have distinct reproductive cycles.
In the case of *Hipposideros caffer*, the dual activity could be explained by the presence of a
cryptic species complex within this acoustic group, as suggested in previous studies (Wright,
2009). Alternatively, the observed extended annual phenology of bat activity could reflect a
somewhat flexible reproductive period. Although individual females may start reproduction at
slightly different times, parturition generally occurs during the peak of insect availability in the
rainy season. This partial synchrony allows offspring to benefit from abundant food resources
while maintaining flexibility in reproductive timing, which can maximize survival (Racey &
Entwistle, 2000; McWilliam, 1988). This flexible reproductive timing likely aligns annual
phenology of bat activity with seasonal resource peaks.

This methodological framework can thus link nocturnal activity patterns with ecological drivers
such as resource availability and rainfall cycle. The most favorable period in terms of resources is
the long rainy season (October - December) and the short rainy season (March - May) for
insectivorous bats (Fenton and Thomas, 1980). These wet periods favour the presence of a great
diversity and abundance of insects (Jenber & Wili, 2021; Wolda, 1988). In mammals, including
bats, this synchronization is important because of the high energy expenditure by lactating
females (Racey & Entwistle, 2000). In addition, the aggregation of individuals can be an
advantage, as females cooperate with each other to raise their young.

The long dry season (June–August) showed reduced nocturnal activity patterns compared with
the other seasons. This suggests that bats congregate less in these habitats due to limited
resources, a pattern that aligns with previous studies (Barclay & Kurta, 2007; Patriquin &

Ratcliffe, 2016; Patterson et al., 2007; Postawa & Gas, 2009). Scattered nocturnal activity
patterns were observed throughout the night during the long dry season (June - August).
Unfortunately, we did not capture in the months of July and August, and could not assess the
reproductive status of the individuals. However, the gestation period observed from late
September to early December in all acoustic groups suggests that copulation likely occurs
towards the end of the long dry season (June–July). This timing aligns with swarming behavior
and is consistent with observations from other countries. For example, *Hipposideros caffer*
typically mates at the end of the first rainy season (June–July) and gives birth during the middle
of the second rainy season (October–November) (Bernard, 1982; Brosset, 1968; Günther et al.,
2016).

The only exception to the general drop-in nocturnal activity patterns was the *Miniopterus* group
in Boundou cave, which remained very active in August (long dry season). The females of this
species may use the cave to gather in a maternity colony before giving birth in September and
October (obs. per ML). This would support Cumming and Bernard (1997) who mentioned that
many bat species give birth to their young about a month before the peak in resource availability.

Finally, variation in roost use by species and cave highlights the complexity of bat ecology in
these systems. *Rhinolophus denti* appeared to use caves solely for rearing young, not for mating.
Conversely, *Hipposideros caffer* used both caves for mating and rearing young. It is noteworthy
that the mating period of *Hipposideros caffer* seemed to vary between cave (January, February,
June in Mont Belo Cave and March and June in Boundou Cave). This variation in breeding
months could potentially be explained by the presence of two cryptic species of Hipposideros in
this study (obs. per ML).

Conservation implications for the caves and application in other biomes

By describing nocturnal activity patterns and annual phenology of bat activity, our results also
highlight the ecological and conservation significance of these caves. Although this study did not
aim to formally calculate the Bat Cave Vulnerability Index (BCVI) (from Tanalgo & Hughes,
2024), a quick assessment of standard criteria suggests differing vulnerability between the two
caves. Mont Belo hosts more species and several maternity colonies. It therefore has the highest
biotic value. Its vulnerability score must then be assessed by considering the different threats
affecting the site (Labadie et al., 2025a). Boundou shows lower vulnerability, with fewer
disturbances and only occasional predation (Labadie et al., 2025a).

Limitation of this study

Information on reproductive status cannot be obtained from acoustic monitoring, it requires the
capture of individuals. We were not able to capture during the long dry season (June–August),
which hampers our understanding of the annual phenology of bat activity of the different species.
Moreover, we did not have the opportunity to validate our acoustic monitoring method with
manual counts, video monitoring, or infrared cameras to verify the number of individuals in each
cave per month due to logistical constraints. These complementary methods would have provided
direct estimates of colony size and temporal fluctuations in bat abundance. Such information
would have allowed us to link nocturnal activity patterns more precisely to actual population
dynamics and to better distinguish between changes in the number of individuals. Acoustic
activity is, however, generally correlated with colony size in caves (Revilla-Martín et al., 2020).
Still, variations in flight behavior—such as increased vocal activity during social interactions or

reduced calling during certain foraging strategies—may lead to an over- or underestimation of
this correlation.

Conclusion

We characterized the variation in activity of acoustic groups in relation to season and breeding
status, providing insights into temporal use of roosts and potential sensitivity to disturbance.
While both caves host similar species groups and share general seasonal trends, Mont Belo shows
more distinct biannual reproductive peaks (especially for Miniopterus and R. landeri–alcyone),
whereas Boundou displays more evenly distributed activity and possibly more stable year-round
occupation (notably for *Hipposideros caffer* and the *Rhinolohus* Macro Tria group).
These patterns suggest functional complementarity between the two caves — Mont Belo cave
likely serving as a breeding and maternity site, and Boundou cave as a multi-species roost used
across seasons. These observations underscore the potential of PAM combined with acoustic
classification to study reproductive behavior in wild populations, a novel application in Central
African caves.

Our study demonstrates that quantitative bat monitoring is possible even in data-deficient areas.
This approach offers a valuable tool for cave conservation, as it enables the identification of
critical periods of bat activity, reproductive timing, and species-specific cave use. This
information is essential for prioritizing protection and managing disturbance of cave bats
(Tanalgo & Hughes, 2024). A major strength of this study lies in its non-invasive approach,
applicable across multiple caves and species, while obtaining both nocturnal activity patterns and
annual phenology of bat activity.

Future improvements could include non-invasive hormonal monitoring by collecting bat urine to
analyze stress levels and reproductive hormones, allowing identification of mating and lactation
periods without capturing bats (Greville et al., 2022; Kumar & Umapathy, 2019; Voigt &
Schwarzenberger, 2008; Wilkinson et al., 2024). Given the importance of bats for the health of
the environment across the world, this study provides new avenues to produce knowledge in data-
deficient and remote areas in order to adapt bat conservation and management.

Appendices

478 **Appendix 1**- Summary table of eight emergence models for each cave (Boundou and Mont Belo) and
479 each acoustics group tested in this study.
480

Model name	Description
Model Hipposideros caffer emergence Boundou Cave	Nb_batpass ~ Month + (1|month-year)
Model Miniopterus group emergence Boundou Cave	Nb_batpass ~ Month + (1|month-year)
Model Rhinolophus denti emergence Boundou Cave	Nb_batpass ~ Month + (1|month-year)
Model Rhinolophus landeri-alcyone emergence Boundou Cave	Nb_batpass ~ Month + (1|month-year)
Model Hipposideros caffer emergence Mont Belo Cave	Nb_batpass ~ Month + (1|month-year)
Model Miniopterus group emergence Mont Belo Cave	Nb_batpass ~ Month + (1|month-year)
Model Rhinolophus denti re-enter Mont Belo Cave	Nb_batpass ~ Month + (1|month-year)
Model Rhinolophus Macro Tria emergence Boundou Cave	Nb_batpass ~ Month + (1|month-year)

Appendix 2 - Number of bat passes for each acoustic group according to month and site (Boundou and
Mont Belo Cave). The reproductive status of the species or acoustic group is indicated by two

pictograms: young rearing (lactation, pregnant) and : mating period/male activity. The months of
May, July and August, which have an *, are month when no captures were made, hence the absence of
data on the reproductive status of the acoustic group.

Appendix 3 - Generalized linear mixed (GLM) with Bonferroni correction describing the relationship between activity and month for the
four acoustic groups in the Boundou Cave during the emergence time of bats.

Predictors	Hipposideros caffer emergence Boundou			Miniopterus group emergence Boundou			Rhinolophus denti emergence Boundou			Rhinolophus Macro Tria group emergence Boundou		
	Estimates	CI	p	Estimates	CI	p	Estimates	CI	p	Estimates	CI	p
Intercept Jan	250.45 ***	192.77 – 325.39	<0.001	522.93 ***	440.28 – 621.08	<0.001	6.71 ***	4.73 – 9.51	<0.001	661.78 ***	488.20 – 897.08	<0.001
Feb	0.68	0.47 – 1.00	0.593	0.96	0.77 – 1.19	1.000	0.68	0.43 – 1.06	1.000	0.81	0.53 – 1.24	1.000
Mar	0.66	0.45 – 0.96	0.382	0.86	0.69 – 1.08	1.000	0.72	0.43 – 1.21	1.000	0.83	0.54 – 1.26	1.000
Apr	0.58	0.37 – 0.89	0.166	0.76	0.61 – 0.95	0.191	0.69	0.44 – 1.07	1.000	0.88	0.54 – 1.45	1.000
May	0.47 *	0.30 – 0.74	**0.012**	0.97	0.78 – 1.20	1.000	0.61	0.36 – 1.03	0.750	0.91	0.55 – 1.48	1.000
Jun	0.31 ***	0.20 – 0.49	<0.001	0.71 *	0.57 – 0.89	**0.038**	0.58	0.37 – 0.91	0.203	0.72	0.44 – 1.19	1.000
Jul	0.22 ***	0.14 – 0.35	<0.001	0.57 ***	0.43 – 0.74	<0.001	0.60	0.33 – 1.08	1.000	0.37 **	0.21 – 0.63	**0.003**
Aug	0.24 ***	0.15 – 0.38	<0.001	0.97	0.79 – 1.21	1.000	0.59	0.34 – 1.01	0.671	0.83	0.50 – 1.36	1.000
Sep	0.26 ***	0.18 – 0.38	<0.001	0.62 ***	0.51 – 0.77	<0.001	0.64	0.41 – 0.99	0.557	0.58	0.38 – 0.89	0.137
Oct	0.30 ***	0.21 – 0.44	<0.001	0.37 ***	0.30 – 0.46	<0.001	1.10	0.76 – 1.58	1.000	0.45 **	0.29 – 0.68	**0.002**
Nov	0.41 **	0.27 – 0.64	**0.001**	0.31 ***	0.23 – 0.42	<0.001	1.75	1.19 – 2.59	0.056	0.94	0.61 – 1.46	1.000
Dec	0.61	0.41 – 0.90	0.166	0.61 ***	0.48 – 0.77	**0.001**	4.47 ***	3.13 – 6.38	<0.001	1.06	0.69 – 1.64	1.000
Random Effects												
σ²	58.54			192.82			7.76			191.71		
τ₀₀	0.03 month2			0.00 month2			0.00 month2			0.04 month2		
ICC	0.00									0.00		
N	19 month2			19 month2			19 month2			19 month2		
Observations	275			275			182			275		
Marginal R² / Conditional R²	0.003 / 0.004			0.001 / NA			0.047 / NA			0.001 / 0.001		
AICc	2867.068			3566.270			900.772			3695.560		

$* p<0.05 \quad ** p<0.01 \quad *** p<0.001$

490 **Appendix 4** - Generalized linear mixed (GLM) with Bonferroni correction describing the relationship between activity and month for the
491 four acoustic groups in the Mont Belo Cave during the emergence time of bats.

Predictors	Hipposideros caffer emergence Mont Belo			Miniopterus group emergence Mont Belo			Rhino landeri-alcyone emergence Mont Belo			Rhinolophus denti emergence Mont Belo		
	Estimates	CI	p	Estimates	CI	p	Estimates	CI	p	Estimates	CI	p
Intercept Jan	9.35 ***	6.82 – 12.82	**<0.001**	737.51 ***	636.15 – 855.01	**<0.001**	499.73 ***	346.52 – 720.68	**<0.001**	1.00	NaN – NaN	NaN
Feb	1.05	0.74 – 1.47	1.000	1.08	0.91 – 1.29	1.000	1.03	0.66 – 1.60	1.000			
Mar	1.34	0.91 – 1.98	1.000	1.15	0.96 – 1.39	1.000	1.14	0.72 – 1.79	1.000			
Apr	1.08	0.72 – 1.63	1.000	1.06	0.86 – 1.31	1.000	1.27	0.76 – 2.11	1.000			
May	1.14	0.79 – 1.64	1.000	1.03	0.85 – 1.26	1.000	1.12	0.68 – 1.86	1.000			
Jun	0.68	0.44 – 1.05	0.958	0.79	0.64 – 0.97	0.316	0.92	0.55 – 1.54	1.000			
Jul	0.83	0.55 – 1.24	1.000	0.74	0.60 – 0.91	0.064	0.85	0.51 – 1.42	1.000			
Aug	0.61	0.37 – 1.00	0.588	0.70 *	0.56 – 0.86	**0.012**	0.77	0.46 – 1.28	1.000			
Sep	0.76	0.52 – 1.11	1.000	0.73 *	0.60 – 0.88	**0.013**	0.63	0.40 – 0.99	0.542	1.00	NaN – NaN	NaN
Oct	0.83	0.58 – 1.18	1.000	1.05	0.88 – 1.25	1.000	1.06	0.68 – 1.65	1.000	1.00	NaN – NaN	NaN
Nov	0.90	0.61 – 1.33	1.000	0.99	0.82 – 1.19	1.000	1.00	0.63 – 1.57	1.000			
Dec	1.85 **	1.34 – 2.56	**0.003**	1.00	0.84 – 1.19	1.000	0.98	0.62 – 1.53	1.000			
Random Effects												
σ^2	13.88			152.17			169.52			0.00		
τ_{00}	0.00 month2			0.00 month2			0.03 month2			0.00 month2		
ICC				0.00			0.00					
N	18 month2			18 month2			18 month2			4 month2		
Observations	271			293			293			4		
Marginal R^2 / Conditional R^2	0.007 / NA			0.000 / 0.000			0.000 / 0.000			0.000 / NA		
AICc	1644.111			3796.908			3859.854			NA		

*p<0.05 **p<0.01 ***p<0.001

 Appendix 5 - Generalized linear mixed (GLM) with Bonferroni correction describing the relationship between activity and month for the
494 four acoustic groups in the Boundou Cave during the enter time of bats.

Predictors	Hipposideros caffer enter Boundou			Miniopterus group enter Boundou			Rhinolophus denti enter Boundou			Rhinolophus Macro Tria group enter Boundou		
	Estimates	CI	p	Estimates	CI	p	Estimates	CI	p	Estimates	CI	p
Intercept Jan	249.41 ***	156.62 – 397.20	**<0.001**	463.67 ***	314.74 – 683.07	**<0.001**	17.20 ***	11.53 – 25.65	**<0.001**	439.89 ***	289.90 – 667.50	**<0.001**
Feb	1.07	0.59 – 1.96	1.000	1.71	1.07 – 2.74	0.290	0.43 *	0.25 – 0.73	**0.020**	1.65	0.96 – 2.85	0.865
Mar	1.57	0.87 – 2.85	1.000	1.63	1.02 – 2.60	0.497	0.55	0.33 – 0.91	0.242	1.54	0.89 – 2.67	1.000
Apr	1.16	0.58 – 2.31	1.000	1.53	0.91 – 2.57	1.000	0.37 **	0.21 – 0.64	**0.005**	1.70	0.90 – 3.20	1.000
May	1.18	0.60 – 2.34	1.000	1.45	0.86 – 2.44	1.000	0.41 *	0.24 – 0.72	**0.021**	1.86	0.99 – 3.50	0.656
Jun	0.75	0.38 – 1.49	1.000	0.76	0.44 – 1.31	1.000	0.29 ***	0.16 – 0.50	**<0.001**	1.26	0.66 – 2.40	1.000
Jul	0.48	0.24 – 0.96	0.457	0.89	0.52 – 1.52	1.000	0.31 ***	0.17 – 0.55	**0.001**	1.50	0.80 – 2.84	1.000
Aug	0.46	0.22 – 0.93	0.377	1.36	0.81 – 2.30	1.000	0.36 **	0.20 – 0.65	**0.007**	1.40	0.74 – 2.65	1.000
Sep	0.49	0.27 – 0.89	0.229	1.17	0.74 – 1.87	1.000	0.29 ***	0.17 – 0.50	**<0.001**	1.02	0.59 – 1.76	1.000
Oct	0.67	0.37 – 1.21	1.000	1.19	0.75 – 1.89	1.000	0.48 *	0.30 – 0.77	**0.028**	0.83	0.48 – 1.44	1.000
Nov	0.69	0.37 – 1.31	1.000	1.12	0.68 – 1.83	1.000	1.15	0.72 – 1.82	1.000	1.39	0.79 – 2.44	1.000
Dec	1.00	0.54 – 1.84	1.000	1.38	0.85 – 2.23	1.000	2.80 ***	1.73 – 4.54	**<0.001**	2.01	1.16 – 3.47	0.154
Random Effects												
σ^2	155.58			202.81			9.54			229.51		
τ_{00}	0.06 month2			0.03 month2			0.02 month2			0.06 month2		
ICC	0.00			0.00			0.00			0.00		
N	19 month2			19 month2			19 month2			19 month2		
Observations	276			277			242			277		
Marginal R^2 / Conditional R^2	0.001 / 0.001			0.000 / 0.000			0.042 / 0.044			0.000 / 0.001		
AICc	3301.278			3769.686			1366.644			3790.904		

*p<0.05   **p<0.01   ***p<0.001

496 **Appendix 6 -** Generalized linear mixed (GLM) with Bonferroni correction describing the relationship between activity and month for the
497 four acoustic groups in the Mont Belo Cave during the enter time of bats.

Predictors	Hipposideros caffer enter Mont Belo			Miniopterus group enter Mont Belo			Rhino landeri-alcyone enter Mont Belo			Rhinolophus denti enter Mont Belo		
	Estimates	CI	p	Estimates	CI	p	Estimates	CI	p	Estimates	CI	p
Intercept Jan	20.98 ***	12.75 – 34.52	**<0.001**	570.35 ***	501.54 – 648.60	**<0.001**	663.65 ***	530.64 – 830.00	**<0.001**	1.17	0.47 – 2.93	1.000
Feb	1.30	0.72 – 2.33	1.000	1.22	1.05 – 1.41	0.086	1.05	0.80 – 1.38	1.000			
Mar	1.41	0.77 – 2.58	1.000	1.17	0.99 – 1.38	0.822	0.92	0.69 – 1.23	1.000			
Apr	1.08	0.54 – 2.15	1.000	1.27	1.07 – 1.50	0.063	1.19	0.87 – 1.62	1.000			
May	0.83	0.42 – 1.63	1.000	1.37 ***	1.18 – 1.59	**0.001**	1.09	0.80 – 1.49	1.000	1.00	0.32 – 3.13	1.000
Jun	0.54	0.27 – 1.08	0.988	1.22	1.04 – 1.43	0.153	0.87	0.64 – 1.19	1.000			
Jul	0.50	0.24 – 1.03	0.705	1.21	1.03 – 1.42	0.229	0.66	0.48 – 0.92	0.155			
Aug	0.38	0.18 – 0.78	0.103	0.94	0.79 – 1.11	1.000	0.52 ***	0.37 – 0.72	**0.001**			
Sep	0.52	0.28 – 0.96	0.453	0.95	0.81 – 1.11	1.000	0.61 **	0.46 – 0.81	**0.007**			
Oct	0.66	0.36 – 1.20	1.000	0.94	0.81 – 1.10	1.000	0.73	0.55 – 0.96	0.279			
Nov	0.64	0.34 – 1.20	1.000	0.79	0.66 – 0.94	0.110	0.63 *	0.47 – 0.84	**0.021**			
Dec	1.22	0.66 – 2.25	1.000	0.98	0.84 – 1.14	1.000	0.97	0.73 – 1.29	1.000	1.55	0.53 – 4.57	1.000
Random Effects												
σ^2	22.89			166.33			157.22					
τ_{00}	0.04 month2			0.00 month2			0.01 month2					
ICC	0.00						0.00					
N	18 month2			18 month2			18 month2			4 month2		
Observations	279			294			294			4		
Marginal R^2 / Conditional R^2	0.007 / 0.009			0.000 / NA			0.000 / 0.000			NA		
AICc	1995.258			3840.501			3818.869			-12.700		

*p<0.05   **p<0.01   ***p<0.001

Appendix 7: Prediction of the activity of acoustic group entries with 95% confidence interval as a function of months and study sites (Boundou and Mont Belo Cave). The Rhinolophus denti group at Mont Belo has not been represented here due to lack of data. The reproductive status of the species or acoustic group is indicated by two pictograms with the color of the acoustics groups: young rearing (lactation, pregnant) and : mating period/male activity. The months of May, July and August, which have an *, are month when no captures were made, hence the absence of data on the reproductive status of the acoustic group.

Acknowledgements

We would also like to extend our warmest thanks to Léa Mariton for her support and advice on the R scripts for the density of bat activity over time. Many thanks to our Congolese partners: the "Laboratoire National de Santé Publique de Brazzaville" (LNSP), the "Direction Générale de l'Elevage" (DGE) and "Ministère de l'économie forestière" for their support in this project, their help with the fieldwork and the applications for the capture and ethics permits. Thanks also to the bat field team who helped collect data: F. Nguilili, C. Bazola, N. Nguimbi (our wizard), R. Dimoukissi and R. Nguimbi. Finally, we would like to thank the local authorities in Dolisie and the local population for their help with this project.

Data, scripts, code, and supplementary information availability

The summary file of species and the scripts used to create the figures can be obtained by following this link: https://doi.org/10.5281/zenodo.15636144

Conflict of interest disclosure

The authors declare that they have no know competing financial interests or personal that could have appeared to influence the work reported in this paper.

Funding

This study and publication were supported by the European Union Delegation under the Agreement [FOOD/2016/379-660], signed between the EU and the WOAH for the implementation of the Action EBO-SURSY.

References

Alcock, J. (2013). Animal Behavior: An Evolutionary Approach (10th edition). Sinauer Associates.

Appel, G., da Rocha, P.A., Cerqueira, P.V. et al. (2025). Using active and passive methods to monitor cave-dwelling bats in Amazonian caves. Biodiversity Conservation, 34, 4429–4449 https://doi.org/10.1007/s10531-025-03160-4

Appel, G., Capaverde, U. D., De Oliveira, L. Q., do Amaral Pereira, L. G., da Cunha Tavares, V., López-Baucells, A., Magnusson, W. E., Baccaro, F. B., & Bobrowiec, P. E. (2021). Use of complementary methods to sample bats in the Amazon. Acta Chiropterologica, 23(2), 499–511. https://doi.org/10.3161/15081109ACC2021.23.2.017

Appel, G., López-Baucells, A., Magnusson, W. E., & Bobrowiec, P. E. D. (2019). Temperature, rainfall, and moonlight intensity effects on activity of tropical insectivorous bats. Journal of Mammalogy, 100(6), 1889–1900. https://doi.org/10.1093/jmammal/gyz140

Azmy, S. N., Sah, S. A. M., Shafie, N. J., Ariffin, A., Majid, Z., Ismail, M. N. A., & Shamsir, M. S. (2012). Counting in the dark: Non-intrusive laser scanning for population counting and identifying roosting bats. Scientific Reports, 2(1), 1–4.https://doi.org/10.1038/srep00524

Balogh, A., Ngo, L., Zigler, K. S., & Dixon, G. (2020). Population genomics in two cave-obligate invertebrates confirms extremely limited dispersal between caves. Scientific Reports, 10(1), 17554.https://doi.org/10.1038/s41598-020-74508-9

Barclay, R. M., & Kurta, A. (2007). Ecology and behavior of bats roosting in tree cavities and under bark. Bats in Forests: Conservation and Management (MJ Lacki, JP Hayes, and A. Kurta Eds.). Johns Hopkins University Press, Baltimore, Maryland, 352, 17–59.

Barnhart, P. R., & Gillam, E. H. (2014). The impact of sampling method on maximum entropy species distribution modeling for bats. Acta Chiropterologica, 16(1), 241–248.https://doi.org/10.3161/150811014X683435

Barnosky, A. D., Matzke, N., Tomiya, S., Wogan, G. O. U., Swartz, B., Quental, T. B., Marshall, C., McGuire, J. L., Lindsey, E. L., Maguire, K. C., Mersey, B., & Ferrer, E. A. (2011). Has the Earth's sixth mass extinction already arrived? Nature, 471(7336), 51–57. https://doi.org/10.1038/nature09678

Barros, J. D. S., Bernard, E., & Ferreira, R. L. (2020). Ecological preferences of neotropical cave bats in roost site selection and their implications for conservation. Basic and Applied Ecology, 45, 31–41. https://doi.org/10.1016/j.baae.2020.03.007

Bates, P. J. J., Cameron, K., Pearch, M. J., & Hayes, B. (2013). A review of the bats (Chiroptera) of the republic of Congo, including eight species new to the country. Acta Chiropterologica, 15(2), 313–340. https://doi.org/10.3161/150811013X678955

Bernard, R. & Meester, J. A. J. (1982). Female reproduction and the female reproductive cycle of Hipposideros caffer caffer (Sundevall, 1846) in Natal, South Africa. Annals of the Transvaal Museum, 33(8), 131–144.

Bion, R. (2023). ggradar: Create radar charts using ggplot2. R Package Version 0.2.

Brooks, M. E., Kristensen, K., Van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Machler, M., & Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. The R Journal, 9(2), 378–400. https://doi.org/10.32614/RJ-2017-066

Brosset, A. (1968). La Permutation du Cycle sexuel saisonnier chez le chiroptère Hipposideros caffer, au voisinage de l'Équateur.

Browning, E., Gibb, R., Glover-Kapfer, P., & Jones, K. E. (2017). Passive acoustic monitoring in ecology and conservation. http://dx.doi.org/10.25607/OBP-876

Bullock, D., Combes, B., Eales, L., & Pritchard, J. (1987). Analysis of the timing and pattern of emergence of the pipistrelle bat (Pipistrellus pipistrettus). Journal of Zoology, 211(2), 267–274. https://doi.org/10.1111/j.1469-7998.1987.tb01533.x

Cabin, R. J., & Mitchell, R. (2000). To Bonferroni or not to Bonferroni: When and how are the questions. ESA Bulletin, 81, 246–248. 10.2307/20168454

Catto, C., Racey, P., & Stephenson, P. (1995). Activity patterns of the serotine bat (Eptesicus serotinus) at a roost in southern England. Journal of Zoology, 235(4), 635–644. https://doi.org/10.1111/j.1469-7998.1995.tb01774.x

Ceballos, G., Ehrlich, P. R., Barnosky, A. D., García, A., Pringle, R. M., & Palmer, T. M. (2015). Accelerated modern human–induced species losses: Entering the sixth mass extinction. Science Advances, 1(5), e1400253. https://doi.org/10.1126/sciadv.1400253

Cumming, G., & Bernard, R. (1997). Rainfall, food abundance and timing of parturition in African bats. Oecologia, 111, 309–317. https://doi.org/10.1007/s004420050240

Deleva, S., Toshkova, N., Kolev, M., & Tanalgo, K. C. (2023). Important underground roosts for bats in Bulgaria: current state and priorities for conservation. Biodiversity Data Journal, 11, e98734.

Dietz, M., & Kalko, E. K. (2006). Seasonal changes in daily torpor patterns of free-ranging female and male Daubenton's bats (Myotis daubentonii). Journal of Comparative Physiology B, 176, 223–231. https://doi.org/10.1007/s00360-005-0043-x

Erkert, H. G. (1982). Ecological Aspects of Bat Activity Rhythms. In T. H. Kunz (Ed.), Ecology of Bats (pp. 201–242). Springer US. https://doi.org/10.1007/978-1-4613-3421-7_5

Flaquer, C., Torre, I., & Arrizabalaga, A. (2007). Comparison of sampling methods for inventory
of bat communities. Journal of Mammalogy, 88(2), 526–533. https://doi.org/10.1644/06-
MAMM-A-135R1.1
Forsyth, D. M., Scroggie, M. P., & McDonald-Madden, E. (2006). Accuracy and precision of
grey-headed flying-fox (Pteropus poliocephalus) flyout counts. Wildlife Research, 33(1), 57–
65. 10.1071/WR05029
Furey, N. M., & Racey, P. A. (2016). Conservation ecology of cave bats. In Bats in the
Anthropocene: Conservation of Bats in a Changing World (pp. 463–500). Springer, Cham.
https://doi.org/10.1007/978-3-319-25220-9_15
Gibb, R., Browning, E., Glover□Kapfer, P., & Jones, K. E. (2019). Emerging opportunities and
challenges for passive acoustics in ecological assessment and monitoring. Methods in Ecology
and Evolution, 10(2), 169–185. https://doi.org/10.1111/2041-210X.13101
Greville, L. J., Bueno, L. M., Pollock, T., & Faure, P. A. (2022). Quantification of Urinary
Sex Steroids in the Big Brown Bat (Eptesicus fuscus). *Physiological and Biochemical*
Zoology, *95*(1), 22-34.
Günther, L., Lopez, M. D., Knörnschild, M., Reid, K., Nagy, M., & Mayer, F. (2016). From
resource to female defence: The impact of roosting ecology on a bat's mating strategy. Royal
Society Open Science, 3(11), 160503. https://doi.org/10.1098/rsos.160503
Happold, D.C.D. and Happold, M. (1990), Reproductive strategies of bats in Africa. Journal of
Zoology, 222: 557-583. https://doi.org/10.1111/j.1469-7998.1990.tb06014.x
Hartig, F., & Hartig, M. (2022). Package 'DHARMa'. R Package. Available Online:
Https://CRAN. R-Project. Org/Package= DHARMa (Accessed on 5 September 2022).
Henry, M., Thomas, D. W., Vaudry, R., & Carrier, M. (2002). Foraging Distances and Home
Range of Pregnant and Lactating Little Brown Bats (Myotis Lucifugus). Journal of
Mammalogy,   83(3),   767–774.   https://doi.org/10.1644/1545-
1542(2002)083<0767:FDAHRO>2.0.CO;2
Hill, A. P., Prince, P., Snaddon, J. L., Doncaster, C. P., & Rogers, A. (2019). AudioMoth: A low-
cost acoustic device for monitoring biodiversity and the environment. HardwareX, 6, e00073.
https://doi.org/10.1016/j.ohx.2019.e00073
Hughes, A. C. (2017). Understanding the drivers of Southeast Asian biodiversity loss. Ecosphere,
8(1), e01624. https://doi.org/10.1002/ecs2.1624
IPBES (2019): Global assessment report on biodiversity and ecosystem services of the
Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. E. S.
Brondizio, J. Settele, S. Díaz, and H. T. Ngo (editors). IPBES secretariat, Bonn, Germany. 1148
pages. https://doi.org/10.5281/zenodo.3831673
IUCN, N. and I. (International U. for C. of N. (2023). The IUCN Red List of Threatened Species.
Version 2022-2. IUCN Red List of Threatened Species. https://www.iucnredlist.org/en
Jenber, A. J., & Wili, G. G. (2021). Insect diversity in wet and dry seasons at hursa forest, central
Ethiopia. International Journal of Entomology Research, 6(4), 59–63.
Jones, G., & Rydell, J. (1994). Foraging strategy and predation risk as factors influencing
emergence time in echolocating bats. Philosophical Transactions of the Royal Society of
London.   Series   B: Biological   Sciences,   346(1318),   445–455.
https://doi.org/10.1098/rstb.1994.0161
Keeley, B. W., & Tuttle, M. D. (1999). Bats in American bridges. Bat Conservation International
Incorporated.
Koger, B., Hurme, E., Costelloe, B. R., O'Mara, M. T., Wikelski, M., Kays, R., & Dechmann, D.
K. N. (2023). An automated approach for counting groups of flying animals applied to one of
the world's largest bat colonies. Ecosphere, 14(6), e4590. https://doi.org/10.1002/ecs2.4590

Kumar, V., & Umapathy, G. (2019). Non-invasive monitoring of steroid hormones in wildlife for conservation and management of endangered species-A review. *Indian J Exp Biol*, 57, 307-314.

Kunz, T. H. (1982). Roosting ecology of bats. In Ecology of bats (pp. 1–55). Springer.

Kunz, T. H., Lumsden, L. F., & Fenton, M. (2003). Ecology of cavity and foliage roosting bats. Bat Ecology, 1, 3–89. http://doi.org/10.5281

Kurta, A., Bell, G. P., Nagy, K. A., & Kunz, T. H. (1989). Energetics of pregnancy and lactation in freeranging little brown bats (Myotis lucifugus). Physiological Zoology, 62(3), 804–818. https://doi.org/10.1111/j.1469-7998.1992.tb04433.x

Labadie, M., Morand, S., Bourgarel, M., Niama, F. R., Nguilili, F., N'Kaya, T., Caron, A., De Nys, H., (2025a). PeerJ. Habitat sharing and interspecies interactions in caves used by bats in the Republic of Congo 13:e18145 https://doi.org/10.7717/peerj.18145

Labadie, M., Morand, S., Caron, A., De Nys, H. M., Niama, F. R., Nguilili, F., … Roemer, C. (2025b). Passive acoustic monitoring of cave-dwelling bats with a sonotype classifier. Bioacoustics, 1–29. https://doi.org/10.1080/09524622.2024.2438600

Labadie, M. (2025c). Complete results of acoustic data collected in the Republic of the Congo (Tadarida classifier). Zenodo. https://doi.org/10.5281/zenodo.15636144

Lewis, S. E. (1995). Roost Fidelity of Bats: A Review. Journal of Mammalogy, 76(2), 481–496. https://doi.org/10.2307/1382357

Maier, C. (1992). Activity patterns of pipistrelle bats (Pipistrellus pipistrellus) in Oxfordshire. Journal of Zoology, 228(1), 69–80.

Mariton, L., Le Viol, I., Bas, Y., & Kerbiriou, C. (2023). Characterising diel activity patterns to design conservation measures: Case study of European bat species. Biological Conservation, 277, 109852. https://doi.org/10.1016/j.biocon.2022.109852

Mcwilliam, A.N. (1988), The reproductive cycle of male long-fingered bats, Miniopterus minor (Chiroptera: Vespertilionidae), in a seasonal environment of the African Tropics. Journal of Zoology, 216: 119-129. https://doi.org/10.1111/j.1469-7998.1988.tb02419.x

Medellin, R. A., Wiederholt, R., & Lopez-Hoffman, L. (2017). Conservation relevance of bat caves for biodiversity and ecosystem services. Biological Conservation, 211, 45–50.

Meierhofer, M. B., Johnson, J. S., Perez-Jimenez, J., Ito, F., Webela, P. W., Wiantoro, S., Bernard, E., Tanalgo, K. C., Hughes, A., Cardoso, P., Lilley, T., & Mammola, S. (2024). Effective conservation of subterranean-roosting bats. Conservation Biology, 38(1), e14157. https://doi.org/10.1111/cobi.14157

Mering, E. D., & Chambers, C. L. (2014). Thinking outside the box: A review of artificial roosts for bats. Wildlife Society Bulletin, 38(4), 741–751. 10.1002/wsb.461

Mickleburgh, S. P., Hutson, A. M., & Racey, P. A. (2002). A review of the global conservation status of bats. Oryx, 36(1), 18–34. doi:10.1017/S0030605302000054

Millon, L., Julien, J.-F., Julliard, R., & Kerbiriou, C. (2015). Bat activity in intensively farmed landscapes with wind turbines and offset measures. Ecological Engineering, 75, 250–257. https://doi.org/10.1016/j.ecoleng.2014.11.050

Monadjem, A., Taylor, P. J., & Schoeman, M. C. (2020). Bats of southern and central Africa: A biogeographic and taxonomic synthesis. Wits University Press.

Moutaouakil, S., Souza-Silva, M., Oliveira, L. F., Ghamizi, M., & Ferreira, R. L. (2024). A cave with remarkably high subterranean diversity in Africa and its significance for biodiversity conservation. *Subterranean Biology*, *50*, 1-28.

Nkoana, T. T. (2020). Temporal changes in food resource availability between two karst bat assemblages (Doctoral dissertation, University of Pretoria).

Neuhuber, S., Plan, L., Gier, S., Hintersberger, E., Lachner, J., Scholz, D., Lüthgens, C., Braumann, S., Bodenlenz, F., & Voit, K. (2020). Numerical age dating of cave sediments to

quantify vertical movement at the Alpine-Carpathian transition in the Plio-and Pleistocene.
Geologica Carpathica, 71(6), 539–557. https://doi.org/10.31577/GeolCarp.71.6.5

Okonkwo, E. E., Afoma, E., & Martha, I. W. (2017). Cave Tourism and its Implications to
Tourism Development in Nigeria: A Case Study of Agu-Owuru Cave in Ezeagu. International
Journal of Research, 3, 16–24. doi:dx.doi.org/10.20431/2455-0043.0303003

Ormsbee, P. C., Kiser, J. D., & Perlmeter, S. I. (2007). Importance of night roosts to the ecology
of bats. Bats in Forests: Conservation and Management. Johns Hopkins University Press,
Baltimore, MD, 129–151.

Patriquin, K. J., & Ratcliffe, J. M. (2016). Should I stay or should I go? Fission–fusion dynamics
in bats. Sociality in Bats, 65–103. https://doi.org/10.1007/978-3-319-38953-0_4

Patterson, B. D., Dick, C. W., & Dittmar, K. (2007). Roosting habits of bats affect their
parasitism by bat flies (Diptera: Streblidae). Journal of Tropical Ecology, 23(2), 177–189.
doi:10.1017/S0266467406003816

Pretorius, M., Markotter, W. & Keith, M. Assessing the extent of land-use change around
important bat-inhabited caves. BMC Zool 6, 31 (2021). https://doi.org/10.1186/s40850-
021-00095-5

Postawa, T., & Gas, A. (2009). Do the thermal conditions in maternity colony roost determine the
size of young bats? Comparison of attic and cave colonies of Myotis myotis in Southern
Poland. Folia Zoologica, 58(4), 396.

R Core Team. (2023). R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing. https://www.R-project.org/

Racey, P. A., & Entwistle, A. C. (2000). Life-history and Reproductive Strategies of Bats. In
Reproductive Biology of Bats (pp. 363–414). Elsevier. https://doi.org/10.1016/B978-
012195670-7/50010-2

Revilla-Martín, N., Budinski, I., Puig-Montserrat, X., Flaquer, C., & López-Baucells, A. (2020).
Monitoring cave-dwelling bats using remote passive acoustic detectors: A new approach for
cave monitoring. Bioacoustics, 00(00), 1–16. https://doi.org/10.1080/09524622.2020.1816492

Roemer, C., Julien, J. F., & Bas, Y. (2021). An automatic classifier of bat sonotypes around the
world. Methods in Ecology and Evolution, 12(12), 2432–2444. https://doi.org/10.1111/2041-
210X.13721

Russo, D., & Ancillotto, L. (2015). Sensitivity of bats to urbanization: A review. Mammalian
Biology, 80(3), 205–212. https://doi.org/10.1016/j.mambio.2014.10.003

Racey, P.A. & Entwistle, A.C. (2000). Conservation Ecology of Cave Bats. In: Conservation of
Tropical Biodiversity. Springer.

Rydell, J. (1993). Variation in Foraging Activity of an Aerial Insectivorous Bat during
Reproduction. Journal of Mammalogy, 74(2), 503–509. https://doi.org/10.2307/1382411

Sabol, B. M., & Hudson, M. K. (1995). Technique using thermal infrared-imaging for estimating
populations of gray bats. Journal of Mammalogy, 76(4), 1242–1248.
https://doi.org/10.2307/1382618

Simmons, N. B., & Cirranello, A. L. (2025). Bat Species of the World: A taxonomic and
geographic database. https://batnames.org/query.html

Simons, J. W. (1998). Guano mining in Kenyan lava tunnel caves. International Journal of
Speleology, 27, 4. https://api.semanticscholar.org/CorpusID:56146083

Struebig, M. J., Kingston, T., Zubaid, A., Le Comber, S. C., Mohd-Adnan, A., Turner, A., Kelly,
J., Bożek, M., & Rossiter, S. J. (2009). Conservation importance of limestone karst outcrops
for Palaeotropical bats in a fragmented landscape. Biological Conservation, 142(10), 2089–
2096. https://doi.org/10.1016/j.biocon.2009.04.005

Sugai, L. S. M., Desjonquères, C., Silva, T. S. F., & Llusia, D. (2020). A roadmap for survey designs in terrestrial acoustic monitoring. Remote Sensing in Ecology and Conservation, 6(3), 220–235. https://doi.org/10.1002/rse2.131

Swift, S. M. (1980). Activity patterns of pipistrelle bats (Pipistrellus pipistrellus) in north east Scotland. Journal of Zoology, 190(3), 285–295. https://doi.org/10.1111/j.1469-7998.1980.tb01428.x

Swift, S. M. (1997). Roosting and foraging behaviour of Natterer's bats (Myotis nattereri) close to the northern border of their distribution. Journal of Zoology, 242(2), 375–384. https://doi.org/10.1111/j.1469-7998.1997.tb05809.x

Tanalgo, K. C., Oliveira, H. F., & Hughes, A. C. (2022). Mapping global conservation priorities and habitat vulnerabilities for cave-dwelling bats in a changing world. Science of The Total Environment, 843, 156909.

Tanalgo, K. C., & Hughes, A. C. (2024). Bat Cave Vulnerability Index 3.0 (BCVI-S): An integrative and scalable tool to prioritise bat caves for conservation. Global Ecology and Conservation, 57, e03396. https://doi.org/10.1016/j.gecco.2024.e03396

The Protection of Bat Roost Guidelines Subcommittee, Sheffield, S. R., Shaw, J. H., Heidt, G. A., & McClenaghan, L. R. (1992). Guidelines for the protection of bat roosts. Journal of Mammalogy, 707–710. https://doi.org/10.2307/1382051

Thieurmel, B., Elmarhraoui, A., & Thieurmel, M. B. (2022). Package 'suncalc'. R Package Version 0.5. https://cran.r-project.org/package=suncalc

Thomas, D. W., & West, S. D. (1989). Sampling methods for bats. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. https://doi.org/10.2737/pnw-gtr-243

Voigt, C. C., & Schwarzenberger, F. (2008). Reproductive endocrinology of a small tropical bat (female Saccopteryx bilineata; Emballonuridae) monitored by fecal hormone metabolites. *Journal of Mammalogy*, *89*(1), 50-57.

Westcott, D. A., & McKeown, A. (2004). Observer error in exit counts of flying-foxes (Pteropus spp.). Wildlife Research, 31(5), 551–558. 10.1071/WR03091

Whiting, J. C., Doering, B., Aho, K., & Rich, J. (2021). Long-term patterns of cave-exiting activity of hibernating bats in western North America. Scientific Reports, 11(1), 8175.

Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York. https://ggplot2.tidyverse.org

Wilkinson, G. S., Adams, D. M., & Rayner, J. G. (2024). Sex, season, age and status influence urinary steroid hormone profiles in an extremely polygynous neotropical bat. *Hormones and Behavior*, *164*, 105606.

Wolda, H. (1988). Insect seasonality: Why? Annual Review of Ecology and Systematics, 19(1), 1–18. https://doi.org/10.1146/annurev.es.19.110188.000245

Wolkovich, E. M., Cook, B. I., McLauchlan, K. K., & Davies, T. J. (2014). Temporal ecology in the Anthropocene. Ecology Letters, 17(11), 1365–1379. https://doi.org/10.1111/ele.12353

Wright, G. S. (2009). Hipposideros caffer (Chiroptera: Hipposideridae). Mammalian Species, 845, 1–9. https://doi.org/10.1644/845.1.Key