Convergence in amino acid outsourcing between animals and predatory bacteria

Niko Kasalo[1], Mirjana Domazet-Lošo[2] and Tomislav Domazet-Lošo[1,3,*]

[1]Laboratory of Evolutionary Genetics, Division of Molecular Biology, Ruđer Bošković

Institute, Bijenička cesta 54, HR-10000 Zagreb, Croatia

[2]Department of Applied Computing, Faculty of Electrical Engineering and Computing,

University of Zagreb, Unska 3, HR-10000 Zagreb, Croatia

[3]School of Medicine, Catholic University of Croatia

*Correspondence: T.D.-L. tdomazet@irb.hr

Abstract

All animals have outsourced about half of the 20 proteinogenic amino acids (AAs). We recently

demonstrated that the loss of biosynthetic pathways for these outsourced AAs is driven by

energy-saving selection. Paradoxically, these metabolic simplifications enabled animals to use

costly AAs more frequently in their proteomes, allowing them to explore sequence space more

freely. Based on these findings, we proposed that environmental AA availability and cellular

respiration mode are the two primary factors determining the evolution of AA auxotrophies in

animals. Remarkably, our recent analysis showed that bacterial AA auxotrophies are also

governed by energy-related selection, thereby roughly converging with animals. However,

bacterial AA auxotrophies are highly heterogeneous and scattered across the bacterial

phylogeny, making direct ecological and physiological comparisons with the animal AA

outsourcing model challenging. To better test the universality of our model, we focused on

Bdellovibrionota and Myxococcota—two closely related bacterial phyla that, through aerobic

respiration and a predatory lifestyle, best parallel animals. Here, we show that Bdellovibrionota,

driven by energy-related selection, outsourced a highly similar set of AAs to those in animals.

This sharply contrasts with Myxococcota, which exhibit far fewer AA auxotrophies and rarely

show signatures of energy-driven selection. These differences are also reflected in

Bdellovibrionota proteomes, which are substantially more expensive than those of

Myxococcota. Finally, we found evidence that the expression of costly proteins plays a crucial

role in the predatory phase of the Bdellovibrio life cycle. Together, our findings suggest that

Bdellovibrionota, through their obligate predatory lifestyle, exhibit the closest analogy to the
AA auxotrophy phenotype observed in animals. In contrast, facultative predation, as seen in
Myxococcota, appears to substantially limit the evolution of AA auxotrophies. These cross-
domain convergences strongly support the general validity of our AA outsourcing model.

Keywords: bacteria; predation; amino acids; animals; auxotrophy; energetics; selection;
evolution

Introduction

The biosynthesis of amino acids (AAs) is a fundamental biochemical process essential for
sustaining all life on Earth. However, despite the universal metabolic demand for all 20
proteinogenic AAs, some organisms have lost the ability to synthesize certain AAs
endogenously. The most well-known example is animals [1–3], which have lost the capacity to
produce approximately half of the complete AA set. A similar pattern of auxotrophies has
independently emerged in other eukaryotic groups, such as certain amoebae and euglenozoans
[1,2]. While most bacteria remain fully prototrophic [4], many bacterial lineages display
varying degrees of AA auxotrophy [4–6].

It has long been speculated that the ability to acquire AAs from the environment influences the
evolution of AA auxotrophies [2]. For instance, experiments have shown that bacteria
auxotrophic for an externally supplemented AA can gain a selective advantage over fully
prototrophic counterparts [7]. However, a robust theoretical framework for this phenomenon
remained elusive. In our previous work, we addressed this gap by proposing a model that
explains the evolution of AA outsourcing through several key factors: an AA is more likely to
be lost if (i) its biosynthesis is highly energy-demanding, (ii) it has a low pleiotropic effect, (iii)
it is abundantly available in the environment, and (iv) the organism relies on efficient aerobic
respiration for energy production [1].

Most importantly, we demonstrated that there is a constant selective pressure to outsource the
synthesis of energetically costly AAs to the environment. Surprisingly, this leads to an
increased usage of these expensive AAs in proteomes, allowing animal proteins to explore
sequence space more freely [1,8]. To determine whether these global patterns are also valid in
bacteria, we investigated AA auxotrophies across bacterial phylogeny and found that energy-
related selection also plays an important role in shaping AA outsourcing in bacteria [5].

However, bacteria exhibit far greater metabolic and ecological diversity than animals [4,9,10],

which necessitates testing our AA outsourcing model in specific bacterial groups whose

lifestyles more closely parallel those of animals.

One such group is the phylum Bdellovibrionota, which includes several aerobic or

microaerophilic species with an obligate predatory lifestyle [11–13]. Within this phylum,

predation occurs via two distinct strategies: epibiotic predation, where the bacterium attaches

to the prey cell and leeches nutrients from its surface, and endobiotic predation, where the

bacterium enters the prey cell and forms a bdelloplast, within which it feeds, grows, and divides

[11–13]. The life cycle of Bdellovibrionota consists of two transcriptionally distinct phases: the

attack phase, during which the bacterium seeks and attaches to prey, and the feeding phase,

characterized by growth and replication [12,14].

Another bacterial group exhibiting animal-like behaviors is the phylum Myxococcota, a close

relative of Bdellovibrionota [15,16]. Myxococcota are aerobes known for their complex

multicellular behaviors, including social movement in coordinated "wolf packs," fruiting body

formation, and predation [17]. However, unlike Bdellovibrionota, Myxococcota are not

obligate predators—they can scavenge nutrients from dead organic matter and survive periods

of starvation through sporulation [18–20].

The predatory lifestyle evolved independently in Bdellovibrionota and Myxococcota [16],

leading to vastly different phenotypic outcomes [11–13,18–20]. Thus, these two bacterial

groups provide an ideal system to test the predictive power of our AA outsourcing model, as

they allow us to directly link the evolution of AA auxotrophies in bacteria to distinct ecological

and physiological traits [1,5,8].

Here, we demonstrate that Bdellovibrionota evolved a remarkably animal-like set of AA

auxotrophies, accompanied by an increase in relative proteome costs. Our findings indicate that

energy-related selection played a key role in shaping these auxotrophies and that the attack and

feeding phases of their life cycle exhibit distinct energy dynamics at the transcriptomic level.

Surprisingly, this pattern does not hold for Myxococcota, which show fewer auxotrophies and

lower proteome costs, suggesting a fundamental difference in how different reliance on a

predatory lifestyle shapes metabolic evolution across these two groups.

Results

We suspected that the animal-like ecophysiology of Bdellovibrionota and Myxococcota imposes energy-related selection on their AA metabolism, leading to reductions in AA biosynthetic pathways similar to those observed in animals. To test this hypothesis, we first estimated the completeness of AA biosynthesis pathways in 89 Bdellovibrionota and 203 Myxococcota high-quality proteomes, which we retrieved from NCBI (Table S1). To assess AA biosynthesis pathway completeness, which represents the likelihood of a given pathway being present, we used the MMseqs2 clustering approach [5,21]. Heatmap representations clearly revealed significantly lower AA completeness scores in Bdellovibrionota compared to Myxococcota (Figure 1, File S1).

Most Bdellovibrionota showed reductions in pathway completeness scores for nine amino acids (AAs) that fall on the expensive end of the biosynthesis cost distribution (Figure 1, File S1). This set of AAs with low completeness scores largely overlaps with those that are auxotrophic in animals [1], with one notable exception—lysine, which appears to be prototrophic in Bdellovibrionota (Figure 1, File S1). In contrast, the pattern of AA biosynthesis pathway reduction in Myxococcota is markedly different. Most Myxococcota remain prototrophic for the majority of AAs, with only valine, leucine, and isoleucine biosynthesis pathways consistently exhibiting reductions (Figure 1, File S1). An exception to this trend is observed in two Myxococcota species—*Vulgatibacter incomptus* and *Pajaroellobacter abortibovis*—which display a substantial number of AA auxotrophies (Figure S1). Taken together, these findings suggest that while some reductions of expensive AA occur in Myxococcota, some ecological factors likely prevent them from outsourcing most of their costly AA biosynthesis pathways.

Figure 1. Completeness score of AA biosynthesis pathways in Bdellovibrionota and Myxococcota. We created a database of 89 Bdellovibrionota and 203 Myxococcota proteomes to get a comprehensive overview of AA dispensability in these groups. Full figure is shown in File S1. We retrieved all enzymes involved in AA biosynthesis from the KEGG and MetaCyc databases (reference collection) and searched for their homologs within our

Bdellovibrionota/Myxococcota database using MMseqs2 (see Methods). For each AA, we

showed a completeness score, which represents the percentage of enzymes within a pathway

that returned significant sequence similarity matches to our reference collection of AA

biosynthesis enzymes. In the case of AAs with multiple alternative pathways, we showed the

results only for the most complete one.

To globally test whether energy-related selection influences the observed reductions in AA

biosynthesis pathways of Bdellovibrionota and Myxococcota, we correlated the average AA

auxotrophy index with opportunity costs calculated under high respiration mode [5], a metric

that estimates the impact of AA biosynthesis on the cell's energy budget. To obtain the average

AA auxotrophy index for a given AA, we first subtracted the completeness score from 1

(Material and Methods, Equation 1) and then averaged the resulting AI values across all

considered proteomes. We detected a significant correlation between higher biosynthesis costs

and the loss of AA biosynthetic ability in Bdellovibrionota (Figure 2A). This suggests that

selection driven by energy management shaped the global pattern of auxotrophies in

Bdellovibrionota. As might be expected considering heatmap pattern (Figure 1), this broad-

scale analysis did not detect a positive correlation in Myxococcota, which are prototrophic for

most AAs (Figure 2B).

Figure 2. Correlation between AA biosynthesis cost and the average AA auxotrophy index. We estimated the AA auxotrophy index (AI), a measure which equals one minus completeness score (see Materials and Methods, Equation 1), for 89 Bdellovibrionota (a) and 203 Myxococcota (b) proteomes and calculated the average value for every AA. We correlated this value with the opportunity cost of each AA, calculated for high respiration mode (see Materials and Methods). Pearson correlation coefficient and p-value are shown on the graph.

To further evaluate whether the observed reductions in AA biosynthesis pathways in Bdellovibrionota and Myxococcota result from energy-optimizing selection, we analyzed AA auxotrophy patterns at the species level. To achieve this, we first explicitly determined the AA auxotrophy status of each species by transforming the completeness score to binary values (auxotrophic/prototrophic) (see Materials and Methods, Table S2). Although this procedure

reduces the information contained in the completeness scores, it allowed us to test the impact of energy-related selection more directly by explicitly defining auxotrophic and prototrophic AAs. In principle, the loss of even a single enzyme within a biochemical pathway could render that pathway nonfunctional. Thus, we assigned auxotrophy status to any AA whose biosynthesis pathway was incomplete. Using this binary-transformed dataset, we statistically compared opportunity costs between auxotrophic and prototrophic AAs within each species [1]. In addition, we applied a permutation-based selection test to assess the probability that the observed constellation of auxotrophic AAs evolved under energy-related selection [1].

On average, Bdellovibrionota species exhibited 7.89 auxotrophic AAs per species (Table S2). The comparison between auxotrophic and prototrophic AA sets using the Mann-Whitney non-parametric test shows that auxotrophic AAs have significantly higher opportunity costs in 92% of the 89 tested Bdellovibrionota species (Table S2; File S2). For illustration, we singled out the results of this comparison for the type strains of *Bdellovibrio bacteriovorus* and *Pseudobdellovibrio exovorus*, representing endobiotic and epibiotic lifestyles, respectively (Figure 3). It is evident that, regardless of feeding ecology, auxotrophic AAs are significantly more expensive than prototrophic ones (Figure 3A,B). The permutation-based selection test revealed similar global trends, detecting that energy-related selection impacted the observed distribution of auxotrophic AAs in 61% of Bdellovibrionota species (Table S2; File S2). For instance, the average opportunity cost of the auxotrophic AA sets observed in *Bdellovibrio bacteriovorus* and *Pseudobdellovibrio exovorus* falls at the right end of the distribution of all possible permutations (Figure 3C,D). This indicates, just like in animals, that energy-optimizing selection governed the outsourcing of auxotrophic AAs in these bacteria.

The trends are quite different in Myxococcota, where an average of 2.79 AAs are auxotrophic (Table S2). The Mann-Whitney non-parametric test shows that auxotrophic AAs have significantly higher opportunity costs in only 9% of the 203 tested species (Table S2; File S2). In comparison, the permutation-based selection test revealed similar results, indicating that energy-related selection impacted AA auxotrophies in only 7% of Myxococcota species (Table S2). A prominent representative of Myxococcota that shows the impact of energy-related selection on AA auxotrophies is *Pajaroellobacter abortibovis*, a species whose pathogenic ecology differs from the facultative predatory lifestyle of most other Myxococcota (File S1, Table S2, File S2). Taken together, these findings suggest that energy-optimizing selection related to AA auxotrophies is rather rare among Myxococcota compared to Bdellovibrionota.

Figure 3. Comparisons of auxotrophic and prototrophic AA sets in *Bdellovibrio bacteriovorus* (A, C) and *Pseudobdellovibrio exovorus* (B, D). (A, B) The comparison of opportunity costs in high respiration mode between auxotrophic and prototrophic AA groups was tested by the Mann-Whitney U test with continuity correction. The corresponding W-value, p-value, and effect size (r) are depicted in each panel. The X symbol represents the mean. Individual AAs are shown by one-letter symbols. (C, D) Permutation analyses of opportunity costs (selection tests) were performed by calculating the average opportunity cost of every possible permutation for the number of auxotrophic AAs identified in a given species (n = 8 *B. bacteriovorus*, n = 9 *P. exovorus*). The proportions of these averages are shown in histograms. The obtained distribution represents the empirical probability mass function (PMF). The value in red denotes the average opportunity cost value of the EAA sets observed in nature. The p-value was calculated by summing the proportions that correspond to average opportunity cost values equal

to, or more extreme, than the observed value. Low p-values indicate a high probability that energy-related selection drove the loss of auxotrophic AA biosynthesis capability.

The fact that Bdellovibrionota converge to an animal-like set of amino acid (AA) auxotrophies, while closely related Myxococcota remain mainly prototrophic, allows us to further test the predictions of our model [1]. In our previous work, we showed that animals code for significantly more expensive proteomes, compared to Choanoflagellates, their sister group which is mainly prototrophic [1]. We proposed that animals have more expensive proteomes than Choanoflagellates, likely because they consume less energy on AA biosynthesis and are thus able to maintain a larger number of expensive auxotrophic AAs in their proteomes [1]. Using a non-parametric test to compare the opportunity costs (high respiration mode) of an average AA in bacterial proteomes (Material and Methods, Equation 2), we recovered an analogous result: the proteomes of the more auxotrophic Bdellovibrionota are significantly more expensive than those of the more prototrophic Myxococcota (Figure 4), underscoring the universality of energy-related selection on proteome composition.

Figure 4. The comparison of the opportunity cost (OC) of an average AA in proteomes of Bdellovibrionota and Myxococcota. The opportunity cost under high respiration mode (OC) of an average AA in a proteome represents a weighted mean of AA biosynthesis energy costs where the frequencies of twenty AAs in the proteome act as weights. The differences in energy costs between the two groups were shown by boxplots and the significance of these differences

was tested by the Mann-Whitney U test with continuity correction. We depicted the corresponding W-value, p-value, and effect size (r) in each panel. The X symbol represents the mean. The list of Bdellovibrionota (89) and Myxococcota (203) whose proteomes are included in calculations is available in Supplementary Data X.

Finally, we used the published transcriptome data to examine the energetics of the two distinct phases in the life cycle of *Bdellovibrio bacteriovorus*: the attack phase and the growth phase [14]. For each transcript, we calculated its frequency in the transcriptome at a given life cycle phase [22]. We then multiplied this transcript frequency by the opportunity cost of an average amino acid (AA) encoded by that transcript ($OC_{protein}$) (Equations 3 and 4). This measure, which we named the transcript energy score (TES), couples transcript levels in the cell with the encoded protein costs (see Material and Methods). Higher TES values reflect greater impact, while lower TES values indicate a smaller impact of a given transcript on the total energy budget of the cell. We performed the TES-based analysis in two ways: by including all genes expressed per phase and by considering only those which were exclusively expressed in one phase (Figure 5).

The attack phase is generally characterized by a much smaller number of expressed genes, which are more evenly distributed across the narrower range of TES values (Figure 5). In contrast, the TES values of the growth phase are predominantly grouped at the lower end of the TES range. This suggests that the attack phase of predatory *B. bacteriovorus* is underpinned by high transcription of a relatively small number of genes many of which encode expensive proteins. In contrast, the growth phase is characterized by relatively low transcription of many genes that encode cheaper proteins. These differences are even more apparent in the analysis of phase-specific genes (Figure 5B). Together, this suggests that the active predation in *B. bacteriovorus* requires proteins which are composed of expensive AAs. As an analogy, we previously speculated that similar phenotypes might exist in the context of animal predation [1].

Figure 5. The comparison of transcriptome energy scores (TES) between two phases in the life cycles of *B. bacteriovorus*. Calculations were performed (a) for all genes that are expressed per phase and (b) only for phase-specific genes. The TES value represents the average opportunity cost of a gene product multiplied by its frequency in the transcriptome at a given life cycle phase. The significance of the difference between the two phases was tested by the Mann-Whitney U test with continuity correction. We depicted the corresponding W-value, p-value, and effect size (r). Outliers were removed from the graph for the clarity of presentation, but were included in the calculation of statistics. The X symbol represents the mean, and the horizontal line within a violin-plot represents the median. Transcription data was retrieved from Karunker et al. 2013 (Karunker et al., 2013).

Discussion

Functional outsourcing presumes that genes that support essential functions can be lost from the genome if their activity can be substituted through environmental interactions [8]. We have successfully applied this concept to the evolution of AA auxotrophies in animals and bacteria [1,5]. We found that all animals and some bacterial groups which are capable of harvesting a sufficient amount of AAs from their ecosystem lose the ability to produce expensive AAs on their own [1,5,8]. This AA outsourcing is at least partially driven by energy-optimizing selection, which not only favors the loss of the ability to synthesize expensive AAs but also allows for more frequent usage of expensive AAs in the proteome [1,5]. We proposed that these phenotypes could have been triggered in animals by predation and aerobic respiration [1]. If

these processes are indeed selection-driven, it would be expected that they occur convergently under similar ecological pressures across the tree of life.

Unlike most other bacteria, Bdellovibrionota and Myxococcota are aerobic predators; they hunt and consume their bacterial prey under aerobic or microaerobic conditions [11–13,23]. As an independent evolutionary event, these behaviors were likely crucial for the outsourcing of AA production in animals [1]. We have shown here that obligate aerobic predation left an astonishingly similar metabolic impact on Bdellovibrionota, resulting in the largely overlapping set of AA auxotrophies compared to animals. In contrast, very few auxotrophies can be observed in Myxococcota, which are facultative predators [18,20].

There are likely multiple factors that influenced the vast differences in AA biosynthesis capabilities between Bdellovibrionota and Myxococcota. However, the most apparent and potentially vital factor is that Myxococcota are only facultatively predatory [18], making them unable to consistently obtain AAs in large quantities. This is supported by the observation that Bdellovibrionota grow and assimilate carbon at higher rates than Myxococcota [18]. Furthermore, it has been observed that Bdellovibrionota are significantly more abundant in aerobic environments than in anaerobic ones, in contrast to Myxococcota, which show no apparent preference [23]. This might indicate that the metabolism of Bdellovibrionota is more dependent on efficient respiration, which could also drive them toward heightened AA auxotrophy levels [1]. Another important factor might be feeding efficiency—Bdellovibrionota are always physically connected to their prey, while Myxococcota secrete hydrolytic enzymes around their prey, which carries the risk of diffusion, leading to a lower return of the energy expended on predation [24].

The reduction in the ability to synthesize expensive AAs is directly correlated with the increased usage of expensive AAs in the proteomes [1,5]. Animals use more frequently expensive AAs in their proteomes than their sister group, choanoflagellates, and here we showed that the same is true for Bdellovibrionota when compared to their related group, Myxococcota [15]. Although Myxococcota have larger proteomes than Bdellovibrionota [11,24], their encoded AAs are on average cheaper by a large margin, which suggests that their complex lifestyle requires a wide range of functions that do not require proteins with very expensive AAs.

On the other hand, the ecology of Bdellovibrionota is relatively simple, consisting of two distinct phases: the attack phase, during which cells actively hunt, and the growth phase, during which cells consume their prey and subsequently divide [11,12,14]. A previous study produced

transcriptome data for these two phases in *B. bacteriovorus* [14], providing a unique
opportunity to examine the effects of energy-related selection on the *B. bacteriovorus* life cycle.
We used a novel measure, the transcript energy score (TES), which combines the energy cost
of a coding gene with its level of transcription, producing higher scores for more expensive and
highly expressed transcripts. Using this metric, we found that the growth phase is underlined
by a much larger set of transcripts with relatively similar expression levels that encode for
cheaper AAs, while the attack phase consists of a relatively small subset of transcripts with a
wide range of expression levels that encode for more expensive AAs. This supports the
possibility that the energy saved through the outsourcing of AAs was invested in the attack
phase, enabling more efficient predatory behavior and thus ensuring a more consistent influx
of AAs from the environment, creating a positive feedback loop.

In conclusion, it is evident that every level of biological organization—from metabolism and
proteome composition to the regulation of gene transcription—is intimately tied to the
organism's energy budget. The repeated convergent evolution of similar sets of AA
auxotrophies across eukaryotes and prokaryotes can be attributed to obligate predation under
aerobic conditions, which itself goes hand in hand with the evolution of novel and expensive
predation-related functions. Deeper understanding of the evolutionary pressures leading to
predatory behavior in bacteria is especially important, as it has implications not only for
understanding the regulation of ecological networks [18] but also for potential medical
applications [25].

Materials and Methods

All proteomes used in this study were retrieved from the NCBI GenBank. We acquired the
highest-quality Bdellovibrionota (89) and Myxococcota (203) proteomes by using the NCBI
filter to exclude atypical genomes, metagenome-assembled genomes, and genomes from large
multi-isolate projects.

We conducted the detection of amino acid (AA) biosynthesis pathway completeness using the
protocol described in our earlier publication [5]. Briefly, we prepared a reference database of
387,892 enzyme sequences across 2,095 bacterial species, with each enzyme annotated with
the biosynthetic pathway it is involved in [5]. We combined the reference database with our

proteomes and clustered the sequences using MMseqs2 [21] with the following parameters: -cluster-mode 0, -cov-mode 0, -c 0.8, and -e 0.001. We then functionally annotated all members of a cluster based on the presence of enzymes from the reference database in that cluster.

For each AA biosynthesis pathway and species in the database, we calculated a pathway completeness score (i.e., prototrophy index) by dividing the number of detected enzymes by the total number of enzymes in that pathway,s resulting in values ranging from 0 to 1. If a species contained alternative biosynthetic pathways for an AA, the pathway with the highest completeness score was selected. We then calculated the AA auxotrophy index of the i-th amino acid (AI_i) by subtracting the completeness score from 1 [5]:

$$AI_i = 1 - completness\ score \tag{1}$$

We used the energy costs of AA biosynthesis as described in our earlier publications [1,5]. The opportunity cost reflects the impact of AA synthesis on the cell's energy budget and is calculated as the sum of the energy lost in the synthesis of AAs (direct cost) and the energy that would have been produced if a cell catabolized precursors instead of making AAs. Using the AA opportunity cost, we also calculated the opportunity cost of an average AA in each proteome ($OC_{proteome}$) using the following equation:

$$OC_{proteome} = \frac{\sum_{i=1}^{n=20} OC_i \times N_i}{\sum_{i=1}^{n=20} N_i} = \sum_{i=1}^{n=20} OC_i \times F_i \tag{2}$$

In this equation, OC_i represents the opportunity cost of the i-th AA, N_i denotes the total number of occurrences of this AA in the entire proteome, and F_i represents the frequency of the AA in the proteome.

The permutation analyses were performed separately for each species by first determining the number of auxotrophic AAs and then generating all possible permutations for that number of auxotrophic AAs. For each permutation, we then calculated the average opportunity cost of the auxotrophic AAs. For example, if a species was found to be auxotrophic for 10 AAs, we found all possible combinations of 10 AAs and calculated the average opportunity cost of each. Since there is a limited number of possible average values, each value was treated as a bin. We calculated the proportion of permutations within a bin by dividing the number of elements in that bin by the total number of permutations. The obtained distribution represents empirical probability mass function (PMF) which was then used to calculate the probability that the observed set of auxotrophies in a given species is a result of a random process. We calculated

p-values by summing the proportions of permutation in the range from the actual value observed in nature to the most extreme value at the closest distribution tail [1].

For transcriptome analysis, we obtained data on differential gene expression in the two distinct life cycle phases of *Bdellovibrio bacteriovorus* from an earlier study [14]. For each gene (i), we calculated its proportion in a given transcriptome phase by dividing its transcription value (t_i) by the sum of all transcription values in that phase. This proportion in essence represents the frequency of each transcript in the transcriptome phase ($f_{transcript}$). We then multiplied this number by the opportunity cost (high respiration mode) of the protein encoded by that gene ($OC_{protein}$) to obtain a transcript energy score (TES). This score is meant to represent the energy impact of each transcript on the total energy budget of a transcriptome phase. TES was calculated using the following equation:

$$TES_i = \frac{t_i}{\sum_i^n t_i} \times OC_{protein} = f_{transcript} \times OC_{protein} \qquad (3)$$

The opportunity cost of proteins was calculated using the following equation:

$$OC_{protein} = \frac{\sum_{i=1}^{n=20} OC_i \times n_i}{\sum_{i=1}^{n=20} n_i} = \sum_{i=1}^{n=20} OC_i \times f_i \qquad (4)$$

In this equation, OC_i represents the opportunity cost of the i-th AA, n_i denotes the total number of occurrences of this AA in the protein, and f_i represents the frequency of the AA in the protein.

We used the Mann-Whitney U test with continuity correction to compare the opportunity costs of an average AA in proteomes ($OC_{proteome}$) of Bdellovibrionota and Myxococcota using the package rcompanion (https://CRAN.R-project.org/package=rcompanion). We used the same test to compare transcript energy scores (TES) of different life cycle phases in *B. bacteriovorus*. To calculate correlations, we used the cor.test() function in the R stats (v. 3.6.2) package. The heatmap was visualized using the ComplexHeatmap package [26].

Supplementary Materials: File S1: Full pathway completeness heatmap; File S2: Per-species statistics on binary encoded auxotrophies; Table S1: Database with pathway completeness detection results.xlsx; Table S2: Summary of statistics based on binary encoding of auxotrophies

Author Contributions: Conceptualization, N.K., M.D.-L. and T.D.-L.; methodology, N.K., M.D.-L. and T.D.-L.; software, N.K. and M.D.-L.; validation, N.K., M.D.-L. and T.D.-L.; formal analysis, N.K., M.D.-L. and T.D.-L.; writing—original draft preparation, N.K., M.D.-L.

and T.D.-L.; writing—review and editing, N.K., M.D.-L. and T.D.-L.; visualization, N.K. and M.D.-L.; supervision, M.D.-L. and T.D.-L. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the European Regional Development Fund KK.01.1.1.01.0009 DATACROSS (M.D.-L., T.D.-L.).

Data Availability Statement: All data are available in the supplementary materials.

Acknowledgments: We thank M. Futo, A. Tušar, S. Koska, D. Franjević and G. Klobučar for discussions. We used the computational resources of the University Computing Center - SRCE (Padobran) and the Institute Ruđer Bošković.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Kasalo, N.; Domazet-Lošo, M.; Domazet-Lošo, T. Massive Outsourcing of Energetically Costly Amino Acids at the Origin of Animals. *bioRxiv* 2024, https://doi.org/10.1101/2024.04.18.590100.

2. Payne, S.H.; Loomis, W.F. Retention and Loss of Amino Acid Biosynthetic Pathways Based on Analysis of Whole-Genome Sequences. *Eukaryot. Cell* 2006, *5*, 272–276, doi:10.1128/ec.5.2.272-276.2006.

3. Trolle, J.; McBee, R.M.; Kaufman, A.; Pinglay, S.; Berger, H.; German, S.; Liu, L.; Shen, M.J.; Guo, X.; Martin, J.A.; et al. Resurrecting Essential Amino Acid Biosynthesis in Mammalian Cells. *eLife* 2022, *11*, e72847, doi:10.7554/eLife.72847.

4. Ramoneda, J.; Jensen, T.B.N.; Price, M.N.; Casamayor, E.O.; Fierer, N. Taxonomic and Environmental Distribution of Bacterial Amino Acid Auxotrophies. *Nat. Commun.* 2023, *14*, 7608, doi:10.1038/s41467-023-43435-4.

5. Kasalo, N.; Domazet-Lošo, T.; Domazet-Lošo, M. Bacterial Amino Acid Auxotrophies Enable Energetically Costlier Proteomes. *bioRxiv* 2025, https://doi.org/10.1101/2025.01.24.634666.

6. Starke, S.; Harris, D.M.M.; Zimmermann, J.; Schuchardt, S.; Oumari, M.; Frank, D.; Bang, C.; Rosenstiel, P.; Schreiber, S.; Frey, N.; et al. Amino Acid Auxotrophies in Human Gut

Bacteria Are Linked to Higher Microbiome Diversity and Long-Term Stability. *ISME J.* 2023, *17*, 2370–2380, doi:10.1038/s41396-023-01537-3.

7. D'Souza, G.; Waschina, S.; Pande, S.; Bohl, K.; Kaleta, C.; Kost, C. Less Is More: Selective Advantages CAN Explain the Prevalent Loss of Biosynthetic Genes in Bacteria. *Evolution* 2014, *68*, 2559–2570, doi:10.1111/evo.12468.

8. Domazet-Lošo, M.; Široki, T.; Šimičević, K.; Domazet-Lošo, T. Macroevolutionary Dynamics of Gene Family Gain and Loss along Multicellular Eukaryotic Lineages. *Nat. Commun.* 2024, *15*, 2663, doi:10.1038/s41467-024-47017-w.

9. Lozupone, C.A.; Knight, R. Global Patterns in Bacterial Diversity. *Proc. Natl. Acad. Sci.* 2007, *104*, 11436–11440, doi:10.1073/pnas.0611525104.

10. Price, M.N.; Zane, G.M.; Kuehl, J.V.; Melnyk, R.A.; Wall, J.D.; Deutschbauer, A.M.; Arkin, A.P. Filling Gaps in Bacterial Amino Acid Biosynthesis Pathways with High-Throughput Genetics. *PLOS Genet.* 2018, *14*, e1007147, doi:10.1371/journal.pgen.1007147.

11. Sockett, R.E. Predatory Lifestyle of *Bdellovibrio Bacteriovorus*. *Annu. Rev. Microbiol.* 2009, *63*, 523–539, doi:10.1146/annurev.micro.091208.073346.

12. Wang, Z.; Kadouri, D.E.; Wu, M. Genomic Insights into an Obligate Epibiotic Bacterial Predator: Micavibrio Aeruginosavorus ARL-13. *BMC Genomics* 2011, *12*, 453, doi:10.1186/1471-2164-12-453.

13. Ye, X.-S.; Chen, M.-X.; Li, H.-Y.; He, X.-Y.; Zhao, Y. Halobacteriovorax Vibrionivorans Sp. Nov., a Novel Prokaryotic Predator Isolated from Coastal Seawater of China. *Int. J. Syst. Evol. Microbiol.* 2019, *69*, 3917–3923, doi:10.1099/ijsem.0.003703.

14. Karunker, I.; Rotem, O.; Dori-Bachash, M.; Jurkevitch, E.; Sorek, R. A Global Transcriptional Switch between the Attack and Growth Forms of Bdellovibrio Bacteriovorus. *PLoS ONE* 2013, *8*, e61850, doi:10.1371/journal.pone.0061850.

15. Davis, S.C.; Cerra, J.; Williams, L.E. Comparative Genomics of Obligate Predatory Bacteria Belonging to Phylum *Bdellovibrionota* Highlights Distribution and Predicted Functions of Lineage-Specific Protein Families. *mSphere* 2024, *9*, e00680-24, doi:10.1128/msphere.00680-24.

16. Waite, D.W.; Chuvochina, M.; Pelikan, C.; Parks, D.H.; Yilmaz, P.; Wagner, M.; Loy, A.; Naganuma, T.; Nakai, R.; Whitman, W.B.; et al. Proposal to Reclassify the Proteobacterial

Classes Deltaproteobacteria and Oligoflexia, and the Phylum Thermodesulfobacteria into Four Phyla Reflecting Major Functional Capabilities. *Int. J. Syst. Evol. Microbiol.* 2020, *70*, 5972–6016, doi:10.1099/ijsem.0.004213.

17. Cao, P.; Dey, A.; Vassallo, C.N.; Wall, D. How Myxobacteria Cooperate. *J. Mol. Biol.* 2015, *427*, 3709–3721, doi:10.1016/j.jmb.2015.07.022.

18. Hungate, B.A.; Marks, J.C.; Power, M.E.; Schwartz, E.; Van Groenigen, K.J.; Blazewicz, S.J.; Chuckran, P.; Dijkstra, P.; Finley, B.K.; Firestone, M.K.; et al. The Functional Significance of Bacterial Predators. *mBio* 2021, *12*, e00466-21, doi:10.1128/mBio.00466-21.

19. Kaiser, D.; Robinson, M.; Kroos, L. Myxobacteria, Polarity, and Multicellular Morphogenesis. *Cold Spring Harb. Perspect. Biol.* 2010, *2*, a000380–a000380, doi:10.1101/cshperspect.a000380.

20. Reichenbach, H. The Ecology of the Myxobacteria. *Environ. Microbiol.* 1999, *1*, 15–21, doi:10.1046/j.1462-2920.1999.00016.x.

21. Steinegger, M.; Söding, J. MMseqs2 Enables Sensitive Protein Sequence Searching for the Analysis of Massive Data Sets. *Nat. Biotechnol.* 2017, *35*, 1026–1028, doi:10.1038/nbt.3988.

22. Domazet-Lošo, T.; Tautz, D. A Phylogenetically Based Transcriptome Age Index Mirrors Ontogenetic Divergence Patterns. *Nature* 2010, *468*, 815–818, doi:10.1038/nature09632.

23. Zhang, L.; Huang, X.; Zhou, J.; Ju, F. Active Predation, Phylogenetic Diversity, and Global Prevalence of Myxobacteria in Wastewater Treatment Plants. *ISME J.* 2023, *17*, 671–681, doi:10.1038/s41396-023-01378-0.

24. Muñoz-Dorado, J.; Marcos-Torres, F.J.; García-Bravo, E.; Moraleda-Muñoz, A.; Pérez, J. Myxobacteria: Moving, Killing, Feeding, and Surviving Together. *Front. Microbiol.* 2016, *7*, doi:10.3389/fmicb.2016.00781.

25. Sockett, R.E.; Lambert, C. Bdellovibrio as Therapeutic Agents: A Predatory Renaissance? *Nat. Rev. Microbiol.* 2004, *2*, 669–675, doi:10.1038/nrmicro959.

26. Gu, Z. Complex Heatmap Visualization. *iMeta* 2022, *1*, e43, doi:10.1002/imt2.43.