Habitat complexity reduces feeding strength of freshwater predators

Mireia Aranbarri[1], Lorea Flores[1], Ioar de Guzmán[1], Aitor Larrañaga[1], Arturo Elosegi[1],
Björn C. Rall[2] and Julia Reiss[3]*

[1]*Laboratory of Stream Ecology; Dept. of Plant Biology and Ecology, Fac. of Science and Technology;
University of the Basque Country, UPV/EHU PO Box 644; 48080 Bilbao; Spain*

[2]*Aquatic Ecology and Evolution Group, Limnological Institute, University of Konstanz, Mainaustraße
252, 78464 Konstanz/Egg, Germany*

[3]*Centre for Pollution Research and Policy, Brunel University of London, Uxbridge, UB8 3PH, UK.*

* corresponding author
Email: julia.reiss@roehampton.ac.uk (JR)

ORCID: MA* [0009-0001-3506-0914], LF [0000-0002-0082-4072], IdG [0000-0001-8894-8477],
AL [0000-0002-0185-9154], AE [0000-0001-8809-8484], BCR [0000-0002-3191-8389], and JR*
[0000-0002-3740-0046].

Abstract

The physical structure of an environment potentially influences feeding interactions among organisms by providing refuges for prey. We examined how habitat complexity affects the functional feeding response of an ambush predator (damselfly larvae *Ischnura elegans*) and a pursuit predator (backswimmer *Notonecta glauca*) feeding on the isopod *Asellus aquaticus*. We run experiments in aquatic microcosms with an increasing number of structural elements (0, 2, or 3 rings of plastic plants in different spatial configurations), resulting in five habitat complexity levels. Across these levels, predators were presented with different prey densities to determine their functional response. The experimental design and statistical analysis allowed us to test for effects of structure presence, amount, and complexity level on functional response in one pass, without confounding predictors. The feeding for both predators across all complexity levels was best described by a Type II functional response model and habitat drove feeding strength. Regarding the latter, the predators showed different responses to the complexity treatments. The overall feeding rate of *I. elegans* was mainly driven by the absence vs. presence of structure. Yet, in the case of *N. glauca* feeding rate was strongly dependent on habitat complexity with the predator showing unique maximum feeding rates (i.e. the inverse of the handling time) for each complexity level and a decreasing attack rate with increasing amount of habitat. Overall, prey consumption was less than half for both predators when complex structures were present, compared to the no habitat structure environment. Our findings demonstrate that habitat complexity dampens feeding rates and therefore plays a key role in the stability of freshwater ecosystems.

Keywords: habitat complexity, functional response, trophic interactions, predation. stability

Introduction

The physical structure of an environment is a crucial factor in shaping community composition and the ecological processes driven by living organisms. Habitat complexity has been shown to influence population density, body size, and species richness of invertebrates in both terrestrial and aquatic ecosystems (Flores et al. 2016, White and Walsh 2020, Soukup et al. 2022). Furthermore, habitat complexity is also likely to influence the interactions among organisms by providing more refuge, thereby reducing predation risk for prey (Hauzy et al. 2010, Vucic-Pestic et al. 2010, Vucic-Pestic et al. 2010, Orrock et al. 2013, Barrios-O'Neill et al. 2015). Indeed, previous studies have demonstrated that increasing habitat complexity can lower predation pressure on prey, thus reducing the top-down control exerted by predators (Kalinkat et al. 2013, Mocq et al. 2021, Chang and Todd 2023). Yet habitat complexity could also aid predators in their search for prey, particularly those operating in three-dimensional environments (Pawar et al. 2012), as observed in the case of a freshwater copepod preying on ciliates (Reiss and Schmid-Araya 2011). Therefore, habitat complexity can potentially modify or shape energy flows through food webs, influencing feeding relationships between resources and consumers and among predators and prey.

In general, feeding rates are determined by the time required for predators to find, attack, capture, handle, and ingest their prey (Holling 1959a, 1959b). Testing how refuge availability and habitat complexity alter trophic relationships requires careful experimental design to avoid confounding the amount of structure with habitat complexity (St. Pierre and Kovalenko 2014, Flores et al. 2016). For instance, performing a laboratory experiment, Kalinkat et al. (2013) attributed the observed changes in feeding rates to a dilution effect (i.e. higher amount of structure, leaves in that case, had a dilution effect on prey density) and observed that attack

rates by the predator did not change per unit surface. Nevertheless, complexity can also increase

through changes in the arrangement of a fixed number of structures rather than simply adding

more, which must be considered in experimental designs.

Functional response models are a suitable tool when it comes to addressing the effects

of habitat complexity on species interactions, as they describe the relationship between prey

density and the intake of prey by a predator (Holling, 1959a, b). The feeding rate usually follows

either a hyperbolic or s-shaped (sigmoidal) relationship with increasing prey density (see e.g.

(Kalinkat et al. 2023), also known as Type II and Type III functional response, respectively (we

describe the models in the methods section, also see (Rall et al. 2025).

The hyperbolic shape of a type II functional response is determined by an initial increase

of the feeding rate, driven by the search speed (a.k.a. attack rate, capture rate, filtration rate,

and others) and the handling time, which controls satiation of the response (Holling 1959a). In

the case of a type III functional response (Holling 1959a), the attack rate itself is a function of

prey density, which is either linear or a power law function (Jeschke et al. 2002, Vucic-Pestic

et al. 2010, Kalinkat et al. 2013, Pritchard et al. 2017, Rosenbaum and Rall 2018, Juliano et al.

2022, Rall et al. 2025). In fact, habitat complexity has been put forward as driving a type III

response (Vucic-Pestic et al. 2010, Vucic-Pestic et al. 2010, Kalinkat et al. 2013, Barrios-

O'Neill et al. 2015).

In addition to habitat complexity, the predation strategy of the predator is an important

factor determining the interaction strength and by extension the stability of food webs (Almany

2004, Schmitz 2017). These strategies include "ambush" (also known as "sit-and-wait" tactics)

and "pursuit" (active chasing), and these traits can determine spatial utilization and interactions

with environmental structures (Pawar et al. 2012, Schmitz 2017). For instance, ambush

predators, exemplified in aquatic systems by species like dragonfly and damselfly nymphs, occur in complex habitats that provide ample hiding spots, enhancing their ability to remain concealed and ambush prey (Mocq et al. 2021, Soukup et al. 2022, Chang and Todd 2023). Conversely, pursuit predators, such as predatory fish, certain types of predatory shrimp or backswimmers require more open and less obstructed spaces to effectively chase down their prey and dense structures within complex habitat can impede their navigation and reduce hunting efficiency (Almany, 2004; Warfe & Barmuta, 2006). Consequently, habitat complexity is likely to have varying impacts on predatory success depending on the predator's hunting strategy. For instance, in aquatic environments, a more complex habitat may favour ambush predators by providing increased hiding opportunities, while posing challenges for active swimming predators, thereby potentially reducing their hunting efficiency(Almany 2004, Dunn and Hovel 2020, Froneman and Cuthbert 2022). Studying how functional response of different predators responds to habitat structure is crucial to understand trophic interactions and energy transfer through real food webs in nature.

Our aim was to explore how two different predators with contrasting hunting strategies are affected by habitat complexity in their search and consumption of prey. Specifically, we tested how habitat complexity (measured as (1) the presence of structure, (2) the amount of structure and (3) the spatial arrangement of structure) influenced the functional response parameters, addressing three hypotheses:

1. Habitat complexity alters the shape of the functional response curve, as more structure can create refuges for prey and makes detection of prey more challenging, particularly at low prey densities.

2. Feeding rates generally decrease with increasing amount of habitat structure resulting in altered functional response parameters.

119   3.   The pursuit predator (*N. glauca*) will show a different response to habitat structure

120   compared to the ambush predator (*I. elegans*).

Material and Methods

Experimental set-up

The experiment was carried out using individual microcosms (9.5 cm diameter and 11.5 cm

high circular beakers). The prey (water hog-louse *Asellus aquaticus*) was provided with leaves

to feed on. Hence, air-dried alder leaves (*Alnus glutinosa* L.) were weighed, added to each

microcosm (2 g each) and conditioned in 500 mL water (1 part of filtered pond water to 5 parts

of tap water) for 7 days. The water was renewed when the prey was introduced to the

microcosms. Every microcosm was served by one air-stone, and it was sealed with cling film

to prevent evaporation. The experiment was run in a controlled-temperature room (15 ℃) with

a 14 h light:10 h dark cycle.

When manipulating habitat complexity, it was important to disentangle the effects of

complexity *per se* from the effects of amount of structure in the microcosm (see Kalinkat et al.

2013 for reasoning). We used plastic rings to create both structure and different levels of

complexity. These rings were artificial plastic plants mimicking the aquatic plant

Ceratophyllum spp. (Code No. FRF 491, Fish are Fun®). Habitat complexity was created in

three ways: 1. as the presence vs. absence of habitat structure; 2. as the amount of structure (0,

2 or 3 of plastic plant rings); and 3. as two spatial configurations of structures in the case of 2

and 3 rings (see Figs. 1&2 and Flores et al. 2016), resulting in 5 complexity levels. The order

of these five levels (from 'simple' to 'complex' [levels 0, 1, 2, 3 and 4] was determined by

measuring the fractal dimension in the feeding arena (see Flores et al. 2016).

Feeding functional response

We tested the feeding functional response of the ambush predator *Ischnura elegans* (van der

Linden) (16.8 mm average length and 3.5 mm average head width) and a pursuit predator

Notonecta glauca (L.) (14.7 mm average length and 4.9 mm wide) preying on *Asellus* (6.6 mm

average body length).

The predators were able to get into the structures added to the microcosms, which thus,

did not function as a full refuge for prey. The predators were allowed to feed for 24 hours. We

used time blocks to run a replicate each, which were all started two days apart from each other

because of the time needed to sort *A. aquaticus* individuals. Further, the blocks had the

advantage that this approach allowed us to adjust the very high densities after running the first

replicate and to further adjust the number of replicates needed after observing feeding during

the first time block. Prey densities for *I. elegans* were 1, 3, 5, 10, 30, 80 and 120 individuals per

microcosm. Prey densities for *N. glauca*, were 1, 3, 5, 10, 30, 80, 120 and 180 individuals per

microcosm 1, 3, 5, 10, 30, 80, 120, and 180 individuals per microcosm. In total we run 297

microcosms. The data are available in Flores et al. (2025)(Flores et al. 2025).

Data Analysis

We first investigated if the shape of the functional response was a hyperbolic type II or s-shaped

type III for all ten combinations of complexity treatment and predator species. This was

achieved using a phenomenological test (Pritchard et al. 2017, Juliano 2020) and validated by

fitting a so-called generalized functional response model allowing for a smooth shift between

type II and type III functional responses (Rosenbaum and Rall 2018). A full supplemental

statistics report (Rall et al. 2025) and the code are available.

Holling (1959a) described the mechanics of predation mathematically:

$$F = \frac{aN}{(1+aT_h N)}$$

where the feeding rate, F, is a hyperbolic function in which the attack rate, a, drives the initial

T_h slope, and the handling time, h, controls the satiation and N is the density of prey. While we

discuss and explain all data analysis in-depth in Rall et al. (2025), it is worth noting that, as an

'updated approach' to Holling, the generalized functional response model (Real 1977, Williams

and Martinez 2004, Vucic-Pestic et al. 2010, Rosenbaum and Rall 2018) is:

$$F = \frac{F_{max}N^{1+q}}{N_{half}^{1+q}+N^{1+q}} = \frac{bN^{1+q}}{1+bT_h N^{1+q}} \tag{1}$$

where the feeding rate F depends on the prey density N. F_{max} is maximum feeding rate (Real

1977), N_{half} is the half saturation density (Real 1977), T_h is the handling time (Holling 1959b),

b is the attack coefficient (Vucic-Pestic et al. 2010) and q is the shape parameter (Williams and

Martinez 2004, Vucic-Pestic et al. 2010, Rosenbaum and Rall 2018). If q is „0", the function

becomes a type II functional response (Holling 1959b), and if q is (significantly) greater than

„0", the function becomes s-shaped with $q = 1$ being a 'strict' type III functional response often

used if analyzing functional response experiments (e.g. Kalinkat et al. 2023). We will refer to

the models with the parameters attack coefficient and handling type as Holling type of

functional response, and with the parameters maximum feeding rate and half saturation density

as Real type of functional response. Both versions can be translated into each other by

$$F_{max} = \frac{1}{T_h} \text{ and } N_{half} = \frac{1}{(bT_h)^{1/1+q}} \tag{2}.$$

The attack rate a, which is often used in functional response analyses is a function of prey

density if $q > 0$ and can be described by $a = bN^q$ (Vucic-Pestic et al. 2010). The attack rate,

a, becomes the attack coefficient b if $q = 0$ (Type II functional response).

After determining the shape of the functional response, we tested a suite of functional

response models to address habitat effects on the functional response parameters (see the

supplemental statistical report, Rall et al. 2025, for details). This suite of models tested for 1)

the assumption that habitat does not influence feeding, 2) an effect of no structure vs. structure,

3) for an effect caused by the amount of structure and 4) for effects due to the complexity of

the structure (while being related to 'amount'). The models targeted two parameters (case type

II or strict type III) or three parameters (case generalized functional response) at once (Table

1). We dubbed the former 'Holling' and the latter 'Real' as these authors were the first to

introduce these parameters (Holling 1959a, 1959b, Real 1977). Testing four assumptions on

two parameters (type II and strict type III functional response) in a fully factorial fashion gives

a suite of 16 models for the 'Holling' (1H to 16H) and the 'Real' type (1R to 16R) of functional

responses each (Table 1). Consequently, models 1R to 16R targeted F_{max} and N_{half} (Real type)

and models 1H to 16H targeted parameters T_h and a. In case we detected any type III functional

response (see above) the full-factorial scheme would need 64 models for each functional

response formulation (see the supplemental statistics report, Rall et al. 2025, for details).

This approach allowed testing for these different habitat predictors on all functional

response parameters in one pass and avoided confounding 'amount of structure' with

'complexity', based on the rationale developed in Flores et. al. (2016). We compared the models

by using both the Akaike's Information Criterion (AIC) and the Bayesian information criterion

(BIC), where the lowest AIC and BIC indicates the best (most parsimonious) model. The output

includes a point estimate for the parameter and lower and upper confidence intervals (CIs).

207   Hence, the functional response models provide information about the overall amount of

208 prey eaten, driven by the maximum feeding rate or its inverse, the handling time. This effect is

209 most obvious at high prey densities where the curve reaches its asymptote. The attack rate, and

210 the half saturation density predominantly control the feeding rate at lower densities, and we

211 were able to compare those parameters between habitat treatments.

Results

The shape of the functional response curves for both *N. glauca* and *I. elegans* consistently fit best to a type II model for all five complexity levels and this result held for both the classical phenomenological test and for fitting the generalized functional response model directly to data (see Supplementary Information Tables 1 and 2).

Therefore, we tested a total of 32 type II functional response models (Table 1) to test for the effect of 'habitat'. For the functional response exhibited by *Ischnura elegans,* the best fitting model (both lowest AIC and BIC, Table 2) was 'model 5R' (Table 1, Figure 1). This model assumed that the effect of structure presences vs. absence overrides all other possible predictors. In the '5R' model, F_{max} takes a value of 28 prey eaten in 24 h when no structure is present (CI: 19.4 to 40.8). However, when structure is present, then F_{max} is 14.6 (CI: 10.7 to 29). N_{half} is not affected by the habitat and takes the value of 57 prey items per arena (CI: 34 to 94).

The best fitting (most parsimonious) model (both lowest AIC and BIC) for *Notonecta glauca* was 'model 15H' and this model assumed that each complexity level has a unique handling time, T_h, and that attack rate, a, decreases with increasing amount of habitat structure. Hence, the parameters that affect feeding rate are more complex in this case. Maximum feeding rate (from the handling time ($F_{max} = {}^1\!/_{T_h}$)) was 31 (CI: 26 to 35) prey eaten in 24 h for complexity level 0 (no structure, level 1), 18 (CI: 15-21), for level 2, 28 (CI: 24-33), 21 (CI: 17 to 27) for level 3 and 19 (CI: 15 to 23) for level 4.The attack rate decreased with increasing number of habitat structural elements (rings, $log_{10}(a) = a_{intercept} + a_{slope}N_{rings}$), indicated by a negative slope (-0.144, CI: -0.191 to -0.097).

The reason why this more complex model was significant (for the maximum feeding

rate) over the very similar, but simpler, 'presence of structure' model, was that 'unexpected'

responses were observed when two plastic plant rings had been added to the microcosms

(Figure 3). A possible explanation is that the configuration of 2 rings into two different shapes

creates very different habitat types and that this is not the case when three rings were used (the

latter essentially created an ideal hiding spot in an any case, a 'dense ball'). For two rings, it is

likely that one of the configurations left more space for the agile pursuit predator *Notonecta* to

navigate in.

Overall, fitting these models showed that the amount of prey eaten was generally higher

without habitat compared to all high habitat complexity levels. On average, prey consumption

was less than half for both predators when complex structures were present, compared to the

no habitat structure environment. Both predators consumed significantly more prey in the

absence of structures, with average consumption reaching 5-6 times their body weight in 24

hours.

Discussion

Our findings suggest that habitat complexity affects predator-prey interactions in freshwater

ecosystems. Results from our feeding experiments showed that feeding, i.e. the functional

response type, did not significantly vary with habitat complexity (all feeding was consistent

with a type II response). However, we found support for the hypothesis that increased habitat

complexity would decrease feeding rates overall. Both, ambush predator (damselfly larvae

Ischnura elegans) and pursuit predators (backswimmer *Notonecta glauca*) showed higher

feeding rates in microcosms without structures compared to those with structure. This suggests

that while habitat complexity reduced feeding rates by affecting maximum feeding rate, attack

rate and handling time, the overall mechanisms (shape of the functional response curve)

remained consistent across different complexity levels.

Generally, it is known that complex structures offer refuge for prey, thereby reducing

the efficiency of predators (Hauzy et al. 2010, Mocq et al. 2021, Froneman and Cuthbert 2022).

This reduction in feeding rates due to increased habitat complexity implies a decrease in the

top-down control of predators on prey populations(Chang and Todd 2023). In our study, the

reduction in feeding rates was associated with both a reduction in attack rate and an increase in

handling time (i.e. an increase in maximum feeding rate while the half saturation density is

relatively constant). However, the magnitude of these changes varied across different

complexity levels. Previous studies, conducted using microcosm approaches, have shown that

attack rate and handling time can respond differently to habitat complexity. For instance, a

study on zooplankton prey *Aracartia longipatella* and predator *Mesopodopsis wooldridgei*

reports stronger effects on attack rates (Froneman and Cuthbert 2022); while the dragonfly

larvae *Aeshna cyanea* preying on *Chaoborus obscuripes* larvae changes handling time with

complexity (Mocq et al. 2021). Soil mites preying on collembolans displayed varying response

in both attack rate and handling time with habitat structure (Hauzy et al. 2010).

We hypothesized that at lowest habitat complexity (i.e. microcosms without added

structures), the shape of the functional response would best fit to a type II functional response,

because feeding rates would be mainly limited by the time required by the predator to kill and

eat its prey. With increasing habitat complexity levels, we expected sigmoid functional response

(i.e. type III functional response), as at low prey densities the efficiency of the predator would

be limited by the structures offering multiple hiding places (Vucic-Pestic et al. 2010a,b,

Kalinkat et al. 2023). However, as prey density increases and the amount of refuge become

saturated, the efficiency of the predator would again increase at intermediate prey densities.

Nevertheless, functional response of both predators was consistently best described by a type

II model across all the levels of habitat complexity. For *Ischnura*, (Mikolajewski et al. 2015)

highlighted a phenotypic plasticity, allowing it to adapt to environmental changes while

maintaining a Type II functional response. Moreover, our design of the habitat structure allowed

both predators to enter the potential prey refuge which may be also a reason for not finding a

type III response. One prey per predator was the lowest density in our experiment, which

corresponds to a density of 141.1 individuals per square metre, but in nature, lower abundances,

can be observed for many taxa (Larrañaga et al. 2009). Indeed, not considering low prey in

laboratory set-ups can hinder the detection of a type III response (Sarnelle and Wilson 2008).

Although the prevalence of type III functional response has been postulated as the way to

maintain stability of prey populations and high diversity (Murdoch and Oaten 1975, Rall et al.

2008, Kalinkat et al. 2023), type II is still the most common functional response in previous

studies (Dunn and Hovel 2020).

Although the type III functional response was not supported by our data, we saw a hint

that the interaction of both predators and its prey was stabilizing as feeding rates generally

decreased with the presence and with increasing complexity of the habitat structure. The general

decrease of interaction strength is associated with more stable population dynamics and higher

species richness in model food webs (May 1972, Yodzis and Innes 1992, Rall et al. 2008).

The two predators in our experiment differed in terms of their predatory strategies and

the functional response parameters highlighted some of those differences. The pursuit predator

(the backswimmer *N. glauca*) exhibited three times the attack rate of the ambush predator (the

damselfly *I. elegans*), although a similar handling time when no structures were present in their

environment. Also, as an open-water hunter, *N. glauca* is more likely to operate in a 3D environment compared to *I. elegans* that sits within/ on habitat surface (see Pawar et al. 2012). Indeed, dimensionality of consumer search space is probably a major driver of species coexistence, and the stability and abundance of populations (Pawar et al. 2012). As predicted, the effect of habitat complexity was stronger for the pursuit predator than the ambush predator. An explanation is the barrier effect that the structures can create for the pursuit predator, reducing the visibility of prey and the capture success(Grabowski and Kimbro 2005). In agreement with(Svensson et al. 2000), habitat heterogeneity significantly reduced predation efficiency in *N. glauca* by providing refuges that lower prey visibility and decrease attack rates, highlighting the vulnerability of pursuit predators to habitat complexity. Previous studies have also shown that *Notonecta* spp. are more successful in open environments(Gittelman 1974), and that factors such as water depth and refuge availability influence its hunting behaviour (Cockrell 1984).

Our short-term experiment with three species certainly demonstrated that habitat complexity is an important factor to be considered when it comes to trophic interactions. Both predators were more successful in environments without structure and were able to feed on substantially more prey without structure.

Conclusion

We conclude that habitat complexity has the potential to shape species interactions and carbon flow within the food web. The results of this study demonstrate that increased habitat complexity significantly reduces feeding rates in both *Ischnura elegans* and *Notonecta glauca*. The reduction in predation efficiency, driven by decreases in attack rates and increases in handling times, supports the idea that structural complexity provides refuges for prey, limiting

predator success. Despite these changes in feeding rates, both species maintained a Type II functional response across all levels of habitat complexity, indicating that while habitat structure affects predator efficiency, it does not fundamentally alter the shape of the functional response curve. Overall, these findings underscore the critical role of habitat complexity in shaping trophic interactions and sustain energy flow. In more complex, natural ecosystems, habitat complexity will likely drive species coexistence, stability of food webs and diversity of communities.

Funding Information

MA was funded by the Investigo Programm funded by the NextGenerationEU initiative, LF was funded by a grant by the Spanish Ministry of Education and Culture, IdG was funded by the Spanish Ministry of Science, Innovation and Universities (TED2021-129966B-C31). JR was supported by a Royal Society of London Starting Grant. Björn C. Rall gratefully acknowledges the funding by the German Science Foundation (DFG) to the Research Unit DynaSym (FOR 5064).

Acknowledgements

We thank the WWT London for the permission to sample damselflies and backswimmers on their grounds and Szymon Szary for his assistance in the field and laboratory.

References

Almany, G. R. 2004. Does increased habitat complexity reduce predation and competition in coral reef fish assemblages? Oikos 106:275–284.

Barrios-O'Neill, D., J. T. A. Dick, M. C. Emmerson, A. Ricciardi, and H. J. MacIsaac. 2015. Predator-free space, functional responses and biological invasions. Functional Ecology 29:377–384.

Chang, C., and P. A. Todd. 2023. Reduced predation pressure as a potential driver of prey diversity and abundance in complex habitats. npj Biodiversity 2.

Cockrell, B. J. 1984. Effects of water depth on choice of spatially separated prey by Notonecta glauca L. Oecologia 62:256–261.

Dunn, R. P., and K. A. Hovel. 2020. Predator type influences the frequency of functional responses to prey in marine habitats. Biology Letters 16:20190758.

Flores, L., R. A. Bailey, A. Elosegi, A. Larrañaga, and J. Reiss. 2016. Habitat complexity in aquatic microcosms affects processes driven by detritivores. PLoS One 11:e0165065.

Flores, L., J. Reiss, A. Larrañaga, B. C. Rall, M. Aranbarri, and I. de Guzmán. 2025. Complexity reduces feeding strength of freshwater predators (CRITTER) - Data.

Froneman, P. W., and R. N. Cuthbert. 2022. Habitat complexity alters predator-prey interactions in a shallow wate r ecosystem. Diversity 14.

Gittelman, S. H. 1974. Locomotion and predatory strategy in backswimmers (Hemiptera: Notonectidae). American Midland Naturalist:496–500.

Grabowski, J. H., and D. L. Kimbro. 2005. Predator-avoidance behavior extends trophic

cascades to refuge habitat s. Ecology 86:1312–1319.

Hauzy, C., T. Tully, T. Spataro, G. Paul, and R. Arditi. 2010. Spatial heterogeneity and

functional response: an experiment in microcosms with varying obstacle densities.

Oecologia 163:625–636.

Holling, C. S. 1959a. The components of predation as revealed by a study of small-mammal

predation of the European Pine Sawfly. The Canadian Entomologist 91:293–320.

Holling, C. S. 1959b. Some characteristics of simple types of predation and parasitism. The

Canadian Entomologist 91:385–398.

Jeschke, J. M., M. Kopp, and R. Tollrian. 2002. Predator functional responses: discriminating

between handling and digesting prey. Ecological monographs 72:95–112.

Juliano, S. A. 2020. Nonlinear curve fitting: predation and functional response curves. Pages

159–182 Design and analysis of ecological experiments. Chapman and Hall/CRC.

Juliano, S. A., J. A. Goughnour, and G. D. Ower. 2022. Predation in many dimensions: spatial

context is important for meaningful functional response experiments. Frontiers in Ecology

and Evolution 10:845560.

Kalinkat, G., U. Brose, and B. C. Rall. 2013. Habitat structure alters top-down control in litter

communities. Oecologia 172:877–887.

Kalinkat, G., B. C. Rall, S. F. Uiterwaal, and W. Uszko. 2023. Empirical evidence of type III

functional responses and why it remains rare. Frontiers in Ecology and Evolution

11:1033818.

Larrañaga, A., A. Basaguren, A. Elosegi, and J. Pozo. 2009. Impacts of Eucalyptus globulus

plantations on Atlantic streams: change s in invertebrate density and shredder traits.

Fundamental and Applied Limnology 175:151–160.

May, R. M. 1972. Will a large complex system be stable? Nature 238:413–414.

Mikolajewski, D. J., A. Conrad, and G. Joop. 2015. Behaviour and body size: plasticity and

genotypic diversity in larval *Ischnura elegans* as a response to predators (Odonata:

Coenagrio nidae). International Journal of Odonatology 18:31–44.

Mocq, J., P. R. Soukup, J. Näslund, and D. S. Boukal. 2021. Disentangling the nonlinear effects

of habitat complexity on functiona l responses. Journal of Animal Ecology 90:1525–1537.

Murdoch, W. W., and A. Oaten. 1975. Predation and population stability. Pages 1–131

Advances in ecological research. Elsevier.

Orrock, J. L., E. L. Preisser, J. H. Grabowski, and G. C. Trussell. 2013. The cost of safety:

refuges increase the impact of predation risk in aquatic systems. Ecology 94:573–579.

Pawar, S., A. I. Dell, and V. M. Savage. 2012. Dimensionality of consumer search space drives

trophic interaction strengths. Nature 486:485–489.

St. Pierre, J. I., and K. E. Kovalenko. 2014. Effect of habitat complexity attributes on species

richness. Ecosphere 5:1–10.

Pritchard, D. W., R. A. Paterson, H. C. Bovy, and D. Barrios-O'Neill. 2017. Frair: an R package

for fitting and comparing consumer functional responses.

Rall, B. C., M. Aranbarri, L. Flores, I. de Guzmán, A. Larrañaga, and J. Reiss. 2025. Complexity

reduces feeding strength of freshwater predators (CRITTER) - Supplemental Statistics

Report.

Rall, B. C., C. Guill, and U. Brose. 2008. Food-web connectance and predator interference

dampen the paradox of enrichment. Oikos 117:202–213.

Real, L. A. 1977. The kinetics of functional response. The American Naturalist 111:289–300.

Reiss, J., and J. M. Schmid-Araya. 2011. Feeding response of a benthic copepod to ciliate prey

type, prey concentration and habitat complexity. Freshwater Biology 56:1519–1530.

Rosenbaum, B., and B. C. Rall. 2018. Fitting functional responses: Direct parameter estimation

by simulating differential equations. Methods in Ecology and Evolution 9:2076–2090.

Sarnelle, O., and A. E. Wilson. 2008. Type III functional response in Daphnia. Ecology

89:1723–1732.

Schmitz, O. 2017. Predator and prey functional traits: understanding the adaptive machin ery

driving predator–prey interactions. F1000Research 6:1767.

Soukup, P. R., J. Näslund, J. Höjesjö, and D. S. Boukal. 2022. From individuals to communities:

Habitat complexity affects all levels of organization in aquatic environments. John Wiley

and Sons Inc.

Svensson, B. G., B. Tallmark, and E. Petersson. 2000. Habitat heterogeneity, coexistence and

habitat utilization in five backswimmer species (Notonecta spp.; Hemiptera,

Notonectidae). Aquatic Insects 22:81–98.

Vucic-Pestic, O., K. Birkhofer, B. C. Rall, S. Scheu, and U. Brose. 2010. Habitat structure and

prey aggregation determine the functional response in a soil predator–prey interaction.

Pedobiologia 53:307–312.

Vucic-Pestic, O., B. C. Rall, G. Kalinkat, and U. Brose. 2010. Allometric functional response

model: body masses constrain interaction strengths. Journal of Animal Ecology 79:249–

256.

White, J. Y., and C. J. Walsh. 2020. Catchment-scale urbanization diminishes effects of habitat

complexity on instream macroinvertebrate assemblages. Ecological Applications

30:e02199.

Williams, R. J., and N. D. Martinez. 2004. Stabilization of chaotic and non-permanent food-

web dynamics. The European Physical Journal B 38:297–303.

Yodzis, P., and S. Innes. 1992. Body size and consumer-resource dynamics. The American

Naturalist 139:1151–1175.

Tables

Table 1: Overview of models tested on 2 parameters at once to show if the feeding rate was affected by habitat complexity. Four parameters are tested, yet only 2 of them in one pass as this reduced the testing from a possible 128 combinations to 32 models. Also, the two parameters correspond to different simulation models (after Real 1977 and Holling 1959 respectively). Levels (used to calculate df of the models) are for hypothesis: One: presence has effect—2 levels (present and absent); Two: amount has effect—3 levels, zero, 2 and 3 plastic plant rings and Three: complexity level (fractal dimension) has effect —5 levels: 0-4.

	'Holling (1959) approach'								'Real (1977) approach'							
Parameter	T_h				a				F_{max}				N_{half}			
Hypothesis	zero	one	two	three	zero	one	two	three		zero	one	two	three	zero	one	two	three
	habitat has no effect	presence has effect	amount has effect	complexity level has effect	habitat has no effect	presence has effect	amount has effect	complexity level has effect		habitat has no effect	presence has effect	amount has effect	complexity level has effect	habitat has no effect	presence has effect	amount has effect	complexity level has effect
Model									**Model**								
1H	1	0	0	0	1	0	0	0	1R	1	0	0	0	1	0	0	0
2H	1	0	0	0	0	1	0	0	2R	1	0	0	0	0	1	0	0
3H	1	0	0	0	0	0	1	0	3R	1	0	0	0	0	0	1	0
4H	1	0	0	0	0	0	0	1	4R	1	0	0	0	0	0	0	1
5H	0	1	0	0	1	0	0	0	5R	0	1	0	0	1	0	0	0
6H	0	1	0	0	0	1	0	0	6R	0	1	0	0	0	1	0	0
7H	0	1	0	0	0	0	1	0	7R	0	1	0	0	0	0	1	0
8H	0	1	0	0	0	0	0	1	8R	0	1	0	0	0	0	0	1
9H	0	0	1	0	1	0	0	0	9R	0	0	1	0	1	0	0	0
10H	0	0	1	0	0	1	0	0	10R	0	0	1	0	0	1	0	0
11H	0	0	1	0	0	0	1	0	11R	0	0	1	0	0	0	1	0
12H	0	0	1	0	0	0	0	1	12R	0	0	1	0	0	0	0	1
13H	0	0	0	1	1	0	0	0	13R	0	0	0	1	1	0	0	0
14H	0	0	0	1	0	1	0	0	14R	0	0	0	1	0	1	0	0
15H	0	0	0	1	0	0	1	0	15R	0	0	0	1	0	0	1	0
16H	0	0	0	1	0	0	0	1	16R	0	0	0	1	0	0	0	1

1= model tests for this, 0= model does not test for this

449 **Table 2**: The six best models that describe the functional response parameter F_{max} and N_{half};

450 and T_h and a, for predators *Ischnura elegance* and *Notonecta glauca*. The full suite of models

451 can be found in Rall et al. (2025). The best fit of the models was tested using AIC (AIC

452 generally performs better for models that have more parameters) and BIC. A low value indicates

453 the best fit.

AIC			BIC		
Model	**df**	**dAIC**	**Model**	**df**	**dBIC**
Ischnura elegance					
Model 5R	**3**	**0.000**	**Model 5R**	**3**	**0.000**
Model 7R	4	0.265	Model 5H	3	1.583
Model 6H	4	0.952	Model 7R	4	2.731
Model 6R	4	0.952	Model 6H	4	3.418
Model 5H	3	1.583	Model 6R	4	3.418
Model 13R	6	2.107	Model 7H	4	4.718
Notonecta glauca					
Model 15H	**7**	**0.000**	**Model 15H**	**7**	**0.000**
Model 16H	10	1.274	Model 9R	3	1.004
Model 16R	10	1.274	Model 11H	4	1.485
Model 15R	7	1.990	Model 11R	4	1.485
Model 14R	7	4.189	Model 13R	6	1.507
Model 13R	6	4.854	Model 15R	7	1.990
Model 12H	7	8.352	Model 7H	4	2.516

Figure captions

Figure 1. Functional response curves for the ambush predator *I. elegans* for microcosm without structure and with structure (plastic plants).

Figure 2. Functional response curves for the pursuit predator *N. glauca* for each habitat complexity level (a-e are levels 0 to 4 respectively). Habitat complexity levels 1 and 2 (b, c) were created with two rings of plastic plants, while levels 3 and 4 (d, e) had three rings of plastic plants.

Figure 3. The functional feeding response parameter F_{max} of *Notonecta glauca* feeding on *Asellus aquaticus*. Amount of structure (number of plastic plant rings) is regressed against the feeding rate, and this shows that when two rings are added, there is an 'unexpected response' explaining why a more complex model that assumed an effect of fractal dimension is more parsimonious.

473 **Figures**

475   Figure 1

477   Figure 2

479   Figure 3

Supplementary Information

The full data and statistics can be found here:

1. Flores, L., J. Reiss, A. Larrañaga, B. C. Rall, M. Aranbarri, and I. de Guzmán. 2025. Complexity reduces feeding strength of freshwater predators (CRITTER) - Data. https://doi.org/10.5281/ZENODO.14891981

2. Rall, B. C., Aranbarri, M., Flores, L., de Guzmán, I., Larrañaga, A., & Reiss, J. (2025). Complexity reduces feeding strength of freshwater predators (CRITTER) - Supplemental Statistics Report. Zenodo. https://doi.org/10.5281/zenodo.14898819

3. GitHub Repository: https://github.com/b-c-r/CRITTERstatistics

SI Table 1: Testing for functional response type for two predators feeding at 5 complexity

levels (0 to 4), fitting a Polynomial to Proportion Feeding. The functional response pattern

consistently fits a type II pattern. Testing for functional response type using 'Juliano approach'.

We observed a declining proportion of eaten prey consistent with a type II functional responses

(i.e. both a negative linear and a positive quadratic term. There were three exceptions of this

pattern – at the highest complexity level, the results for I. elegans suggest a hump shaped

functional response pattern as the linear term is positive and the quadratic term is negative.

However, both estimates are not significant. Further, the N. glauca functional response for

complexity levels 3 and 4 show a negative quadratic term, which could be indicative of a type

III functional response, yet both results are not significant.

Predator	Complexity	Linear (Q)	Quadratic (Q)	Linear (L)	Type
Ischnura elegans	0	-(n.s.)	-(n.s.)	-(*)	II
Ischnura elegans	1	-(n.s.)	-(n.s.)	-(*)	II
Ischnura elegans	2	-(n.s.)	-(n.s.)	-(***)	II
Ischnura elegans	3	-(n.s.)	-(n.s.)	-(**)	II
Ischnura elegans	4	-(n.s.)	-(n.s.)	-(**)	II
Notonecta glauca	0	-(***)	-(***)	-(***)	II
Notonecta glauca	1	-(***)	-(***)	-(***)	II
Notonecta glauca	2	-(***)	-(**)	-(***)	II
Notonecta glauca	3	-(n.s.)	-(n.s.)	-(***)	II
Notonecta glauca	4	-(*)	-(n.s.)	-(***)	II

503 **SI Table 2**: Testing for functional response type using q-exponent (known as Hill exponent)

504 approach. The significance of the shape parameter q of all ten separately fitted generalized

505 functional response models. None of the q values is significantly different from zero, hence the

506 feeding rate follows a functional response type II, confirming results from SI Table 1.

Predator	Complexity	q	Significance	Type
Ischnura elegans	0	0.116	n.s.	II
Ischnura elegans	1	0.047	n.s.	II
Ischnura elegans	2	-0.001	n.s.	II
Ischnura elegans	3	-0.001	n.s.	II
Ischnura elegans	4	0.193	n.s.	II
Notonecta glauca	0	-0.011	n.s.	II
Notonecta glauca	1	-0.006	n.s.	II
Notonecta glauca	2	0.194	n.s.	II
Notonecta glauca	3	-0.002	n.s.	II
Notonecta glauca	4	-0.001	n.s.	II