The evolution of host exploitation by parasitoids:

the timing of attack and consumption

Ryuichiro Isshiki[1, *]

Ryosuke Iritani[2, 3]

[1]Research Center for Integrative Evolutionary Science, SOKENDAI (The Graduate University for Advanced Studies), Shonan Village, Hayama, Kanagawa 240-0193 Japan

[2]RIKEN Center for Interdisciplinary Theoretical and Mathematical Sciences, 2-1 Hirosawa, Wako, Saitama, 351-0198, Japan

[3]Department of Biology, Faculty of Sciences, The University of Tokyo, Japan

[*]e-mail: isshiki_ryuichiro@soken.ac.jp (Isshiki is corresponding author)

Country affiliation for the authors: Japan

ORCID: 0009-0006-5011-4657, 0000-0002-2396-1109

Acknowledgement

We thank Troy Day, Koichi Ito, Bo-Moon Kim, Raiki Nakano, Stuart West, and Atsushi Yamauchi for useful comments.

17 **Author contributions**

18 Ryuichiro Isshiki conceived the study, developed the theoretical framework, performed the

19 mathematical analyses, and drafted the manuscript. Ryosuke Iritani provided supervision,

20 critical feedback, and editorial revisions. All authors read and approved the final version of

21 the manuscript.

Abstract

Parasitoid wasps exhibit remarkable diversity in life-history traits and are categorized into two major groups based on the timing of emergence: those that begin consuming host tissues immediately after hatching ("idiobiont") and those that can delay the emergence depending on host maturation ("koinobiont"). Although delayed emergence allows parasitoids to exploit unparasitized, young hosts, it incurs a mortality cost during the waiting period. While numerous empirical studies have examined the adaptive significance of this trait, how host stage-structure and ecological factors jointly shape the evolution of emergence timing remains poorly understood in a formalized context. Here, we develop a stage-structured mathematical model to analyze the evolutionary dynamics of delayed emergence and its consequences for host exploitation. By explicitly incorporating host developmental stages, our model explores the joint evolution of two traits: the preference for young versus old host, and the timing of emergence. Our results reveal that the optimal strategy is determined by the relative reproductive value of host stages, which depends on the balance between host growth and mortality. We demonstrate that delayed emergence evolves when the future value of a growing host outweighs the immediate cost of mortality, thereby driving the divergence between idiobiont and koinobiont strategies. These findings provide a theoretical framework that unifies the diversity of parasitoid life histories.

Keywords: life-history traits, optimal foraging, parasitoid, reproductive value, stage structure

Significance Statement

Parasitoids exhibit striking diversity in life-history traits such as body size, fecundity, and clutch size, reflecting adaptations to host exploitation. A key behavioral distinction is whether parasitoids consume the host immediately (idiobionts) or can delay emergence while the host continues to grow (koinobionts). Many empirical studies have suggested and tested some hypothesis. On the other hand,

the adaptive significance of delayed emergence have not been formulated yet, so the quantitative

relationship between selection pressure and life history traits is yet to be better understood. Here, we

develop a mathematical model that integrates host growth with parasitoid emergence timing. Our

analysis supports the adaptive significance of delayed emergence and shows how delayed emergence can

promote diversification of parasitoid life-history strategies. These results suggest that delayed emergence

is a key driver of idiobiont–koinobiont dichotomy, providing new insights into the evolutionary basis of

parasitoid–host interactions and aligning directly with the central aims of behavioral ecology.

Introduction

Resource exploitation strategies, specifically their timing and allocation, are fundamental

life-history traits. Parasitoids, particularly wasps, serve as an excellent model system for

studying life-history traits, with their variations in parasitoids particularly pronounced

(Godfray 1994; Mayhew and Blackburn 1999; Jervis et al. 2008; Wajnberg et al. 2008;

Jervis and Ferns 2011; Quicke 2014): for example, the order *Hymenoptera* comprises at

least 200,000 species by a conservative estimate (Pennacchio and Strand 2006), probably

each utilizing different host species (mostly arthropods). Also, wasp species exhibit life-

history trait variations, by a factor of 18 in body-size (Jervis et al. 2003), more than one

hundred in clutch size and lifetime potential fecundity (Jervis et al. 2008), with these

variations presumably resulting from adaptation to host species. Parasitoid life-history traits

have significant impacts on numerous processes such as parasitoid-prey interactions

(Lafferty et al. 2008), policy making in pest management (Wang et al. 2019), and

coevolution (Amoroso and Antonovics 2020). Hence, improving our understanding of the

evolution of parasitoids provides crucial implications for biodiversity and its applications.

In parasitoid wasps, life-history trait variations are typically found in sequential timings of

host exploitation, from attacking, hatching to emerging (Alonzo and Kindsvater 2008). For

example, some species emerge immediately after oviposition (e.g., *Ichneumon*

promissorius; Carpenter 1995), while others emerge some time after (e.g., *Venturia*

canescens; Beckage & Riddiford 1978; Sequeira and Mackauer, 1992a; Harvey, 1994;

Harvey & Vet, 1996; Harvey, 2000; Castelo et al., 2010; Harvey et al., 2012). Such

differences in the exploitation timings determine the degree to which parasitoids acquire

resources from younger (and thus smaller) versus older (thus larger) hosts, which as a

consequence influences the population dynamics of younger and older hosts. In other

words, the evolution of exploitation timings strongly influences evolutionary and ecological

aspects of parasitoid lives.

The delayed emergence, resulting from parasitoids not consuming their hosts immediately

after the attack, is a key factor in creating diversity in the life-history characteristics of

parasitoids, including larval mortality, fecundity, and egg size (Jervis and Ferns 2011).

Parasitoid wasps can be categorized into two groups according to this trait; namely,

"idiophytic" species ('idio' means 'own/individual/personal' descending from Greek,

indicating that hosts are personalized or monopolized by a single parasitoid individual) that

consume the host immediately after the attack, and "koinophytic" species ('koino' means

'common', also from Greek, indicating parasitoids live with a living host) that allow hosts

to continue to feed and grow after parasitization (Haeselbarth 1978; Askew &Shaw 1986).

By applying this criterion for a wider range of parasitoid wasp species, the major

dichotomy is proposed in the larval developmental pattern of parasitoid wasps: idiobiont-

koinobiont extremes (Askew&Shaw 1986). The delayed emergence can therefore

characterize two major categories of parasitoids (Askew&Shaw 1986; Godfray 1994) with

different life-history characteristics. The dichotomy is often used as a basis to understand

the life-history traits in parasitoid wasps (Godfray, 1994; Jervis, 2008; Quicke, 2014),

although in many koinobiont species, they exhibit delayed emergence with various

durations depending on the host stage (Harvey et al., 1994; Harvey et al., 2000; Harvey &

Strand, 2002; Harvey et al., 2016).

Studies testing the dichotomous hypothesis examine the relationship between delayed

emergence trait and other traits. Statistical analyses that control for phylogenetic

correlations also provide some support for the dichotomy hypothesis, suggesting possible

mechanisms for the observed pattern of association between life-history traits (Mayhew and

Blackburn 1999). For example, koinobionts exhibit higher larval mortality, greater

fecundity, and smaller egg sizes compared to idiobionts (Mayhew and Blackburn 1999).

Mayhew and Blackburn (1999) suggest that these trait associations are explained by both

selective pressures and developmental constraints: the higher mortality rate may result from

the increased exposure to predators or immaturity of the host. This in turn favours higher

fecundity but at the cost of smaller egg sizes (Mayhew and Blackburn 1999). Furthermore,

idiobionts require a substantial reproductive investment for egg production because females

must package all the nutrients – primarily proteins – necessary for the completion of

embryogenesis into these eggs (Jervis & Kidd, 1986). Consequently, protein-rich

"anhydropic" eggs have evolved. Due to the large size of these eggs, most idiobiont females

cannot carry many eggs at one time and the fecundity tends to be low (Jervis & Kidd,

1986). In contrast, "hydropic" eggs that contain little protein have evolved in koinobionts,

likely resulting from the evolution of an extra-embryonic membrane that enables the

embryo to uptake proteins and other nutrients directly from the host haemolymph (Jervis &

Kidd, 1986). This adaptation allows koinobiont females to produce many small eggs and

the handling time for oviposition is often shorter than idiobiont's (Jervis & Kidd, 1986).

Therefore, various life-history traits are subject to the different selective pressures

depending on the host's developmental stage.

Previous studies have identified the adaptive significance of parasitizing early-stage hosts

with delayed emergence (Slansky, 1986). First, young hosts are more abundant (and thus

more available) than older hosts and thus are easier for wasps to locate (Price 1972).

Second, young hosts are more susceptible to parasitism due to their immature defense

mechanisms, such as thinner cuticles and weaker immune responses (Beckage and

Riddiford 1978; Vinson and Iwantsch 1980b). Third, attacking young hosts can repel

subsequent, heterospecific competitors attempting to attack the same hosts (e.g., in

Parasetigena silvestris and *Blepharipa pratensis* parasitizing; Godwin and Odell 1984).

Furthermore, delayed emergence may potentially increase their fecundity of the parasitoid,

because it allows parasitoids to get more resource from their hosts (Harvey et al. 1994,

2004). On the other hand, delayed emergence is subject to a cost of increased mortality

during the period for the host to grow (Price 1972; c.f., the "Slow-Growth High-Mortality

hypothesis"; Clancy and Price, 1987; Benrey and Denno 1997). Godfray (1994)

hypothesizes that the divergent selective pressures imposed by parasitizing early and late-

stage hosts lead to the polarization implied by the idiobiont-koinobiont dichotomy.

Consistent evidence is found in species, e.g., *Hyposoter horticola*, with a minute ovipositor

to lay eggs on the host (*Melitaea cinxia*) while pausing host development for approximately

one year until the host becomes the seventh instar (Castelo et al., 2010). Thus, the adaptive

significance of delayed emergence is qualitatively understood.

On the other hand, it remains unclear which factors such as host mortality, growth rates,

and fecundity with host stage determine the directional selection for koinobiont versus

idiobiont. Price (1972) constructed a framework in which the offspring number (F) is

determined by the product of the number of offspring per female (N) and the survival

probability of parasitoid larvae on host (P); namely, $F = N \cdot P$. Price's (1972) formula is a

general but verbal model in the sense that ecological or physiological determinants of N and

P are left unspecified, which limits the quantitative predictions of the direction of natural

selection. Hackett-Jones et al. (2011) formulated the adaptive significance of delayed

emergence. However, their model does not explicitly incorporate the stage-dependent traits

of the host such as density ratio and parasitoid fecundities, the mortalities. These factors are

critical for understanding the koinobiont strategy, since early-stage hosts often lack

sufficient resources to support immediate emergence. Therefore, the key driving factor of

evolution of koinobiont or idiobiont is yet to be better understood.

In this study, we use mathematical models to explore the adaptive evolution of delayed

emergence. We focus on explaining the evolutionary dynamics of the dichotomy, assessing

what conditions favor koinobiont versus idiobiont. To simplify the conceptualization of the

results, we take operational definitions of idiobionts which attack late-stage hosts and

consume immediately. We consider the evolutionary dynamics of two traits: (i) the

parasitoids' preference for hosts' developmental stages, and (ii) the degree of delayed

emergence for those parasitoids attacking young hosts. Using adaptive dynamics theory

(Dieckmann and Law 1996), we determine the directions of natural selection on these traits

and the stability of evolutionary consequences (i.e., the attainable strategies that no other

rare mutant with small phenotypic differences can increase in frequency). As the host

population is physiologically structured, we use Fisher's reproductive values to interpret the

optimal timing of emergence for parasitoids (Fisher 1930).

Model

Fig. 1 Diagram of host and parasitoids. Symbols: E for the density of early-stage hosts, L for the density of late-stage hosts, A for the density of adult hosts, and K for the density of parasitoids. We use subscript 0 for unparasitized and 1 for parasitized state, respectively. Arrows with black points represent death

We first construct the dynamical equations of stage-structured host-parasitoid populations. We incorporate delayed emergence using a stage-structured model consisting of early-stage hosts, late-stage hosts, and adult hosts. Either stage of the stage hosts, but not adults, may be attacked by the parasitoids and subsequently transits to the parasitized stage (Fig 1 for a

diagram depicting the patterns of life cycles). The variables and parameters are summarized

in Table 1.

We consider two independent traits subject to natural selection. One is preference for early-

stage hosts, x, and the other is probability of delayed emergence, y, with $0 \leq x \leq 1$ and

$0 \leq y \leq 1$. For example, the case with $x = 1$ (attacking early-stage hosts only) and $y = 0$

(emerging only from early-stage hosts) corresponds to the life cycle graph in Fig 2A, which

thus omits the lifecycle pathway between early-stage and late-stage hosts. We do not

consider the evolution of host traits.

We use ordinary differential equations to describe the population dynamics of hosts and

parasitoids:

185
$$
\begin{aligned}
\dot{E}_0 &= rA - a_0 E_0 - \lambda x E_0 K - d_E E_0 \\
\dot{L}_0 &= a_0 E_0 - \lambda(1 - x)L_0 K - b_0 L_0 - d_L L_0 \\
\dot{A} &= b_0 L_0 - d_A A \\
\dot{E}_1 &= \lambda x E_0 K - y a_1 E_1 - ((1 - y)\sigma_E + d_E + v_E)E_1 \\
\dot{L}_1 &= y a_1 E_1 + \lambda(1 - x)L_0 K - (\sigma_L + d_L + v_L)L_1 \\
\dot{K} &= f_E(1 - y)\sigma_E E_1 + f_L \sigma_L L_1 - d_K K
\end{aligned}
\tag{1}
$$

(with dot representing the time derivative). Here, from top to bottom, the ordinary

differential equations represent the dynamics of non-parasitized early-stage host (E_0), non-

parasitized late-stage hosts (L_0), adult hosts (A), parasitized early-stage host (E_1),

parasitized late-stage host (L_1), and adult parasitoids (emerging from either early or late

hosts; K). The key ingredients are the transitions in states due to (i) parasitism (e.g., from

E_0 to E_1), and (ii) ontogeny (e.g., from E to L) (Fig 1). Surviving late-stage hosts may

develop into adults (with density A).

The first three lines in Equation (1) describe the population dynamics of non-parasitized

hosts. With non-parasitized early-stage hosts parasitized at rate $\lambda x K$ with searching

efficiency of parasitoids λ and the probability of preferentially parasitizing early-stage

hosts x, the non-parasitized early-stage hosts (with density E_0) are produced by adults at

rate r, develop into late-stage hosts at rate a_0 and die at rate d_E; they are parasitized by

wild-type parasitoids at rate λx. Non-parasitized late-stage hosts (with density L_0) develop

from non-parasitized early-stage hosts but may be parasitized at rate $\lambda(1 - x)K$, develop

into adults at rate b_0, or die at rate d_L.

The middle two lines in Equation (1), on the other hand, describe the population dynamics

of parasitized hosts. Parasitized early-stage hosts (with density E_1), after transiting from the

parasitism of early-stage hosts (E_0), develop into late-stage hosts, or otherwise die due

either to (i) parasitoid emergence, at rate $(1 - y)\sigma_L$ (ii) natural mortality at rate d_E, or (iii)

virulence at rate v_E (increased mortality due to parasitism). Parasitized late-stage hosts

(with density L_1) develop from parasitized early-stage hosts due to ontogeny or transit from

non-parasitized late-stage hosts due to parasitism but die at rate σ_L due to emergence of the

parasitoid, or at rate $d_E + \gamma_E$ due to natural mortality or virulence. We assume that the

parasitized hosts are incapable of producing offspring.

The last line in Equation (1) describes the population dynamics of adult parasitoids. Adult

parasitoids emerge from either parasitized early-stage hosts (at rate f_E) or late-stage hosts

(at rate f_L), where f_E or f_L represents the fecundity of adult parasitoids (per capita)

emerging from early larvae or late larvae, respectively.

Table 1 Definitions of the parameters

Parameter	Definition
x	Preference for unparasitized early-stage hosts in parasitoids (evolving trait)
y	Preference for delayed emergence in parasitoids (evolving trait)
λ	Searching efficiency of parasitoids
r	Birth rate of hosts
a_0	Growth rate of unparasitized early-stage hosts
b_0	Growth rate of unparasitized late-stage hosts
a_1	Growth rate of parasitized early-stage hosts
d_E	Mortality rate of unparasitized early-stage hosts
d_L	Mortality rate of unparasitized late-stage hosts
d_A	Mortality rate of adult hosts
d_K	Mortality rate of parasitoids
v_E	virulence against early-stage hosts
v_L	virulence against late-stage hosts
σ_E	Consumption rate of early-stage hosts by parasitoids
σ_L	Consumption rate of late-stage hosts by parasitoids

Parameter	Definition
f_E	Per capita fecundity of parasitoids emerging from early-stage hosts
f_L	Per capita fecundity of parasitoids emerging from late-stage hosts

To investigate the long-term evolutionary dynamics of the preference for host stages in

parasitoids and the preference for delayed emergence in parasitoids, we apply the adaptive

dynamics theory (Dieckmann and Law 1996). We first suppose that the resident population

dynamics, Equation (1), have reached an endemic equilibrium, and introduce a rare mutant

of the parasitoid attempting to invade the resident equilibrium. The resident population has

a unique equilibrium (Appendix D). To determine whether the mutant parasitoid can invade

the resident population, we derive the invasion fitness proxy R_m by using the next-

generation theorem (Van den Driessche and Watmough 2002; Hurford et al. 2010):

$$R_\mathrm{m}(x_\mathrm{m}, y_\mathrm{m}; x, y) = \underbrace{\frac{\lambda x_\mathrm{m} E_0^*}{d_K} \cdot \frac{f_E(1 - y_\mathrm{m})\sigma_E}{y_\mathrm{m} a_1 + d_E + v_E + (1 - y_\mathrm{m})\sigma_E}}_{\text{path}(A)}$$

224
$$+ \underbrace{\frac{\lambda x_\mathrm{m} E_0^*}{d_K} \cdot \frac{y_\mathrm{m} a_1}{y_\mathrm{m} a_1 + d_E + v_E + (1 - y_\mathrm{m})\sigma_E} \cdot \frac{f_L \sigma_L}{d_L + v_L + \sigma_L}}_{\text{path }(B)} + \underbrace{\frac{\lambda(1 - x_\mathrm{m})L_0^*}{d_K} \cdot \frac{f_L \sigma_L}{d_L + v_L + \sigma_L}}_{\text{path }(C)}$$

$$\tag{2}$$

where E_0^* and L_0^* represent the equilibrium density of unparasitized early-stage host and

unparasitized late-stage host, respectively. In Equation (2), we partition the reproductive

success of the mutant individual into multiple pathways (Fig 2). The first, path (A) term

represents the reproductive success by attacking and consuming early-stage hosts (Fig 2A).

The path (B) term represents the reproductive success by attacking on early-stage hosts and

consuming late-stage hosts (Fig 2B). The path (C) term represents the reproductive success

by attacking and consuming late-stage hosts (Fig 2C).

The rare mutant can invade the resident population when $R_m > 1$. By using this criterion,

we examine a candidate evolutionary consequence, its evolutionary stability (i.e., whether

the rare mutant can invade a population dominated by the wild type; Maynard Smith 1982),

and also convergence stability (i.e., evolutionary dynamics convergence of the traits; Eshel

et al. 1983).

237

Fig. 2 Three pathways of reproductive success: (A) attacking the early-stage hosts and

immediately preying on them; (B) attacking the early-stage hosts and waiting until they

grow to late-stage hosts, and then preying on them; (C) attacking the late-stage host and

immediately preying on them

Although the results in the main text all assume the exponential growth of the parasitoid-

free host population (as this assumption immensely simplify the analyses), we also analyze

the case where host density follows logistic growth (i.e., density-dependent regulation of

host fecundity). To check the robustness of the results in the main text, we conduct

numerical, deterministic simulations (Appendix E).

Results

Evolution of delayed emergence

251   First, we present one of the key results that the delayed emergence trait y is polarized; i.e.,

252   natural selection pushes the population towards either $y = 0$ (immediate emergence) or

253    $y = 1$ (complete delayed emergence). We assess the directional selection on the

254   preference for delayed emergence y, by using the selection gradient (Appendix B) given

255   by:

256
$$\left.\frac{\partial R_\mathrm{m}\ (x_\mathrm{m}, y_\mathrm{m}; x, y)}{\partial y_\mathrm{m}}\right|_{x_\mathrm{m}=x, y_\mathrm{m}=y} = (\alpha F_L - F_E) \times (\text{positive term}) \qquad (3)$$

which is positive (or negative) when natural selection favours delayed (or early) emergence

(respectively), depending on the sign of the first term, $(\alpha F_L - F_E)$. In Equation (3), $\alpha =$

$a_1/(a_1 + d_E + v_E)$ is the probability of successful development from early to late-stage for

parasitized host, $F_L = f_L \sigma_L/(d_L + v_L + \sigma_L)$ or $F_E = f_E \sigma_E/(d_E + v_E + \sigma_E)$ is the product

of: (1) the probability of consumption of the host and (2) the fecundity (i.e., the expected

conversion rate of parasitoids preying on early or late-stage hosts, respectively). Although

the selection gradient per se depends on both of the evolutionary traits x and y, the sign is

determined only by non-evolving parameters such as α and F_L. Thus, the preference

evolves monotonically to either 0 or 1 (i.e. immediate or delayed emergence) depending on

the sign of $\alpha F_L - F_E$. We also find that the evolutionary outcome is always evolutionarily

and convergence stable (Appendix C), and therefore Equation (3) predicts that the long-

term evolutionary outcome is either immediate emergence or completely delayed

emergence (Otto and Day 2007).

We can interpret the monotonicity result of Equation (3) using the Fisher's (1930)

reproductive values. Equation (3) represents the relationship between the reproductive

value of early-stage hosts when the parasitoids consume immediately versus when they

wait for the host to develop into late-stage hosts before feeding. Here, the parameters $\alpha F_L -$

F_E quantify the balance between reproductive gains of parasitoids from these two

strategies: αF_L represents the expected number of adult parasitoids when the host survives

the initial attack, grows to the late-stage stage, and is consumed at that stage (pathway B in

Fig 2), whereas F_E measures the number of adult parasitoids that can be produced when an

early-stage host is consumed immediately (pathway A in Fig 2). The preference for delayed

emergence is likely to be favoured when the product of (1) the growth probability of the

parasitized early-stage hosts and (2) the average conversion rate when parasitoids consume

the late-stage host, is higher than (3) the conversion rate of the early-stage host. These

conditions mean that delayed emergence is favoured when late-stage hosts are overall more

valuable as resource for parasitoids than early-stage hosts. Taken together, our models

predict that the delayed emergence trait is polarized, leading to a clear division of

parasitoids into koinobionts and idiobionts depending on these conditions.

Fig. 3 The evolutionary consequences of the preference for delayed emergence ($\alpha =$

$a_1/(a_1 + d_E + v_E)$: the growth probability of the parasitized early-stage host, $F_E =$

$f_E\sigma_E/(d_E + v_E + \sigma_E)$: the average fecundity rate of parasitoids preying on the early-stage

host and $F_L = f_L\sigma_L/(d_L + v_L + \sigma_L)$: that on the late-stage host)

Patterns of life cycles

Next we examine the evolution of the preference to attack early-stage hosts (x). The

selection gradient of the preference for host stages x is given by:

$$\left.\frac{\partial R_{\mathrm{m}}\left(x_{\mathrm{m}}, y_{\mathrm{m}}; x, y\right)}{\partial x_{\mathrm{m}}}\right|_{x_{\mathrm{m}}=x, y_{\mathrm{m}}=y} = \frac{\lambda E_0^*}{d_K}\left((p_{E\to L}F_L + p_{E\to K}f_E) - \frac{L_0^*}{E_0^*}(x)F_L\right) \quad (4)$$

(Appendix B), where the compound parameter $p_{E\to L} = ya_1/(ya_1 + d_E + v_E + (1 -$

$y)\sigma_E)$ represents the survival probability (transition) from parasitized early-stage hosts to

late ones and $p_{E\to K} = (1 - y)\sigma_E/(ya_1 + d_E + v_E + (1 - y)\sigma_E)$ represents the probability

300   of parasitoid emergence among dead hosts The sign of Equation (4) depends on both x and

301   y; note that according to the previous section, y evolves to either 0 or 1 regardless of the

302   evolution of x .

303   Now we substitute $y = 0$ or 1 considering the result in the previous section. The selection

304   gradients on the preference for early-stage hosts for both cases are given by:

305
$$\left. \frac{\partial R_m\ (x_m, y_m; x, y)}{\partial x_m} \right|_{x_m=x, y_m=y=0} \propto F_E - \frac{L_0^*}{E_0^*}(x)F_L, \quad (5)$$

306
$$\left. \frac{\partial R_m\ (x_m, y_m; x, y)}{\partial x_m} \right|_{x_m=x, y_m=y=1} \propto \left(\alpha - \frac{L_0^*}{E_0^*}(x) \right) F_L \quad (6)$$

where $F_E = f_E \sigma_E / (d_E + v_E + \sigma_E)$ is the product of (1) the probability of consumption of

early-stage hosts and (2) the fecundity of them (i.e., the expected fecundity of parasitoids

preying on early-stage hosts) and $F_L = f_L \sigma_L / (d_L + v_L + \sigma_L)$ is that on late-stage host, $\alpha =$

$a_1 / (a_1 + d_E + v_E)$ is the probability of successful development of parasitized early-stage

host to late-stage host. We also find that the evolutionary outcome is always evolutionarily

and convergence stable (Appendix C), and therefore Equation (5) and (6) predicts the long-

term evolutionary outcome (Appendix E). A unique candidate evolutionary consequence x,

obtained analytically from Equations (5) and (6), is defined as the singular point where the

selection gradient becomes zero. When the candidate evolutionary consequence falls

outside the interval between 0 and 1, we carry out case-analyses to locate the interior

singular point (Appendix B.3). The evolutionary consequence in each case is shown in Fig

4. This result suggests that delayed emergence trait can generate diverse life cycles of

parasitoids, including idiobiont (Fig 4 (D- v)) koinobiont (Fig 4 (D-ii) and (D-iv)) (see

below). Figs 4 (D-i) and (D-iii) exhibit strategies which immediate exploit early-stage

hosts. These patterns are biologically implausible since early-stage hosts provide

insufficient resources for parasitoid development.

Fig. 4 The patterns of delayed emergence resulting from our model. (A) Only early hosts

are attacked (the singular point is 0) when the growth probability of the parasitized early-

stage host α is larger than the maximum of the density ratio of the unparasitized late-stage

host to the early-stage hosts L_0^*/E_0^*. (B) Both early and late hosts are attacked (the singular

point is between 0 and 1) when the growth probability of the parasitized early-stage host α

is between the minimum and maximum of L_0^*/E_0^*). (C) Only late hosts are attacked (the

singular point is 1) when α is smaller the minimum of L_0^*/E_0^*). (D) The five panels show

distinct evolutionary outcomes in parasitoid life cycles resulting from variation in

emergence timing and host stage exploitation, referred to as (ⅰ)-(ⅴ) in the graphs. *E* or *L*

represents early-stage hosts or late-stage hosts (respectively). (ⅰ) only early hosts are

attacked and they are consumed immediately. (ⅱ) only early hosts are attacked and they

are not consumed immediately. (ⅲ) both early and late hosts are attacked and they are

consumed immediately. (ⅳ) both early and late hosts are attacked and late ones are

consumed. (ⅴ) late hosts are attacked and they are consumed immediately

Evolution of idiobiont and koinobiont

Following Godfray (1994), we define idiobiont as the parasitoids attacking unparasitized

late-stage hosts and prey on the host immediately. Similarly, we define koinobiont as the

parasitoids that attack an unparasitized early-stage host and prey on the host after the host

grows up to the late-stage stage (Godfray, 1994). Therefore, Fig 4 (D-ii) represents

koinobiont and Fig 4 (D-v) represents idiobiont. The evolution of koiobionts is more likely

when the survival probability of parasitized early-stage hosts to late ones is high, and the

relative density of late-stage hosts to early-stage hosts is small. Fig (D-iv) provides a clear

illustration of the strategy depending on host stage at parasitism in koinobionts. While

koinobionts can attack hosts at various developmental stages, they differentiate their

approach—either allowing further host growth or initiating immediate consumption—

depending on the specific stage encountered. The evolution of idiobionts is more likely

when the survival probability of parasitized early-stage hosts to late ones is low, and the

relative density of late-stage hosts to early-stage hosts is large.

Finally, we check the robustness of the results based on the exponential growth of hosts in

the absence of parasitoid. Numerical simulations (assuming the host population exhibits

logistic growth due to density-dependent regulation) confirmed the robustness of our results

against the logistic growth case (Appendix E).

Discussion

In this study, we investigated the effect of stage-structure and host ecology on the evolution

of life-history traits in parasitoids. Our primary achievement is the formulation of the

selection pressure on the delayed emergence using the concept of reproductive value

(Fisher 1930). Specifically, if the host's developmental growth does not increase the

expected conversion efficiency (a measure of nutritional quality), where $F_L \leq F_E$, the

parasitoids are then selected to emerge from the host immediately (Fig 3). Otherwise, either

early or late emergence is selected for (Fig 3). The specific outcome of this selection is

determined by comparing the reproductive value of the host for parasitoids emerging early

(F_E) with that for those waiting for host growth (αF_L) (Fig 3). Moreover, despite its

fundamental structure, our model can predict diverse evolutionary outcomes of parasitoid

life cycles (Fig 4). To incorporate delayed emergence into the model, we considered two

developmental stages of hosts available for parasitism, allowing parasitized early-stage

hosts to continue developing into later stages. As a result, the host utilization strategies of

parasitoids exhibit five distinct patterns (Fig 4D). This result indicates that the evolution of

emergence timing leads to the observed diversity of parasitoid life cycles, including both

idiobiont- and koinobiont- strategies. Notably, our model can account for the strategy

observed in many koinobionts, which utilize both early- and late-stage hosts (Fig 4 (D-iv)).

When parasitoids attack early-stage hosts, the model predicts that they delay emergence to

maximize resource acquisition from host growth. Conversely, when parasitoids attack late-

stage hosts, our model predicts immediate consumption. This result aligns with the patterns

in which the duration of delayed emergence is shortened when the host already has high

resource value. This strategy involves a dynamic preference between host stages, driven by

their relative abundance (density ratio). Our model explores how the trade-offs between

host stages shift depending on the density ratio. This finding provides a theoretical

explanation for why koinobionts do not always delay their emergence but instead finetune

the timing based on the host's value at the time of parasitism.

Our predictions are overall consistent with a previous theoretical study on stage-structured

host populations. Iritani et al. (2019) considered stage-structured disease dynamics and

showed that the evolutionary consequences of virulence toward larval hosts depend on the

interaction between host maturation rate and the infection pathway. Specifically, virulence

toward larval hosts evolves to be higher when either (i) the adult stage is relatively long and

the transmission across host stages mostly occurs, or (ii) the larval stage is relatively long

and the transmission across the same stage mostly occurs (Iritani et al. 2019). These

findings align with our results: koinobiont strategies evolve when early-stage hosts are

abundant and maturation under infection is likely, whereas the idiobiont strategy evolves

when late-stage hosts are abundant and maturation is unlikely. Hackett-Jones et al. (2011)

investigated the evolution of delayed emergence in parasitoids using a discrete-time model,

showing that delayed emergence is under balancing selection between two forces. First,

delayed emergence allows parasitoids to attain a larger adult body size, which is favored by

the increased searching efficiency associated with larger size. Second, delayed emergence

is costly because of the host mortality during the period of delayed emergence. In this

context, our models are a simpler generalization of their model for two reasons. Compared

to Hackett-Jones et al.'s (2011) model assuming the host population is homogeneous (thus

with no structure), we explicitly consider host stage-structure. Also, the authors' analytical

formula involves host density in a complex mathematical form, making biological

interpretation difficult. Even more crucially, their model does not account for two key

biological facts: (1) young hosts are more abundant than older hosts (Price, 1972), and (2)

young hosts are easier to parasitize due to weaker resistance, such as lower physical

defense or immunity (Beckage and Riddiford 1978; Vinson and Iwantsch 1980b). Our

model incorporates these essential factors to derive a biologically interpretable formula of

selection pressure for delayed emergence, revealing the optimal strategy is determined by

the relative reproductive value of host stages depending on the balance between host

growth, mortality, and nutritional quality.

Our model predicts that host density (density ratio) has strong impact on the evolution of

delayed emergence. In environments where early-stage hosts are relatively abundant

compared to late-stage hosts, natural selection favours delayed emergence which exploits

the high encounter rate of early-stage hosts and waits for growth (Slansky 1986). In nature,

early-stage hosts are more numerous than late-stage hosts. Therefore, even if there is a

mortality cost while waiting for growth within the host, the benefits provided by the high

encounter rate can outweigh this cost. Conversely, idiobionts tend to evolve in

environments where the natural mortality of the host is low and late-stage hosts are nearly

as abundant as early-stage hosts. This allows them to efficiently avoid the cumulative

mortality associated with the period required for early-stage hosts to mature (Price 1972).

Host density best illuminates ecological characteristics of the host (host ecology). We

discuss this point focusing on host natural mortality caused by scavengers and other

predators. Idiobionts tend to attack concealed hosts, such as those within plant tissues or

cocoons (Price 1972; Gauld, 1988; Quicke, 2014). This preference arises because

permanent paralysis by idiobiont renders the host entirely defenseless. In an exposed

environment, such an incapacitated host and the parasitoid larva feeding on it would be

highly vulnerable to natural mortality from scavengers and predators (Gauld, 1988).

Idiobionts target hosts in concealed niches where this natural mortality v_L is sufficiently

small (Gauld, 1988). In addition, late-stage hosts are likely to be relatively abundant in

these protected environments because natural mortality due to predation throughout the

developmental period d_E is relatively low (Price, 1972). These aspects of host ecology

support the association between immediate emergence and the parasitization of late-stage

hosts, under the selection pressures on parasitoids defined by Equation (3) for delayed

emergence and Equation (4) for host stage, respectively. In contrast, koinobionts often

attack exposed hosts that feed on surfaces such as leaves (Price 1972; Gauld, 1988; Quicke,

2014). Paralyzing hosts in such a location would lead to an increase in natural mortality, as

the host and the parasitoid larva feeding on it would be easily discovered and preyed upon

(Gauld, 1988). Therefore, maintaining host mobility is an essential benefit to ensure the

parasitoid's survival (Gauld, 1988). Koinobionts maintain host mobility so that this natural

mortality v_E remains sufficiently small even in exposed niches (Gauld, 1988). Furthermore,

in these exposed environments, late-stage hosts are rarely abundant because high natural

mortality d_E throughout development limits the number of individuals reaching maturity

(Price 1972). These aspects of host ecology support the association between delayed

emergence and the parasitization of early-stage hosts in koinobionts, under the selection

pressures defined by Equation (3) for delayed emergence and Equation (4) for host stage,

respectively. These formulations explain how host ecology, specifically stage-specific

mortality, leads to the polarization between idiobiont and koinobiont.

The present results predict that the increase in fecundity as the host matures favours

delayed emergence. In fact, the positive correlation between host size and fitness-related

traits such as adult size, egg count, and longevity is particularly pronounced, because they

halt host growth (Salt 1941, Arthur and Wylie 1959, Vinson 1972, Sandlan 1979, Strand et

al. 1988, King 1989). However, in koinobionts, a larger host size does not necessarily

translate into a positive impact on fitness-related traits (Harvey, 2016). The host is not

merely food; the parasitoid must highly synchronize its growth with the host's

physiological and developmental stage (Harvey, 2016). For example, *Apanteles carpatus*

attacks hosts (*Tineola bisselliella*) which are as small as 3% (30–40 mg) of its own body

weight (1–1.5 mg). While it takes about 30 days for the parasitoid to develop from a fifth-

instar host to adult stage, it takes up to 87 days for the parasitoid to develop from a first–

third-instar host to adult stage (Harvey, 2000). On the other hand, there are negative factors

such as the failure to pupate when they cannot fully consume excessively large hosts

(Harvey 1996) and the host's strong immune response in older hosts (Harvey et al. 2004).

That is, host's value to koinobionts must be understood not as being directly proportional to

host size at the time of oviposition (Sequeira and Mackauer 1992a,b; Harvey 2012).

Therefore, the idiobiont-koinobiont dichotomy should be considered as a continuum

because koinobionts can flexibly adjust the degree of delayed emergence depending on the

host's developmental stage at the time of parasitism. Thus, future theoretical studies would

benefit from taking these processes into account to explain the life-history variations in

parasitoid wasps.

We made some simplifying assumptions. For instance, the possibility of multiple

parasitisms is omitted from the model, despite its occurrence in nature (Price, 1972; Harvey

et al., 2013; Ode et al., 2022). Multiple parasitisms are associated with competition, which

drives temporal niche differentiation in host developmental stages (Price, 1972). This niche

differentiation promotes the evolution of delayed emergence (Price, 1972). Future

theoretical studies may incorporate the effects of such competition, which may provide

testable predictions for the evolution of life-history variations in parasitoids. Moreover,

hosts can also evolve defense traits such as resistance, potentially leading to coevolution.

Age structure of hosts may further shape these dynamics, as shown in a model of tolerance–

virulence coevolution (Buckingham and Ashby 2024). Future extensions of the present

model may incorporate host evolution to develop a useful theoretical framework for

understanding how host–parasitoid coevolution drives variation in emergence timing and

associated life-history traits .

In summary, this study rigorously formulated the selection pressure on delayed emergence

in parasitoids using the concept of reproductive value, providing a theoretical foundation to

make testable predictions for existing empirical findings. The primary contribution of this

model lies in providing measurable quantitative indices for the idiobiont-koinobiont

dichotomy. Future studies may extend our model to incorporate the adjustment of delayed

emergence depending on the host's developmental stage in koinobionts to understand the

multi-layered dichotomy.

Appendix

A. Next generation matrix and invasion fitness proxy

493 To study life-history evolution, we assess the invasion possibility of a rare mutant into the

494 resident population assuming that the ecological dynamics has reached an equilibrium. The

495 population dynamics of the rare mutant is given by:

$$\frac{d\mathbf{X}_m}{dt} = (\mathbf{F}_m - \mathbf{V}_m)\mathbf{X}_m \quad \text{(A-1)}$$

497 where

$$\mathbf{X}_m = \begin{pmatrix} E_{1,m} \\ L_{1,m} \\ K_m \end{pmatrix}$$

$$\mathbf{F}_m - \mathbf{V}_m = \begin{pmatrix} -(y_m a_1 + d_E + v_E + (1-y_m)\sigma_E) & 0 & \lambda x_m E_0^* \\ y a_1 & -(d_L + v_L + \sigma_L) & \lambda(1-x_m)L_0^* \\ (1-y_m)f_E\sigma_E & f_L\sigma_L & -d_K \end{pmatrix}$$

$$\mathbf{F}_m = \begin{pmatrix} 0 & 0 & \lambda x_m E_0^* \\ 0 & 0 & \lambda(1-x_m)L_0^* \\ (1-y_m)f_E\sigma_E & f_L\sigma_L & 0 \end{pmatrix}$$

$$\mathbf{V}_m = \begin{pmatrix} y_m a_1 + d_E + v_E + (1-y_m)\sigma_E & 0 & 0 \\ -y_m a_1 & d_L + v_L + \sigma_L & 0 \\ 0 & 0 & d_K \end{pmatrix}$$

(A-

499　2)

500　The Next-Generation Theorem (Van den Driessche and Watmough 2002) states that if the

501　dominant eigenvalue of the next-generation matrix $\mathbf{G}_m = \mathbf{F}_m \cdot \mathbf{V}_m^{-1}$ exceeds 1, then the

502　mutant can invade and drive the resident individuals extinct. The squared, dominant

503　eigenvalue $\rho(\mathbf{G}_m)$ is given by:

$$\begin{aligned}
(\rho(\mathbf{G}_m))^2 &= \frac{\lambda x_m E_0^*}{d_K} \cdot \frac{y_m a_1}{y_m a_1 + d_E + v_E + (1-y_m)\sigma_E} F_L \\
&+ \frac{\lambda x_m E_0^*}{d_K} \cdot \frac{f_E(1-y_m)\sigma_E}{y_m a_1 + d_E + v_E + (1-y_m)\sigma_E} \\
&+ \frac{\lambda(1-x_m)L_0^*}{d_K} F_L
\end{aligned}$$

(A-3)

505　where

$$\mathbf{G_m} = \begin{pmatrix} 0 & 0 & \frac{\lambda x_m E_0^*}{d_K} \\ 0 & 0 & \frac{\lambda(1-x_m)L_0^*}{d_K} \\ p_{m,E\to L}F_L + p_{m,E\to K}f_E & F_L & 0 \end{pmatrix} \quad \text{(A-4)}$$

$$p_{m,E\to L} = \frac{y_m a_1}{y_m a_1 + d_E + v_E + (1-y_m)\sigma_E}$$

$$p_{m,E\to K} = \frac{(1-y_m)\sigma_E}{(y_m a_1 + d_E + v_E + (1-y_m)\sigma_E)}$$

where $p_{m,E\to L}$ represents the probability that mutant parasitoids' larvae can transit from

parasitized early-stage hosts to late ones and $p_{E\to K}$ represents the probability that mutant

parasitoids' larvae consume the host. Since $\rho^2 \gtrless 1$ is equivalent to $\rho \gtrless 1$, we use ρ^2 for

the invasion proxy.

B. Selection gradient

In this section, we derive the stable class distribution, reproductive values and selection

gradient using the next-generation matrix.

B-1. Stable class distribution

The right eigenvector associated with the dominant eigenvalue of the next-generation

matrix gives the stable class distribution (Taylor 1990; Otto and Day 2007), which

represents the asymptotic distribution of mutant alleles. It is given by:

$$\mathbf{G}\vec{u} = \rho(\mathbf{G})\vec{u} \qquad (A-5)$$

520 where

$$\vec{u} = \begin{pmatrix} \dfrac{\lambda x E_0^*}{d_K} \\ \dfrac{\lambda(1-x)L_0^*}{d_K} \\ \rho(\mathbf{G}) \end{pmatrix} \tag{A-6}$$

523 represents the stable distribution of the mutant allele.

B-2. Reproductive value of each class

525 The left eigenvector associated with the dominant eigenvalue of the next-generation matrix

526 gives the reproductive values of each of these classes (Fisher 1930; Otto and Day 2007).

527 The reproductive values represent the contributions of individuals in each class to future

528 reproduction after sufficiently long generations, given by:

$$\vec{v}^{\mathsf{T}}\mathbf{G} = \rho(\mathbf{G})\vec{v}^{\mathsf{T}} \tag{A-7}$$

531 where

$$\vec{v}^{\mathsf{T}} = \begin{pmatrix} p_{E \to L}F_L + p_{E \to K}f_E \\ F_L \\ \rho(\mathbf{G}) \end{pmatrix}$$

$$p_{E \to L} = \frac{ya_1}{ya_1 + d_E + v_E + (1-y)\sigma_E}$$

$$p_{E \to K} = \frac{(1-y)\sigma_E}{ya_1 + d_E + v_E + (1-y)\sigma_E}$$

532 (A-8)

533 $p_{E \to L}$ represents the probability that parasitoids' larvae can transit from parasitized early-

534 stage hosts to late ones and $p_{E \to K}$ represents the probability that mutant parasitoids' larvae

535 consume the host.

B-3. Selection gradient of x

537 From the calculations above, we have $\vec{u}\vec{v} = 2\rho^2(\mathbf{G}) = 2$. Let $c_i = u_i v_i$ be class

538 reproductive value of class i; the selection gradient of x is given by:

539
$$
\begin{aligned}
\frac{\partial}{\partial x_{\mathrm{m}}}\left(\frac{\vec{v}\mathbf{G}_{\mathrm{m}}\vec{u}}{\vec{v}\vec{u}}\right)\Bigg|_{x_{\mathrm{m}}=x} &= \frac{1}{2}c_1\left(\frac{\partial g_{11}}{\partial x_{\mathrm{m}}} + \frac{v_2}{v_1}\cdot\frac{\partial g_{21}}{\partial x_{\mathrm{m}}} + \frac{v_3}{v_1}\cdot\frac{\partial g_{31}}{\partial x_{\mathrm{m}}}\right) \\
&+ \frac{1}{2}c_2\left(\frac{v_1}{v_2}\cdot\frac{\partial g_{12}}{\partial x_{\mathrm{m}}} + \frac{\partial g_{22}}{\partial x_{\mathrm{m}}} + \frac{v_3}{v_2}\cdot\frac{\partial g_{32}}{\partial x_{\mathrm{m}}}\right) \\
&+ \frac{1}{2}c_3\left(\frac{v_1}{v_3}\cdot\frac{\partial g_{13}}{\partial x_{\mathrm{m}}} + \frac{v_2}{v_3}\cdot\frac{\partial g_{23}}{\partial x_{\mathrm{m}}} + \frac{\partial g_{33}}{\partial x_{\mathrm{m}}}\right) \\
&= \frac{1}{2}c_3\left(\frac{v_1}{v_3}\cdot\frac{\partial g_{13}}{\partial x_{\mathrm{m}}} + \frac{v_2}{v_3}\cdot\frac{\partial g_{23}}{\partial x_{\mathrm{m}}}\right)
\end{aligned}
\tag{A-9}
$$

541 where

$$\left.\frac{\partial \mathbf{G}_\mathrm{m}}{\partial x_\mathrm{m}}\right|_{x_\mathrm{m}=x} = \left.\begin{pmatrix} \dfrac{\partial g_{11}}{\partial x_\mathrm{m}} & \dfrac{\partial g_{12}}{\partial x_\mathrm{m}} & \dfrac{\partial g_{13}}{\partial x_\mathrm{m}} \\[2mm] \dfrac{\partial g_{21}}{\partial x_\mathrm{m}} & \dfrac{\partial g_{22}}{\partial x_\mathrm{m}} & \dfrac{\partial g_{23}}{\partial x_\mathrm{m}} \\[2mm] \dfrac{\partial g_{31}}{\partial x_\mathrm{m}} & \dfrac{\partial g_{32}}{\partial x_\mathrm{m}} & \dfrac{\partial g_{33}}{\partial x_\mathrm{m}} \end{pmatrix}\right|_{x_\mathrm{m}=x}$$

$$= \begin{pmatrix} 0 & 0 & \dfrac{\lambda E_0^*}{d_K} \\[2mm] 0 & 0 & -\dfrac{\lambda L_0^*}{d_K} \\[2mm] 0 & 0 & 0 \end{pmatrix} \tag{A-10}$$

$$c_3 = \rho^2(\mathbf{G}) = 1$$
$$v_1 = p_{E \to L} F_L + p_{E \to K} f_E$$
$$v_2 = F_L$$
$$v_3 = \rho(\mathbf{G}) = 1$$

543 ## B-4. Evolution of preference for host stages when immediate emergence

544 occurs.

545 The selection gradient on the preference for early-stage hosts is

$$546 \qquad \left.\frac{\partial R_\mathrm{m}(x_\mathrm{m}, y_\mathrm{m}; x, y)}{\partial x_\mathrm{m}}\right|_{x_\mathrm{m}=x, y_\mathrm{m}=y=0} \propto F_E - \frac{L_0^*}{E_0^*}(x)F_L. \tag{A-11}$$

where $F_E = f_E \sigma_E / (d_E + v_E + \sigma_E)$ is the expected fecundity of parasitoids preying on

early-stage host and $F_L = f_L \sigma_L / (d_L + v_L + \sigma_L)$ is that on late-stage host. A unique

candidate evolutionary consequence x^* can be obtained analytically. The point is always

convergence stable because L_0^*/E_0^* is monotonically increasing function of x^* (Appendix

D). Evolutionary consequences of the preference for host stages are always continuously

stable in the case that parasitoids consume hosts immediately after they attack the hosts.

According to result section, we obtain the evolutionary consequences in three cases (Fig 5).

Equation Equation (A-11) represents the relationship between the class reproductive value

of unparasitized early-stage hosts and that of late ones when the parasitoids prey

immediately. We can interpret Equation Equation (A-11) as the evolution of delayed

emergence such that the hosts' class reproductive value maximizes. When parasitoids

consume hosts immediately after they attack hosts, the preference for early-stage hosts is

likely to evolve if the average conversion rate and the density of late-stage hosts are higher

than that of early-stage hosts. In addition, the lower the growth probability of parasitized

early-stage hosts, the greater the diversity of host stages can evolve.

Fig. 5 Evolutionary consequences of the preference for host stages when parasitoids

emerge immediately. Immediately emerge ($y = 0$) is convergence stable in the white area,

but not in the gray area. Parasitoids attack only early-stage hosts in the "early" area, attack

both of early and late-stage hosts in the "both" area, and attack late-stage hosts in the "late"

area. (a) The singular point is 0 (the growth probability of the parasitized early-stage host α

is larger than the maximum of the density ratio of the unparasitized late-stage host to the

early one L_0^*/E_0^*). (b) The singular point is between 0 and 1 (α is between the minimum and

maximum of L_0^*/E_0^*). (c) The singular point is 1 (α is smaller the minimum of L_0^*/E_0^*)

B-5. Evolution of preference for host stages when delayed emergence occurs.

The selection gradient on the preference for early-stage hosts is

575
$$\left. \frac{\partial R_m(x_m, y_m; x, y)}{\partial x_m} \right|_{x_m=x, y_m=y=1} \propto \left(\alpha - \frac{L_0^*}{E_0^*}(x) \right) F_L. \qquad (A-12)$$

where $\alpha = a_1/(a_1 + d_E + v_E)$ is the growth probability of parasitized early-stage host,

$F_L = f_L \sigma_L/(d_L + v_L + \sigma_L)$ is the expected fecundity rate of parasitoids preying on late-

stage host. A unique candidate evolutionary consequence x^* can be obtained analytically.

The point is always convergence stable because L_0^*/E_0^* is monotonically increasing function

of x^* (Appendix D). Evolutionary consequences of the preference for host stages are

always continuously stable in the case that parasitoids do not consume hosts immediately

after they attack the hosts. According to result section, we obtain the evolutionary

consequences in three cases (Fig 6). Equation Equation (A-12) represents the relationship

between the class reproductive value of unparasitized early-stage hosts and that of late ones

when parasitoids wait for their maturation. We can interpret Equation Equation (A-12) as

the evolution of delayed emergence such that the hosts' class reproductive value

maximizes. Therefore, in the case that parasitoids do not consume hosts immediately after

they attack the hosts, the preference for the unparasitized early-stage host is likely to evolve

under two conditions: (1) the growth probability of the parasitized early-stage host is high,

and (2) the density of the late-stage host is lower than that of early-stage hosts.

Fig. 6: The evolutionary consequences of preference for host stages when delayed

emergence occurs. Delayed emergence ($y = 1$) is convergence stable in the white area, but

not in the gray area. Parasitoids attack only early-stage hosts in the "early" area, attack both

early and late-stage hosts in the "both" area, and attack late-stage hosts in the "late" area

B-6. Selection gradient of y

598 Selection gradient of y is given by:

$$\frac{1}{2}\vec{v}\frac{\partial \mathbf{G}_m}{\partial y_m}\Big|_{y_m=y}\vec{u} = \frac{1}{2}c_1\left(\frac{\partial g_{11}}{\partial y_m} + \frac{v_2}{v_1}\cdot\frac{\partial g_{21}}{\partial y_m} + \frac{v_3}{v_1}\cdot\frac{\partial g_{31}}{\partial y_m}\right)$$
$$+ \frac{1}{2}c_2\left(\frac{v_1}{v_2}\cdot\frac{\partial g_{12}}{\partial y_m} + \frac{\partial g_{22}}{\partial y_m} + \frac{v_3}{v_2}\cdot\frac{\partial g_{32}}{\partial y_m}\right)$$
$$+ \frac{1}{2}c_3\left(\frac{v_1}{v_3}\cdot\frac{\partial g_{13}}{\partial y_m} + \frac{v_2}{v_3}\cdot\frac{\partial g_{23}}{\partial y_m} + \frac{\partial g_{33}}{\partial y_m}\right)$$
$$= \frac{1}{2}c_1\frac{v_3}{v_1}\cdot\frac{\partial g_{31}}{\partial y_m} \qquad (A-13)$$

600   where

$$\frac{\partial \mathbf{G}_m}{\partial y_m}\Big|_{y_m=y} = \begin{pmatrix} \dfrac{\partial g_{11}}{\partial y_m} & \dfrac{\partial g_{12}}{\partial y_m} & \dfrac{\partial g_{13}}{\partial y_m} \\[2mm] \dfrac{\partial g_{21}}{\partial y_m} & \dfrac{\partial g_{22}}{\partial y_m} & \dfrac{\partial g_{23}}{\partial y_m} \\[2mm] \dfrac{\partial g_{31}}{\partial y_m} & \dfrac{\partial g_{32}}{\partial y_m} & \dfrac{\partial g_{33}}{\partial y_m} \end{pmatrix}\Big|_{y_m=y}$$

$$= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \dfrac{a_1(d_E + v_E + \sigma_E)}{(ya_1 + d_E + v_E + (1-y)\sigma_E)^2)}F_L - \dfrac{f_E\sigma_E(a_1 + d_E + v_E)}{(ya_1 + d_E + v_E + (1-y)\sigma_E)^2} & 0 & 0 \end{pmatrix}$$

$$c_1 = \frac{\lambda x E_0^*}{d_K}(p_{E\to L}F_L + p_{E\to K}f_E)$$
$$v_1 = p_{E\to L}F_L + p_{E\to K}f_E$$
$$v_3 = \rho(\mathbf{G}) = 1 \qquad (A-14)$$

C. Evolutionary stability

604   By examining the sign of the second derivative given that the candidate evolutionary

605   consequence is in the interior of the phenotypic space (which is the square phenotypic

606   space spanned by the two traits (x, y), we assess the evolutionary stability of x using the

607   adaptive dynamics theory. We have now written as

608   The second derivative with respect to x_m is given by:

609
$$\left. \frac{\partial^2 R_m(x_m, y_m; x, y)}{\partial x_m^2} \right|_{x_m = x = x^*, y_m = y} = 0. \qquad (A-15)$$

This follows from the equality $w(x_m, x_m) = w(x^*, x_m)$. In this case, by definition, for the

evolutionarily singular strategy (x^*) to be evolutionarily stable, the condition $w(x^*, x) >$

$w(x, x)$ must be satisfied.

For simplicity, let us express $R_m(x_m, y_m; x, y)$ as $R_m(x_m, x) = A(x)x_m + B(x)(1 -$
$x_m)$. We guarantee the evolutionary stability by proving that $R_m(x_m, x) > R_m(x, x)$.

$R_m(x^*, x) - R_m(x, x) = \{A(x)x^* + B(x)(1 - x^*)\} - \{A(x)x + B(x)(1 - x)\}$

$\qquad\qquad = \{A(x) - B(x)\}(x^* - x)$

Here, equations (A-6) and (A-7) imply convergence stability. Since the selection gradient is
written as $A(x) - B(x)$,

If $x^* < x$, $A(x) - B(x) < 0$

If $x^* > x$, $A(x) - B(x) > 0$

Therefore, the inequality $R_m(x^*, x) - R_m(x, x) > 0$ always holds.

On the other hand, we can examine the evolutionary stability of the preference for delayed

emergence y without taking the second derivative because the selection gradient of y is

independent of frequency. The selection gradient of y is given by:

625
$$\left. \frac{\partial R_m(x_m, y_m; x, y)}{\partial y_m} \right|_{x_m = x, y_m = y} \propto \alpha F_L - F_E. \qquad (A-16)$$

The selection pushes the trait to the boundary of the phenotypic space, causing evolution to

stop there. Thus, when the sign is negative, $y^* = 0$ is evolutionarily stable (i.e., the

strategies that no other rare mutant with small phenotypic differences can increase in

frequency) because it attains a local maximum. We can apply a similar argument to the case

where $y^* = 1$, and find that $y^* = 1$ is also evolutionarily stable when the sign is positive,

because it attains a local maximum.

D. An unique equilibrium of the population dynamics

633   We can obtain the density ratio of L_0^* to E_0^* as follows:

634
$$\frac{L_0^*}{E_0^*}(x) = \frac{-d_A(b_0 + d_L)x + d_A(a_0 + d_E)(1-x) + \sqrt{\left(d_A(b_0 + d_L)x + d_A(a_0 + d_E)(1-x)\right)^2 + 4rd_A a_0 b_0 x(1-x)}}{2rb_0(1-x)}$$

$$(A-19)$$

636   which is a monotone increasing function of x. Here, we will prove the uniqueness of the

637   equilibrium of the population dynamics (Equation (1)). From the third line, we obtain

638
$$A^* = \frac{b_0}{d_A} L_0^*$$

639   From the third line, we obtain K^* analytically

640
$$K^* = \frac{1}{\lambda x}\left(-(a_0 + d_E) + r\frac{L_0^*}{E_0^*}\right)$$

641 From the fourth line and the fifth line, we obtain

642
$$L_1^* = \frac{1}{\sigma_L + d_L + v_L} \left(\frac{\lambda x y a_1}{y a_1 + (1-y)\sigma_E + d_E + v_E} + \lambda(1-x) \right) K^* E_0^*$$

643 From the sixth line, we obtain

644
$$E_0^* = \frac{d_K}{\frac{\lambda x f_E (1-y)\sigma_E}{y a_1 + (1-y)\sigma_E + d_E + v_E} + \frac{f_L \sigma_L}{\sigma_L + d_L + v_L} \left(\frac{\lambda x y a_1}{y a_1 + (1-y)\sigma_E + d_E + v_E} + \lambda(1-x) \right)}$$

645 From this equation, we can obtain all the rest equilibrium L_0^*, A^*, E_1^*.

646
$$L_0^* = \frac{L_0^*}{E_0^*} E_0^*$$
$$A^* = \frac{b_0}{d_A} \frac{L_0^*}{E_0^*} E_0^*$$
$$E_1^* = \frac{\lambda x}{y a_1 + (1-y)\sigma_E + d_E + v_E} K^* E_0^*$$

E. Simulation

We conduct deterministic evolutionary simulations and then compare the results of

exponential growth with those of logistic growth. First, we divide the parasitoids

population into 25 strains based on the evolutionary traits and construct a mathematical

model that represents the population dynamics of these strains. Second, we numerically

solve the population dynamics to determine which strain survives over an extended period.

Finally, we compare the outcomes of evolutionary simulations of the case where the host

has logistic growth with that of the case where the host has exponential growth. We show

that our analytical results are robust against this modified assumption.

(a) Time variation of frequency of each strain   (c) Time variation of frequency of each strain

(b) Time variation of frequency of each strain   (d) Time variation of frequency of each strain

Fig. 7 Evolutionary simulation: (a) The host density follows linear growth. (b) The host

density follows logistic growth (carrying capacity $c = 100$). Both represent the case where

parasitoids evolve to idiobiont. Parasitoids attack late-stage hosts when $x = 0$, so the

preference for delayed emergence y is neutral ($\lambda = 1, r = 4, a_0 = 1.5, b_0 = 0.5, d_E = 1,$

$d_L = 1, d_A = 0.5, d_K = 0.5, v_E = 0.5, v_L = 0.5, \sigma_E = 0.5, \sigma_L = 1, f_E = 0.5, f_L = 0.6$).

(c)The host density follows linear growth. (d)The host density follows logistic growth (The

carrying capacity $c=100$). Both of Fig7 (c) and Fig7 (d) represent the case that parasitoids

evolve to koinobiont ($\lambda = 0.1, r = 0.5, a_0 = 0.16, b_0 = 0.2, d_E = 0.1, d_L = 0.1, d_A =$

$0.1, d_K = 0.1, v_E = 0.1, v_L = 0.1, \sigma_E = 1, \sigma_L = 1, f_E = 1, f_L = 2$)

Declarations

Funding

This work was supported by JSPS KAKENHI (grant numbers 20K15882, 24H02291, and

24H01528 to RIri); also by Mathematics-Based Creation of Science Program (MACS

program; Kyoto University).

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

672   References

Alonzo SH, Kindsvater HK (2008) Life-history patterns. In: Encyclopedia of ecology, five-

volume set. Elsevier Inc.

Amoroso CR, Antonovics J (2020) Evolution of behavioural resistance in host–pathogen

systems. Biol Lett 16: 20200508. https://doi.org/10.1098/rsbl.2020.0508

Arthur, A. P., & Wylie, H. G. (1959). Effects of host size on sex ration, development time

and size of Pimpla turionellae (L.)(Hymenoptera: Ichneumonidae). *Entomophaga*, *4*(4),

297-301. https://doi.org/10.1007/BF02373365

Askew RR, Shaw MR (1986) Parasitoid communities: their size, structure and

development. In: Insect parasitoids, pp 225–264.

Beckage, N. E., & Riddiford, L. M. (1978). Developmental interactions between the

tobacco hornworm Manduca sexta and its braconid parasite Apanteles congregatus

1. *Entomologia Experimentalis et Applicata*, *23*(2), 139-151.

https://doi.org/10.1111/j.1570-7458.1978.tb03016.x

Benrey, B., & Denno, R. F. (1997). The slow-growth–high-mortality hypothesis: a test

using the cabbage butterfly. *Ecology*, *78*(4), 987-999. **https://doi.org/10.1890/0012-**

9658(1997)078[0987:TSGHMH]2.0.CO;2

Buckingham LJ, Ashby B (2024) Coevolution of age-structured tolerance and virulence.

Bull Math Biol 86: 62. https://doi.org/10.1007/s11538-024-01292-2

Castelo, M. K., Nouhuys, S. V., & Corley, J. C. (2010). Olfactory attraction of the larval

parasitoid, Hyposoter horticola, to plants infested with eggs of the host butterfly, Melitaea

cinxia. *Journal of Insect Science*, *10*(1), 53. https://doi.org/10.1673/031.010.5301

Clancy, K. M., & Price, P. W. (1987). Rapid herbivore growth enhances enemy attack:

sublethal plant defenses remain a paradox. *Ecology*, *68*(3), 733-737.

https://doi.org/10.2307/1938479

Dieckmann U, Law R (1996) The dynamical theory of coevolution: a derivation from

stochastic ecological processes. J Math Biol 34: 579–612.

https://doi.org/10.1007/BF02409751

Eshel I, Akin E, et al. (1983) Coevolutionary instability of mixed Nash solutions. J Math

Biol 18: 123–133. https://doi.org/10.1007/BF00280661

Fisher RA (1930) The genetical theory of natural selection. Oxford.

GAULD, I. D. (1988). Evolutionary patterns of host utilization by ichneumonoid

parasitoids (Hymenoptera: Ichneumonidae and Braconidae). *Biological Journal of the*

Linnean Society, *35*(4), 351-377. https://doi.org/10.1111/j.1095-8312.1988.tb00476.x

Godfray HCJ (1994) Parasitoids: behavioral and evolutionary ecology. Vol. 12. Princeton

University Press.

Godwin, P. A., & Odell, T. M. (1984). Laboratory study of competition between

Blepharipa pratensis and Parasetigena silvestris (Diptera: Tachinidae) in Lymantria dispar

(Lepidoptera: Lymantriidae). *Environmental Entomology*, *13*(4), 1059-1063.

https://doi.org/10.1093/ee/13.4.1059

Hackett-Jones E, White A, Cobbold CA (2011) The evolution of developmental timing in

natural enemy systems. J Theor Biol 275: 1–11. https://doi.org/10.1016/j.jtbi.2010.12.040

Haeselbarth E (1978) Zur parasitierung der puppen von Forleule (Panolis flammea

[Schiff.]), Kiefernspanner (Bupalus piniarius [L.]) und Heidel-beerspanner (Boarmia

bistortata [Goeze]) in bayerischen kiefernwäldern 1: teil 1. Z Angew Entomol 87: 186–202.

https://doi.org/10.1111/j.1439-0418.1978.tb02442.x

Harvey, J. A. (2000). Dynamic effects of parasitism by an endoparasitoid wasp on the

development of two host species: implications for host quality and parasitoid

fitness. *Ecological Entomology*, *25*(3), 267-278. **https://doi.org/10.1046/j.1365-**

2311.2000.00265.x

Harvey JA (2005) Factors affecting the evolution of development strategies in parasitoid

wasps: the importance of functional constraints and incorporating complexity. Entomol Exp

Appl 117: 1–13. https://doi.org/10.1111/j.1570-7458.2005.00348.x

Harvey, J. A., & Strand, M. R. (2002). The developmental strategies of endoparasitoid

wasps vary with host feeding ecology. *Ecology*, *83*(9), 2439-2451.

https://doi.org/10.1890/0012-9658(2002)083[2439:TDSOEW]2.0.CO;2

Harvey, J. A., Bezemer, T. M., Elzinga, J. A., & Strand, M. R. (2004). Development of the

solitary endoparasitoid Microplitis demolitor: host quality does not increase with host age

and size. *Ecological Entomology*, *29*(1), 35-43. **https://doi.org/10.1111/j.0307-**

6946.2004.00568.x

Harvey, J. A., Gols, R., Vet, L. E., & Kruidhof, H. M. (2012). Development of a

hyperparasitoid wasp in different stages of its primary parasitoid and secondary herbivore

hosts. *Journal of insect physiology*, *58*(11), 1463-1468.

https://doi.org/10.1016/j.jinsphys.2012.08.013

Harvey, J. A., Harvey, I. F., & Thompson, D. J. (1994). Flexible larval growth allows use

of a range of host sizes by a parasitoid wasp. *Ecology*, *75*(5), 1420-1428.

https://doi.org/10.2307/1937465

Harvey, J. A., Jervis, M. A., Gols, R., Jiang, N., & Vet, L. E. (1999). Development of the

parasitoid, Cotesia rubecula (Hymenoptera: Braconidae) in Pieris rapae and Pieris brassicae

(Lepidoptera: Pieridae): evidence for host regulation. *Journal of Insect Physiology*, *45*(2),

173-182. https://doi.org/10.1016/S0022-1910(98)00113-9

Harvey, J. A., & Malcicka, M. (2016). Nutritional integration between insect hosts and

koinobiont parasitoids in an evolutionary framework. *Entomologia experimentalis et*

applicata, *159*(2), 181-188. **https://doi.org/10.1111/eea.12426**

Harvey, J. A., Poelman, E. H., & Tanaka, T. (2013). Intrinsic inter-and intraspecific

competition in parasitoid wasps. *Annual review of entomology*, *58*(1), 333-351.

https://doiorg/10.1146/annurev-ento-120811-153622

Harvey, J. A., & Vet, L. E. (1997). Venturia canescens parasitizing Galleria mellonella and

Anagasta kuehniella: differing suitability of two hosts with highly variable growth

potential. *Entomologia experimentalis et applicata*, *84*(1), 93-100.

https://doi.org/10.1046/j.1570-7458.1997.00202.x

Hurford A, Cownden D, Day T (2010) Next-generation tools for evolutionary invasion

analyses. J R Soc Interface 7: 561–571. https://doi.org/10.1098/rsif.2009.0448

Iritani R, Visher E, Boots M (2019) The evolution of stage-specific virulence: differential

selection of parasites in juveniles. Evol Lett 3: 162–172. https://doi.org/10.1002/evl3.105

Jervis M, Ferns P (2011) Towards a general perspective on life-history evolution and

diversification in parasitoid wasps. Biol J Linn Soc 104: 443–461.

https://doi.org/10.1111/j.1095-8312.2011.01719.x

Jervis MA, Ellers J, Harvey JA (2008) Resource acquisition, allocation, and utilization in

parasitoid reproductive strategies. Annu Rev Entomol 53: 361–385.

https://doi.org/10.1146/annurev.ento.53.103106.093433

Jervis MA, Ferns PN, Heimpel GE (2003) Body size and the timing of egg production in

parasitoid wasps: a comparative analysis. Funct Ecol 17: 375–383.

https://doi.org/10.1046/j.1365-2435.2003.00742.x

Jervis, M. A., & Kidd, N. A. C. (1986). Host-feeding strategies in hymenopteran

parasitoids. *Biological Reviews*, *61*(4), 395-434. **https://doi.org/10.1111j.1469-**

185X.1986.tb00660.x

King, B. H. (1989). Host-size-dependent sex ratios among parasitoid wasps: does host

growth matter?. *Oecologia*, *78*(3), 420-426. https://doi.org/10.1007/BF00379119

Lafferty KD, Allesina S, Arim M et al. (2008) Parasites in food webs: the ultimate missing

links. Ecol Lett 11: 533–546. https://doi.org/10.1111/j.1461-0248.2008.01174.x

Mayhew PJ, Blackburn TM (1999) Does development mode organize life-history traits in

the parasitoid Hymenoptera? J Anim Ecol 68: 906–916. https://doi.org/10.1046/j.1365-

2656.1999.00338.x

Maynard Smith J (1982) Evolution and the theory of games. In: Did Darwin get it right?

Essays on games, sex and evolution. Springer.

Ode, P. J., Vyas, D. K., & Harvey, J. A. (2022). Extrinsic inter-and intraspecific

competition in parasitoid wasps. *Annual Review of Entomology*, *67*(1), 305-328.

https://doi.org/10.1146/annurev-ento-071421-073524

Otto SP, Day T (2007) A biologist's guide to mathematical modeling in ecology and

evolution. Princeton University Press.

Pennacchio F, Strand MR (2006) Evolution of developmental strategies in parasitic

Hymenoptera. Annu Rev Entomol 51: 233–258.

https://doi.org/10.1146/annurev.ento.51.110104.151029

Price, P. W. (1972). Parasitiods utilizing the same host: adaptive nature of differences in

size and form. *Ecology*, *53*(1), 190-195. **https://doi.org/10.2307/1935729**

Quicke DLJ (2014) The braconid and ichneumonid parasitoid wasps: biology, systematics,

evolution and ecology. John Wiley & Sons.

Salt, G. (1941). The effects of hosts upon their insect parasites.

https://doi.org/10.1111/j.1469-185X.1941.tb01103.x

Sandlan, K. (1979). Sex ratio regulation in Coccygomimus turionella Linnaeus

(Hymenoptera: Ichneumonidae) and its ecological implications. **10.1111/j.1365-**

2311.1979.tb00596.x

Sequeira, R., & Mackauer, M. (1992). Covariance of adult size and development time in the

parasitoid wasp Aphidius ervi in relation to the size of its host, Acyrthosiphon

pisum. *Evolutionary Ecology*, *6*(1), 34-44. https://doi.org/10.1007/BF02285332

Slansky Jr, F. (1986). Nutritional ecology of endoparasitic insects and their hosts: an

overview. *Journal of Insect Physiology*, *32*(4), 255-261. https://doi.org/10.1016/0022-

1910(86)90036-3

Strand, M. R., Johnson, J. A., & Culin, J. D. (1988). Developmental interactions between

the parasitoid Microplitis demolitor (Hymenoptera: Braconidae) and its host Heliothis

virescens (Lepidoptera: Noctuidae). *Annals of the Entomological Society of America*, *81*(5),

822-830. https://doi.org/10.1093/aesa/81.5.822

Taylor PD (1990) Allele-frequency change in a class-structured population. Am Nat 135:

95–106. https://doi.org/10.1086/285034

Thompson MJ, Jiggins CD (2014) Supergenes and their role in evolution. Heredity 113: 1–

8. https://doi.org/10.1038/hdy.2014.20

Van den Driessche P, Watmough J (2002) Reproduction numbers and sub-threshold

endemic equilibria for compartmental models of disease transmission. Math Biosci 180:

29–48. https://doi.org/10.1016/S0025-5564(02)00108-6

Vinson, S. B. (1972). Effect of the parasitoid, Campoletis sonorensis, on the growth of its

host, Heliothis virescens. *Journal of Insect Physiology*, *18*(8), 1509-1514.

https://doi.org/10.1016/0022-1910(72)90229-6

Vinson, S. B., & Iwantsch, G. F. (1980). Host suitability for insect parasitoids.

10.1146/annurev.en.25.010180.002145

Wajnberg E, Bernstein C, Van Alphen J (2008) Behavioral ecology of insect parasitoids:

from theoretical approaches to field applications. John Wiley & Sons.

Wang ZZ, Liu YQ, Shi M, Huang JH, Chen XX (2019) Parasitoid wasps as effective

biological control agents. J Integr Agric 18: 705–715. https://doi.org/10.1016/S2095-

3119(18)62078-7