An updated catalogue of split-GAL4 driver lines for descending neurons in *Drosophila melanogaster*

Jessica L. Zung[1,*], Shigehiro Namiki[2,*†], Geoffrey W. Meissner[2], Marta Costa[3], Katharina Eichler[3,4], Tomke Stürner[3,5], Gregory S.X.E. Jefferis[3,5], Claire Managan[2], FlyLight Project Team[2], Wyatt Korff[2,#], Gwyneth M. Card[1,#]

[1]Howard Hughes Medical Institute and Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY, USA
[2]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA, USA
[3]Drosophila *Connectomics Group, Department of Zoology, University of Cambridge, Cambridge, UK*
[4]Genetics Department, Leipzig University, Leipzig, Germany
[5]Neurobiology Division, MRC Laboratory of Molecular Biology, Cambridge, UK
[†]Present address: Research Center for Advanced Science and Technology, University of Tokyo, Tokyo, Japan

*co-first authors
[#]co-corresponding authors; WK: korffw@janelia.hhmi.org; GMC: gc3017@columbia.edu

Abstract

Descending neurons (DNs) occupy a key position in the sensorimotor hierarchy, conveying signals from the brain to the rest of the body below the neck. In *Drosophila melanogaster* flies, approximately 480 DN cell types have been described from electron-microscopy image datasets. Genetic access to these cell types is crucial for further investigation of their role in generating behaviour. We previously conducted the first large-scale survey of *Drosophila melanogaster* DNs, describing 98 unique cell types from light microscopy and generating cell-type-specific split-Gal4 driver lines for 65 of them. Here, we extend our previous work, describing the morphology of 137 additional DN types from light microscopy, bringing the total number DN types identified in light microscopy datasets to 235, or nearly 50%. In addition, we produced 500 new sparse split-Gal4 driver lines and compiled a list of previously published DN lines from the literature for a combined list of 738 split-Gal4 driver lines targeting 171 DN types.

Introduction

A common feature of nervous systems across most of the animal kingdom is the presence of a brain, which coordinates most sensory processing and decision making, and a separate nerve cord—a ventral nerve cord or spinal cord—that performs final coordination of motor output. Communication from the brain to the nerve cord is relayed via a relatively small population of descending neurons (DNs). The privileged position of these neurons in the sensorimotor hierarchy makes them prime targets for the study of behaviour and motor control.

In *Drosophila melanogaster*, genetic resources and stereotypy of neurons across individuals allow us to generate cell-type-specific driver lines that help reveal the morphology of individual neurons and enable targeted manipulation or measurement of neurons. We previously created 178 sparse split-Gal4 driver lines targeting 65 DN types and described an additional 33 DN types labelled by more broadly expressing Gal4 lines (Namiki et al. 2018). This covered roughly half of the ~700 total DNs we estimated at the time based on counting DN cell bodies using a pan-neuronally expressed photoactivatable GFP.

Recent analyses based on electron-microscopy (EM) images have since revealed that there are in fact ~1300 DNs split across ~480 types in *Drosophila melanogaster* males (Cheong et al. 2024) and a similar number in females (Stuerner et al. 2024). These exhaustive catalogues underscore the value of EM-based connectomes for delivering a more complete picture of animal nervous systems. However, genetic driver lines are still required for *in vivo* manipulation or measurement of neurons. Indeed, our previous collection of descending-neuron lines has already been used to study *Drosophila* escape behaviour (Dombrovski et al. 2023), saccades during flight (Ros et al. 2024), walking initiation (Bidaye et al. 2020), steering during flight (Namiki et al. 2022) and walking (Chen et al. 2018; Rayshubskiy et al. 2024), freezing (Zacarias et al. 2018), landing (Ache et al. 2019), grooming (Guo et al. 2022), and diverse other behaviours (Cande et al. 2018). Here, we extend our previous work, describing an additional 137 DN types from light-level microscopy data and producing 500 additional DN driver lines. We also survey the literature for previously published DN lines to compile a combined list of 738 split-Gal4 driver lines targeting 171 DN types.

Results

Identification of new DN types

Our first goal was to individually characterize the morphology of descending neurons using a large database of light-microscopy (LM) brain and ventral nerve cord (VNC) images (Meissner et al. 2023). Although new EM datasets show neuron morphology with greater spatial resolution, all EM datasets published to date terminate at the neck (Stuerner et al. 2024), limiting their utility for describing complete DN morphology. We previously described 98 DN types from an initial survey of this LM database (Namiki et al. 2018) and now describe the results of our completed search below.

We manually searched publicly available images in Janelia's "Gen1" collection of broadly expressing Gal4 driver lines (Jenett et al. 2012). We specifically searched for lines that showed expression in the brain and VNC, using the MultiColor FlpOut (MCFO) technique to label neurons sparsely enough for easy visualization (Meissner et al. 2023). We identified DNs from these images, operationally defining them as in Namiki et al. (2018) as neurons with a soma in the brain and processes extending to the VNC (Fig. 1a,b). Using these criteria, we characterized the morphology of 137 new DN types from light-level microscopy and named them following previous convention (Namiki et al. 2018). Systematic names were generated based on the location of the soma in the central brain (Fig. 1c) (DNa: anterior dorsal, DNb: anterior ventral, DNc: pars intercerebralis, DNd: outside cell cluster on the anterior surface, DNg: gnathal ganglion, DNp: posterior surface of the brain, DNx: outside the brain) and assigned sequential numbers within these groups. We describe 6 new 'DNa' types, 3 new 'DNb' types, 2 new 'DNd' types, 85 new 'DNg' types, 40 new 'DNp' types, and 1 new 'DNx' type. DNs were also categorized by the VNC neuropils they innervated, following the convention from analysis of an EM VNC dataset (Cheong et al. 2024) (Fig. 1d) (fl: foreleg, ml: midleg, hl: hindleg, xl: multiple leg neuropils, nt: neck tectulum, wt: wing tectulum, ht: haltere tectulum, ut: upper tectulum, it: intermediate tectulum, lt: lower tectulum, ad: abdomen, xn: multiple neuropils).

We have now described a total of 235 DN types from LM images—approximately 50% of the types known from EM datasets (Fig. 1e,f). The morphologies of these neurons are shown in Figures 2 through 25. Nearly all of these have been matched to neurons in existing EM datasets (FAFB, MANC, FANC) by Stuerner et al. (2024).

Updates to 2018 eLife *paper*

While characterizing new DN types, we discovered subtle but consistent morphological variation among neurons within four DN types we previously defined in Namiki et al. (2018): DNb01, DNg05, DNg16, and DNp13. These variants have now been defined as new types. In each case that follows, the first name of each pair formerly encompassed both variants, but both names now describe distinct types: DNb01/DNb09, DNg05/DNg126, DNg16/DNg69, and DNp13/DNp62.

In addition, we note that the morphologies of several DN types described in Namiki et al. 2018 were atypical, due to developmental abnormalities or misidentifications in the image examples used to describe them. Figures 2 through 25 include revised images for all these cases. Our ground truth was the split-Gal4 lines annotated with a given DN type (if available) or the images found in the figure supplements to Figure 2 in Namiki et al. (2018). A full list of the revisions may be found in Supplementary Table 1.

We also update annotations for split-Gal4 lines produced by (Namiki et al. 2018), generally adding unannotated types and in a few cases removing annotations for types where we no longer see convincing evidence that they are labelled (see Supplementary Table 2).

New DN split-Gal4 lines

Our next goal was to create new cell-type-specific driver lines for as many DN types as possible using the intersectional split-Gal4 technique (Luan et al. 2006). Briefly, this method involves choosing pairs of enhancers whose expression pattern overlaps in the target cell type and using them to drive expression of the two halves of the GAL4 transcription factor, such that a functional GAL4 protein is expressed only in cell types labelled by both enhancers. Using the Janelia FlyLight image database (Meissner et al. 2023) and "split half" lines (Tirian and Dickson 2017; Dionne et al. 2018), we identified candidate enhancer pairs and crossed them to produce 500 new split-Gal4 driver lines that sparsely label DNs (Supplementary Files 1,2).

We gave each line an overall quality score from 'A' (best) to 'C' (worst) based on their specificity and consistency in labelling the annotated DN type(s). We used stringent criteria for lines given an 'A' score (see Methods for details), stricter than we used previously (Namiki et al. 2018). For these lines, optogenetic activation induced by illuminating the entire animal is expected to reliably activate the target neuron(s) with virtually no off-target effects in the rest of the central nervous system. For many other applications such as imaging or targeted optogenetic illumination, 'B' or even 'C' lines should be more than suitable. Examples of lines given different scores are shown in Supplementary Figure 1. In total, we scored 97 lines 'A', 325 'B', and 78 'C'. Supplementary Table 3 lists lines by DN type and quality score.

Compilation of previously published DN split-Gal4 lines

Many other research groups have created split-Gal4 driver lines that label descending neurons. We have compiled a list of the 60 lines we found in the literature and collated it with our own collection (Supplementary File 2; lines also included in Supplementary Table 3). While our search was by no means exhaustive, our list now represents the most comprehensive collection to date of DN driver lines, comprising 738 lines targeting 171 DN types.

Discussion

Genetic access is often a key prerequisite to understanding the function of a target cell type. Here, we provide a significantly expanded catalogue of driver lines that enable genetic access to DNs in *Drosophila melanogaster*. DNs are relatively few in number yet are the sole conduits of nervous signals from the brain to the ventral nerve cord and its downstream motor circuits. Thus, DNs are well poised to provide insight into the neural control of behaviour.

Optogenetic activation of DNs can be especially informative, since these neurons transmit information to motor neurons via a relatively short path, in some cases forming direct synapses. Indeed, activation of single bilateral DN pairs has been shown to drive whole coordinated motor programs such as forwards walking (Bidaye et al. 2020; Rayshubskiy et al. 2024), backwards walking (Bidaye et al. 2014), takeoff (von Reyn et al. 2014), landing (Ache et al. 2019), courtship song (Ding et al. 2019; Lillvis et al. 2024), and grooming (Hampel et al. 2015; Guo et al. 2022), reinforcing the idea that DNs convey high-level commands driving complex behaviours rather than lower-level coordination of individual muscle movements. There is also mounting evidence that subsets of DNs coordinate behaviour as a population. Namiki et al. (2018) noted that the number of individual neurons in a DN type, as defined by neurons with nearly identical morphology, typically ranges from two (a single left-right pair) to a couple dozen. Those found to activate coordinated motor programs thus far have mostly been types with only a single left-right pair, or "unique" DNs. In contrast, a "population" DN type with ~13 neurons per hemisphere represents the gain of the fly's wingstroke as the fraction of the population activated (Namiki et al. 2022). Further supporting population representation, activation of different DN types can evoke similar behaviours (Cande et al. 2018), and calcium-imaging experiments reveal that many DNs are active during multiple behaviours (Aymanns et al. 2022). Activation of a single DN type has also been shown to directly recruit other DNs through collateral connections (Braun et al. 2024). These observations suggest that many DNs may not act as "command neurons" that uniquely signal a specific behaviour, but instead encode activity as a population. In any case, results from optogenetic experiments must be interpreted with caution. While optogenetic activation of single types may indicate involvement in a particular behaviour, it is unlikely to recapitulate natural endogenous activation.

Our driver lines also enable recording of neural activity from the same identified DNs across multiple animals. Imaging using genetically encoded calcium indicators has revealed important insights into DN activity (Chen et al. 2018; Namiki et al. 2022; Aymanns et al. 2022). However, several studies have shown that precise spike timing is important for DN signalling (von Reyn et al. 2014; Rayshubskiy et al. 2024). The relatively low temporal resolution of calcium imaging may therefore obscure important biological details. Electrophysiological recordings allow much higher temporal resolution, and *in vivo* whole-cell patch clamp recordings in behaving flies is a well-established technique (Wilson et al. 2004; Maimon et al. 2010). DNs often have large cell bodies that are amenable to electrophysiology, and our collection of driver lines now enable many of them to be targeted effectively by expressing fluorescent markers. We also anticipate that our driver lines will be useful for voltage imaging as new improvements in genetically encoded voltage sensors and optical methods make this method more feasible.

For all experiments using these lines, researchers are strongly encouraged to examine expression patterns themselves to evaluate line cleanliness and consistency. Ideally, the expression pattern should be visualized using the actual effector gene(s) used in experiments, as the pattern may be effector-specific (Meissner et al. 2025). We have also occasionally observed changes in expression pattern after many generations of stock maintenance, perhaps due to epigenetic changes or mutation. Furthermore, it is good practice to replicate results using two or more lines to reduce or eliminate the confounding effects of off-target expression. Our collection includes two or more high-quality 'A' lines

for only 34 DN types (Fig. 1e,f), but some lines with lower specificity may still provide useful validation if lines do not contain overlapping off-target expression.

Rough estimates suggested that Janelia's Gen1 MCFO database should enable the creation of cell-type-specific lines for 50–80% of cell types (Meissner et al. 2023). However, after screening this database and surveying the literature for additional DN split-Gal4 driver lines, we report lines for only 171 (36%) of all DN types. We estimate this corresponds to ~550 (42%) of the roughly 1300 DNs counted from EM data. This lower-than-expected rate may partly reflect the database's slightly lesser focus on the VNC compared to the central brain. It may also suggest that cell-type-specific lines labelling new DNs can still be produced using existing images and split hemidrivers. We expect this may be feasible for some of the 64 (roughly 13% of) DN types that we have identified from broadly expressing Gal4 lines but that still lack specific split-Gal4 drivers. However, many of these were identified from only a single image. Using the split-Gal4 technique to generate additional cell-type-specific lines would require LM identification in at least one other driver line with expression in the desired DN type.

Approximately 240 DN types, or 50% of the total, have only been identified from EM data. While an intense, targeted effort may still uncover instances of these neurons in the Gen1 MCFO database, we believe our search was reasonably thorough. Further efforts using the methods we employ here are likely to yield diminishing returns, barring large-scale generation of new enhancer lines or at least new MCFO images of existing lines.

With the advent of connectomic data that enable precise identification of neurons involved in a given circuit, it is becoming increasingly urgent to gain genetic access to specific neurons of interest. New methods may help increase the efficiency of successfully generating desired split-GAL4 lines. For example, single-cell transcriptomic data enable the selection of gene pairs that often uniquely specify a single cell type. Knocking the two split-Gal4 halves into these loci are likely to produce a cell-type-specific line—a method successfully demonstrated by Chen et al. (2023). Spatial transcriptomics (Janssens et al. 2024) may be an important tool in facilitating this approach, allowing researchers to discover genes expressed in cell types that are identified morphologically from connectome data.

In summary, we complete our survey of existing resources for producing cell-type-specific DN lines and report a total of 738 lines covering roughly 36% of DN types. We anticipate that these lines will be useful for understanding the neural control of behaviour in *Drosophila melanogaster*.

Acknowledgements

This work was supported by the FlyLight Project Team at Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA. During this time, the FlyLight Project Team consisted of: Megan Atkins, Kari Close, Gina DePasquale, Zachary Dorman, Kaitlyn Forster, Theresa Gibney, Joanna Hausenfluck, Yisheng He, Kristin Henderson, Nirmala Iyer, Jennifer Jeter, Lauren Johnson, Rebecca Johnston, Rachel Lazarus, Kelley Lee, Hsing Hsi Li, Hua-Peng Liaw, Oz Malkesman, Geoffrey Meissner, Brian Melton, Scott Miller, Alexandra Novak, Omotara Ogundeyi, Alyson Petruncio, Jacquelyn Price, Sophia Protopapas, Susana Tae, Allison Vannan, Rebecca Vorimo, Brianna Yarbrough, and Kevin Xiankun Zeng, and the Steering Committee consisted of: Gerry Rubin, Barry Dickson, Reed George, Gwyneth Card, Wyatt Korff, Geoffrey Meissner. We also thank Gudrun Ihrke, Manager of Project Technical Resources, for helping to coordinate additional imaging and supervision of Claire Managan's efforts in performing neuron segmentations from light-level data.

References

Ache JM, Namiki S, Lee A, et al (2019) State-dependent decoupling of sensory and motor circuits underlies behavioral flexibility in Drosophila. Nat Neurosci 22:1132–1139. https://doi.org/10.1038/s41593-019-0413-4

Aymanns F, Chen C-L, Ramdya P (2022) Descending neuron population dynamics during odor-evoked and spontaneous limb-dependent behaviors. eLife 11:e81527. https://doi.org/10.7554/eLife.81527

Bidaye SS, Laturney M, Chang AK, et al (2020) Two brain pathways initiate distinct forward walking programs in *Drosophila*. Neuron 108:469-485.e8. https://doi.org/10.1016/j.neuron.2020.07.032

Bidaye SS, Machacek C, Wu Y, Dickson BJ (2014) Neuronal control of *Drosophila* walking direction. Science 344:97–101. https://doi.org/10.1126/science.1249964

Braun J, Hurtak F, Wang-Chen S, Ramdya P (2024) Descending networks transform command signals into population motor control. Nature 630:686–694. https://doi.org/10.1038/s41586-024-07523-9

Cande J, Namiki S, Qiu J, et al (2018) Optogenetic dissection of descending behavioral control in Drosophila. eLife 7:e34275. https://doi.org/10.7554/eLife.34275

Chen C-L, Hermans L, Viswanathan MC, et al (2018) Imaging neural activity in the ventral nerve cord of behaving adult Drosophila. Nat Commun 9:4390. https://doi.org/10.1038/s41467-018-06857-z

Chen Y-CD, Chen Y-C, Rajesh R, et al (2023) Using single-cell RNA sequencing to generate predictive cell-type-specific split-GAL4 reagents throughout development. Proc Natl Acad Sci 120:e2307451120. https://doi.org/10.1073/pnas.2307451120

Cheong HSJ, Eichler K, Stürner T, et al (2024) Transforming descending input into behavior: The organization of premotor circuits in the Drosophila Male Adult Nerve Cord connectome. 2023.06.07.543976

Ding Y, Lillvis JL, Cande J, et al (2019) Neural evolution of context-dependent fly song. Curr Biol 29:1089–1099. https://doi.org/10.1016/j.cub.2019.02.019

Dionne H, Hibbard KL, Cavallaro A, et al (2018) Genetic reagents for making Split-GAL4 lines in *Drosophila*. Genetics 209:31–35. https://doi.org/10.1534/genetics.118.300682

Dombrovski M, Peek MY, Park J-Y, et al (2023) Synaptic gradients transform object location to action. Nature 613:534–542. https://doi.org/10.1038/s41586-022-05562-8

Guo L, Zhang N, Simpson JH (2022) Descending neurons coordinate anterior grooming behavior in *Drosophila*. Curr Biol 32:823-833.e4. https://doi.org/10.1016/j.cub.2021.12.055

Hampel S, Franconville R, Simpson JH, Seeds AM (2015) A neural command circuit for grooming movement control. eLife 4:e08758. https://doi.org/10.7554/eLife.08758

Janssens J, Mangeol P, Hecker N, et al (2024) Spatial transcriptomics in the adult Drosophila brain and body. 2023.10.06.561233

Jenett A, Rubin GM, Ngo T-TB, et al (2012) A GAL4-Driver Line Resource for *Drosophila* Neurobiology. Cell Rep 2:991–1001. https://doi.org/10.1016/j.celrep.2012.09.011

Lillvis JL, Wang K, Shiozaki HM, et al (2024) Nested neural circuits generate distinct acoustic signals

during Drosophila courtship. Curr Biol 34:808-824.e6. https://doi.org/10.1016/j.cub.2024.01.015

Luan H, Peabody NC, Vinson CR, White BH (2006) Refined Spatial Manipulation of Neuronal Function by Combinatorial Restriction of Transgene Expression. Neuron 52:425–436. https://doi.org/10.1016/j.neuron.2006.08.028

Maimon G, Straw AD, Dickinson MH (2010) Active flight increases the gain of visual motion processing in *Drosophila*. Nat Neurosci 13:393–399. https://doi.org/10.1038/nn.2492

Meissner GW, Nern A, Dorman Z, et al (2023) A searchable image resource of Drosophila GAL4 driver expression patterns with single neuron resolution. eLife 12:e80660. https://doi.org/10.7554/eLife.80660

Meissner GW, Vannan A, Jeter J, et al (2025) A split-GAL4 driver line resource for Drosophila neuron types. eLife 13:RP98405. https://doi.org/10.7554/eLife.98405

Namiki S, Dickinson MH, Wong AM, et al (2018) The functional organization of descending sensory-motor pathways in Drosophila. eLife 7:e34272. https://doi.org/10.7554/eLife.34272

Namiki S, Ros IG, Morrow C, et al (2022) A population of descending neurons that regulates the flight motor of Drosophila. Curr Biol 32:1189-1196.e6. https://doi.org/10.1016/j.cub.2022.01.008

Nern A, Pfeiffer BD, Rubin GM (2015) Optimized tools for multicolor stochastic labeling reveal diverse stereotyped cell arrangements in the fly visual system. Proc Natl Acad Sci 112:E2967–E2976. https://doi.org/10.1073/pnas.1506763112

Rayshubskiy A, Holtz SL, Bates A, et al (2024) Neural circuit mechanisms for steering control in walking Drosophila. 2020.04.04.024703

Ros IG, Omoto JJ, Dickinson MH (2024) Descending control and regulation of spontaneous flight turns in Drosophila. Curr Biol 34:531-540.e5. https://doi.org/10.1016/j.cub.2023.12.047

Schlegel P, Yin Y, Bates AS, et al (2024) Whole-brain annotation and multi-connectome cell typing of Drosophila. Nature 634:139–152. https://doi.org/10.1038/s41586-024-07686-5

Stuerner T, Brooks P, Capdevila LS, et al (2024) Comparative connectomics of the descending and ascending neurons of the Drosophila nervous system: stereotypy and sexual dimorphism. 2024.06.04.596633

Tirian L, Dickson BJ (2017) The VT GAL4, LexA, and split-GAL4 driver line collections for targeted expression in the *Drosophila* nervous system. 198648

von Reyn CR, Breads P, Peek MY, et al (2014) A spike-timing mechanism for action selection. Nat Neurosci 17:962–970. https://doi.org/10.1038/nn.3741

Wilson RI, Turner GC, Laurent G (2004) Transformation of olfactory representations in the *Drosophila* antennal lobe. Science 303:366–370. https://doi.org/10.1126/science.1090782

Zacarias R, Namiki S, Card GM, et al (2018) Speed dependent descending control of freezing behavior in Drosophila melanogaster. Nat Commun 9:3697. https://doi.org/10.1038/s41467-018-05875-1

Methods

Identification of DN types and creation of split-Gal4 lines

We generally employed the same methods as in Namiki et al. (2018), except that we occasionally used DN annotations from newly available EM datasets (Cheong et al. 2024; Stuerner et al. 2024; Schlegel et al. 2024) to help clarify morphology and aid in segmentation.

In brief: We manually searched the Janelia Gen1 MCFO database (Meissner et al. 2023) to identify distinct descending neuron types. For newly identified types, we selected a representative image from either a Gen1 Gal4 line or a sparsely labelled split-Gal4 line and semi-manually produced the segmentations shown in Fig. 2–25. Image signal was thresholded to create a mask using the Amira 'Interactive Thresholding Function' or the Fiji 'Threshold' function, then manually corrected using other LM images and/or matching EM neurons as a reference. We then applied this mask to the original images to produce the final images shown in Fig. 2–25.

We created split-Gal4 driver lines by crossing hemidriver lines (containing either the GAL4 activation domain (AD) or DNA binding domain (DBD)) from the collections described in (Tirian and Dickson 2017) and (Dionne et al. 2018). Candidate combinations of split-Gal4 halves were chosen based on images in the Janelia Gen1 MCFO database (Meissner et al. 2023). Combinations producing a line that sparsely labelled the target cell type were made into stable stocks (i.e., homozygous or carrying balancer chromosomes). Final expression patterns were evaluated by crossing stable driver lines to a line containing 20XUAS-CsChrimson-mVenus in attP18. Select lines were also crossed to a reporter line containing 5XUAS-IVS-myr::smFLAG in VK00005, pJFRC51-3XUAS-IVS-Syt::smHA in su(Hw)attP1 to obtain polarity information. For lines labelling multiple cell types, we often also used the MCFO technique (Nern et al. 2015) to discern the morphology and identity of individual neurons.

Scoring of line specificity and consistency

Line specificity and consistency was evaluated using all available Janelia images that did not use stochastic effectors (i.e., all non-MCFO images).

We used the following criteria for line quality scores:

A - Target cell type is consistently expressed bilaterally. Background expression is zero or in very few cell types. Any background is extremely faint and typically also stochastic relative to expression of the target cell type.

B - Fails one of the three criteria for A. E.g., may have low/no background, but stochastic expression; sparse but bright background expression; or faint, diffuse background expression.

C - Target cell type is very inconsistently expressed and/or there is high background expression, especially background that overlaps spatially with the target cell type.

Figure 1. (**a–b**) Upper right schematics show in blue the locations of the brain (a; frontal head view) and VNC (b; top view of head and thorax) within a fly body. Lower right schematics show the general anatomy of a DN, including a soma in the brain (red dotted circle) (a) and processes in the VNC (b). Fly cartoons from Wikimedia Commons. (**c**) Schematic showing the approximate location of DN somas according to the naming scheme used in (Namiki et al. 2018). For clarity, the left-hand illustration shows locations of each soma group on only one side of the bilaterally symmetric brain. DNx somas (not shown) are outside the brain. (**d**) Schematic showing the approximate location of VNC neuropils by which DNs were categorized systematically in (Cheong et al. 2024). As in (b), the dorsal/ventral views often show neuropil location on only one side of the bilaterally symmetric VNC. 'xn' DN types (not shown) innervate multiple neuropils. (**e**) Bar plot showing the number of DN types in each soma-location category. Coloured bars indicate the number of types for which there exist at least 2 highly specific ('A') lines (teal), 1 highly specific line (dark blue), only non-specific or inconsistently expressing lines ('B' or 'C') (light blue), or no lines (grey). DN types with no corresponding split-Gal4 lines are further separated by whether they were ever identified from light microscopy images in a very broadly expressing Gal4 line (dark grey) or identified only from electron microscopy images (light grey). (**f**) Bar plot showing the number of DN types innervating each VNC neuropil. As in (c), coloured bars indicate the number of types for which there exist a given number of specific or non-specific lines.

Figures 2–25. Morphology of descending neurons identified in Namiki et al. 2018 and in the present study. (See following pages.)

Figure 2

Figure 4

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

Figure 12

Figure 13

Figure 14

Figure 15

Figure 16

Figure 17

Figure 18

Figure 19

Figure 20

Figure 21

Figure 22

Figure 23

Figure 24

No data

Figure 25

DNx01

DNx02

Supplementary Figure 1. Maximum-intensity projections showing DNp17 lines representative of 'A', 'B+', 'B', and 'C' scores for cleanliness/consistency. Colour indicates depth in the Z axis (violet→pink = anterior→posterior in the brain and ventral→dorsal for the VNC).

Supplementary Table 1. List of updates to the morphology of DNs described in Namiki et al. 2018.

Supplementary Table 2. List of annotation updates for split-Gal4 lines described in Namiki et al. 2018. Cell-type annotations were removed if images available online (at https://splitgal4.janelia.org/ and https://flylight-raw.janelia.org/) did not show the cell type listed or showed it was extremely faintly expressed.

Supplementary Table 3. List of all DN types identified from light-level microscopy and split-Gal4 lines that label them (from this study or any previously published source we could find). Lines were given quality scores from 'A' (best) to 'C' (worst) for their specificity and consistency. If any type had fewer than two 'A'-quality lines, we bumped 'B' lines to a 'B+' category if they met a certain minimum standard. Asterisks mark lines with tentative identifications for a given DN type.

Supplementary File 1. One representative maximum-intensity projection showing the expression pattern in the brain and VNC for each new split-Gal4 line produced in this study. Colour indicates depth in the Z axis (violet→pink = anterior→posterior in the brain and ventral→dorsal for the VNC). Images are aligned to a unisex template.

Supplementary File 2. Spreadsheet listing all split-Gal4 lines labelling descending neurons that were produced in this study, produced in Namiki et al. 2018, or found in other previously published work.