Machine Learning-Assisted Decoding of Temporal Transcriptional Dynamics via

Fluorescent Timer

Nobuko Irie[1], Naoki Takeda[2], Yorifumi Satou[1], Kimi Araki[2,3], Masahiro Ono[1,4]

1 The Joint Research Center for Human Retrovirus Infection, Kumamoto University,

Kumamoto, Japan

2 Institute of Resource Development and Analysis, Kumamoto University, Kumamoto,

Japan

3 Center for Metabolic Regulation of Healthy Aging, Kumamoto University, Kumamoto,

Japan.

4 Department of Life Sciences, Imperial College London, London, United Kingdom

Running Title: ML-Decoding of Transcriptional Dynamics

Abstract:

Investigating the temporal dynamics of gene expression is crucial for understanding

gene regulation across various biological processes. Using the Fluorescent Timer

protein (Timer), the Timer-of-cell-kinetics-and-activity (Tocky) system enables analysis of

transcriptional dynamics at the single-cell level. However, the complexity of Timer data

has limited its broader application. Here, we introduce an integrative approach

combining molecular biology and machine learning to elucidate Foxp3 transcriptional

dynamics through flow cytometric Timer analysis. We have developed a Convolutional

Neural Networks (ConvNet) approach that incorporates image conversion and Gradient-

weighted Class Activation Mapping (Grad-CAM) for class-specific feature identification at

the single-cell level. Biologically, we developed a novel CRISPR mutant of Foxp3-Tocky

lacking the Conserved Non-coding Sequence 2 (CNS2), which has successfully

elucidated CNS2-dependent Foxp3 transcription dynamics, revealing novel roles of

CNS2 in regulating Foxp3 transcription frequency under specific conditions.

Furthermore, generating new data from WT Foxp3 Tocky mice at various ages, the

Grad-CAM methods successfully revealed distinct dynamics of Foxp3 expression from

neonatal to aged mice, highlighting prominent thymus-like features of neonatal splenic

Foxp3$^+$ T cells. In conclusion, our study uncovers previously unrecognised Foxp3

transcriptional dynamics, establishing a proof-of-concept for integrating CRISPR, Tocky,

and machine learning methods as advanced techniques to understand transcriptional

dynamics in vivo. (200 words)

Keywords: Fluorescent Timer protein, Flow cytometry, Random Forest, Convolutional

Neural Network (ConvNet), Gradient-weighted Class Activation Mapping (Grad-CAM),

Transcription, Foxp3

Introduction:

Understanding the temporal dynamics of gene expression is fundamental to

comprehending gene regulation's functional significance, cellular differentiation, and

development. Single-cell level analyses of diverse tissues including differentiating cells [1,

2] and developing or functioning T cells [3, 4, 5] have utilised trajectory analyses to

reconstruct in vivo activation or differentiation processes. However, trajectory analysis

inherently relies on similarity analysis through dimensional reduction, which is not a

direct measurement of temporal dynamics in vivo. Thus, the current challenge lies in

accurately capturing and measuring temporal elements of these processes within

individual cells.

Fluorescent Timer proteins offer a unique solution to this challenge. Previous studies

include a mathematical modelling approach to cell population dynamics [6]. However, the

full integration of single-cell techniques with a Timer-based approach has yet to be fully

realised. We previously developed the Timer-of-cell-kinetics-and-activity (Tocky) system

for single-cell flow cytometric analysis of cellular activities and transcription [7]. Tocky

uses a mutant mCherry fluorescent timer protein, Fast-FT, which spontaneously and

irreversibly changes its chromophore from blue to a mCherry-type red form with the

maturation half-life 4.1 hours [8]. The mature red-form Timer protein is stable and its

decay rate is 122 hours [9]. Using this Tocky approach, we have developed Foxp3-Tocky

and Nr4a3-Tocky mice, enabling us to analyse the temporal dynamics of Foxp3

transcription and transcriptional activities downstream of T cell receptor (TCR) signalling,

respectively [7].

While Foxp3 has traditionally been designated as the lineage-specific transcription factor

for regulatory T cells (Tregs), implying a stable and continuous expression as its default

state [10], it is increasingly recognised as a dynamically regulated gene within CD4$^+$ T cells

[11, 12, 13]. The Foxp3-Tocky system has significantly contributed to our understanding by uncovering the highly dynamic expression of Foxp3 during both inflammatory and homeostatic states [9]. Key activating signals for Foxp3 include TCR, Interleukin-2 (IL-2) and transforming growth factor-beta (TGF-β). In addition, Foxp3 expression is regulated by an autoregulatory loop, where Foxp3 protein impacts its own transcription [9]. Intriguingly, the interaction between Foxp3 and Runx1 not only controls Foxp3 function [14] but also affects Foxp3 transcription [15]. Along with Stat5, NFAT [16], CREB, and Ets-1 [17, 18, 19], Foxp3 and Runx1 bind to Conserved Non-Coding Sequence 2 (CNS2) [20], which is a key enhancer in the Foxp3 gene and is essential for maintaining Foxp3 expression via epigenetic modifications [16, 20].

Previous studies have provided foundational insights into CNS2 and its role in sustaining Foxp3 expression post-cell division and potential roles in reactive expression in response to TCR and IL-2 signals [16, 20, 21]. Specifically, these studies demonstrated that, in CNS2 KO T cells, Foxp3 expression is lost as cells divide in culture [20, 21], and under inflammatory conditions, Foxp3$^+$ T cells can lose Foxp3 expression [21]. However, past research has relied predominantly on bulk analyses of cell populations from CNS2 KO mice, where the CNS2 region of the endogenous Foxp3 gene is deleted. Such approaches have not pinpointed specific cells in which CNS2 is actively functioning. Thus, it remains unresolved how a functioning CNS2 controls Foxp3 transcription in real time at single-cell resolution, what types of temporal dynamics of transcription active CNS2 induces, and whether and how IL-2 and TCR signal downstream genes are induced under intact CNS2 activity. This gap in understanding underscores the need for methodologies that can dissect these temporal dynamics at a granular, single-cell level, to better elucidate the functional mechanisms of CNS2.

Meanwhile, the influence of developmental and ageing processes on Foxp3 transcription remains largely unknown. Previous studies often concentrate on thymic neonatal stages alone, without offering a comprehensive spleen-thymus comparison [22]. Compounding this issue, controversial evidence regarding Treg developmental dynamics in neonates has circulated widely, creating substantial confusions in the research field [23]. While Foxp3-expressing T cells are known to accumulate in aged individuals, the biological significance of this accumulation remains unclear [24]. Moreover, studies examining aged spleen and thymus typically involve only two time points—adult and aged [25, 26]. Accordingly, our current study aims to develop standard quantitative data on the dynamic regulation of Foxp3 transcription across tissues and throughout both development and ageing.

To address our biological aims, we have developed a novel data-driven framework integrating advanced molecular biology techniques, including Foxp3-Tocky and CRISPR-mediated mutagenesis within the reporter construct of the Tocky system. Given the continuous nature of Timer fluorescence distribution [9], it is essential to eliminate manual gating methods from our research framework. Manual gating, constrained by manually pre-determined gates, is vulnerable to arbitrariness and subjectivity, leading to non-reproducibility [27], and is particularly problematic given the continuous nature of Timer Blue and Red fluorescence data. Additionally, while flow cytometry provides high-throughput single-cell analysis, its static snapshots of dynamic processes, combined with the limited dynamic range of Timer fluorescence, present significant challenges [28]. Moreover, there is no universal mathematical approach to transforming Timer fluorescence data back into transcriptional kinetics.

To overcome these limitations, we have developed a suite of machine learning (ML) methodologies specifically tailored for flow cytometric Timer analysis. These

methodologies directly analyze the spatial patterns in two-dimensional flow cytometric

Timer data, enabling a comprehensive capture of group-specific Timer dynamics. By

eliminating the arbitrariness and subjectivity associated with manual gating and

obviating the need for rigid assumptions about Foxp3 transcriptional dynamics or Timer

profiles, this approach ensures that the outputs are data-driven and reproducible. The

effectiveness of our methodologies is confirmed by quantitative model performance

metrics, underscoring their reliability.

Supervised ML approaches including Random Forest (RF) and Convolutional Neural

Networks (ConvNets) have been successfully applied to genomic data [29, 30] and

multidimensional flow cytometric data [31, 32, 33, 34, 35]. However, methods designed for

multi-marker datasets can introduce bias when applied directly to Timer fluorescence

data [36]. Thus, we introduce two complementary approaches: TockyKmeansRF, which

integrates clustering with RF analysis, and TockyConvNet, a ConvNet framework

employing a novel image conversion technique and Gradient-weighted Class Activation

Mapping (Grad-CAM). This toolkit moves beyond conventional gating and unsupervised

clustering in flow cytometry, enabling a more sophisticated and data-oriented analysis of

Foxp3 transcriptional dynamics at the single-cell level.

Results

Overview of Novel ML Approaches to Analyse Flow Cytometric Tocky Data

Figure 1a outlines the significant pitfalls and risks associated with 'manual gating', the

most prevalent method in immunology for identifying cells of interest in cytometry

analysis. The most widely used gating methods are rectangle, polygon and ellipse gates,

all of which are arbitrarily hand-drawn and highly problematic. While these gates allow

immunologists "flexible" identification of cell populations to test a hypothesis, they

inherently introduce substantial arbitrariness and subjectivity, increasing variability and

reducing transparency in data analysis [37, 38]. Gates depend on predefined features of cell

populations (e.g., a rectangle gate for $CD25^{high}$ $Foxp3^{high}$), embedding various hidden

and ambiguous assumptions and potential biases into the analysis. Designed to isolate

'populations of interest' to test a 'hypothesis', such arbitrariness makes the analytical

process prone to *'confirmation bias'*—the tendency to cherry-pick data that supports pre-

existing beliefs [29]. These arbitrary and subjective elements undermine the transparency

and rigor of data analysis, contributing to the reproducibility crisis, which is widespread in

life sciences and preclinical studies [39], including the research fields using flow cytometric

analysis [27].

In our strategic shift from manual gating to a data-oriented approach, we have

developed a new research framework that utilizes ML to enable coherent two-

dimensional analysis of Timer Blue and Red fluorescence, departing from conventional

two-variable analysis and significantly enhancing the power of Tocky. **Figure 1b** depicts

a workflow within this framework, aimed at unravelling transcriptional dynamics in a

functional system through ML-assisted identification of group-specific features. Using the

Tocky system, transcriptional dynamics influenced by an enhancer are investigated by

CRISPR-induced mutation of the enhancer within the Foxp3 Timer transgene.

Independent experiments are conducted to generate training and test datasets through

flow cytometric analysis of the Timer fluorescence profile of T cells. ML models are then

trained to classify samples into experimental groups using the Timer fluorescence data.

Subsequently, model performance analysis is conducted using the test data to obtain

model performance metrics and validate the trained model. The behavior of the trained

model is then analyzed by tailored methods, which enables the identification of group-

specific '*feature cells*.' These feature cells represent the group-specific *features* that

assist in the classification of samples across datasets. Importantly, our approach is

designed to perform cross-dataset analysis by applying the trained ML model to new data inputs to dynamically identify group-specific feature cells. The robustness of the identified group-specific features is quantitatively supported by the model performance metrics, enhancing transparency and rigor of the data analysis process.

Under this novel research framework, we developed and implemented two primary ML approaches, *TockyKmeansRF* and *TockyConvNet*, each tailored to identify feature cells within Timer fluorescence data (**Figure 1c**). Both methods are used to train ML models as classifiers, which are monitored and analyzed to identify group-specific features. TockyKmeansRF integrates k-means clustering with Random Forest (RF) analysis, using the mean decrease Gini index to monitor model behaviour. Meanwhile, TockyConvNet converts this data into two-dimensional "images" for ConvNet analysis, employing Grad-CAM to identify key regions influencing network predictions [40]. In addition, we show the capability of TockyConvNet to establish a continuous scoring system for quantitatively analyze cellular phenotype.

To use these models, flow cytometric Timer Blue and Red fluorescence data are pre-processed and transformed into Timer Angle and Timer Intensity, as previously described [7] and implemented as a computational tool [41]. The 'Timer Angle' is measured from the y-axis, represented by Timer Blue fluorescence, towards the x-axis, represented by Timer Red fluorescence. 'Timer Intensity' is the magnitude (or *norm*) of the vector formed by these fluorescence values (**Figure 1c**).

A Novel Experimental Tool to Investigate the Roles of Conserved Non-Coding Sequence 2 (CNS2) in Regulating Temporal Dynamics of Foxp3 transcription

To identify a biologically significant enhancer sequence and establish a prototypic approach to studying Foxp3 transcriptional dynamics, we analysed Chromatin

Immunoprecipitation sequencing (ChIP-seq) data. Our analysis demonstrated that both Foxp3 and Runx1 proteins uniquely bound to the CNS2 region of the Foxp3 gene (**Figure 2a**) as reported previously [20]. Importantly, our investigations using Foxp3-Tocky revealed that Foxp3 protein is required for sustaining Foxp3 transcription [9] and that the CNS2 region is actively demethylated at the moment when Foxp3 transcription is sustained and persistent in the CD4 single-positive thymocytes [7]. Therefore, we hypothesised that CNS2 functions as a platform for critical transcription factors, including Foxp3 itself and Runx1, to dynamically regulate the Foxp3 transcriptional activities. Deleting the CNS2 sequence should therefore elucidate the temporal phases of Foxp3 transcriptional regulation that are dependent on CNS2 (**Figure 2b**).

The deletion of a sequence within a bacterial artificial chromosome (BAC) transgene could be done in vitro, followed by the creation of a new mouse strain. However, such an approach is susceptible to between-founder variations, an inherent issue in BAC reporter systems [42]. Meanwhile, modifying the endogenous Foxp3 sequence could make any output reporter measurement secondary to the modified dynamics of the Foxp3 protein [43]. Therefore, it was essential to delete the CNS2 sequence within the BAC Foxp3-Tocky transgene only, without disturbing the endogenous Foxp3 gene.

We achieved this by using a CRISPR KO method combined with a dedicated breeding strategy. Fertilised eggs from Foxp3-Tocky mice underwent CRISPR-based electroporation (Materials and Methods). To facilitate the deletion of CNS2, a single-stranded oligodeoxynucleotide carrying homology arms was used to enable homologous recombination and replace the CNS2 region with a short oligo (**Figure 2c**). Critically, we established two distinct PCR assays: a CNS2 deletion-specific PCR to detect both the endogenous and BAC Foxp3-Timer loci ("*Common CNS2 Deletion*") and Foxp3 Timer-specific PCR to discriminate WT Foxp3 Timer and CNS2 KO Foxp3 Timer ("*Foxp3*

Timer-Specific Discrimination PCR", **Figure 2d**). These assays identified founder mouse

#87, which carried the CNS2 KO Foxp3 Timer without any evidence of CRISPR editing

in the endogenous Foxp3 gene (**Figure 2e–2f**). Sanger sequencing further confirmed

the successful deletion and homologous recombination of the CNS2 locus in the BAC

transgene (**Figure 2g**).

The founder mouse #87 was used to establish a breeding line through successive

matings with WT mice. Over multiple generations spanning more than two years, we

selectively bred progeny that expressed the CNS2 KO Foxp3 Timer, consistently

backcrossing them to the B6 background. This extensive breeding and selection process

confirmed that the CNS2 KO Foxp3-Tocky transgene was stably inherited in a

Mendelian manner (**Figure 2h**), reassuring that the modification involved a single

transgene in an autosomal chromosome. Our breeding strategy and the backcrossing

ensured the CNS2 deletion was specific to the Foxp3 Timer transgene, thereby

effectively eliminating any possibility of CRISPR-induced alterations to the endogenous

Foxp3 locus. The specificity of the Foxp3 Timer-Specific Discrimination PCR was further

validated by Sanger sequencing (**Figure 2i–2j**). Based on these validations, we used

hemizygous Foxp3-Tocky and hemizygous CNS2 KO Foxp3-Tocky mice as parents in

all experiments, excluding double transgenics, and ensured that littermate analysis was

consistently employed throughout the study.

Analysis of CNS2 KO Foxp3-Tocky Using Established Methods

After establishing the CRISPR mutant strain, we first examined the effects of CNS2

deletion with conventional flow cytometric analyses (**Figure 2k**). Mean fluorescence

intensity (MFI) measurements revealed only a moderate, albeit significant, decrease in

Timer Red fluorescence in KO T cells (p < 0.01), with Timer Blue largely unchanged

(**Figure 2l**). We next applied a trigonometric transformation [7, 44], converting Timer

fluorescence into Timer Angle and Intensity. Although the Timer Locus categorisation

method [44] indicated that active Foxp3 transcription, identified as *NPt*, *Persistent*, and *PAt*

categories, was reduced in KO T cells (**Figure 2m**), these existing approaches rely on

predominantly one-dimensional analyses and failed to capture the nuanced dynamics of

CNS2-mediated Foxp3 transcription. This limitation underscored the need for more

comprehensive methods to dissect Foxp3 transcriptional regulation.

TockyKmeansRF: Clustering and Random Forest Analysis of Timer Fluorescence

Figure 3a illustrates the TockyKmeansRF implementation, a combinatorial ML method

that integrates k-means clustering with RF classification. TockyKmeansRF constructs an

RF model using training flow cytometric Timer fluorescence data, which is subsequently

tested on an independent test dataset. Initially, TockyKmeansRF applies k-means

clustering separately to both training and test datasets, producing two tables that show

the percentage of cells in each cluster. Clusters between these datasets are matched

based on the Euclidean distances between them (Materials and Methods). The cluster

percentage table from the training set is then utilised to build an RF model, which is

evaluated using the corresponding table from the test set. The mean decrease Gini

(MDG) index helps identify significant clusters and, consequently, feature cells within the

original Tocky data [45].

To validate the effectiveness of the ML approaches in the current study, we generated

two independent datasets by conducting flow cytometric analysis on lymph node

samples from WT Foxp3 Tocky and CNS2 KO Foxp3 Tocky littermates. This process

produced the initial training and test datasets discussed in the following sections (**Figure

3b**).

First, we assessed the robustness of the TockyKmeansRF model by varying the number

of clusters in k-means clustering and the number of trees in the Random Forest model.

We confirmed stable performance across a range of cluster and tree numbers by

conducting area under the curve (AUC) analysis (**Figure 3c**). The TockyKmeansRF

model demonstrated a commendable classification accuracy. Constructed using training

data with 18 clusters and subsequently tested on an independent dataset, the model

achieved an out-of-bag (OOB) error rate of 7.69%. The confusion matrix from the test

dataset indicates a high predictive accuracy, with an overall accuracy of 91.18%. This

performance underscores the model's robustness and its capability to distinguish

effectively between the KO and WT classes within the testing framework, supporting the

significance of the feature cells as follows.

Using the MDG index as an importance score, CNS2-dependent feature cells were

identified among individual single cells within the test dataset, specifically in the Timer

Angle and Intensity space and within the original Timer fluorescence data (**Figure 3d**).

Density-based clustering of these feature cells revealed three distinct clusters (**Figure**

3e). Cluster 1 was predominantly found in KO mice, whereas Clusters 2 and 3 were

more prevalent in WT mice in the training data (**Figure 3f**). Notably, cells in Cluster 2

from WT Foxp3 Tocky mice exhibited high expression levels of CD25 (Interleukin-2

receptor alpha chain) and PD-1. Meanwhile, cells in Cluster 3 showed increased

expression of CD69 and CD44 (**Figure 3g**). Thus, each cluster had a unique activation

profile and CNS2 KO T cells shifted from Clusters 2 and 3 to Cluster 1, markedly

reducing Timer Intensity and approaching to the Timer Angle 90, which indicates the

arrested transcription [7].

Computational performance metrics, such as runtime and memory usage, were

evaluated during the execution of TockyKmeansRF on the scaled CNS2 KO dataset

(**Figure 3h**). During training, the maximum runtime was approximately 3 seconds per training session, and peak memory usage did not exceed 30 MB, even as the sample size increased to 136. Notably, the number of trees in the RF model did not significantly impact either runtime or memory usage.

These findings underscore the efficacy of TockyKmeansRF in identifying unique CNS2-dependent feature cells and classifying CNS2 KO-specific patterns. This demonstrates the model's robustness and precision in analysing complex dynamics of Timer fluorescence profiles, which are crucial for establishing a data-oriented approach to studying temporal transcriptional dynamics using the Tocky system.

TockyConvNet: A ConvNet Approach Using Grad-CAM for Discriminating WT and CNS2 KO Foxp3 Timer

The successful deployment of TockyKmeansRF, coupled with the optimisation of a relatively high number of clusters, indicates that transforming Timer fluorescence data into image data could open new avenues for ML applications, particularly by leveraging ConvNet technologies. The conversion process involved binning the data into 100 x 100 pixel images (**Figure 4a**), effectively preserving the essential visual characteristics of the Timer data after conversion (**Figure 4b**).

To prevent overfitting, for TockyConvNet we designed a compact ConvNet model consisting of two convolutional layers. Each layer features a sigmoid-activated pointwise convolution, termed "Spatial Attention", enhancing spatially relevant feature extraction (Materials and Methods). The architecture includes two dense layers (**Figure 4c**). The model successfully learnt through three-fold cross-validation using a relatively small number of training epochs (**Figure 4d**).

The model's efficacy was validated through ROC analysis on an independent test dataset. Furthermore, we benchmarked the TockyConvNet model against traditional manual gating methods employed in Fluorescent Timer analysis. These manual methods include Quadrant gating for distinguishing Timer Blue and Red positivity, and Polygons for cells with high Blue levels above and below the diagonal line between Timer Blue and Red, *Polygon-Blue(high)* and *Polygon-Red(high)*, respectively. TockyConvNet achieved excellent performance metrics, with both the Area Under the Curve (AUC) of the ROC and Average Precision scoring 1.0. In contrast, manual gating methods demonstrated significantly lower performance: the polygon gate scored 0.87 for AUC and 0.76 for Average Precision, while the quadrant gate scored 0.5 for AUC and 0.41 for Average Precision (**Figure 4e–4f**).

Grad-CAM Analysis of CNS2-Dependent Foxp3 Timer Dynamics

To elucidate CNS2-dependent Foxp3 transcription dynamics, Grad-CAM was applied across various convolutional layers to visualize transcriptional features that distinguish two genotypes. Precisely, for each layer, gradients for each pixel across all feature maps were calculated, globally averaged, and weighted. These weighted feature maps were then transformed into a single heatmap through pixel-wise summation, retaining only positive activations using the ReLU function (Materials and Methods).

To analyse the features of CNS2-dependency effectively, we generated differential heatmaps from Grad-CAM outputs of WT and KO samples using each convolutional layer of our model (**Figure 4g**). These heatmaps identified pixels critical for classification, illustrating how Grad-CAM progressively reveals these pixels across convolutional layers. The outputs were then reverse-mapped to their respective positions in the Timer Angle-Intensity space and to their original locations in the raw Timer Blue and Red fluorescence space (**Figure 4h**).

Quantitative comparisons between the two genotypes were conducted by analyzing cells

in the top 90th percentile as WT CNS2 feature cells (CNS2-dependent), and those in the

bottom 10th percentile as KO feature cells (CNS2-independent). This analysis revealed

that while all convolutional layers discerned differences between the two Foxp3 Tocky

variants, the most pronounced differences were observed in the last convolutional layer,

Attention2-Conv (**Figure 4i**). In contrast, the cells increased in the KO group were

predominantly captured by the first three convolutional layers (**Figure 4j**).

The CNS2-dependent feature cells identified across all convolutional layers, as shown in

Figure 4i, exhibited higher CD44 expression compared to other Timer+ cells (**Figure**

4k). Notably, the cells highlighted by the Attention2-Conv layer displayed significantly

elevated expression of CD69 and PD1, distinguishing them from other Timer$^+$ cells.

Gene Regulation in CNS2-Dependent Feature Cells

To further explore the biological significance of CNS2-dependent feature cells in relation

to the temporal regulation of Foxp3 transcription, we analyzed RNA-seq data from flow-

sorted Timer-positive cells from WT Foxp3 Tocky mice, fractionated into B1, B2, R1, and

R2 fractions [9] (**Figure 5a**). Utilizing *TockyPrep* for data preprocessing [41], we converted

Timer fluorescence data from these sorted cells into Timer Angle and Intensity values.

This conversion facilitated the application of Grad-CAM analyses in **Figure 4g** to

dynamically identify CNS2-dependent and independent cells as CNS2 WT and KO

feature cells, based solely on Timer distribution within the flow cytometric data linked to

RNA-seq data.

Precisely, using the extensive Grad-CAM analyses shown in Figures 4g, 4h, 4i, and 4j,

CNS2-dependent cells were identified using *WT Feature* of Attention-Conv2, while

CNS2-independent, inactive cells were pinpointed using *KO Feature* of Conv2 (**Figure**

5b). This cross-dataset analysis revealed that the fraction B2 was highly enriched with CNS2-dependent cells, comprising over 50% of the cells in this fraction, while the other fractions contained only a few such cells (**Figure 5c**). Conversely, CNS2-independent cells were predominantly found in the fraction R2, representing over 50% of the cells, while more than 30% in the fraction R1 as well.

The CNS2-dependent fraction B2 were characterized by uniquely high expression of NFAT genes (Nfatc1 and Nfatc2), distinctly among TCR signal downstream genes. In contrast, the CNS2-independent fractions R1 and R2 highly expressed other TCR signal downstream genes including Egr1, Nr4a1, Nr4a3, Rel, and Rela, but notably not the NFAT genes (**Figure 5e**). These findings suggest that each fraction is associated with unique TCR signal dynamics, which may be also influenced by additional signaling pathways. Importantly, Foxp3 expression was highest in fraction B2, aligning with the high-frequency transcriptional dynamics observed by Foxp3 Tocky. In addition, CNS2-dependent B2 cells also showed elevated expression of genes associated with Foxp3 function, such as Tnfrsf4, Tnfrsf18, Ctla4, Icos, and the TGF-β receptor component Tgfbr1. Intriguingly, the dynamics of IL-2 signal-related genes in the B2 fraction showed a distinctive pattern, with upregulation of IL-2 receptors (Il2ra and Il2rb) and repression of Stat5a (**Figure 5f**).

Collectively, our results confirm that CNS2-dependent cells exhibit the highest Foxp3 expression, aligned with the highest-frequency of Foxp3 transcription revealed by Tocky (Figures 4c, 5b, and 5g). In addition, the gene expression profile of CNS2-dependent cells supports that CNS2 orchestrates Foxp3 transcription under finely tuned and unique TCR signaling dynamics predominantly mediated by NFAT. The downstream activities of TCR signaling notably exclude other well-characterized genes downstream of TCR

signaling such as NF-kB and Nr4a genes, and are potentially influenced by unique

dynamics of IL-2 signaling as well (**Figures 4c and 5g**).

Expanding the Application of TockyConvNet to Understand Developmental and Ageing

Foxp3 Transcription Dynamics

Having demonstrated the utility of TockyConvNet with the CNS2 KO datasets, we aimed

to further generalize the ConvNet method by generating and analysing independent flow

cytometric datasets using Foxp3-Tocky mice. To this end, we analysed CD4$^+$ T cells

from the spleen and the thymus of WT Foxp3-Tocky mice across various ages, including

neonates and aged mice, to capture the full spectrum of Foxp3 Timer dynamics

throughout the mouse lifespan.

We discovered notable variations in Foxp3 Timer profiles, influenced by age and organ

(**Figure 6a**). Thymic T cells displayed new and active Foxp3 transcription, which

decreased over time, resembling splenic Timer profiles in older mice. Particularly in

neonates at days 3 and 4 post-birth, we observed high levels of new Foxp3 transcription

in both thymus and spleen. On days 1 and 2 post-birth, the thymus included substantial

numbers of CD4-single positive cells, whereas the spleen had too few CD4+ T cells to

permit meaningful analysis.

The flow cytometric Foxp3 Timer data were normalised and transformed into Timer

Angle and Intensity formats to quantitatively analyse the datasets (**Figure 6b–6c**),

revealing dynamic and gradual changes between the tissues across different ages.

Notably, as mice aged, Timer Blue fluorescence in both splenic and thymic Foxp3

transcription diminished, while Timer expressing cells accumulated, particularly those

with low Blue and high Red fluorescence (**Figure 6a–6b**) with high Timer Angles (**Figure 6c**).

Thus, we generated independent training and test datasets (designated as "Foxp3 Neonatal-to-Ageing Benchmarking Data". The aim of the analysis was to understand the tissue-specific and age-dependent dynamics of Foxp3 Timer profiles. Traditional manual gating methods, including quadrant and polygon gates, along with mean Timer Angle, captured some aspects of these dynamics, especially when the age scale is transformed logarithmically (**Figures 6d–6e**). These observations support significant changes in Foxp3 transcription dynamics from early life into old age (**Figures 6d–6e**). In aged mice, Timer$^+$ cells accumulated, reflecting the impact of ageing on Foxp3 transcription.

These nuanced, continuous, and dynamic changes in Timer profiles across a broad range of samples make them ideal targets for analysis using TockyConvNet. To further validate this ConvNet method, we generated a training dataset and an independent test dataset, both consisting of splenic and thymic T cells from mice of various ages.

Development of the TockyConvNet for Quantitative Assessment of Thymus and Spleen Characteristics in Foxp3 Transcriptional Dynamics

First, we aimed to develop a TockyConvNet model that captures the features of thymic Foxp3 dynamics and enables classification of samples given the age of mice. Accordingly, we adapted the TockyConvNet model to classify spleen and thymus samples by including age as an input for model training (**Figure 7a**). This model effectively learnt the training dataset using three-fold cross-validation and showed high performance metrics including Area under curve of ROC 0.9 and Average Precision 0.95 (**Figure 7b**). Furthermore, by transferring the learned parameters from all layers and

removing the softmax activation from the final dense layer, we constructed a continuous

score model that captures the smooth and continuous dynamics of Foxp3 transcription

(Thymus-Spleen model score, **Figure 7c**).

The Thymus-Spleen model score data fit better with a quadratic regression using logged

age rather than raw age (**Figure 7c**), suggesting that changes are more pronounced in

early developmental stages and diminish in adult mice. Interestingly, splenic T cells from

neonatal mice, particularly at 3–4 days postpartum, showed high Thymus-Spleen model

scores, comparable to those of adult thymic T cells, indicating that their Timer profiles

closely resemble those found in thymic T cells. In contrast, thymic T cells from aged

mice, especially those older than 10 weeks, demonstrated Timer profiles more

characteristic of spleen-like T cells (**Figure 7c).** These dynamics, reflecting gradual

changes observed in two-dimensional plots of the raw data (**Figure 6)**, illustrate the

model's capability to capture the spectrum of changes from neonatal stages through to

and ageing process.

Optimising and Benchmarking of the TockyConvNet Four-Class Classifier

The successful development of the TockyConvNet classifier and the continuous model

scoring system demonstrates that specific patterns of Timer fluorescence dynamics are

associated with Foxp3 transcriptional dynamics in both the spleen and thymus, adjusted

for the age of mice. This is particularly noteworthy as the thymus in adult and aged mice,

typically over 6-7 weeks old [46], may include peripheral T cells that have recirculated into

the organ. Despite this, the model has successfully classified aged thymus samples

correctly, confirming that T cells within the thymus of aged mice exhibit distinctive Foxp3

transcriptional dynamics, distinct from both young thymic T cells and aged splenic T

cells. To comprehensively capture the compositions and real-time transcriptional

dynamics across the lifetime of mice in these two major immunological organs—and to

fully utilize this resource to benchmark the TockyConvNet approach—we extended the

TockyConvNet approach from two-class classifier into four-class-classifier categorizing

samples by both organ type (thymus and spleen) and age (young, <30 days postpartum;

aged, ≥30 days), aligned with standard definition of young adult thymus [47].

Consequently, the benchmarking dataset has been biologically optimized and exhibits

balanced class distribution for ML modeling and analysis.

We assessed three distinct ConvNet architectures varying in the number of convolutional

layer blocks (one, two, or three), resulting in the development of Conv1-Layer, Conv2-

Layers, and Conv3-Layers TockyConvNet 4-Classifiers, respectively. Each convolutional

layer block consists of a 3x3 convolutional layer followed by a 1x1 convolutional layer

serving as a spatial attention mechanism, similar to the structure used in the

TockyConvNet model for CNS2 KO Foxp3 Tocky data (Materials and Methods section).

Among these, the Conv3-Layers model (**Figure 8a**) demonstrated superior performance,

as evidenced by ROC and Precision-Recall analyses (**Figure 8b**). While the Conv2-

Layers model exhibited comparable, albeit slightly reduced performance, the Conv1-

Layer model showed a marked decrease in effectiveness.

Next, we examined the impact of data preprocessing on model performance. Converting

Timer Fluorescence into Timer Angle and Intensity was crucial as raw fluorescence data

was challenging for the ConvNet to process, leading to suboptimal performance.

Additionally, experimenting with different data resolutions, we found that neither low (25

$\times$ 25) nor high (400 $\times$ 400) resolution grids were effective. This suggests that a resolution

of 100 x 100 strikes the optimal balance by maintaining sufficient detail for accurate

feature extraction and ensuring a good density of cells across pixels, which is critical for

reducing meaningless variability in the data.

Finally, we benchmarked the TockyConvNet models against conventional manual gating

methods, specifically polygon and quadrant gates, which have been widely utilised in

prior studies using Fluorescent Timer proteins (**Figure 8c**) [48] [49]. These traditional

methods yielded satisfactory results for categorising Thymus-Young and Spleen-Aged

samples, yet they faltered in accurately classifying the other two classes. Although the

TockyKmeansRF method demonstrated notable efficiency with Thymus-Young and

Spleen-Aged samples, it was less effective for Spleen-Young. Notably, TockyConvNet

consistently outperformed all the other methods in both ROC and Precision-Recall

analyses (**Figure 8b**), establishing the TockyConvNet approach as a robust classifier

using image-converted Timer data.

Optimising Grad-CAM Method for Analysing Foxp3 Timer Dynamics

To further explore and develop the Grad-CAM method for flow cytometric Timer data, we

analysed each convolutional layer's output by Grad-CAM using the optimised

TockyConvNet model (**Figure 9**). Heatmaps of the Grad-CAM outputs illustrate these

distinctions, showing the progression of feature capture across layers (**Figure 9a**). ROC

analysis identified the efficacy of each convolutional layer in differentiating the four

classes. The Thymus-Young and Spleen-Aged classes were distinctly recognised

throughout the layers, while and Thymus-Aged and Spleen-Aged classes were the most

distinctly recognised in the Conv2 layer (**Figure 9b**).

Visualisation of the most informative convolutional layers in both Timer Angle and

Intensity space, as well as the original Timer fluorescence space, provided further

insights into important features in the transcriptional dynamics of different classes with

single-cell granularity (**Figure 9c**). Notably, Grad-CAM high cells in the Spleen-Aged

class were predominantly located on the lower edge within the Timer Blue-Red space,

with high Timer Intensity within the high 80˚–90˚ Timer Angle range (**Figure 9c**),

mirroring the marked accumulation of cells within Blue$^-$Red$^+$ (**Figure 6e**). This suggests

their substantially attenuated and infrequent Foxp3 transcription after culminating Foxp3

proteins. In contrast, Grad-CAM high cells in the Spleen-Young class included both new

Timer expression (Timer Blue$^+$Red$^-$, Angle = 0 ˚) and low Timer Intensity within the ~90˚

Timer Angle arrested class, suggesting that some spleen cells newly and moderately

express Timer protein but may rapidly transition to arrested transcription states without

sustained Foxp3 transcription. Meanwhile, the Thymus classes are both characterised

by newly induced Foxp3 transcription, with Thymus-Young cells predominantly featured

by remarkably high new transcription to intermediate Timer Angles with high Timer

Intensity (**Figure 9c**). Thymus-Aged cells are characterised by both new expression and

high Timer Angle cells with high Timer Intensity, suggesting their ability to activate Foxp3

transcription as well as the accumulation of spleen-like cells, presumably due to the

recirculation of peripheral T cells in aged mice [46].

Runtime and Memory Usage for TockyConvNet model learning

Lastly, the computational efficiency of TockyConvNet model learning was evaluated by

analysing the runtime and memory usage. For the CNS2 KO Foxp3 Tocky data,

processed with the established Conv 2-Layer model, the runtime ranged from

approximately 1 to 6 seconds, and memory usage spanned from about 2 GB using

sample sizes between 52 and 416. Similarly, using the Developmental and Ageing WT

Foxp3 Tocky dataset to train the Conv 3-Layers model, runtime and memory usage also

increased with larger sample sizes, ranging from around 10 to 70 seconds for runtime

and significantly higher memory usage from approximately 1.5 to 3 GB. Together with

1   the model complexity information, the analysis highlights the scalability of the

2   TockyConvNet model, albeit with increasing computational demands for larger datasets.

3
4
5
6

Discussion

This study establishes a proof-of-concept for dissecting physiological transcriptional

dynamics at the single-cell level, introducing two principal ML methods—

TockyKmeansRF and TockyConvNet—that represent a significant departure from

traditional manual gating approaches. While conventional gating typically confirms

known cell populations using fixed strategies, our ML approaches dynamically identify

feature cells, which represent group-specific effects crucial for classification, based

solely on the patterns of flow cytometric Timer data. TockyKmeansRF combines

clustering and RF algorithms for feature-based cell identification but remains inherently

cluster-focused. In contrast, TockyConvNet employs ConvNet to achieve single-cell

resolution and uses Grad-CAM to provide visually intuitive, quantitative insights.

Surpassing cluster-level outputs, TockyConvNet captures the finest details of Timer

dynamics, enabling the precise identification of CNS2-dependent cells, as well as

developmentally or ageing-specific transcriptional features of Foxp3 at the single-cell

level. Benchmarking through two independent experiments robustly validates the

performance and generalizability of TockyConvNet. Although TockyConvNet utilizes

relatively compact ConvNet models to reduce the risk of overfitting, the extensive

parameters within ConvNets still pose a risk, which can be mitigated by employing

shallow learning strategies and generating high-quality training data. A summary of each

method's strengths and limitations is presented.

The Tocky ML approaches developed here are anticipated to be broadly applicable to

genes beyond Foxp3 by effectively analyzing Fluorescent Timer fluorescence data,

facilitated by the normalisation and data transformation methods provided by TockyPrep

[50]. By removing the effects of autofluorescence, these methods enable the effective

application of the ML methods across various Fluorescent Timer systems. However, it is

important to acknowledge that adjustments in model architecture and training methods

may be necessary to accommodate different data structures, as broadly recognised

within ML communities [51]. Additionally, the image conversion techniques employed in

our approaches could be effectively adapted for other transcriptional reporter systems,

such as EGFP, when combined with an appropriate temporal marker. This adaptation

would allow the use of the TockyConvNet algorithms, which can provide a more

nuanced understanding of dynamic transcriptional events. Nevertheless, such

adaptations would likely require extensive experimentation and optimisation to tailor the

approach to specific requirements of individual cases.

ConvNet models jointly learn features across all layers, with no single layer solely

responsible for the model's learning [51]. Originally established for image classification

with photographic images, Grad-CAM typically utilizes the last convolutional layer, which

captures abstract and semantic meanings [40]. However, our application of Grad-CAM to

flow cytometric Timer data suggests that it may be necessary to investigate each

convolutional layer to determine the most effective one for summarizing the features of

group-specific cells. The areas highlighted by Grad-CAM likely represent the most

distinct features of cells within a given class, associating high Grad-CAM values with

class-specific transcriptional dynamics. This approach opens new avenues for further

investigating transcriptional dynamics using Grad-CAM outputs in future studies using

Foxp3 Tocky and other Fluorescent Timer reporter systems.

Biologically, by developing cutting-edge technologies such as CRISPR-based

modification of only the reporter, Timer-based single-cell resolution, and machine-

learning-driven data analysis, we demonstrate uniquely high-frequency Foxp3

transcriptional dynamics under functioning CNS2. This is evidenced by cells displaying

low Timer Angles (~40–80°) with high Timer Intensities, which indicate persistent, high-

frequency transcription [7]. This discovery challenges the traditional view that CNS2's

primary role is merely to maintain Foxp3 expression post-cell division. Previous studies,

relying on endpoint measurements of bulk populations from endogenous CNS2 KO mice

[16, 20, 21], were unable to differentiate between direct CNS2 activity and secondary effects

from reduced Foxp3 protein. Although a previous report hinted the link between low

Foxp3 expression and CNS2 activity by demonstrating loss of Foxp3 expression in

relative lower Foxp3 expressor in CNS2 KO [21], all preceding studies failed to pinpoint

specific cells where CNS2 was actively functioning. In contrast, our use of both WT and

CNS2 KO Foxp3 Tocky mice, which retain the intact endogenous Foxp3 gene, has

allowed us to identify specific CNS2-functioning cells and analyse how functioning CNS2

directly controls Foxp3 transcription dynamics, addressing a key aspect of Foxp3

autoregulation [9]. The integration of the innovative CRISPR strategy with sophisticated

ML methods, through comparative analysis of CNS2 WT and KO features, confidently

identify CNS2 functioning and independent cells from WT Foxp3 Tocky, revealing that

intact CNS2 regulates the frequency of Foxp3 transcription bursts within specific CNS2-

functioning cells. This paradigm shift, unachievable with traditional methods, is robustly

supported by our high-performance ML models, unravelling a previously unrecognised

temporal dimension of CNS2 function at the single-cell level.

To further obtain insights into the discovery in the temporal dimension of Foxp3

transcription, our ML-driven analysis using TockyConvNet revealed a nuanced

transcriptional signature of CNS2-dependent cells, characterized by elevated IL-2R

expression, repressed Stat5, and selective modulation of TCR signalling components—

including increased Nfatc1/2, decreased Egr1/3, and relatively low or absent induction of

Nr4a and NF-kB (identified as the 'B2' fraction in **Figure 5**). This pattern indicates that

CNS2 operates under finely tuned TCR and IL-2 signal dynamics. Such transient

signalling might involve *periodic* IL-2 and TCR inputs. Consistent with our previous work

employing Nr4a3 Tocky, which indicated *periodic* and *brief* TCR signals approximately

every week in Foxp3-expressing cells [7], these findings imply CNS2-mediated Foxp3

transcription is regulated through intermittent and short signals rather than constant

stimulation. Similarly, in vivo supply of IL-2 signalling is likely sparse in the body as each

T cell can produce it for a short time only [52]. Thus, although it is well known that IL-2

enhances Foxp3 expression, its physiological transcription via endogenous IL-2 remains

incompletely understood. With our compelling evidence and the established Tocky and

ML tools, future studies on Foxp3 transcriptional regulation can focus on identifying the

finely tuned signalling milieu involving antigen stimulation (TCR) and IL-2 supply and

signalling.

Integrating the findings, we propose a model where CNS2 controls the frequency of

Foxp3 transcriptional activity through integrating transient and periodic signals from IL-2

and TCR inputs, inducing transient bursts of transcription under certain restricted

situations. Future studies can build on our experimental and ML tools to further explore

how the frequency Foxp3 transcription is controlled by active CNS2 – for example, this

may involve integrated analysis of single-cell level methylation for CpGs in CNS2, and/or

single-cell level chromatin configuration analysis, combined with our Tocky and ML

technologies. Doing so, fragmented pieces of evidence regarding Foxp3 transcriptional

regulation may well be integrated into a coherent and dynamic perspective. For

example, CNS2 itself is regulated by Foxp3 protein [9, 13, 20] and downstream factors from

IL-2 signalling (Stat5) and TCR signalling pathways (including NFAT [16], CREB [17] and

Ets-1 [18, 19]). Supporting this, our flow cytometric analysis shows upregulation of PD-1 and

CD25 specifically in CNS2-dependent cells. CD25 induction results from TCR and/or IL-

2 signalling [53] and is further amplified by Foxp3 [14], whereas PD-1 induction via TCR

serves as negative feedback [54].

Further, our study has established that Foxp3 transcription is dynamically modulated by

ageing, producing benchmark data for the developmental and ageing dynamics of Foxp3

transcription. In addition to the generation of the resource data, through TockyConvNet

and Grad-CAM, we uncovered substantial age-dependent variations in Foxp3

transcriptional dynamics across the two major immunological organs, the spleen and the

thymus.

Younger thymic T cells exhibit strong induction of new Foxp3 expression, the dynamics

and trajectories of which have been captured by Grad-CAM (**Figure 9c**). Intriguingly, the

thymic pattern of Foxp3 transcription dynamics, as captured by the TockyConvNet

continuous model, peaks at days 3-4 post-birth (**Figure 7c**). This observation resonates

with the uniqueness of the thymic environment in the first days of life in mice, which

potentially influences negative selection processes [55].

In contrast, older thymic T cells, though still capable of initiating new Foxp3 expression,

primarily accumulate cells with arrested transcription, which likely represents recirculated

aged peripheral Foxp3$^+$ T cells [46]. Recirculation of peripheral Foxp3$^+$ T cells into the

thymus typically starts only after 6 to 7 weeks post-birth [46], and thus, the cut-off of young

and aged organs at 30 days old in our study effectively eliminate the contribution of

recirculating T cells in young thymus. An intriguing future direction is to distinguish the

dynamics of Foxp3 transcription between newly generated thymic T cells and peripheral

T cells that have recirculated into the thymus in adult and aged mice, thereby addressing

the possible impact of the thymic environment on recirculating T cells. Currently, the

experimental identification of recirculating T cells is challenging due to limitations in

available methods, even with a reporter mouse strain such as Rag2p-EGFP. These

methods identify recirculating T cells as EGFP(-) cells, but critically, the expression of

EGFP can be prematurely lost in newly generated thymic T cells following intrathymic

cell divisions before emigration [56, 57]. Therefore, future studies would benefit from

combining our integrated approach involving Tocky and ML. Specifically, the

development of a novel Rag2 Tocky strain, which reports Rag2 gene activity with the

Fluorescent Timer protein, is a promising target. The integrated ML analysis of Timer

Blue with Timer Red using our methods should comprehensively reveal the dynamics of

recently generated thymic T cells as well as recirculating T cells, thus overcoming the

limitations of Rag2p-EGFP. This concept for Rag2 Tocky was initially proposed in our

earlier work on Tocky [7] and the current study has renewed the relevance of this

promising avenue.

In addition, intriguingly, we discovered that neonatal splenic T cells (3–4 days post-birth)

actively initiate new Foxp3 expression, making their Foxp3 Timer profiles resembling to

adult thymic T cells, as demonstrated by the continuous score model (**Figure 7c**). Grad-

CAM analysis revealed low Timer Intensity in Foxp3$^+$ splenic T cells showing a mix of

new and arrested transcription states in young mice. This means that some cells initiate

Foxp3 transcription with relatively low activity, while others terminate Foxp3 transcription

before accumulating substantial Timer proteins. This pattern suggests that new Foxp3

induction in splenic CD4$^+$ T cells in young neonates frequently leads to premature

termination. Traditionally, these initiating cells may be viewed as peripherally-induced

Tregs, or pTregs [58], and the terminating ones as ex-Tregs [59]. However, such

classifications do not capture the nuanced continuous dynamics of Foxp3 transcription

our study has revealed. Future research should conduct quantitative, temporal analyses

of Foxp3 transcriptional activities using data-driven methods, further refining the ML

approaches introduced in this study.

Furthermore, the progressive decline of Foxp3 transcriptional dynamics in splenic T cells

with ageing, as illustrated by the TockyConvNet continuous score (**Figure 7c**),

demonstrates how ageing progressively modifies the temporal dynamics of Foxp3

transcription in peripheral T cells, providing insights into T cell senescence. The arrested

Foxp3 transcription in aged splenic T cells, characterized by high Timer Angle and low

Timer Intensity (**Figure 9c**), indicates that transcriptional activities become infrequent

and weak as mice age. Although previous studies described an increased Foxp3[+] T cells

in the periphery of aged mice [25, 60], the biological significance of this increase remains

unclear. The progressive increase in the percentage of Foxp3 Timer[+] cells (**Figure 6e**)

alongside the steady decline of Foxp3 transcriptional frequency in its homeostatic

regulation indicates that splenic Foxp3 transcriptional dynamics are shaped by

progressive changes induced by ageing, affecting both the quantity and composition of

Foxp3[+] T cell fraction. These changes are thus likely driven by ageing-related epigenetic

modifications such as global shifts in methylation patterns [61] and targeted methylation in

key genes [62]. These modifications may directly impact Foxp3 transcription or indirectly

influence it through signalling molecules and transcription factors involved in Foxp3

transcription, such as TCR, IL-2, and TGF-β signalling downstream [13].

In summary, our study demonstrates how intact CNS2 controls Foxp3 transcription in the

presence of undisturbed Foxp3 protein, addressing a key aspect of Foxp3

autoregulation [9]. It also reveals age- and development-dependent changes in Foxp3

transcription within both the thymus and periphery. These findings are validated by

quantitative model performance metrics, ensuring transparency and reproducibility. Our

study thus establishes a proof-of-concept for modern data-oriented methodologies that

integrate ML with CRISPR and Tocky to unravel dynamic gene regulation at single-cell

resolution, which would not be possible by existing methods. This approach overcomes

the limitations of traditional immunological research practices, particularly the narrative-

driven interpretations of endpoint KO analysis and manual gating. The ambiguity and

subjectivity inherent in these traditional practices likely contribute to the ongoing

reproducibility crisis in scientific research, especially in immunology and flow cytometric

analysis [27, 63].

Materials and Methods

CRISPR-mediated mutagenesis of Foxp3-Tocky

The Foxp3-Tocky mouse strain (BAC Tg (*Foxp3^ΔExon3 FastFT*)) was generated by the Ono

group and reported previously [7]. The CRISPR-mediated mutagenesis of the Foxp3-

Timer transgene in Foxp3-Tocky mice was performed using the previously reported

CRISPR transgenic approach [64, 65]. Briefly, fertilised eggs were obtained from Foxp3-

Tocky and were subjected to electroporation to introduce the Cas9 protein (317-08441,

Nippon Gene, Toyama, Japan), tracrRNA (GE-002, FASMAC, Kanagawa, Japan),

synthetic crRNAs, and a single-stranded oligodeoxynucleotide (ssODN). The CRISPR

strategy was designed to replace the CNS2 sequence with a short insert sequence,

AAGTTTAAAC. Accordingly, the following synthetic crRNAs were used: (1) 5' crRNA

targeting CCTGAGCTCCATTATGACAG (PAM: AGG) and (2) 3' crRNA targeting

AGTTCCACAAGTATTTAAGG (PAM: AGG). The ssODN designed for homology-

directed repair and the insertion of the short nucleotide sequence

AAGTTTAAAC/GTTTAAACTTC was 5'-

CTCTTTATGTTTGGTCAGAACTTATAAGAAATCTCCTCCT**GTTTAAACTTC**AGAGGAT

TGGAAAACCCTCTACTGTCCTGATCTGGGGTC-3'. All animal experiments were

approved by the Animal Welfare and Ethical Review Body at Imperial College London

and the Animal Experiment Committee at Kumamoto University.

ChIP-seq Analysis

ChIP-seq data analysis was performed using the NCBI SRA dataset DRP003376 [66].

Briefly, HISAT2 [67] was used to align sequence reads to the mouse genome (*mm10*),

followed by the data processing using samtools [68] to produce sorted bam files. The

sequence peaks were visualised by Integrative Genomics Viewer [69].

Timer Data Preprocessing

Methods for Timer fluorescence data pre-processing, implemented as the R package

TockyPrep [50], were reported previously [7]. Briefly, immature blue and mature red

fluorescence data are normalised to treat both types of fluorescence equally and

thresholded to remove autofluorescence. This is followed by a trigonometric

transformation of the normalised data, converting Timer fluorescence into polar

coordinates, specifically Timer-Angle and Timer-Intensity.

TockyKmeansRF Implementation

TockyKmeansRF is implemented within the R *TockyRandomForest* package, part of the

TockyMachineLearning package suite. It integrates k-means clustering [70] and RF

classification by *randomForest* [71] utilising the Tocky data preprocessing method

TockyPrep [50]. Briefly, pre-processed training and test datasets undergo k-means

clustering separately. The default setting for k-means is to use the number of random

starts at 1 with the maximum number of iterations set to 10. After k-means clustering, the

percentage of cells within each cluster is computed per sample, forming a table of

cluster percentages for each dataset.

To match the clusters between the training and testing datasets, first, Timer Angle and

Intensity data are standardised. Next, each cluster's representative point is calculated

as:

$$\mathbf{m}_i = \begin{pmatrix} \text{median}(\text{angle}_i) \\ \text{median}(\text{intensity}_i) \end{pmatrix}$$

2   where i indexes the cluster.

3   Using the representative points $\mathbf{m_{train}}$ for the training clusters and $\mathbf{m_{test}}$ for the test

4   clusters respectively, the distance matrix **D** is computed using the Euclidean norm:

$$D_{ij} = \|\mathbf{m}_{\text{train},i} - \mathbf{m}_{\text{test},j}\|$$

where $\mathbf{m_{train,\,i}}$ and $\mathbf{m_{test,\,j}}$ are the representative vectors of the i-th training cluster and the

j-th testing cluster, respectively.

For the optimal assignment of clusters, we address the linear sum assignment problem

(LSAP) using the Hungarian method [72]. This method systematically identifies a

permutation of the test set cluster indices that minimizes the total Euclidean distance

between the corresponding clusters from the training and test datasets. Specifically, the

LSAP seeks a permutation matrix P that reorders the columns of the distance matrix

such that the sum of the diagonal elements of **DP** is minimised.

$$\min_{P} \text{trace}(DP),$$

where **D** is the distance matrix, and **P** is a permutation matrix that reorders the columns

of **D** to optimally align each test cluster with a training cluster. The *solve_LSAP* function

from the R package *clue* is utilised for this purpose [72].

A RF model is then trained using the training data as a cluster percentage table, and the

mean decrease Gini (MDG) index of the RF model is used to assess the importance of

each cluster. The R package randomForest [71] is used in the implementation. The test

data as a cluster percentage table with the matched clusters is used to predict sample

identity, generating model performance metrics. The MDG index is then used to

calculate the importance score of individual cells within the test dataset. Subsequently,

feature cells are identified using a threshold value and are classified into clusters, where

appropriate, using a *density-based spatial clustering of applications with noise* the R

package *dbscan* [73].

Image Conversion Techniques for TockyConvNet

The image conversion and reversion techniques are implemented as integral

components of the TockyConvNetR package, part of the *TockyMachineLearning*

package suite. Following data preprocessing, the conversion of Timer Angle and

Intensity data into image data was performed as follows: First, global minimum and

maximum values for the angle and intensity measurements were computed. These

extremes were used to establish uniform bin edges across all datasets, typically defining

100 equidistant intervals spanning from the global minimum to the global maximum for

each variable. The data were then transformed into pixel-style representations by

mapping the flow cytometric measurements into a 2D matrix. Each matrix element

represents the count of data points within each bin, effectively converting the flow

cytometric data into a 100×100 pixel image, where each pixel corresponds to a bin

count. This method provides a consistent and precise pixelated representation of the

flow cytometric data for each sample. To revert image data back to original Timer

fluorescence data, parameters defining the pixels, such as global minimum and

maximum values along with bin widths, are used.

ConvNet Architectures and Model Training for TockyConvNet

TensorFlow and Keras [74] were used to construct ConvNet models for TockyConvNet and implemented as the python TockyConvNetPy package, part of the *TockyMachineLearning* package series. The input layer accepts batch image data of size 100×100 pixels with a single channel representing cell density, as pre-processed by TockyConvNetR as detailed above.

In our convolutional network architecture, each convolutional operation with a 3×3 kernel with a ReLU activation is immediately followed by a sigmoid-activated pointwise convolution, which serves as a simple attention block within the network [75]. The primary purpose of this layer is to generate an attention map A:

$$A = \sigma(\mathrm{Conv2D}(X, (1,1)))$$

where σ represents the sigmoid activation function, and X denotes the input to the attention layer. The resulting attention map A, which has a single channel with spatial dimensions equal to X, is broadcast across all channels of the feature map X to match its dimensions. The broadcasted attention map A' is then used for element-wise multiplication with X to modulate the feature response:

$$Y = X \odot A'$$

where Y is the output feature map after applying the attention, $\odot$ denotes element-wise multiplication, and A' is the broadcasted version of A. Thus, it applies an element-wise multiplication to the preceding feature map, enhancing or suppressing features based on their spatial importance. A subsequent max pooling layer reduces spatial dimensions, followed by a dropout layer. This configuration, including a convolutional layer, an attention block, max pooling, and dropout, is repeated as indicated.

Finally, the convolutional layers feed into a flatten layer that transitions the data. For classification, this is followed by a fully connected dense layer with ReLU activation and a final dropout at the rate of 0.5 before reaching the softmax output layer. For regression, the flattened data is processed through a sigmoid-activated dense layer to generate a continuous score.

For the TockyConvNet application to Foxp3 Timer Neonatal-to-Ageing Data, the ConvNet model was configured to also receive %total Timer$^+$ cells as numeric data. This inclusion ensures a fair and accurate comparison between TockyConvNet and traditional gating methods, which similarly analyse the percentage of cells within the parent population.

Model training employed the Adam optimizer and used a three-fold cross-validation approach. For each fold, models were compiled using either binary cross-entropy or categorical cross-entropy as the loss function, with accuracy as the evaluation metric. Training was typically conducted over 10 to 12 epochs with batch sizes ranging from 4 to 8, and learning rates set between 0.001 and 0.002. Validation data specific to each fold were used to monitor and assess model performance throughout the training process.

Grad-CAM

The Grad-CAM methodology [40] with an aggregation method was used to identify the regions of input images that most strongly influence the prediction of ConvNet for a chosen target class. We used the prediction score y_c for class c, which is the logit score prior to the softmax function, and each of available convolutional layers whose feature maps A_k are denoted by k. Using TensorFlow's gradient tape function, we computed the gradient of the class score y_c with respect to the feature map activations:

$$\frac{\partial y_c}{\partial A_{ij}^k}$$

Here, i and j indicate the i-th and the j-th pixel within the k-th feature map. Next, all the

feature matrices were pixel-wise aggregated by global average pooling over i,j. First, the

scalar weight α_c^k, as a weighted factor for the k-th feature map, was obtained:

$$\alpha_c^k = \frac{1}{Z} \sum_i \sum_j \frac{\partial y_c}{\partial A_{ij}^k}$$

Note that Z is the total number of pixels in the feature map (for instance, 10,000 for the

resolution 100×100). Subsequently, the ReLU was applied to retain the feature

contributions that positively influence y_c:

$$L_{ij}^{\text{Grad-CAM}(c)} = \text{ReLU}\left(\sum_k \alpha_c^k A_{ij}^k \right)$$

Next, the outputs from all the samples (N_c samples) belonging to the class c will be

summed and aggregated.

$$\bar{L}_{ij}^{\text{Grad-CAM}(c)} = \sum_{n=1}^{N_c} L_{ij}^{\text{Grad-CAM}(c,n)}$$

The heatmap is then normalised to have the range of [0,1]. After generating individual

heatmaps for each image in the KO and WT groups, the heatmaps were summed within

each group to aggregate activation signals across multiple images. Each aggregated

heatmap then underwent Gaussian smoothing to suppress noise and enhance the

visibility of predominant activation patterns. These smoothed and aggregated heatmaps

were used for the analysis of the ontogeny data. For the analysis of CNS2 KO data, a

differential matrix was calculated by subtracting the KO group's Grad-CAM heatmap

from the WT group's Grad-CAM heatmap. To implement Grad-CAM, we developed a

new gradient model for each convolutional layer by retrieving the layer outputs and

utilizing the *tf.GradientTape* function in TensorFlow (version 2.10.0) [76] on Jupyter Lab

(3.6.3, https://jupyter.org/).

RNA-seq Data Analysis

The RNA-seq data analyzed in Figure 5 was previously reported [7] and is deposited to NCBI GEO with the accession number GSE89481. The R package DESeq2 [77] was used to perform a Wald test and p-value adjustment with the Benjamini & Hochberg method. In the cross-dataset analysis of flow cytometric Timer data presented in Figure 5, WT Feature cells were identified as the top 90th percentile cells based on Attention-Conv2 Grad-CAM scores. KO Feature cells were identified as the bottom 25th percentile cells based on Conv2 Grad-CAM scores. Both sets of Grad-CAM scores were derived from the model trained in Figure 4.

Logistic Regression Model and Other Statistical Methods

To quantitatively evaluate the efficacy of manual gating methods, a logistic regression model was developed to predict group classifications based on variables derived from manual gating techniques using the glm function from the R package stats [70], specifying a binomial family. The model was trained on a training dataset and validated against an independent test dataset. Predicted probabilities were converted to class labels using a threshold of 0.5. The model's performance was visualised using the R package pROC [78].

Cluster percentage data were analysed by a Kruskal test followed by post-hoc Dunn's test using the R package *dunn.test* with p-value adjustment using Bonferroni's correction. Pair-wise comparisons of percentage data were conducted using the Mann-Whitney U test. Mean Fluorescence Intensity (MFI) data were obtained from logarithmically transformed values to achieve normality, and the Student's t-test was applied. Where required, p-values obtained from these tests were adjusted for multiple comparisons using the Benjamini-Hochberg method.

Computational Performance

The computational performance was assessed using a system with an Apple M2 Ultra processor, 128 GB of memory, and running macOS Sonoma 14.7.1. Runtime was assessed using the *system.time* function in R or the time package in python. The memory usage during the execution was monitored using the R package *profmem* and Python's *memory_profiler*.

Acknowledgements: MO was supported by a CRUK Programme Foundation Award (DCRPGF\100007) and the MRC grant (MR/S000208/1). This research was also supported by KAKENHI research grants from the Japan Society for the Promotion of Science (JSPS) (JP21K07082, JP21H00433, and JP24K10259 to MO), JSPS Core-to-core program to MO and YS (JPJSCCA2020008 to MO and YS), and Japan Agency for Medical Research and Development (24gm1810001s0103 to MO), Advanced Animal Model Support (AdAMS) (JP16H0627601 and JP22H04922 to KA).

Author Contributions: MO conceived the study and all computational and ML methodologies, designed model architectures, wrote computational codes, and performed all computational analysis. NI, YS, KA, and MO designed experiments. MO and KA designed the CRISPR mutagenesis. NT and KA performed transgenesis. NI and MO established the CRISPR mutant colony and performed all flow cytometric experiments. YS, KA, and MO secured funding. MO wrote the manuscript.

Conflicts of Interest: A patent associated with the ML method in this study has been filed.

Data Availability: The TockyMachineLearning toolkit developed in this study is available on GitHub at https://MonoTockyLab.github.io/TockyMachineLearning.

References

1. Petropoulos, S. *et al.* Single-Cell RNA-Seq Reveals Lineage and X Chromosome Dynamics in Human Preimplantation Embryos. *Cell* **165**, 1012-1026 (2016).

2. Treutlein, B. *et al.* Dissecting direct reprogramming from fibroblast to neuron using single-cell RNA-seq. *Nature* **534**, 391-395 (2016).

3. Park, J.E. *et al.* A cell atlas of human thymic development defines T cell repertoire formation. *Science* **367** (2020).

4. Bradley, A., Hashimoto, T. & Ono, M. Elucidating T cell activation-dependent mechanisms for bifurcation of regulatory and effector T cell differentiation by multidimensional and single-cell analysis. *Frontiers in immunology* **9**, 1444 (2018).

5. Tan, B.J.Y. *et al.* HTLV-1 infection promotes excessive T cell activation and transformation into adult T cell leukemia/lymphoma. *The Journal of Clinical Investigation* **131** (2021).

6. Barry, J.D., Donà, Erika, Gilmour, D. & Huber, W. TimerQuant: a modelling approach to tandem fluorescent timer design and data interpretation for measuring protein turnover in embryos. *Development* **143**, 174-179 (2016).

7. Bending, D. *et al.* A timer for analyzing temporally dynamic changes in transcription during differentiation in vivo. *Journal of Cell Biology* **217**, 2931-2950 (2018).

8. Subach, F.V. *et al.* Monomeric fluorescent timers that change color from blue to red report on cellular trafficking. *Nat Chem Biol* **5**, 118-126 (2009).

9. Bending, D. *et al.* A temporally dynamic Foxp3 autoregulatory transcriptional circuit controls the effector Treg programme. *The EMBO journal* **37**, e99013 (2018).

10. Bailey-Bucktrout, S.L. & Bluestone, J.A. Regulatory T cells: stability revisited. *Trends Immunol* **32**, 301-306 (2011).

11. Overacre, A.E. & Vignali, D.A.A. Treg stability: to be or not to be. *Current Opinion in Immunology* **39**, 39-43 (2016).

12. Barbi, J., Pardoll, D. & Pan, F. Treg functional stability and its responsiveness to the microenvironment. *Immunol Rev* **259**, 115-139 (2014).

13. Ono, M. Control of regulatory T‐cell differentiation and function by T‐cell receptor signalling and Foxp3 transcription factor complexes. *Immunology* **160**, 24-37 (2020).

14. Ono, M. *et al.* Foxp3 controls regulatory T-cell function by interacting with AML1/Runx1. *Nature* **446**, 685-689 (2007).

15. Kitoh, A. *et al.* Indispensable role of the Runx1-Cbfβ transcription complex for in vivo-suppressive function of FoxP3+ regulatory T cells. *Immunity* **31**, 609-620 (2009).

16. Li, X., Liang, Y., LeBlanc, M., Benner, C. & Zheng, Y. Function of a Foxp3 cis-element in protecting regulatory T cell identity. *Cell* **158**, 734-748 (2014).

17. Kim, H.-P. & Leonard, W.J. CREB/ATF-dependent T cell receptor–induced FoxP3 gene expression: a role for DNA methylation. *Journal of Experimental Medicine* **204**, 1543-1551 (2007).

18. Mouly, E. *et al.* The Ets-1 transcription factor controls the development and function of natural regulatory T cells. *Journal of Experimental Medicine* **207**, 2113-2125 (2010).

19. Polansky, J.K. *et al.* Methylation matters: binding of Ets-1 to the demethylated Foxp3 gene contributes to the stabilization of Foxp3 expression in regulatory T cells. *Journal of Molecular Medicine* **88**, 1029-1040 (2010).

20. Zheng, Y. *et al.* Role of conserved non-coding DNA elements in the Foxp3 gene in regulatory T-cell fate. *Nature* **463**, 808-812 (2010).

21. Feng, Y. *et al.* Control of the inheritance of regulatory T cell identity by a cis element in the Foxp3 locus. *Cell* **158**, 749-763 (2014).

22. Fontenot, J.D., Dooley, J.L., Farr, A.G. & Rudensky, A.Y. Developmental regulation of Foxp3 expression during ontogeny. *J Exp Med* **202**, 901-906 (2005).

23. Ono, M. & Tanaka, R.J. Controversies concerning thymus‐derived regulatory T cells: fundamental issues and a new perspective. *Immunology and cell biology* **94**, 3-10 (2016).

24. Garg, S.K. *et al.* Aging is associated with increased regulatory T-cell function. *Aging Cell* **13**, 441-448 (2014).

25. Lages, C.S. *et al.* Functional Regulatory T Cells Accumulate in Aged Hosts and Promote Chronic Infectious Disease Reactivation1. *The Journal of Immunology* **181**, 1835-1848 (2008).

26. Sharma, S., Dominguez, A.L. & Lustgarten, J. High Accumulation of T Regulatory Cells Prevents the Activation of Immune Responses in Aged Animals1. *The Journal of Immunology* **177**, 8348-8355 (2006).

27. Brinkman, R.R. Improving the Rigor and Reproducibility of Flow Cytometry-Based Clinical Research and Trials Through Automated Data Analysis. *Cytometry A* **97**, 107-112 (2020).

28. Ono, M. Unraveling T-cell dynamics using fluorescent timer: Insights from the Tocky system. *Biophys Physicobiol* **21**, e211010 (2024).

29. Ono, M., Tanaka, R.J. & Kano, M. Visualisation of the T cell differentiation programme by Canonical Correspondence Analysis of transcriptomes. *BMC genomics* **15**, 1-15 (2014).

30. Libbrecht, M.W. & Noble, W.S. Machine learning applications in genetics and genomics. *Nature Reviews Genetics* **16**, 321-332 (2015).

31. Duetz, C. *et al.* Computational flow cytometry as a diagnostic tool in suspected-myelodysplastic syndromes. *Cytometry A* **99**, 814-824 (2021).

32. Arvaniti, E. & Claassen, M. Sensitive detection of rare disease-associated cell subsets via representation learning. *Nature Communications* **8**, 14825 (2017).

33. Hu, Z., Tang, A., Singh, J., Bhattacharya, S. & Butte, A.J. A robust and interpretable end-to-end deep learning model for cytometry data. *Proceedings of the National Academy of Sciences* **117**, 21373-21380 (2020).

34. Dinalankara, W., Ng, D.P., Marchionni, L. & Simonson, P.D. Comparison of three machine learning algorithms for classification of B-cell neoplasms using clinical flow cytometry data. *Cytometry Part B: Clinical Cytometry* **n/a**.

35. Botta, C. *et al.* FlowCT for the analysis of large immunophenotypic data sets and biomarker discovery in cancer immunology. *Blood Advances* **6**, 690-703 (2022).

36. Ono, M. & Crompton, T. A multidimensional toolkit for elucidating temporal trajectories in cell development in vivo. *Development* **151**, dev204255 (2024).

37. Aghaeepour, N. *et al.* Critical assessment of automated flow cytometry data analysis techniques. *Nat Methods* **10**, 228-238 (2013).

38. Fujii, H. *et al.* Regulatory T cells in melanoma revisited by a computational clustering of FOXP3+ T cell subpopulations. *The Journal of Immunology* **196**, 2885-2892 (2016).

39. Freedman, L.P., Cockburn, I.M. & Simcoe, T.S. The Economics of Reproducibility in Preclinical Research. *PLOS Biology* **13**, e1002165 (2015).

40. Selvaraju, R.R. *et al.* Grad-CAM: Visual Explanations from Deep Networks via Gradient-Based Localization. *International Journal of Computer Vision* **128**, 336–359 (2019).

41. Ono, M. TockyPrep: data preprocessing methods for flow cytometric fluorescent timer analysis. *BMC Bioinformatics* **26**, 44 (2025).

42. Van Keuren, M.L., Gavrilina, G.B., Filipiak, W.E., Zeidler, M.G. & Saunders, T.L. Generating transgenic mice from bacterial artificial chromosomes: transgenesis efficiency, integration and expression outcomes. *Transgenic Res* **18**, 769-785 (2009).

43. Schallenberg, S., Petzold, C., Tsai, P.Y., Sparwasser, T. & Kretschmer, K. Vagaries of fluorochrome reporter gene expression in Foxp3+ regulatory T cells. *PLoS One* **7**, e41971 (2012).

44. Ono, M. TockyLocus: Quantitative Analysis Methods for Flow Cytometric Fluorescent Timer Data. *arXiv [q-bio.QM]* (2024).

45. Neto, N.G.B. *et al.* Non-invasive classification of macrophage polarisation by 2P-FLIM and machine learning. *eLife* **11**, e77373 (2022).

46. Thiault, N. *et al.* Peripheral regulatory T lymphocytes recirculating to the thymus suppress the development of their precursors. *Nat Immunol* **16**, 628-634 (2015).

47. Rowell, J. *et al.* Distinct T-cell receptor (TCR) gene segment usage and MHC-restriction between foetal and adult thymus. *Elife* **13** (2024).

48. Yau, B. *et al.* A fluorescent timer reporter enables sorting of insulin secretory granules by age. *Journal of Biological Chemistry* **295**, 8901-8911 (2020).

49. Eastman, A.E. *et al.* Resolving Cell Cycle Speed in One Snapshot with a Live-Cell Fluorescent Reporter. *Cell Reports* **31** (2020).

50. Ono, M. TockyPrep: Data Preprocessing Methods for Flow Cytometric Fluorescent Timer Analysis. *arXiv [q-bio.QM]* (2024).

51. Chollet, F. *Deep Learning with Python*. Manning Publications Co., 2017.

52. Sojka, D.K., Bruniquel, D., Schwartz, R.H. & Singh, N.J. IL-2 Secretion by CD4+ T Cells In Vivo Is Rapid, Transient, and Influenced by TCR-Specific Competition. *The Journal of Immunology* **172**, 6136-6143 (2004).

53. Ono, M. & Satou, Y. Spectrum of Treg and Self-Reactive T cells: Single Cell Perspectives from Old Friend HTLV-1. *Discovery Immunology* (2024).

54. Shimizu, K. *et al.* PD-1 Imposes Qualitative Control of Cellular Transcriptomes in Response to T Cell Activation. *Mol Cell* **77**, 937-950 e936 (2020).

55. Dong, M. *et al.* Alterations in the Thymic Selection Threshold Skew the Self-Reactivity of the TCR Repertoire in Neonates. *J Immunol* **199**, 965-973 (2017).

56. Kirberg, J., Bosco, N., Deloulme, J.C., Ceredig, R. & Agenès, F. Peripheral T lymphocytes recirculating back into the thymus can mediate thymocyte positive selection. *J Immunol* **181**, 1207-1214 (2008).

57. Hale, J.S., Boursalian, T.E., Turk, G.L. & Fink, P.J. Thymic output in aged mice. *Proceedings of the National Academy of Sciences* **103**, 8447-8452 (2006).

58. Bilate, A.M. *et al.* Tissue-specific emergence of regulatory and intraepithelial T cells from a clonal T cell precursor. *Science Immunology* **1**, eaaf7471-eaaf7471 (2016).

59. Saxena, V., Lakhan, R., Iyyathurai, J. & Bromberg, J.S. Mechanisms of exTreg induction. *Eur J Immunol* **51**, 1956-1967 (2021).

60. Chiu, B.C., Stolberg, V.R., Zhang, H. & Chensue, S.W. Increased Foxp3(+) Treg cell activity reduces dendritic cell co-stimulatory molecule expression in aged mice. *Mech Ageing Dev* **128**, 618-627 (2007).

61. Maegawa, S. *et al.* Widespread and tissue specific age-related DNA methylation changes in mice. *Genome Res* **20**, 332-340 (2010).

62. Mi, T. *et al.* Conserved epigenetic hallmarks of T cell aging during immunity and malignancy. *Nature Aging* **4**, 1053-1063 (2024).

63. Boulbes, D.R. *et al.* A Survey on Data Reproducibility and the Effect of Publication Process on the Ethical Reporting of Laboratory Research. *Clin Cancer Res* **24**, 3447-3455 (2018).

64. Suzuki, H., Kinoshita, G., Tsunoi, T., Noju, K. & Araki, K. Mouse Hair Significantly Lightened Through Replacement of the Cysteine Residue in the N-Terminal Domain of Mc1r Using the CRISPR/Cas9 System. *Journal of Heredity* **111**, 640-645 (2020).

65. Takemoto, K. *et al.* Meiosis-Specific C19orf57/4930432K21Rik/BRME1 Modulates Localization of RAD51 and DMC1 to DSBs in Mouse Meiotic Recombination. *Cell Reports* **31** (2020).

66. Kawakami, R. *et al.* Distinct Foxp3 enhancer elements coordinate development, maintenance, and function of regulatory T cells. *Immunity* **54**, 947-961.e948 (2021).

67. Kim, D., Paggi, J.M., Park, C., Bennett, C. & Salzberg, S.L. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* **37**, 907-915 (2019).

68. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079 (2009).

69. Robinson, J.T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29**, 24-26 (2011).

70. Team, R.C. R: A Language and Environment for Statistical Computing. (2024).

71. Liaw, A. & Wiener, M. Classification and Regression by randomForest. *R News* **2**, 18-22 (2002).

72. Hornik, K. A CLUE for CLUster Ensembles. *Journal of Statistical Software* **14**, 1 - 25 (2005).

73. Hahsler, M., Piekenbrock, M. & Doran, D. dbscan: Fast Density-Based Clustering with R. *Journal of Statistical Software* **91**, 1 - 30 (2019).

74. Chollet, F. Keras. 2015.

75. D, S.C. & Clement, J.C. G-Net: Implementing an enhanced brain tumor segmentation framework using semantic segmentation design. *PLoS One* **19**, e0308236 (2024).

76. Developers, T. TensorFlow. v2.18.0 ed: Zenodo; 2024.

77. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

78. Robin, X. *et al.* pROC: an open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics* **12**, 77 (2011).

Figure Legends

Figure 1. Research Framework and Overview of Machine Learning Methods

(a) Pitfalls and risks in manual gating are schematically presented, highlighting the major pitfalls and risks associated with manual gating, emphasizing the hand-drawn nature of the methodology that introduces bias and undermines reproducibility.

(b) Proposed research framework for Machine Learning (ML)-assisted decoding of transcriptional dynamics. This schematic outlines the comprehensive workflow employed to unravel transcriptional dynamics of Foxp3 within a functional system. It covers the experimental design, generation of independent training and test datasets, training of ML models, performance evaluation, and data-driven identification of group-specific feature cells through model behavior analysis.

(c) Implementation of the research framework as *TockyMachineLearning*, a novel machine learning suite designed for this study. Data preprocessing, performed by *TockyPrep*, normalizes and transforms flow cytometric Timer data into standardised Timer Angle and Timer Intensity data. This pre-processed data then feeds into the TockyMachineLearning toolkit. Within this toolkit, TockyKmeansRF combines k-means clustering with Random Forest (RF) analysis, utilizing the mean decrease Gini index to identify feature cells. TockyConvNet transforms Timer Angle and Intensity data into 2D grayscale images representing cell density. These images are batch-processed by ConvNet, with model behaviours monitored using Grad-CAM to enable identification of feature cells at the single-cell level.

Figure 2. Development of the CRISPR-mediated CNS2 KO Foxp3-Tocky Mouse Model.

(a) Aligned sequence reads from ChIP-seq experiments (DRP003376 [66]) using anti-Runx1 antibody in Foxp3-negative CD4$^+$ T cells and Foxp3-expressing CD4$^+$ T cells, as well as anti-Foxp3 antibody in Foxp3-expressing CD4$^+$ T cells.

(b) The working model for CNS2-mediated Foxp3 transcriptional regulation involving Foxp3 and Runx1 proteins.

(c) CRISPR-Cas9 strategy employed to specifically target and delete the CNS2 sequence. The upper panel shows the architecture of the Foxp3-Timer locus. The lower panel shows the targeting oligonucleotide with homology arms and a short oligo sequence designed to replace the CNS2 sequence.

(d) PCR Strategies for Detecting CNS2 KO: Two PCR approaches were utilised to detect CNS2 deletion. The first set of primers (CNS2Del_F and CNS2Del_R) detects CNS2 deletion in both the BAC Foxp3 Timer and the endogenous Foxp3 gene, referred to as "*Common CNS2 Deletion*." The second set (DelCommon_F and Timer_74R as shown in (c)) allows discrimination of WT Foxp3 Timer and CNS2 KO Foxp3 Timer loci, termed as "*Foxp3 Timer-Specific Discrimination PCR*."

(e–f) Selection and Validation of CNS2 KO Founder Mouse, Specifically Founder #87 by (e) Common CNS2 Deletion PCR and (f) Foxp3 Timer-Specific Discrimination PCR.

(g) Sanger sequencing of the CNS2 region in founder mice derived from CRISPR-edited fertilised eggs.

(h) Breeding strategy illustrating how CNS2 KO Foxp3-Tocky mice were successively bred with WT animals over several generations to establish a stable mouse colony while ensuring that the endogenous Foxp3 gene remains free from mutations.

(i) Established genotyping PCR for discriminating CNS2 KO Foxp3-Tocky and WT Foxp3-Tocky mice.

(j) Sanger sequencing analysis to confirm the specificity of the Foxp3 Timer-specific discrimination genotyping PCR shown in (h). (Upper panel): Sequence reads from the WT Foxp3 Timer amplicon were aligned to the WT Foxp3 locus sequence. (Lower panel): Sequence reads from the CNS2 KO Foxp3 Timer amplicon were aligned to the WT Foxp3 Timer locus.

(k) Representative flow cytometric 2D plots showing Timer Blue and Timer Red fluorescence (upper panel) and Timer Angle and Intensity plots (lower panel) from WT Foxp3-Tocky and CNS2 KO Foxp3-Tocky samples.

(l) Flow cytometric analysis of Timer fluorescence in CRISPR CNS2KO Foxp3 Tocky. Box plots showing the mean fluorescence intensity (MFI) of Timer Red and Timer Blue fluorescence in CD4$^+$ T cells from WT Foxp3-Tocky and CNS2 KO Foxp3-Tocky. Statistical significance ($p < 0.01$) is indicated by asterisks and was assessed using Student's t-test.

(m) The percentage of cells within each Timer locus in the parent population in CD4$^+$ T cells from the superficial lymph nodes. Timer locus analysis categorises Timer Angle into the five loci: New (0˚); New-to-Persistent-transitioning (NPt, 0˚ - 30˚); Persistent (30˚ - 60˚); Persistent-to-Arrested-transitioning (PAt) (60˚ - 90˚); and Arrested (90˚). Statistical significance ($p < 0.05$) is indicated by asterisks and was assessed using the Mann-Whitney test, with a p-value adjustment.

Figure 3. TockyKmeansRF: Combinatorial Approach using Clustering and

Random Forest

(a)   Schematic diagram illustrating the TockyKmeansRF implementation, integrating

k-means clustering and Random Forest (RF) classification methodologies applied to

flow cytometric Timer fluorescence data for developing predictive models.

(b)   Training and test datasets generated from flow cytometric analysis of lymph node

samples from WT Foxp3 Tocky and CNS2 KO Foxp3 Tocky littermates.

(c)   Area Under the Curve (AUC) analysis of the TockyKmeansRF model across

various numbers of clusters (upper) and trees (lower) within the RF model.

(d)   Visualization of CNS2-dependent feature clusters identified in the test dataset

within Timer Angle and Intensity space (left panels) and within the original Timer

fluorescence space (right panels). The upper panels show the importance score by

Mean Decrease Gini (MDG) index, while lower panels show feature cells as identified

by the 60th percentile of MDG.

(e)   Results from the density-based clustering of feature cells.

(f)   Percentage of cells in each cluster in each sample, comparing CNS2 KO Foxp3

Tocky (KO) and WT Foxp3 Tocky (WT) mice.

(g)   Mean Fluorescence Intensity (MFI) of CD25, CD44, PD-1, and CD69 in the two

clusters identified and the rest of the cells (others) from WT Foxp3 Tocky mice.

Statistical analysis used the Kruskal-Wallis test to assess differences among groups,

followed by post-hoc Dunn's test with Bonferroni correction for p-value adjustment.

Note that Timer-negative cells are included only for baseline comparison and were not

considered in the statistical analysis.

(h)   Computational performance metrics of the TockyKmeansRF, including runtime

and memory usage, observed while incrementally increasing the CNS2 KO training

data by factors of 2, 3, and 4 to create proportionally larger datasets for analysis.

Figure 4. TockyConvNet: ConvNet Modelling Using Image Conversion and Gradient Analysis.

(a) Schematic representation of the image conversion method.

(b) Dot plots (left) and pseudocolour plots (right) showing image-converted Timer data.

(c) ConvNet architecture for TockyConvNet highlighting convolutional layers utilised for Gradient-weighted Class Activation Mapping (Grad-CAM) in (g-h).

(d) Model learning curve through a three-fold cross-validation method.

(e-f) Receiver Operator Characteristics (ROC) analysis (e) and Precision-Recall analysis (f) for benchmarking TockyConvNet against common manual gating approaches.

(g-h) Differential heatmaps are presented across convolutional layers of the ConvNet model. Heatmap values were determined by the differential of Grad-CAM outputs from WT and CNS2 KO Foxp3 Tocky samples. These heatmaps are presented in the Timer Angle-Intensity space (g) and the Timer Blue-Red space (h).

(i-j) Violin plots depict the distribution of CNS2 feature cells. In (i), WT CNS2 feature cells, or CNS2-dependent cells, are identified as those within the top 90th percentile of the differential heatmaps in (g) and (h). Conversely, in (j), KO CNS2 feature cells are identified as those below the 10th percentile. Statistical significance of differences between distributions is assessed using the Mann-Whitney test. Statistical significance levels indicated at $p < 0.01$ (**), $p < 0.001$ (****).

(k)   Mean Fluorescence Intensity (MFI) analysis of indicated markers for WT feature cluster, other Timer[+] cells, and Timer-negative cells using WT samples in the test dataset. Statistical analysis used the Kruskal-Wallis test to assess differences among groups, followed by post-hoc Dunn's test with Bonferroni correction for p-value adjustment. Note that Timer-negative cells are included only for baseline comparison and were not considered in the statistical analysis. Statistical significance levels indicated at $p < 0.01$ (**), $p < 0.005$ (***), $p < 0.001$ (****).

Figure 5. Gene Expression Analysis of CNS2-Dependent Feature Cells by Cross-Analysis of Grad-CAM Output from TockyConvNet and RNA-seq Data.

(a) Timer expression profile (Upper) and Timer Angle and Intensity profile (Lower) of pre-sort CD4$^+$ T cells and fractionated Foxp3 Timer$^+$ cells from WT Foxp3 Timer mice.

(b) Grad-CAM heatmap using the TockyConvNet, trained as shown in Figure 4, applied to RNA-seq flow cytometric data. (Upper) Visualization of WT feature analyzed via Attention-Conv2 Grad-CAM, highlighted by red on heatmap (Upper); (Lower) KO feature cells visualized through Conv2 Grad-CAM, highlighted with a blue on heatmap.

(c) Bar charts showing the percentage of WT feature cells (i.e. CNS2-dependent cells, left) and KO feature cells (i.e. CNS2-independent cells, right) in pre-sort CD4$^+$ T cells and fractionated Foxp3 Timer$^+$ cells.

(e-f) Expression dynamics of key genes in fractionated Foxp3 Timer$^+$ cells: (e) transcription factors downstream of TCR signalling; (f) genes associated with IL-2 and TGF-β signalling, along with prototypic upstream and downstream Foxp3 genes. P-values were obtained by the Wald test of the R package DESeq2 and adjusted by the Benjamini & Hochberg method. Asterisks indicate statistical significance, with an adjusted p-value < 0.05.

Figure 6. Generation of 'Foxp3 Timer Neonatal-to-Ageing Benchmarking Data' Across Developmental and Ageing Stages in WT Foxp3 Tocky Mice

(a) Timer Blue and Timer Red expression in CD4+ T cells from the thymus and spleen of WT Foxp3 Timer mice at various ages. All samples within the training dataset were concatenated per group and shown as pseudocolour plots. Days and weeks since birth are indicated.

(b) Normalised Timer fluorescence data from the flow cytometric analysis in (a).

(c) Timer Angle and Intensity transformed from the normalised Timer fluorescence

data in (b).

(d) Percentage of CD4$^+$ T cells in each of the indicated gates or the mean Timer

Angle.

(e) Percentage of mean Timer Angle plotted against logarithmically transformed age,

with axis labels indicating actual age in days. Each line represents a quadratic

regression model for each organ's data.

Figure 7. Development of the TockyConvNet for Quantitative Assessment of

Thymus and Spleen Characteristics in Foxp3 Transcriptional Dynamics

(a) Diagram of the ConvNet architecture for the age-adjusted TockyConvNet classifier

and continuous score models.

(b) ROC and Precision-Recall curve analysis using the TockyConvNet classifier.

(c) Thymus-Spleen Continuous Score data from the TockyConvNet continuous score

model analysing an independent test dataset, using linear age values (left) and log2-

transformed age values (right).

Figure 8. Optimizing and Benchmarking of the TockyConvNet Four-Class

Classifier

(a) Diagram of the 3 Conv-Layer TockyConvNet architecture as a four-class

classifier for classifying the two organs (Spleen vs Thymus) and stratified ages

(Young [< 30 days old]; vs Aged (> 30 days old))

(b) Optimisation of the TockyConvNet architecture, comparing models with one, two,

and three convolutional layer blocks by ROC and Precision-Recall curve analysis.

(c) Benchmarking results of TockyConvNet against other methods including

TockyKmeansRF and manual gating strategies in differentiating the four classes

using ROC (left) and Precision-Recall curves (right). The data generated in

Figure 6 were used.

Figure 9. Optimizing Grad-CAM Method for Analysing Foxp3 Timer Dynamics

(a) Heatmaps of Grad-CAM for each convolutional layer, visualising pixels used for

discriminating each indicated class.

(b) ROC curves showing class-discriminative capacity of each of the convolutional layers

by Grad-CAM.

(c) Heatmap visualisation of the Grad-CAM outputs for each class in both Timer Angle

and Intensity space, and original Timer fluorescence space.

Figure 1

a

Pitfalls and Risks of 'Manual Gating' in Flow Cytometry Data Analysis

b

c

a

Runx1 ChIP (Foxp3−)
Runx1 ChIP (Foxp3+)
Foxp3 ChIP (Foxp3+)
Foxp3 (RefSeq)

CNS2

b

Modulation

Transcription

Foxp3
Runx1

Foxp3

Exon 1   CNS2   Exon 2   Exon 3
CDS

c

Foxp3 Tocky:
BAC Tg (Foxp3 ^{Timer})

5' Homology Arm   3' Homology Arm
CNS2 (to be deleted)
PAM, 5' crRNA   3' crRNA,PAM
Exon   Fast-FT
Timer_74R

CNS2 KO Foxp3 Tocky:
BAC Tg (Foxp3 $^{Timer \Delta CNS2 KO}$)

Exon   Fast-FT
Timer_74R

Inserted Sequence
5 homology arm   3 homology arm
DelCommon_F
PAM   PAM
Targeting Oligonucleotide

d

CNS2   3 homology arm   Foxp3 Timer Locus
PAM, 5' crRNA   3' crRNA,PAM   Fast-FT
5' homology arm   exon   Foxp3   WT Foxp3 Locus
DelCommon_F   CNS2Del_R
CNS2Del_F   Timer_74R
Common CNS2 Deletion
Foxp3 Timer-Specific Discrimination PCR between WT Foxp3 Timer and CNS2 KO Foxp3 Timer

e

PCR: Common CNS2 Deletion

M 81 82 83 84 85 86 87 88 89 90 91 92

WT: 1141 bp
KO: 286 bp

1.5k
500
1k
300
400

f

Foxp3 Timer-Specific Discrimination PCR

M 82 81 87 90

WT: 2.8k bp
KO: 2k bp

3k
2.5k
2k

g

5,750   5,760   5,770   5,780   5,790   5,800

Foxp3
5' crRNA

Inserted Sequence

h

Foxp3 Tocky   Fertilised eggs   CRISPR   CNS2 KO-specific Foxp3 Tocky Founder
Founder Screening

Colony Establishment
CNS2 KO Foxp3 Tocky Colony

x

WT
Successively bred with WT

i

Foxp3 Timer-Specific Discrimination PCR

M 1 2

WT: 2.8k bp
KO: 2k bp

3k
2.5k
2k

j

Sequence Sample: WT Foxp3 Timer
Reference: WT Foxp3 Locus

CNS2 (to be deleted)
5' read   3' read

Sequence Sample: CNS2 KO Foxp3 Timer
Reference: WT Foxp3 Locus

CNS2 (to be deleted)
5' read
CNS2 (to be deleted)
3' read

k

WT Foxp3 Tocky   CNS2 KO Foxp3 Tocky

Timer Blue vs Timer Red
Intensity vs Angle

l

Timer Blue   Timer Red
ns   **
Mean Fluorescence Intensity (MFI)
CNS2 KO   WT   CNS2 KO   WT

m

Percentage (%)
Genotype
CNS2 KO Foxp3 Tocky
WT Foxp3 Tocky

New   NPt   Persistent   PAt   Arrested
Locus

New   Arrested
NPt   Persistent   PAt
0   30   60   90

Figure 3

Figure 4

Figure 6

Figure 7

Figure 8

a

b

	ROC Curve	Precision-Recall Curve

TockyCNN

Conv 3-Layers

ThymusYoung | ThymusAged | SpleenYoung | SpleenAged

Conv 2-Layers

ThymusYoung | ThymusAged | SpleenYoung | SpleenAged

Conv 1-Layer

ThymusYoung | ThymusAged | SpleenYoung | SpleenAged

c

	ROC Curve	Precision-Recall Curve

Polygon: Red (high)

ThymusYoung | ThymusAged | SpleenYoung | SpleenAged

Q3: Blue-Red+

ThymusYoung | ThymusAged | SpleenYoung | SpleenAged

TockyKmeansRF

ThymusYoung | ThymusAged | SpleenYoung | SpleenAged

Figure 9

a

	Spleen-Aged	Spleen-Young	Thymus-Aged	Thymus-Young
Conv1	SpleenAged_conv1	SpleenYoung_conv1	ThymusAged_conv1	ThymusYoung_conv1
Attention1-Conv	SpleenAged_attention1_conv	SpleenYoung_attention1_conv	ThymusAged_attention1_conv	ThymusYoung_attention1_conv
Conv2	SpleenAged_conv2	SpleenYoung_conv2	ThymusAged_conv2	ThymusYoung_conv2
Attention2-Conv	SpleenAged_attention2_conv	SpleenYoung_attention2_conv	ThymusAged_attention2_conv	ThymusYoung_attention2_conv
Conv3	SpleenAged_conv3	SpleenYoung_conv3	ThymusAged_conv3	ThymusYoung_conv3
Attention3-Conv	SpleenAged_attention3_conv	SpleenYoung_attention3_conv	ThymusAged_attention3_conv	ThymusYoung_attention3_conv

Timer-Intensity (y-axis)

Timer-Angle (x-axis)

b

	Spleen-Aged	Spleen-Young	Thymus-Aged	Thymus-Young
Conv1	TPR vs FPR			
Attention1-Conv				
Conv2				
Attention2-Conv				
Conv3				
Attention3-Conv				

Percentile
- 50%
- 60%
- 70%
- 80%
- 90%

c

Spleen, Aged: Attention2-Conv

Spleen, Young: Conv2

Thymus, Aged: Conv2

Thymus, Young: Attention1-Conv