High genomic stability of *w*Mel *Wolbachia* after introgression into three geographically distinct *Aedes aegypti* populations

Kimberley R. Dainty[1][†], Jane Hawkey[2], Endah Supriyati[3], Louise M. Judd[2], Alexander Uribe[4], Iván D. Vélez[4,5], Dang D. Anh[6], Eggi Arguni[3,7], Warsito Tantowijoyo[3], Scott L. O'Neill[8], Kathryn E. Holt[2,9], Cameron P. Simmons[1,8], Heather A. Flores[10*]

1 Department of Microbiology, Monash University, Melbourne, Victoria, Australia

2 Department of Infectious Diseases, School of Translational Medicine, Monash University, Melbourne, Victoria, Australia

3 World Mosquito Program Yogyakarta, Centre for Tropical Medicine, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

4 World Mosquito Program, Medellín, Colombia

5 PECET, U de Antioquia, Medellín, Colombia

6 National Institute of Hygiene and Epidemiology, Hanoi, Vietnam

7 Department of Child Health, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

8 World Mosquito Program, Monash University, Melbourne, Victoria, Australia

9 Department of Infection Biology, London School of Hygiene and Tropical Medicine, London, UK

10 School of Biological Sciences, Monash University, Melbourne, Victoria, Australia

[†] Present address: Department of Biochemistry, University of Otago, Dunedin, New Zealand

* heather.flores@monash.edu

Abstract

The introgression of *w*Mel *Wolbachia* into *Aedes aegypti* populations is being used for the biocontrol of arboviruses such as dengue and chikungunya in 14 countries to date. A *w*Mel infection in *Ae. aegypti* both reduces the transmission of viruses by the mosquito and causes a reproductive manipulation that aids *w*Mel introgression into naïve populations. However, a critical concern is whether *w*Mel could evolve over time, potentially diminishing these desired phenotypes. Here, we investigated the stability of the *w*Mel genome in *Ae. aegypti* released for biocontrol in Colombia, Indonesia, and Vietnam. We sequenced the *w*Mel genome at the start of releases and up to six years after *w*Mel introgression into each population. Our study identifies very few genomic changes, suggesting the *w*Mel genome is not rapidly evolving despite its release into three geographically different field sites and subsequent exposure to novel environments. These results align with previous *w*Mel sequencing studies from Australia and provide strong evidence for the long-term genomic stability of *w*Mel, reinforcing its potential as a reliable biocontrol tool against *Ae. aegypti*-transmitted arboviruses.

Introduction

Human pathogenic viruses such as dengue, Zika, and yellow fever are spread primarily by *Aedes aegypti*. The burden of these diseases has increased substantially over the past 50 years due to increasing globalisation, the rise in population-dense cities, and expansion of the geographical breeding range of *Ae. aegypti* (Gubler 2002; Bonizzoni et al. 2013; Mayer et al. 2017). To reduce the burden of these viruses, a public health intervention utilising the bacterial endosymbiont, *Wolbachia pipientis* (*w*Mel strain), has been deployed in 14 countries to date (World Mosquito Program, n.d.).

Stable *w*Mel infection of *Ae. aegypti* reduces the ability of the mosquito to transmit multiple human pathogenic viruses such as dengue, chikungunya and Zika (Aliota, Peinado, et al., 2016; Aliota, Walker, et al., 2016; Carrington et al., 2018; Pinto et al., 2021; Rocha et al., 2019; Tan et al., 2017; Utarini et al., 2021; van den Hurk et al., 2012; Ivan Dario Velez et al., 2023; Walker et al., 2011). Effective biocontrol of these pathogens requires the spread and persistence of a *w*Mel infection in *Ae. aegypti*. This is facilitated by both *w*Mel maternal transmission, and the *w*Mel-induced reproductive modification cytoplasmic incompatibility

(CI), which aids the introgression of *w*Mel through naïve populations (Ant et al., 2018;

Walker et al., 2011).

With an endorsement from the World Health Organisation and recent investment of funds

geared towards expansion of this biocontrol method, it is likely that the next decade will see

further global deployment of *w*Mel and other *Wolbachia* strains for use in *Ae. aegypti* and

other mosquito species to reduce the disease burdens. One risk of using an endosymbiont as a

biocontrol tool is the potential for this endosymbiont to undergo genomic changes which

limit the phenotypic effects the endosymbiont has on its host. In this instance, the cessation of

virus inhibition or the ability for *w*Mel to spread through or remain sustained in field

populations would halt the effectiveness of the biocontrol. The genomic variation of *w*Mel

has been examined in Australian *Ae. aegypti.* Studies compared the *w*Mel genome from *Ae.*

aegypti in Cairns, Australia when first released and up to nine years post release to that of the

reference *w*Mel genome sequenced directly from *D. melanogaster* (Wu et al., 2004). These

studies found very little genomic variation after transinfection and field establishment,

suggesting the *w*Mel genome is stable in Australian *Ae. aegypti* (Dainty et al., 2021; Huang et

al., 2020; Ross et al., 2022).

After the success of Australian releases, *w*Mel was introgressed into *Ae. aegypti* populations

in a further 13 countries, including Indonesia, Vietnam, and Colombia (Hien et al., 2021;

Tantowijoyo et al., 2020; Iván Darío Velez et al., 2023; World Mosquito Program, n.d.).

Exposure of *w*Mel to geographically diverse environments in their novel *Ae. aegypti* host

again prompted concerns this would result in the local adaptation of *w*Mel. Studies of *w*Mel

in *D. melanogaster* have identified strong geographical haplotypes (Early & Clark, 2013;

Richardson et al., 2012) which have been hypothesised to be influenced by selection and

local adaptation rather than genetic drift alone (Early & Clark, 2013). To date there is no

conclusive field data indicating the crucial phenotypic effects *w*Mel induces in *Ae. aegypti*

have been impacted by *w*Mel evolution. However, *w*Mel introgression and stability in fields

sites has been highly variable. While many operational factors related to field releases such as

number, life stage and method of mosquitoes released as well as local climates and host

fitness may underlie this variation, changes to the *w*Mel genome may also explain the

variability observed. One way to investigate this and the future potential for this to occur is to

understand the breadth and spread of genomic changes occurring to the *w*Mel genome when

used as a biocontrol agent.

Here we examined the genomic changes to the *w*Mel genome after introgression into three

geographically and environmentally diverse *Ae. aegypti* populations in Indonesia (first

released January 2014), Vietnam (first released May 2014), and Colombia (first released June

2015). Overall, we found that the *w*Mel genome is highly stable in the 4-6 years of sampling

post release in each population, suggesting the use of this endosymbiont as a biocontrol agent

is not inducing rapid evolution that may lead to the loss of critical phenotype effects.

Methods

<u>Sample collection and processing</u>

Colombian mosquitoes

Colombian mosquito samples used for sequencing were collected from releases performed in

the París comuna of Bello, Antioquia. Pilot releases in the central part of París took place

between June and December of 2015, and a phase 2 release throughout the remaining París

comuna took place between July and August 2016 (Iván Darío Velez et al., 2023). Female

*w*Mel-infected *Ae. aegypti* mosquitoes were collected on March 19th 2015 from an outbred

pre-release colony which was used for subsequent field releases that commenced in June. Ten

individual mosquitoes were chosen for sequencing, and *w*Mel genomes represent the 2015

early-release *w*Mel genomes from Colombia. Female *w*Mel-infected *Ae. aegypti* mosquitoes

were collected from the pilot release region of París on October 29th 2019 via BG Sentinel

traps (Biogents AG, Germany). These 24 *w*Mel genomes sequenced were from mosquitoes

collected across 5 individual BG Sentinel traps and represent the 2019 recent-collection *w*Mel

genomes from Colombia.

Indonesian mosquitoes

Indonesian mosquito samples used for sequencing were collected from the Sleman and

Bantul districts of Yogyakarta, where *w*Mel releases took place between January-June 2014

and November 2014-May 2015, respectively. Female *w*Mel-infected *Ae. aegypti* mosquitoes

were collected from Bantul district and Sleman district on February 23rd 2015 via BG

Sentinel traps as described in (Tantowijoyo et al., 2020). These 9 *w*Mel genomes sequenced

were from mosquitoes collected from 9 individual BG Sentinel traps and represent the 2015

early-release *w*Mel genomes from Indonesia. Female *w*Mel-infected *Ae. aegypti* mosquitoes

were collected again from these two same districts on February 20[th] 2020 via BG Sentinel

traps. These 30 *w*Mel genomes sequenced were from mosquitoes collected across 26

individual BG Sentinel traps and represent the 2020 recent-collection *w*Mel genomes from

Indonesia.

Vietnamese mosquitoes

Vietnamese mosquito samples used for sequencing were collected from Tri Nguyen Island,

where *w*Mel releases took place over a 27-week period from May to November 2014. Female

*w*Mel-infected *Ae. aegypti* mosquitoes were collected from the central and southern regions

of Tri Nguyen Island (see Figure 1, Hien et al., 2021), via BG Sentinel traps between the 27[th]

October 2015 and the 3[rd] September 2015. Mosquitoes were collected exclusively from the

central and southern end of Tri Nguyen Island as this was where *w*Mel infection prevalence

was highest. These 9 *w*Mel genomes sequenced were from mosquitoes collected from 8

individual BG Sentinel traps and represent the 2015 early-release *w*Mel genomes from

Vietnam. Female *w*Mel-infected *Ae. aegypti* mosquitoes were collected from the southern

end of Tri Nguyen Island where *w*Mel was still established via BG Sentinel traps between the

28[th] May 2019 and the 4[th] June 2019. These 18 *w*Mel genomes sequenced were from

mosquitoes collected from 6 individual BG Sentinel traps and represent the 2019 recent-

collection *w*Mel genomes from Vietnam.

All mosquitoes

DNA was extracted from the abdomens of single mosquitoes and used for sequencing as this

would enrich for *w*Mel which is at higher density in the ovaries and Malpighian tubules

(Amuzu & McGraw, 2016; Fraser et al., 2020). Sequencing libraries were also performed

from single DNA extracts, and thus, each genome represents the diversity present within an

individual mosquito. Single abdomens were homogenised using a hand-pestle in Buffer ATL

before DNA was extracted using the MagAttract HMW DNA Kit (QIAGEN) with a 50 ⌈L

elution in nuclease-free water following manufacturer's instructions. Libraries were

generated for each single mosquito sample using the Illumina DNA Prep (M) Tagmentation

kit (Illumina®) using unique index tags. All Illumina DNA Prep libraries were made

according to manufacturer's directions, excepting all reactions were scaled to 25% of

recommended volumes and 25ng of template DNA was used. Samples were sequenced via

the Illumina platform using NovaSeq 6000 S4 Reagent Kit v1.5 (300 cycles) (Illumina®),

generating 150-base-paired-end reads.

<u>Analysis</u>

Illumina reads belonging to *Ae. aegypti* (accession NC_035107.1) were identified using

BioBloom Tools v2.3.2 (Chu et al., 2014) using default parameters and excluded from

downstream analysis. The remaining Illumina reads, not trimmed before analysis, were

mapped to the *D. melanogaster w*Mel reference genome (Wu et al., 2004) (NC_002978.6)

and single nucleotide variants identified using the RedDog V1b.11 pipeline

(https://github.com/katholt/reddog) using standard parameters and following the developers'

guidelines. Briefly, this pipeline maps reads to the reference sequence using Bowtie2 version

2.2.3 (Langmead & Salzberg, 2012) before SNP's with QUAL values ε 30 are called with

SAMtools version 1.9 (Li, 2011). Mean Illumina read depth for Colombian samples was

262.49x, for Indonesian samples was 846.51x, and for Vietnamese samples was 61.63x.

Coverage was 100% for all samples.

Low frequency variants were detected using LoFreq V2.1.3.1 (Wilm et al., 2012) with default

parameters. Variants were removed from the final data set if strand bias was ε 10. Variants

were annotated with SnpEff V5.2, using the classical style of annotation and the database

Wolbachia endosymbiont of *Drosophila melanogaster* GCA 000008025 (Cingolani et al.,

2012). Gene copy number variation was analysed by normalising the sequencing depth of

coverage of each gene in sequenced genomes by dividing the mean sequence depth for each

gene by the mean sequence depth of the whole genome.

Insertion sequence elements were identified by using the *w*Mel reference genome

(NC_002978.6) to search against the ISfinder (Siguier et al., 2006) database via the ISsaga

(Varani et al., 2011) web server available at http://issaga.biotoul.fr/issaga_index.php. IS

queries that were identified to have greater than 80% similarity (Supplementary Table 3)

were mapped to the short-read sequencing data for each sample using ISMapper (Hawkey et

al., 2015) to identify IS insertion sites with default parameters. Insertion or deletion of IS

elements identified as imprecise (*) or uncertain (?) were considered mapping artefacts.

Results

No novel genomic changes occurred in the establishment of country-specific release lines
195

We sequenced *w*Mel from 9 Indonesian (Yogyakarta providence), 9 Vietnamese (Tri Nguyen
196

Island), and 10 Colombian (París) individual mosquitoes collected just prior to or soon after
197

introgression of *w*Mel (Figure 1). To generate these release strains, *w*Mel-infected Australian
198

Ae. aegypti mothers were backcrossed to *Ae. aegypti* males from Indonesian, Vietnamese,
199

and Colombian populations. *w*Mel from Australian *Ae. aegypti* used in these backcrosses has
200

previously shown to differ by only one SNP different from the *D. melanogaster w*Mel
201

reference sequence (NC_002978.6) (Dainty et al., 2021). Therefore, we compared their
202

genomes to the *D. melanogaster w*Mel reference sequence (NC_002978.6) to identify
203

changes that occurred during the establishment of country-specific release lines.
204

Across all pre/early release samples, only a single SNP was identified. We identified
206

SNP_1174712 in 4/10 *w*Mel genomes collected from Colombia in 2015 (Figure 2). This SNP
207

causes a synonymous change of base T->C in the WD_1228 gene, the product of which is a
208

hypothetical protein (Table 1). Previously, this SNP has been identified in *w*Mel from
209

Australian *Ae. aegypti* in up to 30% of individuals (Dainty et al., 2021; Huang et al., 2020;
210

Ross et al., 2022), suggesting it was likely introduced into the Colombian background during
211

backcrossing. SNP_1174712 was not found in the Indonesian or Vietnamese populations.
212

These two *w*Mel-infected *Ae. aegypti* strains were generated via backcrossing of the same
213

Australian population known to carry the SNP. Since this SNP was not fixed in the Australian
214

*w*Mel population, it is possible this SNP was not inherited by the final Indonesian and
215

Vietnamese colonies or inherited in low prevalence such that it was not sampled.
216

Alternatively, we can't discount the possibility that this SNP may cause a fitness
217

disadvantage in these populations and thus has been selected against prior to the first
218

collections and sequencing of these populations.
219

Table 1. SNPs present in *w*Mel *Wolbachia* sequenced from *Ae. aegypti* collected in
221

Colombia, Indonesia, and Vietnam compared to the *D. melanogaster w*Mel reference
222

sequence.
223

SNP Position	Population	Reference nucleotide	Alternate nucleotide	Change outcome	Gene	Ancestral codon	Derived codon	Ancestral amino acid	Derived amino acid	Gene product
547257	Colombia 2019 1/24	T	G	Intergenic	-	-	-	-	-	-
591822	Indonesia 2020 1/30	G	A	Non-synonymous	WD_0610	ACA	ATA	T	I	Helicase, SNF2 family
1116668	Vietnam 2019 1/18	T	C	Synonymous	WD_1170	CCT	CCC	P	P	Enoyl-(acyl-carrier-protein) reductase II
1174712	Colombia 2015 4/10 Colombia 2019 13/24	T	C	Synonymous	WD_1228	CCT	CCC	P	P	Hypothetical protein

wMel genome is highly stable up to 6 years post-field release across all three field sites

We next sequenced *w*Mel genomes from Indonesian, Vietnamese, and Colombian *Ae. aegypti*

populations 4-6 years post-release, to determine genomic changes influenced by field

establishment and differing environments. Across the *w*Mel genomes sequenced here (30

from Indonesia, 18 from Vietnam, 24 from Colombia), one recurrent and three unique single

nucleotide polymorphisms were identified in comparison to early field caught mosquitoes

from each respective country. Of these three unique SNP's, none were present in across

multiple locations, and all three were found only once.

SNP_1174712 was again observed only in Colombian mosquitoes. Similar to Australian

populations, we again observed that this SNP did not significantly increase nor decrease in

frequency (4/10 to 13/24), suggesting it is not under selection in the Colombian population.

SNP_547257 was found in one *w*Mel genome collected from a Colombian *Ae. aegypti* in

2019 (Figure 2). This SNP causes an intergenic change of T->G at position 547257 (Table1)

and is thus unlikely to be functionally impactful. Next, SNP_591822 present in one *w*Mel

genome collected from an Indonesian *Ae. aegypti* in 2020 causes a non-synonymous change

at position 591822 in the genome (Figure 2). This SNP constitutes a base change from G->A

resulting in an amino acid change from threonine to isoleucine in gene WD_0610, the

product of which is a SNF2 family helicase (Table1). This family of proteins have been

shown to be responsible for directing energy from ATP hydrolysis into the remodelling of

chromatin structures as well as processing of DNA damage repair, and transcriptional

replication (Eisen et al., 1995; D. P. Ryan & Owen-Hughes, 2011). As this SNP was only

found in 1/30 genomes sequenced from Indonesia in 2020, this indicates either this SNP has

been of recent occurrence, or it is providing no evolutionary benefit and thus not spreading

throughout the population. Monitoring the prevalence of this SNP in the future will allow for

distinguishing between these two possibilities. Lastly, SNP_1116668 was found in one *w*Mel

genome from *Ae. aegypti* collected in Vietnam in 2019 (Figure 2). This SNP causes a

synonymous base change from T->C in the gene WD_1170 which produces an enoyl-(acyl-

carrier-protein) reductase II product (Table1). Again, as this SNP was found in a single

genome only, and is causing a synonymous change, it is unlikely to impact the gene function.

In addition to these three polymorphisms and the previously described SNP_1174712, one

further SNP was identified in all 100 genomes sequenced when compared to the *w*Mel

reference genome, a SNP at position 1097797. However, this SNP has previously been shown

to likely be an error in the *D. melanogaster w*Mel reference sequence (Dainty et al., 2021)

rather than a novel genomic change, and is thus not included here. Additionally, when

compared to the *D. melanogaster w*Mel reference genome, seven indels were identified in the

genomes sequenced here. These seven indels were previously described in the *Ae. aegypti*

*w*Mel genomes sequenced in (Dainty et al., 2021), and again shown to be errors in the *D.*

*melanogaster w*Mel reference sequence.

Gene copy number variation was analysed in each genome by normalising gene mean

sequencing depth to the mean depth coverage of the entire genome. No genes showed distinct

copy number variation in any of the genomes (Supplementary Figure 1). The sequenced

genomes were also analysed to assess insertion sequence (IS) movement, using the ISsaga

and ISMapper programs. However, no evidence for movement, deletion, or novel insertion of

any IS element was identified in any of the genomes. Due to the limitations of short-read

sequencing, large scale structural changes cannot be ruled out in these genomes. However,

given the lack of identified SNPs and indels, and the previously documented stability of the

*w*Mel genome in Australian *Ae. aegypti,* it is unlikely that these genomes have undergone

such structural changes.

Evidence of low frequency wMel genomic heteroplasmy within a mosquito

The above results looked at population level variation of *w*Mel genomes across individual *Ae.*

aegypti females. However, our sequencing was performed on single female samples allowing

us to investigate *w*Mel heteroplasmy, i.e. *w*Mel variants *within* a single mosquito. We found a

number of low frequency variants within each individual mosquito; however, the majority of

these were at frequencies less than 10% (Supplementary Table 1). Averaged across the

collected populations, the number of variants with frequencies between 10-20% varies

between 41.88 and 163.3 variants per genome. The number of variants with frequencies

above 20% ranges between 1.75 and 27 variants per genome (Table 2). Interestingly, the

average level of *w*Mel heteroplasmy within a mosquito decreased in each population in later

collections despite many generations in the field. The number of variants did not correlate

with either *w*Mel density or sequencing depth (Supplementary Table 2). Collectively, this

suggests there are low levels of standing variation in these genomes that have not

substantially increased in frequency nor yet fixed in the population.

Table 2. Mean number of low-frequency SNPs at frequencies of 10-20% and >20%

identified per genome from each population. Mean number of variants predicted to cause

missense, synonymous, or other impacts for each frequency is also reported.

Population Collection (Sample size)	Mean # of SNPs at 10-20%	Mean # of SNPs with predicated impact variants at 10-20% (%)		Mean # of SNPs >20%	Mean # of SNPs with predicated impact variants at >20% (%)	
Indonesia 2015 (9)	99.44	Missense	11.51	5.44	Missense	20.69
		Synonymous	15.86		Synonymous	12.07
		Other	72.63		Other	67.24
Indonesia 2020 (30)	85.1	Missense	11.99	7.3	Missense	6.83
		Synonymous	13.79		Synonymous	2.81
		Other	74.22		Other	90.36
Vietnam 2015 (9)	121.1	Missense	19.86	27.0	Missense	14.78
		Synonymous	34.41		Synonymous	22.60
		Other	45.73		Other	62.61
Vietnam 2019 (18)	43.93	Missense	13.34	19.74	Missense	8.38
		Synonymous	22.28		Synonymous	6.09
		Other	64.38		Other	85.53

Colombia 2015 (10)	163.3	Missense	15.87	30	Missense	7.23
		Synonymous	22.39		Synonymous	7.23
		Other	61.75		Other	85.53
Colombia 2019 (24)	41.88	Missense	13.83	1.75	Missense	6.67
		Synonymous	18.51		Synonymous	21.33
		Other	67.66		Other	72.00

Discussion

Wolbachia biocontrol is transforming the fight against *Ae. aegypti*-transmitted arboviruses

with the *w*Mel strain being deployed across diverse geographic and ecological landscapes.

Given the evolutionary pressures imposed by these novel environments, we hypothesized that

*w*Mel would be undergoing adaptive genomic changes, which could potentially explain

variation in *w*Mel introgression and stability observed across field sites. Our study however

identified only minor genomic changes present at low frequency in the three populations

studied, thus suggesting that despite environmental pressures and novel hosts, the *w*Mel

genome remains remarkably resilient, reinforcing its robustness as a long-term biocontrol

strategy.

The evolutionary rate of *w*Mel in its native host *D. melanogaster* has been estimated to be

low compared to other bacteria (Duchêne et al., 2016), at 6.8×10^{-10} substitutions per site per

D. melanogaster generation (Early & Clark, 2013; Richardson et al., 2012). Assuming this

evolutionary rate is similar in *Ae. aegypti*, combined with the relatively short time *w*Mel has

been established in these populations (~5 years, 12 generations per year), this rate predicts

approximately 0.05 substitutions to have occurred per genome. Thus, the level of variation

reported here is consistent with this estimation suggesting *w*Mel is evolving similar to levels

observed in its natural host.

Although low levels of fixed SNPs were identified in the studied *w*Mel genomes, we did

identify higher levels of low frequency polymorphisms, indicating there is mutational input to

these genomes. Previous *w*Mel sequencing from Australian mosquitoes was performed on

pooled samples (Dainty et al., 2021). Our sequencing here was performed on single female

mosquitoes allowing us to attribute this variation to heteroplasmy of wMel within single

mosquitoes. Our analysis shows that on average 40-190 variants occur per genome (between

10-50% frequency), with this number reaching more than 400 in some genomes. Clearly,

however, these polymorphisms are not being transmitted at high levels, as evidenced by the

low levels of SNP fixation detail in this study. Severe bottlenecks such as those that occur

during maternal transmission have been suggested to contribute to evolution of insect

endosymbionts by altering the efficacy of purifying selection (Moran, 1996; Wernegreen,

2002). Additionally, our sequencing included wMel found in somatic tissues within the

abdomen, a dead end for this variation, as *Wolbachia* is maternally transmitted through the

germline. Our data however suggest there are reasonable amounts of standing genetic

variation present in wMel populations that may allow wMel to respond to various selective

pressures it may experience in its mosquito host and/or environment.

These findings in combination with previous reports (Dainty et al., 2021; Huang et al., 2020;

Ross et al., 2022), show the wMel genome to be stable in four geographically distinct *Ae.*

aegypti populations. Stability of the desired wMel-induced phenotypes is critical for the

success of the *Wolbachia* biocontrol methods, thus these findings in combination with

entomological studies reflect positively on its continued success (Ary A. Hoffmann et al.,

2014; Ross et al., 2022). However, wMel is only one element of the tripartite interaction in

this biocontrol method. In addition to *Wolbachia*, the *Ae. aegypti* host or the viruses that

Wolbachia inhibit may evolve resistance, potentially reducing the efficacy of the method

(Bull & Turelli, 2013; Edenborough et al., 2021; Ritchie et al., 2018). Sequencing of *Ae.*

aegypti up to 8 years post release in Cairns, Australia, revealed that host genomes remained

mostly stable (Lau et al., 2021), however further work is needed to understand if this

translates to other populations. Previous laboratory studies using artificial selection have

suggested *Ae. aegypti* genetic variation can impact the strength of wMel-mediated virus

blocking (Ford et al., 2019), indicating the need to monitor changes to this genome in

addition to critical *Wolbachia*-induced phenotypes over time. The evolution of viruses in

wMel-infected hosts has not been extensively studied. RNA viruses are known to have a high

mutation rate but are also exposed to strong purifying selection, in particular, as arboviruses

must replicate in both human and insect hosts. Multiple lab evolution studies have shown that

DENV and Drosophila C virus did not evolve to evade inhibition by *Wolbachia* (Koh et al.,

2019; Martinez et al., 2019). However, a recent study showed that selective pressure from

*w*Mel resulted in positive selection on DENV (Thi Hue Kien et al., 2023). The viral kinetics of the positively selected SNP have not been evaluated in *w*Mel-infected mosquitoes, so it is unclear whether this change results in evasion of *w*Mel inhibition. Overall, these data highlight the importance of monitoring all three aspects of this tripartite interaction.

Here, we report a multi-country sequencing analysis of 100 individual *w*Mel genomes in *Ae. aegypti*. We assessed what changes occurred to the *w*Mel genome after introgression into three global *Ae. aegypti* populations, and post-field release up to six years. We show *w*Mel genome stability in Colombian, Indonesian, and Vietnamese *Ae. aegypti* field populations, finding limited genomic changes. Despite the different geographical locations, these *w*Mel populations show little divergence since introduction, indicating that *w*Mel has not undergone selection in response to local, country-specific selective pressures. This indicates the use of *w*Mel as a biocontrol agent in novel global *Ae. aegypti* populations has had little effect on the *w*Mel genome in the timeframe of this study. These findings provide reassurance toward the potential of the endosymbiont *w*Mel to remain a successful public-health biocontrol method against arboviruses globally.

Acknowledgements

We would like to thank I'ah Donovan-Banfield for technical assistance.

Figure 1. Timeline of the *w*Mel genome sampling. Release and establishment of *w*Mel-
infected *Ae. aegypti* in Indonesia, Vietnam, and Colombia are indicated by yellow, orange,
and green bars respectively, with the start of each colour block indicating initiation of *w*Mel-
infected mosquito releases. Time points at which *w*Mel-infected *Ae. aegypti* were collected
for sampling from each site are indicated by black mosquitoes. Time periods between
collection times are displayed in black text.

Figure 2. SNPs present in *w*Mel *Wolbachia* collected from *Ae. aegypti* compared to the
*w*Mel reference sequence from *D. melanogaster*. Colours corresponding to the following:
blue bars represent forward strand gene, green bars represent reverse strand genes, orange
segments represent phage regions (in clockwise order WO-A, pyocin-like, WO-B, WO-B.
SNPs are represented by black dots with corresponding information in connected boxes. Box

information includes SNP position, mutation outcome, and population and number of
genomes that contained the SNP.

References

Aliota, M. T., Peinado, S. A., Velez, I. D., & Osorio, J. E. (2016). The wMel strain of Wolbachia Reduces Transmission of Zika virus by Aedes aegypti. *Scientific Reports*, *6*(1), 28792.

Aliota, M. T., Walker, E. C., Yepes, A. U., Velez, I. D., Christensen, B. M., & Osorio, J. E. (2016). The wMel Strain of Wolbachia Reduces Transmission of Chikungunya Virus in Aedes aegypti. *PLoS Neglected Tropical Diseases*, *10*(4), e0004677.

Amuzu, H. E., & McGraw, E. A. (2016). Wolbachia-Based Dengue Virus Inhibition Is Not Tissue-Specific in Aedes aegypti. *PLoS Neglected Tropical Diseases*, *10*(11), e0005145.

Ant, T. H., Herd, C. S., Geoghegan, V., Hoffmann, A. A., & Sinkins, S. P. (2018). The Wolbachia strain wAu provides highly efficient virus transmission blocking in Aedes aegypti. *PLoS Pathogens*, *14*(1), e1006815.

Bull, J. J., & Turelli, M. (2013). Wolbachia versus dengue: Evolutionary forecasts. *Evolution, Medicine, and Public Health*, *2013*(1), 197–207.

Carrington, L. B., Tran, B. C. N., Le, N. T. H., Luong, T. T. H., Nguyen, T. T., Nguyen, P. T., Nguyen, C. V. V., Nguyen, H. T. C., Vu, T. T., Vo, L. T., Le, D. T., Vu, N. T., Nguyen, G. T., Luu, H. Q., Dang, A. D., Hurst, T. P., O'Neill, S. L., Tran, V. T., Kien, D. T. H., … Simmons, C. P. (2018). Field- and clinically derived estimates of Wolbachia-mediated blocking of dengue virus transmission potential in Aedes aegypti mosquitoes. *Proceedings of the National Academy of Sciences*, *115*(2), 361–366.

Chu, J., Sadeghi, S., Raymond, A., Jackman, S. D., Nip, K. M., Mar, R., Mohamadi, H., Butterfield, Y. S., Robertson, A. G., & Birol, I. (2014). BioBloom tools: fast, accurate and memory-efficient host species sequence screening using bloom filters. *Bioinformatics* , *30*(23), 3402–3404.

Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., Land, S. J., Lu, X., & Ruden, D. M. (2012). A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. *Fly*, *6*(2), 80–92.

Dainty, K. R., Hawkey, J., Judd, L. M., Pacidônio, E. C., Duyvestyn, J. M., Gonçalves, D. S., Lin, S. Y., O'Donnell, T. B., O'Neill, S. L., Simmons, C. P., Holt, K. E., & Flores, H. A. (2021). wMel Wolbachia genome remains stable after 7 years in Australian Aedes

aegypti field populations. *Microbial Genomics*, *7*(9).

https://doi.org/10.1099/mgen.0.000641

Duchêne, S., Holt, K. E., Weill, F.-X., Le Hello, S., Hawkey, J., Edwards, D. J., Fourment,

M., & Holmes, E. C. (2016). Genome-scale rates of evolutionary change in bacteria.

Microbial Genomics, *2*(11), e000094.

Early, A. M., & Clark, A. G. (2013). Monophyly of Wolbachia pipientis genomes within

Drosophila melanogaster : geographic structuring, titre variation and host effects

across five populations. *Molecular Ecology*, *22*(23), 5765–5778.

Edenborough, K. M., Flores, H. A., Simmons, C. P., & Fraser, J. E. (2021). Using Wolbachia

to Eliminate Dengue: Will the Virus Fight Back? *Journal of Virology*, *95*(13),

e0220320.

Eisen, J. A., Sweder, K. S., & Hanawalt, P. C. (1995). Evolution of the SNF2 family of

proteins: subfamilies with distinct sequences and functions. *Nucleic Acids Research*,

23(14), 2715–2723.

Ford, S. A., Allen, S. L., Ohm, J. R., Sigle, L. T., Sebastian, A., Albert, I., Chenoweth, S. F.,

& McGraw, E. A. (2019). Selection on Aedes aegypti alters Wolbachia-mediated

dengue virus blocking and fitness. *Nature Microbiology*, *4*(11), 1832–1839.

Fraser, J. E., O'Donnell, T. B., Duyvestyn, J. M., O'Neill, S. L., Simmons, C. P., & Flores,

H. A. (2020). Novel phenotype of Wolbachia strain wPip in Aedes aegypti challenges

assumptions on mechanisms of Wolbachia-mediated dengue virus inhibition. *PLoS*

Pathogens, *16*(7), e1008410.

Frentiu, F. D., Zakir, T., Walker, T., Popovici, J., Pyke, A. T., van den Hurk, A., McGraw, E.

A., & O'Neill, S. L. (2014). Limited Dengue Virus Replication in Field-Collected

Aedes aegypti Mosquitoes Infected with Wolbachia. *PLoS Neglected Tropical*

Diseases, *8*(2), e2688.

Gloria-Soria, A., Chiodo, T. G., & Powell, J. R. (2018). Lack of Evidence for Natural

Wolbachia Infections in Aedes aegypti (Diptera: Culicidae). *Journal of Medical*

Entomology, *55*(5), 1354–1356.

Hawkey, J., Hamidian, M., Wick, R. R., Edwards, D. J., Billman-Jacobe, H., Hall, R. M., &

Holt, K. E. (2015). ISMapper: identifying transposase insertion sites in bacterial

genomes from short read sequence data. *BMC Genomics*, *16*(1), 667.

Hien, N. T., Anh, D. D., Le, N. H., Yen, N. T., Phong, T. V., Nam, V. S., Duong, T. N.,

Nguyen, N. B., Huong, D. T. T., Hung, L. Q., Trinh, C. N. T., Hoang, N. V., Mai, V.

Q., Nghia, L. T., Dong, N. T., Tho, L. H., Kutcher, S., Hurst, T. P., Montgomery, J.

L., … Ryan, P. A. (2021). Environmental factors influence the local establishment of

Wolbachia in Aedes aegypti mosquitoes in two small communities in central

Vietnam. *Gates Open Research*, *5*, 147.

Hoffmann, A. A., Montgomery, B. L., Popovici, J., Iturbe-Ormaetxe, I., Johnson, P. H.,

Muzzi, F., Greenfield, M., Durkan, M., Leong, Y. S., Dong, Y., Cook, H., Axford, J.,

Callahan, A. G., Kenny, N., Omodei, C., McGraw, E. A., Ryan, P. A., Ritchie, S. A.,

Turelli, M., & O'Neill, S. L. (2011). Successful establishment of Wolbachia in Aedes

populations to suppress dengue transmission. *Nature*, *476*(7361), 454.

Hoffmann, Ary A., Iturbe-Ormaetxe, I., Callahan, A. G., Phillips, B. L., Billington, K.,

Axford, J. K., Montgomery, B., Turley, A. P., & O'Neill, S. L. (2014). Stability of the

wMel Wolbachia Infection following Invasion into Aedes aegypti Populations. *PLoS*

Neglected Tropical Diseases, *8*(9), e3115.

Huang, B., Yang, Q., Hoffmann, A. A., Ritchie, S. A., van den Hurk, A. F., & Warrilow, D.

(2020). Wolbachia genome stability and mtDNA variants in Aedes aegypti field

populations eight years after release. *IScience*, 101572.

Koh, C., Audsley, M. D., Di Giallonardo, F., Kerton, E. J., Young, P. R., Holmes, E. C., &

McGraw, E. A. (2019). Sustained Wolbachia-mediated blocking of dengue virus

isolates following serial passage in Aedes aegypti cell culture. *Virus Evolution*, *5*(1),

vez012.

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*

Methods, *9*(4), 357–359.

Lau, M.-J., Schmidt, T. L., Yang, Q., Chung, J., Sankey, L., Ross, P. A., & Hoffmann, A. A.

(2021). Genetic stability of Aedes aegypti populations following invasion by wMel

Wolbachia. *BMC Genomics*, *22*(1), 894.

Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association

mapping and population genetical parameter estimation from sequencing data.

Bioinformatics , *27*(21), 2987–2993.

Martinez, J., Bruner-Montero, G., Arunkumar, R., Smith, S. C. L., Day, J. P., Longdon, B., &

Jiggins, F. M. (2019). Virus evolution in Wolbachia-infected Drosophila.

Proceedings. Biological Sciences / The Royal Society, *286*(1914), 20192117.

Moran, N. A. (1996). Accelerated evolution and Muller's rachet in endosymbiotic bacteria.

Proceedings of the National Academy of Sciences, *93*(7), 2873–2878.

O'Neill, S. L., Ryan, P. A., Turley, A. P., Wilson, G., Retzki, K., Iturbe-Ormaetxe, I., Dong,

Y., Kenny, N., Paton, C. J., Ritchie, S. A., Brown-Kenyon, J., Stanford, D.,

Wittmeier, N., Jewell, N. P., Tanamas, S. K., Anders, K. L., & Simmons, C. P.
(2018). Scaled deployment of Wolbachia to protect the community from dengue and
other Aedes transmitted arboviruses. *Gates Open Research*, *2*, 36.

Pinto, S. B., Riback, T. I. S., Sylvestre, G., Costa, G., Peixoto, J., Dias, F. B. S., Tanamas, S.
K., Simmons, C. P., Dufault, S. M., Ryan, P. A., O'Neill, S. L., Muzzi, F. C., Kutcher,
S., Montgomery, J., Green, B. R., Smithyman, R., Eppinghaus, A., Saraceni, V.,
Durovni, B., … Moreira, L. A. (2021). Effectiveness of Wolbachia-infected mosquito
deployments in reducing the incidence of dengue and other Aedes-borne diseases in
Niterói, Brazil: a quasi-experimental study. *MedRxiv*, 2021.01.31.21250726.

Richardson, M. F., Weinert, L. A., Welch, J. J., Linheiro, R. S., Magwire, M. M., Jiggins, F.
M., & Bergman, C. M. (2012). Population Genomics of the Wolbachia Endosymbiont
in Drosophila melanogaster. *PLoS Genetics*, *8*(12), e1003129.

Ritchie, S. A., van den Hurk, A. F., Smout, M. J., Staunton, K. M., & Hoffmann, A. A.
(2018). Mission Accomplished? We Need a Guide to the 'Post Release' World of
Wolbachia for Aedes-borne Disease Control. *Trends in Parasitology*, *34*(Bull. World
Health Organ. 95 2017), 217–226.

Rocha, M. N., Duarte, M. M., Mansur, S. B., Silva, B. D. M., Pereira, T. N., Adelino, T. É.
R., Giovanetti, M., Alcantara, L. C. J., Santos, F. M., de Melo Costa, V. R., Teixeira,
M. M., de Melo Iani, F. C., Costa, V. V., & Moreira, L. A. (2019). Pluripotency of
Wolbachia against Arbovirus: the case of yellow fever. *Gates Open Research*, *3*(161),
161.

Ross, P. A., Ritchie, S. A., Axford, J. K., & Hoffmann, A. A. (2019). Loss of cytoplasmic
incompatibility in Wolbachia-infected Aedes aegypti under field conditions. *PLoS
Neglected Tropical Diseases*, *13*(4), e0007357.

Ross, P. A., Robinson, K. L., Yang, Q., Callahan, A. G., Schmidt, T. L., Axford, J. K.,
Coquilleau, M. P., Staunton, K. M., Townsend, M., Ritchie, S. A., Lau, M.-J., Gu, X.,
& Hoffmann, A. A. (2022). A decade of stability for wMel Wolbachia in natural
Aedes aegypti populations. *PLoS Pathogens*, *18*(2), e1010256.

Ryan, D. P., & Owen-Hughes, T. (2011). Snf2-family proteins: chromatin remodellers for
any occasion. *Current Opinion in Chemical Biology*, *15*(5), 649–656.

Ryan, P. A., Turley, A. P., Wilson, G., Hurst, T. P., Retzki, K., Brown-Kenyon, J., Hodgson,
L., Kenny, N., Cook, H., Montgomery, B. L., Paton, C. J., Ritchie, S. A., Hoffmann,
A. A., Jewell, N. P., Tanamas, S. K., Anders, K. L., Simmons, C. P., & O'Neill, S. L.
(2019). Establishment of wMel Wolbachia in Aedes aegypti mosquitoes and reduction

of local dengue transmission in Cairns and surrounding locations in northern

Queensland, Australia. *Gates Open Research*, *3*, 1547.

Siguier, P., Perochon, J., Lestrade, L., Mahillon, J., & Chandler, M. (2006). ISfinder: the

reference centre for bacterial insertion sequences. *Nucleic Acids Research*, *34*(90001),

D32–D36.

Tan, C. H., Wong, P. J., Li, M. I., Yang, H., Ng, L. C., & O'Neill, S. L. (2017). wMel limits

zika and chikungunya virus infection in a Singapore Wolbachia-introgressed Ae.

aegypti strain, wMel-Sg. *PLoS Neglected Tropical Diseases*, *11*(5), e0005496.

Tantowijoyo, W., Andari, B., Arguni, E., Budiwati, N., Nurhayati, I., Fitriana, I., Ernesia, I.,

Daniwijaya, E. W., Supriyati, E., Yusdiana, D. H., Victorius, M., Wardana, D. S.,

Ardiansyah, H., Ahmad, R. A., Ryan, P. A., Simmons, C. P., Hoffmann, A. A.,

Rancès, E., Turley, A. P., … O'Neill, S. L. (2020). Stable establishment of wMel

Wolbachia in Aedes aegypti populations in Yogyakarta, Indonesia. *PLoS Neglected

Tropical Diseases*, *14*(4), e0008157.

Thi Hue Kien, D., Edenborough, K., da Silva Goncalves, D., Thuy Vi, T., Casagrande, E.,

Thi Le Duyen, H., Thi Long, V., Thi Dui, L., Thi Tuyet Nhu, V., Thi Giang, N., Thi

Xuan Trang, H., Lee, E., Donovan-Banfield, I., Thi Thuy Van, H., Minh Nguyet, N.,

Thanh Phong, N., Van Vinh Chau, N., Wills, B., Yacoub, S., … Simmons, C. (2023).

Genome evolution of dengue virus serotype 1 under selection by Wolbachia pipientis

in Aedes aegypti mosquitoes. *Virus Evolution*, *9*(1), vead016.

Utarini, A., Indriani, C., Ahmad, R. A., Tantowijoyo, W., Arguni, E., Ansari, M. R.,

Supriyati, E., Wardana, D. S., Meitika, Y., Ernesia, I., Nurhayati, I., Prabowo, E.,

Andari, B., Green, B. R., Hodgson, L., Cutcher, Z., Rancès, E., Ryan, P. A., O'Neill,

S. L., … AWED Study Group. (2021). Efficacy of Wolbachia-Infected Mosquito

Deployments for the Control of Dengue. *The New England Journal of Medicine*,

384(23), 2177–2186.

van den Hurk, A. F., Hall-Mendelin, S., Pyke, A. T., Frentiu, F. D., McElroy, K., Day, A.,

Higgs, S., & O'Neill, S. L. (2012). Impact of Wolbachia on Infection with

Chikungunya and Yellow Fever Viruses in the Mosquito Vector Aedes aegypti. *PLoS

Neglected Tropical Diseases*, *6*(11), e1892.

Varani, A. M., Siguier, P., Gourbeyre, E., Charneau, V., & Chandler, M. (2011). ISsaga is an

ensemble of web-based methods for high throughput identification and semi-

automatic annotation of insertion sequences in prokaryotic genomes. *Genome

Biology*, *12*(3), R30.

Velez, Ivan Dario, Tanamas, S. K., Arbelaez, M. P., Kutcher, S. C., Duque, S. L., Uribe, A., Zuluaga, L., Martínez, L., Patiño, A. C., Barajas, J., Muñoz, E., Mejia Torres, M. C., Uribe, S., Porras, S., Almanza, R., Pulido, H., O'Neill, S. L., Santacruz-Sanmartin, E., Gonzalez, S., … Anders, K. L. (2023). Reduced dengue incidence following city-wide wMel Wolbachia mosquito releases throughout three Colombian cities: Interrupted time series analysis and a prospective case-control study. *PLoS Neglected Tropical Diseases*, *17*(11), e0011713.

Velez, Iván Darío, Uribe, A., Barajas, J., Uribe, S., Ángel, S., Suaza-Vasco, J. D., Mejia Torres, M. C., Arbeláez, M. P., Santacruz-Sanmartin, E., Duque, L., Martínez, L., Posada, T., Patiño, A. C., Gonzalez, S. M., Velez, A. L., Ramírez, J., Salazar, M., Gómez, S., Osorio, J. E., … O'Neill, S. L. (2023). Large-scale releases and establishment of wMel Wolbachia in Aedes aegypti mosquitoes throughout the Cities of Bello, Medellín and Itagüí, Colombia. *PLoS Neglected Tropical Diseases*, *17*(11), e0011642.

Walker, T., Johnson, P. H., Moreira, L. A., Iturbe-Ormaetxe, I., Frentiu, F. D., McMeniman, C. J., Leong, Y. S., Dong, Y., Axford, J., Kriesner, P., Lloyd, A. L., Ritchie, S. A., O'Neill, S. L., & Hoffmann, A. A. (2011). The wMel Wolbachia strain blocks dengue and invades caged Aedes aegypti populations. *Nature*, *476*(7361), 450–453.

Wernegreen, J. J. (2002). Genome evolution in bacterial endosymbionts of insects. *Nature Reviews. Genetics*, *3*(11), 850–861.

Wilm, A., Aw, P. P. K., Bertrand, D., Yeo, G. H. T., Ong, S. H., Wong, C. H., Khor, C. C., Petric, R., Hibberd, M. L., & Nagarajan, N. (2012). LoFreq: a sequence-quality aware, ultra-sensitive variant caller for uncovering cell-population heterogeneity from high-throughput sequencing datasets. *Nucleic Acids Research*, *40*(22), 11189–11201.

World Mosquito Program. (n.d.). *World Mosquito Program*. World Mosquito Program. Retrieved January 23, 2023, from https://www.worldmosquitoprogram.org/

Wu, M., Sun, L. V., Vamathevan, J., Riegler, M., Deboy, R., Brownlie, J. C., McGraw, E. A., Martin, W., Esser, C., Ahmadinejad, N., Wiegand, C., Madupu, R., Beanan, M. J., Brinkac, L. M., Daugherty, S. C., Durkin, A. S., Kolonay, J. F., Nelson, W. C., Mohamoud, Y., … Eisen, J. A. (2004). Phylogenomics of the Reproductive Parasite Wolbachia pipientis wMel: A Streamlined Genome Overrun by Mobile Genetic Elements. *PLoS Biology*, *2*(3), e69.

623