Lithium partially rescues gene expression and enhancer activity from heterozygous knockout of *AKAP11* while inducing novel differential changes

Nargess Farhangdoost[1,2], Calwing Liao[3,4,5], Yumin Liu[2,6], Daniel Rochefort[2,6], Alessia Pietrantonio[2,6], Martin Alda[7], Patrick A. Dion[2,6], Boris Chaumette[8,9*], Anouar Khayachi[2,6*], and Guy A. Rouleau[1,2,6*]

Department of Human Genetics, McGill University, Montreal, Quebec, Canada
Montreal Neurological Institute-Hospital, Montreal, Quebec, Canada
Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA
Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA
Department of Medicine, Harvard Medical School, Boston, MA, USA
Department of Neurology & Neurosurgery, McGill University, Montreal, Quebec, Canada
Department of Psychiatry, Dalhousie University, Halifax, Nova Scotia, Canada
Université Paris Cité, Institute of Psychiatry and Neuroscience of Paris (INSERM U1266), Institut Pasteur (CNRS UMR3571), GHU Paris Psychiatrie et Neurosciences, Paris, France
Department of Psychiatry, McGill University, Montreal, Quebec, Canada

* Correspondence: guy.rouleau@mcgill.ca, anouar.khayachi@mcgill.ca, boris.chaumette@inserm.fr

Abstract

Bipolar disorder (BD) is a complex psychiatric condition usually requiring long-term treatment. Lithium (Li) remains the most effective mood stabilizer for BD, yet it benefits only a subset of patients, and its precise mechanism of action remains elusive. Exome sequencing has identified *AKAP11* (A-kinase anchoring protein 11) as a shared risk gene for BD and schizophrenia (SCZ). Given that both the AKAP11-Protein Kinase A (PKA) complex and Li target and inhibit Glycogen Synthase Kinase-3 beta (GSK3B), we hypothesize that Li may partially normalize the transcriptomic and/or epigenomic alterations observed in heterozygous *AKAP11*-Knockout (Het-*AKAP11*-KO) iPSC-derived neurons. In this study, we employed genome-wide approaches to assess the effects of Li on the transcriptome and epigenome of human iPSC-derived Het-*AKAP11*-KO neuronal culture. We show that chronic Li treatment in this cellular model upregulates key pathways that are downregulated by Heterozugous AKAP11 knockout, some of which have also been reported to be downregulated in synapses of BD and SCZ post-mortem brain tissues. Moreover, we demonstrated that Li treatment partially rescues certain transcriptomic alterations resulting from Het-*AKAP11*-KO, bringing them closer to the WT state. We suggest two possible mechanisms underlying these transcriptomic effects: (1) Li modulates histone H3K27ac levels at intergenic and intronic enhancers, influencing enhancer activity and transcription factor binding, and (2) Li enhances GSK3B serine 9 phosphorylation, impacting WNT/β-catenin signaling and downstream transcription. These findings underscore Li's potential as a therapeutic agent for BD and SCZ patients carrying *AKAP11* loss-of-function variants or exhibiting similar pathway alterations to those observed in Het-*AKAP11*-KO models.

Introduction

BD is a severe psychiatric disorder characterized by recurrent episodes of mania or hypomania and depression[1,2]. It affects about 2.4% of the global population and is linked to reduced functioning, cognitive impairment, and a diminished quality of life[3,4]. BD patients are at a high risk of suicide with about a 20-30-fold increase compared to the general population[5]. The mood stabilizer Li has been shown to significantly lower the risk of illness recurrences, as well as decrease the risk of suicide and overall mortality rates[6]. However, only about 30% of BD patients are fully responsive to Li[7]. BD is highly heritable (~70-90%[8-10]) but it is also very heterogeneous and has considerable genetic overlap with other psychiatric disorders, such as SCZ[11,12]. The largest-to-date whole-exome sequencing collaborative project for BD has identified *AKAP11* as a rare-variant large-effect risk gene shared with SCZ, showing an excess of rare protein-truncating variants (PTVs) in *AKAP11* in BD and SCZ cases compared to controls (combined odds ratio [OR] = 7.06, p = 2.83 × 10E−9)[13]. The *AKAP11* gene is under evolutionary constraint (LOEUF = 0.32, pLI =1 on gnomAD v4.0) and encodes AKAP11 protein, a membrane-associated and vesicular anchoring protein that is highly expressed in brain tissue[14-16]. The AKAP11 protein is known to bind to the regulatory subunits of cAMP-dependent PKA, directing it to different substrates for phosphorylation[17]. Specifically, AKAP11 functions as a scaffold protein to facilitate the phosphorylation of GSK3B, a key enzyme of the Wnt/β-catenin pathway[18], via PKA[15,19]. Notably, GSK3B is a target of Li--the gold standard treatment for stabilizing the mood in BD patients[20-24].

Previously, our team investigated the impact of heterozygous *AKAP11* loss-of-function (LoF) mutations on the gene expression landscape and profiled the epigenomic modifications, including histone H3 Lysine 27 acetylation, a marker of active enhancers[25-28]. We found widespread transcriptional dysregulation of genes primarily involved in DNA-binding transcription factor activity, actin binding and cytoskeleton regulation, as well as cytokine receptor binding functions. We specifically highlighted the role of aberrant intergenic and intronic enhancer activity in driving these gene expression changes[28]. Furthermore, we highlighted the significant downregulation of pathways related to ribosome structure and function, a pathway also altered in synapse proteomics from BD and SCZ post-mortem brain tissues, as well as heterozygous *Akap11*-KO mice[28,29]. However, the impact of Li on such transcriptomic and epigenomic changes is not yet understood. Given the known role of AKAP11-PKA protein complex as well as Li in inhibiting GSK3B through increasing serine 9 (S9) phosphorylation levels[15,19,30,31], we postulated that further treating the Het-*AKAP11*-KO human iPSC-derived neurons with Li might reverse some of the transcriptomic and/or epigenomic modifications previously observed in Het-*AKAP11*-KO compared to wild-type (WT) neurons. In fact, Palmer et al.[13] reported that, out of 11 BD cases carrying *AKAP11* PTVs with available clinical data on Li response, 7 were good responders, while 4 were not[13]. Notably, they showed that the percentage of good responders to Li among *AKAP11* PTV carriers (63.6%) was slightly higher than the overall response rate in BD patients in their study (52%)[13]. Although the sample size is too small to draw firm conclusions regarding *AKAP11* PTV carriers' response to Li, these findings further emphasize the importance of investigating the direct molecular impact of Li in Het-*AKAP11*-

KO models. Such research is a critical step toward identifying effective therapeutic strategies for this genetically and clinically heterogeneous disorder.

In this study, using RNA sequencing (RNA-seq), H3K27ac chromatin immunoprecipitation followed by sequencing (ChIP-seq), we first investigate the direct transcriptomic and epigenomics effects of Li treatment in Het-*AKAP11*-KO human iPSC-derived neurons compared to the untreated state. Subsequently, we compare the gene expression and intergenic and intronic H3K27ac profile of Het-*AKAP11*-KO-Li-treated (Het-*AKAP11*-KO-Li) vs. WT-untreated (WT-UNT) with Het-*AKAP11*-KO-untreated (Het-*AKAP11*-KO-UNT) vs. WT-untreated (WT-UNT) to uncover the transcriptomics and epigenomic effects of Li, including any rescue or novel changes, relative to previous findings with untreated Het-*AKAP11*-KO vs. WT[28]. We show that chronic Li treatment of Het-*AKAP11*-KO iPSC-derived neurons results in a limited number of differentially expressed genes (DEGs), most notably the downregulation of the glutamate ionotropic receptor N-methyl-d-aspartate (NMDA) type subunit 2A (*GRIN2A*). This gene has been identified as a candidate in a genome-wide association study (GWAS) investigating the lithium response in BD. Moreover, it has been associated with BD in a GWAS[32] and is also a high-confidence SCZ risk gene identified in the largest schizophrenia sequencing study (OR: 24.1, P = 7.18 x 10-06)[33]. We next show that Li not only rescues some of the transcriptomic consequences of the heterozygous knockout of *AKAP11* but also induces new DEGs previously absent in Het-*AKAP11*-KO vs. WT. Finally, we noted two potential mechanisms via which Li may rescue or induce those DEGs and impact the transcriptome of Het-*AKAP11*-KO vs. WT: 1) by modifying H3K27ac levels at intergenic and intronic enhancers, influencing enhancer activity and transcription factor binding strength compared to the WT system; 2) by increasing the GSK3B S9 phosphorylation, thereby impacting WNT/β-catenin signaling pathway and the downstream genes regulated by this pathway.

By systematically comparing these different conditions, we pave the way for unraveling the complex transcriptomic and epigenomic impacts of chronic Li treatment in the Het-*AKAP11*-KO human iPSC-derived neuronal model. This approach sheds light on the underlying biological mechanisms and potential therapeutic opportunities for *AKAP11* LoF variant carriers, and possibly other BD or SCZ cohorts with similarly disrupted pathways as seen in Het-*AKAP11*-KO models.

Results

The effect of Lithium on the gene expression profile of Heterozygous *AKAP11*-KO iPSC-derived neurons

After chronically treating the 3 Het-*AKAP11*-KO clones chronically with Li (1.5 mM, 11 days), we first examined the direct gene expression modifications resulting from Li treatment in the Het-*AKAP11*-KO model (Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT). We profiled the differentially expressed genes (DEGs) at adjusted P- value (p.adj) < 0.05 and | log2FC| > 0.25 and detected 33 total DEGS: 28 downregulated genes and 5 upregulated genes (Fig. 1A, B; Supplementary Table 1). Using a Z-score hierarchical clustering heatmap of DEGs, we determined the distinct averaged gene expression patterns of Het-*AKAP11*-KO clones with and without Li treatment (Fig. 1C; Supplementary Fig. 1C). Standing out among the DEGs was downregulation of *GRIN2A*, which encodes the GluN2A subunit of NMDA receptors--glutamate-gated cation channels characterized by high calcium permeability[34] (Fig. 1B, C). To reveal functional differences between the Li-treated and untreated Het-*AKAP11*-KO groups and detect statistically enriched gene sets in our transcriptomic data, we carried out gene set enrichment analysis (GSEA)[35,36] using Gene Ontology(GO)[37,38] molecular function database. The top 10 most significantly enriched molecular functions were all upregulated in Het-*AKAP11*-KO-Li compared to *AKPA11*-KO-UNT (Fig. 1D). These functions include those related to RNA binding, processing, and protein synthesis (RNA binding, catalytic activity acting on RNA, snRNA binding, ribonucleoprotein complex binding, sequence-specific DNA binding, transcription regulator activity), ribosome regulation (structural constituent of ribosome, including both ribosomal protein and mitochondrial ribosomal protein genes), and transcription regulation (transcription regulator activity, sequence-specific DNA binding; Fig. 1D, Supplementary Table 2). Interestingly, our group previously reported that several of these molecular functions upregulated following Li treatment in Het-*AKAP11*-KO neurons—including RNA binding, structural constituent of ribosome, sequence-specific DNA binding, and transcription regulator activity—were significantly downregulated (p.adj < 0.05) in Het-*AKAP11*-KO-UNT vs. WT-UNT gene expression GSEA[28] (Supplementary Fig. 1D, Supplementary Table 3). Specifically, pathways related to the structural constituent of ribosome, sequence-specific DNA binding, and transcription regulator activity have been also previously reported to be significantly downregulated (p.adj < 0.1) in the GSEA of synapse proteomics from post-mortem brain samples of BD and SCZ patients, compared to controls[29].

Li treatment resulted in more DEGs in the WT (WT-Li vs. WT-UNT, 87 DEGs, p.adj < 0.05 and |log2FC| > 0.25, Supplementary Fig. 1A, B; Supplementary Table 1) compared to the Het-*AKAP11*-KO neurons (Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT, 33 DEGS). Interestingly however, Li impacted the gene expression profile of the WT neurons markedly differently with only 5 DEGs shared between the Li treatment within the WT and the KO models: *ADAMTS9*, *ATF4*, *BMP8A*, *LINC00365*, and *RFLNB* (Supplementary Table 1). These distinct responses to Li may be influenced by the baseline gene expression profiles of WT and Het-*AKAP11*-KO neurons, suggesting that different active pathways are directly targeted by Li. Nonetheless, we cannot interpret the results in relation to the quality of the therapeutic response to Li, as our investigation was conducted using iPSCs derived from a clinically unaffected individual.

Fig. 1. Li-dependent transcriptomic changes in Het-*AKAP11*-KO **iPSC-derived neuronal model. A)** Number of up and downregulated DEGs in Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT (P- value < 0.05 and |log2FC| > 0.25). blue: downregulated, red: upregulated. **B)** Volcano plot of differentially expressed genes in Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT (P- value < 0.05 and |log2FC| > 0.25). Red: upregulated, blue: downregulated, grey: nonDEG. **C)** Hierarchical clustering heatmap of differentially expressed genes (rows) in Het-*AKAP11*-KO-Li clones vs. Het-*AKAP11*-KO-UNT reps (averaged). The expression levels across samples were standardized by the Z-Score method. Red shows the highest (2) and blue the lowest (-2) Z-score values. **D)** GSEA[35,36] showing the enrichment of the 10 most significant

GO[37,38] molecular function gene sets for the Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT comparison; P.adj<0.05. For the full list refer to Supplementary Table 2. The size of the circles represents gene count (count of core enrichment genes); the colors represent the strength of -log (p-adj value), with red showing the most significant p-adj value, and gene ratio represents (count of core enrichment genes) / (count of pathway genes).

Lithium treatment of Heterozygous *AKAP11*-KO iPSC-derived neurons partially rescues KO-induced DEGs and induces novel DEGs

To identify the compensatory transcriptomic effects of chronic Li treatment in Het-*AKAP11*-KO neurons relative to WT-UNT, we first obtained DEGs from Het-*AKAP11*-KO-UNT vs. WT-UNT as well as Het-*AKAP11*-KO-Li vs. WT-UNT at p-adj < 0.05 and |log2FC| > 0.25 (Supplementary Fig. 2A, Supplementary Table 1). We observed an overall reduction in the number of DEGs, in both up-and downregulated directions, following chronic Li treatment of Het-*AKAP11*-KO neurons (Fig. 2A, Supplementary Fig. 2A, Supplementary Table 1). Li appeared to lessen the extent of differential gene expression that resulted from the heterozygous knockout of *AKAP11*, suggesting a partially compensating effect of Li on some of the Het-*AKAP11*-KO-induced gene expression changes. We then plotted a log2 fold change (Log2FC) hierarchical clustering heatmap showing the genes that were differentially expressed in Het-*AKAP11*-KO-UNT vs. WT-UNT but lost their DEG status upon Li treatment of the Het-*AKAP11*-KO (|log2FC difference|>0.6) hereafter referred to as Li-rescued DEGs (Fig. 2B, Supplementary Table 4). The heatmap shows that a relatively large portion of upregulated DEGs in the Het-*AKAP11*-KO vs. WT was rescued by treatment of Het-*AKAP11*-KO with Li (Fig. 2B). Among all the Li-rescued DEGs, we identified those encoding proteins involved in a variety of functions including DNA-binding transcription factor activity (*MEOX1, FOXC1, FOSL2, NKX3-2, IRX4, HOXD11, SHOX2, IRX3*), G protein-coupled receptor (GPCR) activity (*GAL, FZD10, RAMP3, SPHK1*), transmembrane transporter activity (*SLC7A8, ABCC3, SLC14A2, CHRND, GLRA1, RYR3*), excitatory extracellular ligand-gated ion channel activity (*CHRND, GLRA1*), growth factor binding (*FGFR3, DUSP1, FGFBP1, FGFBP2*), and oxidoreductase activity (*BCO1, HPD*) (Supplementary Fig. 2B). The most significantly enriched GO[37,38] molecular function was glycosaminoglycan binding (*VTN, SMOC2, FGFBP1, HAPLN1, EGFLAM*; enrichment FDR=5.86E-02; Fig. 2D, Supplementary Fig. 2B). Subsequently, we showed that Li treatment of the Het-*AKAP11*-KO iPSC-derived neurons resulted in differential expression of a group of genes that were not differentially expressed in the Het-*AKAP11*-KO-UNT vs. WT-UNT comparison (|log2FC difference|> 0.6), hereafter referred to as Li-induced DEGs (Fig. 2C, Supplementary Table 4). Most of these genes went from being non-DEG in Het-*AKAP11*-KO-UNT vs WT-UNT to being differentially downregulated upon treatment of Het-*AKAP11*-KO with Li. ORA using the GO[37,38] molecular function database (FDR < 0.1) revealed enrichment of extracellular matrix (ECM) structural constituent (*TNC, TFPI2, FMOD, LUM*), signaling receptor regulator activity (*FGF10, ANGPT4, WNT3A, NPPC, CALCB*), calcium channel activity (*TRPC7, OPRM1, TRPM2*), and metal ion transmembrane transporter activity (*TRPC7, SLC6A4, OPRM1, TRPM2, KCNJ16*) within these genes (Fig. 2E).

It is important to emphasize the presence of the key WNT/β-catenin-pathway genes, such as *FZD10* and *WNT3A* (Fig. 2B, C), as well as the enrichment of β-catenin-regulated TCF4, LEF1, and PITX2 binding sites within the Li-responsive genes (Li-rescued and-induced DEGs). The latter was determined by ORA using the Transcription Factor Target Genome Browser Position Weight Matrix database (PWMs; enrichment FDR<0.05; Fig. 2F, Supplementary Table 5). This suggests that the dysregulation of GSK3B phosphorylation at the S9 residue—targeted by both AKAP11-PKA and Li—may play a role in Li-response-related transcriptomic modifications. These modifications include partial rescuing

of gene expression changes caused by the heterozygous knockout of *AKAP11*, as well as the induction of novel gene expression changes that were previously absent. To test this hypothesis, we carried out immunoblotting of phospho-GSK3B (phGSK3B, S9) in all our samples, Li-treated and untreated, and quantified the bands (Fig. 2G, top and bottom; Supplementary Fig. 2C; Supplementary Table 6). Although none of the comparisons were statistically significant, we observed a trend: the heterozygous knockout of *AKAP11* resulted in a slightly decreased phosphorylation of GSK3B, and further treatment of the Het-*AKAP11*-KO with Li increased these levels relative to both Het-*AKAP11*-KO and the WT baseline (UNT) (Fig. 2G, top).

Fig. 2. The differential gene expression rescuing and inducing effects of Li treatment in Het-*AKAP11*-KO **relative to WT-UNT iPSC-derived neurons. A**) Number of up and downregulated DEGs in Het-*AKAP11*-KO-UNT vs. WT-UNT and Het-*AKAP11*-KO-Li vs. WT-UNT. P.adj < 0.05 and |log2FC| > 0.25. Red: upregulated, blue: downregulated. **B**) Log2FC hierarchical clustering heatmap demonstrating the DEGs from Het-*AKAP11*-KO-UNT vs. WT-UNT that became nonDEGs (rescued) by Li treatment of the Het-*AKAP11*-KO (Het-*AKAP11*-KO-Li vs. WT-UNT). p-adj < 0.05 and |log2FC| > 0.25; DEG cutoff: P-adj < 0.05 and |log2FC| > 0.25. A minimum log2FC difference of |0.6| between the two sets of comparisons was set as a condition for [(Het-*AKAP11*-KO-Li vs. WT-UNT) − (Het-*AKAP11*-KO-UNT vs. WT-UNT)]. **C**) Log2FC hierarchical clustering heatmap demonstrating genes that were nonDEGs in Het-*AKAP11*-KO-UNT vs. WT-UNT and became DEGs (induced) in Het-*AKAP11*-KO-Li vs. WT-UNT. p-adj < 0.05 and |log2FC| > 0.25; A minimum log2FC difference of |0.6| between the two sets of comparisons was set as a condition (Het-*AKAP11*-KO-Li vs. WT-UNT) − (Het-*AKAP11*-KO-UNT vs. WT-UNT). **D, E**) Over-representation analysis (ORA) of rescued and newly induced DEGs, respectively, following Li treatment of Het-*AKAP11*-KO, using all genes from Fig.s 2B and 2C heatmaps. GO[37,38] Molecular function database using ShinyGO 0.77[39]; FDR < 0.1. The background genes used were all the 19,568 genes that were considered expressed (see Methods). Shown here are the top

10 pathways with redundancy removed. **F**) Top 5 (based on adj p-value) Transcription Factor Target Genome Browser PWMs (FDR<0.05) enrichment results used to detect which binding sites were most significantly associated with the Li-responsive DEGs. Determined using ShinyGo 0.77[39]. For the full list of the top 20 motifs refer to Supplementary Table 5. **G**) Bottom: Immunoblot showing the levels of phGSK3B (S9) and GAPDH (loading control) in the Het-*AKAP11*-KO clones and WT replicates with and without Li treatment. Only the bands indicated with the arrows were quantified. Top: A two-way ANOVA followed by Tukey's multiple comparison test was conducted to determine the statistical significance of differences in phosphorylation levels of GSK3B among the Het-*AKAP11*-KO and WT groups. Data are represented as mean ± SEM. Statistical significance is indicated by NS (not significant), *p<0.05, **p<0.01, ***p<0.001.

Impact of lithium on differential intergenic and intronic H3K27ac modifications resulting from heterozygous knockout of *AKAP11* relative to WT: implications for regulation of lithium-rescued and induced gene expression changes

We next aimed to determine whether modifications in the H3K27ac mark, a hallmark of active enhancers[25-27], are involved in modulating gene expression changes resulting from chronic Li treatment in the Het-AKAP11-KO system. With this purpose in mind, we first profiled the intergenic and intronic H3K27ac differential peaks, which are the primary sites of enhancers previously dysregulated with the heterozygous knockout of *AKAP11*, in Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT at p.adj < 0.05 and |log2FC| > 0.25 and detected 39 intergenic and 44 intronic differential peaks; Supplementary Fig. 3A, B; Supplementary Table 7). However, within the WT system (WT-Li vs. WT-UNT) there were no differential H3K27ac peaks with the same cutoffs of p.adj < 0.05 and |log2FC| > 0.25, highlighting that the H3K27ac modifications observed above are Het-*AKAP11*-KO-specific (Supplementary Table 8). We then associated the differential H3K27ac peaks from Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT with the nearest TSS of known genes —within the distance of 100 kb from the TSS to the peak center—and obtained a total of 28 different genes associated with intergenic and intronic differential H3K27ac peaks and marked their gene expression status (Supplementary Table 7); however, there was no overlap between them and the 33 DEGs from gene expression analysis of Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT at p.adj < 0.05 and |log2FC| > 0.25.

Next, we sought to investigate whether Li treatment rescues or further modifies the Het-*AKAP11*-KO-induced differential H3K27ac peaks in Het-*AKAP11*-KO iPSC-derived neurons relative to WT. We specifically aimed to detect modifications that aligned with Li-rescued or Li-induced DEGs, with respect to the direction of change (up/down) and the type of effect (induced/rescued). Like our gene expression data, we observed an overall reduction in the number of both differential intergenic and intronic H3K27ac peaks in Het-*AKAP11*-KO-Li vs. WT-UNT relative to Het-*AKAP11*-KO-UNT vs. WT-UNT, in both up and down directions (Fig. 3A, B, Supplementary Table 8). We specifically searched for the intergenic and intronic differential H3K27ac peaks in Het-*AKAP11*-KO vs. WT that lost their differential status (Li-rescued peaks) as well as intergenic and intronic differential H3K27ac peaks that newly appeared in Het-*AKAP11*-KO-Li vs. WT-UNT (Li-induced peaks; p.adj < 0.05 and |log2FC| > 0.25 for both, |log2FC difference| > 0.3; Supplementary Table 9). We then associated these Li-responsive peaks (both rescued and Li-induced peaks) with the nearest TSS of known genes within the distance of 100 kb from the TSS to the peak center. Subsequently, we integrated the H3K27ac with gene expression data and detected 8 intergenic peaks (associated with 8 genes) and 11 intronic peaks (associated with 9 genes) among the Li-rescued and Li-induced peaks that were overlapping our Li-rescued or Li-induced DEGs, respectively, with the same direction of change (Fig. 3C, D, Supplementary Table 9). In particular, one gene (*PLEKHG4B*) has both intergenic and intronic Li-rescued H3K27ac peaks. Although a few Li-induced DEGs were noted among these genes, most were Li-rescued DEGs (Fig. 3C, D). Among all these Li-induced and Li-rescued changes in enhancer activity that were linked to Li-induced and Li-rescued gene expression changes in the same direction, there were several ncRNA genes (including lncRNA, miRNA, and anti-sense RNA). In addition, we noted

several coding genes playing roles in SLC-mediated transmembrane transport (*SLC7A8, SLC14A2*) and MAPK family signaling cascades (*FGFR3, DUSP1*), calcitonin-like ligand receptors (*RAMP3*) among others (Supplementary Fig. 10). Finally, we demonstrated the gene expression and H3K27ac averaged tracks of Li-treated and -untreated Het-*AKAP11*-KO clones, as well as untreated WT replicates, using the UCSC genome browser. The example tracks highlight the gene *RAMP3* which demonstrates an intergenic differentially elevated H3K27ac peak in Het-*AKAP11*-KO vs. WT, marking increased enhancer activity and differential binding, along with upregulation of gene expression, both of which are then lowered to a non-differential state in Het-*AKAP11*-KO-Li vs. WT-UNT, and thereby rescued (Fig. 3E).

These findings suggest that one mechanism by which Li rescues some of the DEGs resulting from the heterozygous knockout of *AKAP11* or induces new DEGs is by modulating intergenic and intronic H3K27ac levels in a concordant direction. This indicates that Li may play a role in modifying enhancer activity and transcription factor binding at intergenic and intronic regions, thereby rescuing or inducing gene expression patterns compared to the WT system.

Fig. 3. Li-rescued/induced H3K27ac peaks and their association with Li-rescued/induced gene expression changes: differential H3K27ac and gene expression modifications in Het-*AKAP11*-KO-Li vs. WT-UNT and Het-*AKAP11*-KO-UNT vs. WT-UNT. **A, B**) Number of differentially elevated and decreased H3K27ac peaks in intergenic and intronic regions, respectively, in Het-*AKAP11*-KO-UNT vs. WT-UNT and Het-*AKAP11*-KO-Li vs. WT-UNT; p.adj < 0.05 and |log2FC| > 0.25. Red: differentially increased peaks, blue: differentially decreased peaks. **C, D**) Tables showing Li-rescued/induced differential H3K27ac peaks that changed in the same direction as the Li-rescued/induced DEGs, respectively, following chronic Li treatment of Het-*AKAP11*-KO (Het-*AKAP11*-KO-Li vs. WT-UNT relative to Het-*AKAP11*-KO-UNT vs. WT-UNT). For both gene expression and H3K27ac: differentially marked (P.adj < 0.05 and |log2FC| > 0.25) in one comparison but not the other (i.e., in either Het-*AKAP11*-KO-UNT vs. WT-UNT or Het-*AKAP11*-KO-Li vs. WT-UNT); |log2FC difference| > 0.3 between the two sets of comparisons [(Het-*AKAP11*-KO-Li vs. WT-UNT) − (Het-*AKAP11*-KO-UNT vs. WT-UNT)] was also set as a condition. Red circle shows a negative value of Log2FC difference as a result of (Het-*AKAP11*-KO-Li vs. WT-UNT) − (Het-*AKAP11*-KO-UNT vs. WT-UNT); green circle shows a positive value of Log2FC difference as a result of [(Het-*AKAP11*-KO-Li vs. WT-UNT) − (Het-*AKAP11*-KO-UNT vs. WT-UNT)]. Li-induced (non-differential-> differential) DEG/differential peaks are colored in orange and Li-rescued (differential->non-differential) are colored in blue. **E**) UCSC

Genome Browser H3K27ac and gene expression averaged tracks of Het-*AKAP11*-KO clones and WT replicates, Li-treated and untreated. Demonstrating *RAMP3*, as an example, with Li-rescued expression profile and an intergenic Li-rescued H3K27ac peak. Blue rectangles highlight enhancers with differential binding strength and red squares mark gene expression profiles of the genes.

Discussion

BD is a highly heritable and heterogeneous psychiatric condition, and Li remains one of the most effective mood stabilizers for its treatment[6]. Studying the molecular effects of Li in Het-*AKAP11*-KO neuronal culture could contribute to refining treatment strategies and improving outcomes for this rare LoF risk gene associated with both BD and SCZ. Given the identification of *AKAP11* as a rare PTV large-effect risk gene shared between BD and SCZ[13], and its involvement at the protein level in GSK3B regulation—a pathway influenced by Li[13,18]—in this study we: 1) used bulk RNA-seq data to first investigate how chronic Li treatment of Het-*AKAP11*-KO iPSC-derived neurons impacts the gene expression landscape within this model. Subsequently, we performed ChIP-seq of H3K27ac mark, a histone modification indicative of enhancer activity and transcription factor binding, to determine if changes in this histone mark are associated with the dysregulation of gene expression exerted by Li. 2) used bulk RNA-seq data to explore the Li-rescued and-induced differential gene expression by comparing Het-*AKAP11*-KO-UNT vs. WT-UNT DEG profile with that of Het-*AKAP11*-KO-Li vs. WT-UNT. Similarly, we then utilized the H3K27ac mark to determine the Li-rescued and -induced differential peaks and if such peaks were associated with the Li-rescued and -induced DEGs, respectively, in the same direction.

Among the transcriptomic effects of Li in the Het-*AKAP11*-KO iPSC-derived neuronal culture (Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT), we found the downregulation of *GRIN2A*, which encodes the GluN2A subunit of NMDA receptors. *GRIN2A* has been reported to be associated with BD in GWAS[32,40] and is a high-confidence LoF risk gene for schizophrenia[33,41,42]. Furthermore, it has been detected as a candidate gene in a GWAS of BD lithium response[43], suggesting a potential link between *GRIN2A* and Li treatment response in BD patients[43]. The encoded protein is involved in excitatory neurotransmission and synaptic plasticity[44,45], and its expression regulation has been proposed as one of the therapeutic effects of Li in bipolar disorder, potentially due to Li's role in intracellular calcium regulation[46]. Interestingly, we showed that Li affected the gene expression profile of Het-*AKAP11*-KO distinctly from the WT model, with *GRIN2A* expression dysregulation being specific to Li treatment in Het-*AKAP11*-KO. These findings suggest that Li's therapeutic effects in BD may be partly mediated by its ability to downregulate *GRIN2A*. This modulation may influence glutamatergic neurotransmission, which is reported to be dysregulated in BD[47,48].

The absence of H3K27ac peaks linked to differential gene expression following Li treatment in the Het-*AKAP11*-KO iPSC-derived neuronal model indicates that Li's impact on chromatin (H3K27ac) does not directly influence gene expression through enhancer activity in this context. This could imply that Li's effects on the H3K27ac landscape are independent of transcriptional changes in the genes proximal to those enhancers in the Het-*AKAP11*-KO model. Instead, these H3K27ac peaks might affect the expression of the distal target genes rather than the neighboring genes. So, the absence of such a correlation does not necessarily mean a lack of functional significance for these differential H3KK27ac peaks.

The molecular functions related to the structural constituent of ribosomes and transcription regulation following chronic Li treatment in Het-*AKAP11*-KO were previously shown to be downregulated as a result of heterozygous knockout of *AKAP11*[28]. This downregulation also aligned with certain synapse proteomics GSEA findings from post-mortem dorsolateral prefrontal cortex samples of BD and SCZ patients compared to controls[29], suggesting that Li may target and reverse key pathways commonly disrupted in Het-*AKAP11*-KO iPSC-derived neuronal cultures and in post-mortem brain tissues from BD and SCZ patients. Firstly, it is notable that there is a growing interest in studying the role of ribosome dysregulation induced by stress as a potential mechanism for neuropsychiatric disorders, particularly mood disorders[49-51]. Aberrant protein translation, possibly linked to decreased neurotransmission and synaptic plasticity[52,53], is associated with the pathophysiology of SCZ[52,53]. In addition, dysfunctional stress response in the endoplasmic reticulum, which hosts a portion of the cytoplasmic ribosomes, is associated with BD[54,55]. Both in cell and animal models, Li has been shown to reverse stress-induced aberrant protein synthesis by decreasing the phosphorylation of a key modulator of mRNA translation eukaryotic elongation factor-2 (eEF2) at its inhibitory site, thereby increasing its activity, even under stress conditions such as nutrient and serum deprivation[56]. Secondly, mitochondrial abnormalities have been linked to major psychiatric disorders[57,58]. Mitochondria are crucial for neuronal functions, particularly in energy production, cellular respiration, and calcium signaling[59]. Evidence suggests that Li enhances mitochondrial respiration, possibly by influencing the mitochondrial oxidative phosphorylation (OXPHOS) pathway[59]. Consequently, the upregulation of mitochondrial ribosomal protein genes within the "structural constituent of ribosome" function observed after chronic Li treatment in the Het-*AKAP11*-KO model, may contribute to improved mitochondrial respiration. Lastly, Li can enhance the transcription and DNA binding through different paths. One possibility is via targeting the CREB (cAMP-response-element-binding protein)-co-activator TORC (transducer of regulated CREB) followed by enhancing CREB-directed gene transcription[60]. Other paths might involve Li targeting GSK3B, inositol mono phosphatase (IMPase), and/or inositol polyphosphate-1-phosphatase (IPPase)[61,62] and affecting the downstream signaling and gene expression regulation.

Het-*AKAP11*-KO-Li vs. WT-UNT has a smaller number of DEGs than Het-*AKAP11*-KO-UNT vs. WT-UNT, under the same p.adj value and log2FC cutoffs, indicating that Li rescues some of the gene expression changes associated with the heterozygous knockout of *AKAP11*, bringing the Het-*AKAP11*-KO gene expression profile closer to that of the WT. This suggests a possible partial therapeutic or normalizing effect of Li on the Het-*AKAP11*-KO model. Chronic Li treatment of Het-*AKAP11*-KO rescued genes involved in various key roles, particularly glycosaminoglycan binding. Glycosaminoglycans are linear polysaccharides that exist on the cell surface in the ECM and covalently bind to core proteins to form proteoglycans[63]. Proteoglycans are one of the main components of the ECM. The Li-rescued genes associated with glycosaminoglycans binding function (*VTN, SMOC2, FGFBP1, HAPLN1, EGFLAM*) are primarily involved in the ECM organization, cell adhesion, and migration. The Li-induced DEGs are also involved in the function and structure of the ECM, signaling receptor regulator activity, calcium channel activity, and metal ion transmembrane transporter activity. Aberrant gene expression in ECM-related genes is noted in both Li-induced and

Li-rescued genes, implying that Li's therapeutic effects may involve complex regulation of ECM processes, which are crucial for neuronal activity, neuroplasticity, and cellular homeostasis[64]. Interestingly, alterations in ECM, such as degradation, overproduction, and altered composition have been reported in several neuropsychiatric disorders including bipolar disorder, depression, and schizophrenia[64-66].

We noted that the Li-responsive DEGs in Het-*AKAP11*-KO vs. WT, including both Li-rescued and Li-induced DEGs, were enriched in TCF4, LEF1, and PTX2 transcription factor binding sites. This suggests that one of the mechanisms by which Li may alter the expression of these genes is via its action on the GSK3B/β-catenin signaling pathway, which operates upstream of these transcription factors and is a well-established target of Li[21]. Immunoblotting of S9 GSK3B showed a pattern of a slight decrease in GSK3B's phosphorylation levels at S9, which was followed by an increase upon chronic Li treatment in the Het-*AKAP11*-KO relative to both untreated state Het-*AKAP11*-KO and WT. Although none of these changes were statistically significant, the pattern of increase in phosphorylation of S9 GSK3B upon chronic Li treatment in Het-*AKAP11*-KO confirms that Li exerts its expected effect on S9 GSK3B phosphorylation. Some of the Li-rescued/induced gene expression changes we observed could be explained by these alterations in GSK3B phosphorylation levels. There are several possible reasons why the GSK3B phosphorylation levels at S9 are not statistically significant 1) for Het-*AKAP11*-KO vs. WT: there are kinases other than AKAP11-dependant PKA that can phosphorylate GSK3B at S9 to inhibit it, e.g. Akt, protein kinase C (PKC), and p90Rsk. Thus, the impact of the heterozygous KO of *AKAP11* may have been compensated for by other mechanisms. 2) for Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT: A statistically significant increase may require a higher cellular dose of Li (we used 1.5mM, to be close to the therapeutic dose in serum concentrations). In addition, chronic Li treatment allows time for possible compensatory processes to take place to reduce the S9 phosphorylation to a level that is statistically non-significant. However, the transcriptomic modifications observed may still be a consequence of this statistically non-significant increase or a transient more elevated increase in S9 phosphorylation of GSK3B immediately after treatment, even if the significant increase is not detectable at this stage.

Lastly, we showed that chronic Li treatment of Het-*AKAP11*-KO rescues a subset of DEGs resulting from heterozygous knockout of *AKAP11* and induces DEGs previously absent in Het-*AKAP11*-KO vs. WT iPSC-derived neurons. This effect partially occurs via Li's impact on intergenic and intronic H3K27ac peaks, which mark active enhancers[25-27], driving changes in the same direction as observed in gene expression. These particularly include non-coding RNA genes (lncRNAs and miRNAs), and coding genes involved in MAPK signaling cascades (e.g., *FGFR3*, *DUSP1*). *DUSP1* encodes MAP kinase phosphatase 1 (MKP1) which belongs to a family of proteins that dephosphorylate both threonine and tyrosine residues, acting as a negative regulator of the MAPK cascade[67]—a significant signaling pathway involved in neuronal plasticity, function, and survival[68,69]. Elevated *DUSP1* mRNA expression is associated with the pathophysiology of depressive behavior[70,71]. More specifically, elevated mRNA levels of *DUSP1* have been observed in the hippocampus of both patients with depression and stressed rats[70]. Overall, our findings suggest that Li can modify a subset of the gene expression changes that occur because of heterozygous knockout of *AKAP11* by altering

H3K27ac levels at specific intergenic and intronic enhancers, thereby influencing enhancer activity and transcription factor binding affinity.

Altogether, our findings increase our understanding of how Li may act as a therapeutic, particularly for individuals with BD and/or SCZ who carry *AKAP11* LoF variants. Additionally, Li may also benefit BD and SCZ patients who do not have this specific rare variant but exhibit alterations of similar pathways resulting from the heterozygous knockout of *AKAP11*, such as those related to ribosomes, proteoglycans, and ECM dynamics. These pathways were shown here to be influenced by Li and may be also critical to the phenotype. Overall, these findings highlight the complexity of gene expression regulation in response to Li treatment and emphasize that a specific genetic variation, such as heterozygous LoF mutations in *AKAP11*, may affect therapeutic outcomes. Although Li is not generally recommended for SCZ, this study opens the door to exploring its efficacy in a subset of patients based on their genetic background. Further research with larger sample sizes, greater genetic diversity, and functional studies is needed to deepen our understanding of heterozygous *AKAP11* knockout in cell and animal models, as well as the associated response to Li in the context of BD and SCZ.

Materials / Subjects and Methods

Ethics statement

The cell line and protocols in the present study were used in accordance with guidelines approved by the institutional human ethics committee guidelines and the Nova Scotia Health Authority Research Ethics Board (REB # 1020604).

RNP-CRISPR-Cas9 heterozygous knockout of *AKAP11* in induced pluripotent stem cells

The iPSCs used in this study were reprogrammed from the blood sample of a consenting individual as described previously (informed consent was obtained) [72]. The protocol for the generation of Het-*AKAP11*-KO s with the RNP-CRISPR-Cas9 technique and the validation of the KOs using RT-qPCR are described in our previous work [28]. The same WT replicates (SBP009, male, age 53, Caucasian; Table 1) and Het-*AKAP11*-KO clones (Het-AKAP11-KO-Clone 21: 4 bp insertion CAGT, Het-AKAP11-KO-Clone 16: 11bp deletion ACACCTTGGAC, Het-AKAP11-KO-Clone 20: 10 bp deletion GGTACACCTT) as our previous study [28] are utilized here. Het-*AKAP11*-KO*s*: 3 clones each carrying a different frameshift mutation, all derived from SBP009; WT: 3 SBP009 replicates, each with a different passage number, cultured and differentiated to neurons separately.

Sample	Source	Reprogramming	Identifier	Donor information
Induced Pluripotent Stem Cell (iPSC)	Human lymphocytes	Reprogrammed into iPSC by Sanford Burnham, as described previously[72].	SBP009	Control, Unaffected individual, Male, 51, Caucasian

Table 1. iPSC Sample Information.

iPSC and NPC differentiation and maintenance and neuronal cell culture

iPSCs were maintained on a Matrigel (Corning, Cat. #08-774-552) coated plate with mTeSR™1 medium (STEMCELL Technologies, Cat. #85870) along with Y-27632 (ROCK inhibitor, STEMCELL Technologies, Cat. # 72307) after each passage. During the course of induction of iPSC to NPCs (20 days), STEMdiff™ SMADi Neural Induction Kit (STEMCELL Technologies, Cat. #08581) was used. The method for induction of iPSC to NPCs was based on the STEMCELL™ neural induction EB-based method and was performed following the manufacturer's protocol. On Day 0, a single-cell suspension of iPSC was generated using Gentle Cell Dissociation Reagent (STEMCELL Technologies, Cat. # 100-0485), and 10,000 cells per microwell were seeded in an ultralow attachment 96-well plate to generate Embryoid bodies (EBs), cultured in STEMdiff™ Neural Induction Medium + SMADi (STEMdiff™ SMADi Neural Induction Kit, Cat. # 08582) + 10 µM Y-27632. From Day 1 to Day 4, a daily partial (3/4) medium change was performed using STEMdiff™ Neural Induction Medium + SMADi. On Day 5, EBs were harvested using a wild-board 1ml serological pipettes and 40µm strainer and transferred to a single well of a 6-well plate coated with Poly-L-ornithine hydrobromide (PLO) and

laminin. A daily full medium change was performed from Day 6 to Day 11. After the neural induction efficiency was determined higher than 75%, neural rosettes were manually selected using STEMdiff™ Neural Rosette Selection Reagent (STEMCELL Technologies, Cat. # 05832) on Day 12 and replated onto a single well of PLO/Laminin coated 6-well plate. With continuous daily full medium change, selected rosette-containing clusters attached, and NPC outgrowths formed a monolayer between the clusters. The NPCs were ready for passage 1 when cultures were approximately 80- 90% confluent (typically on Day 19). NPCs were maintained and expanded (7-10 days) using DMEM/F12 with Glutamax (ThermoFisher, Cat. #10565042), N2 1x (ThermoFisher, Cat. # 17502001), B27 1x (ThermoFisher, Cat. #17504001), FGF2 20 ng/ml (PEPROTECH, Cat. #100-18B), and EGF 20 ng/ml (Gibco, Cat. # AF-100-15-1MG). For the final differentiation of NPCs into neurons, NPCs were detached using Accutase (STEMCELL Technologies, Cat. #7922) and then differentiated onto PLO/laminin-coated plates with final differentiation media of BrainPhys™ Neuronal Medium (STEMCELL Technologies, Cat. #05790) supplemented with Glutamax 1x (ThermoFisher, Cat. #35050061), N2 1x (ThermoFisher, Cat. #17502001), B27 1x (ThermoFisher, Cat. #17504001), ascorbic acid 200nM (STEMCELL Technologies, Cat. #72132), cyclic AMP 500 µg/ml (Sigma-Aldrich, Cat #A9501-1G), brain-derived neurotrophic factor 20 ng/ml (BDNF, GIBCO, Cat. #PHC7074), Wnt3a 10 ng/ml (R&D Systems, Cat. #5036-WN-500), and laminin 1 µg/ml (Gibco, Cat. #23017015) for 14 days (medium change of 3 times per week). Subsequently, the cells were maintained in STEMdiff™ Forebrain Neuron Maturation Kit for 11 days (STEMCELL Technologies, Cat. #08605) in the same dishware with media change occurring every 4 days. At this stage and in parallel, a subset of the Het-*AKAP11*-KO dishes was treated with 1.5 mM Li included in the STEMdiff™ Forebrain Neuron Maturation Kit media. They were also maintained in the Li-containing media for 11 days, to mimic chronic Li treatment, with media change every 4 days. The culture composition was previously shown to be predominantly neuronal cells with glial cells in lower proportions[28].

RNA-seq library Preparation and Sequencing

RNA was extracted from bulk neuronal culture 25 days post-differentiation using RNeasy Mini Kit (Cat. #74104). Library preparation from total RNA was performed using NEB rRNA-depleted (HMR) stranded library preparation kit according to the manufacturer's instructions and sequencing was carried out using Illumina NovaSeq 6000 (100 bp paired-end).

Immunoblotting

Cells were homogenized in lysis buffer (10 mM Tris–HCl pH7.5, 10 mM EDTA, 150 mM NaCl, 1% Triton X100, 0.1% SDS) in the presence of a mammalian protease inhibitor cocktail (Sigma, Cat #P8340) and phosphoSTOP (Sigma, Cat. #4906845001) to protect proteins from dephosphorylation. Protein extracts (40µg) were resolved by SDS–PAGE, transferred onto PVDF membrane (Millipore IPFL00010), and immunoblotted with the primary antibody: rabbit polyclonal phospho-GSK-3 (Ser9; 1:1000, Cell Signaling Technology, Cat# 9336, RRID: AB_331405). Standard GAPDH loading controls were included using a mouse monoclonal anti-GAPDH antibody (1:1000, Novus, Cat. #NB300-221,

RRID: AB_10077627). Subsequently, the membrane was revealed using the appropriate LI-COR fluorophore-conjugated secondary antibodies. Images were acquired using a LI-COR Odyssey Infrared image system. Fluorescence intensity values for each protein of interest were normalized to GAPDH signal from the same gel. Band quantification was carried out using Fiji[73] software. Full-size blots for cropped gels can be found in the supplementary material. The two-way ANOVA followed by Tukey's multiple comparison test was performed using GraphPad Prism version 10.2.3 (403) for Windows, GraphPad Software; www.graphpad.com.

Cross-linking, ChIP-seq library preparation, and sequencing

About 10 million neuronal cells per sample (3 Het-*AKAP11*-KO clones and 3 WT replicates) were grown and directly crosslinked on the plate with 1% formaldehyde (Sigma) for 10 minutes at room temperature and the reaction was stopped using 125nM Glycine for 5 minutes. Fixed cell preparations were washed with ice-cold PBS, scraped off the plate, pelleted, washed twice again in ice-cold PBS, and the flash-frozen pellets were stored at −80°C. Thawed pellets were resuspended in 500ul cell lysis buffer (5 mM PIPES-pH 8.5, 85 mM KCl, 1% (v/v) IGEPAL CA-630, 50 mM NaF, 1 mM PMSF, 1 mM Phenylarsine Oxide, 5 mM Sodium Orthovanadate, EDTA-free Protease Inhibitor tablet) and incubated 30 minutes on ice. Samples were centrifugated and pellets resuspended in 500ul of nuclei lysis buffer (50 mM Tris-HCl pH 8.0, 10 mM EDTA, 1% (w/v) SDS, 50 mM NaF, 1 mM PMSF, 1 mM Phenylarsine Oxide, 5 mM Sodium Orthovanadate and EDTA-free protease inhibitor tablet) and incubated 30 minutes on ice. Sonication of lysed nuclei was performed on a BioRuptor UCD-300 at max intensity for 45 cycles, 10 s on 20 s off, centrifuged every 15 cycles, and chilled by a 4°C water cooler. Samples were checked for sonication efficiency using the criteria of 150–500bp by gel electrophoresis of a reversed cross-linked and purified aliquot. After the sonication, the chromatin was diluted to reduce the SDS level to 0.1% and concentrated using Nanosep 10k OMEGA (Pall). ChIP reaction for histone modifications was performed on a Diagenode SX-8G IP-Star Compact using Diagenode automated Ideal ChIP-seq Kit for Histones. Dynabeads Protein A (Invitrogen) were washed, then incubated with specific antibodies (rabbit polyclonal anti-H3K27ac Diagenode Cat. #C15410196, RRID: AB_2637079), 1 million cells of sonicated cell lysate, and protease inhibitors for 10 hr, followed by 20 min wash cycle using the provided wash buffers (Diagenode Immunoprecipitation Buffers, iDeal ChIP-seq kit for Histone). Reverse cross-linking took place on a heat block at 65°C for 4 hr. ChIP samples were then treated with 2ul RNase Cocktail at 65°C for 30 min followed by 2ul Proteinase K at 65°C for 30 min. Samples were then purified with QIAGEN MinElute PCR purification kit (QIAGEN) as per manufacturers' protocol. In parallel, input samples (chromatin from about 50,000 cells) were reverse crosslinked, and DNA was isolated following the same protocol. Library preparation was carried out using Kapa Hyper Prep library preparation reagents (Kapa Hyper Prep kit, Roche 07962363001) following the manufacturer's protocol. ChIP libraries were sequenced using Illumina NovaSeq 6000 at 100bp paired-end reads.

RNA-seq data processing and analysis

Adaptor sequences and low-quality score bases (Phred score < 30) were first trimmed using Trimmomatic[74]. The resulting reads were aligned to the GRCh38 human reference genome assembly, using STAR[75]. Read counts were obtained using HTSeq[76] with parameters-m intersection-nonempty-stranded=reverse, using version Ensembl112 GTF annotation. For all downstream analyses, we excluded lowly-expressed genes with an average read count lower than 20 across all samples and unannotated genes (with ID names starting with ENSG), resulting in 19,568 genes. Raw counts were normalized using edgeR's TMM algorithm[77] and were then transformed to log2-counts per million (logCPM) using the voom function implemented in the limma R package[78]. To assess differences in gene expression levels between the different conditions (3 KO clones versus 3 WT replicates), we fitted a linear model using limma's lmfit function with parameter method="robust". For the Li-plus vs Li-minus analyses, we used a paired statistical test to account for the sample matchings between groups. Nominal p-values were corrected for multiple testing using the Benjamini-Hochberg method. The Z-score heatmap was constructed using the R package ComplexHeatmap[79]; KO vs WT analysis: P-adj (adjusted p-value) < 0.05 & |log2FC| > 0.25. Gene set enrichment analysis (GSEA)[35,36] based on a pre-ranked gene list by t-statistic was performed using the R package fgsea (http://bioconductor.org/packages/fgsea), P-adj < 0.05. Over-representation analysis (ORA) was performed using ShinyGO 0.77[39] (http://bioinformatics.sdstate.edu/go). The background genes used for ORA were all the 19,568 genes that were considered expressed. FDR is calculated based on the nominal P-value from the hypergeometric test. Fold Enrichment is the percentage of genes in the input gene list belonging to a pathway divided by the corresponding percentage in the background. For visualization on the heatmaps, gene expression data was corrected for batch effects using ComBat[80]. Significantly differentially expressed genes in KO-Li vs KO-UNT comparison were obtained using the cutoffs P.adj < 0.05 and |log2FC| > 0.25. Li-induced/rescued genes were obtained using the criteria below:

- Significantly differentially expressed (P.adj < 0.05 and |log2FC| > 0.25) in one comparison but not the other (i.e., in either Het-*AKAP11*-KO-UNT vs. WT-UNT or Het-*AKAP11*-KO-Li vs. WT-UNT)
- |diff_log2FC| > 0.6

ChIP-seq data processing and analysis

ChIP-seq reads were first trimmed for adapter sequences and low-quality score bases using Trimmomatic[74]. The resulting reads were mapped to the human reference genome (GRCh38) using BWA-MEM[81] in paired-end mode at default parameters. Only reads that had a unique alignment (mapping quality > 20) were retained and PCR duplicates were marked using Picard tools (http://broadinstitute.github.io/picard/). Peaks were called using MACS2[82]. To detect changes in histone modifications, a consensus peak set for H3K27ac was first generated by merging ChIP-seq peaks across samples using bedtools merge (https://bedtools.readthedocs.io/). A peak must be present in at least one sample in either condition. Read counts were obtained within these genomic regions using HOMER. Differential peak analysis was performed using the R package limma[78]. Nominal p-values were corrected for multiple testing using the

Benjamini-Hochberg method. Peaks were associated with the nearest TSS of genes using the annotatePeaks command from HOMER software suite[83]. Enrichment ORA of genes assigned to differential peaks was performed using ShinyGO 0.77[39] (http://bioinformatics.sdstate.edu/go). The background genes used for ORA were all the 19,568 genes that were considered expressed. FDR is calculated based on the nominal P-value from the hypergeometric test. Fold Enrichment is the percentage of genes in the input gene list belonging to a pathway divided by the corresponding percentage in the background. Bigwig coverage files were created with the HOMER makeUCSCfile command and bedGraphToBigWig utility from UCSC. Data were normalized so that each value represents the read count per base pair per 10 million reads. UCSC Genome Browser (http://genome.ucsc.edu/) was implemented for track visualization. Significantly differentially marked peaks in KO-Li vs KO-UNT comparison were obtained using the cutoffs P.adj < 0.05 and |log2FC| > 0.25. Li-induced/rescued peaks were obtained using the criteria below:

- Significantly differentially marked (P.adj < 0.05 and |log2FC| > 0.25) in one comparison but not the other (i.e., in either Het-*AKAP11*-KO-UNT vs. WT-UNT or Het-*AKAP11*-KO-Li vs. WT-UNT)
- |diff_log2FC| > 0.3

Acknowledgments

The work in G.A.R.'s lab is supported by the ERA-PerMed in partnership with FRQS (JTC2018); 280240. B.C.'s lab is supported by ERAPerMed PLOT-BD, a grant from the Fondation Bettencourt Schueller, and a government grant managed by the Agence Nationale de la Recherche under the France 2030 program (ANR-22-EXPR-0013). We also acknowledge the financial support from the Brain & Behavior Research Foundation Young Investigator Award to A.K; 30822. N.F. is supported by the studentship award Fonds de Recherche du Québec – Santé (FRQS). C.L. is supported by the CIHR Banting Fellowship. Computational analysis, data pre-processing, and adapted bioinformatics pipelines for analyses were performed by the Canadian Centre for Computational Genomics (C3G)-Montréal Node (Alain Pacis) using infrastructure provided by Compute Canada and Calcul Quebec. ChIP-seq library preparation and sequencing were performed at the McGill Genome Center platform.

Author contributions

N.F. conceived, designed, and carried out the laboratory experiments. A.K. and Y.L. gave guidance on the iPSC, NPC, and neuronal culture maintenance. N.F. designed and C3G carried out the computational analysis of the data with guidance from N.F. and A.K. D.R. and A.P. assisted in the laboratory work. C.L., M.A., P.A.D., G.A.R., B.C., C.L., and A.K. gave valuable scientific advice during the course of this study. N.F. wrote the manuscript.

Additional Information

Declaration of interests

The authors declare no competing interests.

Lead contacts

Additional information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contacts Drs. Rouleau, Khayachi, and Chaumette.

Materials availability

This study did not generate new unique reagents.

Supplementary Material

Supplementary Figures

Supplementary Fig. 1. (Related to Fig. 1). Transcriptomic effects of Li within the Het-*AKAP11*-KO and WT iPSC-derived neuronal models. A) Number of up and downregulated DEGs in WT-Li vs. WT-UNT; p.adj < 0.05 and |log2FC| > 0.25. Red: upregulated, blue: downregulated. **B)** Volcano plot of differentially expressed genes from WT-Li vs. WT-UNT comparison; p-value < 0.05 and |log2FC| > 0.25. Red: upregulated, blue: downregulated, grey: nonDEG. **C)** Hierarchical clustering heatmap of differentially expressed genes (rows) from each heterozygous Het-*AKAP11*-KO-Li clones vs. Het-*AKAP11*-KO-UNT reps. The expression levels across samples were standardized by the Z-Score method. Red shows the highest (3) and blue the lowest (-3) Z-score values. **D)** Heatmap of normalized enrichment scores (NES) of the overlapping GSEA[35,36] GO[37,38] molecular functions between Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT and Het-*AKAP11*-KO-UNT vs. WT-UNT from our previous study[28]. The functions marked with * are those that were previously

shown to be also significantly downregulated (p.adj < 0.1) in the GSEA of synapse proteomics from post-mortem brain samples of BD and SCZ patients relative to controls[29] as well as in our iPSC-derived Het-*AKAP11*-KO neuronal culture[28].

Supplementary Fig. 2. (Related to Fig. 2). Li's role in rescuing some of the DEGs obtained from Het-*AKAP11*-KO-UNT vs. WT-UNT and inducing new DEGs previously absent in Het-*AKAP11*-KO-UNT vs. WT-UNT. A) Volcano plot of DEGs from Het-*AKAP11*-KO-UNT vs. WT-UNT (top) and Het-*AKAP11*-KO-Li vs. WT-UNT (bottom) comparisons; p-value < 0.05 and |log2FC| > 0.25. Red: upregulated, blue: downregulated, grey: non-DEG. **B)** ORA of rescued DEGs following Li treatment of Het-*AKAP11*-KO (using all genes from **Fig. 2B** heatmap). GO[39] Molecular function database using ShinyGO 0.77[39]; Included in this table are FDR < 0.2. The yellow highlight indicates the most significant GO term. The background genes used were all the 19,568 genes that were considered expressed (see Methods). Shown here are the top 10 pathways, with redundancy removed. **C)** Full-size blots including all 12 samples; top: phGSK3B (S9), bottom: GAPDH loading control.

A Intergenic

B Intronic

Het-AKAP11-KO-Li vs. Het-AKAP11-KO-UNT   Het-AKAP11-KO-Li vs. Het-AKAP11-KO-UNT

Supplementary Fig. 3. (Related to Fig. 3). Differential intergenic and intronic H3K27ac peaks from Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT comparison. A) Differential intergenic peaks from *AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT comparison; p.adj < 0.05 and |log2FC| > 0.25; red: differentially elevated peaks, blue: differentially decreased peaks. **B)** Differential intronic peaks from *AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT comparison; P.adj < 0.05 and |log2FC| > 0.25; red: differentially elevated peaks, blue: differentially decreased peaks.

Supplementary Tables

Supplementary Table 1. DEG results of all the comparisons. We used DEGs with the cutoff of p-value < 0.05 and | log2FC| > 0.25 in this study.

Supplementary Table 2. GSEA[35,36] GO[37,38] molecular function of Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT; P.adj<0.05.

Supplementary Table 3. Overlapping GSEA[35,36] GO[37,38] molecular functions between Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNTand Het-*AKAP11*-KO-UNT vs. WT-UNT from our previous study[28].

Supplementary Table 4. Li-rescued and Li-induced DEG list. This table shows those genes that are significantly differentially expressed (P.adj < 0.05 and |log2FC| > 0.25) in one comparison but not the other (in either Het-*AKAP11*-KO-UNT vs. WT-UNT or Het-*AKAP11*-KO-Li vs. WT-UNT); |diff_log2FC| > 0.6.

Supplementary Table 5. Transcription factor Target Genome Browser PWMs generated using ShinyGo 0.77[39]. The table includes the top 20 most significant enriched motifs with FDR<0.1.

Supplementary Table 6. Immunoblot quantification table Het-*AKAP11*-KO-UNT, WT-UNT Het-*AKAP11*-KO-Li, WT-Li, determined using Fiji[73] software.

Supplementary Table 7. Het-*AKAP11*-KO-Li vs. Het-*AKAP11*-KO-UNT intergenic and intronic differential H3K27ac peaks and associated genes within the distance of 100 kb from the TSS to the peak center along with their gene expression levels.

 Supplementary Table 8. Differential (p.adj < 0.05 and |log2FC| > 0.25) intergenic and intronic H3K27ac peaks annotated results for Het-*AKAP11*-KO-UNT vs WT-UNT and Het-*AKAP11*-KO-Li vs. WT-UNT, along with all other comparisons.

Supplementary Table 9. Li-responsive differential intergenic and intronic H3K27ac peaks. Significantly differentially marked (P.adj < 0.05 and |log2FC| > 0.25) in one comparison but not the other (in either Het-*AKAP11*-KO-UNT vs. WT-UNT or Het-*AKAP11*-KO-Li vs. WT-UNT; |diff_log2FC| > 0.3). The H3K27ac peak-associated genes that overlapped with all Li-rescued or Li-induced DEGs are noted at the top. Yellow highlights indicate Li-rescued and Li-induced peaks that were linked to Li-rescued or Li-induced DEGs, respectively, with the same direction of change.

References

Grande, I., Berk, M., Birmaher, B. & Vieta, E. Bipolar disorder. *Lancet* **387**, 1561-1572 (2016). https://doi.org/10.1016/S0140-6736(15)00241-X

Kato, T. Current understanding of bipolar disorder: Toward integration of biological basis and treatment strategies. *Psychiatry Clin Neurosci* **73**, 526-540 (2019). https://doi.org/10.1111/pcn.12852

Bonnin, C. D. M. *et al.* Improving Functioning, Quality of Life, and Well-being in Patients With Bipolar Disorder. *Int J Neuropsychopharmacol* **22**, 467-477 (2019). https://doi.org/10.1093/ijnp/pyz018

Zhong, Y. *et al.* Global, regional and national burdens of bipolar disorders in adolescents and young adults: a trend analysis from 1990 to 2019. *Gen Psychiatr* **37**, e101255 (2024). https://doi.org/10.1136/gpsych-2023-101255

Miller, J. N. & Black, D. W. Bipolar Disorder and Suicide: a Review. *Curr Psychiatry Rep* **22**, 6 (2020). https://doi.org/10.1007/s11920-020-1130-0

Cipriani, A., Hawton, K., Stockton, S. & Geddes, J. R. Lithium in the prevention of suicide in mood disorders: updated systematic review and meta-analysis. *BMJ* **346**, f3646 (2013). https://doi.org/10.1136/bmj.f3646

Tighe, S. K., Mahon, P. B. & Potash, J. B. Predictors of lithium response in bipolar disorder. *Ther Adv Chronic Dis* **2**, 209-226 (2011). https://doi.org/10.1177/2040622311399173

Edvardsen, J. *et al.* Heritability of bipolar spectrum disorders. Unity or heterogeneity? *J Affect Disord* **106**, 229-240 (2008). https://doi.org/10.1016/j.jad.2007.07.001

McGuffin, P. *et al.* The heritability of bipolar affective disorder and the genetic relationship to unipolar depression. *Arch Gen Psychiatry* **60**, 497-502 (2003). https://doi.org/10.1001/archpsyc.60.5.497

Richards, A. L. *et al.* Genetic Liabilities Differentiating Bipolar Disorder, Schizophrenia, and Major Depressive Disorder, and Phenotypic Heterogeneity in Bipolar Disorder. *JAMA Psychiatry* **79**, 1032-1039 (2022). https://doi.org/10.1001/jamapsychiatry.2022.2594

O'Connell, K. S. & Coombes, B. J. Genetic contributions to bipolar disorder: current status and future directions. *Psychol Med* **51**, 2156-2167 (2021). https://doi.org/10.1017/S0033291721001252

Bipolar, D., Schizophrenia Working Group of the Psychiatric Genomics Consortium. Electronic address, d. r. v. e., Bipolar, D. & Schizophrenia Working Group of the Psychiatric Genomics, C. Genomic Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes. *Cell* **173**, 1705-1715 e1716 (2018). https://doi.org/10.1016/j.cell.2018.05.046

Palmer, D. S. *et al.* Exome sequencing in bipolar disorder identifies AKAP11 as a risk gene shared with schizophrenia. *Nat Genet* **54**, 541-547 (2022). https://doi.org/10.1038/s41588-022-01034-x

Lester, L. B., Coghlan, V. M., Nauert, B. & Scott, J. D. Cloning and characterization of a novel A-kinase anchoring protein. AKAP 220, association with testicular peroxisomes. *J Biol Chem* **271**, 9460-9465 (1996). https://doi.org/10.1074/jbc.271.16.9460

Whiting, J. L. *et al.* Protein Kinase A Opposes the Phosphorylation-dependent Recruitment of Glycogen Synthase Kinase 3beta to A-kinase Anchoring Protein 220. *J Biol Chem* **290**, 19445-19457 (2015). https://doi.org/10.1074/jbc.M115.654822

Consortium, G. T. The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science* **369**, 1318-1330 (2020). https://doi.org/10.1126/science.aaz1776

Nygren, P. J. & Scott, J. D. Therapeutic strategies for anchored kinases and phosphatases: exploiting short linear motifs and intrinsic disorder. *Front Pharmacol* **6**, 158 (2015). https://doi.org/10.3389/fphar.2015.00158

Meffre, D. *et al.* Wnt and lithium: a common destiny in the therapy of nervous system pathologies? *Cell Mol Life Sci* **71**, 1123-1148 (2014). https://doi.org/10.1007/s00018-013-1378-1

Tanji, C. *et al.* A-kinase anchoring protein AKAP220 binds to glycogen synthase kinase-3beta (GSK-3beta) and mediates protein kinase A-dependent inhibition of GSK-3beta. *J Biol Chem* **277**, 36955-36961 (2002). https://doi.org/10.1074/jbc.M206210200

Freland, L. & Beaulieu, J. M. Inhibition of GSK3 by lithium, from single molecules to signaling networks. *Front Mol Neurosci* **5**, 14 (2012). https://doi.org/10.3389/fnmol.2012.00014

Kishore, B. K. & Ecelbarger, C. M. Lithium: a versatile tool for understanding renal physiology. *Am J Physiol Renal Physiol* **304**, F1139-1149 (2013). https://doi.org/10.1152/ajprenal.00718.2012

Alda, M. Lithium in the treatment of bipolar disorder: pharmacology and pharmacogenetics. *Mol Psychiatry* **20**, 661-670 (2015). https://doi.org/10.1038/mp.2015.4

Jope, R. S. Lithium and GSK-3: one inhibitor, two inhibitory actions, multiple outcomes. *Trends Pharmacol Sci* **24**, 441-443 (2003). https://doi.org/10.1016/S0165-6147(03)00206-2

Khayachi, A., Schorova, L., Alda, M., Rouleau, G. A. & Milnerwood, A. J. Posttranslational modifications & lithium's therapeutic effect-Potential biomarkers for clinical responses in psychiatric & neurodegenerative disorders. *Neurosci Biobehav Rev* **127**, 424-445 (2021). https://doi.org/10.1016/j.neubiorev.2021.05.002

Creyghton, M. P. *et al.* Histone H3K27ac separates active from poised enhancers and predicts developmental state. *Proc Natl Acad Sci U S A* **107**, 21931-21936 (2010). https://doi.org/10.1073/pnas.1016071107

Rada-Iglesias, A. *et al.* A unique chromatin signature uncovers early developmental enhancers in humans. *Nature* **470**, 279-283 (2011). https://doi.org/10.1038/nature09692

Melgar, M. F., Collins, F. S. & Sethupathy, P. Discovery of active enhancers through bidirectional expression of short transcripts. *Genome Biol* **12**, R113 (2011). https://doi.org/10.1186/gb-2011-12-11-r113

Farhangdoost, N. *et al.* Transcriptomic and epigenomic consequences of heterozygous loss of function mutations in AKAP11, the first large-effect shared risk gene for bipolar disorder and schizophrenia. *bioRxiv*, 2024.2003.2014.584883 (2024). https://doi.org/10.1101/2024.03.14.584883

Aryal, S. *et al.* Deep proteomics identifies shared molecular pathway alterations in synapses of patients with schizophrenia and bipolar disorder and mouse model. *Cell Rep* **42**, 112497 (2023). https://doi.org/10.1016/j.celrep.2023.112497

Zhang, F., Phiel, C. J., Spece, L., Gurvich, N. & Klein, P. S. Inhibitory Phosphorylation of Glycogen Synthase Kinase-3 (GSK-3) in Response to Lithium: EVIDENCE FOR AUTOREGULATION OF GSK-3 *. *Journal of Biological Chemistry* **278**, 33067-33077 (2003). https://doi.org/10.1074/jbc.M212635200

Snitow, M. E., Bhansali, R. S. & Klein, P. S. Lithium and Therapeutic Targeting of GSK-3. *Cells* **10** (2021). https://doi.org/10.3390/cells10020255

Stahl, E. A. *et al.* Genome-wide association study identifies 30 loci associated with bipolar disorder. *Nat Genet* **51**, 793-803 (2019). https://doi.org/10.1038/s41588-019-0397-8

Singh, T. *et al.* Rare coding variants in ten genes confer substantial risk for schizophrenia. *Nature* **604**, 509-516 (2022). https://doi.org/10.1038/s41586-022-04556-w

Iacobucci, G. J. & Popescu, G. K. Ca(2+)-Dependent Inactivation of GluN2A and GluN2B NMDA Receptors Occurs by a Common Kinetic Mechanism. *Biophys J* **118**, 798-812 (2020). https://doi.org/10.1016/j.bpj.2019.07.057

Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550 (2005). https://doi.org/10.1073/pnas.0506580102

Mootha, V. K. *et al.* PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* **34**, 267-273 (2003). https://doi.org/10.1038/ng1180

Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* **25**, 25-29 (2000). https://doi.org/10.1038/75556

Gene Ontology, C. *et al.* The Gene Ontology knowledgebase in 2023. *Genetics* **224** (2023). https://doi.org/10.1093/genetics/iyad031

Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals and plants. *Bioinformatics* **36**, 2628-2629 (2020). https://doi.org/10.1093/bioinformatics/btz931

Forstner, A. J. *et al.* Identification of shared risk loci and pathways for bipolar disorder and schizophrenia. *PLoS One* **12**, e0171595 (2017). https://doi.org/10.1371/journal.pone.0171595

Schizophrenia Working Group of the Psychiatric Genomics, C. Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511**, 421-427 (2014). https://doi.org/10.1038/nature13595

Trubetskoy, V. *et al.* Mapping genomic loci implicates genes and synaptic biology in schizophrenia. *Nature* **604**, 502-508 (2022). https://doi.org/10.1038/s41586-022-04434-5

Amare, A. T. *et al.* Association of polygenic score and the involvement of cholinergic and glutamatergic pathways with lithium treatment response in patients with bipolar disorder. *Mol Psychiatry* **28**, 5251-5261 (2023). https://doi.org/10.1038/s41380-023-02149-1

Bhattacharya, S. *et al.* Triheteromeric GluN1/GluN2A/GluN2C NMDARs with Unique Single-Channel Properties Are the Dominant Receptor Population in Cerebellar Granule Cells. *Neuron* **99**, 315-328 e315 (2018). https://doi.org/10.1016/j.neuron.2018.06.010

Franchini, L., Carrano, N., Di Luca, M. & Gardoni, F. Synaptic GluN2A-Containing NMDA Receptors: From Physiology to Pathological Synaptic Plasticity. *Int J Mol Sci* **21** (2020). https://doi.org/10.3390/ijms21041538

Akkouh, I. A. *et al.* Exploring lithium's transcriptional mechanisms of action in bipolar disorder: a multi-step study. *Neuropsychopharmacology* **45**, 947-955 (2020). https://doi.org/10.1038/s41386-019-0556-8

Ghasemi, M. *et al.* The role of NMDA receptors in the pathophysiology and treatment of mood disorders. *Neurosci Biobehav Rev* **47**, 336-358 (2014). https://doi.org/10.1016/j.neubiorev.2014.08.017

Khayachi, A. *et al.* Akt and AMPK activators rescue hyperexcitability in neurons from patients with bipolar disorder. *EBioMedicine* **104**, 105161 (2024). https://doi.org/10.1016/j.ebiom.2024.105161

Sharma, V., Swaminathan, K. & Shukla, R. The Ribosome Hypothesis: Decoding Mood Disorder Complexity. *Int J Mol Sci* **25** (2024). https://doi.org/10.3390/ijms25052815

Ino, H. *et al.* Glutamatergic Neurometabolite Levels in Bipolar Disorder: A Systematic Review and Meta-analysis of Proton Magnetic Resonance Spectroscopy Studies. *Biol Psychiatry Cogn Neurosci Neuroimaging* **8**, 140-150 (2023). https://doi.org/10.1016/j.bpsc.2022.09.017

Jacoby, A. S., Vinberg, M., Poulsen, H. E., Kessing, L. V. & Munkholm, K. Increased DNA and RNA damage by oxidation in patients with bipolar I disorder. *Transl Psychiatry* **6**, e867 (2016). https://doi.org/10.1038/tp.2016.141

Laguesse, S. & Ron, D. Protein Translation and Psychiatric Disorders. *Neuroscientist* **26**, 21-42 (2020). https://doi.org/10.1177/1073858419853236

Forrest, M. P., Parnell, E. & Penzes, P. Dendritic structural plasticity and neuropsychiatric disease. *Nat Rev Neurosci* **19**, 215-234 (2018). https://doi.org/10.1038/nrn.2018.16

Pfaffenseller, B. *et al.* Impaired endoplasmic reticulum stress response in bipolar disorder: cellular evidence of illness progression. *Int J Neuropsychopharmacol* **17**, 1453-1463 (2014). https://doi.org/10.1017/S1461145714000443

Liu, Z. S. J. *et al.* Effects of Psychotropic Drugs on Ribosomal Genes and Protein Synthesis. *Int J Mol Sci* **23** (2022). https://doi.org/10.3390/ijms23137180

Karyo, R. *et al.* Identification of eukaryotic elongation factor-2 as a novel cellular target of lithium and glycogen synthase kinase-3. *Mol Cell Neurosci* **45**, 449-455 (2010). https://doi.org/10.1016/j.mcn.2010.08.004

Stork, C. & Renshaw, P. F. Mitochondrial dysfunction in bipolar disorder: evidence from magnetic resonance spectroscopy research. *Mol Psychiatry* **10**, 900-919 (2005). https://doi.org/10.1038/sj.mp.4001711

Mertens, J. *et al.* Differential responses to lithium in hyperexcitable neurons from patients with bipolar disorder. *Nature* **527**, 95-99 (2015). https://doi.org/10.1038/nature15526

Osete, J. R. *et al.* Lithium increases mitochondrial respiration in iPSC-derived neural precursor cells from lithium responders. *Mol Psychiatry* **26**, 6789-6805 (2021). https://doi.org/10.1038/s41380-021-01164-4

Heinrich, A., Boer, U., Tzvetkov, M., Oetjen, E. & Knepel, W. Stimulation by lithium of the interaction between the transcription factor CREB and its co-activator TORC. *Biosci Rep* **29**, 77-87 (2009). https://doi.org/10.1042/BSR20080116

Brown, K. M. & Tracy, D. K. Lithium: the pharmacodynamic actions of the amazing ion. *Ther Adv Psychopharmacol* **3**, 163-176 (2013). https://doi.org/10.1177/2045125312471963

Serretti, A., Drago, A. & De Ronchi, D. Lithium pharmacodynamics and pharmacogenetics: focus on inositol mono phosphatase (IMPase), inositol poliphosphatase (IPPase) and glycogen sinthase kinase 3 beta (GSK-3 beta). *Curr Med Chem* **16**, 1917-1948 (2009). https://doi.org/10.2174/092986709788186101

Kamerling, J. P., Kamerling, J. P., Boons, G.-J. & ScienceDirect. *Comprehensive glycoscience : from chemistry to systems biology*. 1st edition edn, (Elsevier, 2007).

Soles, A. *et al.* Extracellular Matrix Regulation in Physiology and in Brain Disease. *Int J Mol Sci* **24** (2023). https://doi.org/10.3390/ijms24087049

Dityatev, A., Seidenbecher, C. & Morawski, M. Brain extracellular matrix: An upcoming target in neurological and psychiatric disorders. *Eur J Neurosci* **53**, 3807-3810 (2021). https://doi.org/10.1111/ejn.15336

Berretta, S. Extracellular matrix abnormalities in schizophrenia. *Neuropharmacology* **62**, 1584-1597 (2012). https://doi.org/10.1016/j.neuropharm.2011.08.010

Jeffrey, K. L., Camps, M., Rommel, C. & Mackay, C. R. Targeting dual-specificity phosphatases: manipulating MAP kinase signalling and immune responses. *Nat Rev Drug Discov* **6**, 391-403 (2007). https://doi.org/10.1038/nrd2289

Grewal, S. S., York, R. D. & Stork, P. J. Extracellular-signal-regulated kinase signalling in neurons. *Curr Opin Neurobiol* **9**, 544-553 (1999). https://doi.org/10.1016/S0959-4388(99)00010-0

Fukunaga, K. & Miyamoto, E. Role of MAP kinase in neurons. *Mol Neurobiol* **16**, 79-95 (1998). https://doi.org/10.1007/BF02740604

Duric, V. *et al.* A negative regulator of MAP kinase causes depressive behavior. *Nat Med* **16**, 1328-1332 (2010). https://doi.org/10.1038/nm.2219

Perez-Sen, R. *et al.* Dual-Specificity Phosphatase Regulation in Neurons and Glial Cells. *Int J Mol Sci* **20** (2019). https://doi.org/10.3390/ijms20081999

Stern, S. *et al.* Neurons derived from patients with bipolar disorder divide into intrinsically different sub-populations of neurons, predicting the patients' responsiveness to lithium. *Mol Psychiatry* **23**, 1453-1465 (2018). https://doi.org/10.1038/mp.2016.260

Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012). https://doi.org/10.1038/nmeth.2019

Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120 (2014). https://doi.org/10.1093/bioinformatics/btu170

Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013). https://doi.org/10.1093/bioinformatics/bts635

Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015). https://doi.org/10.1093/bioinformatics/btu638

Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140 (2010). https://doi.org/10.1093/bioinformatics/btp616

Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47 (2015). https://doi.org/10.1093/nar/gkv007

Gu, Z. Complex heatmap visualization. *iMeta* **1**, e43 (2022). https://doi.org/https://doi.org/10.1002/imt2.43

Leek, J. T., Johnson, W. E., Parker, H. S., Jaffe, A. E. & Storey, J. D. The sva package for removing batch effects and other unwanted variation in high-throughput experiments. *Bioinformatics* **28**, 882-883 (2012). https://doi.org/10.1093/bioinformatics/bts034

Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754-1760 (2009). https://doi.org/10.1093/bioinformatics/btp324

Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008). https://doi.org/10.1186/gb-2008-9-9-r137

Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-589 (2010). https://doi.org/10.1016/j.molcel.2010.05.004