Profiling brain morphology for autism spectrum disorder with two cross-culture large-scale consortia

Xue-Ru Fan [1], **Ye He** [2], **Yin-Shan Wang** [1], **Lei Li** [3], **for the Lifespan Brain Chart Consortium (LBCC)** [*], **for the China Autism Brain Imaging Consortium (CABIC)** [**], **Xujun Duan** [3,***], **Xi-Nian Zuo** [1,***]

[1] State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal University, Beijing, 100875, China.
[2] School of Artificial Intelligence, Beijing University of Posts and Telecommunications, Beijing, 100876, China.
[3] Sichuan Provincial Center for Mental Health, Sichuan Provincial People's Hospital, School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu, 610054, China.
[*] A list of LBCC authors and their affiliations appears at the end of the paper. More information about LBCC are available at http://www.brainchart.io.
[**] A list of CABIC authors and their affiliations appears at the end of the paper. More information about CABIC are available at https://php.bdnilab.com.
[***] Corresponding author: Xi-Nian Zuo & Xujun Duan; Emails: xinian.zuo@bnu.edu.cn, duanxujun@uestc.edu.cn

ABSTRACT

We explore neurodevelopmental heterogeneity in Autism Spectrum Disorder (ASD) through normative modeling of cross-

cultural cohorts. Leveraging large-scale datasets from Autism Brain Imaging Data Exchange (ABIDE) and China Autism Brain

Imaging Consortium (CABIC), the model identifies two ASD subgroups with distinct brain morphological abnormalities:

subgroup "L" is characterized by generally smaller brain region volumes and higher rates of abnormality, while subgroup

"H" exhibits larger volumes with less pronounced deviations in specific areas. Key areas, such as the isthmus cingulate and

transverse temporal gyrus, were identified as critical for subgroup differentiation and ASD trait correlations. In subgroup H,

the regional volume of the isthmus cingulate cortex showed a direct correlation with individuals' autistic mannerisms, po-

tentially corresponding to its slower post-peak volumetric declines during development. These findings offer insights into

the biological mechanisms underlying ASD and support the advancement of subgroup-driven precision clinical practices.

INTRODUCTION

Autism Spectrum Disorder (ASD), or autism, is a lifelong neurodevelopmental condition characterized by impairments in social com-

munication and the presence of repetitive, unusual sensorimotor behaviors [1,2]. The global prevalence of ASD is estimated to be

approximately 1% in children [3], with different rates reported in specific regions, such as 2.8% in children aged 8 years in the United

States [4] and 0.7% in children aged 6 to 12 years in China [5]. Several factors, including improved survey methodologies and diagnos-

tic practices, increased public awareness and access to services, have contributed to this increase in prevalence [3,6]. Over time, un-

derstanding of ASD has developed from a categorical diagnosis to a dimensional perspective of neurodiversity [7]. This change rede-

fined autism as a spectrum that includes a wide range of characteristics within a unified framework [8]. However, expanded diagnos-

tic criteria, actually too low [9], have introduced significant heterogeneity within ASD. And this heterogeneity suggests that the term

"spectrum" encompasses multiple subgroups with distinct etiological phenotypes [10,11].

Current treatments for ASD, such as behavioral and speech therapies, provide benefits to some individuals but show limited overall

effectiveness [12]. In fact, there is limited evidence that any type of early intervention has produced significant and broad effects on

the main autistic outcomes [13]. Traditional case-control neuroimaging studies, which assume ASD and typical developing groups as

homogeneous entities, often fail to account for subgroup differences [14]. This has led to inconsistent findings across brain regions

and participants, further complicating efforts to identify reliable brain-behavior biomarkers. The decreasing effect sizes in group

comparison studies over time [15] highlight the difficulty of constructing mechanistic models of ASD. Data-driven clustering tech-

niques have emerged as a promising approach to address this complexity by leveraging cognitive and behavioral profiles alongside

patterns of brain morphology and function. Various clustering methods [16–20] have been used to identify meaningful subgroups, in

order to uncover the neurobiological mechanisms of ASD and provide insights into subgroup-specific traits [8,21]. Despite decades

of effort, inconsistencies in sample selection, classification methods, and feature selection between studies remain significant chal-

lenges [21]. These issues have limited the ability of clustering techniques to consistently identify robust and replicable subgroups.

Inconsistencies in ASD studies may be due to altered neurodevelopmental adaptations [22] and selection biases [23]. These limitations

highlight the need to group variables with strong biological relevance for the ASD subgroups [24]. Individual differences in sex, in-

telligence, medication use, and comorbidities [25,26], along with methodological variability [27] in data acquisition and analysis, still

complicate the interpretation of the results of neuroimaging studies. In response to these challenges, normative modeling [28] tech-

niques offer a promising framework by capturing the full range of neurobiological variation [14]. This approach enables standardized

assessments by comparing an individuals brain morphology to population-based references [29]. In ASD research, references are of-

ten built on internal or public datasets [18,20,30,31], but the relatively small sample size limits the generalizability and reliability of the

results. Recently, the Lifespan Brain Chart Consortium (LBCC, https://github.com/brainchart/lifespan) provided neurodevelopmental

variations throughout life through the most inclusive data available [32], providing a robust framework for us to achieve percentile-

based comparisons of ASD with considerable statistical power.

This study adjusted the LBCC normative models in two large-scale cross-cultural ASD consortia: the China Autism Brain Imaging

Consortium (CABIC) [33] and the Autism Brain Imaging Data Exchange (ABIDE) [34,35]. Focusing on childhood development, we used

the largest-scale brain charts [32] to explore neurodevelopmental heterogeneity in ASD before adolescence. We classified ASD into

subgroups using spectral clustering based on Out-of-Sample (OoS) centile scores, a biological measure which quantified brain mor-

phological deviations of ASD from normative growth. A Support Vector Machine (SVM) with Recursive Feature Elimination and Cross-

Validation (RFECV) identified key brain regions driving subgroup classification. The ABIDE-based classifier was then applied to CABIC

to identify similar subgroups. Brain-behavior analyses were performed separately for each dataset to identify reproducible subgroup-

specific correlations. From a perspective of brain morphology, this work disentangles the mechanisms of mixed neurosubtypes and

provides new insight into the biological complexity of ASD.

RESULTS

Brain Morphological Profiles Reveal Two Distinct Autism Subgroups

We first analyzed the ABIDE dataset. The distributions of the OoS centile scores across 34 bilateral Desikan-Killiany [36] cortical vol-

umes ranged from 0 to 1 (Fig. S1a in the Supplementary Information). OoS scores estimate the morphological variations of each

participant in 34 brain regions compared to the median regional volumes of the population (50th centile, equivalent to a 0.5 OoS

score). Relative to 0.5, individuals with more atypical phenotypes have more extreme scores. A ShapiroWilk test on the 34 OoS scores

indicated that none followed a normal distribution (Table 1, column 2). Using spectral clustering on the OoS scores, we identified

two distinct clusters with the highest silhouette coefficient among cluster solutions ranging from two to ten (Figure 1a). All p-values

for between-subgroup comparisons (two-sample t-tests for the paracentral and pars orbitalis regions, and rank-sum tests for all

other regions) were < 0.001). One subgroup, referred to as "L", exhibited generally smaller OoS scores, while the other, referred to

as "H", displayed larger scores (Table 1, columns 5 and 7).

Certain regions of the brain showed significant differences in their OoS score distributions between the two subgroups (Figure 1g,

Fig. S1b in the Supplementary Information. For CABIC results, please see Fig. S2a and S2b in the Supplementary Information.).

Subgroup H exhibited greater variability in the occipital lobe, particularly in the lingual gyrus, pericalcarine cortex, and cuneus. In

contrast, subgroup L showed more pronounced reductions in these regions compared to the normative benchmark (0.5). Other re-

gions, such as the pars opercularis and superior temporal gyrus, also exhibited subgroup-specific changes. The results of the Shapiro-

Wilk test for OoS scores (Table 1, columns 4 and 6) indicated that more regions of the brain in subgroup H followed a normal distri-

bution compared to subgroup L. Broader variability was observed in regions such as the entorhinal cortex in subgroup L and the

pericalcarine in subgroup H, highlighting further subgroup-specific clustering.

Distinct Structural and Age Differences Between Subgroups

The median OoS scores before clustering (Figure 1b; Table 1, column 3) reveal a mixed pattern of deviations across brain regions.

Most regions exhibit reduced volumes, particularly in the occipital lobe. However, some regions, such as the precuneus, paracentral

lobule, transverse temporal gyrus, and insula, show slightly higher OoS scores. This combined pattern reflects overlapping structural

deviations from both subgroup H (Figure 1c, top) and L (Figure 1c, bottom), obscuring the distinct morphological differences that

become evident after clustering. For instance, although subgroup L shows smaller volumes in the insula and precuneus, the larger

volumes in subgroup H predominantly drive the combined result in the whole population. In contrast, smaller volumes of the lin-

gual gyrus and the middle temporal gyrus in subgroup L have a stronger influence on the overall result. Some regions, such as the

Figure 1: Clustering-derived subgroups reveal distinct brain morphological patterns in ASD from ABIDE dataset.
a, Silhouette scores across different clustering solutions (2 to 10 clusters). The highest silhouette score indicates two distinct subgroups. **b,** Median OoS scores of 34 cortical regions before clustering. **c,** Median OoS scores of ASD subgroup H (top) and L (bottom). **d,** SHAP summary plot displaying the top 10 brain regions with the highest contributions to the SVM model's predictions. **e,** The mean SHAP values across the 29 selected cortical regions contributed to the classification. **f,** Prevalence maps depicting the proportion of participants with extreme (2.5% for subgroup L, 97.5% for subgroup H) structural anomalies. **g,** OoS scores across 34 cortical regions (top) and global measures (bottom) for subgroup H (orange) and L (green). TCV, total cortical volume; WMV, total white matter volume; GMV, total cortical gray matter volume; sGMV, total subcortical gray matter volume; CSF, total ventricular cerebrospinal fluid volume; mCT, mean cortical thickness; tSA, total surface area; Dots indicate the mean value of OoS scores, bars indicate the standard deviation. See Supplementary Information Fig. S1b and Fig. S1d for detailed density plots. Distribution of participant ages (**h**); IQ scales (**i**); ADI-R (**j**); PDD category (**k**, top left), MRI scanner model (**k**, top middle), manufacturer (**k**, top right), and data collection site (**k**, bottom); SRS (**l**); and ADOS (**m**) across the two subgroups, with subgroup L participants being significantly younger than those in subgroup H ($p = 0.02$). Note, the left hemispheres are plotted in **b, c, e, and f** just for visualization purposes. For plots **h, i, j, l and m**, the center line shows the median; the box limits represent the 25th and 75th percentiles; the whiskers show the minimum and maximum values; and the dots represent potential outliers.

pericalcarine cortex, are almost entirely influenced by subgroup L. Meanwhile, regions like the inferior and superior temporal gyrus

exhibit volumes close to the normative average because of the contrasting characteristics of both subgroups effectively canceling

each other out.

To identify the regions of the brain most associated with morphological abnormalities, we calculated the prevalence of abnormali-

ties for each region in the two subgroups. For subgroup L, the prevalence represents the percentage of participants with OoS scores

below 0.025 (2. 5% centile), while above 0.975 (97. 5% centile) for subgroup H. Subgroup L (Figure 1f, bottom; Table 1, column 8)

exhibited the highest prevalence of abnormalities in regions such as the middle and inferior temporal gyrus, pericalcarine cortex,

frontal pole, and medial orbital frontal gyrus. In contrast, subgroup H (Figure 1f, top; Table 1, column 9) generally exhibited less se-

vere abnormalities, with a concentration in the insula, transverse temporal gyrus, and caudal anterior cingulate. Participants in sub-

group L were significantly younger than those in subgroup H (Figure 1h, $p = 0.02$, Table S1 in Supplementary Information). How-

ever, no significant differences were observed in the category PDD $p = 0.32$, the MRI scanner model $p = 0.07$ and the manufacturer

$p = 0.86$, the data collection site $p = 0.16$ (Figure 1k), IQ (Figure 1i, see Table S1 in Supplementary Information for details), or scores

from ADI-R (Figure 1j), SRS (Figure 1l), and ADOS-2 (Figure 1m). The corresponding results for CABIC are shown in Supplementary

Information Fig. S3.

Machine Learning Reveals Subgroup Morphological Features

The optimized SVM model for the identification of subgroups demonstrated robust predictive performance. It used a polynomial

kernel with a regularization parameter $C = 0.1$ and a kernel coefficient $\gamma = 1$. The model achieved a high classification accuracy

$(0.95, p < 0.001)$ and a F1 score $(0.94, p < 0.001)$ in five cross-validation folds. 1d shows the SHAP summary plot for the top 10 se-

lected features. 1e and Table S2 in Supplementary Information summarize the mean SHAP values for all 29 selected regions. SHAP

values quantify the contribution of individual brain regions to the predictions of the model, with higher values indicating stronger

influence. Among the regions, the isthmus cingulate, entorhinal cortex, precuneus, and middle temporal gyrus emerged as the

most predictive features. Higher SHAP values for these regions were strongly associated with subgroup H, while lower values cor-

responded to subgroup L. These findings indicate different volumetric patterns in the cingulate, temporal, and parietal areas, effec-

tively distinguishing the two subgroups.

Distinct Structural Covariance Patterns Across Subgroups

Structural covariance analysis on OoS scores revealed significant correlations between brain regions, with distinct patterns observed

across ASD subgroups and control group (2c). These patterns were reproduced in both the ABIDE and CABIC datasets. In subgroup

H, enhanced covariation was identified between the isthmus cingulate and the caudal middle frontal region (ABIDE: $z = 2.88, p =$

0.00; CABIC: $z = 2.25, p = 0.02$). In subgroup L, the parahippocampal gyrus showed reduced covariation with the entorhinal cortex

(ABIDE: $z = -2.07, p = 0.04$; CABIC: $z = -3.25, p = 0.00$). Similarly, reduced covariation was observed between the posterior cin-

gulate and the entorhinal cortex (ABIDE: $z = -2.12, p = 0.03$; CB: $z = -2.64, p = 0.01$) and between the insula and the entorhinal

cortex (ABIDE: $z = -3.30, p = 0.00$; CABIC: $z = -2.18, p = 0.03$). An increased covariation was observed in subgroup L between the

parahippocampal gyrus and the lateral occipital cortex (ABIDE: $z = 1.97, p = 0.05$; CABIC: $z = 2.04, p = 0.04$).

Brain-Behavior Correlations in Subgroup H

We examined correlations between cortical region volumes (OoS scores) and clinical measures in the ABIDE and CABIC datasets,

identifying significant associations that were consistent between both datasets. All significant correlations were found within sub-

group H. The transverse temporal gyrus showed positive correlations with ADOS Total and Social Affect scores (2a). In the CABIC

dataset, moderate correlations were observed for ADOS Total ($r(198) = 0.17, p = 0.02$) and Social Affect ($r(191) = 0.21, p = 0.00$).

Stronger correlations were identified in the ABIDE dataset for ADOS Total ($r(93) = 0.24, p = 0.01$) and Social Affect ($r(93) = 0.20$,

$p = 0.04$). The inferior temporal gyrus was positively correlated with ADOS RRB, with a slightly stronger correlation in ABIDE ($r(94) =$

$0.19, p = 0.05$) compared to CABIC ($r(189) = 0.17, p = 0.02$). The volumetric centile of the isthmus cingulate is correlated with SRS

Autistic Mannerisms, showing a higher correlation in ABIDE ($r(58) = 0.33, p = 0.01$) than in CABIC ($r(192) = 0.14, p = 0.04$). No signifi-

cant correlations were identified between structural covariance patterns and cognitive behavior outcomes.

Figure 2: Brain-behavior correlations and structural covariations of Out-of-Sample scores.
a, Reproducible correlations between brain region volumes, measured as Out-of-Sample (OoS) scores, and clinical measures across ABIDE and CABIC datasets (subgroup H only). **b,** Mediation models for significant brain-behavior associations identified in subgroup H (ABIDE cohort only). Black solid arrows represent significant effects, while gray arrows indicate non-significant ones. Top: transverse temporal; bottom left: inferior temporal; bottom right: isthmus cingulate. **c,** Significant structural covariations across ABIDE and CABIC datasets. Positive z-values (red lines) indicate stronger covariance in ASD participants compared to controls, while negative z-values (blue lines) reflect weaker covariance.

As a common comorbidity of ASD, intellectual disability is closely associated with atypical brain morphological devel­opmental pat-
terns that accompany individual differences in intellectual functioning. To examine whether Full-Scale Intelligence Quotient (FIQ)
mediates the relationships between brain morphology and cognitive behaviors within clusters, mediation analyses were conducted
on the significant correlations we found in subgroup H. Results revealed direct effects for specific cortical regions (2b). The trans-
verse temporal gyrus showed a direct effect on ADOS Total ($C' = 0.21, p = 0.03$), while the isthmus cingulate exhibited a direct effect
on SRS Autistic Mannerisms ($C' = 0.32, p = 0.01$). No mediation effects through FIQ were observed. Neither direct nor indirect ef-
fects were significant for correlations to ADOS RRB or Social Affect.

DISCUSSION

This study used the ABIDE and CABIC datasets to discover distinct brain morphological subgroups of ASD within a young male co-
hort. We identified two subgroups characterized by significant differences in OoS scores in 34 cortical regions. Machine learning
models showed a high predictive accuracy in distinguishing these subgroups based on their brain morphology. Furthermore, struc-
tural covariance and brain behavior correlation analysis illustrated different patterns of morphological relationships and their associ-
ations with clinical measures, especially in subgroup H. The consistent correlations across both the ABIDE and CABIC datasets high-
light the robustness and reproducibility of our results across diverse cohorts. We will now discuss these findings from several neuro-
biological and neurodevelopmental perspectives to enhance our understanding of the distinct morphological patterns identified.

Two Subgroups Suggest Different Neurobiological Basis

Studies have reported extremely large or small head circumferences in individuals with ASD, possibly related to different subgroups.
Early brain overgrowth is one of the most consistent findings in ASD research [37]. Abnormal increases in brain size during early de-
velopment suggest that atypical cell proliferation significantly contributes to ASD symptoms [38]. Ultrasound studies have detected
head enlargement during the second trimester in individuals later diagnosed with ASD [39]. iPSC-based studies reveal enlarged em-

bryonic stage brain cortical organoids (BCOs) in babies with ASD, with larger BCOs associated with more severe social symptoms[40].

MRI scans of these individuals show overgrowth in primary auditory and somatosensory cortices, while undergrowth in the visual

cortex[40]. This neuronal overproduction, caused by the acceleration of the cell cycle, results in impaired differentiation and ultimately

disrupts neural functions[41].

Smaller brain volumes are also frequently reported in ASD. A meta-analysis[42] identified volumetric reductions in a large cluster

of regions, including the parahippocampal gyrus and entorhinal cortex. Higher levels of autistic traits have been associated with a

smaller total cortical volume (TCV), a lower cortical thickness, a smaller surface area and a lower gyrification[43,44]. One potential ex-

planation for this global brain underdevelopment is impairment caused by insufficient blood circulation and oxygen saturation[45].

For example, children with complex congenital heart disease (CHD) also demonstrate cognitive difficulties similar to those observed

in ASD and have an increased likelihood of developing ASD[46]. Similarly, head enlargement also coincide with greater increases in

height often[44]. Therefore, brain overgrowth could be part of the broader physical growth dysregulation[47] too. These evidences pro-

vide the distinct neurobiological basis for the subgroups identified in our study. The different underlying physiological mechanisms

may explain the inconsistent effectiveness of commonly used pharmacological treatments.

Hypothesis of Abnormal Brain Morphology Specific Functional Impairment Pathways

Abnormal neural migration and cortical laminar organization in ASD[48] suggest that structural abnormalities are unlikely to be lo-

calized. Postmortem studies have identified focal laminar disorganization and mis-migrated neurons in the prefrontal and tempo-

ral cortex, particularly in layers 2, 3, and 4[48]. Layers 2 and 3 support information exchange between cortical regions, while layer 4

receives sensory inputs[49]. Disruptions in these layers can cause profound miscommunication between the sensory and higher-

order cognitive regions. Key regions in our clustering analysis (such as the precuneus, isthmus cingulate, and entorhinal cortex; 3a,

a~c) are primarily involved in higher-order association networks (3a). This aligns with the concepts of functional information flow in

the brain[50], suggesting different pathways of impairment as the basis for the heterogeneity of ASD. Inconsistencies in information

flow reported by previous ASD studies[21,51] may arise from two distinct structural impairment patterns identified in our study.

The regional volume abnormalities identified in our study involve distinct functional brain regions in the two subgroups. Subgroup

L showed a notable prevalence in the middle temporal gyrus, inferior temporal gyrus, pericalcarine cortex and frontal pole (3a, d~g),

while subgroup H in the caudal anterior cingulate, transverse temporal cortex, and insula (3a, h~j). Drawing on the spatial distribu-

tions[52] of 7 large-scale functional networks[53], we constructed a schematic (3b) of the latest 15 large-scale functional networks of

the brain estimated from individuals[54] (3a). This perspective revealed different pathways of functional impairment in the regions

with a high prevalence of abnormalities for each subgroup. Subgroup L exhibited an atypical morphology mainly affecting Visual –

Dorsal Attention – Frontoparietal – Language and Default networks (DN) (3b, green path), while subgroup H affected the Auditory

and Somatomotor – Cingulo-Opercular and Salience/Parietal Memory (SAL/PMN) – Frontoparietal networks (3b, orange path), indi-

cating disturbances in peculiar unimodal sensory integration and multimodal cortical functions. This phenomenon may hint at the

presence of distinct abnormal brain functional circuits from first- to third-order networks in the subgroups.

For example, in adults without ASD[55], increased gray matter volume in regions of the somatomotor network was associated with

greater attention to detail, while changes in regions of the visual network were associated with poor imagination. These findings

suggest potential cognitive patterns in two subgroups of ASD that have yet to be discovered. In typically developing children, the

principal functional gradient[52] of the cortical organization reflects differentiation within unimodal sensory areas, progressing from

the somatomotor and auditory cortex to the visual cortex. The transition from childhood to adolescence extends this organization

to higher-order association cortices[56]. Brain maturation appears to involve a shift from local to distributed network architecture.

The distinct morphological abnormalities in these two ASD subgroups, observed before this transition begins, raise a critical ques-

tion: How do these "morphological foundations" disrupt their development from childhood to adolescence? These structural im-

pairment pathways in childhood are likely to contribute further to heterogeneity in ASD during and after adolescence.

Figure 3: The theoretical brain functional network impairment model of two ASD subgroups.
a, A brain map combining the 34 Desikan regions[36] with the latest 15 large-scale brain functional networks estimated from individuals[54]. *a*, precuneus; *b*, isthmus cingulate; *c*, entorhinal; *d*, middle temporal; *e*, inferior temporal gyrus; *f*, pericalcarine; *g*, frontal pole; *h*, caudal anterior cingulate; *i*, transverse temporal; and *j*, insula. **b,** A schematic of the potential spatial distributions of distinct functional impairment networks inferred from 34 cortical regions with high abnormal prevalence for two subgroups. VIS, Visual (C, Central; P, Peripheral); SMOT, Somatomotor; AUD, Auditory; dATN, Dorsal Attention; PM-PPr, Premotor-Posterior Parietal Rostral; CG-OP, Cingulo-Opercular; SAL/PMN, Salience/Parietal Memory; FPN, Frontoparietal; LANG, Language; DN, Default networks.

Understanding the Heterogeneity from a Developmental Perspective

Different network-based vulnerabilities suggest that interventions targeting specific impairments within each subgroup could be

more effective. Interestingly, we found no significant differences in cognitive or behavioral measures between the subgroups. Al-

though there distinct patterns of structural covariation between subgroups, but no significant correlations with autism-related be-

haviors. Structural covariance reflects long-term developmental changes. Reduced structural covariance may indicate disruptions

in shared developmental pathways, while enhanced covariance could result from compensatory adaptations in response to atypical

neurodevelopmental trajectories in ASD. This implies that morphological deficiencies may activate compensatory mechanisms dur-

ing development, leading to similar cognitive and behavioral outcomes. The observed age difference between subgroups, though

minor, could have significant implications during sensitive developmental periods. Lifespan trajectories[32] of 34 brain regions show

an early steep increase in volume, followed by a near-linear decline. The middle temporal gyrus peaks at 7.8 years, while the caudal

anterior cingulate peaks at 9.2 years[32], possibly related to the result that subgroup L is generally younger than subgroup H. These

distinct structural abnormalities likely reflect differences in the timing of atypical developmental processes.

The isthmus cingulate, a key region in subgroup clustering, typically peaks in volume at 3.8 years. The lack of age-related decline

in cortical volume was consistently found in subgroup H in both the ABIDE and CABIC datasets (See Supplementary Analysis and

Fig. S1f). However, subgroup H aligns with normative references before age 5, while subgroup L shows smaller initial volumes (Fig.

S1e). In subgroup H, an increase in structural covariance between the caudal middle frontal cortex (peaking at 7.75 years) and the

isthmus cingulate (2c) suggests synchronized developmental trajectories. This may reflect precocious caudal middle frontal de-

velopment, delayed isthmus cingulate development, or both regions following similarly abnormal patterns[22]. Correlation analy-

ses link larger isthmus cingulate volumes in subgroup H directly with more pronounced autistic mannerisms (2a). These evidence

hint at delayed development of the isthmus cingulate in subgroup H. With a normal OoS distribution ($p = 0.06$; Table 1, column

6) in this subgroup, this morphological change appears to represent a homogeneous and quantifiable pattern. In subgroup L, the

parahippocampal gyrus (peaking at 10.63 years) shows reduced covariance with the entorhinal cortex (peaking at 22.67 years) but

increased covariance with the lateral occipital cortex (peaking at 5.21 years). This pattern suggests a precocious development of the

parahippocampal gyrus and/or the delayed lateral occipital cortex in this subgroup. Enhanced structural covariances may also re-

flect compensatory adaptations to mitigate structural disadvantages inherent in abnormal developmental patterns. This highlights

the need for longitudinal studies to explore how brain maturation influences behavior at different stages of development in ASD.

Reproducibility across Datasets and Implications for Biomarker Development

One key contribution of this study is the reproducibility of brain-behavior associations across the ABIDE and CABIC datasets. Cross-
dataset validation confirms that the findings reflect stable neurobiological characteristics. In subgroup H, an enlarged transverse
temporal gyrus may impair precise auditory processing. This disruption can affect auditory responsiveness and the ability to inte-
grate and understand information during social interactions. It may also damage early language perception and acquisition, partic-
ularly before a diagnosis of ASD is made. Research has found that when ASD is accompanied by cognitive learning needs, head en-
largement becomes even more pronounced[44]. This suggests a shared mechanism linking brain overgrowth with impaired intellec-
tual functioning. Alternatively, cognitive impairments might mediate the relationship between this overgrowth and ASD. Although
intelligence moderates the relationship between ASD traits (ADOS total score), transverse temporal gyrus abnormality remains a
direct indicator (2b). This suggests impaired primary sensory perceptual function that is directly disrupted by abnormal sensory
input. The inferior temporal gyrus (ITG) is also uniquely associated with restricted interests and repetitive behaviors (ADOS RRB),
despite the low prevalence of structural abnormalities in subgroup H. This region plays a crucial role in object recognition[57]. Dis-
ruptions in these functions can lead to a tendency to focus repetitively on objects or details. The involvement of IFG in perceptual
and cognitive processes can also influence how individuals respond to environmental stimuli, further impacting RRB. In particular,
the correlation between ITG and RRB was no longer significant after controlling for FIQ. This suggests that variations in intelligence
may obscure the relationship. ITG might be specifically associated with certain dimensions of RRB while individuals with higher FIQ
could develop other cognitive strategies to regulate RRB, reducing the apparent role of ITG in these behaviors.

We classified ASD subtypes solely based on brain morphology, assuming a causal relationship between brain structure and cogni-
tion. However, this relationship is inherently complex, thus interpreting the brain in regions goes against its interconnected nature.
Although subgroup L exhibited a higher prevalence of structural abnormalities, suggesting more extensive morphological varia-
tions, we did not find reproducible brain-behavior associations between cohorts. Several factors may explain these findings. First,
cognitive impairments related to structural abnormalities in subgroup L might be more sensitive to cultural differences. Second,
the validity of ASD diagnoses in subgroup L might be questionable. The best predictor of a DSM-IV diagnosis has been found to be
often the specific clinic attended, rather than any defining characteristic of the individual[58]. This indicates that current diagnostic
tools may not capture the nuanced behaviors directly related to brain abnormalities in subgroup L. Considering global reduction in
brain volume, which may result from cardiogenic etiological damage, our findings highlight the urgent need for new methods and
paradigms to enable deep phenotyping and reduce the reliance on behavioral scales for differentiation. Third, heterogeneity within
subgroups may also contribute to the limited reproducibility findings. Behavioral paradigms targeting brain regions with a normal
distribution of OoS scores could better reveal subgroup-specific relationships between brain morphology and cognitive behavior.

Limitations & Future Direction

Some limitations in this study should be noted. First, previous research highlights atypical structural connectome asymmetry in
ASD[59], while we could not fully evaluate hemispheric differences within subgroups using LBCC trajectories. Second, the cross-sectional
nature of the data limits causal inferences about how these abnormalities of the brain morphology arise and their relationships
with ASD symptoms. Future research should thoroughly investigate brain function and cognitive differences between these sub-
groups while exploring how demographic, environmental, and developmental factors shape brain-based subtypes. Combining lon-
gitudinal data $invivo$ and $invitro$[41] could expand cohort diversity and provide deeper insights into the neurobiological and molec-
ular mechanisms underlying different subtypes. When designing research cohorts to study further, focus on more representative
ASD prototypes[60] could help reduce heterogeneity within subgroups. More attention should be paid to people with more severe
ASDs, such as minimally verbal autism, who remain underrepresented in brain imaging studies[61], partly due to the challenges of
collecting neuroimaging data, especially in young children. Our research strategy uses structural magnetic resonance imaging, fea-
sible in natural sleep or sedation, with significant clinical potential. However, multimodal approaches with higher ecological valid-
ity are essential to identify neurosubtypes that include these samples. At the same time, replication with different clustering algo-

rithms is required to confirm the stability of these findings. More nuanced clustering strategies should be explored, as disease vari-

ations may exist within the range of normal variation [62], and multiple subpopulations may also exist in typically developing popula-

tions. Finally, translational studies using animal models are essential for understanding the mechanisms underlying these different

subtypes. Such studies can provide critical information on how abnormalities in brain morphology develop [63].

Conclusion

Linking specific brain morphology abnormalities to cognitive or behavioral impairments in ASD is challenging. Different biological

mechanisms can lead to similar behavioral outcomes, complicating the identification of neural correlates. This overlap may explain

the difficulty in replicating neuroimaging findings between studies or cohorts. Using the largest brain normative datasets and two

large-scale cross-cultural consortiums, we applied normative modeling to disentangle the heterogeneity of brain morphology in

ASD. By breaking the ASD population into smaller and more morphologically homogeneous subgroups, we identified two subtypes

of morphological abnormalities. We identified significant correlations between cortical regions and autistic traits across both con-

sortia, with particular emphasis on the isthmus cingulate cortex, as well as the transverse and inferior temporal gyri. Based on the

abnormal prevalence of regional volumes, we found two structural impairment pathways that could disrupt sensory to higher cog-

nitive functions. These findings highlight the importance of untangling mixed biological mechanisms and offer insight to develop

effective subgroup-driven individualized interventions. In conclusion, our study provides a reference for elucidating the etiological

mechanisms of neurodevelopmental disorders and advancing future subgroup-driven precision clinical practice.

MATERIALS AND METHODS

Participants and Data Preprocessing

Participants were selected from the Autism Brain Imaging Data Exchange (ABIDE) cohort [34,35]. The original dataset includes 1,833

autistic and 1,838 non-autistic individuals aged 264 years from 56 sites. All diagnoses were based on the DSM-IV or DSM-V criteria.

To align with the objectives of our study, we focus on male participants under 13 years of age. This age range reflects our interest in

brain development during childhood and early adolescence.

We used T1-weighted MRI data from the ABIDE cohort. The acquisition parameters are publicly available at ABIDE website. Data

quality was assessed through a combination of automated checks using (MRIQC) [64] (version 22.0.6) and manual visual inspections.

Images with severe motion artifacts, segmentation errors, or structural abnormalities were excluded from further analysis. Pre-processing

performed through the Connectome Computation System (CCS) pipeline [65,66]. FreeSurfer (toolbox: version 6.0.0) was used for skull

removal, cortical reconstruction, segmentation, and registration in the Montreal Neurological Institute (MNI) template. Each T1-

weighted image was segmented into gray matter (GM), white matter (WM), and cerebrospinal fluid (CSF).

The total cortical volume (TCV) was calculated as the sum of the GM and WM volumes, following the LBCC method [32]. The mean

cortical thickness (CT) was calculated as a weighted average of the thicknesses of the left and right hemispheres. The total surface

area (SA) was determined by summing the surface areas of white matter of both hemispheres. Regional brain volumes for 34 bilat-

erally averaged cortical regions were evaluated using Desikan-Killiany parcellation [36]. To reduce site-related variability, we applied

ComBat harmonization [67]. This process was implemented using (*neuroCombat*) R package.

Normative Scoring and Clustering Analysis

To assess how the brain morphology of each participant aligned with age-standardized normative trajectories, we used the out-of-

sample (OoS) centile score [32]. First, all non-ASD individuals from the site-harmonized ABIDE data set were employed to estimate

cohort-specific statistical offsets. The centile scores of each brain measurement were then estimated for all individuals with ASD

compared to the offset trajectory (see *Methods: Centile scoring of new MRI data* in the original LBCC work [32]). This approach takes

advantage of the generalized additive models for location, scale, and shape (GAMLSS) [68] framework to estimate statistical offsets

such as mean, variance, and skewness. These offsets capture individual deviations from typical brain development. OoS scores are

comparable between age and cohort. The scoring process was conducted with the publicly available code from Lifespan Charts

website.

Using OoS scores from 34 brain regions as classification features, we applied spectral clustering[69] among individuals with ASD in

the ABIDE cohort. Spectral clustering was selected for its ability to detect nonlinear data structures, which are common in brain

imaging datasets. Unlike k-means clustering, which relies on centroid-based optimization, spectral clustering transforms the data

into a lower-dimensional space using the eigenvalues of a similarity matrix. Before clustering, the Jarque-Bera test was performed

to determine whether OoS scores follow a normal distribution.

Clustering analysis was implemented using the *kernlab* package R. We tested clusters of 2 to 10 to identify the optimal solution.

The quality of clustering was evaluated using the Silhouette Coefficient (SC)[70], which assesses both within-cluster cohesion and

between-cluster separation. SC values range from -1 to 1, with higher scores indicating better defined clusters. The optimal num-

ber of clusters was decided by the highest SC score.

Key Brain Regions for Subgroup Classification

We developed a predictive model using machine learning with individual OoS scores from 34 regions of the brain as input charac-

teristics. We trained a Support Vector Machine (SVM) classifier[71] with Recursive Feature Elimination and Cross-Validation (RFECV).

This was implemented using the *scikit-learn* library in Python. Combining recursive feature elimination (RFE)[72] with cross-validation

allows the model to iteratively identify the most informative features while maintaining generalizability and robustness. The data

set was divided into 80% training sets and 20% testing sets. Using a linear SVM as the base estimator, RFECV removed less impor-

tant features in a stepwise manner, with performance monitored through 5-fold cross-validation. This process ensured that the final

feature set balanced predictive power and reduced over-fitting.

Following feature selection, we fine-tuned the SVM model by performing a grid search to optimize key hyperparameters, includ-

ing regularization strength (C) and kernel coefficients (γ). The model was evaluated across different kernel functions, such as linear,

polynomial, and radial basis functions, to determine the best configuration. Once optimized, the model was trained on the selected

features to achieve optimal classification performance. SHapley Additive exPlanations (SHAP)[73] were applied to quantify the contri-

bution of each brain region to the model predictions.

Structural Covariance Network Analysis

314 To investigate morphological relationships of a specific cluster, we constructed structural covariance analysis[74] on OoS scores of

315 each two regions. Pearson's correlation analysis was performed on residuals obtained from linear regression models that controlled

316 for site effects and TCV OoS scores. To enable statistical comparison, correlation coefficients were normalized using Fishers r-to-z

317 transformation, defined as:

$$Z = \frac{1}{2} \log_e \left(\frac{1+r}{1-r} \right), \tag{1}$$

318 where r represents the Pearson correlation coefficient between two regions. The difference (cluster i/ii vs. normal control) in trans-

319 formed correlation coefficients was calculated using:

$$z = \frac{Z_1 - Z_2}{\sqrt{\frac{1}{n_1-3} + \frac{1}{n_2-3}}}, \tag{2}$$

320 where n_1 and n_2 are the two sample sizes being compared. The resulting z-values were converted to p-values using the cumulative

321 distribution function of the standard normal distribution.

Cluster Differences and Brain-Behavior Analysis

We investigated differences between clusters and demographic, technical and clinical variables. For categorical variables, such as

subtypes of perinatal developmental disorders (PDD), scanner models, and manufacturers, Chi-square tests were employed. For

data collection sites, Fishers exact test was used with Monte Carlo simulations to approximate p values. To ensure robust analysis,

sites with fewer than 10 participants were excluded. For other continuous variables, statistical methods were selected based on

data distribution: the Wilcoxon rank sum test was used for nonnormally distributed data, and the two-sample t-test for data shows

normal distribution. The effect sizes were calculated using Cramér's V for categorical variables and Cohens d for continuous vari-

ables. Statistical significance was set at $p < 0.05$.

To further explore brain-behavior relationships within clusters, we performed correlation analyses using Full-Scale Intelligence Quo-

tient (FIQ), ADOS-2 (Social Affect and Restricted and Repetitive Behaviors, RRB), ADI-R (Social Interaction and RRB), and SRS sub-

scales. Pearson's correlation was applied to most measures, while Spearman's correlation was used for ADOS-2 RRB scores due to

their limited range. In these analyzes, the effects of the site and the OoS scores of TCV were controlled.

For each group, we also examined partial correlations to investigate structural covariance between 34 brain regions and their asso-

ciation with cognitive behaviors. For each participant, the partial covariance scores of two regions were calculated as

$$\text{partial\_p} = \text{residual}_{region1} \times \text{residual}_{region2},$$

336 The resulting partial covariance scores were then correlated with the same measurements with the same correlated methods.

337 Finally, we performed mediation analyzes to examine whether FIQ mediates the relationships between brain morphology and cog-

338 nitive behaviors within clusters. Linear models were used to estimate direct and indirect effects, with adjustments for site and OoS

339 scores of TCV. All brain morphology and cognitive behavior variables were standardized to ensure comparability among participants.

340 The significance of indirect effects was assessed using the Sobel test.

341 Detection of Reproducible Results Between Datasets

We applied the ABIDE-based classifier to an independent dataset from the China Autism Brain Imaging Consortium (CABIC) [33],

which is a multicenter collaboration among clinical and research institutions in China. It is designed to collect neuroimaging, de-

mographic, and behavioral data on children and adolescents with ASD. Diagnoses in the CABIC cohort were based on DSM-IV, DSM-

V, or ICD-11 criteria and validated using ADOS and ADI-R. The dataset includes 1,400 participants, comprising 983 autistic individ-

uals and 514 typically developing controls. Detailed MRI acquisition parameters are publicly accessible at CABIC Site.

CABIC participants, aged 1 to 13 years, tend to exhibit more severe ASD symptoms, resulting in lower variability within clusters. Al-

though this homogeneity is clinically valuable, it poses challenges in detecting subtle brain-behavior relationships. In contrast, the

ABIDE cohort provides greater behavioral differences, enhancing the ability to identify nuanced patterns in brain structure. By lever-

aging these complementary datasets, our goal was to identify results that are reproducible across both cohorts. We repeated the

statistical, structural covariance, and correlation analyses on CABIC cohort. However, the limited FIQ data in CABIC restricted the

replication of mediation analysis with sufficient statistical power. Therefore, mediation analysis was conducted on the ABIDE cohort

only.

ACKNOWLEDGMENTS

The STI 2030 - the major projects of the Brain Science and Brain-Inspired Intelligence Technology (2021ZD0200500), the Start-up Funds for Leading Talents at Beijing Normal University, the National Natural Science Foundation of China (82102134, 82322035, 62273076), the Key-Area Research and Development Program of Guangdong Province (2019B030335001), the Beijing Municipal Science and Technology Commission (Z161100002616023, Z181100001518003), the Major Project of National Social Science Foundation of China (20&ZD296), the CAS-NWO Programme (153111KYSB20160020), the National Basic Research (973) Program (2015CB351702), the Chinese Academy of Sciences Key Research Program (KSZD-EW-TZ-002) and the Major Fund for International Collaboration of National Natural Science Foundation of China (81220108014) provide funding supports.

AUTHOR CONTRIBUTIONS

Conception and Design: X-R.F., X-N.Z. Data Analysis: X-R.F. MRI Data Preprocessing: Y.H., CABIC. Initial Drafting of the Manuscript: X-R.F. Critical Review and Editing of the Manuscript: All authors contributed to the critical review and editing of the manuscript.

AUTHOR COMPETING INTERESTS

Competing interests

The authors declare no competing interests.

DATA AVAILABILITY

The datasets analyzed in this study are publicly available. The ABIDE dataset can be accessed from the Autism Brain Imaging Data Exchange repository at ABIDE website. The CABIC dataset is available upon request through the CABIC Data Sharing Initiative at CABIC website.

CODE AVAILABILITY

All code used in the analyses is available at https://github.com/XueruFan/ASD-Project. For detailed information on how to use these codes, please contact the first author by email.

For the Lifespan Brain Chart Consortium (LBCC)

Chris Adamson[4,5], Sophie Adler[6], Aaron F. Alexander-Bloch[7,8,9], Evdokia Anagnostou[10,11], Kevin M. Anderson[12], Ariosky Areces-Gonzalez[13,14], Duncan E. Astle[15], Bonnie Auyeung[16,17], Muhammad Ayub[18,19], Jong Bin Bae[20], Gareth Ball[4,21], Simon Baron-Cohen[17,22], Richard Beare[4,5], Saashi A. Bedford[17], Vivek Benegal[23], Richard A.I. Bethlehem[17,24], Frauke Beyer[25], John Blangero[26], Manuel Blesa Cábez[27], James P. Boardman[27], Matthew Borzage[28], Jorge F. Bosch-Bayard[29,30], Niall Bourke[31], Edward T. Bullmore[32], Vince D. Calhoun[33], Mallar M. Chakravarty[34,30], Christina Chen[35], Casey Chertavian[9], Gaël Chetelat[36], Yap S. Chong[37,38], Aiden Corvin[39], Manuela Costantino[40,41], Eric Courchesne[42,43], Fabrice Crivello[44], Vanessa L. Cropley[45], Jennifer Crosbie[46], Nicolas Crossley[47,48], Marion Delarue[36], Richard Delorme[49,50], Sylvane Desrivieres[51], Gabriel Devenyi[52,53], Maria A. Di Biase[45,54], Ray Dolan[55,56,57], Kirsten A. Donald[58,59], Gary Donohoe[60], Lena Dorfschmidt[7,8,9], Katharine Dunlop[61], Anthony D Edwards[62,63,64], Jed T. Elison[65], Cameron T. Ellis[12,66], Jeremy A. Elman[67], Lisa Eyler[68,69], Damien A. Fair[65], Paul C. Fletcher[70,71], Peter Fonagy[72,73], Carol E. Franz[74], Lidice Galan-Garcia[75], Ali Gholipour[76], Jay Giedd[77,78], John H. Gilmore[79], David C. Glahn[80,81], Ian M. Goodyer[32], P. E. Grant[82], Nynke A. Groenewold[83,59], Shreya Gudapati[7,8,9], Faith M. Gunning[84], Raquel E. Gur[7,9], Ruben C. Gur[7,85], Christopher F. Hammill[46,86], Oskar Hansson[87,88], Trey Hedden[89,90], Andreas Heinz[91], Richard N. Henson[15,92], Katja Heuer[93], Jacqueline Hoare[94], Bharath Holla[95,96], Avram J. Holmes[97], Hao Huang[98], Jonathan Ipser[99], Clifford R. Jack Jr[100], Andrea P. Jackowski[101,102], Tianye Jia[103,104,105], David T. Jones[106,107], Peter B. Jones[32,108], Rene S Kahn[109,110], Hasse Karlsson[111,112], Linnea Karlsson[111,112], Ryuta Kawashima[113], Elizabeth A. Kelley[114], Silke Kern[115,116], Ki-Woong Kim[117,118,119,120], Manfred G. Kitzbichler[32], William S. Kremen[67], François Lalonde[121], Brigitte Landeau[36], Jason Lerch[122,123,124], John D. Lewis[125], Jiao Li[126], Wei Liao[126], Conor Liston[127], Michael V. Lombardo[128,17], Jinglei Lv[129,45], Travis T. Mallard[130], Machteld Marcelis[131], Samuel R. Mathias[80], Bernard Mazoyer[44,132], Philip McGuire[133], Michael J. Meaney[134], Andrea Mechelli[133], Bratislav Misic[135], Sarah E Morgan[136,137,138], David Mothersill[139,140,141], Cynthia Ortinau[142], Rik Ossenkoppele[143,144], Minhui Ouyang[98], Lena Palaniyappan[145], Leo Paly[36], Pedro M. Pan[146,147], Christos Pantelis[148,149,150], Min Tae M. Park[151], Tomas Paus[152,153], Zdenka Pausova[46,154], Deirel Paz-Linares[13,155], Alexa Pichet Binette[156,157], Karen Pierce[158], Xing Qian[159], Anqi Qiu[160], Armin Raznahan[121], Timothy Rittman[161], Amanda Rodrigue[80], Caitlin K. Rollins[162,163], Rafael Romero-Garcia[32,164], Lisa Ronan[32], Monica D. Rosenberg[165], David H. Rowitch[166], Giovanni A. Salum[167,168], Theodore D. Satterthwaite[169,7], H. Lina Schaare[170,171], Jenna Schabdach[7,8,9], Russell J. Schachar[46], Michael Schöll[172,173,174], Aaron P. Schultz[175,176,81], Jakob Seidlitz[7,8,9], David Sharp[177,178], Russell T. Shinohara[35,179], Ingmar Skoog[115,116], Christopher D. Smyser[180], Reisa A. Sperling[175,181,81], Dan J. Stein[182], Aleks Stolicyn[183], John Suckling[32,184], Gemma Sullivan[27], Benjamin Thyreau[113], Roberto Toro[93,185], Nicolas Traut[186,187], Kamen A. Tsvetanov[161,188], Nicholas B. Turk-Browne[12,189], Jetro J. Tuulari[190,191,192], Christophe Tzourio[193], Étienne Vachon-Presseau[194,195,196], Mitchell J. Valdes-Sosa[75], Pedro A. Valdes-Sosa[197,198], Sofie L. Valk[199], Therese van Amelsvoort[200], Simon N. Vandekar[201,202], Lana Vasung[203], Petra E Vértes[32,204], Lindsay W. Victoria[84], Sylvia Villeneuve[157,205,156], Arno Villringer[25,206], Jacob W. Vogel[169,7], Konrad Wagstyl[207], Yin-Shan Wang[1,208,209,210], Simon K. Warfield[76], Varun Warrier[211], Eric Westman[212], Margaret L. Westwater[32], Heather C. Whalley[183,213], Simon R. White[32,214], A. Veronica Witte[206,25,215], Ning Yang[1,208,209,210], B.T. Thomas Yeo[216,217,218,219], Hyuk Jin Yun[220], Andrew Zalesky[221], Heather J Zar[83,222], Anna Zettergren[115], Juan H. Zhou[159,223,216], Hisham Ziauddeen[32,224,2], Dabriel Zimmerman[7,8,9], Andre Zugman[226,227,147], Xi-Nian Zuo[1,208,209,228].

LBCC Affiliations

[4]Developmental Imaging, Murdoch Children's Research Institute, Melbourne, Victoria, Australia
[5]Department of Medicine, Monash University, Melbourne, Victoria, Australia
[6]UCL Great Ormond Street Institute for Child Health, 30 Guilford St, Holborn, London WC1N 1EH
[7]Department of Psychiatry, University of Pennsylvania, Philadelphia, PA 19104
[8]Department of Child and Adolescent Psychiatry and Behavioral Science, The Children's Hospital of Philadelphia, Philadelphia, PA 19104
[9]Lifespan Brain Institute, The Children's Hospital of Philadelphia, Philadelphia, PA 19104
[10]Department of Pediatrics University of Toronto
[11]Holland Bloorview Kids Rehabilitation Hospital, Toronto, Canada
[12]Department of Psychology, Yale University, New Haven, CT, USA
[13]The Clinical Hospital of Chengdu Brain Science Institute, MOE Key Lab for NeuroInformation, University of Electronic Science and Technology of China, No. 2006, Xiyuan Ave., West Hi-Tech Zone, Chengdu, 611731, China
[14]University of Pinar del Río Hermanos Saiz Montes de Oca, Cuba
[15]MRC Cognition and Brain Sciences Unit, University of Cambridge, Cambridge UK
[16]Department of Psychology, School of Philosophy, Psychology and Language Sciences, University of Edinburgh, Edinburgh, United Kingdom
[17]Autism Research Centre, Department of Psychiatry, University of Cambridge, Cambridge, CB2 0SZ, UK
[18]Queen's University, Department of Psychiatry, Centre for Neuroscience Studies, Kingston, Ontario, Canada
[19]University College London, Mental Health Neuroscience Research Department, Division of Psychiatry, London UK
[20]Department of Neuropsychiatry, Seoul National University Bundang Hospital, Seongnam, Korea
[21]Department of Paediatrics, University of Melbourne, Melbourne, Victoria, Australia
[22]Cambridge Lifetime Asperger Syndrome Service (CLASS), Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge, United Kingdom
[23]Centre for Addiction Medicine, National Institute of Mental Health and Neurosciences (NIMHANS), Bengaluru, India 560029
[24]Brain Mapping Unit, Department of Psychiatry, University of Cambridge, Cambridge, CB2 0SZ, UK
[25]Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, 04103, Germany
[26]Department of Human Genetics, South Texas Diabetes and Obesity Institute, University of Texas Rio Grande Valley
[27]MRC Centre for Reproductive Health, University of Edinburgh, UK
[28]Fetal and Neonatal Institute, Division of Neonatology, Children's Hospital Los Angeles, Department of Pediatrics, Keck School of Medicine, University of Southern California, Los Angeles, California USA
[29]McGill Centre for Integrative Neuroscience, Ludmer Centre for Neuroinformatics and Mental Health, Montreal Neurological Institute
[30]McGill University
[31]Department of Brain Sciences, Imperial College London, London UK & Care Research & Technology Centre, UK Dementia Research Institute
[32]Department of Psychiatry, University of Cambridge, Cambridge, CB2 0SZ, UK
[33]Tri-institutional Center for Translational Research in Neuroimaging and Data Science, Georgia State University, Georgia Institute of Technology, and Emory University, Atlanta, GA, USA
[34]Computational Brain Anatomy (CoBrA) Laboratory, Cerebral Imaging Centre, Douglas Mental Health University Institute
[35]Penn Statistics in Imaging and Visualization Center, Department of Biostatistics, Epidemiology, and Informatics, Perelman School

of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[36]Normandie Univ, UNICAEN, INSERM, U1237, PhIND "Physiopathology and Imaging of Neurological Disorders", Institut Blood and Brain @ Caen-Normandie, Cyceron, 14000 Caen, France

[37]Singapore Institute for Clinical Sciences, Agency for Science, Technology and Research, Singapore

[38]Department of Obstetrics and Gynaecology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

[39]Department of Psychiatry, Trinity College, Dublin, Ireland

[40]Cerebral Imaging Centre, Douglas Mental Health University Institute, Verdun, Canada

[41]Undergraduate program in Neuroscience, McGill University, Montreal, Canada

[42]Department of Neuroscience, University of California, San Diego, San Diego, CA 92093, USA

[43]Autism Center of Excellence, University of California, San Diego, San Diego, CA 92037, USA

[44]Institute of Neurodegenerative Disorders, CNRS UMR5293, CEA, University of Bordeaux

[45]Melbourne Neuropsychiatry Centre, University of Melbourne, Melbourne, Australia

[46]The Hospital for Sick Children, Toronto, Canada

[47]Department of Psychiatry, School of Medicine, Pontificia Universidad Católica de Chile, Diagonal Paraguay 362, Santiago 8330077, Chile

[48]Department of Psychiatry, University of Oxford OX3 7JX

[49]Child and Adolescent Psychiatry Department, Robert Debré University Hospital, AP-HP, F-75019, Paris France

[50]Human Genetics and Cognitive Functions , Institut Pasteur, F-75015, Paris France

[51]Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom

[52]Cerebral Imaging Centre, Douglas Mental Health University Institute, Montreal, QC, Canada, McGill Department of Psychiatry, Montreal, QC, Canada

[53]Department of Psychiatry, McGill University, Montreal, QC, Canada

[54]Department of Psychiatry, Brigham and Womens Hospital, Harvard Medical School, Boston, Massachusetts, United States

[55]Max Planck UCL Centre for Computational Psychiatry and Ageing Research, University College London, London, UK

[56]Wellcome Centre for Human Neuroimaging, University College London, London, UK

[57]Wellcome Centre for Human Neuroimaging, 12 Queen Square, London WC1N 3AR

[58]Division of Developmental Paediatrics, Department of Paediatrics and Child Health, Red Cross War Memorial Children's Hospital, Klipfontein Road/Private Bag, Rondebosch, 7700/7701, Cape Town, South Africa

[59]Neuroscience Institute, University of Cape Town, Cape Town, South Africa

[60]Center for Neuroimaging, Cognition & Genomics (NICOG), School of Psychology, National University of Ireland Galway, Galway, Ireland

[61]Weil Family Brain and Mind Research Institute, Department of Psychiatry, Weill Cornell Medicine

[62]Centre for the Developing Brain, King's College London, London, UK

[63]Evelina London Children's Hospital

[64]MRC Centre for Neurodevelopmental Disorders, London

[65]Institute of Child Development, Department of Pediatrics, Masonic Institute for the Developing Brain, University of Minnesota, Minneapolis, MN, United States

[66]Haskins Laboratories, New Haven, CT, USA

[67]Department of Psychiatry, Center for Behavior Genetics of Aging, University of California, San Diego, La Jolla, CA

[68]Desert-Pacific Mental Illness Research Education and Clinical Center, VA San Diego Healthcare, San Diego, CA, USA

[69]Department of Psychiatry, University of California San Diego, Los Angeles, CA, USA

[70]Department of Psychiatry, University of Cambridge, and Wellcome Trust MRC Institute of Metabolic Science, Cambridge Biomedical Campus, Cambridge, United Kingdom

[71]Cambridgeshire and Peterborough NHS Foundation Trust

[72]Department of Clinical, Educational and Health Psychology, University College London, London, UK

[73]Anna Freud National Centre for Children and Families, London UK

[74]Department of Psychiatry, Center for Behavior Genetics of Aging, University of California, San Diego, La Jolla, CA 92093

[75]Cuban Center for Neuroscience, La Habana, Cuba

[76]Computational Radiology Laboratory, Boston Childrens Hospital, Boston, MA 02115

[77]Department of Child and Adolescent Psychiatry, University of California, San Diego, San Diego, CA 92093, USA

[78]Department of Psychiatry, University of California San Diego, San Diego, CA, USA

[79]Department of Psychiatry, University of North Carolina, Chapel Hill, NC, USA

[80]Department of Psychiatry, Boston Childrens Hospital and Harvard Medical School, Boston, MA 02115

[81]Harvard Medical School, Boston, MA 02115

[82]Division of Newborn Medicine and Neuroradiology, Fetal Neonatal Neuroimaging and Developmental Science Center, Boston Childrens Hospital, Harvard Medical School, Boston, MA 02115, USA

[83]Department of Paediatrics and Child Health, Red Cross War Memorial Children's Hospital, SA-MRC Unit on Child & Adolescent Health, University of Cape Town, South Africa

[84]Weill Cornell Institute of Geriatric Psychiatry, Department of Psychiatry, Weill Cornell Medicine

[85]Lifespan Brain Institute, The Children's Hospital of Philadelphia, Philadelphia, PA 19105

[86]Mouse Imaging Centre, Toronto, Canada

[87]Clinical Memory Research Unit, Department of Clinical Sciences Malmö, Lund University, Malmö, Sweden

[88]Memory Clinic, Skåne University Hospital, Malmö, Sweden

[89]Department of Neurology, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

[90]Athinoula A. Martinos Center for Biomedical Imaging, Department of Radiology, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02129, USA

[91]Department of Psychiatry and Psychotherapy, Charite University Hospital Berlin, Berlin, Germany

[92]Department of Psychiatry, University of Cambridge, Cambridge, UK

[93]Institut Pasteur, Université Paris Cité, Unité de Neuroanatomie Appliquée et Théorique, F-75015 Paris, France

[94]Department of Psychiatry, University of Cape Town, Cape Town, South Africa

[95]Department of Integrative Medicine, NIMHANS, Bengaluru-560029, India

[96]Accelerator Program for Discovery in Brain disorders using Stem cells (ADBS), Department of Psychiatry, NIMHANS, Bengaluru-

560029, India

[97] Department of Psychiatry, Brain Health Institute, Rutgers University, Piscataway, NJ, USA

[98] Department of Radiology, Children's Hospital of Philadelphia and University of Pennsylvania, Philadelphia, PA 19104

[99] Department of Psychiatry and Mental Health, Clinical Neuroscience Institute, University of Cape Town

[100] Department of Radiology, Mayo Clinic, Rochester, MN 55905, USA

[101] Department of Psychiatry, Universidade Federal de São Paulo

[102] National Institute of Developmental Psychiatry, CNPq

[103] Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University, Shanghai, 200433, China

[104] Key Laboratory of Computational Neuroscience and BrainInspired Intelligence (Fudan University), Ministry of Education, Shanghai, China

[105] Centre for Population Neuroscience and Precision Medicine (PONS), Institute of Psychiatry, Psychology and Neuroscience, SGDP Centre, King's College London, London SE5 8AF, UK

[106] Department of Neurology, Mayo Clinic, Rochester, MN, USA

[107] Department of Radiology, Mayo Clinic, Rochester, MN, USA

[108] Cambridgeshire and Peterborough NHS Foundation Trust, Huntingdon, United Kingdom

[109] Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[110] Department of Psychiatry, Icahn School of Medicine, Mount Sinai, New York, USA

[111] Department of Clinical Medicine, Department of Psychiatry and Turku Brain and Mind Center, FinnBrain Birth Cohort Study, University of Turku and Turku University Hospital, Turku, Finland

[112] Centre for Population Health Research, Turku University Hospital and University of Turku, Turku, Finland

[113] Institute of Development, Aging and Cancer, Tohoku University, Seiryocho, Aobaku, Sendai 980-8575, Japan

[114] Queen's University, Departments of Psychology and Psychiatry, Centre for Neuroscience Studies, Kingston, Ontario, Canada

[115] Neuropsychiatric Epidemiology Unit, Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, the Sahlgrenska Academy, Centre for Ageing and Health (AGECAP) at the University of Gothenburg, Sweden

[116] Region Västra Götaland, Sahlgrenska University Hospital, Psychiatry, Cognition and Old Age Psychiatry Clinic, Gothenburg, Sweden

[117] Department of Brain and Cognitive Sciences, Seoul National University College of Natural Sciences, Seoul, Republic of Korea

[118] Department of Neuropsychiatry, Seoul National University Bundang Hospital, Seongnam, Republic of Korea

[119] Department of Psychiatry, Seoul National University College of Medicine, Seoul, Republic of Korea

[120] Department of Brain and Cognitive Science, Seoul National University College of Natural Sciences

[121] Section on Developmental Neurogenomics, Human Genetics Branch, National Institute of Mental Health, Bethesda, MD, USA

[122] Department of Medical Biophysics, University of Toronto, Toronto, ON, Canada

[123] Mouse Imaging Centre, The Hospital for Sick Children, Toronto, ON, Canada

[124] Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical Neuroscience, University of Oxford, Oxford, UK

[125] Montreal Neurological Institute, McGill University, Montreal, Canada

[126] The Clinical Hospital of Chengdu Brain Science Institute, University of Electronic Science and Technology of China, Chengdu 611731, China

[127] Department of Psychiatry and Brain and Mind Research Institute, Weill Cornell Medicine

[128] Laboratory for Autism and Neurodevelopmental Disorders, Center for Neuroscience and Cognitive Systems @UniTn, Istituto Italiano di Tecnologia, Rovereto, Italy

[129] School of Biomedical Engineering & Brain and Mind Centre, The University of Sydney, Sydney, NSW, Australia

[130] Department of Psychology, University of Texas, Austin, Texas 78712, USA

[131] Department of Psychiatry and Neuropsychology, School of Mental Health and Neuroscience, EURON, Maastricht University Medical Centre, PO Box 616, 6200 MD, Maastricht, the Netherlands; Institute for Mental Health Care Eindhoven (GGzE), Eindhoven, the Netherlands

[132] Bordeaux University Hospital

[133] Professor, Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK

[134] Ludmer Centre for Neuroinformatics and Mental Health, Douglas Mental Health University Institute, McGill University, Montreal, Quebec, Canada; Singapore Institute for Clinical Sciences, Singapore

[135] McConnell Brain Imaging Centre, Montreal Neurological Institute, McGill University, Montreal, QC H3A 2B4, Canada

[136] Department of Computer Science and Technology, University of Cambridge, Cambridge CB3 0FD, United Kingdom

[137] Department of Psychiatry, University of Cambridge, Cambridge CB2 0SZ, United Kingdom

[138] The Alan Turing Institute, London, NW1 2DB

[139] Department of Psychology, School of Business, National College of Ireland, Dublin, Ireland

[140] School of Psychology & Center for Neuroimaging and Cognitive Genomics, National University of Ireland Galway, Galway, Ireland

[141] Department of Psychiatry, Trinity College Dublin, Dublin, Ireland

[142] Department of Pediatrics, Washington University in St. Louis, St. Louis, Missouri, United States

[143] Alzheimer Center Amsterdam, Department of Neurology, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam, The Netherlands

[144] Lund University, Clinical Memory Research Unit, Lund, Sweden

[145] Robarts Research Institute & The Brain and Mind Institute, University of Western Ontario,London,Ontario,Canada

[146] Department of Psychiatry, Federal University of Sao Poalo (UNIFESP)

[147] National Institute of Developmental Psychiatry for Children and Adolescents (INPD), Brazil

[148] Melbourne Neuropsychiatry Centre, Department of Psychiatry, The University of Melbourne and Melbourne Health, Carlton South, Victoria, Australia

[149] Melbourne School of Engineering, The University of Melbourne, Parkville, Victoria, Australia

[150] Florey Institute of Neuroscience and Mental Health, Parkville, VIC, Australia

[151] Department of Psychiatry, Schulich School of Medicine and Dentistry, Western University, London, ON, Canada

[152] Department of Psychiatry, Faculty of Medicine and Centre Hospitalier Universitaire Sainte-Justine, University of Montreal, Montreal, Quebec, Canada

[153] Departments of Psychiatry and Psychology, University of Toronto, Toronto, ON, Canada

[154] Departments of Physiology and Nutritional Sciences, University of Toronto, Toronto, Canada

[155] Cuban Neuroscience Center, Havana, Cuba
[156] Department of Psychiatry, Faculty of Medicine, McGill University, Montreal, Qc, H3A 1Y2, Canada
[157] Douglas Mental Health University Institute, Montreal, Qc, H4H 1R3, Canada
[158] Department of Neurosciences, University of California, San Diego La Jolla, CA, USA
[159] Center for Sleep and Cognition, Yong Loo Lin School of Medicine, National University of Singapore, Singapore
[160] Department of Biomedical Engineering, The N.1 Institute for Health, National University of Singapore
[161] Department of Clinical Neurosciences, University of Cambridge, Cambridge UK
[162] Department of Neurology, Harvard Medical School
[163] Department of Neurology, Boston Childrens Hospital, Boston, MA 02115
[164] Instituto de Biomedicina de Sevilla (IBiS) HUVR/CSIC/Universidad de Sevilla, Dpto. de Fisiología Médica y Biofísica, Spain
[165] Department of Psychology, Neuroscience Institute, University of Chicago
[166] Department of Paediatrics and Wellcome-MRC Cambridge Stem Cell Institute, University of Cambridge, Hills Road, Cambridge, UK
[167] Department of Psychiatry, Universidade Federal do Rio Grande do Sul (UFRGS)
[168] National Institute of Developmental Psychiatry (INPD)
[169] Lifespan Informatics & Neuroimaging Center, University of Pennsylvania, Philadelphia, PA 19104
[170] Otto Hahn Group Cognitive Neurogenetics, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany
[171] Institute of Neuroscience and Medicine (INM-7: Brain and Behaviour), Research Centre Juelich, Juelich, Germany
[172] Wallenberg Centre for Molecular and Translational Medicine, University of Gothenburg, Gothenburg, Sweden
[173] Department of Psychiatry and Neurochemistry, University of Gothenburg, Sweden
[174] Dementia Research Centre, Queen's Square Institute of Neurology, University College London, UK
[175] Harvard Aging Brain Study, Department of Neurology, Massachusetts General Hospital, Boston, MA 02114
[176] Athinoula A. Martinos Center for Biomedical Imaging, Department of Radiology, Massachusetts General Hospital, Charlestown, MA 02129, USA
[177] Department of Brain Sciences, Imperial College London, London UK
[178] Care Research & Technology Centre, UK Dementia Research Institute
[179] Center for Biomedical Image Computing and Analytics, Department of Radiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
[180] Departments of Neurology, Pediatrics, and Radiology, Washington University School of Medicine, St. Louis, United States
[181] Center for Alzheimer Research and Treatment, Department of Neurology, Brigham and Womens Hospital, Boston, MA 02115
[182] SA MRC Unit on Risk & Resilience in Mental Disorders, Dept of Psychiatry and Neuroscience Institute, University of Cape Town, Cape Town, South Africa
[183] Division of Psychiatry, Centre for Clinical Brain Sciences, University of Edinburgh, UK
[184] Cambridge and Peterborough Foundation NHS Trust
[185] Université de Paris, Paris, France
[186] Department of Neuroscience, Institut Pasteur, Paris, France
[187] Center for Research and Interdisciplinarity (CRI), Université Paris Descartes, Paris, France
[188] Department of Psychology, University of Cambridge, Cambridge, UK
[189] Wu Tsai Institute, Yale University, New Haven, CT, USA
[190] Department of Clinical Medicine, Department of Psychiatry, FinnBrain Birth Cohort Study, University of Turku, Turku, Finland
[191] Department of Clinical Medicine, University of Turku, Turku Finland
[192] Turku Collegium for Science, Medicine and Technology, University of Turku, Turku, Finland
[193] Univ. Bordeaux, Inserm, Bordeaux Population Health Research Center, U1219, CHU Bordeaux, F-33000 Bordeaux, France
[194] Faculty of Dental Medicine and Oral Health Sciences, McGill University, Montreal, Qc, H3A 1G1, Canada
[195] Faculty of Dentistry, McGill University, Montreal, Qc, H3A 1G1, Canada
[196] Alan Edwards Centre for Research on Pain (AECRP), McGill University, Montreal, Qc, H3A 1G1, Canada
[197] Joint China-Cuba Lab, University of Electronic Science and Technology, Chengdu China/Cuban Center for Neuroscience, La Habana, Cuba
[198] University of Electronic Science and Technology of China/Cuban Center for Neuroscience
[199] Institute for Neuroscience and Medicine 7, Forschungszentrum Juelich; Max Planck Institute for Human Cognitive and Brain Sciences
[200] Department of Psychiatry & Neurosychology, Maastricht University, Maastricht, The Netherlands
[201] Department of Biostatistics, Vanderbilt University, Nashville, Tennessee, USA
[202] Department of Biostatistics, Vanderbilt University Medical Center, Nashville, Tennessee, USA
[203] Division of Newborn Medicine, Fetal Neonatal Neuroimaging and Developmental Science Center, Department of Pediatrics, Boston Childrens Hospital, Boston, MA 02115
[204] The Alan Turing Institute, London NW1 2DB, UK
[205] McConnell Brain Imaging Center, Montreal Neurological Institute, McGill University, Montreal, Quebec, Canada
[206] Clinic for Cognitive Neurology, University of Leipzig Medical Center, Leipzig, 04103, Germany
[207] Wellcome Centre for Human Neuroimaging, Institute of Neurology, University College London, WC1N 3AR
[208] Developmental Population Neuroscience Research Center, IDG/McGovern Institute for Brain Research, Beijing Normal University, Beijing 100875, China
[209] National Basic Science Data Center, Beijing 100190, China
[210] Research Center for Lifespan Development of Brain and Mind, Institute of Psychology, Chinese Academy of Sciences, Beijing 100101, China
[211] Department of Psychiatry, University of Cambridge, Cambridge, CB2 0SZ, UK
[212] Division of Clinical Geriatrics, Center for Alzheimer Research, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Stockholm, Sweden
[213] Generation Scotland, University of Edinburgh
[214] MRC Biostatistics Unit, University of Cambridge, Cambridge, England
[215] Faculty of Medicine, CRC 1052 Obesity Mechanisms, University of Leipzig, Leipzig, 04103, Germany
[216] Department of Electrical and Computer Engineering, National University of Singapore, Singapore
[217] Centre for Sleep & Cognition and Centre for Translational MR Research, Yong Loo Lin School of Medicine, National University of

Singapore, Singapore
[218]N.1 Institute for Health & Institute for Digital Medicine, National University of Singapore, Singapore
[219]Integrative Sciences and Engineering Programme (ISEP), National University of Singapore, Singapore
[220]Fetal Neonatal Neuroimaging and Developmental Science Center, Division of Newborn Medicine, Boston Childrens Hospital, Harvard Medical School, Boston, MA 02115, USA
[221]Melbourne Neuropsychiatry Centre, University of Melbourne, Melbourne, Australia; Department of Biomedical Engineering, University of Melbourne, Melbourne, Australia
[222]SAMRC Unit on Child & Adolescent Health, University of Cape Town, South Africa
[223]Center for Translational Magnetic Resonance Research, Yong Loo Lin School of Medicine, National University of Singapore, Singapore
[224]Wellcome Trust-MRC Institute of Metabolic Science, University of Cambridge, Cambridge, CB2 0SZ
[225]Cambridgeshire and Peterborough Foundation Trust, Cambridge, CB21 5EF
[226]National Institute of Mental Health (NIMH), National Institutes of Health (NIH), Bethesda, Maryland, USA
[227]Department of Psychiatry, Escola Paulista de Medicina, São Paulo, Brazil
[228]Research Center for Lifespan Development of Brain and Mind, Institute of Psychology, Chinese Academy of Sciences, Beijing 100101, China ; Key Laboratory of Brain and Education, School of Education Science, Nanning Normal University, Nanning 530001, China

For the China Autism Brain Imaging Consortium (CABIC)

Miaoshui Bai[229], Jinhua Cai[230], Kelong Cai[231,232], Doudou Cao[233], Xuan Cao[234], Aiguo Chen[232,235], Huafu Chen[236], Jie Chen[237], Xu-jun Duan[236], Xue-Ru Fan[238], Peng Gao[238], Wenjing Gao[239,240], DongZhi He[239], Feiyong Jia[229], Haoxiang Jiang[241], Xi Jiang[229], Jin Jing[242], Lei Li[236], Shijun Li[233], Tingyu Li[237], Xiuhong Li[243], Lizi Lin[242], Yingqiang Liu[236], Zhimei Liu[244], Fanchao Meng[245,246], Litong Ni[233], Ning Pan[247], Qi Qi[233], Bin Qin[230], Xiaolong Shan[236], Xiaojing Shou[238,245,248], Jia Wang[234], Longlun Wang[230], Miaoyan Wang[241], Wei Wang[249], Xin Wang[247], Lijie Wu[234], Dandan Xu[241], Yin Xu[237], Yang Xue[229], Ting Yang[237], Xuntao Yin[239], Rong Zhang[245,248], Yun Zhang[230], Xi-Nian Zuo[238]

CABIC Affiliations

[229] Department of developmental and behavioral pediatrics, Childrens Medical Center, The First Hospital of Jilin University, Jilin university, Changchun 130021, China.
[230] Department of Radiology, Children's Hospital of Chongqing Medical University, Chongqing, 400042, China
[231] College of Physical Education, Yangzhou University, Yangzhou 225127, China
[232] School of Sport and Brain Health, Nanjing Sport Institute, Nanjing 210014, China
[233] Department of Radiology, First Medical Center, Chinese PLA General Hospital, Beijing, 100853, China
[234] Department of Childrens and Adolescent Health, Public Health College of Harbin Medical University, Harbin, 150086, China
[235] Nanjing Sport Institute, Nanjing 210014, China
[236] The Clinical Hospital of Chengdu Brain Science Institute, School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu, 610054, China.
[237] Children's Nutrition Research Center, Ministry of Education Key Laboratory of Child Development and Disorders, National Clinical Research Center for Child Health and Disorders, China International Science and Technology Cooperation Base of Child Development and Critical Disorders, Children's Hospital of Chongqing Medical University, Chongqing, 400042, China
[238] State Key Laboratory of Cognitive Neuroscience and Learning, Developmental Population Neuroscience Research Center at IDG/McGovern Institute for Brain Research, Beijing Normal University, Beijing, 100875, China
[239] Department of Radiology, Guangzhou Women and Childrens Medical Center, Guangzhou Medical University, Guangzhou, 511400, China
[240] School of Public Health, Shenzhen, Sun Yat-sen University, 66 Gongchang Road, Guangming District 518107, Shenzhen, China
[241] Department of Radiology, Affiliated Children's Hospital of Jiangnan University, Wuxi, 214000, China
[242] Department of Maternal and Child Health, Joint International Research Laboratory of Environment and Health, Ministry of Education, Guangdong Provincial Engineering Technology Research Center of Environmental Pollution and Health Risk Assessment, School of Public Health, Sun Yat-sen University, Guangzhou 510080, China
[243] Department of Maternal and Child Health and Aging Health, School of Public Health, Shenzhen, Sun Yat-sen University, 66 Gongchang Road, Guangming District, 518107 Shenzhen, China
[244] Department of Rehabilitation Medicine, Affiliated Hospital of Yangzhou University, Yangzhou, 225001, China
[245] Department of Neurobiology, School of Basic Medical Sciences, Peking University Health Science Center, Beijing 100191, China
[246] The National Clinical Research Center for Mental Disorders & Beijing Key Laboratory of Mental Disorders, Beijing Anding Hospital, Capital Medical University, Beijing, China
[247] Key Laboratory of Brain, Cognition and Education Sciences, Ministry of Education, Institute for Brain Research and Rehabilitation, Guangdong Key Laboratory of Mental Health and Cognitive Science, South China Normal University, Guangzhou 510630, China
[248] Key Laboratory for Neuroscience, Ministry of Education / National Health Commission, Peking University, Beijing 100191, China
[249] Department of Radiology, Affiliated Hospital of Yangzhou University, Yangzhou, 225001, China

Table 1: Group comparisons of regions morphology between overall and two ASD subgroups.

| Region | OoS Scores | | | | | | AbP[1] (%) | |
| | All | | L | | H | | L | H |
	p	Median	p	Median	p	Median		
Superior frontal	0.01	0.48	0.02	0.28	0.37	0.62	5.56	2.70
Rostral middle frontal	0.01	0.46	0.02	0.25	0.32	0.60	7.94	0.68
Caudal middle frontal	0.00	0.42	0.00	0.26	0.05	0.60	6.35	2.70
Pars opercularis	0.00	0.47	0.00	0.23	0.07	0.61	6.35	4.73
Pars triangularis	0.00	0.45	0.02	0.28	0.02	0.60	6.35	1.35
Pars orbitalis	0.01	0.46	0.06	0.33	0.11	0.56	5.56	0.68
Frontal pole	0.00	0.48	0.02	0.34	0.03	0.60	11.90	2.03
Lateral orbital frontal	0.00	0.46	0.01	0.26	0.06	0.63	7.94	2.03
Medial orbital frontal	0.00	0.42	0.00	0.26	0.04	0.61	10.32	2.70
Rostral anterior cingulate	0.00	0.45	0.00	0.25	0.04	0.65	4.76	2.70
Caudal anterior cingulate	0.00	0.47	0.01	0.30	0.00	0.68	3.97	6.76
Precentral	0.00	0.45	0.01	0.26	0.17	0.60	6.35	3.38
Paracentral	0.01	0.52	0.05	0.34	0.10	0.62	4.76	5.41
Postcentral	0.00	0.48	0.01	0.29	0.05	0.64	7.14	4.05
Supramarginal	0.00	0.49	0.01	0.31	0.04	0.65	7.94	3.38
Posterior cingulate	0.00	0.46	0.02	0.29	0.04	0.64	7.94	5.41
Isthmus cingulate	0.00	0.47	0.00	0.27	0.06	0.64	3.97	4.05
Precuneus	0.00	0.52	0.04	0.29	0.02	0.70	5.56	4.73
Superior parietal	0.00	0.44	0.00	0.27	0.08	0.60	7.94	2.70
Inferior parietal	0.00	0.45	0.07	0.30	0.02	0.63	7.94	2.03
Transverse temporal	0.00	0.52	0.02	0.32	0.03	0.68	7.94	6.08
Banks superior temporal	0.00	0.44	0.00	0.26	0.02	0.59	3.97	5.41
Superior temporal	0.00	0.49	0.00	0.26	0.04	0.67	6.35	4.05
Middle temporal	0.00	0.44	0.02	0.24	0.06	0.61	17.46	2.70
Inferior temporal	0.00	0.49	0.05	0.30	0.04	0.62	11.90	0.68
Fusiform	0.00	0.45	0.01	0.26	0.09	0.59	10.32	1.35
Parahippocampal	0.00	0.49	0.02	0.33	0.01	0.63	3.17	2.70
Entorhinal	0.01	0.51	0.05	0.40	0.04	0.62	6.35	0.68
Temporal pole	0.00	0.44	0.03	0.36	0.02	0.58	10.32	1.35
Lateral occipital	0.00	0.46	0.01	0.28	0.02	0.63	9.52	2.70
Lingual	0.00	0.40	0.00	0.24	0.01	0.58	7.94	2.70
Pericalcarine	0.00	0.36	0.00	0.23	0.01	0.53	11.90	2.70
Cuneus	0.00	0.46	0.01	0.25	0.03	0.55	6.35	1.35
Insula	0.00	0.53	0.02	0.31	0.02	0.69	3.17	6.08

---

[1] AbP: Prevalence of abnormalities.

SUPPLEMENTARY INFORMATION

Supplementary Analysis

Brain Growth Charts and Age-Related Changes in OoS Scores for ASD Subgroup

To investigate potential developmental trajectories in ASD subgroups, here we illustrated the normative growth trajectories for volumes of the bilaterally averaged cortical regions with ABIDE dataset. The analysis focused on three brain regionstransverse temporal, inferior temporal, and isthmus cingulatethat exhibited reproducible correlations across both datasets. The subgroup-specific brain charts (Fig. S1e) were upgraded from the LBCC charts and adjusted by leveraging all the samples from the ABIDE. Then, we applied Generalized Additive Mixed Models (GAMM) using the *gamm4* R package to explore the relationship between OoS scores and age within subgroup H across the CABIC and ABIDE datasets.

$$\text{OoS score} \sim s(\text{Age}, \text{bs} = \text{"tp"}, \text{k} = 4) + \text{TCV}$$

Although OoS scores are adjusted for age-related effects, examining residual variability allows us to gain additional insights into brain-cognition relationships within specific subgroups.

In the transverse temporal region (Fig. S1f, left), CABIC presented a non-significant positive trend with age ($R^2 = 0.07, p = 0.56$, $F = 0.57$), while ABIDE displayed a marginally significant trend ($R^2 = 0.06, p = 0.04, F = 4.10$). For the inferior temporal region (Fig. S1f, middle), CABIC showed a moderate but non-significant trend ($R^2 = 0.20, p = 0.22, F = 1.49$), while ABIDE exhibited a flat, non-significant pattern ($R^2 = 0.07, p = 0.46, F = 0.56$). The most robust findings were observed in the isthmus cingulate (Fig. S1f, right). CABIC demonstrated a significant positive association between OoS scores and age ($R^2 = 0.13, p < 0.001, F = 72.69$), suggesting meaningful developmental changes in this region. ABIDE also showed a positive trend, albeit with a smaller effect size ($R^2 = 0.12$, $p = 0.02, F = 5.51$).

Figure S1
a, Distributions of ABIDE Out-of-Sample (OoS) scores across 34 cortical regions before clustering. **b,** Distributions of ABIDE OoS scores across 34 cortical regions after clustering. **c,** Distributions of ABIDE OoS scores across global measures before clustering. **d,** Distributions of OoS scores across global measures after clustering. **e,** ABIDE Subgroup-specific brain charts of cortical regional volumes of transverse temporal (left), inferior temporal (middle), and isthmus cingulate (right). **f,** The developmental effects of OoS scores in individuals from subgroup H were modeled using GAMM across the ABIDE and CABIC datasets. In plots **b**, **d**, and **e**, orange for subgroup H and green for subgroup L. TCV, total cortical volume; WMV, total white matter volume; GMV, total cortical gray matter volume; sGMV, total subcortical gray matter volume; CSF, total ventricular cerebrospinal fluid volume; mCT, mean cortical thickness; tSA, total surface area.

Figure S2. Brain morphology of two CABIC subgroups.
a, Distributions of Out-of-Sample (OoS) scores across 34 cortical regions after clustering. **b,** Distributions of OoS scores across global measures after clustering. TCV, total cortical volume; WMV, total white matter volume; GMV, total cortical gray matter volume; sGMV, total subcortical gray matter volume; CSF, total ventricular cerebrospinal fluid volume; mCT, mean cortical thickness; tSA, total surface area. **c,** Median OoS scores of 34 before cortical regions before clustering. **d,** Median OoS scores of ASD subgroup H (top) and L (bottom). **e,** Prevalence maps depicting the proportion of participants with extreme (top, 97.5% for subgroup H; bottom, 2.5% for subgroup L) structural anomalies. Note, the left hemispheres are plotted in **c, and d** just for visualization purposes.

Figure S3. Differences between two CABIC subgroups.
Distribution of participant ages (**a**); FIQ scales (**b**); ADI-R (**c**); ADOS (**d**); SRS (**e**); and data collection site (**f**) across the two subgroups of CABIC, with no significant difference between them. For plots **a** to **e**, the center line shows the median; the box limits represent the 25th and 75th percentiles; the whiskers show the minimum and maximum values; and the dots represent potential outliers.

Supplementary Tables

Table S1. Statistical comparison between two clustering ASD groups for different measures.

	Phenotype	Group(N)	Median	Mean	SD	Shapiro-Wilk Test	Rank-Sum Test	Two-sample t-Test	Cohen's d
Age		H (148)	10.51	10.25	1.87	0.00	0.02		-0.30
		L(126)	9.94	9.65	2.09	0.00			
IQ	PIQ	H (115)	109	109.43	18.05	0.77		0.11	-0.21
		L(111)	104	105.76	16.21	0.19			
	VIQ	H (116)	110	109.51	17.23	0.70		0.16	-0.19
		L(106)	106	106.29	16.94	0.45			
	FIQ	H (139)	110	108.66	18.39	0.26		0.09	-0.21
		L(117)	104	105.02	15.83	0.28			
ADOS-2	RRB	H (108)	3	3.06	1.84	0.00	0.94		0.02
		L(100)	3	3.09	1.85	0.00			
	Social Affect	H (107)	9	9.36	3.44	0.00	0.11		-0.18
		L(100)	8	8.73	3.76	0.00			
	Total	H (108)	12	11.62	4.75	0.00	0.15		-0.14
		L(100)	11	11.84	4.75	0.00			
	Severity	H (107)	7	7.16	1.78	0.00	0.18		-0.23
		L(100)	7	6.71	2.14	0.00			
ADI-R	RRB	H (120)	5	5.56	2.66	0.01	0.14		0.16
		L(103)	6	5.96	2.38	0.02			
	Nonverbal	H (120)	8	8.12	2.96	0.15	0.14		0.29
		L(28)	9.5	9.07	3.40	0.00			
	Verbal	H (120)	15	15.35	4.33	0.01	0.44		0.09
		L(103)	16	15.75	4.35	0.07			
	Social	H (119)	19	18.84	5.52	0.04	0.77		0.04
		L(103)	19	19.07	5.68	0.07			
SRS	Autistic Mannersims	H (67)	16	16.49	6.96	0.53	0.15		0.22
		L(69)	19	18.10	7.54	0.04			
	Motivation	H (67)	13	13.73	6.38	0.18		0.68	0.07
		L(69)	14	14.17	6.38	0.52			
	Communication	H (67)	29	30.43	10.48	0.89		0.93	-0.01
		L(69)	31	30.28	11.67	0.38			
	Cognition	H (67)	17	16.57	6.60	0.74		0.39	0.15
		L(69)	17	16.57	6.60	0.42			
	Awareness	H (67)	12	12.25	3.84	0.12		0.62	-0.09
		L(69)	11	11.93	3.73	0.39			

Table S2. Mean SHAP values for all 29 selected regions.

Region	Mean SHAP value
Isthmus cingulate	0.05
Entorhinal	0.04
Precuneus	0.04
Middle temporal	0.04
Fusiform	0.03
Transverse temporal	0.03
Rostral middle frontal	0.03
Postcentral	0.03
Inferior temporal	0.03
Rostral anterior cingulate	0.03
Supramarginal	0.02
Insula	0.02
Pericalcarine	0.02
Frontal pole	0.02
Lateral occipital	0.02
Posterior cingulate	0.02
Superior parietal	0.02
Pars opercularis	0.02
Caudal anterior cingulate	0.02
Medial orbital frontal	0.02
Lateral orbital frontal	0.02
Superior frontal	0.02
Inferior parietal	0.02
Caudal middle frontal	0.02
Cuneus	0.01
Pars orbitalis	0.01
Temporal pole	0.01
Precentral	0.01
Lingual	0.01

REFERENCES

[1] Catherine Lord, Mayada Elsabbagh, Gillian Baird, and Jeremy Veenstra-Vanderweele. Autism spectrum disorder. *The Lancet*, 392(10146):508–520, 2018.

[2] Tomoya Hirota and Bryan H. King. Autism Spectrum Disorder A Review. *JAMA*, 329(2):157–168, 2023.

[3] Jinan Zeidan, Eric Fombonne, Julie Scorah, Alaa Ibrahim, Maureen S. Durkin, Shekhar Saxena, Afiqah Yusuf, Andy Shih, and Mayada Elsabbagh. Global prevalence of autism: A systematic review update. *Autism Research*, 15(5):778–790, 2022.

[4] Matthew J. Maenner, Zachary Warren, Ashley Robinson Williams, Esther Amoakohene, Amanda V. Bakian, Deborah A. Bilder, Maureen S. Durkin, Robert T. Fitzgerald, Sarah M. Furnier, Michelle M. Hughes, Christine M. Ladd-Acosta, Dedria McArthur, Elise T. Pas, Angelica Salinas, Alison Vehorn, Susan Williams, Amy Esler, Andrea Grzy-bowski, Jennifer Hall-Lande, Ruby H. N. Nguyen, Karen Pierce, Walter Zahorodny, Allison Hudson, Libby Hallas, Kristen Clancy Mancilla, Mary Patrick, Josephine Shenouda, Kate Sidwell, Monica DiRienzo, Johanna Gutierrez, Margaret H. Spivey, Maya Lopez, Sydney Pettygrove, Yvette D. Schwenk, Anita Washington, and Kelly A. Shaw. Prevalence and Characteristics of Autism Spectrum Disorder Among Children Aged 8 Years - Autism and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2020. *MMWR Surveillance Summaries*, 72(2):1–14, 2023.

[5] Hao Zhou, Xiu Xu, Weili Yan, Xiaobing Zou, Lijie Wu, Xuerong Luo, Tingyu Li, Yi Huang, Hongyan Guan, Xiang Chen, Meng Mao, Kun Xia, Lan Zhang, Erzhen Li, Xiaoling Ge, Lili Zhang, Chunpei Li, Xudong Zhang, Yuanfeng Zhou, Ding Ding, Andy Shih, Eric Fombonne, Yi Zheng, Jisheng Han, Zhongsheng Sun, Yong-hui Jiang, Yi Wang, and LATENT-NHC Study Team. Prevalence of Autism Spectrum Disorder in China: A Nationwide Multi-center Population-based Study Among Children Aged 6 to 12 Years. *Neuroscience Bulletin*, 36(9):961–971, 2020.

[6] Eric Fombonne. Editorial: The rising prevalence of autism. *Journal of Child Psychology and Psychiatry*, 59(7):717–720, 2018.

[7] Francesca Happé and Uta Frith. Annual Research Review: Looking back to look forward – changes in the concept of autism and implications for future research. *Journal of Child Psychology and Psychiatry*, 61(3):218–232, 2020.

[8] Juliette Rabot, Eya-Mist Rodgaard, Ridha Joober, Guillaume Dumas, Danilo Bzdok, Boris Bernhardt, Sebastien Jacquemont, and Laurent Mottron. Genesis, modelling and methodological remedies to autism heterogeneity. *Neuroscience & Biobehavioral Reviews*, 150, 2023.

[9] Olof Arvidsson, Christopher Gillberg, Paul Lichtenstein, and Sebastian Lundström. Secular changes in the symptom level of clinically diagnosed autism. *Journal of Child Psychology and Psychiatry*, 59(7):744–751, 2018.

[10] Daan van Rooij, Evdokia Anagnostou, Celso Arango, Guillaume Auzias, Marlene Behrmann, Geraldo F. Busatto, Sara Calderoni, Eileen Daly, Christine Deruelle, Adriana Di Martino, Ilan Dinstein, Fabio Luis Souza Duran, Sarah Durston, Christine Ecker, Damien Fair, Jennifer Fedor, Jackie Fitzgerald, Christine M. Freitag, Louise Gallagher, Ilaria Gori, Shlomi Haar, Liesbeth Hoekstra, Neda Jahanshad, Maria Jalbrzikowski, Joost Janssen, Jason Lerch, Beatriz Luna, Mauricio Moller Martinho, Jane McGrath, Fil-ippo Muratori, Clodagh M. Murphy, Declan G.M. Murphy, Kirsten O'Hearn, Bob Oranje, Mara Parellada, Alessandra Retico, Pedro Rosa, Katya Rubia, Devon Shook, Margot Taylor, Paul M. Thompson, Michela Tosetti, Gregory L. Wallace, Fengfeng Zhou, and Jan K. Buitelaar. Cortical and Subcortical Brain Morphometry Differences Between Patients With Autism Spectrum Disorder and Healthy Individuals Across the Lifespan: Results From the ENIGMA ASD Working Group. *American Journal of Psychiatry*, 175(4):359–369, 2018.

[11] Michael V. Lombardo, Meng-Chuan Lai, and Simon Baron-Cohen. Big data approaches to decomposing heterogeneity across the autism spectrum. *Molecular Psychiatry*, 24(10):1435–1450, 2019.

[12] Anne Masi, Marilena M. DeMayo, Nicholas Glozier, and Adam J. Guastella. An Overview of Autism Spectrum Disorder, Heterogeneity and Treatment Options. *Neuroscience Bulletin*, 33(2):183–193, 2017.

[13] Micheal Sandbank, Kristen Bottema-Beutel, Shannon Crowley, Margaret Cassidy, Kacie Dunham, Jacob I. Feldman, Jenna Crank, Susanne A. Albarran, Sweeya Raj, Prachy Mahbub, and Tiffany G. Woynaroski. Project AIM: Autism intervention meta-analysis for studies of young children. *Psychological Bulletin*, 146(1):1–29, 2020.

[14] Andre F. Marquand, Seyed Mostafa Kia, Mariam Zabihi, Thomas Wolfers, Jan K. Buitelaar, and Christian F. Beckmann. Conceptualizing mental disorders as deviations from normative functioning. *Molecular Psychiatry*, 24(10):1415–1424, 2019.

[15] Eya-Mist Rødgaard, Kristian Jensen, Jean-Noël Vergnes, Isabelle Soulières, and Laurent Mottron. Temporal Changes in Effect Sizes of Studies Comparing Individuals With and Without Autism: A Meta-analysis. *JAMA Psychiatry*, 76(11):1124–1132, 2019.

[16] Shuting Zheng, Kara A. Hume, Harriet Able, Somer L. Bishop, and Brian A. Boyd. Exploring Developmental and Behavioral Heterogeneity among Preschoolers with ASD: A Cluster Analysis on Principal Components. *Autism Research*, 13(5):796–809, 2020.

[17] Heng Chen, Lucina Q. Uddin, Xiaonan Guo, Jia Wang, Runshi Wang, Xiaomin Wang, Xujun Duan, and Huafu Chen. Parsing brain structural heterogeneity in males with autism spectrum disorder reveals distinct clinical subtypes. *Human Brain Mapping*, 40(2):628–637, 2019.

[18] Xiaolong Shan, Lucina Q. Uddin, Jinming Xiao, Changchun He, Zihan Ling, Lei Li, Xinyue Huang, Huafu Chen, and Xujun Duan. Mapping the Heterogeneous Brain Struc-tural Phenotype of Autism Spectrum Disorder Using the Normative Model. *Biological Psychiatry*, 91(11):967–976, 2022.

[19] Guanlu Liu, Liting Shi, Jianfeng Qiu, and Weizhao Lu. Two neuroanatomical subtypes of males with autism spectrum disorder revealed using semi-supervised machine learning. *Molecular Autism*, 13(1):9, 2022.

[20] Mariam Zabihi, Dorothea L. Floris, Seyed Mostafa Kia, Thomas Wolfers, Julian Tillmann, Alberto Llera Arenas, Carolin Moessnang, Tobias Banaschewski, Rosemary Holt, Simon Baron-Cohen, Eva Loth, Tony Charman, Thomas Bourgeron, Declan Murphy, Christine Ecker, Jan K. Buitelaar, Christian F. Beckmann, Andre Marquand, and The EU-AIMS LEAP Group. Fractionating autism based on neuroanatomical normative modeling. *Translational Psychiatry*, 10(1):384, 2020.

[21] Seok-Jun Hong, Joshua T. Vogelstein, Alessandro Gozzi, Boris C. Bernhardt, B.T. Thomas Yeo, Michael P. Milham, and Adriana Di Martino. Toward Neurosubtypes in Autism. *Biological Psychiatry*, 88(1):111–128, 2020.

[22] Adriana Di Martino, Damien A. Fair, Clare Kelly, Theodore D. Satterthwaite, F. Xavier Castellanos, Moriah E. Thomason, R. Cameron Craddock, Beatriz Luna, Bennett L. Lev-enthal, Xi-Nian Zuo, and Michael P. Milham. Unraveling the Miswired Connectome: A Developmental Perspective. *Neuron*, 83(6):1335–1353, 2014.

[23] Ginny Russell, William Mandy, Daisy Elliott, Rhianna White, Tom Pittwood, and Tamsin Ford. Selection bias on intellectual ability in autism research: A cross-sectional re-view and meta-analysis. *Molecular Autism*, 10(1):9, 2019.

[24] Eric Feczko, Oscar Miranda-Dominguez, Mollie Marr, Alice M. Graham, Joel T. Nigg, and Damien A. Fair. The Heterogeneity Problem: Approaches to Identify Psychiatric Subtypes. *Trends in Cognitive Sciences*, 23(7):584–601, 2019.

[25] Saashi A. Bedford, Min Tae M. Park, Gabriel A. Devenyi, Stephanie Tullo, Jurgen Germann, Raihaan Patel, Evdokia Anagnostou, Simon Baron-Cohen, Edward T. Bullmore, Lindsay R. Chura, Michael C. Craig, Christine Ecker, Dorothea L. Floris, Rosemary J. Holt, Rhoshel Lenroot, Jason P. Lerch, Michael V. Lombardo, Declan G. M. Murphy, Armin Raznahan, Amber N. V. Ruigrok, Elizabeth Smith, Michael D. Spencer, John Suckling, Margot J. Taylor, Audrey Thurm, Meng-Chuan Lai, M. Mallar Chakravarty, and MRC AIMS Consortium. Large-scale analyses of the relationship between sex, age and intelligence quotient heterogeneity and cortical morphometry in autism spectrum disor-der. *Molecular Psychiatry*, 25(3):614–628, 2020.

[26] Meng-Chuan Lai, Caroline Kassee, Richard Besney, Sarah Bonato, Laura Hull, William Mandy, Peter Szatmari, and Stephanie H Ameis. Prevalence of co-occurring mental health diagnoses in the autism population: A systematic review and meta-analysis. *The Lancet Psychiatry*, 6(10):819–829, 2019.

[27] Ye He, Lisa Byrge, and Daniel P. Kennedy. Nonreplication of functional connectivity differences in autism spectrum disorder across multiple sites and denoising strategies. *Human Brain Mapping*, 41(5):1334–1350, 2020.

[28] Andre F. Marquand, Iead Rezek, Jan Buitelaar, and Christian F. Beckmann. Understanding Heterogeneity in Clinical Cohorts Using Normative Models: Beyond Case-Control Studies. *Obsessive-Compulsive Disorder*, 80(7):552–561, 2016.

[29] Huaihou Chen, Clare Kelly, F. Xavier Castellanos, Ye He, Xi-Nian Zuo, and Philip T. Reiss. Quantile rank maps: A new tool for understanding individual brain development. *NeuroImage*, 111:454–463, 2015.

[30] Mariam Zabihi, Marianne Oldehinkel, Thomas Wolfers, Vincent Frouin, David Goyard, Eva Loth, Tony Charman, Julian Tillmann, Tobias Banaschewski, Guillaume Dumas, Rosemary Holt, Simon Baron-Cohen, Sarah Durston, Sven Bölte, Declan Murphy, Christine Ecker, Jan K. Buitelaar, Christian F. Beckmann, and Andre F. Marquand. Dis-secting the Heterogeneous Cortical Anatomy of Autism Spectrum Disorder Using Normative Models. *The Bridging of Scales: Techniques for Translational Neuroscience*, 4(6):567–578, 2019.

[31] Richard A. I. Bethlehem, Jakob Seidlitz, Rafael Romero-Garcia, Stavros Trakoshis, Guillaume Dumas, and Michael V. Lombardo. A normative modelling approach reveals age-atypical cortical thickness in a subgroup of males with autism spectrum disorder. *Communications Biology*, 3(1):486, 2020.

[32] R. A. I. Bethlehem, J. Seidlitz, S. R. White, J. W. Vogel, K. M. Anderson, C. Adamson, S. Adler, G. S. Alexopoulos, E. Anagnostou, A. Areces-Gonzalez, D. E. Astle, B. Auyeung, M. Ayub, J. Bae, G. Ball, S. Baron-Cohen, R. Beare, S. A. Bedford, V. Benegal, F. Beyer, J. Blangero, M. Blesa Cábez, J. P. Boardman, M. Borzage, J. F. Bosch-Bayard, N. Bourke, V. D. Calhoun, M. M. Chakravarty, C. Chen, C. Chertavian, G. Chetelat, Y. S. Chong, J. H. Cole, A. Corvin, M. Costantino, E. Courchesne, F. Crivello, V. L. Cropley, J. Crosbie, N. Crossley, M. Delarue, R. Delorme, S. Desrivieres, G. A. Devenyi, M. A. Di Biase, R. Dolan, K. A. Donald, G. Donohoe, K. Dunlop, A. D. Edwards, J. T. Elison, C. T. Ellis, J. A. Elman, L. Eyler, D. A. Fair, E. Feczko, P. C. Fletcher, P. Fonagy, C. E. Franz, L. Galan-Garcia, A. Gholipour, J. Giedd, J. H. Gilmore, D. C. Glahn, I. M. Goodyer, P. E. Grant, N. A. Groenewold, F. M. Gunning, R. E. Gur, R. C. Gur, C. F. Hammill, O. Hansson, T. Hedden, A. Heinz, R. N. Henson, K. Heuer, J. Hoare, B. Holla, A. J. Holmes, R. Holt, H. Huang, K. Im, J. Ipser, C. R. Jack, A. P. Jackowski, T. Jia, K. A. Johnson, P. B. Jones, D. T. Jones, R. S. Kahn, H. Karlsson, L. Karlsson, R. Kawashima, E. A. Kelley, S. Kern, K. W. Kim, M. G. Kitzbichler, W. S. Kremen, F. Lalonde, B. Landeau, S. Lee, J. Lerch, J. D. Lewis, J. Li, W. Liao, C. Liston, M. V. Lombardo, J. Lv, C. Lynch, T. T. Mallard, M. Marcelis, R. D. Markello, S. R. Mathias, B. Ma-zoyer, P. McGuire, M. J. Meaney, A. Mechelli, N. Medic, B. Misic, S. E. Morgan, D. Mothersill, J. Nigg, M. Q. W. Ong, C. Ortinau, R. Ossenkoppele, M. Ouyang, L. Palaniyappan, L. Paly, P. M. Pan, C. Pantelis, M. M. Park, T. Paus, Z. Pausova, D. Paz-Linares, A. Pichet Binette, K. Pierce, X. Qian, J. Qiu, A. Qiu, A. Raznahan, T. Rittman, A. Rodrigue, C. K. Rollins, R. Romero-Garcia, L. Ronan, M. D. Rosenberg, D. H. Rowitch, G. A. Salum, T. D. Satterthwaite, H. L. Schaare, R. J. Schachar, A. P. Schultz, G. Schumann, M. Schöll, D. Sharp, R. T. Shinohara, I. Skoog, C. D. Smyser, R. A. Sperling, D. J. Stein, A. Stolicyn, J. Suckling, G. Sullivan, Y. Taki, B. Thyreau, R. Toro, N. Traut, K. A. Tsvetanov, N. B. Turk-Browne, J. J. Tuulari, C. Tzourio, É. Vachon-Presseau, M. J. Valdes-Sosa, P. A. Valdes-Sosa, S. Valk, T. van Amelsvoort, S. N. Vandekar, L. Vasung, L. W. Victoria, S. Villeneuve, A. Villringer, P. E. Vértes, K. Wagstyl, Y. S. Wang, S. K. Warfield, V. Warrier, E. Westman, M. L. Westwater, H. C. Whalley, A. V. Witte, N. Yang, B. Yeo, H. Yun, A. Zalesky, H. J. Zar, A. Zettergren, J. H. Zhou, H. Ziauddeen, A. Zugman, X. N. Zuo, C. Rowe, G. B. Frisoni, A. Pichet Binette, E. T. Bullmore, A. F. Alexander-Bloch, 3R-BRAIN, AIBL, Alzheimer's Dis-ease Neuroimaging Initiative, Alzheimer's Disease Repository Without Borders Investigators, CALM Team, Cam-CAN, CCNP, COBRE, cVEDA, ENIGMA Developmental Brain Age Working Group, Developing Human Connectome Project, FinnBrain, Harvard Aging Brain Study, IMAGEN, KNE96, The Mayo Clinic Study of Aging, NSPN, POND, The PREVENT-AD Research Group, and VETSA. Brain charts for the human lifespan. *Nature*, 604(7906):525–533, 2022.

[33] Lei Li, Miaoshui Bai, Kelong Cai, Doudou Cao, Xuan Cao, Jie Chen, Xue-Ru Fan, Peng Gao, Wenjing Gao, Dongzhi He, Fanchao Meng, Xi Jiang, Litong Ni, Xiuhong Li, Lizi Lin, Yingqiang Liu, Zhimei Liu, Ning Pan, Qi Qi, Bin Qin, Xiaolong Shan, Xiaojing Shou, Longlun Wang, Miaoyan Wang, Xin Wang, Dandan Xu, Yin Xu, Yang Xue, Ting Yang, Yun Zhang, Jinhua Cai, Huafu Chen, Aiguo Chen, Feiyong Jia, Haoxiang Jiang, Jin Jing, Tingyu Li, Shijun Li, Wei Wang, Jia Wang, Lijie Wu, Xuntao Yin, Rong Zhang, Xi-Nian Zuo, China Autism Brain Imaging Consortium, and Xujun Duan. China Autism Brain Imaging Consortium: Charting brain growth in chinese children with autism. *bioRxiv*, 2025.02.20.639044, 2025.

[34] A. Di Martino, C-G Yan, Q. Li, E. Denio, F X Castellanos, K. Alaerts, J S Anderson, M. Assaf, S Y Bookheimer, M. Dapretto, B. Deen, S. Delmonte, I. Dinstein, B. Ertl-Wagner, D A Fair, L. Gallagher, D P Kennedy, C L Keown, C. Keysers, J E Lainhart, C. Lord, B. Luna, V. Menon, N J Minshew, C S Monk, S. Mueller, R-A Müller, M B Nebel, J T Nigg, K. O'Hearn, K A Pelphrey, S J Peltier, J D Rudie, S. Sunaert, M. Thioux, J M Tyszka, L Q Uddin, J S Verhoeven, N. Wenderoth, J L Wiggins, S H Mostofsky, and M P Milham. The autism brain imaging data exchange: Towards a large-scale evaluation of the intrinsic brain architecture in autism. *Molecular Psychiatry*, 19(6):659–667, 2014.

[35] Adriana Di Martino, David O'Connor, Bosi Chen, Kaat Alaerts, Jeffrey S. Anderson, Michal Assaf, Joshua H. Balsters, Leslie Baxter, Anita Beggiato, Sylvie Bernaerts, Laura M. E. Blanken, Susan Y. Bookheimer, B. Blair Braden, Lisa Byrge, F. Xavier Castellanos, Mirella Dapretto, Richard Delorme, Damien A. Fair, Inna Fishman, Jacqueline Fitzgerald, Louise Gallagher, R. Joanne Jao Keehn, Daniel P. Kennedy, Janet E. Lainhart, Beatriz Luna, Stewart H. Mostofsky, Ralph-Axel Müller, Mary Beth Nebel, Joel T. Nigg, Kirsten O'Hearn, Marjorie Solomon, Roberto Toro, Chandan J. Vaidya, Nicole Wenderoth, Tonya White, R. Cameron Craddock, Catherine Lord, Bennett Leventhal, and Michael P. Milham. Enhancing studies of the connectome in autism using the autism brain imaging data exchange II. *Scientific Data*, 4(1):170010, 2017.

[36] Rahul S. Desikan, Florent Ségonne, Bruce Fischl, Brian T. Quinn, Bradford C. Dickerson, Deborah Blacker, Randy L. Buckner, Anders M. Dale, R. Paul Maguire, Bradley T. Hyman, Marilyn S. Albert, and Ronald J. Killiany. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*, 31(3):968–980, 2006.

[37] Roberto Sacco, Stefano Gabriele, and Antonio M. Persico. Head circumference and brain size in autism spectrum disorder: A systematic review and meta-analysis. *Psychiatry Research: Neuroimaging*, 234(2):239–251, 2015.

[38] Eric Courchesne, Karen Pierce, Cynthia M. Schumann, Elizabeth Redcay, Joseph A. Buckwalter, Daniel P. Kennedy, and John Morgan. Mapping Early Brain Development in Autism. *Neuron*, 56(2):399–413, 2007.

[39] Fréderique Bonnet-Brilhault, Toky A. Rajerison, Christian Paillet, Marie Guimard-Brunault, Agathe Saby, Laura Ponson, Gabriele Tripi, Joëlle Malvy, and Sylvie Roux. Autism is a prenatal disorder: Evidence from late gestation brain overgrowth. *Autism Research*, 11(12):1635–1642, 2018.

[40] Eric Courchesne, Vani Taluja, Sanaz Nazari, Caitlin M. Aamodt, Karen Pierce, Kuaikuai Duan, Sunny Stophaeros, Linda Lopez, Cynthia Carter Barnes, Jaden Troxel, Kathleen Campbell, Tianyun Wang, Kendra Hoekzema, Evan E. Eichler, Joao V. Nani, Wirla Pontes, Sandra Sanchez Sanchez, Michael V. Lombardo, Janaina S. de Souza, Mirian A. F. Hayashi, and Alysson R. Muotri. Embryonic origin of two ASD subtypes of social symptom severity: The larger the brain cortical organoid size, the more severe the social symptoms. *Molecular Autism*, 15(1):22, 2024.

[41] Eric Courchesne, Tiziano Pramparo, Vahid H. Gazestani, Michael V. Lombardo, Karen Pierce, and Nathan E. Lewis. The ASD Living Biology: From cell proliferation to clinical phenotype. *Molecular Psychiatry*, 24(1):88–107, 2019.

[42] Antonio Del Casale, Stefano Ferracuti, Alessandro Alcibiade, Sara Simone, Martina Nicole Modesti, and Maurizio Pompili. Neuroanatomical correlates of autism spectrum disorders: A meta-analysis of structural magnetic resonance imaging (MRI) studies. *Psychiatry Research: Neuroimaging*, 325:111516, 2022.

[43] Silvia Alemany, Elisabet Blok, Philip R. Jansen, Ryan L. Muetzel, and Tonya White. Brain morphology, autistic traits, and polygenic risk for autism: A population-based neuroimaging study. *Autism Research*, 14(10):2085–2099, 2021.

[44] Sarah A. Ashley, Kate Merritt, Francesca Solmi, Pedro L. Laguna, Abraham Reichenberg, and Anthony S. David. A Longitudinal Study of Head Circumference Trajectories in Autism and Autistic Traits. *Journal of Autism and Developmental Disorders*, 2024.

[45] Michael von Rhein, Andreas Buchmann, Cornelia Hagmann, Hitendu Dave, Vera Bernet, Ianina Scheer, Walter Knirsch, Beatrice Latal, Vera Bernet, Christoph Bürki, Hitendu Dave, René Prêtre, Cornelia Hagmann, Walter Knirsch, Oliver Kretschmar, Christian Kellenberger, Ianina Scheer, Beatrice Latal, Rabia Liamlahi, Michael von Rhein, Barbara R. Plecko, and Felix H. Sennhauser. Severe Congenital Heart Defects Are Associated with Global Reduction of Neonatal Brain Volumes. *The Journal of Pediatrics*, 167(6):1259–1263.e1, 2015.

[46] Eric R. Sigmon, Michael Kelleman, Apryl Susi, Cade M. Nylund, and Matthew E. Oster. Congenital Heart Disease and Autism: A Case-Control Study. *Pediatrics*, 144(5):e20184114, 2019.

[47] Cherie Green, Cheryl Dissanayake, and Danuta Loesch. A review of physical growth in children and adolescents with Autism Spectrum Disorder. *Developmental Review*, 36:156–178, 2015.

[48] Stoner Rich, Chow Maggie L., Boyle Maureen P., Sunkin Susan M., Mouton Peter R., Roy Subhojit, Wynshaw-Boris Anthony, Colamarino Sophia A., Lein Ed S., and Courchesne Eric. Patches of Disorganization in the Neocortex of Children with Autism. *New England Journal of Medicine*, 370(13):1209–1219, 2014.

[49] Telencephalon: Neocortex. In Rudolf Nieuwenhuys, Jan Voogd, and Christiaan van Huijzen, editors, *The Human Central Nervous System*, pages 491–679. Springer Berlin Heidelberg, Berlin, Heidelberg, 2008.

[50] Jorge Sepulcre. Functional Streams and Cortical Integration in the Human Brain. *The Neuroscientist*, 20(5):499–508, 2014.

[51] Kenia Martínez, Magdalena Martínez-García, Luis Marcos-Vidal, Joost Janssen, Francisco X. Castellanos, Clara Pretus, Óscar Villarroya, Laura Pina-Camacho, Covadonga M. Díaz-Caneja, Mara Parellada, Celso Arango, Manuel Desco, Jorge Sepulcre, and Susanna Carmona. Sensory-to-Cognitive Systems Integration Is Associated With Clinical Severity in Autism Spectrum Disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 59(3):422–433, 2020.

[52] Daniel S. Margulies, Satrajit S. Ghosh, Alexandros Goulas, Marcel Falkiewicz, Julia M. Huntenburg, Georg Langs, Gleb Bezgin, Simon B. Eickhoff, F. Xavier Castellanos, Michael Petrides, Elizabeth Jefferies, and Jonathan Smallwood. Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proceedings of the National Academy of Sciences*, 113(44):12574–12579, 2016.

[53] B. T. Thomas Yeo, Fenna M. Krienen, Jorge Sepulcre, Mert R. Sabuncu, Danial Lashkari, Marisa Hollinshead, Joshua L. Roffman, Jordan W. Smoller, Lilla Zöllei, Jonathan R. Polimeni, Bruce Fischl, Hesheng Liu, and Randy L. Buckner. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, 106(3):1125–1165, 2011.

[54] Jingnan Du, Lauren M. DiNicola, Peter A. Angeli, Noam Saadon-Grosman, Wendy Sun, Stephanie Kaiser, Joanna Ladopoulou, Aihuiping Xue, B. T. Thomas Yeo, Mark C. Eldaief, and Randy L. Buckner. Organization of the human cerebral cortex estimated within individuals: Networks, global topography, and function. *Journal of Neurophysiology*, 131(6):1014–1082, 2024.

[55] Yvonne Schröder, Daniela Michelle Hohmann, Tina Meller, Ulrika Evermann, Julia-Katharina Pfarr, Andreas Jansen, Inge Kamp-Becker, Sarah Grezellschak, and Igor Nenadić. Associations of subclinical autistic-like traits with brain structural variation using diffusion tensor imaging and voxel-based morphometry. *European Psychiatry*, 64(1):e27, 2021.

[56] Hao-Ming Dong, Daniel S. Margulies, Xi-Nian Zuo, and Avram J. Holmes. Shifting gradients of macroscale cortical organization mark the transition from childhood to adolescence. *Proceedings of the National Academy of Sciences*, 118(28):e2024448118, 2021.

[57] C Gerlach, C.T Aaside, G.W Humphreys, A Gade, O.B Paulson, and I Law. Brain activity related to integrative processes in visual object recognition: Bottom-up integration and the modulatory influence of stored knowledge. *Neuropsychologia*, 40(8):1254–1267, 2002.

[58] Catherine Lord, Eva Petkova, Vanessa Hus, Weijin Gan, Feihan Lu, Donna M. Martin, Opal Ousley, Lisa Guy, Raphael Bernier, Jennifer Gerdts, Molly Algermissen, Agnes Whitaker, James S. Sutcliffe, Zachary Warren, Ami Klin, Celine Saulnier, Ellen Hanson, Rachel Hundley, Judith Piggot, Eric Fombonne, Mandy Steiman, Judith Miles, Stephen M. Kanne, Robin P. Goin-Kochel, Sarika U. Peters, Edwin H. Cook, Stephen Guter, Jennifer Tjernagel, Lee Anne Green-Snyder, Somer Bishop, Amy Esler, Katherine Gotham, Rhiannon Luyster, Fiona Miller, Jennifer Olson, Jennifer Richler, and Susan Risi. A Multisite Study of the Clinical Diagnosis of Different Autism Spectrum Disorders. *Archives of General Psychiatry*, 69(3):306–313, 2012.

[59] Seulki Yoo, Yurim Jang, Seok-Jun Hong, Hyunjin Park, Sofie L. Valk, Boris C. Bernhardt, and Bo-yong Park. Whole-brain structural connectome asymmetry in autism. *NeuroImage*, 288, 2024.

[60] Laurent Mottron. Progress in autism research requires several recognition-definition-investigation cycles. *Autism Research*, 14(10):2230–2234, 2021.

[61] Allison Jack and Kevin A. Pelphrey. Annual Research Review: Understudied populations within the autism spectrum – current trends and future directions in neuroimaging research. *Journal of Child Psychology and Psychiatry*, 58(4):411–435, 2017.

[62] John N Constantino. The Quantitative Nature of Autistic Social Impairment. *Pediatric Research*, 69(8):55–62, 2011.

[63] Jeremy Veenstra-VanderWeele, Kally C. O'Reilly, Megan Y. Dennis, José M. Uribe-Salazar, and David G. Amaral. Translational Neuroscience Approaches to Understanding Autism. *American Journal of Psychiatry*, 180(4):265–276, 2023.

[64] Oscar Esteban, Daniel Birman, Marie Schaer, Oluwasanmi O. Koyejo, Russell A. Poldrack, and Krzysztof J. Gorgolewski. MRIQC: Advancing the automatic prediction of image quality in MRI from unseen sites. *PLOS ONE*, 12(9):e0184661, 2017.

[65] Xiu-Xia Xing, Ting Xu, Chao Jiang, Yin-Shan Wang, and Xi-Nian Zuo. Connectome Computation System: 2015–2021 updates. *Science Bulletin*, 67(5):448–451, 2022.

[66] Ting Xu, Zhi Yang, Lili Jiang, Xiu-Xia Xing, and Xi-Nian Zuo. A Connectome Computation System for discovery science of brain. *Science Bulletin*, 60(1):86–95, 2015.

[67] Jean-Philippe Fortin, Nicholas Cullen, Yvette I. Sheline, Warren D. Taylor, Irem Aselcioglu, Philip A. Cook, Phil Adams, Crystal Cooper, Maurizio Fava, Patrick J. McGrath, Melvin McInnis, Mary L. Phillips, Madhukar H. Trivedi, Myrna M. Weissman, and Russell T. Shinohara. Harmonization of cortical thickness measurements across scanners and sites. *NeuroImage*, 167:104–120, 2018.

[68] E. Borghi, M. de Onis, C. Garza, J. Van den Broeck, E. A. Frongillo, L. Grummer-Strawn, S. Van Buuren, H. Pan, L. Molinari, R. Martorell, A. W. Onyango, J. C. Martines, and for the WHO Multicentre Growth Reference Study Group. Construction of the World Health Organization child growth standards: Selection of methods for attained growth curves. *Statistics in Medicine*, 25(2):247–265, 2006.

[69] Ulrike von Luxburg. A tutorial on spectral clustering. *Statistics and Computing*, 17(4):395–416, 2007.

[70] Peter J. Rousseeuw. Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. *Journal of Computational and Applied Mathematics*, 20:53–65, 1987.

[71] Corinna Cortes and Vladimir Vapnik. Support-vector networks. *Machine Learning*, 20(3):273–297, 1995.

[72] Isabelle Guyon, Jason Weston, Stephen Barnhill, and Vladimir Vapnik. Gene Selection for Cancer Classification using Support Vector Machines. *Machine Learning*, 46(1):389–422, 2002.

[73] Scott M. Lundberg and Su-In Lee. A unified approach to interpreting model predictions. In *Proceedings of the 31st International Conference on Neural Information Processing Systems*, NIPS'17, page 47684777, Red Hook, NY, USA, 2017. Curran Associates Inc.

[74] Xujun Duan, Runshi Wang, Jinming Xiao, Ya Li, Xinyue Huang, Xiaonan Guo, Jing Cao, Liyao He, Changchun He, Zihan Ling, Xiaolong Shan, Heng Chen, Xiaodong Kang, and Huafu Chen. Subcortical structural covariance in young children with autism spectrum disorder. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 99:109874, 2020.