Biophysical validation of explainable AI for functional brain imaging: bridging cellular

mechanisms and network dynamics

Anthony Strock[1,*], Trang-Anh E. Nghiem[1,*], Alex De Lecea[5], Atif Hassan[6], Srikanth Ryali[1],

Vinod Menon[1,2,3,4]

[1] Department of Psychiatry & Behavioral Sciences, Stanford University, CA 94305, USA.

[2] Department of Neurology & Neurological Sciences, Stanford University, CA 94305, USA.

[3] Wu Tsai Neurosciences Institute, Stanford University, CA 94305, USA.

[4] Graduate School of Education, Stanford University, CA 94305, USA.

[5] Department of Mathematics, University of Washington, Seattle, WA 98195, USA

[6] Department of Artificial Intelligence, IIT Kharagpur, India

[*] Equal contribution

Address correspondence to:

astrock@stanford.edu; menon@stanford.edu

Abstract

Deep neural networks have revolutionized functional neuroimaging analysis but remain "black boxes," concealing which brain mechanisms and regions drive their predictions—a critical limitation for clinical neuroscience. Here we develop and validate an explainable AI (xAI) framework to test whether feature attribution techniques can reliably recover brain regions affected by excitation/inhibition (E/I) imbalance—a fundamental dysregulation implicated in autism, schizophrenia, and other neuropsychiatric disorders. We employed complementary simulation approaches: recurrent neural networks for controlled parameter exploration, and The Virtual Brain simulator incorporating empirically-derived human and mouse connectomes to model E/I balance alterations with unprecedented biological realism. Through systematic validation, we demonstrate that Integrated Gradients and DeepLIFT methods reliably identify brain regions affected by E/I imbalance across challenging conditions, including high noise, low prevalence, and subtle neurophysiological alterations. This performance remains robust across species and anatomical scales, from 68-region human to 426-region mouse connectomes. Application to the ABIDE autism dataset (N=834) reveal convergence between our biophysically grounded simulations and empirical findings, providing computational support for E/I imbalance mechanisms in autism. This work establishes essential tools and data for interpretation of deep learning models in functional neuroimaging, and enables hypothesis-driven analysis of cellular mechanisms across neuropsychiatric disorders.

Introduction

Deep neural networks (DNNs) are increasingly central to human brain research, offering powerful tools for scientific discovery and clinical applications (1, 2, 3, 4, 5). These models have demonstrated remarkable success in identifying patterns and making predictions from complex neuroimaging data (3, 4, 5). However, a fundamental challenge persists: DNN models typically function as "black boxes" offering no insight into the neurobiological and cellular mechanisms underlying their performance. This lack of interpretability poses a significant obstacle in neuroscience, where identifying meaningful biomarkers is essential for advancing our understanding of brain function and dysfunction. Here we develop and validate an explainable AI (xAI) framework for interpreting deep neural networks in functional neuroimaging, enabling hypothesis-driven testing of specific cellular mechanisms.

Excitation/inhibition (E/I) imbalance has emerged as a fundamental framework for understanding various neurological and psychiatric disorders, including autism and schizophrenia (6, 7, 8, 9). This dysregulation manifests through altered glutaminergic and GABAergic signalling and regional changes in neurotransmitter concentrations, as demonstrated by converging evidence from cellular studies and magnetic resonance spectroscopy (9, 10, 11). In autism specifically, disrupted E/I balance appears to be a key mechanism underlying its core symptoms of restricted social interests and circumscribed behaviors (12, 13), with evidence spanning multiple scales of investigation from cellular studies showing altered GABAergic signalling in animal models, to human neuroimaging revealing regional GABA concentration changes (14, 15). These E/I alterations are hypothesized to specifically affect nodes of the default mode network (16), which plays a central role in self-referential processing, social cognition, and autobiographical memory

retrieval - cognitive functions notably impaired in autism, schizophrenia and many other

psychiatric and neurological disorders (16, 17, 18).

However, identifying the specific brain regions or networks where E/I imbalance occurs

remains a critical challenge due to methodological limitations. Current methods to measure

imbalance, such as magnetic resonance spectroscopy, require targeting specific hypothesized

areas a priori. This limitation has crucial consequences in clinical settings, where brain

regional targets must be identified for interventions altering E/I imbalance through localized

brain stimulation. Therefore, providing data-driven, testable hypotheses for E/I imbalance

localization could play a central role in uncovering targets for experimental and therapeutic

protocols.

Functional brain imaging offers a promising approach to probe circuit function and

dysfunction associated with E/I imbalance. Conventional functional MRI analyses rely

heavily on pre-engineered features, focusing primarily on static inter-regional connectivity

measures or predefined activation patterns (19, 20, 21). Recent attempts to incorporate neural

networks, including graph-based approaches (22, 23) and surface-based convolutional

networks (20), still largely reduce temporal dynamics to static representations. However, E/I

imbalance mechanisms manifest through dynamic alterations in neural signalling that may

only be detectable when preserving the full spatiotemporal complexity of brain activity

patterns (24, 25, 26).

Deep neural networks that analyze dynamic fMRI time series data without feature

engineering offer a promising solution to this challenge. Spatiotemporal DNNs (stDNNs), in

particular, address limitations of traditional approaches by learning discriminative features

directly from raw temporal signals, modelling a hierarchy of temporal scales preserving the

full complexity of brain activity patterns (3, 4, 27). These models have shown promise in

detecting aberrant circuit dynamics in psychiatric disorders where E/I imbalance is

prominent, including autism and psychosis (3, 4). Previous applications have demonstrated

that dynamic circuit alterations detected by stDNNs can predict symptom-defined subtypes in

autism (4) and genetically-defined subtypes in psychosis (3), with findings showing

robustness across multisite datasets (3, 4). However, the critical question remains: can we

validate that these models are indeed able to detect E/I imbalance mechanisms?

This validation challenge is particularly acute because xAI methods themselves must be

rigorously tested with known ground truth before we can confidently use them to guide

mechanistic understanding or treatment strategies. Without such validation, it would not be

possible to distinguish between genuine biological insights and algorithmic artifacts. This

interpretability gap is particularly problematic for clinical applications, where understanding

the biological basis of model decisions is essential for developing targeted interventions (25,

28). This reflects a fundamental tension in clinical machine learning between achieving high

predictive accuracy and gaining mechanistic insights that can inform therapeutic development

(28, 29).

To address these challenges, we developed and validated an xAI framework using two

complementary mechanistic simulation approaches to test whether attribution methods can

reliably recover brain regions affected by E/I imbalance. A key goal was to develop recurrent

neural network (RNN) and The Virtual Brain (TVB) models that capture essential brain

dynamics and E/I balance alterations, enabling systematic testing of whether stDNN-based

xAI methods can accurately identify affected regions under controlled conditions with known

ground truth – something impossible with real neuroimaging data alone. First, we employ

RNNs to generate synthetic fMRI data with precisely controlled parameters. Second, we

utilized TVB simulator (30) to create biophysically informed simulations incorporating

realistic brain connectivity from both human (31) and mouse connectomes (32). This dual

approach enables validation across different scales and species while maintaining known

ground truth.

In both approaches, we systematically investigated how various neurophysiological factors –

including noise levels, prevalence of brain regions affected by E/I imbalance, and severity of

E/I alterations – impact feature attribution in neuroimaging analyses. By manipulating these

parameters, we could simulate different scenarios of E/I imbalance, which is crucial for

understanding neurological disorders such as autism (13) and schizophrenia (33).

Our study addresses four key research questions that establish validation protocols for

neuroimaging xAI. First, can feature attribution methods reliably localize regions with altered

E/I balance when clinical differences arise from these cellular-level mechanisms? We test this

using carefully constructed RNN simulations with known ground truth, systematically

varying the prevalence of affected regions (1 to 10% of nodes), severity of E/I alterations

($\delta = 0.1$-0.5), and signal-to-noise ratios (-10dB to 10dB). Second, how robust is this

localization across different biological scales and anatomical complexity at the whole-brain

level? To address this, we extend validation using biophysically realistic TVB simulations

incorporating empirically-derived connectomes from both human (68 regions) and mouse

(426 regions) brains. Third, how does attribution accuracy depend on method choice and

parameter selection? We systematically compare multiple xAI approaches to gradient-based

feature attribution— Integrated Gradients (34), DeepLIFT (35), Input X Gradient (36),

Guided Backprop (37)— that are computationally tractable and inexpensive but whose

relative performance remains actively contested even in foundational domains like visual

image categorization (36, 38). We also examine how baseline choice affects attribution in

dynamic brain imaging, a crucial methodological question unique to time series data. Fourth,

what is the interpretability of attribution maps when applied to real clinical data? We apply

our validated methods to the multisite ABIDE dataset (N = 834)(4), testing whether regions

identified by xAI methods align with known neurobiological theories of E/I imbalance in

autism, enabling us to assess whether our biophysically grounded validation framework

provides sufficient confidence for interpreting deep learning models in clinical neuroimaging.

Our comprehensive framework provides methodological guidelines for xAI applications in

clinical neuroimaging and generates testable hypotheses about target sites and mechanisms to

guide future experimental and therapeutic efforts.

Results

RNN simulations and validation

We first evaluate xAI methods using recurrent neural network (RNN) simulations, which

allowed systematic control over key parameters while maintaining temporal dynamics

characteristic of fMRI data. These simulations generated synthetic fMRI time series from two

groups: one with typical network dynamics and another with hyperexcitable nodes,

mimicking localized E/I imbalance.

Model design and parameter space

The RNN model generated time series data for 100 regions, with adjustable parameters

controlling: signal-to-noise ratio (SNR: -10dB to 10dB), prevalence of affected regions (1 to

10% of nodes), and severity of hyperexcitability (δ: 0.1 to 0.5). Network connectivity was

defined using small-world topology to reflect brain-like architecture, while regional dynamics

were modelled using nonlinear activation functions convolved with a canonical

hemodynamic response (see Methods).

Classifier performance

Our stDNN classifier demonstrated robust performance across simulation conditions

(**Fig S1**). Even at high noise levels, with 5% prevalence, and small hyperexcitability

($\delta = 0.1$), the classifier achieved perfect discrimination (accuracy = 100%) (-10dB SNR)

(**Fig S1A**). Performance remained robust down to 1% prevalence, though convergence

required more training epochs (**Fig S1B**). The classifier successfully detected group

differences across hyperexcitability levels above $\delta = 0.1$, with faster convergence at higher

severities (**Fig S1C**).

xAI analysis

IG successfully identified hyperexcited nodes across all simulation conditions (**Fig 2**).To

establish a consistent threshold across xAI methods and simulation parameters, we

normalized all attributions by dividing it by its maximum value across ROIs. With a

normalized attribution threshold of 0.5, IG achieved perfect detection (F1 score = 100%) of

affected network nodes even under challenging conditions: high noise (-10dB SNR, **Fig 2A-**

C), low prevalence (1%, **Fig 2D-F**), and subtle hyperexcitability ($\delta = 0.1$, **Fig 2G-I**).

Remarkably, IG's performance remained stable across these conditions, suggesting robust

feature detection even with minimal signal.

These RNN simulation results established that IG can reliably identify affected brain regions

across a wide range of simulation conditions.

Human TVB Simulations

We next evaluated xAI methods using biophysically informed simulations with The Virtual

Brain (TVB), incorporating realistic human brain connectivity and dynamics. These

simulations provided a more stringent test of our approach while maintaining known ground

truth alterations in specific brain regions.

Biophysical model implementation

Using structural connectivity from human diffusion imaging (68-region Desikan-Killiany

atlas(39)), we simulated whole-brain dynamics with explicit modelling of excitatory and

inhibitory populations in each region. We created two groups by introducing E/I imbalance in

one group through altered inhibitory synaptic conductance (ΔQi = 0.1-0.5nS): first targeting

only bilateral posterior cingulate cortex (PCC), then progressively including precuneus

(PCun) and angular gyrus (AnG) to test effects of anatomical extent on classification and IG

performance.

Classifier performance

The stDNN classifier achieved robust discrimination between groups across simulation

conditions (**Fig S2**). For large imbalance in PCC-only, classification accuracy remained

above 90% for SNR levels down to -10dB (**Fig S2A**). Convergence time decreased with

increasing anatomical extent of E/I imbalance, with fastest classification when alterations

included PCC, PCun, and AnG (**Fig S2B**). The classifier showed sensitivity to E/I imbalance

severity, with reliable discrimination (accuracy > 90%) for $\Delta Qi \geq 0.3nS$ (**Fig S2C**).

xAI results

IG successfully identified regions with altered E/I balance across conditions (**Fig 3**). With

PCC-only alterations, IG showed high attributions specifically localized in the bilateral PCC

across SNR levels (**Fig 3A-C**). When additional regions were affected, IG accurately detected

the expanded anatomical extent of alterations (**Fig 3D-F**), maintaining precise localization

even as the number of affected regions increased. Attribution patterns remained stable across

E/I imbalance severities (**Fig 3G-I**), though stronger alterations ($\Delta Qi = 0.5nS$) produced more

pronounced attribution scores. Importantly, regions without simulated E/I alterations

consistently showed attribution scores below our normalized attribution threshold of 0.5,

demonstrating high specificity in feature detection.

These TVB simulation results demonstrate that xAI methods can reliably identify

biologically realistic E/I alterations in the context of whole-brain dynamics and authentic

human brain connectivity. The successful detection of affected regions across varying

anatomical distributions and E/I imbalance severities supports the method's potential utility

for identifying circuit alterations in clinical populations.

Mouse TVB simulations

To test the generalizability of our approach across species and scales, we conducted parallel

analyses using the mouse connectome. Unlike human diffusion imaging-based connectivity,

the Allen Brain Atlas provides invasively validated, directional connectivity data across 426

regions, offering higher spatial resolution and more direct measurement of anatomical

connections.

Model implementation

We adapted our TVB simulations to the mouse brain, maintaining the same biophysical

parameters for excitatory and inhibitory populations while incorporating mouse-specific

structural connectivity. E/I imbalance was introduced in the retrosplenial cortex (RSC), the

key DMN hub in rodents analogous to the human PCC. We systematically varied inhibitory

conductance (ΔQi: 0.1 to 0.5nS) to parallel our human simulations.

Classifier performance

The stDNN classifier showed robust but distinct performance patterns with mouse data

(**Fig S3**). Classification accuracy remained above 90% for moderate noise levels (0dB SNR)

with RSC-only alterations (**Fig S3A**). Performance improved when E/I imbalance extended to

additional regions (RSC+Cg, RSC+Cg+PrL; **Fig S3B**), though requiring more training

epochs than for human simulations. The classifier maintained sensitivity to E/I imbalance

severity, achieving reliable discrimination for $\Delta Qi \geq 0.3nS$ (**Fig S3C**).

xAI results

IG successfully identified affected regions of the mouse brain, though with some species-

specific patterns (**Fig 4**). With RSC-only alterations, IG showed enhanced attributions

precisely localized in all RSC subregions only for SNR levels down to 0dB (**Fig 4A-C**).

When E/I imbalance extended to additional regions, IG accurately detected the broader

anatomical distribution (**Fig 4D-F**), maintaining precision and recall above 80%. Hierarchical

attribution strengths, varying across subregions, were observed here due to more structurally

distinct regions affected here than in human TVB simulations. This pattern held across E/I

imbalance severities (**Fig 4G-I**), suggesting robust detection of regional hierarchies in the

mouse brain.

The successful cross-species validation, particularly the detection of analogous regions

(RSC/PCC) despite different scales and connectivity patterns, demonstrates the robustness of

our xAI approach. The higher spatial resolution of the mouse connectome suggests the

potential relevance of the stDNN and IG approach to high-dimensional data and to reveal

additional insights about subregional specificity while maintaining reliable detection of E/I

alterations.

ABIDE fMRI dataset analysis

After validating our approach using simulated data, we applied these methods to the Autism

Brain Imaging Data Exchange (ABIDE) dataset to identify brain regions that reliably

distinguish individuals with autism from neurotypical controls. This multi-site dataset

(N=834; 419 autism, 415 controls) provided a robust test of our methods in real-world

clinical data.

Classifier performance

As reported previously (4), our stDNN achieved significant classification accuracy across the

ABIDE dataset (78.2 ± 2.84%, using 5-fold cross validation), reliably distinguishing autism

from control participants. Performance remained stable across different site splits and cross-

validation folds, demonstrating robustness to data collection variability. Notably,

classification accuracy was maintained even with highly heterogeneous symptomatology

across autistic individuals and multi-site imaging data (4).

xAI analysis

xAI analysis revealed a consistent set of discriminative brain regions (**Fig 5**). IG identified

the bilateral posterior cingulate cortex (PCC) and right precuneus as the regions most

strongly contributing to classification, with attribution scores above our normalized

attribution threshold of 0.5. This finding was remarkably stable across multiple analyses as

described below.

The convergence between our empirical findings and simulation results provides

computational support for the role of E/I imbalance in autism, particularly within the DMN.

The identification of these regions is consistent with previous studies showing altered DMN

function in autism and adds mechanistic insight by suggesting these alterations may reflect

underlying E/I imbalance.

Comparison of xAI methods

To establish the reliability of our approach, we systematically compared multiple xAI

methods across both simulated and empirical data. We evaluated four methods: Integrated

Gradients (IG), DeepLIFT, Input X Gradient (IXG), and Guided Backpropagation (GB), each

representing different theoretical approaches to explaining model decisions.

RNN simulations

In our controlled RNN simulations, IG, DeepLIFT, and IXG showed remarkably consistent

performance (**Fig 6A**). These methods demonstrated high correlations in their attribution

patterns ($r > 0.99$) and achieved similar accuracy as measured by F1score in identifying

hyperexcitable nodes (**Fig S4A**). In contrast, GB showed weaker correlations with other

methods ($r \leq 0.42$) and impaired F1 scores (**Fig S4A**) due to reduced precision (**Fig S4B**) but

not recall (**Fig S4C**) in detecting affected regions.

Human TVB simulations

In human TVB data, IG, DeepLIFT, and IXG maintained strong correlations ($r > 0.97$,

Fig 6B) in identified regions with affected E/I balance. Weaker correlations were seen

between attribution profiles of GB and other methods ($r \leq 0.39$). Poorer F1 scores (**Fig S4D**)

at identifying affected nodes due to lower precision (**Fig S4E**) but not recall (**Fig S4F**) was

also observed for GB compared to other methods.

Mouse TVB simulations

Similar patterns emerged in mouse TVB simulations, where these methods showed high

agreement ($r > 0.99$, **Fig 6C**) despite the increased anatomical complexity of the mouse

connectome. In contrast, GB showed weaker correlations with other methods ($r = 0.77$) and

reduced F1 score due to low precision in detecting affected regions (**Fig S4G-I**).

ABIDE fMRI data

In the ABIDE dataset, IG, DeepLIFT, and IXG identified highly consistent sets of

discriminative regions ($r > 0.98$, **Fig 6D**). All three methods highlighted the bilateral PCC

and right precuneus as key regions distinguishing autism from controls. GB showed markedly

different attribution patterns (r < 0.06), suggesting potential limitations in its clinical

applications.

These results demonstrate that IG, DeepLIFT, and IXG provide consistent and reliable

feature attribution across different experimental conditions and scales of brain organization,

while GB shows divergent patterns and reduced accuracy, suggesting it may be less suitable

for neuroimaging applications.

Impact of baseline choice in Integrated Gradients

The choice of baseline represents a critical methodological consideration when applying

Integrated Gradients to neuroimaging data. Unlike static image analysis where a black (zero)

image serves as a natural baseline, the optimal reference point for dynamic brain data is not

immediately obvious. We therefore systematically evaluated multiple baseline choices

including the zero, mean, median across the ROIs of the input timeseries, and random across

our simulation frameworks and empirical data.

RNN simulations

In RNN simulations, zero, mean, median, and random baselines showed remarkably high

correlations in their attribution patterns (r ≥ 0.99, **Fig 7A**) and achieved equivalently good

performance in identifying affected regions (**Fig S5A-C**).

Human TVB simulations

This consistency extended to the more complex TVB simulations, where these three baselines

maintained strong correlations despite added biological complexity (r ≥ 0.94 for human,

Fig 7B). The preservation of attribution patterns across baseline choices suggests robust

feature detection rather than methodological artifacts. In contrast, random baselines showed

weaker correlations with other methods ($r < 0.49$) and reduced F1 scores in detecting affected

regions (**Fig S5D**) due to both impaired precision (**Fig S5E**) and recall (**Fig S5F**) compared

to other baseline choices.

Mouse TVB simulations

On higher-dimensional mouse TVB simulations yielded high correlations between mean and

median baselines ($r = 0.93$) as well as between zero and random normal baselines ($r = 0.82$).

Weaker correlations ($r \leq 0.64$) were found between other baseline pairs (**Fig 7C**). As with

RNN simulations, no significant differences across baseline choices were found as to their

accuracy scores at identifying affected regions (**Fig S5G-I**).

ABIDE fMRI data

Baseline robustness was further confirmed in empirical data, with zero, mean, and median

baselines showing highly correlated attribution patterns in the ABIDE dataset ($r > 0.99$,

Fig 7D). Importantly, random baselines consistently showed poorer performance across all

conditions, with significantly lower correlations with other baselines (typically $r \leq 0.96$).

These results demonstrate that zero, mean, and median baselines all provide stable and

consistent feature attribution in neuroimaging applications, but random baselines should be

avoided due to their poor reproducibility.

Discussion

DNN models have emerged as powerful tools for analyzing functional MRI data, offering

unprecedented accuracy in predicting cognitive states and clinical outcomes (3, 4). However,

their inherent complexity and "black box" nature have limited our ability to determine which

neuroanatomical features and neurophysiological mechanisms drive their predictions. This

opacity poses a significant challenge in neuroscience, where interpretability is crucial for

advancing our understanding of brain function and dysfunction and for developing targeted

therapeutic interventions. The core challenge is that without ground-truth validation, we

cannot assess whether xAI attribution methods identify genuine neurobiological mechanisms

or algorithmic artifacts. This validation gap has severely limited our understanding of xAI

attributions, particularly for clinical applications where identifying the biological basis of

model decisions is essential for guiding mechanistic hypotheses and treatment development.

Here we addressed this fundamental challenge by developing and validating an xAI

framework to test whether attribution methods can reliably recover brain regions affected by

excitation/inhibition (E/I) imbalance, a well-established cellular mechanism implicated across

multiple neuropsychiatric disorders (6, 7, 8, 9). A key innovation was creating biophysically

realistic simulations using RNN and TVB models that capture essential brain dynamics and

E/I balance alterations with unprecedented biological realism, enabling systematic hypothesis

testing under controlled conditions with known ground truth – something impossible with

real neuroimaging data alone. Our findings demonstrate that xAI methods can reliably

identify brain regions affected by E/I imbalance across a wide range of conditions

establishing the first validated framework for biological interpretation of deep learning

models in functional neuroimaging.

Feature attribution methods reliably localize regions with altered E/I balance

RNN simulations provided the first systematic answer to whether feature attribution methods can reliably localize regions with pathophysiology by generating synthetic fMRI data with precisely controlled E/I alterations. We demonstrated that Integrated Gradients can reliably identify affected brain regions across challenging conditions that mirror real-world neuroimaging scenarios (**Fig 2**). The robustness across extreme conditions is particularly important for clinical applications, where signal quality varies significantly between individuals and disease-related alterations may be subtle or spatially sparse. Our systematic validation across prevalence rates (1 to 10% of nodes), severity levels (δ: 0.1 to 0.5), and noise conditions (SNR: -10dB to 10dB) provides performance benchmarks that enable researchers to assess the reliability of xAI results in their own datasets. The robustness across extreme conditions is particularly important for clinical applications, where signal quality varies significantly between individuals and disease-related alterations may be subtle or spatially sparse.

Our results demonstrate that xAI methods can reliably bridge the gap between cellular-level mechanisms and network-level patterns, making xAI findings more interpretable from a cellular mechanism standpoint in that xAI can localize E/I imbalance.

Robustness of E/I imbalance localization across different biological scales and anatomical complexity

A critical question for any neuroimaging method is whether findings generalize across different brain organizations, spatial scales, and species. Validation must account for substantial differences in connectivity patterns, anatomical resolution, and measurement approaches between human and animal models. Our TVB simulations incorporating

empirically-derived human and mouse connectomes demonstrated remarkable robustness

across biological scales (**Figs 3 and 4**). Despite substantial differences in brain parcellation,

connectivity patterns, and measurement approaches (DTI-based vs viral tracing), xAI

methods successfully identified altered E/I balance in analogous brain regions—the posterior

cingulate cortex in humans and retrosplenial cortex in mice.

The human TVB simulations, incorporating structural connectivity from the Human

Connectome Project (31), allowed us to model E/I imbalance within realistic brain network

architectures. IG successfully identified regions with altered inhibitory conductance across

different anatomical extents and severity levels, maintaining precise localization even as the

number of affected regions increased. The mouse simulations provided validation at higher

spatial resolution using the Allen Brain Atlas's experimentally validated connectivity data

(32), demonstrating that our approach scales effectively to finer anatomical detail.

This cross-species validation represents a significant advance by providing computational

support for translational research using mouse models to study E/I imbalance mechanisms

relevant to human disorders. The consistency between species despite different scales and

connectivity patterns demonstrates the robustness of our xAI approach and suggests broad

applicability across neuroimaging applications. This finding is particularly important as it

establishes that xAI methods detect biological mechanisms rather than artifacts of specific

connectome properties, enabling translation between preclinical models and human studies.

Robustness of xAI methods choice and parameter selection on attribution accuracy

A critical unresolved question in explainable AI has been whether different attribution

methods yield consistent and biologically meaningful results. Even in foundational domains

like visual image categorization, the relative performance of these methods remains actively

contested despite their widespread use (36, 38). For neuroimaging, this uncertainty is

particularly problematic because inconsistent results across methods would undermine

confidence in any biological interpretation. Additionally, methodological parameters like

baseline selection pose unique challenges for dynamic functional brain imaging that have not

been systematically addressed.

Our systematic comparison of four prominent xAI methods – Integrated Gradients (34),

DeepLIFT (DL) (35), Input X Gradient (IXG) (36), Guided Backprop (GB) (37) – revealed

striking patterns of consistency and divergence (**Fig 5**). IG, DeepLIFT, and IXG showed

remarkably high correlations (r > 0.95) across all simulation conditions, from simple RNN

models to complex TVB simulations with both human and mouse connectomes. This

consistency suggests these methods capture fundamental features of network dynamics rather

than methodological artifacts. In contrast, Guided Backpropagation showed notably weaker

correlations with other methods (r < 0.4 in human data, r < 0.77 in mouse data) and reduced

accuracy in identifying ground truth affected regions.

Regarding parameter selection, we addressed the critical question of baseline choice for

dynamic brain imaging (**Fig 6**). Unlike static image analysis where zero baselines are natural,

temporally complex fMRI signals require careful consideration of what constitutes "absence

of signal." Our systematic evaluation revealed that zero, mean, and median baselines yield

highly correlated attribution patterns (r > 0.94) across simulation conditions and real data,

providing methodological flexibility while maintaining interpretive validity. However,

random baselines showed poor reproducibility and should be avoided.

These results provide the first systematic guidelines for method and parameter selection in

neuroimaging xAI. The high consistency between IG, DeepLIFT, and IXG establishes that

reliable biological interpretation is achievable when appropriate methods are selected,

providing methodological clarity and validated protocols for future studies seeking to

uncover mechanistic alterations in autism and other neuropsychiatric disorders.

Clinical implications and future applications

Our validated xAI framework enables several important clinical applications and testable

hypotheses. Most directly, our findings E/I imbalance disruptions can be manifest as

perturbations in xAI attribution maps. This interpretation can be empirically tested by

targeting magnetic resonance spectroscopy measurements to quantify GABA concentrations

in regions identified by feature attribution in individual patients' neuroimaging data.

The demonstrated ability of xAI methods to reliably localize E/I imbalance across different

severity levels and anatomical distributions opens new possibilities for clinical stratification.

Our framework could potentially identify subgroups of individuals with autism based on the

regional patterns and severity of E/I disruptions revealed through attribution maps. This

approach is particularly relevant given the substantial heterogeneity in autism symptom

profiles and severity, which may require different treatment strategies depending on

underlying neurobiological subtypes.

However, several important limitations must be acknowledged before clinical

implementation. Our validation was performed on a single disorder and dataset, requiring

replication across independent cohorts and multiple neuropsychiatric conditions.

Additionally, the relationship between xAI attributions and actual GABA concentrations

measured through spectroscopy remains to be empirically established. Future studies should

directly correlate attribution scores with biochemical measurements and clinical outcomes to

validate the biological relevance of these computational findings.

Beyond autism, our validation framework can be extended to other disorders where E/I

imbalance is hypothesized, such as schizophrenia and epilepsy. The biophysical simulation

approach can also be adapted to test other cellular mechanisms including neuromodulation

and synaptic plasticity, enabling systematic validation of xAI methods across different

neurobiological hypotheses. This represents a general framework for hypothesis-driven

interpretation of deep learning models in clinical neuroscience.

Conclusion

We developed and validated a biophysically realistic simulation framework for testing

whether explainable AI methods can reliably identify cellular-level mechanisms underlying

neuropsychiatric disorders. Our systematic validation demonstrates that Integrated Gradients

and DeepLIFT methods can accurately localize regions affected by excitation/inhibition

imbalance, maintain robust performance across species and anatomical scales, and translate

meaningfully to clinical populations.The scientific significance of this work extends beyond

methodological validation. By establishing that explainable AI methods can reliably recover

known E/I alterations in controlled simulations, we provide the neuroimaging community

with essential tools for confident biological interpretation of deep learning models. This work

represents a foundation for interpretable and biologically informed applications of artificial

intelligence across neuroimaging research.

Materials and Methods

We evaluate the accuracy of the relevant inputs retrieved by xAI methods using two different

simulation models for synthesizing fMRI timeseries. The first simulation model is a simple

recurrent neural network model where the differences between two classes is simulated using

different functional connectivity between the brain regions as well as the local

excitatory/inhibitory (E/I) imbalances in some subset of brain regions. The primary purpose

of this simple model is to investigate the performance of the xAI methods as a function of

prevalence rates and SNRs and to find the optimal thresholds on the xAI scores to separate

the most relevant brain features from the irrelevant ones. The second model is a more realistic

biophysical model for autism which is implemented using The Virtual Brain stimulator. The

differences between the two classes is created by introducing differences in the local E/I

imbalances in some set of brain regions. These regions are chosen based on our recent

findings (4, 40) in the autism population. We simulate timeseries of nodes (or ROIs)

consisting of two classes which differ either with respect to the connectivity of ROIs in the

two classes or differ with respect to the local excitability of some chosen nodes. We then

train a classifier to predict the class label given the timeseries of the nodes. Finally, we

examine how each xAI method discovers important nodes or features in each class. We

hypothesize that these xAI methods would correctly identify the nodes that differ in

functional connectivity or and their local excitability between the classes.

Simulated fMRI data with a recurrent neural network model

We simulated the whole-brain fMRI timeseries by the time series of 100 nodes or regions of

interest (ROIs). To generate simulated fMRI data of two classes, we assumed that the

difference between the two classes lies in their local excitability on some chosen nodes. We

assumed that the network consists of 100 nodes defined using an adjacency matrices A of a

randomly sampled connected Watts–Strogatz small-world graphs (41), each of the 100 nodes

joined with its $k = 10$ nearest neighbors in a ring topology with a probability $p = 0.1$ of

rewiring. The local level of excitability for the two classes, W_0 and W_1, are defined as $W_k = $

$\Lambda_k A^T$, where $\Lambda_0 = \frac{1}{\rho} I_n$, $\Lambda_1 = \begin{pmatrix} \left(\frac{1+\delta}{\rho}\right) I_a & 0 \\ 0 & \frac{1}{\rho} I_{n-a} \end{pmatrix}$, I_a and I_{n-a} are respectively the $a \times a$

and the $(n - a) \times (n - a)$ the identity matrices, $n = 100$ is the total number of nodes, a is

the number of nodes affected by the difference in local excitability, ρ is the spectral radius of

the matrix A, and $\delta > 0$ measures the amplitude of the difference between the two classes.

We simulate several fMRI datasets with varying degrees of differences in their local

excitability. We quantify these differences using (1) a prevalence rate that describes the ratio

of the number of nodes that differ in local excitability to the overall number of nodes (i.e. the

ratio $\frac{a}{n}$), and (2) the severity of the hyper-excitability defined as the ratio of local difference in

excitability in affected nodes to the level of excitability (i.e. the ratio $\frac{\frac{1+\delta}{\rho} - \frac{1}{\rho}}{\frac{1}{\rho}} = \delta$). We

hypothesize that the difficulty of finding the discriminating nodes increases when the

prevalence rate or the hyper-excitability level drops.

We simulated fMRI data using a Recurrent Neural Network (RNN) convolved with a

canonical hemodynamic response function (HRF), which can be described with the following

dynamical system:

$$x_t = tanh(W x_{t-1} + \epsilon_t) \tag{1}$$

$$y = HRF * x + \omega \tag{2}$$

Where $W \in \{W_0, W_1\}$ is the recurrent weight matrix, y_t represents the BOLD fMRI signal of

multiple regions of interests (ROIs) at time t, x_t represents hidden neural variables giving

rise to the BOLD signal at time t, $\epsilon_t = N(0, \sigma_\epsilon)$ represents the intrinsic neural noise, $\omega_t =$

$N(0, \sigma_\omega)$ represents the measurement noise which is independent across regions, HRF is a

canonical hemodynamic response function with a sampling period of 1s (42).

We chose σ_ω to achieve some fixed level of Signal to Noise Ratio (SNR) as follows:

592
$$\sigma_\omega = \frac{\rho}{10^{0.1 * SNR}} \tag{3}$$

where σ_ω is the variance of the noise and ρ is the maximal variance of the signal across ROIs

that we empirically computed after simulating x_t dynamics. An example of this simulated

data for each of the two classes over time can be seen in **Fig 1**.

We investigated the performance of various xAI methods with the prevalence rates ranging

from 1 to 10% in steps of 1%, the hyper-excitability amplitude ranging from 0.0 to 0.5 in

steps of 0.1, and the SNR ranging from -10dB to 10dB in steps of 10dB. For each set of

parameters, 400 random samples were generated, with 80% used for training and 20% held

out for testing.

Simulated fMRI data with The Virtual Brain Simulator

We used a more realistic biophysical model to generate synthetic fMRI datasets for two

classes representing the neurotypical and autism populations to investigate the performance

of the xAI methods. Specifically, we simulated fMRI data using a model of whole-brain

dynamics where biophysical parameters can be varied at the regional level. The model was

implemented in The Virtual Brain (TVB) simulator (30, 43) in previous work (44). Briefly,

each simulated region comprises two neural populations, one of excitatory (E) and one of

inhibitory (I) neurons, whose mean firing rates across E and I populations are described by

the following differential equations:

611
$$\tau \frac{dx_i^E}{dt} = -x_i^E + F_E(Q_E(x_i^E + J_i), x_i^I) + v_i^E \tag{4}$$

612
$$\tau \frac{dx_i^I}{dt} = -x_i^I + F_I\big(Q_E(x_i^E + RJ_i), x_i^i\big) + v_i^I \tag{5}$$

613
$$J_i(t) = \sum_{j \neq i} W_{ij} x_j^E \left(t - \frac{D_{ij}}{s}\right) \tag{6}$$

614
$$y_i = HRF * (rx_i^E + (1 - r)x_i^I) + \omega_i \tag{7}$$

Where x_i^E and x_i^I are respectively the firing rates of excitatory and inhibitory populations in

region i, J_i represents the input into region i of other regions long-range excitatory

connections between regions, $R = 1.4$ is the ratio between the conductance of long-range

excitatory-to-excitatory and excitatory-to-inhibitory synapses, W and D characterize

respectively the number and the length of white-matter fibres between regions between

regions both empirically derived from synaptic tracing in the mouse brain, $s = 3m/s$

represents the speed of transmission between regions, $Q_E = 1\,nS$ and Q_I represents

respectively the synaptic strength of excitation and inhibition, v_E and v_I are standard normal

integration noise, the transfer functions F_E and F_I which relates each population's output rate

to the input it receives from itself and other populations were fitted to reproduce the input-

output relations of integrate-and-fire neurons in spiking simulations (45), y represents the

BOLD fMRI signal of multiple regions of interests (ROIs), $\omega_i = N(0, \sigma_\omega)$ represents the

measurement noise in region i, HRF is a canonical hemodynamic response function. As

previously we chose σ_ω to achieve some fixed level of Signal to Noise Ratio (SNR) as

follows:

630
$$\sigma_\omega = \frac{\rho}{10^{0.1*SNR}} \tag{8}$$

where σ_ω is the variance of the noise and ρ is the maximal variance of the signal across ROIs

that we empirically compute after simulation.

For the human brain, tract weights and lengths were obtained with diffusion tensor imaging

of a healthy adult's brain (44) (data publicly available at https://zenodo.org/records/4263723).

The connectome was partitioned according to the commonly used 68-ROI Desikan-Killiany-

Tourville atlas (46). For the mouse brain, the structural connectome was taken from a tracer-

derived structural connectome of the mouse brain from the Allen Brain Institute (32) with a

426-ROI parcellation. Inter-regional synaptic delays were modelled based on the matrix of

tract lengths from the same source (32). Synaptic weights and delays were taken to be the

same from the left and right brain hemispheres.

The datasets for the autism class was generated by introducing E/I imbalance in a subset of

ROIs. Specifically, the biophysical parameter Q_I capturing inhibitory synaptic conductance

was modified specifically in regions of the default mode network where excitation/inhibition

imbalance has been observed in humans and rodent models of autism and hypothesized to

underlie symptoms. For the human brain, Q_I was altered bilaterally progressively in the

posterior cingulate cortex, in the precuneus (PCun) and in the angular gyrus (AnG). In the

mouse brain, Q_I was locally modified progressively in the bilateral agranular, ventral, and

dorsal subdivisions of the retrosplenial cortex (RSC), in the cingulate cortex (Cg) and in the

prelimbic cortex (PrL).

We simulated time series of 100 s for all ROIs for each connectome. This procedure was

repeated with 100 different random seeds that affected initial conditions and random noise

realizations, 80 seeds were used for training and 20 held out for testing. To simulate fMRI

data, the firing rate time series were then convolved with a canonical hemodynamic response

function with TR of 1s. To test the robustness of the xAI methods across different noise

levels, in both human and mouse models, measurement noise was added to the ROI

660 timeseries generated using the above simulation models. To quantify the impacts of severity

661 of E/I imbalance, inhibitory conductance in affected regions was varied between 4.5 nS and

662 4.9 nS to obtain severity varying from $\Delta Q_i = 0.1nS$ to $\Delta Q_i = 0.5nS$ between affected and

663 unaffected regions. To investigate the effects of prevalence, imbalance was implemented in

664 different subsets of DMN regions of varying size – in the human model, we compared

665 imbalance bilaterally in the PCC only, in the PCC and PCun only, and in the PCC, PCun and

666 AnG, while in the mouse model, we compared imbalance bilaterally in the RSC only, the

667 RSC and Cg only, and the RSC, Cg, and PrL.

669 ***ABIDE dataset***

670 We leveraged neuroimaging and phenotypic data from the Autism Brain Imaging Data

671 Exchange (ABIDE; http://fcon_1000.projects.nitrc.org/indi/abide/) (1, 2). The dataset

672 consists 419 subjects with autism and 415 typically developing subjects from 14 sites. We

673 used the same preprocessed resting-state fMRI dataset as in previous work (4), excluding the

674 same subject identified as having missing or poor-quality imaging data. The Brainnetome

675 Atlas was applied, comprising 246 ROIs included in the analysis.

677 ***stDNN classifier model for the simulated fMRI datasets and for the ABIDE dataset***

678 We used a two-layer 1D convolutional neural network model for classifying the two groups

679 both simulated fMRI datasets and in the ABIDE dataset, integrating spatial ROI information

680 at each convolutional layer through summation across channels. For simulated fMRI datasets,

681 as shown in **Fig 1C**, our stDNN processes simulated fMRI time series with a 1D convolution,

682 followed by a ReLU nonlinearity and average pooling. It then transforms the features with a

683 second 1D convolution and ReLU, followed by adaptive average pooling that reduces each

684 feature to a single time step. The output is flattened and passed through a linear layer for

classification. For ABIDE dataset, following original paper (4), our stDNN processes fMRI

time series with two successive 1D convolution–ReLU–max-pooling blocks, averages

features across time, and applies dropout. The resulting representation is concatenated with

an auxiliary covariate encoding, then passed through a fully connected layer and sigmoid

activation for binary classification. We used 80% of the data to train the model, the remaining

20% for testing the model. The code for the model implemented in PyTorch is available at

https://github.com/scsnl/Strock_Nghiem_bioRxiv_2025.

XAI methods

We obtained the discriminating features from each simulated subject by using a xAI method

which determines how much each nodes or ROI contributes to the decision made by our

classifier for this subject. The simulated fMRI data being temporal, each of these xAI

methods provides us with a value for each ROI and for each time step of simulation. We

obtain the influence of an ROI by taking the absolute value of the median of the xAI across

time step. To evaluate the performance of the attribution method we normalize the attribution

by dividing it by the maximum threshold across ROI, and compare ROI with attribution

above 0.5 with ROI affected E/I imbalance using standard recall, precision and F1 metrics.

We first evaluated the performance of Integrated Gradient (IG) (34) on the simulated

datasets. We examined how IG performance may depend on the choice of baseline which is

required for evaluating the path integral for the feature attribution. It is not clear what an

optimal baseline is for the neuroimaging applications. Therefore, we investigated several

baselines including zero, random, mean, and median of all fMRI time series across the

subjects. We finally compared the performance of other gradient based xAI methods such as

DeepLIFT (DL) (35), Input X Gradient (IXG) (36), Guided Backprop (GB) (37) implemented

in the Captum python library.

Integrated Gradients

713   IG directly uses the gradient of the classifier with respect to the input ROIs to define the xAI.

714   This method can be used for any already trained DNN model with a single backpropagation

715   pass. It integrates this gradient over a path in the fMRI input space going to the subject's

716   simulated fMRI data from a baseline of the same dimension. Formally, IG attribution method

717   can be written as:

718
$$\phi(x) = (x - x') \times \int_{\alpha=0}^{1} \frac{\partial F(x' + \alpha(x - x'))}{\partial x_i} \, d\alpha \qquad (9)$$

where x is the original input, x' is the chosen baseline from which we start our integration, F

represents the function associating fMRI input to its clinical category, and ϕ is the function

giving the xAI score for a given node x. IG overcomes the problems associated with the

simple gradient based approaches such as the noisy gradients and gradient saturation.

As IG requires a baseline, we compared the xAI obtained for various baselines: (1) the

median of the signal across nodes and samples, (2) the mean of the signal across nodes and

samples, (3) the default all-zero baseline, and (4) a random sample drawn from a normal

distribution centred about zero with the variance reflecting the variance of the input.

Deep learning important features (DeepLIFT)

As IG, DeepLIFT is xAI method that can applied to any pretrained DNN model with a single

backpropagation pass. This method computes the importance of each input in the prediction

of the output. This importance for each input is computed in terms of the difference between

the output for a reference or the baseline and the given input. As in the IG, the baseline needs

to be chosen based on the domain. It is developed to address the issues with the basic gradient

based xAI approaches. As in the case of IG, we evaluate the performance of DeepLIFT for all

the baselines as defined above.

Input X Gradient (IXB)

IXB is also a gradient based attribution method used for obtaining xAIs. This is a very basic

method where gradient of the input with respect to the target is computed and then the

gradient is multiplied with the input to obtain the xAI scores. If both the gradient of the input

and also its magnitude is high then that feature's attribution is high. Both IG and DeepLIFT

methods are refinements over this approach.

Guided Backpropagation (GB)

This method is developed for explaining the decisions taken by a CNN model by computing

the gradient of the output with respect to the input. This method requires a modification of

the gradient in the ReLU layers where the gradient to these layers are made zero if the input

to the ReLU during the forward pass is negative. This approach is applicable to only the

convolutional networks.

Statistical analysis

Relationships between variables were quantified using Pearson's correlation coefficient (r),

and group differences were assessed using two-tailed t-tests. All statistical tests were

implemented in Python using the SciPy library, with significance evaluated at $P < 0.05$ (*),

$P < 0.01$ (**), or $P < 0.001$ (***).

Data Availability

This study used only existing, publicly available neuroimaging data from the Autism Brain

Imaging Data Exchange (ABIDE), hosted on the International Neuroimaging Data-Sharing

Initiative (INDI) platform (http://fcon_1000.projects.nitrc.org/indi/abide/). The data were

collected and shared by the ABIDE consortium; no new data were collected by the authors.

All analyses were purely computational. The code used for the computational model and

analyses is available at https://github.com/scsnl/Strock_Nghiem_bioRxiv_2025.

6. References

1.   Durstewitz D, Koppe G, Meyer-Lindenberg A. Deep neural networks in psychiatry.

Mol Psychiatry. 2019;24(11):1583-98.

2.   Kriegeskorte N, Golan T. Neural network models and deep learning. Curr Biol.

2019;29(7):R231-R6.

3.   Supekar K, de Los Angeles C, Ryali S, Kushan L, Schleifer C, Repetto G, et al.

Robust and replicable functional brain signatures of 22q11.2 deletion syndrome and

associated psychosis: a deep neural network-based multi-cohort study. Mol Psychiatry.

2024;29(10):2951-66.

4.   Supekar K, Ryali S, Yuan R, Kumar D, de Los Angeles C, Menon V. Robust,

Generalizable, and Interpretable Artificial Intelligence-Derived Brain Fingerprints of Autism

and Social Communication Symptom Severity. Biol Psychiatry. 2022;92(8):643-53.

5.   Vieira S, Pinaya WH, Mechelli A. Using deep learning to investigate the

neuroimaging correlates of psychiatric and neurological disorders: Methods and applications.

Neurosci Biobehav Rev. 2017;74(Pt A):58-75.

6.   Breakspear M. Dynamic models of large-scale brain activity. Nat Neurosci.

2017;20(3):340-52.

7.   Foss-Feig JH, Adkinson BD, Ji JL, Yang G, Srihari VH, McPartland JC, et al.

Searching for Cross-Diagnostic Convergence: Neural Mechanisms Governing Excitation and

Inhibition Balance in Schizophrenia and Autism Spectrum Disorders. Biol Psychiatry.

2017;81(10):848-61.

8.   Sohal VS, Rubenstein JLR. Excitation-inhibition balance as a framework for

investigating mechanisms in neuropsychiatric disorders. Mol Psychiatr. 2019;24(9):1248-57.

9.   Gao R, Penzes P. Common mechanisms of excitatory and inhibitory imbalance in

schizophrenia and autism spectrum disorders. Current molecular medicine. 2015;15(2):146-

67.

10.   Tang X, Jaenisch R, Sur M. The role of GABAergic signalling in neurodevelopmental

disorders. Nat Rev Neurosci. 2021;22(5):290-307.

11.   Lányi O, Koleszár B, Schulze Wenning A, Balogh D, Engh MA, Horváth AA, et al.

Excitation/inhibition imbalance in schizophrenia: a meta-analysis of inhibitory and excitatory

TMS-EMG paradigms. Schizophrenia. 2024;10(1):56.

12.   Nelson SB, Valakh V. Excitatory/Inhibitory Balance and Circuit Homeostasis in

Autism Spectrum Disorders. Neuron. 2015;87(4):684-98.

13.   Rubenstein JL, Merzenich MM. Model of autism: increased ratio of

excitation/inhibition in key neural systems. Genes Brain Behav. 2003;2(5):255-67.

14.   Hollestein V, Poelmans G, Forde NJ, Beckmann CF, Ecker C, Mann C, et al.

Excitatory/inhibitory imbalance in autism: the role of glutamate and GABA gene-sets in

symptoms and cortical brain structure. Transl Psychiatry. 2023;13(1):18.

15.   Horder J, Petrinovic MM, Mendez MA, Bruns A, Takumi T, Spooren W, et al.

Glutamate and GABA in autism spectrum disorder-a translational magnetic resonance

spectroscopy study in man and rodent models. Transl Psychiatry. 2018;8(1):106.

16.   Padmanabhan A, Lynch CJ, Schaer M, Menon V. The Default Mode Network in

Autism. Biol Psychiatry Cogn Neurosci Neuroimaging. 2017;2(6):476-86.

17.   Hu ML, Zong XF, Mann JJ, Zheng JJ, Liao YH, Li ZC, et al. A Review of the

Functional and Anatomical Default Mode Network in Schizophrenia. Neurosci Bull.

2017;33(1):73-84.

18.   Menon V. 20 years of the default mode network: A review and synthesis. Neuron.

2023;111(16):2469-87.

19.   Gadgil S, Zhao Q, Pfefferbaum A, Sullivan EV, Adeli E, Pohl KM. Spatio-Temporal

Graph Convolution for Resting-State fMRI Analysis. Med Image Comput Comput Assist

Interv. 2020;12267:528-38.

20.   Ngo GH, Khosla M, Jamison K, Kuceyeski A, Sabuncu MR. Predicting individual

task contrasts from resting-state functional connectivity using a surface-based convolutional

network. Neuroimage. 2022;248:118849.

21.   Zhang Y, Farrugia N, Bellec P. Deep learning models of cognitive processes

constrained by human brain connectomes. Med Image Anal. 2022;80:102507.

22.   Lei D, Qin K, Pinaya WHL, Young J, Van Amelsvoort T, Marcelis M, et al. Graph

Convolutional Networks Reveal Network-Level Functional Dysconnectivity in

Schizophrenia. Schizophr Bull. 2022;48(4):881-92.

23.   Wang L, Li K, Hu XP. Graph convolutional network for fMRI analysis based on

connectivity neighborhood. Netw Neurosci. 2021;5(1):83-95.

24.   Ryali S, Supekar K, Chen T, Kochalka J, Cai W, Nicholas J, et al. Temporal

Dynamics and Developmental Maturation of Salience, Default and Central-Executive

Network Interactions Revealed by Variational Bayes Hidden Markov Modeling. PLoS

Comput Biol. 2016;12(12):e1005138.

25.   Lurie DJ, Kessler D, Bassett DS, Betzel RF, Breakspear M, Kheilholz S, et al.

Questions and controversies in the study of time-varying functional connectivity in resting

fMRI. Network Neuroscience. 2020;4(1):30-69.

26.   Calhoun VD, Miller R, Pearlson G, Adali T. The Chronnectome: Time-Varying

Connectivity Networks as the Next Frontier in fMRI Data Discovery. Neuron.

2014;84(2):262-74.

27.   Ryali S, Zhang Y, de Los Angeles C, Supekar K, Menon V. Deep learning models

reveal replicable, generalizable, and behaviorally relevant sex differences in human

functional brain organization. Proc Natl Acad Sci U S A. 2024;121(9):e2310012121.

28.   Eitel F, Schulz MA, Seiler M, Walter H, Ritter K. Promises and pitfalls of deep neural

networks in neuroimaging-based psychiatric research. Exp Neurol. 2021;339.

29.   Ghassemi M, Oakden-Rayner L, Beam AL. The false hope of current approaches to

explainable artificial intelligence in health care. Lancet Digit Health. 2021;3(11):e745-e50.

30.   Sanz-Leon P, Knock SA, Woodman MM, Domide L, Mersmann J, McIntosh AR, et

al. The Virtual Brain: a simulator of primate brain network dynamics. Front Neuroinform.

2013;7:10.

31.   Glasser MF, Smith SM, Marcus DS, Andersson JLR, Auerbach EJ, Behrens TEJ, et

al. The Human Connectome Project's neuroimaging approach. Nature Neuroscience.

2016;19(9):1175-87.

32.   Oh SW, Harris JA, Ng L, Winslow B, Cain N, Mihalas S, et al. A mesoscale

connectome of the mouse brain. Nature. 2014;508(7495):207-+.

33.   Uhlhaas PJ, Singer W. Neuronal dynamics and neuropsychiatric disorders: toward a

translational paradigm for dysfunctional large-scale networks. Neuron. 2012;75(6):963-80.

34.   Sundararajan M, Taly A, Yan QQ. Axiomatic Attribution for Deep Networks. Pr

Mach Learn Res. 2017;70.

35. Shrikumar A, Greenside P, Kundaje A. Learning Important Features Through

Propagating Activation Differences. Pr Mach Learn Res. 2017;70.

36. Kindermans P-J, Hooker S, Adebayo J, Alber M, Schütt KT, Dähne S, et al. The (un)

reliability of saliency methods. Explainable AI: Interpreting, explaining and visualizing deep

learning. 2019:267-80.

37. Springenberg J, Dosovitskiy A, Brox T, Riedmiller M, editors. Striving for

Simplicity: The All Convolutional Net. ICLR (workshop track); 2015.

38. Gevaert A, Rousseau AJ, Becker T, Valkenborg D, De Bie T, Saeys Y. Evaluating

feature attribution methods in the image domain. Mach Learn. 2024;113(9):6019-64.

39. Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. An

automated labeling system for subdividing the human cerebral cortex on MRI scans into

gyral based regions of interest. Neuroimage. 2006;31(3):968-80.

40. Supekar K, Uddin LQ, Khouzam A, Phillips J, Gaillard WD, Kenworthy LE, et al.

Brain Hyperconnectivity in Children with Autism and its Links to Social Deficits. Cell Rep.

2013;5(3):738-47.

41. Watts DJ, Strogatz SH. Collective dynamics of 'small-world' networks. Nature.

1998;393(6684):440-2.

42. Ryali S, Supekar K, Chen T, Menon V. Multivariate dynamical systems models for

estimating causal interactions in fMRI. Neuroimage. 2011;54(2):807-23.

43. Sanz-Leon P, Knock SA, Spiegler A, Jirsa VK. Mathematical framework for large-

scale brain network modeling in The Virtual Brain. Neuroimage. 2015;111:385-430.

44. Goldman JS, Kusch L, Aquilue D, Yalcinkaya BH, Depannemaecker D, Ancourt K, et

al. A comprehensive neural simulation of slow-wave sleep and highly responsive

wakefulness dynamics. Front Comput Neurosc. 2023;16.

45.   Volo MD, Romagnoni A, Capone C, Destexhe A. Biologically Realistic Mean-Field

Models of Conductance-Based Networks of Spiking Neurons with Adaptation. Neural

Comput. 2019;31(4):653-80.

46.   Klein A, Tourville J. 101 labeled brain images and a consistent human cortical

labeling protocol. Front Neurosci-Switz. 2012;6.

Figures

889
Figure 1. Overview of explainable AI framework for E/I imbalance detection. Simulated
multivariate fMRI datasets consisting of two classes are generated using two complementary
simulation approaches: (1) recurrent neural networks (RNNs) for controlled parameter
exploration, and (2) biophysically realistic whole-brain simulations using The Virtual Brain
(TVB) simulator, which incorporates empirically-derived structural connectomes and models
neural dynamics through coupled excitatory and inhibitory populations in each brain region.
Both simulation approaches generate two classes that share identical structural connectivity
(adjacency matrix), but differ in excitation/inhibition balance—with some nodes (highlighted
in red) exhibiting altered E/I balance in one population through modified inhibitory
conductance parameters. A spatiotemporal deep neural network (stDNN) consisting of 1D
convolutional layers is trained to classify the two populations based on their dynamic fMRI
signals. Feature attribution methods are then applied to the trained model to identify which
brain regions drive classification decisions, testing the hypothesis that these methods can
reliably recover the known locations of E/I alterations.

Figure 2. Validation of Integrated Gradients for detecting E/I imbalance in RNN simulations. Performance of Integrated Gradients (IG) in identifying regions with altered excitation/inhibition balance in small-world RNN simulations across varying signal-to-noise ratios (SNR), prevalence rates, and E/I imbalance severity. Regions affected by E/I alterations (highlighted in red) were reliably identified with perfect accuracy (F1-score = 1.0) across all tested conditions. (**A-C**) IG attribution scores for SNRs of 10, 0 and -10 dB with 5% prevalence and severity δ = 0.3. (**D-F**) IG scores for prevalence rates of 1%, 5% and 10% with SNR of 0 dB and severity δ = 0.3. (**G-I**) IG scores for E/I imbalance severity δ = 0.5, δ = 0.3, and δ = 0.2 with SNR of 0 dB and 5% prevalence.

Figure 3. Cross-species validation using human connectome TVB simulations.
Performance of Integrated Gradients (IG) in detecting E/I imbalance in biophysically realistic
whole-brain simulations using human structural connectivity (68-region Desikan-Killiany
atlas). E/I imbalance was modeled through altered inhibitory conductance (ΔQi) in default
mode network regions. Affected regions (highlighted in red) were consistently identified
across all conditions. (**A-C**) IG scores for SNRs of 10, 0 and -10 dB with E/I imbalance
isolated to bilateral posterior cingulate cortex (PCC), ΔQi = 0.5nS. (**D-F**) IG scores for
progressive anatomical extent: PCC only, PCC+precuneus (PCun), and PCC+PCun+angular
gyrus (AnG) with ΔQi = 0.5nS and SNR of 0 dB. (**G-I**) IG scores for varying E/I imbalance
severity in PCC: ΔQi = 0.5nS, 0.4nS, and 0.3nS with SNR of 0 dB.

927
Figure 4. Cross-species validation using mouse connectome TVB simulations.
Performance of Integrated Gradients (IG) in detecting E/I imbalance in high-resolution
mouse brain simulations using Allen Brain Atlas connectivity (426 regions). E/I imbalance
was introduced in retrosplenial cortex (RSC), the rodent analog of human posterior cingulate
cortex. (**A-C**) IG scores for SNRs of 10, 0 and -10 dB with E/I imbalance in RSC,
$\Delta Q_i = 0.5nS$. (**D-F**) IG scores for progressive anatomical extent: RSC only, RSC+cingulate
cortex (Cg), and RSC+Cg+prelimbic cortex (PrL) with $\Delta Q_i = 0.5nS$ and SNR of 0 dB. (**G-I**)
IG scores for varying E/I imbalance severity in RSC: $\Delta Q_i = 0.5nS$, 0.4nS, and 0.3nS with
SNR of 0 dB.

Figure 5. Application to autism neuroimaging data reveals default mode network involvement. Feature attribution scores from the ABIDE autism dataset (N = 834) using validated xAI methods. (**A**) Integrated Gradients (IG), (**B**) DeepLIFT, and (**C**) Input X Gradient (IXG) show highly consistent attribution patterns, with bilateral posterior cingulate cortex (PCC) and right precuneus (PCun) exceeding the validated threshold of 0.5. (**D**) Guided Backpropagation (GB) shows diffuse, less discriminative attributions across many regions, with lower scores for PCC and precuneus.

947
Figure 6. Systematic comparison of feature attribution methods across validation
conditions. Correlation matrices showing agreement between different attribution methods
across simulation and empirical data. (**A-D**) Pairwise correlations between attribution
patterns from Integrated Gradients (IG), DeepLIFT (DL), Input X Gradient (IXG), and
Guided Backpropagation (GB) in: (**A**) RNN simulations, (**B**) human TVB simulations, (**C**)
mouse TVB simulations, and (**D**) ABIDE autism data. High correlations (r > 0.95) between
IG, DeepLIFT, and IXG demonstrate method consistency, while Guided Backpropagation
shows divergent patterns across all conditions.

957
Figure 7. Robustness of Integrated Gradients to baseline selection in dynamic brain
imaging. Correlation matrices evaluating the impact of baseline choice on IG attribution
patterns. (**A-D**) Pairwise correlations between attribution maps generated using median,
mean, zero, and random noise baselines in: (**A**) RNN simulations, (**B**) human TVB
simulations, (**C**) mouse TVB simulations, and (**D**) ABIDE data. High correlations between
median, mean, and zero baselines (r > 0.94) demonstrate robustness, while random baselines
show lower consistency, indicating they should be avoided in neuroimaging applications.

Supplementary Figures

Figure S1. Training and classification performance on simulated data generated using RNNs.
(**A-C**) Classification accuracy as a function of training epochs (**A**) for SNRs ranging from -10 dB to
10 dB with a fixed prevalence rate of 5% and hyper-excitability severity of $\delta = 0.1$, (**B**) for prevalence
rates ranging from 1 to 10% with fixed SNR of -10 dB hyper-excitability severity of $\delta=0.1$, and (**C**)
for hyper-excitability severity varying from $\delta = 0.0$ to $\delta = 0.5$ with fixed SNR of -10 dB and
prevalence rate of 5%. (**D**) Classification accuracy at epoch 149 for hyper-excitability severity of
$\delta = 0.0$ to $\delta > 0.0$. (**E-F**) Training epoch during which a 90% accuracy threshold is passed (**E**) and
convergence epoch (**F**) as a function of prevalence rate and hyper-excitability severity for an SNR of
10 dB. *** $P < 0.001$.

Figure S2. Training and classification performance on whole-brain TVB simulations with a human connectome. (**A-C**) Classification accuracy as a function of training epochs (**A**) for SNRs ranging from -10 dB to 10 dB when only the PCC is affected by E/I imbalance of severity ΔQ_i = 0.5nS (**B**) for only the PCC, the PCC and PCun, and the PCC, PCun, and AnG affected by E/I imbalance of severity ΔQ_i = 0.5nS with fixed SNR of -10 dB, and (**C**) for E/I imbalance of severity varying from ΔQ_i = 0nS to ΔQ_i = 0.5nS in the PCC with fixed SNR of -10. (**D**) Classification accuracy at epoch 149 for E/I imbalance severity of ΔQ_i = 0nS and ΔQ_i > 0nS. (**E-F**) Training epoch during which a 90% accuracy threshold is passed (**E**) and convergence epoch (**F**) as a function of regions affected and E/I imbalance severity for an SNR of 10 dB. *** P < 0.001.

Figure S3. Training and classification performance on whole-brain TVB simulations with a mouse connectome. (**A-C**) Classification accuracy as a function of training epochs (**A**) for SNRs ranging from -10 dB to 10 dB when only the RSC is affected by E/I imbalance of severity ΔQi = 0.5nS (**B**) for only the RSC, the RSC and Cg, and the RSC, Cg, and PrL affected by E/I imbalance of severity ΔQi = 0.5nS with fixed SNR of -10 dB, and (**C**) for E/I imbalance of severity varying from ΔQi = 0nS to ΔQi = 0.5nS in the PCC with fixed SNR of -10. (**D**) Classification accuracy at epoch 149 for E/I imbalance severity of ΔQi = 0nS and ΔQi > 0nS. (**E-F**) Training epoch during which a 90% accuracy threshold is passed (**E**) and convergence epoch (**F**) as a function of regions affected and E/I imbalance severity for an SNR of 10 dB. *** P<0.001.

1000
1001 **Figure S4. Performance of four different feature attribution methods in detecting affected**
1002 **ROIs.** Classifier accuracy quantified by F1score (**A**, **D**, **G**), precision (**B**, **E**, **H**), and recall (**C**, **F**, **I**) in
1003 RNN simulations (**A-C**) as well as TVB simulations with human (**D-F**) and mouse (**G-I**) connectomes
1004 across feature attribution methods IXG, DLIFT, IG, and GB. * P < 0.05, *** P < 0.001.

1005

1006
1007 **Figure S5. Performance of IG methods with four different baselines in detecting affected ROIs.**
1008 Classifier accuracy quantified by F1score (**A**, **D**, **G**), precision (**B**, **E**, **H**), and recall (**C**, **F**, **I**) in RNN
1009 simulations (**A**-**C**) as well as TVB simulations with human (**D**-**F**) and mouse (**G**-**I**) connectomes
1010 across median, mean, zero, and random normal baselines. *** P < 0.001

1011

Supplementary Tables

	Statistic *t*	Effect size *Cohen's d*	P	Degree of freedom
RNN	-73.26	-23.92	1.47e-36	31
TVB (Human)	-11.41	-4.81	3.95e-7	10
TVB (Mouse)	-9.85	-1.69	1.98e-13	51

Table S1. Comparison of models prediction with vs without E/I imbalance. Corresponding to the test performed in the Figure S1D, Figure S2D and Figure S3D.

	Metrics	Comparison	Statistic |t|	Effect size Cohen's |d|	P	Degree of freedom
RNN	F1	(IG|DLIFT|IXG)	NA	NA	NA	NA
		(IG|DLIFT|IXG) vs GB	27.68	3.20	1.33e-60	149
	Precision	(IG|DLIFT|IXG)	NA	NA	NA	NA
		(IG|DLIFT|IXG) vs GB	33.83	3.91	8.37e-72	149
	Recall	(IG|DLIFT|IXG|GB)	NA	NA	NA	NA
TVB (H)	F1	(IG|DLIFT|IXG)	< 0.10	< 0.02	> 0.92	[87, 88]
		(IG|DLIFT|IXG) vs GB	> 18.31	> 3.86	< 4.96e-26	[59, 61]
	Precision	(IG|DLIFT|IXG)	< 0.07	< 0.02	> 0.95	[87, 88]
		(IG|DLIFT|IXG) vs GB	> 19.13	> 5.03	< 2.68e-24	[48, 49]
	Recall	(IG|DLIFT|IXG)	< 0.35	< 0.08	> 0.73	[87, 88]
		(IG|DLIFT|IXG) vs GB	> 2.58	> 0.54	1.32e-02	44
TVB (M)	F1	(IG|DLIFT|IXG)	< 0.23	< 0.05	> 0.82	[87, 88]
		(IG|DLIFT|IXG) vs GB	> 6.66	> 1.40	< 2.73e-8	[46, 47]
	Precision	(IG|DLIFT|IXG)	< 0.25	< 0.06	> 0.81	[87, 88]
		(IG|DLIFT|IXG) vs GB	> 6.96	> 1.46	< 1.20e-8	[44, 45]
	Recall	(IG|DLIFT|IXG|GB)	0<1.10	< 0.24	> 0.27	[87, 88]

1017

1018 **Table S2. Comparison of attribution methods performance.** Corresponding to the test performed
1019 in the Figure S4. NA: Not applicable, comparison of constants (i.e. zero variance) that are equal.

1020

	Metrics	Comparison	Statistic t	Effect size *Cohen's d*	P	Degree of freedom
RNN	F1	(Z|MN|MD)	NA	NA	NA	NA
		(Z|MN|MD) vs RN	1.14	0.13	0.26	149
	Precision	(Z|MN|MD)	NA	NA	NA	NA
		(Z|MN|MD) vs RN	1.19	0.14	0.24	149
	Recall	(Z|MN|MD|RN)	NA	NA	NA	NA
TVB (H)	F1	(Z|MN|MD)	< 0.60	< 0.13	> 0.55	[87, 88]
		(Z|MN|MD) vs RN	> 5.61	> 1.18	< 2.29e-7	[86, 87]
	Precision	(Z|MN|MD)	< 0.96	< 0.03	> 0.34	[86, 88]
		(Z|MN|MD) vs RN	> 4.13	> 0.87	< 8.41e-5	[81, 86]
	Recall	(Z|MN|MD)	< 0.87	< 0.19	> 0.38	[87, 88]
		(Z|MN|MD) vs RN	> 5.78	> 1.21	< 1.36e-7	[80, 82]
TVB (M)	F1	(Z|MN|MD|RN)	< 1.28	< 0.27	> 0.20	[80, 88]
	Precision	(Z|MN|MD|RN)	< 0.57	< 0.12	> 0.57	[87, 88]
	Recall	(Z|MN|MD|RN)	< 1.94	< 0.41	> 0.05	[77, 88]

1021

1022 **Table S3. Comparison of IG baseline performance.** Corresponding to the test performed in the

1023 Figure S5. NA: Not applicable, comparison of constants (i.e. zero variance) that are equal.