Powerful flowers: Public perception of grassland aesthetics is strongly related to

management and biodiversity

Valentin H. Klaus [1,2,3,*], Nathan Fox [4,5], Franziska J. Richter [2,6], Davide Andreatta[7,8,a], Abdesslam Chai-

allah [9,a]

1 Agroscope, Forage Production and Grassland Systems, Zürich, Switzerland

2 ETH Zürich, Institute of Agricultural Sciences, Zürich, Switzerland

3 Ruhr University Bochum, Institute of Geography, Bochum, Germany

4 University of Michigan, School for Environment and Sustainability, Ann Arbor, MI, USA

5 University of Michigan, Michigan Institute for Data & AI in Society, Ann Arbor, MI, USA

6 Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Birmensdorf, Switzerland

7 University of Padova, Department of Agronomy, Food, Natural Resources, Animals and

Environment, Legnaro, Padova, Italy

8 Fondazione Edmund Mach, Research and Innovation Centre, San Michele all'Adige, Trento, Italy

9 University Paris-Est, ESTP, Cachan, France

a Shared last authorship (equal contributions)

* corresponding author: Valentin H. Klaus, University of Bochum, oekologie-geo_AT_ruhr-uni-

bochum.de

ORCiDs

Valentin Klaus: 0000-0002-7469-6800

Davide Andreatta: 0000-0001-6821-1152

Abdesslam Chai-allah: 0000-0002-1787-9627

Nathan Fox: 0000-0002-2816-9751

Franziska Richter: 0000-0001-7865-9903

Abstract (236 words)

Temperate grasslands provide various cultural ecosystem services that are appreciated in diverse

ways. Capturing these diverse appreciations requires different methodological approaches, such as

questionnaire surveys and social media analyses. In this study, we combined the potential of both

approaches to capture two aspects of what people appreciate in agricultural grasslands, i.e., the

aesthetic quality of differently managed plant communities and the objects frequently found in

grassland-based social media images. The two complementary approaches showed that people

preferred colourful flower- and species-rich grasslands over grass-dominated and fertilised swards.

Social media analysis highlighted that people mainly photographed flowers, followed by livestock

and/or wildlife, but this depended also on the social media platform used.

In conclusion, people's appreciation was clearly related to the intensity of grassland management and

to the level of biodiversity, with a preference for extensively managed grasslands with diverse flowers

and wildlife. Yet, we also found significant differences between (i) conservationists and agricultural

professionals in the aesthetic appreciation of the plant communities, and (ii) between common visitors

and naturalists in their social media content. Our results suggest that extensive management and

ecological restoration can be used to increase cultural grassland ecosystem services by enhancing the

richness of forbs, flowers and other attractive wildlife. Thus, targeted management is necessary to

maintain and enhance the attractiveness of grassland landscapes and subsequently increase the health

benefits that can be associated with these cultural grassland ecosystem services and human-nature

contacts.

Keywords (6)

Cultural ecosystem services, landscape aesthetics, biodiversity, flowers, social media, image analysis

1. Introduction

Cultural ecosystem services (CES) and positive human-nature interactions are of great relevance for

human wellbeing and people's perception of nature and nature conservation, but due to

methodological challenges and a lack of data they are widely understudied (Plieninger et al., 2013;

Cheng et al., 2019). Insight into the drivers of CES and how these can be enhanced is required to guide

land management in all major land use types, especially since intensification and global change

threatens CES (Kosanic and Petzold, 2020; Straffelini et al., 2024). Grassland is a major land use type in

many regions around the world and provides a wide range of relevant ecosystem services to local

populations (Bengtsson et al., 2019). Intensification of grassland management as well as climate

change were shown to threaten grassland CES (Allan et al., 2015; Schils et al., 2022; Straffelini et al.,

2024). Yet, grasslands were frequently ignored by international strategies (Bardgett et al., 2021),

highlighting that grassland CES are not only threatened but also widely overlooked. Thus, we urgently

require a better understanding of how grasslands provide CES to be able to develop strategies to

maintain and enhance these. Therefore, we must understand what exactly people find interesting and

attractive in grasslands, and how this can be supported.

A major CES is the aesthetic value of an ecosystem, which is also closely related to other services such

as recreation, tourism, culture and heritage, as well as landscape aesthetics (Bengtsson et al., 2019;

Plieninger et al., 2013). Previous work found grasslands in the temperate zone to be hardly perceived

to contribute to landscape-scale CES (Ridding et al., 2018; Fox et al., 2022a). However, Schils et al.

(2022) found grasslands to contribute more to landscape aesthetics and recreation than forests or

arable land. Furthermore, grasslands were found to be important landscape compartments as they

enable landscape views (Fox et al., 2022b; Chai-allah et al., accepted). In addition, in a previous study,

Richter et al. (2024) found extensive management and use as meadow (versus pasture) to result in

higher perceived aesthetic quality, while organic farming did not have an impact on grassland

aesthetics. Furthermore, a high diversity of plant species was found to be appreciated by visitors of

temperate grasslands (Lindemann-Matthies et al., 2010). Yet, a mechanistic understanding of what

people aesthetically prefer in permanent grasslands and how this can be enhanced by land

management is still missing.

In grasslands, many aspects can contribute to the aesthetic value: the ecosystem itself, i.e., plant and

animal communities and their characteristics, but also land management, such as grazing animals as

well as objects such as single trees or old barns (Van Berkel and Verburg, 2014). Therefore, assessing

the aesthetic value of grasslands requires a comprehensive view on the many characteristics of this

ecosystem, and likely also different methodological approaches that allow insight into the aspects of

aesthetic perception.

The most frequently employed approaches to understanding ecosystem aesthetics comprise

questionnaire surveys that interview people either in front of the ecosystem in question (e.g.,

Lindemann-Matthies et al., 2010) or with images as visual stimuli (e.g., Fischer et al., 2018). The

extraction of specific characteristics of images linked to respondents' appreciation is typically carried

out by hand as well as questionnaires to collect information about the actual appreciation (Richter et

al., 2021; Wang et al., 2024). Examples of this include visually estimating flower cover and other

structural properties in the field. However, this method is time-consuming and challenging to scale and

standardise. Automatic spectral analysis of the images may be able to help get around this restriction

since spectral features of images were shown to be associated with the composition and structure of

plant communities (Torresani et al., 2024). However, the relationship between the spectral

characteristics of grassland images and appreciation remains largely unexplored in the scientific

literature.

More recently, studies used the vast crowdsourced data uploaded to social media and citizen science

platforms (e.g., Havinga et al., 2020; Chai-allah et al., 2023; Ghermandi et al., 2023). Such approaches

benefit from the large amount of data that is being uploaded, such as spatially explicit images, and

employ advanced data science tools for content analysis (Fox et al., 2021; Schirpke et al., 2023).

Combining both questionnaire surveys to indicate 'stated-preferences' and crowdsourced images

analyses to indicate 'revealed preference' has the potential to considerably advance our understanding

of grassland aesthetics (Cheng et al., 2019).

In this work, we explore what grassland characteristics are appreciated by people and how these are

related to management measures (Figure 1). Here, we combine a questionnaire survey, an automatic

spectral image analysis, and a crowdsourced image analysis to understand what people find attractive

and of interest in agriculturally managed grasslands in Switzerland. Further, we conducted (i) an in-

depth analysis of a structured questionnaire survey with people's ratings of 92 grassland images taken

along a land-use intensity gradient (Richter et al., 2024) and (ii) a social media-based image analysis

extracting information on relevant objects that have been photographed in grasslands and uploaded

to two crowdsourced data sources. Thus, we studied two presumably simultaneously relevant aspects

of the aesthetic value of grasslands, i.e., people's perception of the grassland plant community, as

affected by different anthropogenic and environmental drivers, and the objects people photograph

and share online. Some of the results of this questionnaire were used by Richter et al. (2024) to

coarsely characterise the aesthetic value as an aspect of grassland multifunctionality. This analysis,

however, goes much deeper and overcomes the purely descriptive approach of the previous study by

conducting a mechanistic assessment of what drives people's aesthetic rating of the plant

communities.

We defined two sets of hypotheses, as shown in Figure 1.

What grassland characteristics are appreciated by people, and how can these be related to management measures?

Part 1 - The aesthetic value of grassland plant communities

What?

1. People's aesthetic appreciation

2. Prediction of peoples rating from image characteristics

How?

Online questionnaire

Spectral analysis

Part 2 - Objects of interest for photography in grasslands

What?

1. Attractive objects in Flickr & iNaturalist images

2. Attractive wildlife taxa in Flickr & iNaturalist images

How?

Content analysis

Google Vision API

Taxonomic analysis

BioCLIP

Hypothesis 1.1. – Positive predictors of the aesthetic value are related to colorful flowering and high biodiversity, while negative predictors are associated with intensive grassland management.

Hypothesis 1.2. – Standardized automatic image analysis can predict people's aesthetic appreciations via (a) the extraction of flower cover and color richness and (b) the diversity in spectral characteristics of the images.

Hypothesis 2.1 – The most frequent objects in grassland images are easy to find, frequent and unobscured: flowers and livestock.

Hypothesis 2.2 – Besides, images also show familiae and genera of rather rare species with conservational relevance, such as Orchids, with a higher proportion of the latter being uploaded to an ecologically targeted platform (i.e., iNaturalist) than to a common platform (i.e., Flickr).

Figure 1. Conceptual figure highlighting the two approaches used in this study, the respective methods and hypotheses. While part 1 uses images (photographs) taken in a standardised way, the images in part 2 were uploaded by social media (i.e., Flickr and iNaturalist) users in the study region (Figure 2).

2. Materials and methods

Because of the multi-faceted and complex nature of CES, this work combines two complementary approaches assessing two different but presumable simultaneously relevant aspects of the aesthetic attractiveness of grasslands in the northwest of Switzerland (section 2.1). First, we assess the attractiveness of the vegetation matrix, i. e., the grassland plant community, via a photo-based structured survey on people's aesthetic perceptions (section 2.2). Second, we assess objects of interest in grasslands with an analysis of images uploaded to two crowdsourced platforms (section 2.3).

2.1. Study system

We focus our work on the northwest of Switzerland (Figure 2), a grassland-dominated region with a landscape-scale land-use composition comparable to many other temperate European regions. Grassland dominates the two biogeographic subregions studied, i.e., the intensively used plateau, in which grassland is combined especially with arable land, and the mountain range "Swiss Jura",

composed of limestone ridges in which grassland and forest occur. While the plateau stretches

between 400 to 800 m a.s.l., the peaks of the mountainous part reach up to 1200 to 1700 m a.s.l. We

chose this region because its grassland types and their management are typical for central Europe.

These grassland types are (i) productive sown leys (temporary grassland) that can be included in crop

rotations, (ii) productive and intensively managed permanent grasslands and (iii) semi-natural

grasslands, which are usually managed according to agri-environmental schemes that considerably

restrict management intensity, especially fertilisation (Klaus et al., 2023; Lüscher et al., 2019). No

alpine areas and no natural grasslands occur in the study region.

Figure 2. Study region composed of the two biogeographic subregions "Jura" and "Plateau" in
northwestern Switzerland. In the two areas the distribution of grasslands is shown in green according
to Huber et al. (2022). Shaded relief of Switzerland taken from swisstopo (2020). While the
standardised images for part 1 were taken in the canton of Solothurn, indicated by a brown frame,
images taken from social media platforms (part 2) originate from the whole study region.

2.2. Part 1 - structured survey on people's aesthetic perceptions

2.2.1. Questionnaire survey

To conduct a questionnaire survey on people's aesthetic appreciation, we took 92 images of permanent grasslands in the canton of Solothurn, which is located right in the middle of the overall study area and covers both the Jura and the Plateau subregions (Figure 2). Images were taken from mid-May to mid-June 2021. Respective permanent grassland plots were selected along a gradient in management intensity based on the grassland typology provided by official agricultural statistics. The main four grassland types in this typology are meadows (mainly mown) and pastures (mainly grazed), which can be intensively managed (fertilisation allowed) or extensively managed (fertilisation prohibited; management according to the Swiss agri-environmental scheme "extensively managed pastures" and "extensively managed meadows"). For more details on the typology and the management of the grassland types see Klaus et al. (2023).

All images show a close-up of the plant community in an area of approximately 2 m × 2 m from the height of one meter above ground level during peak flowering in late spring, without any objects, such as livestock or other animals, or landscape views, in a resolution of 4032 x 3024 pixels. Most images were taken before harvest, but in twelve cases, grasslands were already grazed during the time of sampling. Eleven out of these twelve plots were pastures, which can generally be grazed early in the season (as from end of March). The resulting "signs of defoliation" were considered by using dummy coded variables in the statistical analysis (see section 2.2.2.). In addition to the permanent grasslands, two images of sown (temporary) grasslands at comparable phenological stage were added to the final set to also consider this grassland type, resulting in a total of 92 images included in the questionnaire.

A German-language online questionnaire was online from the 11th of January to the 7th of March 2022, asking people for their personal perception of the aesthetic quality of grassland plant communities shown on 92 images. Aesthetic quality was available as a 5-point Likert scale from 0 = *unattractive* to 5 = *attractive*. The survey was set up with QuestionPro (QuestionPro Inc, Austin, TX, United States),

with ten of the 92 images randomly chosen per questionnaire. The link to the survey was distributed

via different pathways, i.e., (i) via e-mail to professional stakeholders in agriculture, nature

conservation, tourism, and/or botanical gardens in Switzerland, (ii) via German-language Twitter (now

X) posts and mailing lists, and (iii) further private and professional networks. The survey text and all

pictures are available in supplementary material 1. Since the survey was completely anonymous, it did

not require an approval of the institution's ethical board.

The average rating per plot was computed and used as the indicator for people's aesthetic appreciation

of the shown grassland plant community. We further asked about the professional work of the

respondents, offering the categories "agriculture", "ecology/nature conservation", "both professional

areas", "neither of both, other professions or retired". We therefore additionally computed the

average aesthetic value for all images for three separate groups, omitting the group responding "both

agriculture and ecology/nature conservation" because we expected them to overlap with the other

categories (and due to very few responses in the category, see below). A linear mixed effects model

was used to test for significant differences in aesthetic ratings among the professional groups, with

plot as random factor.

2.2.2. Explanatory variables of plant communities

To understand the drivers of the aesthetic appreciation of people, we recorded information on plant

community composition and the agricultural management of the grasslands, and we extracted visible

and spectral characteristics of the images. Together with taking the 92 images in permanent

grasslands, we at the same time conducted vegetation records in two 2 m × 2 m areas in each site. We

visually assessed all vascular plant species and their ground cover as well as the total ground cover of

cryptogams. The two records were near the area shown on the image but were not necessarily

identical due to logistic reasons. However, since the vegetation of the grasslands was widely

homogeneous and any accidentally occurring clusters of weeds or other atypical vegetation were

avoided, we consider the vegetation records representative for the plant communities shown on the

images. Based on the vegetation records, we calculated three plant diversity metrics (i.e., number of

vascular plant species, evenness and Shannon diversity) as well as the cover sums of all grass and all

forb species, with the latter including legume species. Furthermore, as an indicator of the agricultural

value of a plant community, we calculated abundance-weighted forage quality indicator values

according to Briemle et al. (2002) based on plant indicator values. After the vegetation records, we

sampled the above-ground plant biomass as a measure of productivity (i.e., peak standing crop) and

measured the height of the vegetation by placing a laminated A4 sheet on top of the vegetation,

parallel to the ground and then recording the height of the sheet with a ruler. Height measurements

represent a mean of four replicates per plot taken in each corner of each vegetation record.

Information on topographical features, i.e., slope and elevation of each plot, were extracted from a

Digital Elevation Model (25-m grid resolution, European Union 2018) using QGIS (version 3.22). To

gather detailed information on the specific management of the grasslands, we collected management

data for 2021 and 2022 from farmers. We recorded the fertilisation intensity (sum of available nitrogen

in organic and mineral fertiliser per ha per year applied as fertilizer – not including defecation of

animals grazing on the area), the mowing frequency (number cuts per year), and the grazing intensity

(livestock unit days per ha per year) following Blüthgen et al. (2012). Intensity measures were averaged

for the two years to provide a stable mean. For further details on the images, the questionnaire and

the vegetation records see Richter et al. (2024).

To gather information on the actual plant community visible in the photographs, one person coded

information from all images, i.e., flower cover (coded as % and quasi-metric categories of *1* = 0-1%, *2*

= 1-5%, *3* = 5-15%, *4* = more than 15% of image filled with flowers), excluding green and grassy flowers,

and flower colour richness. We defined the colour classes as follows: yellow flowers (such as

Ranunculus spp., *Rhinantus* spp., *Crepis biennis*, and *Taraxacum* sp.), white flowers (such as

Leucanthemum ircutianum, *Capsella bursa-pastoris*, *Daucus carota*, *Trifolium repens*, *Galium mollugo*,

and *Bellis perennis*), red flowers (such as *Trifolium pratense* and *Onobrychis viciifolia*), violet and

similarly coloured flowers (e.g., *Knautia arvensis, Scabiosa columbaria,* and *Salvia pratensis*), and blue

flowers (e.g., *Ajiuga repens*, and *Veronica* spp.). In addition, other image properties were binary coded

(i.e., signs of defoliation (Yes/No), bare soil visible (Yes/No), litter layer visible (Yes/No), agricultural

weeds prominently visible (Yes/No), and vegetation structure (Light/Dense)). Agricultural weeds found

on the images were *Rumex obtusifolius*, *Colchicum autumnale*, *Senecio* spp., *Rhinanthus* spp., *Carlina*

spp., *Cirsium* spp., *Anthriscus sylvestris*, and *Heracleum sphondylium*. To relate all these characteristics

of the plant communities and associated measures to the aesthetic ratings, we used Spearman

correlations for metric and quasi-metric data as well as ANOVAs for categorical data. These analyses

were run for all responses together and separately for the professional groups. Diagnostic plots were

used to ensure model assumptions were met.

To get insights into the most common taxa found in the different grasslands in the study area, and to

compare this with the Results related to hypothesis 2.2, we in a first step used the "classification"

function from the package "taxsize" (v. 0.9.102, Chamberlain and Szocs, 2013) to extract the higher-

level taxonomy from the species present in the vegetation records. We then used this information to

calculate the cover of a) different plant families and b) different plant genera per grassland (using the

mean covers of the two plots per grassland), as well as the number of grasslands in which a) different

plant families and b) different plant genera were present in the vegetation records. These results were

visualized in the form of barplots using ggplot2 (v. 3.5.1, Wickham, 2016). This analysis was carried out

in R (v. 4.4.2, R Core Team, 2024).

2.2.3. Automatic extraction of flower and spectral diversity metrics

In addition to estimating flower cover and flower colour richness visually, we automatically extracted

flower cover and flower colour richness through supervised image classificationto establish uniform

methods for quantifying floral attributes that relate to people's aesthetic appreciations. First, we

defined the pixel classes as follows: green vegetation and soil, yellow flowers, white flowers, red

flowers, violet flowers, and blue flowers. Then, we developed a labelled dataset by randomly selecting

a 1000 pixels × 1000 pixels image patch (corresponding to around 50 cm × 50 cm at the ground level)

in each image, plotting in RGB colours using the "plotRGB'' function of the "raster" package (Hijmans,

2023), labelling around 50 pixels per patch by clicking on the image to retrieve the X and Y coordinates

using the "locator" function of the "graphics'' package, and assigning to each pixel the class to which

it belongs. Since labelled pixel class frequency was very unbalanced with only a few violet and blue

flower pixels available, additional image patches where violet and blue flowers occurred were

identified in the RGB images, and pixels labelled. Next, we removed pixels too close to each other from

the dataset, i.e., pixels falling in the same pixel after downscaling the images with an aggregation factor

equal to 10. The resulting dataset consisted of 4832 labelled pixels, with 117 to 3315 pixels per class.

The reflectance values in the red, green and blue bands of the images in labelled pixels were extracted

to obtain a dataset reporting image ID, pixel class, XY coordinates, and reflectance values for each

labelled pixel. Afterwards the dataset was split into training and validation by randomly assigning 66%

of images for training, and 33% of images for validation. The training dataset was used to develop a

random forest classifier using the "randomForest'' function of the "randomForest" package (Liaw and

Wiener, 2002) with default parameters. Classifier performance was assessed on the validation dataset

by computing the overall accuracy, the Kappa accuracy, and the mean F1 score of the six classes

(Congalton and Green, 2009). The compiled classifier was used to classify all images except for two,

which were discarded since they had different dimensions. While deriving the flower cover from the

classified image through image statistics was straightforward, defining the flower colour richness

required the definition of the percentage threshold above which a flower class is to be considered as

present in the image. We tested 2000 possible thresholds from 0% to 1% and chose the threshold

resulting in the highest correlation between manually and automatically estimated flower colour

richness.

Since we hypothesised that people's appreciation could not only be affected by flower cover and

flower colour richness, but also by the evenness of the distribution of the pixel classes, with higher

diversity associated with higher appreciation, we applied two diversity metrics widely used in grassland

ecology studies, Shannon and Simpson, to the frequency distribution of the pixel classes in each image.

The six-pixel classes were considered as species, and pixel counts as species abundances. Diversity

metrics were computed using the "diversity" function of the "vegan" package (Oksanen et al., 2022).

The relationships between appreciation and flower cover and between appreciation and diversity metrics were assessed by fitting second-order polynomial regression models, since the relationships were not linear. Finally, since we hypothesised images with more differences in the reflectance spectra between the pixels (e.g., less homogeneously green) to be associated with higher people's aesthetic rating, we assessed the diversity in spectral characteristics of all 92 images of permanent grasslands. The diversity in spectral characteristics of images is not unequivocally defined, but suitable metrics can be found in the field of remote sensing of terrestrial plant biodiversity (Wang and Gamon, 2019). The rationale behind spectral diversity metrics is that the increased spectral variability reflects an increased variety of habitats, species, and organs, since different habitat, species and organs respond in their own way to incoming solar radiation according to their pigment, water, and biochemical content, as well as leaf and canopy structure. The diversity metrics we applied here were: whole image standard deviation, moving window standard deviation, multidimensional Rao index. Since shadowed pixels may lead to noise rather than enhancing the information content at very-high spatial resolution, image resolution was reduced by applying a downscaling factor equal to 10 before computing the diversity metrics, as frequently applied in the optical diversity context (Rocchini et al., 2016).

Standard deviations were separately computed on each of the three bands (red, green, blue) and the resulting statistics were then averaged to obtain a standard deviation (SD) for each image. Computing standard deviations in square moving windows and not only in the whole image allowed us to measure the spatial variability occurring at the scale of interest (i.e., flower scale in our study), and that can therefore be attributed to flowers. A window of 5 pixels × 5 pixels was chosen, corresponding to around 2.5 cm × 2.5 cm. As described in Rocchini et al. (2021), we first computed a diversity map by assigning to each pixel the SD of reflectance values in a square moving window of five pixels, and then obtained a summary metric for each image by averaging the diversity map. Lastly, we computed the Rao's index, which quantifies the difference in reflectance values between two pixels drawn randomly with replacement from a set of pixels by considering their abundance and their pairwise distance (Hauser et al., 2021; Perrone et al., 2024). Rao's index is a multidimensional index, which means that the

diversity in multiple variables can be evaluated at the same time. In our study, the three bands (red,

green, blue) were used. Using the "paRao" function of the "rasterdiv" package, Rao's index was

calculated in square moving windows of five pixels, applying Euclidean distance as a distance metric

(Rocchini et al., 2021).

2.3. Part 2 - Image analysis using crowdsourced platforms

2.3.1. Collection of images and platforms involved

The second part of the study was to identify objects in grasslands that people are interested in. Here,

we assumed that the content of images (objects) uploaded to online platforms is indicative of people's

self-reported appreciation (Lee et al., 2022). To gather publicly available images posted online and

georeferenced within the administrative boundaries of the study region (Figure 2), two crowdsourced

platforms Flickr and iNaturalist were selected. Both platforms are commonly used by people to share

their interactions with plant and animal species (Havinga et al., 2020). Flickr is a social media platform

where photographers share their visual appreciation of nature, including casual interactions with

nature, including but not limited to photographs of species, whereas iNaturalist is a citizen science

platform where users specifically focus on sharing their observations of species (Havinag et al., 2020).

We retrieved all publicly shared georeferenced images and their associated metadata, including user

and image ID, the date of taking the image and a URL link to the image using the "photosearcher"

package in R (Fox et al., 2020) to access the Flickr Application Programming Interface (API) and the

"rinat" package in R (Barve and Hart, 2017) for the iNaturalist API. The data collected covers the years

2004 to 2023 and, after removing duplicate images, results in 1,718,001 unique images from 27,949

users in Flickr and 143,850 images from 3745 users in iNaturalist in the whole study region. These

images were filtered for their location in grasslands using a grassland habitats map from Huber et al.

(2022), resulting in a total of 88,025 georeferenced images by 3,831 Flickr users and 10,772

georeferenced images shared by 725 iNaturalist users. These images were then analysed for their

content, focusing on the objects and especially the biological species depicted, highlighting the self-

reported species of users on both platforms as an indicator of revealed preference (Lee et al., 2022).

2.3.2. Extraction of objects from images

The content analysis consists of two methodological steps. First, we filtered all images for those focusing on objects belonging to nature. Second, we analysed what exactly was on the images in terms of specific plant and animal species. This two-step approach helped to save computational effort, since only the zoomed-in images of natural objects derived by step one were used for the in-depth analysis.

To extract only those images showing nature and natural objects, the content of the images was extracted using Google Cloud Vision (GCV, https://cloud.google.com/vision/) by accessing its API using the R package "imgrec" (Schwemmer, 2019). GCV is a pre-trained machine-learning algorithm that has demonstrated a high ability to detect both natural and non-natural features in social media images (Fox et al., 2021; Wilkins et al., 2022). To extract nature images, we first ran all images through the algorithm using the "label" feature, which returns labels describing the image content, including natural labels (e.g., grassland, landscape, flower) and non-natural labels (e.g., building, road, car). For each image, we extracted up to 20 labels, considering only those with a confidence score of 60% or higher (Gosal et al., 2019) and a total of at least 15 occurrences in the dataset. Second, we adopted the framework proposed by Chai-allah et al. (accepted), classifying the GCV labels into non-natural and natural labels, with the latter being classified as landscape (e.g., landscape, mountain, sky) or objects/species (e.g., flower, bird, insect). The full label classification is listed in Table S3 in supplementary material 2. From nature images (i.e., images with more than 50% nature labels), we selected those with more than 50% object labels to build up our species appreciation dataset. Finally, since geolocations can be prone to error (Zielstra and Hochmair, 2013), we limited our analysis to images in grasslands by selecting only images with the presence of grassland-related labels (i.e., grass, grassland, meadow, and pasture), resulting in a final dataset of 5631 images shared by 808 Flickr users (mean = 6.9) and 4944 shared by 451 iNaturalist users (mean = 10.9).

We ran the GCV again to detect objects using the "object" feature. This feature has better capabilities to accurately detect the species present in the images compared to the "label" feature (https://cloud.google.com/vision/docs/features-list). For each image, we retrieved up to three unique

objects. The confidence score for the retrieved objects ranged from 0.5 to 0.99 (mean = 0.79), which

can be considered high. Thus, we decided to not further exclude any objects based on the confidence

score. However, we limited our analysis to objects that occurred at least twice in total and excluded

generic objects (i.e., animal and plant), objects describing pet and zoo animals (e.g., cat, dog, and

brown bear) and non-natural objects (e.g., person, bicycle and car). The complete list of included and

excluded objects can be found in Table S4 in supplementary material 2.

Since the hierarchical functioning of the GCV algorithm can classify an image using multiple related

objects, such as "bird" as the first object and "swan" as the second object, or one object, simply "bird"

or "swan", a pattern predominantly occurring for birds and insects, we decided to simplify the

classifications by grouping objects. All bird-related objects (i.e., bird, duck, sparrow, goose, swan,

falcon, raven, eagle, magpie, owl, blue jay and woodpecker) were merged into the category "bird" and

all insect-related objects (i.e., insect, moths and butterflies, butterfly, dragonfly, beetle, bee, ladybug

and caterpillar) into the category "insect". We extracted only unique objects returned by GCV, meaning

within an image, three objects belonging to the group "bird" would be reduced to one "bird" object.

This step of the analysis revealed a general overview of the objects on the images taken without diving

into a taxonomic analysis.

2.3.3. Extraction of wildlife from images

To better understand the wildlife shown in the images, we carried out a taxonomy analysis using the

computer vision model BioCLIP (Stevens et al., 2024). It is derived from the CLIP (Contrastive Language–

Image Pre-training) model developed by OpenAI and trained on a large dataset of images including

those from the Encyclopedia of Life project (https://eol.org/), which is a comprehensive collection of

images and information about various species. Since the identification error of the model increases

with increasing taxonomic detail, from kingdom to species (Stevens et al., 2024), we extracted different

taxonomic levels of the objects found in the images. For each retained image, we first performed a

taxonomic analysis at the family level. For this, only families identified with a confidence score of 60%

or higher were considered. Next, we performed a genus-level analysis on these pre-filtered images,

again retaining only those genera identified with a confidence score of 60% or higher.

3. Results

3.1. Predictors of people's aesthetic appreciation of the plant communities (Hypothesis 1.1)

In total, 522 participants answered the questionnaire, which led to, on average, 55.3 ratings per

grassland image (min = 39, max = 73). The questionnaire was answered by 47.9% female and 48.1%

male respondents (1% diverse, 3.1% NA), which were mainly from Switzerland (48.9%) and Germany

(44.3%), plus few from Austria (3.1%) and further countries or NA (3.8%). Age distribution of the

respondents was reported in steps of ten years, with most people being quite evenly distributed

between 20 and 70 years (Figure S1 in supplementary material 1). Yet, few younger (1.1% 18-20 years)

and some older people (3.4% 70-90 years) were also responding (2.5% NA). Of all people responding

on the profession (3.3% NA), 23.2% stated a relation to agriculture, 22.4% to ecology/nature

conservation, 8.6% to both professional areas, and 42.5% to "neither of both, other professions or

retired".

People's ratings of the aesthetic value of the 92 images of grassland plant communities varied widely,

from min 1.79 to max 4.54 (mean = 3.15 and SD = 0.65; allowed range 0 = *unattractive* to 5 = *attractive*;

Figure 3). Basic statistics of the perception of the three professional groups "agriculture",

"ecology/nature conservation", and "neither of both, other professions or retired" (simplified to

"*other*" in what follows) were highly similar (Table S1 in supplementary material 1) and mean values

did generally not significantly differ (linear mixed model, $p > 0.2$). The aesthetic values of all

respondents correlated strongly with those of all separate professional groups ($r_{Spearman}$ 0.87 to 0.96, p

< 0.001; Table 1). This is why the following is focused primarily on all responses together. Yet, ratings

of people working in agriculture and in ecology/nature conservation were less closely correlated

($r_{Spearman}$ 0.64, but $p < 0.001$; Table 1; Figure S2 in supplementary material 1), highlighting that the

aesthetic rating somewhat varied among professional groups. For example, professionals related to ecology/nature conservation rated images of unfertilised grasslands on average a bit higher than agricultural professionals (3.82 vs. 3.42), and vice versa for intensively managed, fertilised grasslands (2.70 vs. 2.96; Figure S2 in supplementary material 1).

Correlation analysis revealed many positive and negative predictions of the aesthetic value of the plant communities (Table 1). The strongest positive predictors based on all responses were flower cover, flower colour richness, stand height, the cover sum of forbs and plant diversity metrics (Shannon diversity and species richness). A weak but significantly positive effect of weeds being visible in the images was found for the professional groups "ecology/nature conservation" and "other", but not for "agriculture" (Table S2 in supplementary material 1). Furthermore, and in contrast to the other professional groups, ratings of people related to agriculture were most positively correlated with stand height but much less so with flowers, forbs, and plant diversity (Table 1). The strongest negative predictors of all ratings were cover sum of grasses, forage value indicator, grazing intensity, and fertilisation intensity. While a high cover of grasses was obviously directly visible in the images, and strongly negatively related with the cover of forbs ($r_{Spearman}$ = -0.57, p < 0.001), the other measures cannot be seen on the images but are indicators for intensive management, which generally reduces positive predictors such as flower cover and plant diversity (e.g., correlations of fertilisation intensity with flower cover and plant species richness: $r_{Spearman}$ = -0.34, p < 0.001 and $r_{Spearman}$ = -0.7, p < 0.001, respectively). However, agricultural professionals´ ratings showed less strong negative relationships with the formerly mentioned factors except for grazing intensity (Table 1). While grazing intensity and, for most groups, also fertilisation intensity were clearly negatively related to aesthetic ratings, cutting frequency was either insignificant or weakly positively related to aesthetic ratings (Table 1). This might, however, be indirectly driven by a strong negative correlation between cutting frequency and grazing intensity ($r_{Spearman}$ = -0.72, p < 0.001), meaning that if grasslands were not intensively grazed, they had to be frequently cut.

In addition, signs of defoliation, bare soil, and a thick litter layer in the images were also strongly

negatively related to the aesthetic ratings of all groups (Table S2 in supplementary material 1). There

was, however, one notable exception: A litter layer did not significantly decrease the aesthetic ratings

of people related to ecology/nature conservation. Note that those twelve images showing signs of

defoliation and potentially a lot of bare ground were mainly pastures, which were stocked particularly

early, explaining the negative effect of grazing intensity on the aesthetic value.

Figure 3. The four images with lowest (upper panel) and highest (lower panel) average ratings
considering all responses (n = 522). Images that show signs of use (defoliation) and bare soil were
perceived as particularly unattractive, while high flower cover and colour richness were perceived as
attractive (Table 1, Table S2 in supplementary material 1).

Table 1. Spearman correlations between aesthetic ratings of different professional groups (in grey above) and with various grassland characteristics. Blue colours show positive predictors and red ones negative. While flower cover and colour richness have been extracted directly from the images used in the survey, all other information is based on field samplings and a land-use questionnaire with farmers. Significant levels: "***" = p < 0.001, "**" = p < 0.01, "*" = p < 0.05, "(.)" = p < 0.1, and "n.s." = p > 0.1.

	Ratings all	Ratings agriculture	Ratings ecology	Ratings other
Ratings agriculture	0.87 ***	-		
Ratings ecology	0.89 ***	0.64 ***	-	
Ratings other	0.96 ***	0.78 ***	0.83 ***	-
Categ. flower cover (picture)	0.66 ***	0.38 **	0.66 ***	0.72 ***
Flower cover perc. (picture)	0.6 ***	0.35 **	0.59 ***	0.65 ***
Number flower colors (picture)	0.6 ***	0.46 ***	0.61 ***	0.56 ***
Stand height	0.6 ***	0.65 ***	0.4 ***	0.6 ***
Cover sum forbs	0.5 ***	0.29 *	0.52 ***	0.52 ***
Number plant species	0.42 ***	0.22 (.)	0.56 ***	0.37 **
Shannon index plants	0.42 ***	0.23 (.)	0.52 ***	0.41 ***
Evenness plants	0.32 *	0.2 n.s.	0.35 **	0.34 **
Cutting frequency	0.18 n.s.	0.27 *	0.02 n.s.	0.21 (.)
Slope	0.12 n.s.	0.02 n.s.	0.21 (.)	0.11 n.s.
Peak standing crop	0.1 n.s.	0.26 (.)	-0.11 n.s.	0.14 n.s.
Cover sum cryptogams	-0.01 n.s.	-0.15 n.s.	0.23 (.)	-0.07 n.s.
Elevation	-0.13 n.s.	-0.08 n.s.	-0.11 n.s.	-0.11 n.s.
Fertilisation intensity	-0.33 *	-0.11 n.s.	-0.51 ***	-0.3 *
Grazing intensity	-0.48 ***	-0.44 ***	-0.43 ***	-0.48 ***
Forage value indicator	-0.48 ***	-0.21 (.)	-0.63 ***	-0.45 ***
Cover sum grasses	-0.49 ***	-0.22 (.)	-0.65 ***	-0.45 ***

3.2. Spectral characteristics related to people's aesthetic perception (Hypothesis 1.2)

The standardised automated process to extract flower cover and flower colour richness produced accurate and insightful results, reflecting the clear correlations with people's aesthetic appreciation. The validation process demonstrated high accuracy of the RF classifier, with an overall accuracy of 0.95, a Kappa accuracy of 0.88, and a mean F1 score of 0.85. The automatically extracted flower cover for each image showed high correlations of 0.79 (p<0.001) when compared to manually coded data (Figure S3a in supplementary material 1). However, the automatically extracted flower cover values were significantly lower than those estimated manually, indicating a perceptual bias in human evaluation. We calculated flower colour richness for each image by counting the number of distinct flower classes that covered more than 0.005% of the image area, i.e., 600 pixels. This specific threshold yielded the highest correlation between manually and automatically estimated flower colour richness ($r_{Pearson}$ = 0.64; p <0.001; Fig. S3b in supplementary material 1). High flower cover was associated with

high appreciation scores, as in the correlation analysis previously shown in Table 1. However, at high levels of flower cover, this was only true when combined with high flower colour richness, as shown both by the manual and automatic estimates (Figure 4a). Images with very high flower cover (over 5%) typically featured uniform yellow blooms (such as dominance of *Taraxacum officinale* or *Ranunculus acris*) and received moderate appreciation scores. In contrast, the most highly appreciated images had moderate flower cover but displayed a flower colour richness between 4 and 5. Shannon and Simpson diversity indices of pixel classes frequency distribution explained a large portion of the variability in aesthetic appreciation (R^2 of 0.42 and 0.43, respectively). The high correlation of these indices with flower cover ($r_{Pearson}$ 0.98 and 1.00, respectively; both with $p<0.001$), confirmed that images with more evenly distributed class frequencies were those with higher flower cover, which were generally more appreciated   by   viewers.

The three spectral diversity metrics were highly correlated with each other ($r_{Pearson}$ between 0.862 and 0.999, $p < 0.001$), uncorrelated to flower colour richness, and significantly positively correlated to flower cover ($r_{Pearson}$ between 0.43 and 0.46; $p <0.001$). However, the spectral diversity metrics were not correlated to people's aesthetic appreciation ($p > 0.05$, Figure 4b).

489   Figure 4. Relationships between aesthetic people's appreciation and image flower presence metrics
490   (a), colour classes diversity indices (b) and spectral diversity metrics (c). Flower cover, colour richness
491   and the Shannon and Simpson indices of colour classes frequency distribution were associated with
492   appreciation (aesthetic ratings), whereas the metrics measuring spectral diversity were not.

3.3. Frequent objects in grassland images (Hypothesis 2.1)

Flowers were the most photographed objects on both platforms, with 2,500 images on iNaturalist (> 56% of all images) and 2,198 images on Flickr (> 46% of all images; Figure 5). Flowers were, for instance, photographed two times more often on iNaturalist and 3.6 times more often on Flickr than insects, the second most popular object on both platforms. Birds appear as the third most photographed object on both platforms, with a similar number of images on both platforms. However, they were photographed 2.8 times less often than insects on iNaturalist (439 versus 1,230) and 1.2 times less often than insects on Flickr (510 versus 612). Following flowers, insects and birds, Flickr users frequently took images of livestock, i.e., cattle, horses, goats and sheep, reflecting a much greater interest in domestic livestock than iNaturalist users. Deer and frogs were similarly photographed on both platforms and on iNaturalist there are also images of mushrooms, lizards, spiders, and snails in the top 10 objects. Over 60% of Flickr users and 64% of iNaturalist users included in our study uploaded at least one image of flower(s), with an average of 5.1 flower images per user on Flickr and 9.5 on iNaturalist. Similarly, the average number of images per user for insects and birds was higher on iNaturalist compared to Flickr.

Images uploaded followed a distinct seasonal distribution. On both platforms, flowers and birds were photographed mainly during spring (March to May), while insects were photographed more often during the summer (June to August). On Flickr, cattle were photographed fairly consistently throughout autumn, spring, and summer, but there was a large drop-off in the winter, with only 14 out of 484 cattle images taken. Cattle were photographed by 25% of all Flickr users, while this was not relevant for iNaturalist users. Meanwhile, on iNaturalist, 76% of mushroom images were taken in autumn (Table S5 in supplementary material 2).

Figure 5. The ten most frequently photographed objects in Flickr and iNaturalist. The complete list of
objects can be found in Table S4 in supplementary material 2. Total number of images included n =
4719 (Flickr) and n = 4464 (iNaturalist).

3.4. Wildlife in grassland images of Flickr vs. iNaturalist users (Hypothesis 2.2)

To understand differences in the content of images uploaded to a common, non-ecologically targeted

platform (Flickr) and an ecologically focused platform (iNaturalist), we extracted familiae and genera

of the wildlife depicted. Plant families make up the majority of the most frequently photographed

wildlife familiae, accounting for 55% of the top 20 most photographed familiae on Flickr and 65% on

iNaturalist (Figure 6). Asteraceae and Orchidaceae were the most photographed plant familiae on both

Flickr and iNaturalist, followed by Rosaceae on Flickr and Fabaceae on iNaturalist. Papaveraceae were

popular on Flickr but not on iNaturalist, whereas Lamiaceae was popular only on iNaturalist. While most top 20 plant familiae were forbs or familiae with mainly woody species (Rosaceae, i.e., roses, and Caprifoliaceae, i.e., honeysuckle), Poaceae (Gramineae, grasses) were also listed in both platforms, showing that not only plants with colourful flowers were among the frequently photographed taxa.

However, compared to the mean cover of plant families from the vegetation records in the canton of Solothurn (grasslands from which the pictures were taken, hypotheses 1.1 and 1.2), Poaceae were highly under-represented in both platforms (supplementary material 2, Figure S4). In the studied grasslands, Poaceae were the family with by far the highest mean cover per plot (61 %), followed by Fabaceae, Asteraceae and Ranunculaceae (all around 6-8%). Orchidaceae, the second most often photographed plant family on both flickr and iNaturalist, showed a very low cover in the studied plots (2.3 %), and was only present in four of the grasslands.

Five insect familiae ranked in the top 20 on both Flickr and iNaturalist, but with a higher proportion of all images on iNaturalist. Among these five popular insect familiae, Nymphalidae (brush-footed butterflies) were the most photographed on both platforms. However, they were about four times more frequent on iNaturalist compared to Flickr. The four other familiae show a difference in interest between Flickr and iNaturalist, with Flickr users photographing various families Syrphidae (hoverflies), Coenagrionidae (damselflies), Mantidae (praying mantis), and Acrididae (grasshoppers and locusts). In contrast, iNaturalist users primarily photographed further butterfly familiae (Geometridae, Pieridae, and Lycaenidae) and dragonflies (Libellulidae).

Vertebrates were rather rare within the top 20 families. Only two bird families made it into the top 20, both on Flickr and iNaturalist. These were Anatidae (water birds including ducks, geese, and swans) and Ardeidae (herons), with Anatidae being photographed more frequently on Flickr than iNaturalist. Frogs (family Ranidae) and deer (family Cervidae) appeared in the top 20 on Flickr, but not on iNaturalist.

The preference of Flickr and iNaturalist users for photographing plant species was also evident at the

genus level (Figure 7). Among the top 20 most popular wildlife genera, about two third or more were

plant genera, i.e., 65% on Flickr and 70% on iNaturalist. Although both Flickr and iNaturalist users

showed a preference for photographing plants, the specific genera differed. Flickr users most often

photographed *Papaver*, *Taraxacum*, and *Prunus*. Two orchid genera (*Ophrys* and *Spiranthes*) were also

popular. In comparison, the most photographed genera on iNaturalist were *Orchis* (an orchid genera),

Veronica, *Cirsium*, *Dactylorhiza*, and *Ranunculus*. Besides the woody species belonging to *Prunus*,

which were ranked third on Flickr, non-forb plant genera, with respect to the species occurring in

grasslands the study region, were quite rare. Yet, *Carex* (sedges) were ranked 11th on iNaturalist.

Compared to the cover of different plant genera from the vegetation records of the grasslands studied

in the canton of Solothurn, grass genera were (like already for the families) underrepresented on Flickr

and iNaturalist. The three grass genera *Poa, Lolium,* and *Festuca* had the highest mean cover

(supplementary material 2, Figure S5). These were followed by *Trifolium* and *Ranunculus*, which were

at the same time present in the highest number of grasslands. However, *Taraxacum* and *Veronica*,

genera that were often photographed on Flickr and iNaturalist, respectively, also had rather high mean

covers (3.8% and 1.3%, within top 10 and top 20, respectively) and were present in many of the

grasslands (81 and 83, respectively).

Bird genera were the in total second most popular group on Flickr, accounting for 25% of the top 20

wildlife genera, while only one insect genus (*Coenagrion*, a genus of damselflies) appeared on this list.

The top five bird genera photographed by Flickr users were mostly large water birds, with *Ardea*

appearing at the top, followed by *Anas* and *Cygnus*. In contrast, insect genera made up 25% of the top

20 on iNaturalist, while in turn only one bird genus (*Ardea*, a genus of herons) made it into the top-20-

list. In addition, the frog genus *Pelophylax* (a genus of true frogs) was among the top 20 wildlife genera

on Flickr. Meanwhile, iNaturalist users showed a clear preference for photographing butterflies, as the

five insect genera among the top 20 were butterfly genera. The top three of these were *Coenonympha*,

Maniola, and *Polyommatus*.

Figure 6. The 20 most frequently photographed familiae in Flickr and iNaturalist wildlife images. The
number given at each bar indicates the total number of identifications of this kind. Total number of
images included n = 2636 (Flickr) and n = 3628 (iNaturalist). The complete list of observed familiae can
be found in Table S6 in supplementary material 2.

Figure 7. The 20 most frequently photographed genera in Flickr and iNaturalist wildlife images. The number given at each bar indicates the total number of identifications of this kind. Total number of images included n = 2251 (Flickr) and n = 3084 (iNaturalist). The complete list of observed genera can be found in Table S7 in supplementary material 2.

4. Discussion

Our study identified clear differences in people's aesthetic appreciation of different plant communities and of different objects (as well as taxonomic units) in permanent grasslands. These different levels of appreciation (or potentially interest, in case of the images from social media and citizen science platforms; Ghermandi et al., 2023) can be seen as indication for differences in the CES and in human-nature interactions taking place within the grasslands studied. Furthermore, we found biodiversity to be directly and indirectly linked to appreciation and CES, which will be discussed in the following.

4.1. Drivers of people's aesthetic perception of grassland plant communities

4.1.1. Positive and negative characteristics of the grasslands and their management (Hypothesis 1.1)

Our study largely confirmed the first hypotheses by proving colourful flower- and forb-rich grasslands to be most attractive to people, while grass-dominated and defoliated swards were not favoured. Our results are in line with previous work showing colourful flowering grasslands and similar grassy habitats to be highly attractive to humans in rural (e.g., Junge et al., 2009; Lindemann-Matthies et al., 2010; Nowak-Olejnik et al., 2020) and urban settings (e.g., Lindemann-Matthies and Brieger, 2016; Southon et al., 2017; Hoyle et al., 2018). Our study extends these findings by adding a more mechanistic land-use and biodiversity context (see also Richter et al., 2024). People's appreciation was clearly linked to the level of management intensity, which strongly determined the perceived attractiveness and, in addition, generally drives the level of biodiversity found in agricultural grasslands. Extensively managed grasslands can harbour not only high numbers of potentially flowering forbs but also a high diversity of other plants and animals, especially when managed at low intensity for decades (Isselstein et al., 2005; Klaus et al., 2023). In contrast, intensively managed and fertilised grasslands, especially when grazed early in the season and dominated by grasses and plants with high forage value, are usually low in biodiversity of many taxa (Allan et al., 2014). Moreover, intensification generally decreases flower colour diversity and usually leads to early defoliation inhibiting a long and diverse flowering aspect (Binkenstein et al., 2013; Johansen et al., 2019). Though there can potentially also be many flowers in such stands, they are usually dominated by single species indicative of nutrient rich conditions and disturbance, such as dandelion (*Taraxcum* spec., yellow flowering aspect; Martinková et al., 2009) or tall growing Apiaceae (e.g., *Heracleum sphondylium*, *Anthriscus sylvestris*, white flowering aspect; Sheppard, 1991). This dominance of one or few highly similar species potentially causing a vast but monochromic flowering aspect is likely to be the factor explaining why at high levels of flower cover people's aesthetic ratings stagnated or even tended to decrease (Figure 4a).

We specifically found signs of disturbance, such as short grass and bare soil, to be perceived as

unattractive. This is in line with Junge et al. (2015) that found early stages (March) and post-harvest

stages (summer) with a lot of bare soil, short grass and a brownish appearance to be strongly disliked

by people. The same study also found that extensively managed grasslands were preferred over

intensively managed ones (Junge et al., 2015). While grazing intensity was strongly negatively related

to perceived aesthetic quality, cutting frequency was not such a strong predictor. This result should be

treated with caution since in our study grasslands with signs of defoliation and bare soil were all

intensively grazed grasslands, while cut grasslands (meadows) were not yet harvested. Thus, the

results likely depend on the seasonality of the images presented and could change if these were taken

a bit later (end of May), when intensive meadows would also be cut, revealing mostly unattractive

short grass and bare ground (Junge et al., 2015).

Differences in the aesthetic ratings of plant communities as derived from different professional groups

were found but did not compromise the general patterns described before. Yet, appreciation was to

some degree also linked to professional expertise. For example, nature conservationists rated

unfertilised grasslands more positively than agricultural professionals, most likely because

conservationists acknowledge the biodiversity value in such unfertilised stands and relate less to

agricultural productivity (Peter et al., 2022). In addition, agricultural professionals were less negative

about grasslands with intensive management than all other groups, potentially because productive

grasslands are still a main pillar of grass-based food production in Europe (Isselstein et al., 2005). On a

different spatial scale, a similar situation was found by Junge et al. (2011), with farmers rating the

aesthetic quality of a more intensively used (grassland) landscape higher than non-farmers. Potentially

surprisingly, when an agriculturally relevant weed species was visible in the image, this was slightly but

significantly positively related to the aesthetic ratings of people belonging to ecology/nature

conservation and also other persons. This was not the case for agricultural professionals. The reason

behind the positive relation between weeds and aesthetics might be that, in addition to their general

value for general biodiversity and pollinators, some agricultural weed species can show intensive and

attractive flowering aspects (e.g., species of Rhinanthus, i.e., rattles, and Cirsium, i.e., thistles; Balfour

and Ratnieks, 2022). This is underlined by the genus of *Cirsium* being among the most frequently

photographed plant genera in the second part of this study (Figure 7). These results highlight the need

to protect colourful flowering extensively managed grasslands in recreational landscapes.

Our study underlines the value of biodiverse grasslands for human-nature interactions. Similar to

Lindemann-Matthies et al. (2010) and Nowak-Olejnik et al. (2020), we found plant diversity to be

directly related to people's aesthetic ratings and, consequently, to the CES provided. While in the urban

context, plant diversity and flower (colour) richness can be decoupled from native biodiversity by using

seed mixtures of (partially) non-native species (Jiang and Yuan, 2017; Hoyle et al., 2017, 2018), this is

not well possible in agriculturally managed permanent grasslands. At least in the atlantic, alpine and

mediterranean biogeographical regions of Europe, the latter are clearly dominated by native species

(Axmanová et al., 2021), of which many became to some degree rare or endangered. Thus, in the rural

context, the aesthetic appreciation clearly favours extensively managed semi-natural grasslands with

high forb richness and, naturally related, high biodiversity. This calls for a protection and, in face of the

EU´s Nature Restoration Law (Regulation 2024/1991), restoration of forb-rich semi-natural grasslands,

especially in areas that are accessible to people.

4.1.2. Spectral characteristics related to people's aesthetic perception (Hypothesis 1.2)

We applied several standardised methods to automatically extract information from images, including

i) automatic flower cover and colour richness classification, ii) pixel class diversity indices, and iii)

spectral diversity metrics, and we evaluated their capability in predicting people's aesthetic

appreciations. These methods worked well, showing a high potential for practical application in similar

cases. Notably, our case study revealed an interplay between flower cover and flower colour richness

in predicting aesthetic ratings, which accounts for the non-linear relationship observed between

flower cover and appreciation. The analysis of the Shannon and the Simpson indices, based on the

frequency distribution of the pixel classes, with aesthetic ratings did not yield additional insight, since

both were strongly correlated with flower cover. As mentioned before, these correlations can be

attributed to the dominance of few flowering species so that increased flower cover led to a more

even distribution of pixels among various classes (i.e., higher Shannon and Simpson indices).

Conversely, low flower cover means most pixels were classified as "green vegetation or soil," leading

to an uneven class distribution (i.e., lower Shannon and Simpson indices).

The spectral diversity metrics were positively but only weakly correlated with flower cover, as in

Perrone et al. (2024), and not significantly related to aesthetic ratings of the grassland plant

communities. This is probably because spectral diversity is predominantly influenced by light

conditions (Arroyo-Mora et al., 2021), grassland vertical structure (Conti et al., 2021), the presence of

dead biomass (litter, Rossi et al., 2022), and differences in the colours of vegetative organs (Binkenstein

et al., 2013). Notably, flowers constituted less than 1% of pixels in over half of the images analysed.

This finding suggests that spectral diversity is no suitable predictor of grassland aesthetic ratings and

cannot be used to ease and standardise the assessment of vegetation aesthetics, falsifying our second

hypothesis. However, the RF model's high accuracy, along with the strong correlation between

automated and manual estimates of flower cover and flower colour richness, confirmed the reliability

of the classification of grassland images. This ability to rapidly classify many grassland images creates

new possibilities for ecological research, allowing for more extensive and frequent site monitoring

(Andreatta et al., 2023; Torresani et al., 2023).

4.2. What people like to photograph in grasslands

4.2.1. Frequent objects in social media derived grassland images (Hypothesis 2.1)

Our results show that flowers are the most frequently photographed objects in grasslands, reflecting

their cultural significance. This was consistent with our third hypothesis and reflects people's deep

connection with flowering plants, which elicit positive emotional responses (Havinga et al., 2024). The

vibrant colours and visual appeal of flowers, especially in contrast to the dominant green of grasslands,

naturally draw attention and contribute to the attractiveness of a habitat (Mou et al., 2023). Flowers

are more abundant and easier to photograph than birds or insects, as they are static and pose no

perceived threat (Austen et al., 2021). Though less photographed, birds and insects still captivate

visitors, highlighting both their ecological importance and the enjoyment people find in spotting them.

In addition to flowers, the presence of different kinds of livestock was appreciated by people,

confirming previous research on people's preferences in agricultural landscapes using surveys (Junge

et al., 2015) or social media (Chai-allah et al., accepted). These studies found cattle to be valued by

people assessing landscape aesthetics. However, the fact that they were photographed exclusively on

Flickr versus iNaturalist highlights that people's interests and motivations often influence what they

share online during outdoor experiences (Di Cecco et al., 2021). It is likely that Flickr users are more

inclined to share photographs of landscapes and aesthetic appeal symbolising rural identity, such as

cows in a field, whereas iNaturalist users are more inclined to share close-up images of specific species

lacking contextual context of the landscape (Lopez et al., 2020). However, beside livestock, the objects

more frequently included in the top 20 in both platforms were highly similar, indicating that availability

(abundance) and commonness of the objects also played a major role in determining their frequency

on the images, a consistent pattern also in previous research (Havinga et al., 2023).

4.2.2. Differences in wildlife taxa photographed by common visitors and naturalists (Hypothesis 2.2)

Despite the general patterns and similarities found before, our analysis also revealed differences

between users of the two platforms, confirming our fourth hypothesis. These differences in occurrence

and abundance ranking imply that commonness is not the only factor driving the frequency of what is

photographed. While commonness is clearly a driver of how often an object is photographed and

uploaded to a social media platform (Havinga et al., 2023), some of the taxa photographed here are

not very common or even particularly rare such as orchid species (Orchidaceae in families and genera

of *Orchis*, *Ophrys*; *Dactylorhiza*, and *Anacamptis*; Djordjević and Tsiftsis, 2022). This could be confirmed

via a comparison with the mean cover as well as occurrence of families and genera on the grasslands

used for the image analysis, where Poaceae were much more dominant, and Orchidaceae on the other

hand were much rarer than the pictures uploaded to Flickr and iNaturalist would suggest.

Analysis at the taxonomic levels of family and genus revealed that common visitors (Flickr users) and

naturalists (iNaturalist users) expressed different appreciations for species, which most likely reflects

their main motivations (Havinga et al., 2023). There is an expected bias for plants with colourful flowers

or leaves (August et al., 2020; White et al., 2023). Flickr images include some more generalist taxa that

are easily recognizable or aesthetically pleasing, such as species of *Prunus* (cherries, plums etc.),

Taraxacum (dandelion), and *Papaver* (poppy), but also ducks and herons. These are quite common and

not of a specific ecological interest, at least in terms of nature conservation. In contrast, iNaturalist

images included species that may not be commonly perceived as particularly beautiful, such as species

of *Cirsium* (thistles) and *Carex* (sedges). This may be because iNaturalist users tend to consider the

visual appeal of species together with their ecological significance. Most notably, there is a high

frequency of orchid species on iNaturalist and, with a lower ranking, also on Flickr. These species can

be seen as quite rare and not very easy to (accidentally) find in the study area. They are also of high

relevance for nature conservation, due to their natural rarity and anthropogenic decline (Kull et al.,

2016). Thus, besides a potentially positive effect of positive interactions with nature on people's

ecological standpoint and pro-environmental behaviour (Mackay and Schmitt, 2019), such eco-

touristic activities can also harm populations of rare and endangered species (Ballantyne and Pickering,

2012). These findings also suggest that exploring users of broader social media, such as of Flickr, can

complement citizen science approaches in the attempt to understand how people interact with certain

species from a conservation perspective, and to identify locations of potentially damaging eco-tourism.

The familiae and genera of grassland-related wildlife that was found to be attractive to people, has

two slightly contradicting implications. First, in line with the results obtained for species-rich plant

communities, many of the species included in the attractive wildlife depend on species-rich,

extensively managed grasslands. Examples include plant species of *Gentiana* (gentian) and all orchids

(Djordjević and Tsiftsis, 2022) but also most butterfly species (Öckinger et al., 2006), which depend on

late (or incomplete) cutting or grazing allowing plants and animals to flower and reproduce. Since most

of these species are directly dependent on grassland habitats, they underline the suitability of our

approach to identify typical grasslands-based wildlife that is attractive to people. On the other hand,

some rather common and not at all grassland-typical wildlife was frequently included, especially in

images from Flickr. Examples are *Papaver* (poppy) species, which in the study area only occur in arable

ecosystems, and classical waterfowl (ducks, swans, herons), which likely belong to ponds surrounded

by grassy strips. The images might originate from people using grasslands only as a spot to observe

neighbouring habitat types. Having both rare and common wildlife frequently photographed shows

that what is considered attractive by people can be very diverse and our landscapes therefore need to

accommodate many different interests and values.

While crowdsourced platforms offer fresh insights into how people value biodiversity, they come with

biases that must be acknowledged (August et al., 2020; Di Cecco et al., 2021). The users of these

platforms may overrepresent certain socio-demographic groups, influenced by technology access or

the users targeted by each platform (Ghermandi et al., 2023; Venter et al., 2023). Though these

platforms are often biased towards accessible areas (e.g. nature trails), this fine-scale spatial bias does

not necessarily mean that recorded taxa are unrepresentative of the wider area (Geurts et al., 2023).

Though season colour variation in plants may impact aesthetic perception (Campbell-Arvai et al.,

2024), our study focused on people's appreciation of species without accounting for seasonal changes.

Future research should explore how seasonality, especially the flowering season, affects people's

appreciation of different kinds of nature.

4.3 Joint discussion and implications for landscape management

We identified different grassland characteristics that contribute to an attractive grassland landscape.

Both approaches used in this study revealed the aesthetic value of extensively managed, species-rich

grasslands that contain various flowers and wildlife. We found high plant diversity and many familiae

and genera that belong to such ancient, nutrient-poor grassland ecosystems to be directly linked to

people's aesthetic preferences and human-nature interactions. This calls for protecting and enhancing

these grasslands in regions where CES and ecotourism need to be increased, e.g., in the surroundings

of a new settlement. Agricultural policies can support such aesthetically attractive but economically

not viable agricultural habitats via different pathways, such as with direct payments for (cultural)

ecosystems services (e.g., Farley and Costanza, 2010) and result-based schemes that account for the

abundance of particularly attractive species such as orchids (e.g., Šumrada et al., 2021) and for

extensive grazing with livestock (Matzdorf et al., 2008). However, as specified above, increased

human-nature interaction can also cause issues, for example, when ecotourism gets in conflict with

the protection of rare and endangered species (Ballantyne and Pickering, 2012).

Since flowers appeared to be a particular part of grassland CES, while signs of harvesting were

particularly unattractive, suitable measures to improve the aesthetic quality of intensively used

grassland landscapes are delaying or diversifying harvest dates, and, even easier to implement, leaving

stripes of vegetation with high forb cover uncut. This doesn't even need to compromise agricultural

production, as shown by Ravetto Enri et al. (2017), which developed a rotational grazing system, which,

during the main flowering period, temporarily excludes a parcel from grazing. They achieved enhanced

flower resources without reducing farm-scale production (Ravetto Enri et al., 2017). In addition,

measures of ecological restoration that increase forb (species) richness in de-intensified but species-

poor plant communities can also increase the aesthetic value of a late-cut grassland, highlighting co-

production of CES and nature conservation (Bullock et al., 2021). However, particularly rare species

such as orchids are usually not directly included in such a restoration approach, highlighting the

importance of prioritising the conservation of ancient species-rich grasslands.

Although our study did not compare the relative importance of the two aspects assessed, i.e., plant

communities and objects/wildlife, we assume that the effect is additive, with attractive objects in

attractive plant communities to be the most valued case. Future research might want to explore this

and the CES of other agricultural habitat types in more detail.

5. Conclusions

Our study has shown several biodiversity-related features of agricultural grasslands to be attractive to

people. This knowledge allows us to specifically manage land for CES and human-nature interactions

by supporting extensive management and boosting the ecological restoration of grasslands. The

attractive features we identified can be used as indicators for grassland CES and inform result-based

agri-environment schemes, for example, to financially remunerated farmers for the ecosystem services

provided by their land. Future research can add to this by extending our approach to other land use

types and landscape-scale evaluations. Since the people's perceptions and interests were quite

diverse, favouring both rare and endangered as well as common wildlife, our landscapes need to be

diverse and accommodate a range of different habitats.

Acknowledgements

814 We are grateful to all people involved in the ServiceGrass project, especially Andreas Lüscher, Nina

815 Buchmann, Nadja El Benni, Pierrick Jan as well as all student helpers involved in the study. We further

816 want to thank all farmers, who granted access to their grassland parcels, all people who answer the

817 online questionnaire, and all Flickr and iNaturalist users that unconsciously supported our study.

Disclosure statement

820 The authors report there are no competing interests to declare.

Funding

823 VHK and FJR further thank the Mercator Foundation Switzerland and the Fondation Sur-la-Croix for

824 funding the ServiceGrass project. VHK acknowledges support by the Agroscope *Indicate* program

825 (project IndiGras). NF acknowledges their support from Schmidt Sciences.

Data availability statement

828 Data will be shared upon request. Code will be made available upon acceptance.

References

1. Allan, E., Bossdorf, O., Dormann, C.F., Prati, D., Gossner, M.M., Tscharntke, T., Blüthgen, N.,

Bellach, M., Birkhofer, K., Boch, S., Böhm, S., Börschig, C., Chatzinotas, A., Christ, S., Daniel, R.,

Diekötter, T., Fischer, C., Friedl, T., Glaser, K., Hallmann, C., Hodac, L., Hölzel, N., Jung, K., Klein,

A.M., Klaus, V.H., Kleinebecker, T., Krauss, J., Lange, M., Morris, E.K., Müller, J., Nacke, H., Pašalić,

E., Rillig, M.C., Rothenwöhrer, C., Schall, P., Scherber, C., Schulze, W., Socher, S.A., Steckel, J.,

Steffan-Dewenter, I., Türke, M., Weiner, C.N., Werner, M., Westphal, C., Wolters, V., Wubet, T.,

Gockel, S., Gorke, M., Hemp, A., Renner, S.C., Schöning, I., Pfeiffer, S., König-Ries, B., Buscot, F.,

Linsenmair, K.E., Schulze, E.-D., Weisser, W.W., Fischer, M., 2014. Interannual variation in land-

use intensity enhances grassland multidiversity. Proc. Natl. Acad. Sci. 111 (1), 308–313.

2. Allan, E., Manning, P., Alt, F., Binkenstein, J., Blaser, S., Blüthgen, N., Böhm, S., Grassein,

F., Hölzel, N., Klaus, V.H., Kleinebecker, T., Morris, E.K., Oelmann, Y., Prati, D., Renner, S.C., Rillig,

M.C., Schaefer, M., Schloter, M., Schmitt, B., Schöning, I., Schrumpf, M., Solly, E., Sorkau,

E., Steckel, J., Steffen-Dewenter, I., Stempfhuber, B., Tschapka, M., Weiner, C.N., Weisser,

W.W., Werner, M., Westphal, C., Wilcke, W., Fischer, M., 2015. Land use intensification alters

ecosystem multifunctionality via loss of biodiversity and changes to functional composition.

Ecol. Lett. 18 (8), 834–843.

3.   Andreatta, D., Bachofen, C., Dalponte, M., Klaus, V.H., Buchmann, N., 2023. Extracting flowering

phenology from grassland species mixtures using time-lapse cameras. Remote Sens. Environ.

298, 113835.

4.   Arroyo-Mora, J.P., Kalacska, M., Løke, T., Schläpfer, D., Coops, N.C., Lucanus, O., Leblanc, G.,

2021. Assessing the impact of illumination on UAV pushbroom hyperspectral imagery collected

under various cloud cover conditions. Remote Sens. Environ. 258.

5.   August, T. A., Pescott, O. L., Joly, A., & Bonnet, P., 2020. AI naturalists might hold the key to

unlocking biodiversity data in social media imagery. Patterns. 1, 100116

6.   Austen, G.E., Dallimer, M., Irvine, K.N., Maund, P.R., Fish, R.D., Davies, Z.G., 2021. Exploring

shared public perspectives on biodiversity attributes. People Nat. *3* (4), 901–913.

7.   Axmanová, I., Kalusová, V., Danihelka, J., Dengler, J., Pergl, J., Pyšek, P., Večeřa, M., Attorre,

F., Biurrun, I., Boch, S., Conradi, T., Gavilán, R.G., Jiménez-Alfaro, B., Knollová, I., Kuzemko,

A., Lenoir, J., Leostrin, A., Medvecká, J., Moeslund, J.E., Obratov-Petkovic, D., Svenning, J.-

C., Tsiripidis, I., Vassilev, K., Chytrý, M., 2021. Neophyte invasions in European grasslands. J. Veg.

Sci. 32 (2), e12994.

8.   Balfour, N.J., Ratnieks, F.L., 2022. The disproportionate value of 'weeds' to pollinators and

biodiversity. J. Appl. Ecol. 59 (5), 1209–1218.

9.   Ballantyne, M., Pickering, C., 2012. Ecotourism as a threatening process for wild orchids. J.

Ecotourism 11 (1), 34–47.

10. Bardgett, R.D., Bullock, J.M., Lavorel, S., Manning, P., Schaffner, U., Ostle, N., Chomel, M.,

Durigan, G., Fry, E, L., Johnson, D., Lavallee, J.M., Le Provost, G., Luo, S., Png, K., Sankaran, M.,

Hou, X., Zhou, H., Ma, L., Ren, W., Li, X., Ding, Y., Li, Y., Shi, H., 2021. Combatting global grassland

degradation. Nat. Rev. Earth Environ. 2, 720–735.

11. Barve, V., Hart, E., 2017. rinat: Access iNaturalist Data Through APIs. R package. https://cran.r-

project.org/package=rinat

12. Bengtsson, J., Bullock, J.M., Egoh, B., Everson, C., Everson, T., O'Connor, T., O'Farrell, P.J., Smith,

H.G., Lindborg, R., 2019. Grasslands—more important for ecosystem services than you might

think. Ecosphere 10 (2), e02582.

13. Binkenstein, J., Renoult, J.P., Schaefer, H.M., 2013. Increasing land-use intensity decreases floral

colour diversity of plant communities in temperate grasslands. Oecologia 173 (2), 461–471.

14. Blüthgen, N., Dormann, C.F., Prati, D., Klaus, V.H., Kleinebecker, T., Hölzel, N., Alt, F., Boch, S.,

Gockel, S., Hemp, A., Müller, J., Nieschulze, J., Renner, S.C., Schöning, I., Schumacher, U., Socher,

S.A., Wells, K., Birkenhofer, K., Buscot, F., Oelmann, Y., Rothenwöhrer, C., Scherber, C.,

Tscharntke, T., Weiner, C.N., Fischer, M., Kalko, E.K.V., Linsenmair, K.E., Schulze, E.-D., Weisser,

W.W., 2012. A quantitative index of land-use intensity in grasslands: Integrating mowing, grazing

and fertilization. Basic Appl. Ecol. 13 (3), 207–220.

15. Briemle, G., Nitsche, S., Nitsche, L., 2002. Nutzungswertzahlen für Gefäßpflanzen des

Grünlandes. Schriftenr. Vegetationskd. 38 (2), 203–225.

16. Bullock, J.M., McCracken, M.E., Bowes, M.J., Chapman, R.E., Graves, A.R., Hinsley, S.A., Hutchins,

M.G., Nowakowski, M., Nicholls, D.J.E., Oakley, S., Old, G.H., Ostle, N.J., Redhead, J.W.,

Woodcock, B.A., Bedwell, T., Mayes, S., Robinson, V.S., Pywell, R.F., 2021. Does agri-

environmental management enhance biodiversity and multiple ecosystem services?: A farm-

scale experiment. Agric. Ecosyst. Environ. 320, 107582.

17. Campbell-Arvai, V., Serrano Vergel, R., Lindquist, M., Fox, N., Van Berkel, D., 2024. Tree selection

for a virtual urban park: Comparing aided and unaided decision-making to support public

engagement in greenspace design. Urban For. Urban Green. 99, 128447.

18. Chai-allah, A., Fox, N., Brunschwig, G., Bimonte, S., Joly, F., 2023. A trail-based approach using

crowdsourced data to assess recreationists' preferences for landscape. Landsc. Urban Plan. 233,

104700.

19. Chai-allah, A., Hermes, J., Zander, V., Joly, F., Brunschwig, G., Bimonte, S., Fox, N., in press.

Assessing recreationists' preferences of the landscape and species using crowdsourced images

and machine learning. Landsc. Urban Plan. Accepted and in press

20. Chamberlain, E. and Szocs, E., 2013. taxize - taxonomic search and retrieval in R. F1000Research,

2:191. http://f1000research.com/articles/2-191/v2.

21. Cheng, X., Van Damme, S., Li, L., Uyttenhove, P., 2019. Evaluation of cultural ecosystem services:

a review of methods. Ecosyst. Serv. 37, 100925. https://doi.org/10.1016/j. ecoser.2019.100925.

22. Congalton, R.G., Green, K., 2009. Assessing the Accuracy of Remotely Sensed Data: Principles

and Practices (2nd ed.). CRC Press.

23. Conti, L., Malavasi, M., Galland, T., Komárek, J., Lagner, O., Carmona, C.P., de Bello, F., Rocchini,

D., Šímová, P., 2021. The relationship between species and spectral diversity in grassland

communities is mediated by their vertical complexity. Appl. Veg. Sci. 24 (3).

24. Di Cecco, G.J., Barve, V., Belitz, M.W., Stucky, B.J., Guralnick, R.P., Hurlbert, A.H., 2021. Observing

the observers: how participants contribute data to iNaturalist and implications for biodiversity

science. Bioscience. 71, 1179–1188.

25. Djordjević, V., Tsiftsis, S., 2022. The role of ecological factors in distribution and abundance of

terrestrial orchids. *Orchids phytochemistry, biology and horticulture: fundamentals and*

applications 3–72. Springer International Publishing.

26. European Union, 2018. Copernicus DEM - Global and European Digital Elevation Model: EEA-10:

European coverage. Copernicus Land Monitoring Service, European Environment Agency (EEA)

https://doi.org/10.5270/ESA-c5d3d65

27. Farley, J., Costanza, R., 2010. Payments for ecosystem services: from local to global. Ecol. Econ.

69 (11), 2060-2068.

28. Fischer, L.K., Honold, J., Cvejić, R., Delshammar, T., Hilbert, S., Lafortezza, R., Nastran, M.,

Nielsen, A.B., Pintar, M., van der Jagt, A.P.N., Kowarik, I., 2018. Beyond green: Broad support for

biodiversity in multicultural European cities. Glob. Environ. Change 49, 35–45.

29. Fox, N., August, T., Mancini, F., Parks, K.E., Eigenbrod, F., Bullock, J. M., Sutter, L., Graham, L.J.,

2020. "photosearcher" package in R: An accessible and reproducible method for harvesting large

datasets from Flickr. SoftwareX, 12, 100624.

30. Fox, N., August, T., Mancini, F., Parks, K.E., Eigenbrod, F., Bullock, J.M., Sutter, L., Graham, L.J.,

2021. Enriching social media data allows a more robust representation of cultural ecosystem

services. Ecosyst. Serv. 50, 101328.

31. Fox, N., Chamberlain, B., Lindquist, M., Van Berkel, D., 2022a. Understanding landscape

aesthetics using a novel viewshed assessment of social media locations within the Troodos

UNESCO Global Geopark, Cyprus. Front. Environ. Sci. 10, 884115.

32. Fox, N., Graham, L.J., Eigenbrod, F., Bullock, J.M., Parks, K.E., 2022b. Geodiversity supports

cultural ecosystem services: an assessment using social media. Geoheritage 14 (1), 27.

33. Geurts, E.M., Reynolds, J.D., Starzomski, B.M., 2023. Turning observations into biodiversity data:

Broadscale spatial biases in community science. Ecosphere 14, e4582.

34. Ghermandi, A., Langemeyer, J., Van Berkel, D., Calcagni, F., Depietri, Y., Vigl, L.E., Fox, N.,

Havinga, I., Jäger, H., Kaiser, N., Karasov, O., McPhearson, T., Podschun, S., Ruiz-Frau, A., Sinclair,

M., Venohr, M., Wood, S.A., 2023. Social media data for environmental sustainability: A critical

review of opportunities, threats, and ethical use. One Earth 6 (3), 236-250.

35. Gosal, A.S., Geijzendorffer, I.R., V´aclavík, T., Poulin, B., Ziv, G., 2019. Using social media, machine

learning and natural language processing to map multiple recreational beneficiaries. Ecosyst.

Serv. 38, 100958.

36. Hauser, L.T., Timmermans, J., van der Windt, N., Sil, Â.F., César de Sá, N., Soudzilovskaia, N.A.,

van Bodegom, P.M., 2021. Explaining discrepancies between spectral and in-situ plant diversity

in multispectral satellite earth observation. Remote Sens. Environ. 265, 112684.

37. Havinga, I., Bogaart, P.W., Hein, L., Tuia, D., 2020. Defining and spatially modelling cultural

ecosystem services using crowdsourced data. Ecosyst. Serv. 43, 101091.

38. Havinga, I., Marcos, D., Bogaart, P., Massimino, D., Hein, L., Tuia, D., 2023. Social media and deep

learning reveal specific cultural preferences for biodiversity. People Nat. 5(3), 981–998.

39. Havinga, I., Marcos, D., Bogaart, P., Tuia, D., Hein, L., 2024. Understanding the sentiment

associated with cultural ecosystem services using images and text from social media. Ecosyst.

Serv. 65, 101581.

40. Hijmans, R.J., 2023. raster: Geographic analysis and modeling with raster data. R package version

3.6-26. https://CRAN.R-project.org/package=raster

41. Hoyle, H., Hitchmough, J., Jorgensen, A., 2017. All about the 'wow factor'? The relationships

between aesthetics, restorative effect and perceived biodiversity in designed urban planting.

Landsc. Urban Plan. 164, 109–123.

42. Hoyle, H., Norton, B., Dunnett, N., Richards, J.P., Russell, J.M., Warren, P., 2018. Plant species or

flower colour diversity? Identifying the drivers of public and invertebrate response to designed

annual meadows. Landsc. Urban Plan. 180, 103–113.

43. Huber, N., Ginzler, C., Pazúr, R., Descombes, P., Baltensweiler, A., Ecker, K., Meier, E., Price, B.,

2022. Distribution maps of permanent grassland habitats for Switzerland. EnviDat.

https://www.doi.org/10.16904/envidat.341.

44. Isselstein, J., Jeangros, B., Pavlu, V., 2005. Agronomic aspects of biodiversity targeted

management of temperate grasslands in Europe–a review. Agron. Res. 3 (2), 139–151.

45. Jiang, Y., Yuan, T., 2017. Public perceptions and preferences for wildflower meadows in Beijing,

China. Urban For. Urban Green. 27, 324–331.

46. Johansen, L., Westin, A., Wehn, S., Iuga, A., Ivascu, C.M., Kallioniemi, E., Lennartsson, T., 2019.

Traditional semi-natural grassland management with heterogeneous mowing times enhances

flower resources for pollinators in agricultural landscapes. Global Ecol. Conserv. 18, e00619.

47. Junge, X., Jacot, K.A., Bosshard, A., Lindemann-Matthies, P., 2009. Swiss people's attitudes

towards field margins for biodiversity conservation. J. Nat. Conserv. 17 (3), 150–159.

48. Junge, X., Lindemann-Matthies, P., Hunziker, M., Schüpbach, B., 2011. Aesthetic preferences of non-farmers and farmers for different land-use types and proportions of ecological compensation areas in the Swiss lowlands. Biol. Conserv. 144 (5), 1430–1440.

49. Junge, X., Schüpbach, B., Walter, T., Schmid, B., Lindemann-Matthies, P., 2015. Aesthetic quality of agricultural landscape elements in different seasonal stages in Switzerland. Landsc. Urban Plan. 133, 67–77.

50. Klaus, V.H., Jehle, A., Richter, F., Buchmann, N., Knop, E., Lüscher, G., 2023. Additive effects of two agri-environmental schemes on plant diversity but not on productivity indicators in permanent grasslands in Switzerland. J. Environ. Manage. 348, 119416.

51. Kosanic, A., Petzold, J., 2020. A systematic review of cultural ecosystem services and human wellbeing. Ecosyst. Serv. 45, 101168.

52. Kull, T., Selgis, U., Peciña, M.V., Metsare, M., Ilves, A., Tali, K., Sepp, K., Kull, K., Shefferson, R.P., 2016. Factors influencing IUCN threat levels to orchids across Europe on the basis of national red lists. Ecol. Evol. 6 (17), 6245–6265.

53. Lee, H., Seo, B., Cord, A.F., Volk, M., Lautenbach, S., 2022. Using crowdsourced images to study selected cultural ecosystem services and their relationships with species richness and carbon sequestration. Ecosyst. Serv. 54, 101411.

54. Liaw, A., Wiener, M., 2002. Classification and Regression by randomForest. R News, 2 (3), 1822.

55. Lindemann-Matthies, P., Junge, X., Matthies, D., 2010. The influence of plant diversity on people's perception and aesthetic appreciation of grassland vegetation. Biol. Conserv. 143 (1), 195–202.

56. Lindemann-Matthies, P., Brieger, H., 2016. Does urban gardening increase aesthetic quality of urban areas? A case study from Germany. Urban For. Urban Green. 17, 33–41.

57. Lopez, B., Minor, E., Crooks, A., 2020. Insights into human-wildlife interactions in cities from bird sightings recorded online. Landsc. Urban Plan. 196, 103742.

58. Lüscher, A., Grieder, C., Huguenin-Elie, O., Klaus, V., Reidy, B., Schneider, M.K., Schubiger, F.,

Suter, D., Suter, M., Kölliker, R., 2019. Grassland systems in Switzerland with a main focus on

sown grasslands. Grassland Sci. Eur. 24, 3–16.

59. Mackay, C.M., Schmitt, M.T., 2019. Do people who feel connected to nature do more to protect

it? A meta-analysis. J. Environ. Psychol. 65, 101323.

60. Martinková, Z., Honek, A., Pekár, S., 2009. Seed availability and gap size influence seedling

emergence of dandelion (Taraxacum officinale) in grasslands. Grass Forage Sci. 64 (2), 160–168.

61. Matzdorf, B., Kaiser, T., Rohner, M.S., 2008. Developing biodiversity indicator to design efficient

agri-environmental schemes for extensively used grassland. Ecol. Indic. 8 (3), 256–269.

62. Mou, N., Wang, J., Zheng, Y., Zhang, L., Makkonen, T., Yang, T., Niu, J., 2023. Flowers as

attractions in urban parks: Evidence from social media data. Urban For. Urban Green. 82,

127874.

63. Nowak-Olejnik, A., Mocior, E., Hibner, J., Tokarczyk, N., 2020. Human perceptions of cultural

ecosystem services of semi-natural grasslands: The influence of plant communities. Ecosyst.

Serv. 46, 101208.

64. Öckinger, E., Eriksson, A.K., Smith, H.G., 2006. Effects of grassland abandonment, restoration

and management on butterflies and vascular plants. Biol. Conserv. 133 (3), 291–300.

65. Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'hara, R.B., Simpson, G.L.,

Solymos, P., Stevens, M.H.H., Wagner, H., 2022. 'vegan', Community Ecology Package.

https://CRAN.R-project.org/package=vegan

66. Perrone, M., Conti, L., Galland, T., Komárek, J., Lagner, O., Torresani, M., Rossi, C., Carmona, C.

P., de Bello, F., Rocchini, D., Moudrý, V., Šímová, P., Bagella, S., Malavasi, M., 2024. "Flower

power": How flowering affects spectral diversity metrics and their relationship with plant

diversity. Ecol. Inform. 81, 102589.

67. Peter, S., Le Provost, G., Mehring, M., Müller, T., Manning, P., 2022. Cultural worldviews

consistently explain bundles of ecosystem service prioritisation across rural Germany. People

Nat. 4 (1), 218–230.

68. Plieninger, T., Dijks, S., Oteros-Rozas, E., Bieling, C., 2013. Assessing, mapping, and quantifying

cultural ecosystem services at community level. Land Use Policy 33, 118–129.

69. Ravetto Enri, S., Probo, M., Farruggia, A., Lanore, L., Blanchetete, A., Dumont, B., 2017. A

biodiversity-friendly rotational grazing system enhancing flower-visiting insect assemblages

while maintaining animal and grassland productivity. Agric. Ecosyst. Environ. 241, 1–10.

70. R Core Team, 2024. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/

71. Richter, F., Jan, P., El Benni, N., Lüscher, A., Buchmann, N., Klaus, V.H., 2021. A guide to assess

and value ecosystem services of grasslands. Ecosyst. Serv. 52, 101376.

72. Richter, F., Suter, M., Lüscher, A., Buchmann, N., El-Benni, N., Feola-Conz, R., Hartmann, M., Jan,

P., Klaus, V.H., 2024. Effects of management practices on the ecosystem service

multifunctionality of temperate grasslands. Nat. Commun. 15, 3829.

73. Ridding, L.E., Redhead, J.W., Oliver, T.H., Schmucki, R., McGinlay, J., Graves, A.R., Morris, J.,

Bradbury, R.B., King, H., Bullock, J.M., 2018. The importance of landscape characteristics for the

delivery of cultural ecosystem services. J. Environ. Manage. 206, 1145–1154.

74. Rocchini, D., Boyd, D.S., Féret, J.B., Foody, G.M., He, K.S., Lausch, A., Nagendra, H., Wegmann,

M., Pettorelli, N., 2016. Satellite remote sensing to monitor species diversity: potential and

pitfalls. Remote Sens. Ecol. Conserv. 2 (1), 25–36.

75. Rocchini, D., Thouverai, E., Marcantonio, M., Iannacito, M., Da Re, D., Torresani, M., Bacaro, G.,

Bazzichetto, M., Bernardi, A., Foody, G. M., Furrer, R., Kleijn, D., Larsen, S., Lenoir, J., Malavasi,

M., Marchetto, E., Messori, F., Montaghi, A., Moudrý, V., Naimi, B., Ricotta, C., Rossini, M., Santi,

F., Santos, M.J., Schaepman, M.E., Schneider, F.D., Schuh, L., Silvestri, S., Šímová, P., Skidmore,

A.K., Tattoni, C., Tordoni, E., Vicario, S., Zannini, P., Wegmann, M., 2021. rasterdiv—An

Information Theory tailored R package for measuring ecosystem heterogeneity from space: To

the origin and back. Methods Ecol. Evol. 12 (6), 1093–1102.

76. Rossi, C., Kneubühler, M., Schütz, M., Schaepman, M.E., Haller, R.M., Risch, A.C., 2022. Spatial

resolution, spectral metrics and biomass are key aspects in estimating plant species richness

from spectral diversity in species-rich grasslands. Remote Sens. Ecol. Conserv. 8 (3), 297–314.

77. Schils, R.L.M., Bufe, C., Rhymer, C.M., Francksen, R.M., Klaus, V.H., Abdalla, M., Milazzo, F., Lellei-

Kovács, E., ten Berge, H., Bertora, C., Chodkiewicz, A., Dămătîrcă, C., Feigenwinter, I., Fernández-

Rebollo, P., Ghiasi, S., Hejduk, S., Hiron, M., Janicka, M., Pellaton, R., Smith, K.E., Thorman, R.,

Vanwalleghem, T., Williams, J., Zavattaro, L., Kampen, J., Derkx, R., Smith, P., Whittingham, M.J.,

Buchmann, N., Price, J.P.N., 2022. Permanent grasslands in Europe: Land use change and

intensification decrease their multifunctionality. Agric. Ecosyst. Environ. 330, 107891.

78. Schirpke, U., Ghermandi, A., Sinclair, M., Van Berkel, D., Fox, N., Vargas, L., Willemen, L., 2023.

Emerging technologies for assessing ecosystem services: A synthesis of opportunities and

challenges. Ecosyst. Serv. 63, 101558.

79. Schwemmer, C., 2019. imgrec: An Interface for Image Recognition. R package version 0.1, 1.

https://github.com/cschwem2er/imgrec

80. Sheppard, A.W., 1991. Heracleum sphondylium L. J. Ecol. 79 (1), 235–258.

81. Southon, G.E., Jorgensen, A., Dunnett, N., Hoyle, H., Evans, K.L., 2017. Biodiverse perennial

meadows have aesthetic value and increase residents' perceptions of site quality in urban green-

space. Landsc. Urban Plan. 158, 105–118.

82. Stevens, S., Wu, J., Thompson, M.J., Campolongo, E.G., Song, C.H., Carlyn, D.E., Dong, L., Dahdul,

W.M., Stewart, C., Berger-Wolf, T., Chao, W.-L., Su, Y., 2024. BioCLIP: A Vision Foundation Model

for the Tree of Life. Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern

Recognition, 19412–19424.

83. Straffelini, E., Luo, J., Tarolli, P., 2024. Climate change is threatening mountain grasslands and

their cultural ecosystem services. Catena 237, 107802.

84. Šumrada, T., Vreš, B., Čelik, T., Šilc, U., Rac, I., Udovč, A., Erjavec, E., 2021. Are result-based

schemes a superior approach to the conservation of High Nature Value grasslands? Evidence

from Slovenia. Land Use Policy, 111, 105749.

85. Swisstopo (Eds.), 2020. The high precision digital elevation model of Switzerland swissALTI3D

(2m).   Swiss   Federal   Office   of   Topography.

www.swisstopo.admin.ch/en/geodata/height/alti3d.html#technische_details

86. Torresani, M., Kleijn, D., de Vries, J.P.R., Bartholomeus, H., Chieffallo, L., Cazzolla Gatti, R.,

Moudrý, V., da Re, D., Tomelleri, E., Rocchini, D., 2023. A novel approach for surveying flowers

as a proxy for bee pollinators using drone images. Ecol. Indic. 149, 110123.

87. Torresani, M., Rossi, C., Perrone, M., Hauser, L.T., Féret, J.-B., Moudrý, V., Simova, P., Ricotta, C.,

Foody, G.M., Kacic, P., Feilhauer, H., Malavasi, M., Tognetti, R., Rocchini, D., 2024. Reviewing the

Spectral Variation Hypothesis: Twenty years in the tumultuous sea of biodiversity estimation by

remote sensing. Ecol. Inform. 82, 102702.

88. Van Berkel, D.B., Verburg, P.H., 2014. Spatial quantification and valuation of cultural ecosystem

services in an agricultural landscape. Ecol. Indic. 37, 163–174.

89. Wang, R., Gamon, J.A., 2019. Remote sensing of terrestrial plant biodiversity. Remote Sens.

Environ. 231.

90. Wang, L., Huang, L., Cao, W., Zhai, J., Fan, J., 2024. Assessing grassland cultural ecosystem

services supply and demand for promoting the sustainable realization of grassland cultural

values. Sci. Total Environ. 912, 169255.

91. Wickham, H., 2016. Ggplot2: Elegant graphics for data analysis (2nd ed.). Springer International

Publishing.

92. Wilkins, E.J., Van Berkel, D., Zhang, H., Dorning, M.A., Beck, S.M., Smith, J.W., 2022. Promises

and pitfalls of using computer vision to make inferences about landscape preferences: Evidence

from an urban-proximate park system. Landsc. Urban Plan. 219, 104315.

93. Zielstra, D., Hochmair, H.H., 2013. Positional accuracy analysis of Flickr and Panoramio images

for selected world regions. J. Spat. Sci. 58, 251–273.

1100

1101   Please note: For the purpose of easing the peer review, we include the supplementary material in the

1102   main file.

1103

Supplementary material 1

1104

Questionnaire text

1105

Questionnaire text translated to English (see below for the original German-language questionnaire

and all pictures included in this study):

Introductory text

"With this questionnaire we would like to understand what characteristics of grassland are

considered beautiful in the sense of contributing to an attractive landscape. Therefore, we would like

you to rate the following ten photos of grasslands in terms of their perceived beauty. Please try to

focus only on the plant communities and not on the technical quality or perspective of the photos.

Thank you very much for your support of our research!

The results of this survey will be used anonymously for the ServiceGrass research project by ETH

Zurich and Agroscope to investigate socially relevant ecosystem services. Ecosystem services are

benefits grassland provides to humans, e.g. it stores carbon, prevents erosion, provides habitat for

animals ... and can look beautiful!

By completing this survey, you agree that your answers may be used anonymously for research in the

ServiceGrass project of ETH Zurich and Agroscope."

1120

1121 *Start of questionnaire*

1122 "Now the photos will follow. Please rate the grasslands freely according to your gut feeling and

1123 personal aesthetic perception.

1124 How attractive do you find this grassland? ("not beautiful" to "very beautiful" as a five-point Likert-

1125 scale question; series of ten pictures)"

1126

End of questionnaire

"Thank you for your evaluation of the grasslands! Now just a few brief personal details:

Which country do you live in? (Switzerland, Germany, Austria, other)

What is your age? (in steps of 10 years)

Which gender do you consider yourself to be? (diverse, female, male)

Is your professional activity related to the following fields? (agriculture, ecology or conservation,

none of these/other/not working)"

1134

1135 **Example of Questionnaire in German language**

1136 Submitted as separate file

1137

1138 **All pictures used for the questionnaire**

1139 Submitted as separate file

1140

1141

1142 **Figure S1**. Distribution of age classes of survey participants (n = 522)

1143

1144

1145 **Figure S2**. Aesthetic ratings of 92 grassland images by agricultural and ecological professionals. Blue

1146 squares depict fertilised and intensively managed grasslands, and green triangles depict extensive

1147 grasslands in which no fertilisation takes place. The black line represents the 1:1-line.

1148

1149 **Figure S3**. Relationships between automatically and manually estimated flower cover and flower

1150 colour richness. Pearson Correlation Coefficients and p-values are reported in the top-left corner of

1151 each panel. A small amount of jittering was applied to reduce overlap.

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

Figure S4. Mean cover of different plant families per grassland (left) and number of grasslands in which different plant families occurred (right) for the 92 grasslands from which the images for the image analysis were taken.

1169
1170
1171

1172

1173

1174

1175

1176

1177

1178

1179

1180   **Figure S5.** Mean cover of different plant genera per grassland (left) and number of grasslands in
1181   which different plant genera occurred (right) for the 92 grasslands from which the images for the
1182   image analysis were taken. Only genera with a cover > 0.1% and genera occurring in > 10 grasslands
1183   are shown, respectively.

1184

1185 **Table S1**. Basic statistics of responses given for all respondents and separated into three groups

1186 related to professional work. Responses with NA and mixed professions were not analysed

1187 separately (n = 62) but are included in all ratings.

	All ratings (n = 522)	Ratings from agriculture (n = 121)	Ratings from ecology/nature conservation (n = 117)	Ratings from neither of both, other professions, retired (n = 222)
Mean value	3.15	3.13	3.12	3.20
Standard deviation	0.65	0.65	0.82	0.70
Min	1.79	1.58	1.59	1.60
Max	4.54	4.50	4.80	4.57

1188

1189

1190 **Table S2**. Results from ANOVAs relating the aesthetic rating (total and of different professional

1191 groups) to different plant community characteristics visible on the images (manually dummy coded).

1192 Significant levels: "***" = p < 0.001, "**" = p < 0.01, "*" = p < 0.05, "." = p < 0.01 (all other p > 0.01).

	Signs of defoliation (yes vs. no)						Bare soil visible (yes vs. no)					
Ratings	Model p	Model R^2	Estimate	SE	t value	Signif.	Model p	Model R^2	Estimate	SE	t value	Signif.
All	< 0.001	0.33	-1.12	0.17	-6.71	***	< 0.001	0.17	-0.61	0.14	-4.34	***
Agriculture	< 0.001	0.23	-0.93	0.18	-5.21	***	< 0.001	0.28	-0.77	0.13	-5.93	***
Ecology	< 0.001	0.22	-1.17	0.23	-5.15	***	0.021	0.06	-0.44	0.19	-2.35	*
Other	< 0.001	0.36	-1.26	0.18	-7.16	***	< 0.001	0.15	-0.61	0.15	-4.00	***
	Weed prominently visible (yes vs. no)						Vegetation structure (light vs. dense)					
All	0.111	0.03	0.23	0.14	1.61		0.078	0.03	0.25	0.14	1.78	.
Agriculture	0.580	0.00	-0.08	0.15	-0.56		0.956	0.00	-0.01	0.14	-0.06	
Ecology	0.007	0.08	0.49	0.18	2.77	**	< 0.001	0.12	0.61	0.17	3.60	***
Other	0.021	0.06	0.36	0.15	2.34	*	0.138	0.02	0.23	0.15	1.50	
	Litter visible (yes vs. no)											
All	< 0.001	0.14	-0.93	0.24	-3.90	***						
Agriculture	< 0.001	0.20	-1.12	0.23	-4.85	***						
Ecology	0.196	0.02	-0.42	0.32	-1.30							
Other	< 0.001	0.17	-1.11	0.25	-4.36	***						

1193

1194

Supplementary material 2

1196

1197 Table S3. Overview of the labels generated by the feature "label" of the Google Cloud Vision algorithm
1198 for Flickr and iNaturalist photographs and their classification. This classification was adapted from Chai-
1199 allah et al. (submitted). Only labels with frequency > 15 were included.

Category	Sub-category	Labels
Nature	Landscape	aeolian landform, agriculture, arroyo, badlands, batholith, beach, bedrock, biome, body of water, cave, chaparral, coast, coastal and oceanic landforms, creek, crop, deciduous, ecoregion, escarpment, fell, fen, field, fluid, fluvial landforms of streams, forest, freshwater marsh, garden, geological phenomenon, geology, glacial landform, glacier, grassland, hay, headland, highland, hill, hinterland, ice cap, jungle, lacustrine plain, lake, landscape, limestone, loch, logging, marsh, massif, meadow, monolith, mountain, mountain range, mountain river, mountainous landforms, natural environment, natural landscape, natural material, nature, nature reserve, northern hardwood forest, ocean, old-growth forest, outcrop, pasture, pebble, plain, plant community, plateau, pond, prairie, reservoir, ridge, riparian forest, riparian zone, river island, rock, rural area, salt marsh, sand, seep, shore, shrubland, slope, snow, soil, speleothem, spruce-fir forest, stalactite, stalagmite, steppe, stream, summit, tarn, temperate broadleaf and mixed forest, temperate coniferous forest, terrain, thatching, tidal marsh, tropical and subtropical coniferous forests, underwater, valley, water, water feature, water resources, watercourse, waterfall, waterway, wave, wetland, wilderness, woodland
	Objects/species	accipitridae, accipitriformes, acmon blue, adonis blue, agaric, agaricaceae, agaricomycetes, aglais, aglais io, algae, allium, alpaca, american black bear, american black duck, american coot, amphibian, anaxyrus, animal migration, annual plant, ant, antelope, anthriscus, antler, apatura, aphantopus, aquatic plant, arachnid, araneus, argiope, argynnis, aricia, arthropod, bald eagle, band winged grasshoppers, beak, bear, bee, bee balm, beetle, berry, bird, bird migration, bird nest, bird of prey, bison, blossom, blowflies, bolete, boloria, bombyliidae, botany, bovine, branch, brimstones, brown argus, brown bear, bud, bufo, bug, bull, bullfrog, bumblebee, bumblebee orchid, burdock, buttercup, butterfly, buzzard, cabbage butterfly, calf, calidrid, camomile, caterpillar, celastrina, chamaemelum nobile, charadriiformes, cherry blossom, chickadee, chicken, ciconiiformes, claw, cleavers, coenagrion, coenonympha tullia, colias, colias croceus, colias hyale, colias sareptensis, coltsfoot, comb, comma, common blue, condor, conifer, corn poppy, cow-goat family, crane-like bird, creeping thistle, crepis paludosa, cretan crocus, cricket-like insect, crow, crow-like bird, cupido (butterfly), cut flowers, dactylorhiza praetermissa, dairy cow, daisy, daisy family, damselfly, dandelion, deer, distaff thistles, domestic pig, dowitcher, dragon lizard, dragonflies and

| | | damseflies, dragonfly, drosophila, duck, ducks, geese and swans, dung beetle, eagle, eastern tent caterpillar, edible mushroom, egret, elapidae, elk, emberizidae, european marsh thistle, european robin, evergreen, falcon, falconiformes, fawn, feather, felidae, feral goat, fern, ferns and horsetails, fin, finch, fire salamander, fish crow, flightless bird, flock, floribunda, flower, flowering plant, fly orchid, forb, forget-me-not, fowl, fox, french lavender, fritillaria, frog, fruit, fungus, fur, galliformes, garden white, gatekeeper (butterfly), gentiana, goat, goat-antelope, goats, gonepteryx cleopatra, gonepteryx rhamni, grapevine family, grass, grass family, grass snake, grasshopper, grazing, great blue heron, great heron, great spangled fritillary, greater scaup, green-veined white, grizzly bear, ground beetle, groundcover, grove, gull, halictidae, hare, harrier, hawk, hawker dragonflies, heath fritillary, hedge, hemp family, hen-of-the-wood, herb, herbaceous plant, herd, heron, hierochloe, high brown fritillary, hipparchia, holly blue, honeybee, horse, horse flies, house fly, hybrid clover, hybrid tea rose, hypericum, iguania, inonotus, insect, invertebrate, khorasan wheat, kingcup, kit fox, ladybug, larch, large skipper, large tortoiseshell, large white, lari, larva, lasiommata, leaf, leafhopper, leptidea, lesser yellowlegs, livestock, lizard, llama, locust, lotus, lotus family, lycaena, lymantriidae, lymnaeidae, lynx, malus, mammal, mane, maniola, mantidae, mantis, mare, marguerite daisy, marine biology, marine invertebrates, marine mammal, marsupial, mayflies, mazarine blue, meadow brown, medicinal mushroom, megachilidae, melanargia, melanargia galathea, melitaea, membrane-winged insect, mole salamander, morning glory, moth, moths and butterflies, mountain cottontail, mushroom, mustelidae, narcissus, natural foods, neotinea ustulata, nest, net-winged insects, new caledonian crow, nightshade family, non-vascular land plant, northern brown argus, northern leopard frog, noxious weed, nymphalis, nymphalis xanthomelas, ochlodes, ophrys, orange sulphur, orb-weaver spider, orchids of the philippines, organism, oriental poppy, osprey, ostrich fern, owl, ox, oxeye daisy, pack animal, papilio, papilio machaon, pararge, parasite, peafowl, pearl-bordered fritillar, pedicel, pelecaniformes, peppered moth, perching bird, perennial plant, pest, petal, phasianidae, piciformes, pincushion flower, plant, plant pathology, plant stem, plebejus, poales, pollen, pollinator, polygonia, polyommatus, polyporales, poultry, produce, pyronia cecilia, rabbit, rabbits and hares, rallidae, raven, red bugs, red fox, reef, rein, reindeer, reptile, rodent, rosa × centifolia, rosa arkansana, rosa canina, rose, rose family, rubus, russula integra, saffron crocus, salamander, salamandra, sandpiper, scaled reptile, scentless plant bugs, sea eagle, sea snail, seabird, seaduck, sharp shinned hawk, sheep, shell, shield bugs, shortstraw pine, shrike, shrub, silver-studded blue, silver-washed fritillary, silvery blue, skink, skipper (butterfly), small blue, small heath (butterfly), small pearl-bordered butterfly, small tortoiseshell, snail, snails and slugs, snout, snow crocus, soldier beetle, songbird, sorrel, southern dogface, southern |

		leopard frog, sow thistles, sparrow, spear thistle, spider, spider web, spotted salamander, spring crocus, squirrel, stallion, stock dove, subshrub, suidae, sunflower, swallowtail butterfly, swan, sweet grass, sweet peas, swift fox, tangle-web spider, teasel, terrestrial animal, terrestrial plant, thistle, thorns, spines, and prickles, thymelicus, tiger salamander, toad, tommie crocus, tortoise, tree, true frog, true salamanders and newts, true toad, trunk, tundra swan, turtle, twig, underwing moths, valerian, vanessa (butterfly), vascular plant, vegetable, vegetation, vertebrate, vicuña, vitis, vulture, wall lizard, wasp, water bird, water lily, waterfowl, weevil, western tiger swallowtail, wheat, whiptail, white-tailed deer, white stork, wild carrot, wildflower, wildlife, wing, wolf, wolf spider, wood, woodpecker, woody plant, working animal
Non-nature	Non-nature	abdomen, accessory fruit, accordion, accordionist, acoustic guitar, acting, active shirt, active shorts, adaptation, adventure, advertising, aerial photography, aerobatics, aerospace engineering, aerospace manufacturer, aerostat, afterglow, agricultural machinery, air racing, air show, air sports, air travel, aircraft, aircraft engine, airline, airliner, airplane, airport, airport apron, airship, alcohol, alcoholic beverage, alley, alligator, alloy wheel, aluminium, amber, american alligator, american crocodile, american crow, ancient history, animal product, animal shelter, animal sports, antique, antique car, apartment, aqua, aquarium, aqueduct, arbor day, arcade, arch, arch bridge, archaeological site, architecture, arctic, arctic ocean, arecales, arête, arm, armored car, army, art, art paint, artifact, artificial flower, artist, artwork, ash, asphalt, astronomical object, astronomy, athlete, athletic shoe, athletics, atlas, atmosphere, atmospheric phenomenon, audience, audio equipment, auto part, auto racing, automotive carrying rack, automotive decal, automotive design, automotive exhaust, automotive exterior, automotive fog light, automotive fuel system, automotive light bulb, automotive lighting, automotive mirror, automotive parking light, automotive side-view mirror, automotive side marker light, automotive tail & brake light, automotive tire, automotive wheel system, automotive window part, aviation, azure, baby, baby & toddler clothing, baby carriage, baby products, backlighting, backpack, bag, baked goods, baking, balance, ball, ball game, balloon, baluster, band plays, bangs, bank, banner, barbed wire, barechested, barefoot, barware, baseball, baseball cap, baseball equipment, baseball field, baseball glove, baseball park, baseball player, baseball protective gear, baseball uniform, basket, bassist, bat-and-ball games, bathroom, batting helmet, beacon, beam, beam bridge, beard, beauty, beef, beer, beer glass, beige, belt, bench, bengal tiger, bermuda shorts, bicycle, bicycle accessory, bicycle chain, bicycle clothing, bicycle fork, bicycle frame, bicycle handlebar, bicycle helmet, bicycle hub, bicycle jersey, bicycle part, bicycle saddle, bicycle seatpost, bicycle stem, bicycle tire, bicycle wheel, bicycle wheel rim, bicycles--equipment and supplies, big cats, billboard, biplane, bird's-eye view, bit, black, black-and-white,

| | | black hair, blazer, blimp, block party, blond, blue, blue-collar worker, board game, boardsport, boardwalk, boat, boats and boating--equipment and supplies, body jewelry, bonfire, book, boot, bottle, bouquet, bowl, box, box girder bridge, brand, brass instrument, brassiere, bread, brick, brickwork, bridal accessory, bridal clothing, bridal party dress, bride, bridge, bridle, bronze, bronze sculpture, brown, brown hair, building, building material, bulldozer, bullet train, bumper, bus, business jet, button, byzantine architecture, cabinetry, cable, cable-stayed bridge, cable car, cage, cake, calabaza, calm, camelid, camera, camera accessory, camera lens, cameras & optics, camouflage, campfire, camping, canal, candle, canopy, canopy walkway, cap, cape, car, car seat, car seat cover, carbon, carbonated soft drinks, cargo aircraft, cargo pants, carmine, carriage, cart, carving, cash crop, castle, ceiling, ceiling fixture, celebrating, celestial event, cello, cemetery, center console, ceramic, cg artwork, chain-link fencing, chair, champagne stemware, championship, chandelier, channel, chapel, charcoal, cheek, cheetah, chemical compound, chess, chessboard, chest, child, chimney, chin, chinese architecture, choreography, christmas, christmas decoration, christmas ornament, christmas tree, church, chute, circle, citrus, city, cityscape, classic, classic car, classical architecture, classical music, classical sculpture, cleanliness, cleat, clock, close-up, clothing, cloud, coat, cobblestone, cockpit, cocktail, coffee cup, coffee table, collaboration, collar, college baseball, college softball, colorfulness, colubridae, column, combat sport, combat vehicle, comfort, comfort food, commemorative plaque, commercial building, commercial vehicle, communication device, community, compact car, compact van, companion dog, competition, competition event, composite material, compost, computer, computer hardware, computer keyboard, computer monitor, concert, concrete, concrete bridge, condominium, cone, confectionery, constellation, construction, construction equipment, construction worker, contact sport, conversation, cooking, cookware and bakeware, cool, corn on the cob, costume, costume hat, cottage, couch, countertop, cowboy hat, crane, crankset, creative arts, crescent, crew, cricket, cricket cap, crocodile, crocodilia, croft, cross, cross-country equestrianism, cross-country skiing, crowd, cuisine, culinary art, cumulus, cup, cupboard, curtain, customer, cutlery, cycle sport, cycling, cycling shoe, cycling shorts, cylinder, cymbal, cynara, dairy, dance, darkness, dawn, daylighting, daytime, decoration, deejay, delicacy, denim, design, desk, dessert, devil's bridge, dew, diagram, digital camera, digital slr, dirt road, dirt track racing, disc dog, disc jockey, dish, dishware, display device, distilled beverage, ditch, diwali, dock, document, dog agility, dog collar, dog sports, dog supply, dome, door, door handle, downhill ski binding, downtown, drawer, drawing, dress, dress shirt, drink, drinking, drinking establishment, drinking water, drinkware, driveway, drizzle, drone, drop, drum, drum stick, drumhead, drummer, drums, dusk, ear, earrings, eclipse, elbow, electric blue, electric guitar, electric locomotive, electrical supply, |

| | | electrical wiring, electricity, electronic device, electronic instrument, electronic keyboard, electronic musical instrument, elephants and mammoths, emblem, emergency service, emergency vehicle, employment, endurance sports, enduro, engineer, engineering, english pleasure, english riding, entertainment, environmental art, equestrian, equestrian helmet, equestrian sport, equestrianism, equitation, erosion, evening, event, eventing, exercise, exhaust system, extreme sport, eye, eye glass accessory, eyebrow, eyelash, eyewear, facade, face, facial expression, facial hair, factory, family car, fan, farmer, farmhouse, farmworker, fashion, fashion accessory, fashion design, fast food, fault, fedora, fence, fender, festival, fictional character, fiddle, fighter aircraft, fighter pilot, film camera, fines herbes, finger, finger food, finial, fire, firefighter, fireplace, fireworks, fixed link, fixture, flag, flag of the united states, flagstone, flame, flap, flash photography, flat panel display, flat racing, flesh, flight, floor, flooring, floral design, floristry, flower arranging, flowerpot, fly, flying boat, flying disc, flying disc freestyle, fog, folding chair, folk instrument, folk wrestling, font, food, food craving, foot, football, football equipment, football player, footwear, forehead, fork, formal wear, formation, fountain, fourball, free reed aerophone, freeway, freezing, freight car, fried food, friendship, frisbee games, frost, fuel tank, full-size car, full moon, fun, function hall, furniture, gadget, galaxy, games, garden buildings, garden roses, garden tool, gardening, garnish, gas, gauge, gazebo, general aviation, gesture, girder bridge, glass, glass bottle, glasses, glider, gliding, glove, gluten, goal, goggles, gold, golf, golf bag, golf ball, golf cart, golf club, golf course, golf equipment, golfer, gothic architecture, government agency, gown, graffiti, grand prix motorcycle racing, graphic design, graphics, grappling, grave, green, grey, grille, grilling, ground attack aircraft, group a, groupset, guard rail, guitar, guitar accessory, guitarist, hair, hairstyle, hall, halter, hand, handbag, handle, handrail, handwriting, hangar, happy, harbor, hard hat, hardtop, hardwood, harvester, hat, haze, head, head restraint, headgear, headlamp, headpiece, headquarters, headstone, hearing, heat, helicopter, helicopter rotor, helmet, high-speed rail, high-visibility clothing, highway, hiking, hiking equipment, hill station, historic site, history, holiday, holy places, home, home appliance, home door, home fencing, hood, horizon, horn, horse and buggy, horse supplies, horse tack, horse trailer, hot air balloon, hot air ballooning, house, household hardware, houseplant, hub gear, hubcap, human, human body, human leg, human settlement, hurdle, hut, idiophone, illustration, individual sports, indoor games and sports, industry, infantry, inflatable, infrastructure, ingredient, interaction, interior design, intersection, intrusion, iris, iron, island, isle of man tt, jacket, japanese architecture, jaw, jazz pianist, jeans, jersey, jet aircraft, jet engine, jewellery, job, jockey, joint, jumping, junk food, kangaroo, keyboard, keyboard player, kick, kit car, kitchen, kitchen & dining room table, kitchen appliance, kitchen stove, kitchen utensil, kite, kite sports, klippe, |

| | | knee, knee pad, knife, knit cap, kodiak bear, ladder, lake district, laminate flooring, lamp, land lot, land vehicle, landing, landmark, landscape lighting, landscaping, lane, lap, laptop, laugh, lavender, law enforcement, lawn, lawn ornament, layered hair, leash, leather jacket, lectern, leg, leisure, lens, lens flare, leopard, light, light aircraft, light commercial vehicle, light fixture, lighthouse, lighting, lightning, line, line art, linens, lion, lip, liqueur, liquid, liver, living room, local food, lockheed martin, locking hubs, locomotive, log cabin, logo, long hair, luggage and bags, lumber, lunar eclipse, luxury vehicle, machine, macro photography, magenta, major appliance, makeover, manor house, map, marathon, marines, market, marketplace, masai lion, mast, match play, material property, measuring instrument, meat, mechanical fan, medieval architecture, meeting, megalith, melting, membranophone, memorial, mesh, metal, meteorological phenomenon, metropolis, metropolitan area, microphone, microphone stand, mid-size car, midnight, military, military aircraft, military camouflage, military helicopter, military officer, military organization, military person, military uniform, military vehicle, milky way, mirror, mirrorless interchangeable-lens camera, mist, mixed-use, mixing bowl, mixture, mobile phone, mode of transport, model aircraft, moisture, molding, monochrome, monochrome photography, monoplane, monument, moon, moonlight, morning, motocross, motor glider, motor vehicle, motorcycle, motorcycle accessories, motorcycle fairing, motorcycle helmet, motorcycle racer, motorcycle racing, motorcycling, motorsport, mountaineer, moustache, mouth, multi-sport event, multimedia, mural, muscle, music, music artist, music stand, music venue, musical ensemble, musical instrument, musical instrument accessory, musical keyboard, musician, mythology, nail, narrow-body aircraft, naval architecture, nebula, neck, necklace, neighbourhood, neon, net, net sports, new year, new year's eve, new years day, nickel, night, non-commissioned officer, nonbuilding structure, northern harrier, nose, notchback, number, nunatak, oar, observation tower, off-road racing, office equipment, office ruler, office supplies, official, one-piece garment, open-wheel car, optical instrument, orange, orator, organ, ornament, outdoor bench, outdoor furniture, outdoor play equipment, outdoor recreation, outdoor shoe, outdoor structure, outdoor table, outer space, outerwear, output device, overhead power line, overpass, paddle, paint, painting, palm tree, paper, paper product, parachute, parachuting, paragliding, parallel, parish, parking, party, party supply, passenger, passenger car, passenger ship, path, pattern, paw, pedestrian, people, people in nature, people on beach, percussion, percussion mallet, percussionist, performance, performance art, performance car, performing arts, performing arts center, peripheral, personal computer, personal luxury car, personal protective equipment, pet supply, photograph, photographer, physical fitness, pianist, piano, picnic table, picture frame, pier, pilot, pink, pink family, pint, pipe, piste, place of worship, plaid, |

| | | planet, plank, plantation, plaster, plastic, plastic bottle, plate, platter, play, player, playground, playing in the snow, playing sports, plucked string instruments, plumbing, plumbing fixture, plywood, podium, point-and-shoot camera, polar ice cap, pole, pollution, polo shirt, pop music, porcelain, porch, pork, portrait photography, poster, pottery, precipitation, precision sports, presentation, primate, product, professional golfer, projection screen, projector accessory, propeller, propeller-driven aircraft, property, public address system, public event, public space, public speaking, public transport, public utility, publication, purple, pyramid, quadrathlon, race car, race track, racing, radiator, radio-controlled aircraft, radio-controlled helicopter, radio-controlled toy, railroad car, railway, rainbow, rallycross, rallying, ranch, rangpur, rapid, real estate, rear-view mirror, recipe, recital, recreation, rectangle, red, red meat, red sky at morning, reed, reed instrument, reflection, reflex camera, relief, religious institute, religious item, residential area, resort town, retail, rice, riding boot, rim, road, road bicycle, road surface, roadster, rock-climbing equipment, rock concert, rolling, rolling stock, romance, roof, room, rope, rose order, rotor, rotorcraft, rubble, rugby, rugby short, ruins, ruler, running, runway, sacred lotus, saddle, sail, sailboat, sailing, saltwater crocodile, sandal, sash window, saxophone, saxophonist, scarf, science, scrap, screenshot, sculpture, security, selfie, serpent, serveware, service, serving tray, shack, shade, shadow, sharing, shed, shelf, shelving, shepherd, shin guard, ship, shirt, shoe, shorts, shoulder, show jumping, siberian tiger, sidewalk, siding, sign, sign language, signage, singer, singing, single-lens reflex camera, sink, sitting, skateboard deck, sketch, ski, ski binding, ski boot, ski equipment, ski helmet, ski pole, skier, skiff, skiing, skin, sky, skyscraper, sled, sledding, sleeve, slug, smile, smoke, snake, snapshot, sneakers, snow leopard, snowboard, snowboarding, soccer, soccer ball, soccer kick, soccer player, social group, sock, soft drink, solar energy, soldier, solution, solvent, song, sound, space, spandex, speaker, spectacle, speech, speed golf, speedometer, sphere, spire, splint boots, split-rail fence, spoke, spokesperson, spoon, sport utility vehicle, sport venue, sporting group, sports, sports car, sports car racing, sports equipment, sports gear, sports jersey, sports toy, sports uniform, sportswear, spring, square, squash, squeezebox, stable, stadium, stage, stage equipment, stained glass, stairs, standing, staple food, star, stately home, statue, steak, steam, steam engine, steel, steeple, steeplechase, steering part, steering wheel, stemware, step cutting, stick and ball sports, still life photography, stock photography, stone carving, stone wall, storm, stove, street, street art, street furniture, street light, street sign, string instrument, string instrument accessory, stunt, stunt performer, style, suburb, sugar house, suit, summer, sun, sun hat, sunglasses, sunlight, sunrise, sunset, superbike racing, superfood, supermoto, supersonic aircraft, suspension bridge, sweatpant, sweetness, swimming pool, swimwear, symbol, symmetry, synthetic rubber, t-shirt, table, tablecloth, tabletop |

		game, tableware, tachometer, tail, takeoff, tap, tar, tarpaulin, tartan, team, team sport, technology, telecommunications engineering, television, temperature, temple, tent, terrace, textile, theater curtain, theatre, thigh, thoroughfare, throat, thumb, thunder, thunderstorm, tie, tied-arch bridge, tiger, tile flooring, tints and shades, tire, tire care, titanium, toddler, tom-tom drum, tool, tooth, tour bus service, touring car racing, tourism, tournament, tower, tower block, town, toy, toy airplane, toy dog, toy vehicle, track, track racing, tractor, tradesman, tradition, traffic sign, trail, trailer, train, train station, trainer, tram, transmission tower, transmitter station, transparency, transparent material, transport hub, travel, travel trailer, tread, triangle, tripod, tropics, trousers, truck, trunks, truss bridge, tunnel, turbine, turboprop, turret, twinjet, ultimate, ultralight aviation, ultramarathon, umbrella, uniform, universe, urban area, urban design, urtica, vacation, van, varnish, vault, vegan nutrition, vehicle, vehicle brake, vehicle door, vehicle registration plate, vest, viaduct, village, vintage car, vintage clothing, viola, violet, violin, violin family, violist, violone, vision care, visual arts, visual effect lighting, volleyball, wagon, waist, walking, walking shoe, walkway, wall, waste, waste container, waste containment, watch, water bottle, water transportation, watercolor paint, watercraft, wedding ceremony supply, wedding dress, wedge, western riding, wheel, whiskers, white, white-collar worker, whole food, whole grain, wide-body aircraft, wind, wind farm, wind instrument, wind turbine, wind wave, windbreaker, windmill, window, window treatment, windscreen wiper, windshield, windsports, wine, wine bottle, wine glass, winter, winter sport, wire, wire fencing, women's football, wood flooring, wood stain, woodwind instrument, woolen, workwear, world, world rally car, woven fabric, wrestling, wrinkle, wrist, writing, yellow, youth, zeppelin

1200

Table S4. Overview of the extracted objects using the feature "object" of the Google Cloud Vision algorithm for Flickr and iNaturalist photographs of species. Only objects with frequency > 1 were included.

Category	Objects
Nature	Bird, Cattle, Centipede, Deer, Fish, Flower, Fox, Frog, Fruit, Goat, Hedgehog, Horse, Insect, Isopod, Lizard, Lynx, Mouse, Mule, Mushroom, Pig, Poultry, Rabbit, Sheep, Snail, Snake, Spider, Squirrel, Tortoise
Non-nature, pet and zoo animals or generic species	Person, Animal, Plant, Antelope, Shoe, Cat, Alpaca, Dog, Flowerpot, Tiger, Food, Footwear, Brown bear, Lion, Hat, Clothing, Poster, Pants, Bridge ,Vegetable, Window, Camel, Penguin, Kangaroo, Squash, Bench, Houseplant, Ruler, Flag, Tire, Lighting, Wheel, Bear, Building, Fire hydrant, Leopard, Apple, Car, Artichoke, Jaguar, Elephant, Vase, Luggage & bags, Top, Chair, Ladder, Outerwear, Red panda, Stool, Picture frame, Pillow, Ball, Figurine, Furniture, Tableware, Grape, Mirror, Sunglasses, Table top, Toy, Door, Dress, Zebra, Belt, Bicycle, Bicycle helmet, Bracelet, Christmas tree, Sculpture, House, Jeans, Packaged goods

1204

1205 Table. S5. Summary of the top 10 most frequently photographed objects, their distribution per user
1206 and their seasonal distribution.

Platform	Species	Number of images	Number of users	Images per user (mean)	Seasonal distribution (N° of images)			
					Spring	Summer	Autumn	Winter
Flickr	Flower	2198	430	5,11	977	729	393	99
	Insect	612	147	4,16	164	331	111	6
	Bird	510	141	3,61	191	163	103	53
	Cattle	484	178	2,71	134	154	182	14
	Horse	352	65	5,41	98	194	41	19
	Goat	118	50	2,36	29	27	40	22
	Sheep	116	50	2,32	21	20	35	40
	Mushroom	80	52	1,53	13	8	45	14
	Deer	67	39	1,71	11	14	33	9
	Frog	61	28	2,17	27	30	4	0
iNaturalist	Flower	2500	263	9,5	1105	981	383	31
	Insect	1230	157	7,83	450	611	156	13
	Bird	439	78	5,62	154	108	97	80
	Mushroom	260	40	6,5	34	23	197	6
	Deer	46	15	3,06	16	7	11	12
	Frog	43	29	1,48	25	10	8	0
	Lizard	41	33	1,24	19	13	8	1
	Spider	33	19	1,73	6	21	6	0
	Snail	32	22	1,45	16	16	0	0
	Fruit	18	11	1,63	1	9	8	0

1207

1208 Table S6. Overview of the extracted families using the BioCLIP algorithm for Flickr and iNaturalist
1209 photographs of the most photographed wildlife species.

Platform	Family
Flickr	Corvidae, Paridae, Phasianidae, Anatidae, Motacillidae, Ciconiidae, Emberizidae, Turdidae, Ardeidae, Phoenicopteridae, Gruidae, Ramphastidae, Rallidae, Troglodytidae, Panuridae, Pandionidae, Muscicapidae, Accipitridae, Cracidae, Scolopacidae, Paradisaeidae, Threskiornithidae, Podicipedidae, Cacatuidae, Meropidae, Laridae, Trochilidae, Laniidae, Pelecanidae, Recurvirostridae, Passeridae, Strigidae, Charadriidae, Columbidae, Hirundinidae, Upupidae, Fringillidae, Sturnidae, Acrocephalidae, Alcedinidae, Phalacrocoracidae, Bombycillidae, Cervidae, Canidae, Dasyproctidae, Dasypodidae, Leporidae, Bovidae, Moschidae, Giraffidae, Ranunculaceae, Fabaceae, Asteraceae, Rosaceae, Papaveraceae, Brassicaceae, Poaceae, Amaryllidaceae, Fagaceae, Plantaginaceae, Liliaceae, Betulaceae, Ulmaceae, Orchidaceae, Nymphaeaceae, Iridaceae, Apiaceae, Sapindaceae, Campanulaceae, Typhaceae, Caprifoliaceae, Asparagaceae, Celastraceae, Arecaceae, Colchicaceae, Salicaceae, Grossulariaceae, Gentianaceae, Orobanchaceae, Lauraceae, Violaceae, Caryophyllaceae, Lamiaceae, Geraniaceae, Hydrangeaceae, Primulaceae, Boraginaceae, Cornaceae, Polygonaceae, Pontederiaceae, Euphorbiaceae, Malvaceae, Convolvulaceae, Cyperaceae, Adoxaceae, Cactaceae, Vitaceae, Onagraceae, Scrophulariaceae,

	Tropaeolaceae, Lentibulariaceae, Lythraceae, Oleaceae, Magnoliaceae, Rubiaceae, Anacardiaceae, Oxalidaceae, Plumbaginaceae, Solanaceae, Polygalaceae, Nelumbonaceae, Santalaceae, Araceae, Nothofagaceae, Cucurbitaceae, Juglandaceae, Eriocaulaceae, Araliaceae, Bignoniaceae, Butomaceae, Rutaceae, Hypericaceae, Tamaricaceae, Saxifragaceae, Cannabaceae, Ranidae, Nyctibatrachidae, Bufonidae, Rhacophoridae, Hylidae, Hemiphractidae, Mantidae, Nymphalidae, Apidae, Aeshnidae, Libellulidae, Platycnemididae, Coenagrionidae, Lestidae, Calopterygidae, Chrysopidae, Carabidae, Tettigoniidae, Saturniidae, Acrididae, Gryllidae, Cerambycidae, Oecophoridae, Pentatomidae, Ascalaphidae, Gomphidae, Coccinellidae, Curculionidae, Polycentropodidae, Geometridae, Pieridae, Gerridae, Acanthosomatidae, Miridae, Noctuidae, Pterophoridae, Elateridae, Tipulidae, Papilionidae, Oedemeridae, Lycaenidae, Panorpidae, Coreidae, Formicidae, Cordulegastridae, Calliphoridae, Muscidae, Tarachodidae, Syrphidae, Nabidae, Tetrigidae, Crambidae, Cicadidae, Lycidae, Aphididae, Corduliidae, Tabanidae, Ephemeridae, Cetoniidae, Tenthredinidae, Conopidae, Scathophagidae, Chrysomelidae, Limoniidae, Lasiocampidae, Cantharidae, Sphingidae, Megachilidae, Bombyliidae, Halictidae, Zygaenidae, Tachinidae, Cleridae, Adelidae, Lacertidae, Emydidae, Fomitopsidaceae, Mycenaceae, Psathyrellaceae, Galeropsidaceae, Bankeraceae, Tricholomataceae, Hygrophoraceae, Agaricaceae, Amanitaceae, Marasmiaceae, Physalacriaceae, Russulaceae, Polyporaceae, Hymenogastraceae, Entolomataceae, Helicidae, Araneidae, Thomisidae, Linyphiidae
iNaturalist	Rallidae, Motacillidae, Fringillidae, Scolopacidae, Passeridae, Paridae, Ardeidae, Alcedinidae, Anatidae, Accipitridae, Phalacrocoracidae, Corvidae, Muscicapidae, Phylloscopidae, Turdidae, Ciconiidae, Charadriidae, Cettidae, Emberizidae, Laridae, Troglodytidae, Phasianidae, Icteridae, Acrocephalidae, Podicipedidae, Pelecanidae, Sturnidae, Columbidae, Picidae, Recurvirostridae, Threskiornithidae, Strigidae, Panuridae, Hirundinidae, Tyrannidae, Laniidae, Regulidae, Cervidae, Bovidae, Leporidae, Asteraceae, Polygonaceae, Boraginaceae, Asparagaceae, Lamiaceae, Celastraceae, Fagaceae, Fabaceae, Onagraceae, Geraniaceae, Orchidaceae, Resedaceae, Orobanchaceae, Plantaginaceae, Brassicaceae, Ranunculaceae, Amaryllidaceae, Salicaceae, Papaveraceae, Violaceae, Primulaceae, Rosaceae, Crassulaceae, Poaceae, Adoxaceae, Coriariaceae, Caryophyllaceae, Euphorbiaceae, Iridaceae, Berberidaceae, Ericaceae, Apiaceae, Thymelaeaceae, Colchicaceae, Caprifoliaceae, Campanulaceae, Cyperaceae, Polygalaceae, Rubiaceae, Apocynaceae, Gentianaceae, Linaceae, Juncaceae, Urticaceae, Hypericaceae, Typhaceae, Droseraceae, Lythraceae, Hamamelidaceae, Saxifragaceae, Melanthiaceae, Arecaceae, Convolvulaceae, Asphodelaceae, Scrophulariaceae, Potamogetonaceae, Penaeaceae, Zingiberaceae, Grossulariaceae, Araceae, Amaranthaceae, Solanaceae, Alismataceae, Betulaceae, Acanthaceae, Malvaceae, Rhamnaceae, Cornaceae, Eupteleaceae, Calceolariaceae, Menyanthaceae, Oleaceae, Liliaceae, Cistaceae, Rhizophoraceae, Lentibulariaceae, Aristolochiaceae, Sapindaceae, Bromeliaceae, Aquifoliaceae, Ceratophyllaceae, Cannabaceae, Tamaricaceae, Butomaceae, Nymphaeaceae, Cucurbitaceae, Ranidae, Salamandridae, Bufonidae, Hylidae, Pelodryadidae, Bombinatoridae, Pentatomidae, Libellulidae, Nymphalidae, Formicidae, Acrididae, Pieridae, Adelgidae, Tettigoniidae, Cetoniidae, Chrysomelidae, Geometridae, Apidae, Bombyliidae, Noctuidae, Megachilidae, Syrphidae, Coccinellidae, Hesperiidae, Crambidae, Gracillariidae, Erebidae, Curculionidae, Ascalaphidae, Lycaenidae, Carabidae,

<table>
<tr><td colspan="2">Pyralidae, Tortricidae, Cantharidae, Cleridae, Cicadidae, Platycnemididae, Calopterygidae, Megalodontesidae, Coenagrionidae, Cerambycidae, Asilidae, Cordulegastridae, Aeshnidae, Aphrophoridae, Lucanidae, Cicadellidae, Miridae, Stratiomyidae, Cecidomyiidae, Lasiocampidae, Pterophoridae, Zygaenidae, Ephemeridae, Sesiidae, Empididae, Melolonthidae, Rhagionidae, Papilionidae, Crabronidae, Lestidae, Corduliidae, Gomphidae, Muscidae, Limnephilidae, Ichneumonidae, Chrysopidae, Flatidae, Vespidae, Cynipidae, Pyrrhocoridae, Tipulidae, Micropezidae, Issidae, Cercopidae, Tarachodidae, Buprestidae, Nitidulidae, Panorpidae, Ectobiidae, Coleophoridae, Stathmopodidae, Saturniidae, Sphingidae, Sapygidae, Coreidae, Conopidae, Eumenidae, Staphylinidae, Pyrgomorphidae, Yponomeutidae, Reduviidae, Sialidae, Meloidae, Scathophagidae, Rutelidae, Mantidae, Geotrupidae, Odontoceridae, Hydrophilidae, Drepanidae, Depressariidae, Tetrigidae, Sarcophagidae, Andrenidae, Adelidae, Lygaeidae, Lacertidae, Anguidae, Scincidae, Dactyloidae, Gekkonidae, Cortinariaceae, Bankeraceae, Clavariaceae, Tricholomataceae, Strophariaceae, Pleurotaceae, Amanitaceae, Boletaceae, Hygrophoraceae, Russulaceae, Morchellaceae, Hymenogastraceae, Meruliaceae, Psathyrellaceae, Agaricaceae, Clavariadelphaceae, Physalacriaceae, Polyporaceae, Bolbitiaceae, Suillaceae, Fomitopsidaceae, Mycenaceae, Ganodermataceae, Hygrophoropsidaceae, Marasmiaceae, Pyronemataceae, Hydnangiaceae, Entolomataceae, Sclerodermataceae, Inocybaceae, Hymenochaetaceae, Omphalotaceae, Helvellaceae, Helicidae, Hygromiidae, Geomitridae, Limacidae, Arionidae, Araneidae, Agelenidae, Thomisidae, Ixodidae, Pisauridae, Sparassidae, Phalangiidae, Lycosidae</td></tr>
</table>

1210

1211 Table S7. Overview of the extracted genus using the BioCLIP algorithm for Flickr and iNaturalist
1212 photographs of the most photographed wildlife species

Platform	Genus
Flickr	Corvus, Periparus, Pavo, Anser, Motacilla, Chrysolophus, Anas, Ciconia, Emberiza, Turdus, Ardea, Cygnus, Phoenicopterus, Grus, Ramphastos, Aramides, Tadorna, Panurus, Pandion, Saxicola, Crax, Egretta, Tringa, Astrapia, Platalea, Podiceps, Cacatua, Fulica, Merops, Larus, Lanius, Pelecanus, Callonetta, Recurvirostra, Balearica, Passer, Bubo, Zoothera, Chroicocephalus, Charadrius, Buteo, Columba, Milvus, Hirundo, Upupa, Aix, Neophron, Gallus, Larvivora, Pica, Vanellus, Gallinago, Sturnus, Acrocephalus, Hydrocoloeus, Gallinula, Coloeus, Ixobrychus, Actitis, Calidris, Alcedo, Alopochen, Anthus, Lamprotornis, Phoenicurus, Rallus, Tachybaptus, Aythya, Lymnocryptes, Erithacus, Circus, Cyanistes, Mareca, Nycticorax, Microcarbo, Phalacrocorax, Sterna, Ichthyaetus, Bombycilla, Fringilla, Carduelis, Garrulus, Corythornis, Phimosus, Spatula, Odocoileus, Canis, Dasyprocta, Cervus, Tolypeutes, Oryctolagus, Dama, Rupicapra, Rucervus, Capra, Moschus, Bos, Capreolus, Muntiacus, Giraffa, Pelea, Kobus, Clematis, Lotus, Helianthus, Malus, Papaver, Chaenomeles, Cirsium, Brassica, Ficaria, Cardamine, Phragmites, Leucojum, Fagus, Carlina, Veronica, Anemone, Globularia, Prunus, Tulipa, Ulmus, Anacamptis, Cosmos, Corylus, Nymphaea, Vicia, Onobrychis, Iris, Angelica, Quercus, Gladiolus, Acer, Campanula, Typha, Dipsacus, Bromus, Muscari, Tussilago, Parnassia, Eranthis, Bellis, Echinops, Taraxacum, Ranunculus, Colchicum, Eschscholzia, Salix, Ribes, Gentiana, Populus, Leucanthemum, Nuphar, Rhinanthus, Rosa, Lolium, Epipactis, Lindera, Crepis, Euonymus,

Allium, Hyalopoa, Corypha, Viola, Narcissus, Orchis, Neottia, Gymnadenia, Hordeum, Tragopogon, Trifolium, Traunsteinera, Dactylorhiza, Ophrys, Crataegus, Limodorum, Himantoglossum, Silene, Pogonia, Neotinea, Helleborus, Pyrus, Phleum, Dactylis, Crocus, Lavandula, Geranium, Aruncus, Cichorium, Eupatorium, Hydrangea, Centaurium, Rudbeckia, Primula, Symphytum, Coreopsis, Gazania, Caltha, Hosta, Trisetum, Monarda, Sinapis, Helosciadium, Cornus, Spiranthes, Bistorta, Knautia, Calendula, Pontederia, Anemonastrum, Pisum, Nigritella, Melampyrum, Centaurea, Pseudocydonia, Erigeron, Euphorbia, Fritillaria, Valeriana, Zinnia, Malva, Salvia, Ornithogalum, Lactuca, Phacelia, Convolvulus, Platanthera, Carex, Ageratum, Calystegia, Viburnum, Tagetes, Tanacetum, Melocactus, Achillea, Vitis, Cotoneaster, Oenothera, Lonicera, Verbascum, Galanthus, Polygonatum, Furcraea, Carpinus, Tropaeolum, Pinguicula, Alopecurus, Lythrum, Aubrieta, Fraxinus, Fragaria, Cephalanthera, Magnolia, Aquilegia, Frasera, Daucus, Sanguisorba, Galium, Camelina, Cotinus, Astranthium, Oxalis, Liriodendron, Sophora, Yucca, Pelargonium, Linaria, Limonium, Solanum, Triticum, Reynoutria, Echium, Polygala, Nelumbo, Viscum, Pulsatilla, Astrantia, Filipendula, Potentilla, Anthriscus, Callistephus, Lemna, Nothofagus, Lathyrus, Cymbalaria, Calamagrostis, Artemisia, Sideritis, Sorghastrum, Rheum, Petasites, Lilium, Bellevalia, Luffa, Aegilops, Sambucus, Parthenocissus, Persicaria, Solidago, Juglans, Arnica, Carya, Wisteria, Rostraria, Dianthus, Phyteuma, Medicago, Melica, Rubus, Betula, Disa, Eriocaulon, Orobanche, Sonchus, Securigera, Hedera, Campsis, Chrysanthemum, Gentianella, Callirhoe, Trachycarpus, Butomus, Lamprocapnos, Ajuga, Tetradium, Rytidosperma, Miscanthus, Succisa, Cicerbita, Herminium, Hypericum, Ipomoea, Tamarix, Myosotis, Alchemilla, Chaerophyllum, Trollius, Carduus, Chrysosplenium, Sabazia, Scilla, Humulus, Gentianopsis, Tilia, Cortaderia, Citrus, Glechoma, Barbarea, Nigella, Pelophylax, Lankanectes, Rana, Bufo, Buergeria, Lithobates, Gastrotheca, Dryophytes, Mantis, Aglais, Apis, Anax, Crocothemis, Platycnemis, Coenagrion, Lestes, Calopteryx, Cicindela, Tettigonia, Attacus, Gryllus, Caligo, Harpella, Nezara, Libelloides, Stethophyma, Erythromma, Sympetrum, Paragomphus, Coccinella, Phaneroptera, Ischnura, Neureclipsis, Graphosoma, Idaea, Orthetrum, Pieris, Gerris, Harmonia, Tenodera, Hipparchia, Acanthosoma, Pseudomantis, Stagmomantis, Aeshna, Oedipoda, Agapanthia, Oxyagrion, Diachrysia, Gampsocleis, Leucorrhinia, Ctenicera, Cephalaeschna, Nephrotoma, Euthystira, Papilio, Oedemera, Phyllobius, Panorpa, Libellula, Leptoglossus, Formica, Anacridium, Psyllobora, Cordulegaster, Erebia, Euchorthippus, Lucilia, Colias, Musca, Polyommatus, Argynnis, Cucullia, Graphium, Tipula, Iris, Parasyrphus, Nabis, Tetrix, Semiothisa, Pseudocatharylla, Pseudopanthera, Vanessa, Sphaerophoria, Cicada, Melitaea, Chrysotoxum, Gonepteryx, Coenonympha, Pyrrhosoma, Agrypnus, Lygistopterus, Pseudochorthippus, Melanargia, Onychogomphus, Maniola, Anatis, Melanostoma, Dasysyrphus, Tuberolachnus, Cordulia, Silvius, Eupeodes, Polydrusus, Indolestes, Paracercion, Ephemera, Ceriagrion, Cetonia, Enallagma, Iphiclides, Oxya, Eristalis, Stenobothrus, Nematus, Pissodes, Helophilus, Cheilosia, Sicus, Scathophaga, Lasius, Carpocoris, Gomphus, Amblyptilia, Somatochlora, Clytra, Ormosia, Siproeta, Malacosoma, Cantharis, Chiasmia, Macrothylacia, Leptopterna, Xanthogramma, Deilephila, Chrysoteuchia, Hierodula, Cyaniris, Adelphocoris, Megachile, Bombylius, Anartia, Bombus, Hybomitra, Heliconius, Halictus, Hemaris, Zygaena, Scaeva, Episyrphus, Myathropa, Leptidea, Trichodes, Aphantopus, Fabriciana, Epistrophe, Volucella, Boloria, Anthocharis, Adalia, Nemophora, Lacerta, Emys,

	Fomitopsis, Mycena, Coprinopsis, Panaeolus, Hydnellum, Hygrocybe, Coprinus, Amanita, Marasmius, Macrolepiota, Flammulina, Russula, Cerioporus, Galerina, Laetiporus, Melanoleuca, Entoloma, Cepaea, Pseudotachea, Gasteracantha, Arachnura, Argiope, Araneus, Ebelingia, Frontinella, Misumena
iNaturalist	Gallinula, Motacilla, Spinus, Limosa, Passer, Parus, Ardea, Alcedo, Anas, Calidris, Buteo, Anser, Amazonetta, Phalacrocorax, Circaetus, Garrulus, Tringa, Erithacus, Corvus, Bubulcus, Gallicrex, Anthus, Phylloscopus, Turdus, Nycticorax, Fringilla, Cygnus, Ciconia, Vanellus, Cettia, Emberiza, Phoenicurus, Saxicola, Gallinago, Luscinia, Troglodytes, Tragopan, Larus, Acrocephalus, Carduelis, Chroicocephalus, Accipiter, Mergus, Sterna, Podiceps, Carpospiza, Pelecanus, Circus, Sturnus, Alopochen, Myrmecocichla, Columba, Jynx, Tadorna, Streptopelia, Himantopus, Plegadis, Numenius, Dendrocygna, Picus, Fulica, Netta, Aythya, Asio, Ixobrychus, Panurus, Hirundo, Molothrus, Milvus, Syrigma, Chlidonias, Mareca, Patagioenas, Xanthocephalus, Lanius, Regulus, Dama, Odocoileus, Capreolus, Rupicapra, Lepus, Mazama, Capra, Hydropotes, Bos, Cirsium, Bistorta, Phacelia, Galinsoga, Agave, Carlina, Thymus, Euonymus, Fagus, Trifolium, Epilobium, Erodium, Dactylorhiza, Gymnadenia, Anthericum, Anacamptis, Echium, Ajuga, Epipactis, Orchis, Quercus, Reseda, Orobanche, Bellis, Veronica, Senecio, Taraxacum, Ficaria, Cardamine, Galanthus, Ornithogalum, Leucojum, Salix, Tussilago, Narcissus, Corydalis, Ranunculus, Glechoma, Viola, Primula, Sonchus, Potentilla, Sedum, Anthoxanthum, Viburnum, Coriaria, Muscari, Lamium, Alopecurus, Prunus, Poterium, Myosotis, Euphorbia, Sisyrinchium, Berberis, Helleborus, Salvia, Rhododendron, Geranium, Silene, Doronicum, Adoxa, Daphne, Polygonatum, Barbarea, Tragopogon, Colchicum, Lotus, Dipsacus, Aquilegia, Triphysaria, Noccaea, Bromus, Rumex, Phyteuma, Carum, Poa, Ophrys, Petasites, Carex, Polygala, Galium, Cerastium, Briza, Vincetoxicum, Gentiana, Foeniculum, Agrostemma, Centaurea, Leopoldia, Himantoglossum, Murbeckiella, Matricaria, Actaea, Vicia, Crataegus, Rhinanthus, Anthyllis, Fragaria, Anemone, Cypripedium, Hippocrepis, Linum, Antennaria, Lathyrus, Luzula, Pulsatilla, Alchemilla, Thalictrum, Trollius, Geum, Leucanthemum, Urtica, Hypericum, Achillea, Pyrola, Typha, Sparganium, Cephalanthera, Traunsteinera, Melilotus, Drosera, Phelipanche, Allium, Tanacetum, Cicerbita, Campanula, Stachys, Persicaria, Coreopsis, Hylotelephium, Goodyera, Pilosella, Milium, Lysimachia, Medicago, Lythrum, Carduus, Onobrychis, Gentianella, Euphrasia, Dianthus, Malus, Linaria, Calamagrostis, Origanum, Teucrium, Crocus, Aster, Filipendula, Papaver, Plantago, Zea, Brassica, Convallaria, Sambucus, Cyclamen, Schoenus, Fothergilla, Iris, Aruncus, Securigera, Chrysosplenium, Aconitum, Festuca, Veratrum, Cruciata, Thlaspi, Capsella, Fumaria, Sinapis, Saxifraga, Valeriana, Licuala, Valerianella, Sherardia, Convolvulus, Legousia, Seseli, Eremogone, Kniphofia, Globularia, Arabis, Hieracium, Oenanthe, Symphytum, Phleum, Bellevalia, Genista, Scrophularia, Lupinus, Alliaria, Soldanella, Platanthera, Cladium, Oenothera, Rosa, Stuckenia, Schoenoplectiella, Sanguisorba, Cota, Brachysiphon, Succisa, Roscoea, Sesleria, Tephroseris, Ribes, Lemna, Setaria, Amaranthus, Panicum, Schoenoplectus, Calystegia, Marrubium, Swertia, Parnassia, Eriophorum, Angelica, Phragmites, Nicandra, Cytisus, Mentha, Symphyotrichum, Digitaria, Erigeron, Hottonia, Echinodorus, Daucus, Corylus, Aegonychon, Scabiosa, Andropogon, Ononis, Knautia, Crepis, Ulex, Artemisia, Melissa, Malva, Frangula, Cornus, Pulmonaria, Euptelea, Calceolaria, Spiranthes, Argyranthemum, Neottia, Lepidium, Vaccinium, Menyanthes, Fraxinus, Laburnum, Nasturtium, Rorippa, Tulipa, Helianthemum, Hudsonia, Borago, Dactylis, Bruguiera, Arenaria, Anemonastrum, Pedicularis, Picris,

| | Utricularia, Galatella, Stellaria, Tilia, Comarum, Betula, Zizia, Betonica, Circaea, Jacobaea, Agrimonia, Asarum, Acer, Cotoneaster, Fagopyrum, Scutellaria, Wendlandia, Centaurium, Gentianopsis, Clematis, Aechmea, Sorbus, Hamamelis, Anthriscus, Moraea, Melittis, Leontodon, Alnus, Holcus, Chaerophyllum, Scorzonera, Argentina, Ainsworthia, Hypochaeris, Tuberaria, Verbascum, Ilex, Heracleum, Clinopodium, Myosoton, Lactuca, Alisma, Cichorium, Aspalathus, Eriocapitella, Conioselinum, Galeopsis, Scorzoneroides, Pastinaca, Solidago, Potamogeton, Ceratophyllum, Tadehagi, Cannabis, Delphinium, Mercurialis, Tamarix, Caltha, Limodorum, Scirpus, Nigella, Digitalis, Pinguicula, Melampyrum, Lonicera, Trisetum, Lolium, Heliophila, Mimulopsis, Lilium, Bupleurum, Juncus, Butomus, Pulicaria, Eupatorium, Halerpestes, Glebionis, Populus, Hypoestes, Sechium, Andromeda, Blackstonia, Neotinea, Calluna, Chaenomeles, Aubrieta, Brunnera, Gagea, Psephellus, Olea, Hibiscus, Symphoricarpos, Spirodela, Lespedeza, Arum, Rubus, Rana, Pelophylax, Neurergus, Bufo, Hyla, Salamandra, Ranoidea, Bombina, Lissotriton, Pseudacris, Atelopus, Sympetrum, Vanessa, Formica, Colias, Anacridium, Adelges, Oxythyrea, Cryptocephalus, Polygonia, Aglais, Bombus, Bombylius, Anthocharis, Gonepteryx, Lordotus, Boloria, Issoria, Osmia, Allograpta, Coccinella, Ematurga, Erynnis, Scopula, Coenonympha, Chiasmia, Chrysoteuchia, Phyllobius, Libelloides, Polyommatus, Cicindela, Platytes, Oncocera, Petrophora, Aricia, Pelochrista, Cantharis, Trichodes, Metaleptea, Epirrhoe, Diacrisia, Gampsocleis, Camptogramma, Euthystira, Idaea, Euclidia, Platycnemis, Pyrgus, Melanargia, Tettigonia, Calopteryx, Orthetrum, Leptinotarsa, Harmonia, Adalia, Maniola, Megalodontes, Pieris, Coenagrion, Libellula, Cordulegaster, Anax, Crocothemis, Aphrophora, Xanthogramma, Sphaerophoria, Lucanus, Minois, Erebia, Argynnis, Pseudochorthippus, Pyrausta, Aeshna, Dryas, Hypochalcia, Lysandra, Stenodema, Nothomyia, Jaapiella, Olethreutes, Graphosoma, Malacosoma, Callophrys, Pararge, Cupido, Autographa, Crambus, Lasiommata, Pyrrhosoma, Zygaena, Hypera, Ephemera, Synanthedon, Cicadella, Empis, Ectinohoplia, Ladona, Dasineura, Rhagio, Lethe, Phyciodes, Papilio, Conocephalus, Cerceris, Erythemis, Zizina, Plebejus, Nomophila, Leptidea, Macrothylacia, Lestes, Somatochlora, Heteronympha, Carcharodus, Orchelimum, Gomphus, Onychogomphus, Austrogynacantha, Coenosia, Lycaena, Udea, Allotruxalis, Scaeva, Scoparia, Chrysopa, Exochomus, Siphanta, Vespa, Ruspolia, Sympecma, Ischnura, Chorthippus, Diplolepis, Chrysochraon, Psophus, Nymphalis, Orthosia, Phytometra, Actinote, Pseudopanthera, Mnais, Micropeza, Propylea, Issus, Cercopis, Siona, Glaucopsyche, Therion, Mythimna, Pontia, Metrioptera, Xylocopa, Leucorrhinia, Melitaea, Cerambyx, Iris, Psiloptera, Erythrodiplax, Cychramus, Miramella, Omocestus, Panorpa, Episyrphus, Ectobius, Coleophora, Stathmopoda, Rivula, Crorema, Chrysotoxum, Saturnia, Deilephila, Sigaus, Aelia, Chrysis, Meconema, Chalcolestes, Dissosteira, Dolycoris, Brintesia, Adelpha, Zicrona, Carabus, Centrocoris, Gnorimus, Nephrotoma, Stenobothrus, Andricus, Tipula, Polistes, Thesium, Maura, Enallagma, Celastrina, Trichius, Erythromma, Stenotus, Calocoris, Eurema, Yponomeuta, Scotopteryx, Rhynocoris, Icaricia, Sialis, Notogomphus, Helophilus, Lytta, Eucera, Nehalennia, Usia, Euphydryas, Scathophaga, Lythria, Diachrysia, Phyllopertha, Celypha, Anania, Acrida, Aphantopus, Stethophyma, Thyanta, Pyronia, Murgantia, Attactagenus, Pero, Sphinx, Gomphocerippus, Timarcha, Pasimachus, Geotrupes, Geometra, Chrysina, Otiorhynchus, Vitisiella, Odontocerum, Decticus, Spulerina, Watsonalla, Carcina, Tetrix, Coenagriocnemis, Sarcophaga, Hyponephele, |

| | Fabriciana, Buprestis, Ochlodes, Nemophora, Spilostethus, Parnassius, Tibicina, Aporia, Eana, Aromia, Anguis, Lacerta, Podarcis, Anolis, Zootoca, Cortinarius, Hydnellum, Clavulinopsis, Clitocybe, Pholiota, Infundibulicybe, Pseudotricholoma, Pleurotus, Saproamanita, Boletus, Sarcodon, Hygrocybe, Gliophorus, Melanoleuca, Russula, Abortiporus, Coprinopsis, Tricholoma, Hypholoma, Amanita, Leucoagaricus, Coprinellus, Clavariadelphus, Flammulina, Lactarius, Morchella, Pycnoporus, Fomes, Macrolepiota, Porpolomopsis, Cuphophyllus, Suillus, Ampulloclitocybe, Postia, Armillaria, Hemimycena, Ganoderma, Hygrophoropsis, Conocybe, Protostropharia, Mycena, Laccaria, Entoloma, Hygrophorus, Psilocybe, Scleroderma, Strobilurus, Galerina, Inocybe, Beenakia, Marasmius, Hebeloma, Coprinus, Trametes, Lycoperdon, Leccinum, Calvatia, Kuehneromyces, Omphalotus, Melanotus, Leucocoprinus, Mycetinis, Helvella, Stropharia, Fomitopsis, Helix, Monacha, Cepaea, Helicella, Eumilax, Arion, Faustina, Trochulus, Argiope, Misumena, Ixodes, Xysticus, Araneus, Malthonica, Larinioides, Micrommata, Pisaura, Aculepeira, Agelena, Phalangium, Araniella, Pardosa |

1213