DuoHexaBody-CD37 induces direct cytotoxic signaling in diffuse large B-cell lymphoma

Simar Pal Singh[1], Michelle D. van den Beukel[1], Sjoerd van Deventer[1], Marije B. Overdijk[2], M. Guy Roukens[2], Kim C.M. Santegoets[2], Esther C.W. Breij[2], Martin ter Beest[1], Annemiek B. van Spriel[1,*]

[1]Department of Medical BioSciences, Radboud Institute for Medical Innovation, Radboud University Medical Center, Nijmegen, the Netherlands.

[2]Genmab, Utrecht, the Netherlands.

*Corresponding author: Annemiek van Spriel (annemiek.vanspriel@radboudumc.nl)

Phone number: +31 24317600

Running title: Cytotoxic signaling by DuoHexaBody-CD37

Abstract

Diffuse large B-cell lymphoma (DLBCL) is a common aggressive form of Non-Hodgkin lymphoma. Tetraspanin CD37 is highly expressed on mature B cells and being studied as a therapeutic target for NHL, including DLBCL. DuoHexaBody-CD37 is a biparatopic antibody with an E430G hexamerization-enhancing mutation targeting two non-overlapping CD37 epitopes shown to promote complement-dependent cytotoxicity. However, the impact of DuoHexaBody-CD37 on direct cytotoxic signaling has not yet been studied. Here we demonstrate that DuoHexaBody-CD37 induces direct cytotoxicity in DLBCL-derived tumor cell lines independent of the subtype. DuoHexaBody-CD37 induced significant CD37 clustering and was retained at the cell surface in contrast to rituximab, which was internalized. Unbiased screening identified the modulation of 26 (phospho)proteins upon DuoHexaBody-CD37 treatment of primary B cells or DLBCL cells. Whereas DLBCL cells predominantly upregulated p-SHP1(Y564) upon DuoHexaBody-CD37 treatment, primary B cells showed significantly increased p-AKT(S473) and MAPK signaling which is linked to cell survival. Studies using CD37-mutants identified the N-terminus to be involved in DuoHexaBody-CD37-induced signaling. Finally, DuoHexaBody-CD37 treatment inhibited cytokine pro-survival signaling in DLBCL cells. These findings provide novel insights into the signaling functions of CD37 upon DuoHexaBody-CD37 treatment, and open up opportunities for developing CD37-immunotherapy in combination with small molecule inhibitors to maximize tumor cell death.

Introduction

Diffuse large B-cell lymphoma (DLBCL), an aggressive type of mature B-cell lymphoma, accounts for approximately one third of Non-Hodgkin lymphoma (NHL) cases. DLBCL arises from mature B cells in the lymph node, and comprises a heterogenous group of tumors that can be classified according to cell-of-origin (germinal-center B-cell (GCB) derived and activated B-cell (ABC) subtypes). In addition, recent comprehensive molecular profiling has identified five to seven genetic subtypes that share similar oncogenic pathways [reviewed in [1]]. A large fraction of DLBCL patients (up to 40%) do not respond or relapse after first line treatment with chemotherapy (cyclophosphamide, hydroxydaunorubicin, vincristine sulfate (Oncovin), and prednisone: CHOP) and immunotherapy (rituximab). Rituximab and also next-generation antibodies (ofatumumab) target CD20 at the plasma membrane of DLBCL [2], however decreased expression of CD20 is related to inferior clinical outcome after treatment with R-CHOP [3,4,5]. Thus, targeting alternative B-cell membrane proteins and development of novel immunotherapies are needed to improve clinical outcomes of patients with relapsed/refractory DLBCL [6,7].

CD37 is a tetraspanin protein with four-transmembrane domains, which is predominantly expressed by mature B cells [8–10]. Tetraspanins interact with immune receptors on the same cell (in cis) and control membrane organization in lymphocytes [11–13]. CD37 is absent on progenitor B cells and terminally differentiated plasma cells making it an ideal target for mature B cell malignancies [14,15,16]. In DLBCL patients, CD37 was reported to be an independent prognostic factor for both GCB and ABC subtypes [4]. In line with these findings, CD37-deficient mice show defects in humoral and cellular immune responses, and spontaneously develop mature B-cell lymphoma, which is dependent on IL-6 [17]. IL-6 signals through the IL-6 receptor complex that activates AKT kinase and STAT3 signaling stimulating

cell survival and proliferation. IL-6 signaling has been reported to be a critical driver in the

tumor microenvironment and a negative prognostic factor in diffuse large B-cell lymphoma

[18].

CD37 is under preclinical and clinical investigation as novel therapeutic target for B-

NHL, including antibody-based and chimeric antigen receptor (CAR) T-cell therapies [19–23,24].

DuoHexaBody-CD37 (GEN3009) is a novel biparatopic immunoglobulin G1 (IgG1) with a point

mutation in the Fc-domain (E430G) that enhances antibody hexamerization upon binding to

the cell surface [25]. This facilitates C1q binding and results in potent complement-dependent

cytotoxicity (CDC). In addition DuoHexaBody-CD37 can induce Fc-gamma receptor (FcγR)

mediated antibody-dependent cellular cytotoxicity (ADCC) and phagocytosis (ADCP) by

effector cells. DuoHexaBody-CD37 mediates superior CDC in patient-derived DLBCL cells ex

vivo compared to CD20 monoclonal antibodies as shown in preclinical studies [26].

In B cells, CD37 has been reported to connect to intracellular signaling pathways via

its two intracellular tails [9,27]. The N-terminal tail of CD37 contains a potential 'ITIM-like' motif

that may induce SHP1-dependent cell-death, whereas the C-terminal tail of CD37 bears a

predicted 'ITAM-like' motif that may stimulate AKT kinase-dependent survival [27]. Here we

investigated whether DuoHexaBody-CD37 is capable of inducing intracellular signaling that

induces cell death of malignant B cells. We report that DuoHexaBody-CD37 mediates direct

CD37-mediated signaling and evokes tumor cytotoxicity in DLBCL-derived tumor cells,

independent of the presence of complement. DuoHexaBody-CD37 modulated the PI3K-AKT

and MAPK signaling pathways, with differences observed between primary B cells and DLBCL

tumor cells. Moreover, DuoHexaBody-CD37 inhibited IL-4-dependent p-STAT6 and IL-6/ IL-21-

dependent p-STAT3 signaling in DLBCL cells. In conclusion, this study shows a novel

mechanism of action of DuoHexaBody-CD37, and provides more insight into CD37 as

therapeutic target for B cell malignancies.

Results

DuoHexaBody-CD37 mediates direct cytotoxicity in DLBCL cell lines

Induction of apoptosis in B cell leukemia and lymphomas has been reported for therapeutic antibodies such as obinutuzumab [28], alemtuzumab [29] and otlertuzumab [30]. To investigate the direct cytotoxic potential of DuoHexaBody-CD37, DLBCL-derived cell lines classified as GCB (Oci-Ly8 and Oci-Ly7), ABC (HBL-1 and U2932) and Burkitt lymphoma-derived cell lines (BJAB, Daudi) were treated with DuoHexaBody-CD37 in the absence of complement, and with or without an Fc-targeting F(ab)2-fragment (a-Fc) to facilitate Fc-mediated crosslinking. Whereas the viability of Burkitt lymphoma-derived cell lines (BJAB and Daudi) was not significantly affected by DuoHexaBody-CD37 treatment with a-Fc, all tested DLBCL-derived cell lines showed >25% reduction in viability after DuoHexaBody-CD37 treatment with a-Fc (Figure 1A). To investigate the physiological relevance of Fc-crosslinking mediated DuoHexaBody-CD37 induced killing, cytotoxicity assays of DLBCL cell lines were performed using peripheral blood mononuclear cells (PBMCs) expressing Fcɣ receptors. In these assays, PBMCs were pre-fixed to prevent FcɣR signaling and to avoid any influence of ADCC/ADCP as previously shown with DuoHexaBody-CD37 [25]. DuoHexaBody-CD37 treatment resulted in significantly increased direct cytotoxicity of DLBCL cells compared to untreated cells in presence of PBMCs as source for Fc-crosslinking for all donors tested (Figure 1B). Collectively, these results show induction of direct cytotoxicity by DuoHexaBody-CD37 treatment with Fc-crosslinking in different DLBCL cells.

DuoHexaBody-CD37 induces CD37 clustering without modulating CD37 cell surface expression

CD37 is involved in the spatial organization of the B-cell plasma membrane by forming

tetraspanin nanodomains [13], therefore we investigated CD37 membrane clustering on

DuoHexaBody-CD37-treated tumor cells compared to tumor cells treated with the IgG1

isotype control antibody. Whereas CD37 surface expression was the same in both samples,

the clustering of CD37 on tumor cells (measured by fluorescence intensity/area) was

significantly higher upon DuoHexaBody-CD37 treatment than isotype control antibody

treatment (Figure 2A-B). Interestingly, DuoHexaBody-CD37-mediated clustering of CD37 was

observed in both DLBCL (Oci-Ly8) and Burkitt cells (BJAB), even in absence of Fc-crosslinker

(Figure 2A-B). Thus, DuoHexaBody-CD37 induces significant CD37 membrane clustering at

the cell surface of malignant B cells which is not dependent on Fc-mediated crosslinking.

Since the efficacy of rituximab can be negatively impacted by induction of CD20

internalization, leading to loss of CD20 expression at the cell surface [31,32], we investigated

whether DuoHexaBody-CD37 affected CD37 cell surface accessibility. We observed no

changes in surface CD37 expression upon DuoHexaBody-CD37 treatment in all cell lines

tested, in contrast to CD20 expression, which was decreased by two-fold on DAUDI, Oci-Ly8

and Oci-Ly7 cells upon rituximab treatment (Figure 2C).

DuoHexaBody-CD37 treatment results in activation of multiple downstream signaling

pathways

Next, we focused on investigating mechanisms underlying the direct cytotoxicity upon

DuoHexaBody-CD37 treatment by analyzing DuoHexaBody-CD37-induced downstream

signaling using high throughput Reverse Phase Protein Array (RPPA) [33]. This technique

enables simultaneous unbiased measurement of 484 proteins (102 phospho targets and 382

total proteins) in multiple samples at once. For this analysis primary B cells (purified CD19+ B

cells from PBMCs) and two DLBCL-derived cell lines (Oci-Ly7 and U2932) were treated with

DuoHexaBody-CD37 in the presence or absence of Fc-crosslinker and compared to untreated

samples from each cell type. Principal component analysis on all tested proteins in RPPA

identified 3 separate clusters correlating with the different cell models (Supplementary

Figure 1). Compared to respective untreated controls, 26 phosphoproteins showed >two-fold

change in either of the three cell models tested upon DuoHexaBody-CD37 treatment (Figure

3A). Pathway analysis on all 26 phospho-sites identified PI3K/AKT/MTOR signaling as the

most significantly enriched pathway upon DuoHexaBody-CD37 treatment (Figure 3B). In

addition, some of these signaling proteins are downstream of B cell receptor (BCR) signaling

and RAS signaling pathways, which have been reported to be essential for B cell survival and

lymphomagenesis [34,35].

Next, the effect of DuoHexaBody-CD37 treatment on primary B cells versus tumor

cells was studied by focusing on those phosphoproteins that showed approximately two-fold

upregulation in signal compared to untreated controls. Primary B cells showed upregulation

of multiple phosphoproteins upon DuoHexaBody-CD37 treatment including p-AKT, p-FAK, p-

Src and p-Shc. In DLBCL cells, different phosphoproteins (p-RPA32, p-DAPK1, p-PAK, p-PI3K

and p-FRS-alpha) were upregulated upon DuoHexaBody-CD37 treatment. The list of targets

and fold changes can be found in Supplementary Table 1.

To validate these results, we analyzed different targets within the three main signaling

pathways that came out of the RPPA analysis: the PI3K/AKT pathway, BCR pathway and

MAPK/ERK pathway using phosphoflow analysis. Anti-BCR (IgM) was used as stimulation

control in the phosphoflow studies. We confirmed specific upregulation of p-AKT(S473) in

primary B cells upon treatment with DuoHexaBody-CD37 in the presence of Fc-crosslinker

compared to untreated cells (Figure 3C). In DLBCL cell lines, upregulation of p-AKT(S473) was

not specific for DuoHexaBody-CD37 treatment as this was already observed in presence of

the Fc-crosslinker only (Figure 3C). These data indicate a differential role of DuoHexaBody-

CD37 mediated signaling in primary B cells versus DLBCL which could be partially explained

by constitutive activation of the PI3K-AKT pathway in DLBCL [36]. As engagement of cell death

pathways in CLL cells was shown to be dependent on SHP1 recruitment and activation via

CD37 N-terminus [27], we additionally assessed p-SHP1(Y564) status upon DuoHexaBody-CD37

treatment of primary B cells and DLBCL cells via phosphoflow. Compared to untreated cells,

DuoHexaBody-CD37 treatment in the presence of Fc-crosslinker induced significant

upregulation of p-SHP1(Y564) in both primary B cells and DLBCL cells (Figure 3C,

Supplementary Figure 2A).

To analyze phosphoproteins downstream of the BCR, we studied p-SYK, p-BTK, p-

PLCγ2 and p-PKC in primary B cells and DLBCL cells by phosphoflow. p-SYK was observed to

be upregulated only after BCR stimulation (anti-IgM), but not after DuoHexaBody-CD37

treatment of primary B cells or DLBCL cells (Figure 3C). On other hand, p-PLCγ2(Y759) and p-

BTK(Y223) were specifically upregulated in DLBCL cells upon treatment with DuoHexaBody-

CD37 in the presence of Fc-crosslinker (Figure 4A, Supplementary Figure 2B). Next, we

analyzed the MAPK signaling pathway and observed p-P38 to be specifically upregulated in

primary B cells, but not in DLBCL cells, upon DuoHexaBody-CD37 treatment in the presence

of Fc-crosslinker (Figure 4B, Supplementary Figure 2C). Similarly, ERK phosphorylation was

upregulated in primary B cells upon DuoHexaBody-CD37 treatment in the presence of Fc-

crosslinker, in contrast to DLBCL cells (Figure 4B, Supplementary Figure 2C). Taken together,

DuoHexaBody-CD37 treatment activates different signaling cascades in primary B cells versus

DLBCL-derived tumor cells. Specific upregulation of p-SHP1 in DLBCL cells could provide an

explanation of the direct cytotoxicity observed upon treatment with DuoHexaBody-CD37.

CD37 N-terminus is involved in DuoHexaBody-CD37-mediated signaling

As tyrosine phosphorylation of the CD37 N-terminus was reported to be crucial in CD37-mediated signaling [27], we examined the involvement of CD37 N-terminus in DuoHexaBody-CD37-induced signaling. In these experiments, B-ALL cells (NALM6) with low endogenous CD37 expression were used to investigate various CD37 mutant constructs [27]. NALM6 cells transfected with wild type CD37 (CD37-WT), or with CD37 carrying either the Tyr13 to phenylalanine mutation (CD37-Y13F) or a deletion of Tyr13 residue (CD37-ΔY13) in the cytosolic region of the CD37 molecule were generated. Cell surface expression of CD37-WT and both mutant variants was validated by flow cytometry (Figure 5A). DuoHexaBody-CD37 treatment in the presence of Fc-crosslinker resulted in significant upregulation of p-AKT(S473) and p-SHP1(Y564) in NALM6 cells expressing CD37-WT compared to treatment with only the crosslinker (Figure 5B, C) in line with findings observed in the DLBCL-derived cell lines (Figure 3C). Interestingly, deleting Tyr13 (CD37-ΔY13) completely impaired p-AKT(S473) and p-SHP1(Y564) upregulation, whereas mutating Tyr13 (CD37-Y13F) did not result in major changes upon DuoHexaBody-CD37 treatment in the presence of Fc-crosslinker (Figure 5B, C). In conclusion, these data validate active CD37 downstream signaling upon target engagement with DuoHexaBody-CD37 in the presence of Fc-crosslinker, and suggest the involvement of the CD37 N-terminus in mediating direct cytotoxic signaling.

DuoHexaBody-CD37 treatment abrogates cytokine mediated pro-survival signaling in DLBCL cells

As cells within the DLBCL tumor microenvironment can stimulate tumor growth by supplying cytokines [37] and CD37 has been reported to inhibit IL-6 signaling [17], we next investigated the

impact of DuoHexaBody-CD37 treatment on cytokine-mediated oncogenic signaling.

Different DLBCL cell lines were treated with DuoHexaBody-CD37 in absence or presence of

Fc-crosslinker, followed by stimulation with either recombinant human IL-4 (rh-IL4), IL-6 (rh-

IL6) or IL-21 (rh-IL21) and analyzed for p-STAT activation by phosphoflow. These cytokines

were selected because of their pro-tumorigenic role in DLBCL and other hematological

cancers [18, 37,38,39,40]. All tested DLBCL-derived tumor cell lines showed a significant increase in

the level of p-STAT6 upon rh-IL-4 stimulation. Pre-treatment with DuoHexaBody-CD37 in the

presence of a crosslinker showed a significant decrease in percentage p-STAT6-positive tumor

cells upon stimulation with rh-IL-4 (Figure 6A) in contrast to controls (untreated or cells

treated with DuoHexaBody-CD37 or Fc-crosslinker alone). Next, IL-6-induced p-STAT3

signaling was investigated in HBL-1 cells, as the other DLBCL-derived tumor cell lines did not

respond to IL-6 (data not shown). A two-fold increase in percentage p-STAT3-positive HBL-1

cells was observed upon rh-IL6 stimulation. Interestingly, treatment with DuoHexaBody-CD37

in the presence of a crosslinker also decreased the percentage of p-STAT3-positive HBL-1 cells

compared to controls (untreated or cells treated with DuoHexaBody-CD37 or crosslinker

alone) although this was not significant (Figure 6B). Finally, IL-21-induced p-STAT3 signaling

was investigated in all four cell lines. Both ABC-DLBCL derived tumor cells (HBL-1 and U2932)

showed a > four-fold increase in the percentage of p-STAT3-positive cells, while the effect

was more modest in the GCB-DLBCL-derived tumor cells (Oci-Ly8) showing a two-fold

increase in p-STAT3 signal. Oci-Ly7 cells did not respond to IL-21 (data not shown).

DuoHexaBody-CD37 treatment in the presence of Fc-crosslinker showed in all three IL-21

responsive cell lines a significant decrease in the percentage of p-STAT3-positive cells upon

rh-IL21 stimulation (Figure 6C) in line with IL-4 and IL-6 stimulation. Collectively, these

experiments show that DuoHexaBody-CD37 treatment in the presence of Fc-crosslinker

inhibits cytokine-mediated pro-survival signaling in the DLBCL tumor microenvironment.

Discussion

A large fraction (~40%) of patients with DLBCL do not respond to or relapse after first-line treatment with standard immunochemotherapy (R-CHOP), emphasizing the unmet medical need to develop new therapies for DLBCL. Tetraspanins have gained strong interest as therapeutic targets in different cancer types due to their capacity to regulate cell proliferation, adhesion and migration. Importantly, many tumor cells have altered cell surface expression of tetraspanins making them attractive targets for the treatment of cancer [16,41]. For example, CD151 is upregulated in carcinomas and CD151 stimulates both primary tumor growth and the metastatic cascade through its interaction with $\beta1$ integrins [41,42]. In B cell lymphoma, CD81 has been reported to be highly expressed by malignant cells from patients with B-NHL, and CD81-specific antibodies have been reported to be cytotoxic for such cells while sparing normal B cells [43]. CD37 represents an alternative therapeutic target for B-NHL because of its specific membrane expression on mature B cells, and its prognostic value in DLBCL and follicular lymphoma [4,44]. DuoHexaBody-CD37 has been previously reported to mediate cytotoxicity of DLBCL cells via different mechanisms of action: complement (CDC) and effector cell-mediated killing (ADCC/ADCP) [25,26]. Here, we report an additional mechanism of action of DuoHexaBody-CD37: CD37-mediated direct cytotoxicity which is independent of complement. Direct antibody-mediated killing may be particularly important in the DLBCL tumor microenvironment when complement is not available or inhibited. Complement has a dual role in tumors including hematological malignancies. The protective role of complement in antibody-mediated CDC has been well-established. However, more recent studies have provided evidence that tumors can hijack the complement components by cell surface expression of C1q, C3a and C5a receptors that promotes tumor growth and progression [45,46]. In addition, lymphoma cells can upregulate

complement regulatory proteins (CRPs: CD46, CD55 and CD59) to escape from antibody-

mediated CDC [47]. Although no direct correlation was found between DuoHexaBody-CD37-

induced CDC and expression of CRPs in vitro [26], expression of CD59 has been shown to

predict clinical outcome in DLBCL patients treated with R-CHOP [48]. In addition, the presence

of regulatory myeloid cells (tumor-associated neutrophils, myeloid-derived suppressor cells)

in the tumor microenvironment has been associated with immune suppression and poor

prognosis in lymphoma (reviewed in [37,49]). These tumor-infiltrating myeloid cells stimulate

tumor growth and may impair ADCC/ADCP activity of therapeutic antibodies [50]. Lastly, DLBCL

cells have been shown to upregulate CD47 to protect themselves from ADCC and ADCP [51].

Taken together, inhibition of complement activation by tumor cells or ineffective ADCC/ADCP

may limit efficacy of antibody-based immunotherapies for hematological malignancies.

Since decreased expression of CD20 is related to inferior clinical outcome of DLBCL

patients after treatment with R-CHOP, DuoHexaBody-CD37 treatment may be more potent

compared to CD20-targeting therapies in DLBCL. Supporting this, we observed no/minimal

internalization of CD37 by DuoHexaBody-CD37 in contrast to rituximab-induced CD20

internalization. We observed that DLBCL cells were more sensitive to DuoHexaBody-CD37-

mediated cytotoxicity than Burkitt cells, a difference that cannot be explained by variations in

CD37 surface expression or clustering. It is possible that Burkitt cells rely on other oncogenic

signaling pathways (i.e. MYC-driven) compared to DLBCL cells.

The activation of multiple intracellular signaling pathways, including PI3K-AKT and

RAS-MAPK that play a central role in cancer, was observed upon DuoHexaBody-CD37

treatment of DLBCL cells. This was dependent on Fc-crosslinking of DuoHexaBody-CD37,

which is in line with former studies using CD37-specific small modular

immunopharmaceuticals (CD37-SMIP) [52,53]. In vivo crosslinking could be facilitated by

effector cells expressing Fc receptors, as evidenced by the observed direct cytotoxicity of

DuoHexaBody-CD37 in tumor cells in the presence of PBMCs. This did not require FcγR-

mediated signaling or effector cell function (degranulation, trogocytosis) as fixed PBMCs

could be used to facilitate the crosslinking. CD37 can activate and inhibit AKT kinase through

its two intracellular tails that contain ITIM-like and ITAM-like motifs [27]. Different mechanisms

of direct tumor cell death by antibodies have been reported, including apoptosis, necrosis,

pyroptosis and necroptosis, and future studies are required to identify which of these

mechanisms underlie DuoHexaBody-CD37-induced killing of DLBCL cells. Interestingly,

increased p-AKT was observed in primary B cells upon DuoHexaBody-CD37 treatment in

contrast to malignant B cells, indicating that primary B cells may be less sensitive to direct

killing than lymphoma cells. Our results show that DuoHexaBody-CD37 directly affects

membrane-proximal signaling proteins (AKT and SHP-1) which provides new opportunities

for combination therapies in DLBCL [54]. For example, studies combining CD37-targeting

antibodies with PI3K inhibitors [55,56] or cell cycle kinase inhibitors [57] demonstrated enhanced

tumor cell death. We also observed p-BTK and p-PLCy upregulation in ABC-DLBCL upon

DuoHexaBody-CD37 treatment, which could provide a rational to combine DuoHexaBody-

CD37 with ibrutinib in ABC-DLBCL where BTK inhibition is already more effective compared to

GBC-DLBCL [58,59].

Moreover, DuoHexaBody-CD37 was found to be particularly effective in

downregulating p-STAT3 and p-STAT6 in presence of IL-6, IL-21 and IL-4, respectively. Within

the DLBCL tumor microenvironment, these cytokines actively contribute to cancer

pathogenesis in ABC-DLBCL [18,37,38,39,40]. IL-6 levels in serum are associated with adverse

clinical outcome in DLBCL, and we previously reported that CD37 inhibits IL-6 receptor

signaling through SOCS3 [17]. In line with this, patients with CD37-negative DLBCL present with

higher IL-6 levels in serum and tumors which is associated with inferior clinical outcome [17].

CD37 is an independent prognostic factor in DLBCL, irrespective of DLBCL subset [4], which has

been confirmed in patients with follicular lymphoma [44]. We anticipate that targeting CD37,

especially in the presence of classical chemotherapy and/or targeted drugs will further

enhance the efficacy of immunotherapy of DLBCL. In conclusion, this study shows

DuoHexaBody-CD37 induces cytotoxic signaling in DLBCL cells, and reduces cytokine-

mediated pro-survival signaling in the DLBCL tumor microenvironment.

Material and Methods

Therapeutic antibodies

DuoHexaBody-CD37 (GEN3009) and the negative control anti-HIV-1 gp120 antibody IgG1-b12 (mentioned in manuscript as IgG1-ctrl) were generated by Genmab (Utrecht, The Netherlands) as previously described [60]. Rituximab anti-CD20 (Mab Thera®) was obtained from Radboudumc (Nijmegen, the Netherlands).

Cell culture

Lymphoma cell lines (GCB-DLBCL: Oci-Ly8, Oci-Ly7; ABC-DLBC: HBL-1, U2932; B-ALL: NALM6, and Burkitt: BJAB, DAUDI) were obtained from DSMZ, ATCC and Blanca Scheijen (Dept. Pathology, Radboudumc) and cell lines were authenticated using STR-analysis. Cells were cultured in RPMI 1640 (Gibco), 1% antibiotic-antimycotic (Gibco) and 10% FBS (Hyclone) (RPMI/10%FCS) and incubated at 37°C with 5% CO_2. For optimum growth, the cell lines were propagated in dilution of 0.5×10^6 cells/ml and the culture media were refreshed twice a week. All experiments were performed one day after refreshing medium and expanding cell culture.

MTS assay

Cell viability was determined using CellTiter 96® AQueous One Solution Reagent from Promega as per manufacturer instructions. Briefly, 2×10^5 cells in RPMI/10%FCS were plated in 96-well plates and treated with 5μg/ml DuoHexaBody-CD37 for 30min at 4°C. Cells were washed and treated with 5μg/ml of goat F(ab')2 anti-human IgG (a-Fc, Southern Biotech) and incubated for 48h at 37°C with 5% CO_2. As control, cells treated with DuoHexaBody-CD37 or a-Fc alone were included. Subsequently, cells were stained with 1:5 dilution of

CellTiter 96® AQueous One Solution Cell Proliferation Assay (MTS, Promega) in media

(RPMI/10%FCS) and incubated at 37°C with 5% CO_2 for 2-3h. Absorbance was recorded at

490nm in a 96-well plate reader (Bio-Rad iMark Microplate Absorbance Reader).

Cytotoxicity assays

Peripheral blood from healthy volunteers was obtained via Sanquin blood bank (Nijmegen,

NL) upon informed consent and anonymized for further use, following the guidelines of the

Institutional Review Board, and in accordance with the declaration of Helsinki. Peripheral

blood mononuclear cells (PBMCs) were isolated using Ficoll Hypaque (GE Healthcare, Little

Chalfont, UK) according to the manufacturer's instructions. For fixation, PBMCs were washed

with PBS and resuspended directly in 1%PFA (filter sterilized) at 7.5×10^6 cells/ml for 20min at

RT (room temperature). Following, cells were washed twice in PBS and resuspended in media

(RPMI/10%FCS). In parallel, 10×10^6 target cells (HBL-1 or U2932) were stained with CellTrace

Violet Cell Proliferation Kit (Invitrogen) at final concentration of 167nM for 20min at 37°C

with 5% CO_2. Subsequently, cells were washed twice and resuspended in media

(RPMI/10%FCS). Next, CTV-stained target cells were treated with 5μg/ml DuoHexaBody-

CD37 for 30min at 4°C and washed with media. In addition, target cells were co-cultured

with fixed PBMCs at 1:20 (target to crosslinker) ratio for 4, 24 or 48h at 37°C in the presence

of 5% CO_2. Subsequently, cells were directly stained with 5μg/ml propidium iodide (Miltenyi

Biotec) for 5min at RT on plate shaker and analyzed using FACS Lyric (BD Biosciences).

Target binding analysis

Briefly, 100,000 lymphoma cells (HBL-1, Oci-Ly7, U2932, BJAB or DAUDI) were treated with

5mg/ml of anti-CD20 Rituximab (Mab Thera®) or DuoHexaBody-CD37 in RPMI/0.2%BSA for

15min at RT in a 96-wells plate. Plates were then directly transferred to incubator (37°C/5%

CO_2) for 0, 2 or 6h. Cells were harvested at respective time points by washing with PBA buffer

(PBS, 1% BSA, azide 0.02%) followed by staining with mouse anti-human IgG1 Fc secondary

antibody conjugated to Alexa Fluor 488 (ThermoFisher) for 30min at 4°C. Next, cells were

washed and stained with Live-dead marker (eFluor 506, eBioscience) and measured using

FACS Lyric (BD Biosciences).

Generation of CD37 mutant cell lines

NALM6 cells were used because of their endogenous low CD37 expression. Cells were grown

in RPMI/10% FBS supplemented with glutamine 2mM. Cells were transfected with either

CD37-sGFP2, CD37 Y13F-sGFP2 or CD37-delta Y13-sGFP2 constructs using the Amaxa 4D

nucleofector. Briefly, for each transfection $3x10^6$ cells were collected and washed with PBS.

Cells were resuspended in 100 µl SF buffer (cat. no. V4XC-2024) containing 3 µg of plasmid

and pulsed with program CV-104. After transfection, cells were transferred to 6-well plates

and incubated overnight in 3 ml of RPMI with 10% FBS without antibiotic-antimycotic and

phenol red. The CD37 constructs were kindly provided by Prof. Dr. N. Muthusamy (Ohio

University, USA) [27], and subcloned in pSGFP-N1 vector using standard techniques. Sequences

were verified via sequencing.

Phosphoflow assays

Lymphoma cells ($2x10^5$ cells/well) or PBMCs ($5x10^5$ cells/well) in RPMI/10%FCS were treated

with 5µg/ml DuoHexaBody-CD37 for 30min at 4°C. Next, cells were washed and treated with

5µg/ml of goat F(ab')2 anti-human IgG (a-Fc) and incubated for 20min at 37°C with 5% CO_2.

As treatment control, cells were treated with DuoHexaBody-CD37 or a-Fc alone. As positive

control of B-cell receptor stimulation, cells were treated with 20μg/ml of goat F(ab')2 anti-

human IgM. Next, cells were spun down and stained at 1:10 dilution with Fixable Viability

stain 510 (BD Biosciences) in RPMI/10%FCS for 10min at 37°C. Thereafter, cells were directly

fixed for 10min at 37°C with 100 μl of the eBioscience FoxP3/Transcription Factor Fixation kit

(Invitrogen) as indicated by the manufacturer. After fixation, samples were washed twice

with eBioscience Permeabilization Wash Buffer (10X diluted in Milli-Q) (Invitrogen). To

identify the B cells within PBMC cultures, cells were stained with antibodies against CD19,

CD3 and IgD diluted in eBioscience wash buffer. To prevent non-specific antibody binding,

FcR-blocking purified rat anti-mouse CD16/CD32 (Mouse BD Fc BlockTM) was included in the

staining. Stained cells were incubated for 30min at 4°C followed by washing in eBioscience

wash buffer and staining with antibodies against phosphorylated proteins; p-SHP1(Y564), p-

AKT(S473), p38(T180/Y182) (all Cell signaling technology), p-SYK(Y348), p-BTK(Y223), p-

PLCγ2(Y759), p-PKCα/βII(T638/641), p-ERK-1/2(T202/Y204) (all BD Biosciences); in wash

buffer for 30min at RT. Unlabeled phospho-antibodies (p-SHP1(Y564), p-AKT(S473),

p38(T180/Y182)) were stained with a secondary PE-labeled F(ab')2 donkey anti-rabbit IgG

(H+L) antibody (Jackson Immuno Research) for 15min at RT. Lastly, the cells were washed and

resuspended in wash buffer before measuring on FACS Lyric.

Cytokine stimulation experiments

To measure p-STAT3 and p-STAT6 responses, cells treated with DuoHexaBody-CD37 and/or

goat F(ab')2 anti-human IgG (a-Fc) for 24h as described above. After washing, cells were

stimulated with recombinant IL-6 (100ng/ml) or IL-4 (20ng/ml) (both from Miltenyi) or IL-21

(Peprotech) (50ng/ml) for 15min at 37°C. Cells were harvested and stained with Fixable

Viability stain 510 (BD Biosciences). Subsequently, cells were fixed with BD Cytofix Fixation

Buffer (BD Biosciences) for 10min at 37°C and washed twice with MACS buffer (PBS/0.5%

BSA/2 mM EDTA). Next, cells were permeabilized using 150 µl Perm Buffer III (BD

Biosciences) at 4°C for 30min and washed twice with MACS buffer, followed by staining with

PE-labeled p-STAT3 (Y705) or p-STAT6 (Y641) (BD Biosciences) antibody diluted (1:8) in MACS

buffer for 30min at RT. Finally, the cells were washed and resuspended in 50 µl MACS buffer

before measuring on FACS Lyric.

Supplementary methods: Microscopy analysis of CD37 clustering and RPPA assay

Acknowledgements

We thank the Radboudumc Microscopy Imaging Center for use of their microscopy facilities, as well as for their support and assistance. We thank Simone Oostindie and Inge Verbrugge for valuable scientific input, and Peter Friedl for facilitating the RPPA studies. The CD37 constructs were kindly provided by Natarajan Muthusamy (Ohio University, USA) [27].

Disclosures

MGR, MBO, KCMS and ECWB are Genmab employees and own Genmab warrants and/or stock. All the other authors have no conflicts of interest to disclose. This research study was funded by Genmab, and ABvS is supported by the Netherlands Organization for Scientific Research (NWO): the Institute of Chemical Immunology (project ICI00023), ZonMW (project 09120012010023), the Dutch Cancer Society (projects 12949 and 14726), and the European Research Council: Consolidator Grant (project 724281) and Proof-of-Concept Grant (project 101112687).

Author contributions

ABvS, SPS, ECWB, MGR, KCMS and MBO provided scientific input and conceptualized the study. SPS developed the methodology and performed the data analysis for the cytotoxicity assays, PBMC assays, phosphoflow experiments, CD37 mutant studies, cytokine studies and RPPA analysis. MDvdB performed the surface accessibility assays and CD37 mutant studies. SvD developed the methodology for the CD37 transfections. MTB developed methodology of CD37 microscopy studies and performed data analysis on CD37 clustering. SPS and ABvS

450   wrote the manuscript. ABvS supervised the study. All authors read, revised, and agreed on

451   the final version of the manuscript.

Figure legends

Figure 1: DuoHexaBody-CD37-induces direct cytotoxicity in DLBCL-derived tumor cell lines

(A) Percentage decrease (mean +/- SD) in viability upon DuoHexaBody-CD37 treatment in presence or absence of goat F(ab')2 anti-human IgG (a-Fc) in indicated cell lines after 48h. Data is shown from at least three independent experiments. Significance was calculated compared to untreated control using Kruskal-Wallis test followed by Dunn' multiple comparison test correction (**$p < 0.01$) **(B)** Percentage increase in cell death (PI-positive) upon co-culturing healthy donor-derived fixed PBMCs with DuoHexaBody-CD37 pre-treated HBL-1 and U2932 cells compared to untreated cells for indicated time points. Duplicates from each individual donor (n=4) are indicated in separate colors. Significance was calculated compared to 4h treated cells using Kruskal-Wallis test followed by Dunn' multiple comparison test correction (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Figure 2: DuoHexaBody-CD37 induces CD37 clustering without modulating CD37 cell surface expression

(A,B) Airyscan images depicting clustering (fluorescence/area) upon DuoHexaBody-CD37 (DHB) treatment or B12 isotype control in BJAB **(A)** and Oci-Ly8 **(B)** cells. Data shown from three independent experiments. Bar is 10 μm. Significance was calculated comparing DuoHexaBody-CD37 treated cells to isotype control antibody using unpaired T-test (****$p < 0.0001$). **(C)** Percentage decrease (mean +/- SD) in cell surface binding by DuoHexaBody-CD37 or rituximab in indicated cell lines as measured by flow cytometry. Data is shown from at least three independent experiments. Significance was calculated comparing respective time point using unpaired T-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Figure 3: DuoHexaBody-CD37 treatment results in activation of different downstream signaling pathways

(**A**) Heatmap depicting log-normalized signal intensities of 26 phosphoproteins with two-fold increase in signal upon DuoHexaBody-CD37 treatment in presence or absence of goat F(ab')2 anti-human IgG (a-Fc) in primary B cells, U2932 and Oci-Ly7. (**B**) Oncogenic hallmark signatures obtained from the MSigDB enriched using 26 phosphoproteins with two-fold increase in signal upon DuoHexaBody-CD37 treatment in presence or absence of goat F(ab')2 anti-human IgG (a-Fc) in primary B cells, U2932 and Oci-Ly7. (**C**) Phosphoflow analysis of p-SYK(Y348), p-SHP1(Y564), p-AKT(S473) on primary B cells (*top row*) or DLBCL cells (*bottom row*) either untreated or treated with anti-BCR (F(ab')2 anti-IgM), DuoHexaBody-CD37 and/or goat F(ab')2 anti-human IgG (a-Fc). Dot plots depict quantification (mean +/- SD) of mean fluorescence intensity (MFI). Each dot represents individual donor (B cells) or experimental replicate (DLBCL). Significance was calculated compared to untreated control using Kruskal-Wallis test followed by Dunn' multiple comparison test correction (**$p<0.01$, ***$p<0.001$, ****$p<0.0001$).

Figure 4: DuoHexaBody-CD37 treatment results in differential activation of BCR and RAS/MAPK downstream signaling proteins in primary B cells and DLBCL

(**A, B**) Phosphoflow analysis of (A) p-BTK(Y223), p-PLCy2(Y759), p-PKCα/βII(T638/641) and (B) p38(T180/Y182), p-ERK-1/2(T202/Y204) on primary B cells (*top row*) or DLBCL cells (*bottom row*) either untreated or treated with anti-BCR (F(ab')2 anti-IgM), DuoHexaBody-CD37 and/or goat F(ab')2 anti-human IgG (a-Fc). Dot plots depict quantification (mean +/- SD) of mean fluorescence intensity (MFI). Each dot represents an individual donor (primary B cells) or experimental replicate (DLBCL tumor cell lines). Significance was calculated compared to

untreated control using Kruskal-Wallis test followed by Dunn' multiple comparison test correction (*p<0.05, **p<0.01, ****p<0.0001).

Figure 5: CD37 N-terminus is involved in DuoHexaBody-CD37-mediated cytotoxic signaling

(**A**) Histogram showing CD37 cell surface expression determined by flow cytometry analysis of NALM6 cell lines expressing wild type CD37 (CD37-WT), or mutations of Tyr13 to phenylalanine (CD37-Y13F), or deletion of Tyr13 (CD37-ΔY13) in the cytosolic regions of the CD37 molecule. (**B**) p-AKT(S473) and (**C**) p-SHP1(Y564) on NALM6 cell lines transfected with CD37-WT, CD37-Y13F or CD37-ΔY13 treated with DuoHexaBody-CD37 (DHB) and goat F(ab')2 anti-human IgG (a-Fc) or crosslinker alone (a-Fc). Dot plots depict p-AKT and p-SHP1 levels in treated (DHB+a-Fc or a-Fc) vs untreated cells in CD37-GFP-positive (transfected) cells using phosphoflow analysis, corrected for background signal from GFP-negative cells. Significance was calculated using Student t-test (*p<0.05). Data shown from three independent experiments.

Figure 6: DuoHexaBody-CD37 treatment decreases cytokine-mediated pro-survival signaling in DLBCL cell lines

(**A**) Percentages of p-STAT6-positive cells (mean and SD) upon recombinant human IL-4 (rh-IL-4) stimulation in DLBCL cells pre-treated with DuoHexaBody-CD37 in presence or absence of goat F(ab')2 anti-human IgG (a-Fc). Data is shown from at least three independent experiments. Percentages of p-STAT3-positive cells (mean +/- SD) upon (**B**) recombinant human IL-6 (rh-IL-6) stimulation or (**C**) recombinant human IL-21 (rh-IL-21) stimulation in DLBCL cells pre-treated with DuoHexaBody-CD37 in presence or absence of a-Fc. Data is shown from at least three independent experiments. (**A-C**) Significance was calculated

compared to untreated control using Kruskal-Wallis test followed by Dunn' multiple

comparison test correction (**p<0.01).

Supplementary Figure 1: Principal component analysis of (phospho)protein expression

profiles after DuoHexaBody-CD37 treatment

Principal component analysis (PCA) of 484 proteins (102 phospho targets and 382 total

proteins) in primary B cells (CD19+ purified) and two DLBCL cell lines (Oci-Ly7 and U2932)

either untreated or treated with DuoHexaBody-CD37 in presence or absence of a-Fc using

reverse phase protein array (RPPA). The first two components of the cluster analysis are

shown.

Supplementary Figure 2: DuoHexaBody-CD37 treatment results in differential activation of

BCR and RAS/MAPK downstream signaling proteins in B cells and DLBCL

(A-C) Phosphoflow analysis of (A) p-SYK(Y348), p-SHP1(Y564), p-AKT(S473); **(B)** p-BTK(Y223),

p-PLCγ2(Y759), p-PKCα/βII(T638/641) and **(C)** p38(T180/Y182), p-ERK-1/2(T202/Y204) on

primary B cells (*top row*) or DLBCL cells (*bottom row*) either anti-IgM treated or treated with

DuoHexaBody-CD37 and/or goat F(ab')2 anti-human IgG (a-Fc). Dot plots depict fold change

(mean +/- SD) of mean fluorescence intensity (MFI) of indicated treatment to untreated

cells. Each dot represents individual donor (B cells) or experimental replicate (DLBCL).

Significance of fold change to untreated control was calculated using Wilcoxon Signed Rank

test (*p<0.05, **p<0.01, ****p<0.0001).

References

1. Morin RD, Arthur SE, Hodson DJ. Molecular profiling in diffuse large B-cell lymphoma: why so many types of subtypes? Br J Haematol. 2022;196(4):814–29.

2. Abou Dalle I, Dulery R, Moukalled N, Ricard L, Stocker N, El-Cheikh J, et al. Bi- and Tri-specific antibodies in non-Hodgkin lymphoma: current data and perspectives. Blood Cancer J. 2024;14(1).

3. Johnson NA, Boyle M, Bashashati A, Leach S, Brooks-Wilson A, Sehn LH, et al. Diffuse large B-cell lymphoma: Reduced CD20 expression is associated with an inferior survival. Blood. 2009;113(16):3773–80.

4. Xu-Monette ZY, Li L, Byrd JC, Jabbar KJ, Manyam GC, De Winde CM, et al. Assessment of CD37 B-cell antigen and cell of origin significantly improves risk prediction in diffuse large B-cell lymphoma. Blood. 2016;128(26):3083–100.

5. Guan XW, Wang HQ, Ban WW, Chang Z, Chen HZ, Jia L, et al. Novel HDAC inhibitor Chidamide synergizes with Rituximab to inhibit diffuse large B-cell lymphoma tumour growth by upregulating CD20. Cell Death Dis. 2020;11(1).

6. Roschewski M, Staudt LM, Wilson WH. Diffuse large B-cell lymphoma—treatment approaches in the molecular era. Nat Rev Clin Oncol. 2013;11(1):12–23.

7. Varma G, Goldstein J, Advani RH. Novel agents in relapsed/refractory diffuse large B-cell lymphoma. Hematol Oncol. 2023;41(S1):92–106.

8. Schwartz-Albiez R, Dörken B, Hofmann W, Moldenhauer G. The B cell-associated CD37 antigen (gp40-52). Structure and subcellular expression of an extensively glycosylated glycoprotein. J Immunol 1988;140(3):905–14.

9. Van Spriel AB, De Keijzer S, Van Der Schaaf A, Gartlan KH, Sofi M, Light A, et al. The tetraspanin CD37 orchestrates the α4β1 integrin-Akt signaling axis and supports long-

lived plasma cell survival. Sci Signal. 2012 ;5(250):ra82.

10.   de Winde CM, Zuidscherwoude M, Vasaturo A, van der Schaaf A, Figdor CG, van Spriel

AB. Multispectral imaging reveals the tissue distribution of tetraspanins in human

lymphoid organs. Histochem Cell Biol. 2015; 144(2):133-46.

11.   Berditchevski F, Rubinstein E. Tetraspanins. 2013. 1–418 p.

http://link.springer.com/10.1007/978-94-007-6070-7

12.   Hemler ME. Targeting of tetraspanin proteins — potential benefits and strategies. Nat

Rev Drug Discov. 2008;7(9):747–58.

13.   Querol Cano L, Dunlock VME, Schwerdtfeger F, van Spriel AB. Membrane organization

by tetraspanins and galectins shapes lymphocyte function. Nat Rev Immunol. 2024

;24(3):193–212.

14.   Barrena S, Almeida J, Yunta M, Lopez A, Fernandez-Mosteirin N, Giralt M, et al.

Aberrant expression of tetraspanin molecules in B-cell chronic lymphoproliferative

disorders and its correlation with normal B-cell maturation. Leukemia.

2005;19(8):1376–83.

15.   Bobrowicz M, Kubacz M, Slusarczyk A, Winiarska M. Cd37 in b cell derived tumors—

more than just a docking point for monoclonal antibodies. Int J Mol Sci. 2020

;21(24):1–16.

16.   Beckwith KA, Byrd JC, Muthusamy N. Tetraspanins as therapeutic targets in

hematological malignancy: A concise review. Front Physiol. 2015 Mar 23;6.

17.   De Winde CM, Veenbergen S, Young KH, Xu-Monette ZY, Wang XX, Xia Y, et al.

Tetraspanin CD37 protects against the development of B cell lymphoma. J Clin Invest.

2016;126(2):653–66.

18.   Hashwah H, Bertram K, Stirm K, Stelling A, Wu C, Kasser S, et al. The IL-6 signaling

complex is a critical driver, negative prognostic factor, and therapeutic target in

diffuse large B-cell lymphoma . EMBO Mol Med. 2019;11(10).

19. Deckert J, Park PU, Chicklas S, Yi Y, Li M, Lai KC, et al. A novel anti-CD37 antibody-drug

conjugate with multiple anti-tumor mechanisms for the treatment of B-cell

malignancies. Blood. 2013;122(20):3500–10.

20. Beckwith KA, Frissora FW, Stefanovski MR, Towns WH, Cheney C, Mo X, et al. The

CD37-targeted antibody-drug conjugate IMGN529 is highly active against human CLL

and in a novel CD37 transgenic murine leukemia model. Leukemia. 2014; 144(2):133-

46.

21. Köksal H, Dillard P, Josefsson SE, Maggadottir SM, Pollmann S, Fane A, et al. Preclinical

development of CD37CAR T-cell therapy for treatment of B-cell lymphoma. Blood Adv.

2019;3(8):1230–43.

22. Imai K, Takeuchi Y, Terakura S, Okuno S, Adachi Y, Osaki M, et al. Dual CAR-T Cells

Targeting CD19 and CD37 Are Effective in Target Antigen Loss B-cell Tumor Models.

Mol Cancer Ther. 2024;23(3):381–93.

23. Caulier B, Joaquina S, Gelebart P, Dowling TH, Kaveh F, Thomas M, et al. CD37 is a safe

chimeric antigen receptor target to treat acute myeloid leukemia. Cell Reports Med.

2024;5(6).

24. Witkowska M, Smolewski P, Robak T. Investigational therapies targeting CD37 for the

treatment of B-cell lymphoid malignancies. Expert Opin Investig Drugs.

2018;27(2):171–7.

25. Oostindie SC, Van Der Horst HJ, Kil LP, Strumane K, Overdijk MB, Van Den Brink EN, et

al. DuoHexaBody-CD37 ® , a novel biparatopic CD37 antibody with enhanced Fc-

mediated hexamerization as a potential therapy for B-cell malignancies. Blood Cancer

J. 2020;10:30.

26.   van der Horst HJ, Oostindie SC, Cillessen SAGM, Gelderloos AT, Overdijk MB, Nijhof IS,

621   et al. Potent Preclinical Efficacy of DuoHexaBody-CD37 in B-Cell Malignancies.

622   HemaSphere. 2021;5(1):e504.

623   27.   Lapalombella R, Yeh YY, Wang L, Ramanunni A, Rafiq S, Jha S, et al. Tetraspanin CD37

624   directly mediates transduction of survival and apoptotic signals. Cancer Cell.

625   2012;21(5):694–708.

626   28.   Fujimura T, Yamashita-Kashima Y, Kawasaki N, Yoshiura S, Harada N, Yoshimura Y.

627   Obinutuzumab in combination with chemotherapy enhances direct cell death in cd20-

628   positive obinutuzumab-resistant non-hodgkin lymphoma cells. Mol Cancer Ther. 2021

629   ;20(6):1133–41.

630   29.   Mone AP, Cheney C, Banks AL, Tridandapani S, Mehter N, Guster S, et al.

631   Alemtuzumab induces caspase-independent cell death in human chronic lymphocytic

632   leukemia cells through a lipid raft-dependent mechanism. Leukemia. 2006;20:272–9.

633   30.   Zhao X, Lapalombella R, Joshi T, Cheney C, Gowda A, Hayden-Ledbetter MS, et al.

634   Targeting CD37-positive lymphoid malignancies with a novel engineered small

635   modular immunopharmaceutical. Blood 2007;110(7):2569–77.

636   31.   Beers SA, French RR, Chan HTC, Lim SH, Jarrett TC, Vidal RM, et al. Antigenic

637   modulation limits the efficacy of anti-CD20 antibodies: Implications for antibody

638   selection. Blood. 2010;115(25):5191–201.

639   32.   Lim SH, Vaughan AT, Ashton-Key M, Williams EL, Dixon S V., Chan HTC, et al. Fc

640   gamma receptor IIb on target B cells promotes rituximab internalization and reduces

641   clinical efficacy. Blood. 2011;118(9):2530–40.

33.   Whittaker KC, Huang RP. Antibody Arrays Methods and Protocols. 2021. 45–54 p.

Available from: http://www.springer.com/series/7651

34. Pal Singh S, Dammeijer F, Hendriks RW. Role of Bruton's tyrosine kinase in B cells and

malignancies. Mol Cancer. 2018;17(1):1–23.

35. Benoit A, Bou-Petit E, Chou H, Lu M, Guilbert C, Luo VM, et al. Mutated RAS-

associating proteins and ERK activation in relapse/refractory diffuse large B cell

lymphoma. Sci Rep. 2022;12(1).

36. Uddin S, Hussain AR, Siraj AK, Manogaran PS, Al-Jomah NA, Moorji A, et al. Role of

phosphatidylinositol 3'-kinase/AKT pathway in diffuse large B-cell lymphoma survival.

Blood. 2006;108(13):4178–86.

37. Höpken UE, Rehm A. Targeting the Tumor Microenvironment of Leukemia and

Lymphoma. Trends in Cancer. 2019;5(6):351–64.

38. Wang Y, Wang C, Cai X, Mou C, Cui X, Zhang Y, et al. IL-21 Stimulates the expression

and activation of cell cycle regulators and promotes cell proliferation in EBV-positive

diffuse large B cell lymphoma. Sci Rep. 2020;10(1):1–13.

39. Jones ELL, Demaria MCC, Wright MDD. Tetraspanins in cellular immunity. Biochem Soc

Trans. 2011;39(2):506–11.

40. Wood B, Sikdar S, Choi SJ, Virk S, Alhejaily A, Baetz T, et al. Leukemia & Lymphoma

Abundant expression of interleukin-21 receptor in follicular lymphoma cells is

associated with more aggressive disease Abundant expression of interleukin-21

receptor in follicular lymphoma cells is associated with more aggressive disease. Leuk

Lymphoma. 2013;54(6):1212–20.

41. Zöller M. Tetraspanins: Push and pull in suppressing and promoting metastasis. Nat

Rev Cancer. 2009;9(1):40–55.

42. Kumari S, Devi G V, Badana anil, ramesh Dasari V, rao malla rama. CD151-A Striking

Marker for Cancer Therapy. Biomark Cancer. 2015;(7):7–7.

43. Vences-Catalán F, Kuo CC, Rajapaksa R, Duault C, Andor N, Czerwinski DK, et al. CD81

is a novel immunotherapeutic target for B cell lymphoma. J Exp Med. 2019

;jem.20190186.

44. Yoshimura T, Miyoshi H, Shimono J, Nakashima K, Takeuchi M, Yanagida E, et al. CD37

expression in follicular lymphoma. Ann Hematol. 2022;101(5):1067–75.

45. Roumenina LT, Daugan M V., Noe R, Petitprez F, Vano YA, Sanchez-Salas R, et al.

Tumor cells hijack macrophage-produced complement C1q to promote tumor growth.

Cancer Immunol Res. 2019;7(7):1091–105.

46. Nemenoff RA, Markiewski MM. The role of complement in the tumor

microenvironment Fac Rev. 2021;10(80).

47. Luo S, Wang M, Wang H, Hu D, Zipfel PF, Hu Y. How Does Complement Affect

Hematological Malignancies: From Basic Mechanisms to Clinical Application. Front

Immunol. 2020;11.

48. Song G, Cho WC, Gu L, He B, Pan Y, Wang S. Increased CD59 protein expression is

associated with the outcome of patients with diffuse large B-cell lymphoma treated

with R-CHOP. Med Oncol. 2014;31(7).

49. Fowler NH, Cheah CY, Gascoyne RD, Gribben J, Neelapu SS, Ghia P, et al. Role of the

tumor microenvironment in mature B-cell lymphoid malignancies. Haematologica.

2016;101(5):531–40.

50. Singhal S, Rao AS, Stadanlick J, Bruns K, Sullivan NT, Bermudez A, et al. Human Tumor–

Associated Macrophages and Neutrophils Regulate Antitumor Antibody Efficacy

through Lethal and Sublethal Trogocytosis. Cancer Res. 2024 ;84(7):1029–47.

51. Bouwstra R, He Y, De Boer J, Kooistra H, Cendrowicz E, Fehrmann RSN, et al. CD47

expression defines efficacy of rituximab with CHOP in non–germinal center B-cell

(Non-GCB) diffuse large B-cell lymphoma patients (DLBCL), but not in GCB DLBCL.

Cancer Immunol Res. 2019 ;7(10):1663–71.

52. Zhao X, Lapalombella R, Joshi T, Cheney C, Gowda A, Hayden-Ledbetter MS, et al.

Targeting CD37-positive lymphoid malignancies with a novel engineered small

modular immunopharmaceutical. Blood. 2007;110(7):2569–77.

53. Rafiq S, Siadak A, Butchar JP, Cheney C, Lozanski G, Jacob NK, et al. Glycovariant anti-

CD37 monospecific protein therapeutic exhibits enhanced effector cell-mediated

cytotoxicity against chronic and acute B cell malignancies. MAbs. 2013;5(5):723–35.

54. Wang J, Xu-Monette ZY, Jabbar KJ, Shen Q, Manyam GC, Tzankov A, et al. AKT

Hyperactivation and the Potential of AKT-Targeted Therapy in Diffuse Large B-Cell

Lymphoma. Am J Pathol 2017;187:1700–16.

55. Arribas A, Napoli S, Munz N, Stussi G, Bertoni F. PI3Kδ activation, IL6 over-expression,

and CD37 loss cause resistance to naratuximab emtansine in lymphomas. Blood Adv.

2024; bloodadvances.2023012291

56. Krause G, Baki I, Kerwien S, Knödgen E, Neumann L, Göckeritz E, et al. Cytotoxicity of

the CD37 antibody BI 836826 against chronic lymphocytic leukaemia cells in

combination with chemotherapeutic agents or PI3K inhibitors. Br J Haematol.

2016;173(5):791–4.

57. Rødland GE, Melhus K, Generalov R, Gilani S, Bertoni F, Dahle J, et al. The Dual Cell

Cycle Kinase Inhibitor JNJ-7706621 Reverses Resistance to CD37-Targeted

Radioimmunotherapy in Activated B Cell Like Diffuse Large B Cell Lymphoma Cell

Lines. Front Oncol. 2019;9:1301.

58. Wilson WH, Young RM, Schmitz R, Yang Y, Pittaluga S, Wright G, et al. Targeting B cell

receptor signaling with ibrutinib in diffuse large B cell lymphoma. Nat Med.

2015;21(8).

59.   Wilson WH, Wright GW, Huang DW, Hodkinson B, Balasubramanian S, Fan Y, et al.

Effect of ibrutinib with R-CHOP chemotherapy in genetic subtypes of DLBCL. Cancer

Cell. 2021;39(12):1643-1653.e3.

60.   Oostindie SC, Taylor RP, Van Der Horst HJ, Lindorfer MA, Cook EM, Tupitza JC, et al.

CD20 and CD37 antibodies synergize to activate complement by Fc-mediated

clustering. Haematologica. 2019 ;104(9):1841.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

B

Figure 6

A

B

C