Maternal Transfer of Oral Vaccine Induced Anti-OspA Antibodies Protects *Peromyscus*

spp Pups from Tick-Transmitted *Borrelia burgdorferi*

Jose F. Azevedo[1], Greg Joyner[1], Suman Kundu[1], Kamalika Samanta[1], and Maria Gomes-Solecki[1]

[1]University of Tennessee Health Science Center, Department of Microbiology, Immunology, and

Biochemistry

Correspondence to: Maria Gomes-Solecki, mgomesso@uthsc.edu

Abstract

The efficacy and duration of passive immunity protection depends on maternal antibody levels and transfer efficiency. We investigated whether oral vaccination of *Peromyscus leucopus* dams with recombinant OspA-expressing *E. coli* could induce maternal transfer of anti-OspA antibodies and protect pups from *Borrelia burgdorferi* challenge. Dams were vaccinated until breeding pairs were created (i), until parturition (ii), and until pups were 2 weeks old (iii). Pups were challenged with nymphal *I. scapularis*-transmitted *B. burgdorferi* at ~ 4 weeks of age. Anti-OspA IgG were quantified in dams and pups, and anti-*B. burgdorferi* IgG were quantified in pups. *B. burgdorferi* burden was assessed by *flaB* qPCR in pups' tissues ~ 4 weeks after tick challenge and viability of *B. burgdorferi* was assessed by culture of heart tissue. *P. leucopus* pups born to dams vaccinated until breeding had low serologic anti-OspA antibody and were not protected from tick transmitted *B. burgdorferi* infection. However, when dams' vaccination extended until parturition and until pups were two weeks old, significant anti-OspA antibody transfer and protection from *B. burgdorferi* infection occurred. This was evidenced by absence of antibody to *B. burgdorferi* PepVF, absence of *B. burgdorferi flaB* DNA in heart and bladder tissues, and absence of *flaB* in culture from heart tissues from pups euthanized >9 weeks after birth. We show that transfer of anti-OspA antibodies from vaccinated *P. leucopus* dams to offspring prevents tick transmission and infection dynamics of *B. burgdorferi* in the major reservoir host of this spirochete in the United States.

Importance: This study contributes to our understanding of how interventions based in reservoir

targeted OspA vaccines designed to block transmission of *B. burgdorferi* from infected *I.*

scapularis ticks may disrupt the enzootic cycle of this spirochete and reduce incidence of Lyme

disease.

Key words: *Peromyscus*, reservoir targeted oral vaccine, transmission blocking, *Ixodes*

scapularis, enzootic cycle, OspA, mouse, *Borrelia burgdorferi*, Lyme disease

Introduction

Lyme disease, caused by *Borrelia burgdorferi*, is the most prevalent vector-borne disease in North America (1). The white-footed mouse (*Peromyscus leucopus*) serves as the primary reservoir host (2), maintaining *B. burgdorferi* in enzootic cycles and facilitating transmission to *Ixodes* ticks that subsequently infect humans and other mammals (3). Unlike humans, reservoir hosts do not develop significant disease following *B. burgdorferi* infection, yet they serve as a crucial source of spirochete acquisition for larval Ixodid ticks (4). Thus, interrupting transmission of *B. burgdorferi* (5) has been proposed as a strategy for controlling Lyme disease (6, 7).

Outer surface protein A (OspA) is a major target for Lyme disease vaccine development (8) due to its essential role in *B. burgdorferi* persistence within the tick vector (9). OspA facilitates spirochete adherence to the tick midgut, allowing the bacterium to survive between blood meals (10). During tick feeding, *B. burgdorferi* downregulates OspA while upregulating OspC (11), which enables migration to the salivary glands and subsequent transmission to the host (12-14). Vaccines targeting OspA aim to elicit antibodies that neutralize spirochetes within the tick gut, preventing their transmission before they reach the mammalian host (15, 16). These vaccines have been explored in multiple contexts, including human vaccination (17, 18), direct vaccination of *P. leucopus* (19), and oral bait delivered reservoir-targeted vaccines (6).

Beyond direct vaccination, maternal transfer of antibody provides an alternative strategy to protect neonates during the critical early stages of life (20). In the context of ecology of vector-borne pathogens, this temporary immunity reduces early-life susceptibility to infection (21). Previous studies have demonstrated that oral vaccination of dams with OspA results in anti-

OspA IgG in offspring (22), though the degree to which this protects against *B. burgdorferi*

transmission remains unknown. In this study, we addressed this critical gap by investigating the

extent of maternal antibody transfer in *P. leucopus* and assessing whether vaccination of dams

with recombinant OspA-expressing *E. coli* confers protection to neonates. We tested whether

vaccination of dams until breeding pairs were created, until parturition and until the pups reached

2 weeks of age influences antibody transfer and protection against tick-mediated *B. burgdorferi*

challenge. Our findings have implications for reservoir-targeted vaccination strategies aimed at

reducing Lyme disease risk by breaking the transmission cycle of *B. burgdorferi*.

Materials and Methods

Ethics Statement. This study was conducted in accordance with the Guide for the Care and Use

of Laboratory Animals of the US National Institutes of Health. The protocol was approved by the

University of Tennessee Health Science Center Institutional Animal Care and Use Committee

(IACUC), protocol #22-0400. *Peromyscus leucopus* were sourced from the Peromyscus Genetic

Stock Center (PGSC) at the University of South Carolina.

Vaccine Preparation. Recombinant *E. coli* BL21(DE3)pLysS encoding *B. burgdorferi* OspA

(*E. coli*-OspA) was produced as previously described (23). Briefly, *E. coli*-OspA was grown in

Terrific Broth (TBY) at 37°C until OD600 ~0.8, induced with 1 mM IPTG for 3 h, transferred to

filter-top flasks and inactivated with 1% β-propiolactone (BPL, 98%, AlfaAesar), incubated at

room temperature at 100 rpm for 24h, and harvested by centrifugation at 2000×g at 4°C. The

bacterial biomass was resuspended in phosphate-buffered saline (PBS) and sprayed onto feed

pellets. Pellets (RTV pellets) were allowed to dry between application layers. Inactivation was

tested by inoculation of BPL-treated culture in TBY (1:200) and incubation overnight at 37°C,

shaking at 250 rpm. The following day, culture inactivation was confirmed by checking turbidity

and recording $OD_{600}=0$.

Vaccination Strategy. Dams (n=9) received RTV pellets daily for 5 consecutive days per week,

with a 2-day rest period over the weekend in which they received regular mouse chow. Control

dams received uncoated pellets. Dams consumed RTV pellets continuously as per vaccination

schedule up to 36 weeks (**Fig 1**) and were assigned to three experimental groups based on

breeding milestones. The level of IgG to OspA was analyzed in dams' blood before selecting

animals with high antibody in serum to create breeding pairs.

Breeding and Litter Management. Breeding pairs were established based on dams having anti-

OspA titers (OD450>0.8) in serum diluted 1:100 assessed by ELISA. Each vaccinated female

was cohoused with an unvaccinated male for 3–4 weeks, after which pregnant females were

single-housed until parturition (**Fig 1**). The first group (Breeding 1) was created with 5 dams

vaccinated for 12 weeks prior to breeding pairs were formed. The second group, Breeding 2, was

created with 4 dams vaccinated for 24 weeks until parturition. The third group, Breeding 3, was

created with the dams who maintained high levels of antibodies to OspA and remained on the

vaccination regimen for 36 weeks, ensuring that immunization continued until their pups reached

two weeks of age. Dams that lost the ability to reproduce or had low levels of serologic antibody

to OspA were euthanized. A group of pups (n=7) born after the 3[rd] breeding were euthanized

immediately after they dropped at parturition and blood was collected. Pups remained with their

dams until 24–25 days of age, at which point they underwent nymphal tick challenge.

Blood Collection. Blood samples were collected via submandibular vein bleed. Dams were sampled at day 0 and before each breeding. Pups were sampled on the day they were born, at day 24 (pre-challenge) and day ~71 (terminal bleed). Samples were allowed to clot for 30 min at room temperature before centrifugation at 2000×g for 10 min at 4°C. Serum was aliquoted and stored at −80°C until analysis.

Tissue Collection and Processing. At necropsy on d71 (~9-10 week old pups), heart and bladder tissues were harvested, transported on dry ice, and stored at −80°C. For organ culture experiments, freshly excised heart tissue was immediately placed into Barbour-Stoenner-Kelly H (BSK-H) medium supplemented with 6% rabbit serum (Sigma-Aldrich, St. Louis, MO) and incubated at 34°C for 3 weeks (on the 7th day of culture, the tissue was removed to avoid putrefaction). On day 21 of culture supernatants were analyzed for motile *B. burgdorferi* detection using Petroff-Hausser chamber (dark-field microscopy, 200× magnification) and for DNA quantification via *flaB* qPCR.

Tick Challenge and Monitoring of *B. burgdorferi* Infection. To assess susceptibility to *B. burgdorferi* infection, pups were challenged with infected nymphal *I. scapularis* ticks at ~ 4 weeks of age. Each pup was exposed to 5–7 infected nymphs, which were allowed to feed for 72 hours in a controlled containment chamber. Flat nymphal ticks were carefully transferred from storage vials using fine forceps and placed at the base of the neck and upper back of gently restrained pups. After placement, pups were monitored to ensure proper tick attachment before being returned to their cages. After 72 hours, all engorged ticks naturally detached and were collected from the bottom of the cages. Detached ticks were stored at −20°C until further analysis for *B. burgdorferi* colonization via dark-field microscopy.

The infected nymphs used in this study were generated by feeding larval *I. scapularis*

(Oklahoma Tick Laboratory) on a mouse infected with a multi-strain culture of *B. burgdorferi*

sensu stricto. This *B. burgdorferi* culture was recovered from heart and bladder tissues of

C3H/HeN mice infected with nymphal ticks collected in Massachusetts in 2020. To maintain

infection across generations, the multi-strain *B. burgdorferi* inoculum was first introduced into

C3H mice, after which successive mice were infected through tick-borne transmission. Larvae

were allowed to feed on infected mice, acquire *B. burgdorferi*, and molt into infected nymphs,

which were then used to challenge mice in a cycle that mimics natural enzootic transmission of

B. burgdorferi.

Detection of Anti-OspA and Anti-PepVF IgG by ELISA. Serum antibody responses to OspA

and to *B. burgdorferi* pepVF were assessed using an enzyme-linked immunosorbent assay

(ELISA). PepVF contains two peptides from *B. burgdorferi* VlsE and FlaB proteins in tandem,

separated by a 3aa glycine linker, and is used in our lab to determine *B. burgdorferi* infection

(24). Presence of antibody to PepVF indicates exposure to *B. burgdorferi*. Purified recombinant

OspA (1 μg/mL) or recombinant PepVF (1 μg/mL) was adsorbed onto Nunc MaxiSorp plates

(Thermo Fisher Scientific, Waltham, MA) by overnight incubation at 4°C. Following antigen

coating, the plates were blocked with 5% bovine serum albumin (BSA) in PBS-Tween (0.05%)

for 1 hour at room temperature to prevent nonspecific binding. Serum samples were diluted

1:100 in blocking buffer and incubated on the plates for 2 hours at 37°C. To detect antigen-

specific IgG, a goat anti-mouse IgG antibody conjugated to horseradish peroxidase (HRP)

(1:6000; Jackson ImmunoResearch, West Grove, PA) was added, followed by a 1-hour

incubation at 37°C. After washing, plates were developed using TMB substrate (Sigma-Aldrich)

for 15 minutes before stopping the reaction with 1N H_2SO_4. Optical density (OD) was measured

at 450 nm and blanked against the control using a SpectraMax Plus ELISA reader (Molecular

Devices, San Jose, CA). For Cutoff determination the positivity threshold was defined as mean +

3 SD of negative controls.

Molecular Detection of *B. burgdorferi* DNA by qPCR. Total DNA was extracted from heart

and bladder tissues and heart culture media using the DNeasy Blood & Tissue Kit (Qiagen,

Germantown, MD) according to the manufacturer's protocol. qPCR was performed to quantify

B. burgdorferi flaB gene copies per mg of tissue or per mL of culture media. Quantitative PCR

(qPCR) was performed to quantify *B. burgdorferi* flaB gene copies in DNA extracted from heart

and bladder tissues, as well as culture supernatants. Reactions were carried out in a 25 μL

volume using 2X TaqMan Master Mix (Thermo Fisher Scientific), with final primer and probe

concentrations of 200 nM. Each reaction contained 5 μL of DNA template, and all samples were

run in duplicate to ensure reproducibility. The primer and probe set used for TaqMan-based

detection of the *flaB* gene was as follows: Forward primer: 5'-AAGCAATCTAGGTCTCAAGC-

3'; Reverse primer: 5'-GCTTCAGCCTGGCCATAAATAG-3'; Probe: 5'-FAM-

AGATGTGGTAGACCCGAAGCCGAG-TAMRA-3'. Amplification was performed using a

QuantStudio 7 Real-Time PCR System (Applied Biosystems) under the following cycling

conditions: an initial denaturation at 95°C for 10 minutes, followed by 45 cycles of 95°C for 15

seconds and 60°C for 1 minute. Fluorescence was measured at each cycle, and *flaB* copy

numbers were determined by comparison to a standard curve generated from known

concentrations of *B. burgdorferi* genomic DNA

Statistical Analysis. Statistical analyses were conducted using GraphPad Prism 9.0 (GraphPad

Software, San Diego, CA). To assess the distribution of the data, the Shapiro-Wilk test was

applied to determine normality. For group comparisons, an unpaired t-test was used when data

followed a normal distribution, while the Mann-Whitney U test was applied to non-normally

distributed data. A significance threshold of $p < 0.05$ was used for all statistical tests.

Results

Transfer of Anti-OspA Antibodies from Dams to Pups. Pups were born to dams undergoing a

continuous vaccination schedule. To determine the extent of maternal antibody transfer, serum

anti-OspA IgG titers were measured in pups when they reached D24 (pre-challenge). Pups born

from dams vaccinated until breeding pairs were formed are grouped under 1st breeding. Pups

born from dams vaccinated until parturition are grouped under 2nd breeding. Pups born from

dams vaccinated until the pups were two weeks old are grouped under 3rd breeding. The results

are shown in **Fig 2**. Anti-OspA IgG was significantly increased in D24 pups born from

vaccinated dams during the 1st, the 2nd and the 3rd breeding, in contrast to pups born to control

unvaccinated dams. Nevertheless, we note that pups born from the 1st breeding of vaccinated

dams had an average anti-OspA IgG OD_{450}~ 0.271 (SD± 0.094); pups born from the 2nd

breeding of vaccinated dams had an average anti-OspA IgG OD_{450}~0.809 (SD± 0.073); and pups

born from the 3rd breeding of vaccinated dams had an average anti-OspA IgG OD_{450}~1.532

(SD± 0.591). The data shows that continued vaccination of the dams past parturition maintained

sufficient levels of protective antibody (OD450>0.8) in blood.

Seroconversion to *B. burgdorferi* in Tick Challenged Pups. To assess whether pups were

protected or infected with *B. burgdorferi* after tick challenge, anti-PepVF IgG was measured in

serum samples collected at euthanasia (~day 71), at least 3 weeks after the last day of challenge

(**Fig 3**). Pups (n=8) from the 1st Breeding of vaccinated dams had equivalent anti-PepVF IgG as

control pups (n=10), confirming infection with tick-transmitted *B. burgdorferi* in both groups (p

= 0.7642) (**Fig 3A**). Although 2/3 pups from the 2nd Breeding of vaccinated dams did not have

anti-PepVF antibodies (**Fig 3B**, circle), overall differences with the control group (n=5) were not

significant (p = 0.9432). Strikingly, pups (n=6) from the 3rd Breeding of vaccinated dams (**Fig

3C**) had anti-PepVF IgG levels below the cutoff in contrast to the control group (n=6), p =

0.0147, indicating absence of infection**.** These results suggest that maternal vaccination that

extended until and beyond parturition conferred anti-OspA passive immunity to pups.

Burden of *B. burgdorferi* in Tissues from Tick Challenged Pups. To provide additional

evidence of *B. burgdorferi* dissemination, we purified DNA from heart and bladder organs from

pups euthanized > 4 weeks after tick challenge (~ day 71) and performed qPCR targeting the *B.*

burgdorferi flaB gene (**Fig 4**). In pups from the 1st Breeding, there are no differences in *flaB*

loads in heart tissues from vaccinated and control groups; in pups from the 2nd Breeding, the

same 2 mice (2/3) in the vaccinated group that did not seroconvert to *B. burgdorferi*, also did not

have *flaB* in heart tissues, whereas 5/5 mice in the control group had *flaB* DNA in heart tissues.

Differences between the groups are not significant (p=0.3393). However, in pups from the 3rd

Breeding, no detectable *flaB* copies were found in bladder tissues from the vaccinated group

(0/6), in contrast to the pups from the control group that were all positive (6/6) for *B. burgdorferi*

flaB (p=0.0022). Absence of *B. burgdorferi flaB* DNA in pups from the 2^{nd} and 3^{rd} Breeding

correlate with absence of PepVF seroconversion, further confirming the protective effect of

maternal antibody transfer when dams were vaccinated until and beyond parturition.

Viability of *B. burgdorferi* in Tissues from Tick Challenged Pups. To further assess *B.*

burgdorferi persistence, heart tissues from pups born from 3^{rd} Breeding dams were cultured in

BSK-H medium at 34°C for 21 days, and live spirochetes were counted using a Petroff-Hausser

chamber under a dark field microscope (**Figure 5A**). Only pups (4/5) from unvaccinated control

dams had detectable motile *B. burgdorferi* in culture, confirming that live spirochetes had

successfully colonized cardiac tissue after tick challenge. The control group exhibited a range of

10^5-10^6 spirochetes per mL of culture, while pups from vaccinated dams had no visible live

bacteria, p=0.0152. qPCR analysis of heart culture supernatants (**Figure 5B**) confirmed that high

flaB copy numbers (mean 10^5) were detected in control heart cultures (5/5), whereas no *B.*

burgdorferi DNA was present in pups from vaccinated dams (0/5), p=0.0022. This indicates that

dams that received vaccine until 2 weeks post-parturition, produced pups that were protected

from cardiac colonization by *B. burgdorferi.*

Absence of OspA IgG in Serum from Pups Euthanized After Birth. To determine whether

maternal antibodies were transferred in utero, OspA ELISA was performed on serum from pups

euthanized immediately after they dropped at birth in comparison to the respective dam (**Fig 6**).

In contrast to the respective dam, no anti-OspA IgG was detected in newborns from the

vaccinated dam, confirming that placental transfer of antibodies was absent.

Anti-OspA IgG depletion from vaccinated dam serum after lactation. To investigate whether

maternal antibody depletion occurred due to antibody transfer to offspring, serum anti-OspA IgG

were measured in dams used for the 3^{rd} breeding at two time points (11Jun and 14Oct) and in the

respective litters, on the latter timepoint, before tick challenge (**Fig 7**). We found that the

reduction in OspA antibody measured between the two time points in the dams corresponds to

the increase in OspA antibody measured in the respective litters in the latter timepoint:

~0.7OD_{450} for Dam 1 (**Fig 7A**) and ~1.8OD_{450} for Dam 2 (**Fig 7B**). These findings suggest that

transfer of antibody from dam to pups may be dependent on the efficiency of the pups' suckling.

Discussion

Our study demonstrates that vaccination of *Peromyscus leucopus* dams against *B. burgdorferi*

lead to transfer of anti-OspA IgG to offspring and that, the extent of transfer and protective

efficacy depends on maintenance of maternal immunization. Pups born to dams vaccinated until

parturition and until two weeks post-parturition exhibited significantly higher anti-OspA IgG

levels than those from dams vaccinated until breeding pairs were created. After challenge with *B.*

burgdorferi infected *I. scapularis* ticks, pups born to vaccinated dams that maintained high

systemic anti-OspA IgG had no anti-*B. burgdorferi* antibody in serum, no *B. burgdorferi* DNA in

bladder or heart tissues and no motile *B. burgdorferi* in heart tissue. Furthermore, we detected no

serologic anti-OspA IgG in offspring from vaccinated dams, euthanized immediately after birth,

before suckling. The data indicates that anti-OspA antibody is transferred from *P. leucopus* dams

to pups via lactation and that it protects against tick-transmitted *B. burgdorferi*.

Vaccination of *P. leucopus* dams until the breeding pairs were created (1st Breeding) did not

protect *P. leucopus* pups from tick-mediated infection due to absence of sufficient levels of anti-

OspA antibodies (OD_{450}<0.8, **Fig 2**) in pups born ~ 10 weeks after the dams were vaccinated.

This was surprising given our previous studies in *P. leucopus* (25) and in C3H/HeN mice (22) in

which we vaccinated mice with the reservoir targeted vaccine (RTV). In the first study, we

vaccinated *P. leucopus* with 20-30 units of live RTV for 1-4 months and found that mice

maintained anti-OspA antibody for up to a year (25). In the second study, we found that

C3H/HeN dams vaccinated with 20 units of live RTV over a period of 8 weeks, produced pups

33-79 days after vaccination that maintained protective titers of OspA antibody from 2-9 weeks

after birth (22). The major difference between the studies is that in the current study we used

RTV inactivated with BPL, whereas in the former studies we used live RTV. This change was

intentional to evaluate a vaccine that mimics the product currently licensed by USDA, which is

inactivated with BPL (26). This data indicates that production of anti-OspA antibody by *P.*

leucopus requires persistent vaccination, if the vaccine formulation is inactivated likely because

it does not colonize the mouse gut.

Previous studies show that transfer of maternal IgG provides passive immunity but does not

necessarily confer sterilizing protection against vector-borne pathogens (20, 21). The degree of

IgG transfer and its impact on early-life immunity can vary depending on maternal antibody

titers, the mechanism of transfer (transplacental vs. lactogenic), the duration of exposure (27) and

neutralizing capability of the antibody (25). In our study, the neutralizing capability of anti-OspA

antibody transferred from *P. leucopus* dams to its pups is evidenced by the absence of *B.*

burgdorferi antibody in serum of pups (2/3 in Breeding 2 and 6/6 in Breeding 3, **Fig 3**), as well

as absence of DNA in heart and bladder tissues (the same mice 2/3 in Breeding 2, and 6/6 in

Breeding 3, **Fig 4**) and complete absence of motile spirochetes in cultures from heart tissue (6/6

in Breeding 3, **Fig 5**). All pups from breeding 1, born to vaccinated and control dams, had

antibodies to *B. burgdorferi* antigens after tick challenge (**Fig 3**) but we did not detect *B.*

burgdorferi DNA in heart tissue of these mice (**Fig 4**). We perform two independent tests

(antibody to *B. burgdorferi* in serum and *B. burgdorferi* DNA in tissues) and we accept one

positive result as evidence of *B. burgdorferi* dissemination (24). This method offsets risks

incurring from tracking all datapoints in experiments that can last up to 1 year. Thus, for

Breeding 1, we consider all mice in this experimental group to be infected and that this was

associated with insufficient production of anti-OspA antibody by the respective vaccinated dams.

Final efficacy of our intervention is determined by culture of *B. burgdorferi* from tissues of tick-

challenged mice. **Fig 5** shows conclusively that pups born to dams vaccinated until 2 weeks post-

parturition did not have viable *B. burgdorferi* in cultures from heart tissue. This shows that for

tick-transmitted *B. burgdorferi,* if *P. leucopus* is persistently vaccinated and produces sufficient

anti-OspA antibody to be transferred to pups, the offspring is protected to at least 4 weeks post

parturition. In addition, we show that immediately after birth, pups do not have anti-OspA

antibodies in serum (**Fig 6**). Thus, we infer that *P. leucopus* pups acquire OspA antibodies during

suckling. This observation aligns with our previous studies in C3H mice demonstrating that anti-

OspA IgG transfer via milk is a primary mechanism of passive immunity (22).

A potential question may arise from differences in **Fig 2** data showing significant increase of

anti-OspA IgG in pups euthanized on the day of birth from dams vaccinated until parturition in

Breeding 2 and data in **Fig 6** showing absence of anti-OspA IgG in pups euthanized immediately

after birth in Breeding 3. This is explained by the fact that pups from the 2nd breeding were born

overnight and were euthanized in the morning. Thus, for pups born from the 3rd breeding and to

confirm if antibody was transferred via lactation, as we showed previously for C3H mice (22),

we euthanized the pups immediately after they dropped to prevent any potential suckling time.

The decline in maternal anti-OspA IgG over time, particularly during lactation (**Fig 7**), suggests

a depletion of circulating IgG that could have impacted the duration of passive protection

provided to pups. This observation is consistent with findings in other models where maternal

antibody levels decrease postnatally as antibodies are transferred to offspring (20).

From a reservoir-targeted vaccine intervention perspective, our findings suggest that if maternal

anti-OspA antibodies reduce tick colonization efficiency or delay early-stage infection in *P.*

leucopus, this may have broader ecological implications by disrupting pathogen transmission

cycles. Further studies are needed to assess the effect of passively transferred anti-OspA

antibody in reduction of nymphal infection prevalence.

Acknowledgments

Research reported in this publication was supported by the National Institute of Allergy and

Infectious Diseases of the National Institutes of Health under Awards Number R01AI139267

(MGS), R43AI155211 (MGS) and R44AI167605 (MGS). The content is solely the responsibility

of the authors and does not necessarily represent the official views of the National Institutes of

Health.

352   **References**

353   1.   Marques AR. 2010. Lyme disease: a review. Curr Allergy Asthma Rep 10:13-20.

354   2.   Levine JF, Wilson ML, Spielman A. 1985. Mice as reservoirs of the Lyme disease

355   spirochete. Am J Trop Med Hyg 34:355-60.

356   3.   Donahue JG, Piesman J, Spielman A. 1987. Reservoir competence of white-footed mice

357   for Lyme disease spirochetes. Am J Trop Med Hyg 36:92-6.

358   4.   Mather TN, Mather ME. 1990. Intrinsic competence of three ixodid ticks (Acari) as

359   vectors of the Lyme disease spirochete. J Med Entomol 27:646-50.

360   5.   Tsao J, Barbour AG, Luke CJ, Fikrig E, Fish D. 2001. OspA immunization decreases

361   transmission of Borrelia burgdorferi spirochetes from infected Peromyscus leucopus mice

362   to larval Ixodes scapularis ticks. Vector Borne Zoonotic Dis 1:65-74.

363   6.   Richer LM, Brisson D, Melo R, Ostfeld RS, Zeidner N, Gomes-Solecki M. 2014.

364   Reservoir targeted vaccine against Borrelia burgdorferi: a new strategy to prevent Lyme

365   disease transmission. J Infect Dis 209:1972-80.

366   7.   Gomes-Solecki M. 2014. Blocking pathogen transmission at the source: reservoir

367   targeted OspA-based vaccines against Borrelia burgdorferi. Front Cell Infect Microbiol

368   4:136.

369   8.   Schaible UE, Kramer MD, Eichmann K, Modolell M, Museteanu C, Simon MM. 1990.

370   Monoclonal antibodies specific for the outer surface protein A (OspA) of Borrelia

371   burgdorferi prevent Lyme borreliosis in severe combined immunodeficiency (scid) mice.

372   Proc Natl Acad Sci U S A 87:3768-72.

9.   Schwan TG, Piesman J, Golde WT, Dolan MC, Rosa PA. 1995. Induction of an outer

surface protein on Borrelia burgdorferi during tick feeding. Proc Natl Acad Sci U S A

92:2909-13.

10.   de Silva AM, Telford SR, 3rd, Brunet LR, Barthold SW, Fikrig E. 1996. Borrelia

burgdorferi OspA is an arthropod-specific transmission-blocking Lyme disease vaccine. J

Exp Med 183:271-5.

11.   Stevenson B, Schwan TG, Rosa PA. 1995. Temperature-related differential expression of

antigens in the Lyme disease spirochete, Borrelia burgdorferi. Infect Immun 63:4535-9.

12.   Schwan TG, Piesman J. 2000. Temporal changes in outer surface proteins A and C of the

lyme disease-associated spirochete, Borrelia burgdorferi, during the chain of infection in

ticks and mice. J Clin Microbiol 38:382-8.

13.   Pal U, Yang X, Chen M, Bockenstedt LK, Anderson JF, Flavell RA, Norgard MV, Fikrig

E. 2004. OspC facilitates Borrelia burgdorferi invasion of Ixodes scapularis salivary

glands. J Clin Invest 113:220-30.

14.   Grimm D, Tilly K, Byram R, Stewart PE, Krum JG, Bueschel DM, Schwan TG,

Policastro PF, Elias AF, Rosa PA. 2004. Outer-surface protein C of the Lyme disease

spirochete: a protein induced in ticks for infection of mammals. Proc Natl Acad Sci U S

A 101:3142-7.

15.   Fikrig E, Barthold SW, Kantor FS, Flavell RA. 1990. Protection of mice against the

Lyme disease agent by immunizing with recombinant OspA. Science 250:553-6.

16.   Fikrig E, Barthold SW, Kantor FS, Flavell RA. 1992. Long-term protection of mice from

Lyme disease by vaccination with OspA. Infect Immun 60:773-7.

17.   Sigal LH, Zahradnik JM, Lavin P, Patella SJ, Bryant G, Haselby R, Hilton E, Kunkel M,

Adler-Klein D, Doherty T, Evans J, Molloy PJ, Seidner AL, Sabetta JR, Simon HJ,

Klempner MS, Mays J, Marks D, Malawista SE. 1998. A vaccine consisting of

recombinant Borrelia burgdorferi outer-surface protein A to prevent Lyme disease.

Recombinant Outer-Surface Protein A Lyme Disease Vaccine Study Consortium. N Engl

J Med 339:216-22.

18.   Steere AC, Sikand VK, Meurice F, Parenti DL, Fikrig E, Schoen RT, Nowakowski J,

Schmid CH, Laukamp S, Buscarino C, Krause DS. 1998. Vaccination against Lyme

disease with recombinant Borrelia burgdorferi outer-surface lipoprotein A with adjuvant.

Lyme Disease Vaccine Study Group. N Engl J Med 339:209-15.

19.   Tsao JI, Wootton JT, Bunikis J, Luna MG, Fish D, Barbour AG. 2004. An ecological

approach to preventing human infection: vaccinating wild mouse reservoirs intervenes in

the Lyme disease cycle. Proc Natl Acad Sci U S A 101:18159-64.

20.   Hasselquist D, Nilsson JA. 2009. Maternal transfer of antibodies in vertebrates: trans-

generational effects on offspring immunity. Philos Trans R Soc Lond B Biol Sci 364:51-

60.

21.   Boulinier T, Staszewski V. 2008. Maternal transfer of antibodies: raising immuno-

ecology issues. Trends Ecol Evol 23:282-8.

22.   Phillip K, Nair N, Samanta K, Azevedo JF, Brown GD, Petersen CA, Gomes-Solecki M.

2021. Maternal transfer of neutralizing antibodies to B. burgdorferi OspA after oral

vaccination of the rodent reservoir. Vaccine 39:4320-4327.

23.   Gomes-Solecki M, Richer L. 2018. Recombinant E. coli Dualistic Role as an Antigen-

adjuvant Delivery Vehicle for Oral Immunization. Methods Mol Biol 1690:347-357.

24.   Gingerich MC, Nair N, Azevedo JF, Samanta K, Kundu S, He B, Gomes-Solecki M.

419   2024. Intranasal vaccine for Lyme disease provides protection against tick transmitted

420   Borrelia burgdorferi beyond one year. NPJ Vaccines 9:33.

421   25.   Meirelles Richer L, Aroso M, Contente-Cuomo T, Ivanova L, Gomes-Solecki M. 2011.

422   Reservoir targeted vaccine for lyme borreliosis induces a yearlong, neutralizing antibody

423   response to OspA in white-footed mice. Clin Vaccine Immunol 18:1809-16.

424   26.   Stafford KC, 3rd, Williams SC, van Oosterwijk JG, Linske MA, Zatechka S, Richer LM,

425   Molaei G, Przybyszewski C, Wikel SK. 2020. Field evaluation of a novel oral reservoir-

426   targeted vaccine against Borrelia burgdorferi utilizing an inactivated whole-cell bacterial

427   antigen expression vehicle. Exp Appl Acarol 80:257-268.

428   27.   Fouda GG, Martinez DR, Swamy GK, Permar SR. 2018. The Impact of IgG

429   transplacental transfer on early life immunity. Immunohorizons 2:14-25.

Figure Legends

Figure 1. Experimental design. A) Oral vaccination schedule of adult female *Peromyscus leucopus*: mice were allowed to eat RTV pellets on vaccination weeks; pellets were exchanged for regular feed during rest weeks; control mice received uncoated pellets; B) Timeline for breeding of dams: dams were bled to evaluate serologic antibody to OspA and three experimental groups were set as 1st, 2nd and 3rd Breeding based on duration of vaccination and breeding milestones; C) Birth of pups, challenge with *B. burgdorferi* infected *I. scapularis* ticks and collection of samples.

Figure 2. Transfer of OspA antibodies from dams to pups. ELISA quantification of pre-challenge anti-OspA IgG in sera from D24 pups born to vaccinated dams. Optical density (OD) at 450nm (blanked) is shown for each group. Each point represents an individual pup, with lines indicating group means ± SD. Statistical analysis by Mann-Whitney test *p<0.05, ** p<0.005, *** p<0.0005.

Figure 3. Seroconversion to *B. burgdorferi* in tick challenged pups. (A–C) ELISA quantification of anti-PepVF IgG in terminal sera from pups born to vaccinated dams. Pups were bled ~ 4 weeks post tick challenge. Optical density (OD) at 450 nm (blanked) is shown for each group. Each data point represents an individual pup, with lines indicating group means ± SD. Statistics by Unpaired t test: ns, not significant, * p<0.05.

Figure 4. Burden of *B. burgdorferi* in tissues from tick challenged pups. (A–C) qPCR quantification of the *flaB* gene in heart and bladder tissues collected from pups ~ 4 weeks post tick

challenge. Each data point represents an individual pup, with lines indicating group means. Statistics

by Mann Whitney test: ns, not significant, **p<0.005.

Figure 5. Viability of *B. burgdorferi* in tissues from tick challenged pups. Quantification of *B.*

burgdorferi in cultures of heart tissue from pups at ~ 4 weeks post-challenge: (A) Enumeration of

motile *B. burgdorferi* using a Petroff-Hausser chamber under a dark field microscope; (B) qPCR

quantification of *B. burgdorferi flaB* in heart culture media. Each data point represents an individual

pup, with lines indicating group means ± SD. Statistics by Mann Whitney test, *p<0.05 and **

p<0.005.

Figure 6. Absence of OspA IgG in serum from pups euthanized after birth. ELISA

quantification of anti-OspA IgG in serum from pups euthanized immediately after birth from dams

vaccinated until two weeks parturition (3rd Breeding). Optical density (OD) at 450 nm (blanked) is

shown. Each data point represents an individual pup, with lines indicating group means. The cutoff

value (dashed line) represents the mean + 3 SD of negative controls.

Figure 7. Anti-OspA IgG depletion from vaccinated dam serum after pup suckling. ELISA

quantification of anti-OspA IgG in serum from two dams used in the 3rd breeding and the respective

litters. Sera from dams were collected at two time-points ~ 4 months apart. Sera from the respective

litters were collected on the latter time point, before tick challenge. Each data point represents an

individual pup, with lines indicating group means ± SD.

A. Dams Vaccination Schedule

Vaccination Rest

Weeks 0 1 2 3 4 5 6 7 8 9 10 11 12 ... etc.

Bleed Bleed Bleed

B. Dams Breeding

9 Dams

1st Breeding
(RTV until pairs were created)

2nd Breeding
(RTV until parturition)

3rd Breeding
(RTV until 2wk breastfeeding)

1st Breeding (RTV until pairs were created): Dams were vaccinated until breeding pairs were created (12 weeks)

2nd Breeding (RTV until parturition): Dams were vaccinated until birth of first pups (24 weeks)

3rd Breeding (RTV until 2wk breastfeeding): Dams were vaccinated until pups reached 2 weeks of age (36 weeks)

C. Pups Birth, Tick Challenge and Euthanasia

Breeding
pairs
created

3-4wk

~ 6 weeks

Pup age (weeks) 0 1 2 3 4 5 6 7 8

Birth

Group euthanized just
after dropping at birth

Pre-challenge
D24 pup bleed

Tick
Challenge

Euthanasia
of pups

Collection of blood,
heart and bladder