Maternal Transfer of Vaccine-Induced Anti-OspA Antibodies in Peromyscus spp Protects

Pups from Tick-Transmitted *Borrelia burgdorferi*

Jose F. Azevedo[1]*, Greg Joyner[1]*, Suman Kundu[1], Kamalika Samanta[1], and Maria Gomes-

Solecki[1]

[1]University of Tennessee Health Science Center, Department of Microbiology, Immunology, and

Biochemistry

*Equal contribution

Correspondence to: Maria Gomes-Solecki, mgomesso@uthsc.edu

Abstract

Maternal antibody transfer provides passive immunity to offspring; however, the efficacy and

duration of this protection depends on maternal antibody levels and transfer efficiency. We

investigated whether oral vaccination of *Peromyscus leucopus* dams with recombinant OspA-

expressing *E. coli* could induce maternal transfer of anti-OspA antibodies and protect pups from

Borrelia burgdorferi challenge. Dams were vaccinated for different durations: (i) until breeding,

(ii) until birth of pups, or (iii) until pups were 2 weeks old. Pups were challenged with *Ixodes*

scapularis nymph-transmitted *B. burgdorferi* at ~ 4 weeks of age. Anti-OspA IgG were

quantified in dams and pups, and *B. burgdorferi* anti-PepVF IgG were quantified in pups.

Furthermore, *B. burgdorferi* burden was assessed by qPCR targeting the *flaB* gene in pups' heart

and bladder tissues > 4 weeks after tick challenge. *P. leucopus* pups born to dams vaccinated

until breeding, exhibited low anti-OspA antibody and were not protected from tick transmitted *B.*

burgdorferi infection. However, when maternal vaccination extended until pups were born and

until pups were two weeks old, significant anti-OspA antibody transfer and protection occurred.

This was evidenced by absence of antibody to *B. burgdorferi* PepVF, absence of *B. burgdorferi*

flaB DNA in heart and bladder tissues, and absence of *flaB* in culture from heart tissues from

pups euthanized >9 weeks after birth. We show that transfer of anti-OspA antibodies from *P.*

leucopus dams to offspring prevents tick transmission of *B. burgdorferi* in the major reservoir

host of this spirochete in the United States.

Importance: This study contributes to our understanding of how reservoir targeted vaccines

designed to block transmission of *B. burgdorferi* from infected *Ixodes scapularis* ticks may

disrupt the enzootic cycle of this spirochete and reduce incidence of Lyme disease.

Key words: *Peromyscus***,** reservoir targeted oral vaccine, block transmission, *Ixodes scapularis*,

enzootic cycle, OspA, mouse, *Borrelia burgdorferi*, Lyme disease

Introduction

Lyme disease, caused by *Borrelia burgdorferi* and transmitted by *Ixodes* ticks, is the most prevalent vector-borne disease in North America (1). The white-footed mouse (*Peromyscus leucopus*) serves as the primary reservoir host, maintaining *B. burgdorferi* in enzootic cycles and facilitating transmission to ticks that subsequently infect humans and other mammals (2). Unlike humans, these reservoir hosts do not develop significant disease following *B. burgdorferi* infection, yet they serve as a crucial source of spirochete acquisition for larval ticks (3). As a result, interrupting this transmission within the reservoir host has been proposed as a strategy for controlling Lyme disease (4).

Outer surface protein A (OspA) is a major target for Lyme disease vaccine development due to its essential role in *B. burgdorferi* persistence within the tick vector (5). OspA facilitates spirochete adherence to the tick midgut, allowing the bacterium to survive between blood meals (6). During tick feeding, *B. burgdorferi* downregulates OspA while upregulating OspC, facilitating migration to the salivary glands and subsequent transmission to the host (7). Vaccines targeting OspA aim to elicit antibodies that neutralize spirochetes within the tick gut, preventing their transmission before they reach the mammalian host (8). OspA-based vaccines have been explored in multiple contexts, including human vaccination (1), direct vaccination of *P. leucopus* (9), and bait-delivered reservoir-targeted vaccines that reduce pathogen burden in wild mouse populations (10).

Beyond direct vaccination, maternal transfer of immunity provides an alternative strategy to protect neonates during the critical early stages of life. (11). In the context of vector-borne pathogens, this temporary immunity may reduce early-life susceptibility to infection (12).

Previous studies have demonstrated that maternal vaccination with OspA results in detectable

anti-OspA IgG in offspring (13), though the degree to which this protects against *B. burgdorferi*

transmission remains unknown.

In this study, we investigate the extent of maternal antibody transfer in *P. leucopus* and assess

whether vaccination of dams with recombinant OspA-expressing *E. coli* can confer protection to

neonates. We test whether different durations of maternal vaccination influence antibody transfer

and protection against tick-mediated *B. burgdorferi* challenge. Our findings have implications

for reservoir-targeted vaccination strategies aimed at reducing Lyme disease risk by breaking the

transmission cycle of *B. burgdorferi*.

Materials and Methods

Ethics Statement. This study was conducted in accordance with the Guide for the Care and Use

of Laboratory Animals of the US National Institutes of Health. The protocol was approved by the

University of Tennessee Health Science Center Institutional Animal Care and Use Committee

(IACUC), protocol #22-0400. *Peromyscus leucopus* were sourced from the Peromyscus Genetic

Stock Center (PGSC) at the University of South Carolina.

Vaccination Strategy and Experimental Design. Dams (n=10) were assigned to three

experimental groups based on the duration of maternal vaccination. In the first group, referred to

as Breeding 1, dams received oral vaccination for 12 weeks prior to mating. The second group,

Breeding 2, was vaccinated for an extended period of 24 weeks, continuing until parturition. The

final group, Breeding 3, remained on the vaccination regimen for 36 weeks, ensuring that

immunization continued until their pups reached two weeks of age.

Vaccine Preparation and Administration. Dams were orally vaccinated with recombinant

OspA-expressing Escherichia coli (RTV pellets) produced as previously described (14),

inactivated by β-propiolactone. Vaccine pellets contained OspA-expressing *E. coli*

BL21(DE3)pLysS grown in Terrific Broth (TBY) at 37°C until OD600 ~0.8, induced with 1 mM

IPTG for 3 h, and harvested by centrifugation at 2000×g at 4°C. The bacterial biomass was

resuspended in phosphate-buffered saline (PBS) and sprayed onto feed pellets. Pellets were

allowed to dry between application layers.

Dams received RTV pellets daily for 5 consecutive days per week, with a 2-day rest period in

which they received regular mouse chow. Control dams received uncoated pellets. The

vaccination schedule was maintained based on experimental grouping (**Figure 1**).

Breeding and Litter Management. Breeding pairs were established based on high anti-OspA

titers [OD450>0.8) assessed by ELISA. Each vaccinated female was cohoused with an

unvaccinated male for 3–4 weeks, after which females were single-housed until parturition. Pups

remained with their dams until 24–25 days of age, at which point they underwent nymphal tick

challenge.

Blood Collection. Blood samples were collected via submandibular vein bleed at day 24 (pre-

challenge) and day 71 (terminal bleed) for pups. Samples were allowed to clot for 30 min at

room temperature before centrifugation at 2000×g for 10 min at 4°C. Serum was aliquoted and

stored at −80°C until analysis.

Tissue Collection and Processing. At necropsy on d71 (~9-10 week old pups), heart and

bladder tissues were harvested, transported on dry ice, and stored at −80°C. For organ culture

experiments, freshly excised heart tissue was immediately placed into Barbour-Stoenner-Kelly H

(BSK-H) medium supplemented with 6% rabbit serum (Sigma-Aldrich, St. Louis, MO) and

incubated at 34°C for 21 days. Culture supernatants were analyzed for: 1. Motile Borrelia

detection using Petroff-Hausser chamber (dark-field microscopy, 200× magnification) and 2.

DNA quantification via *flaB* qPCR.

Tick Challenge and Borrelia Infection Monitoring. To assess susceptibility to *Borrelia*

burgdorferi infection, pups were challenged with infected nymphal *Ixodes scapularis* ticks at ~ 4

weeks of age. Each pup was exposed to 5–7 infected nymphs, which were allowed to feed for 72

hours in a controlled containment chamber. Flat nymphal ticks were carefully transferred from

storage vials using fine forceps and placed at the base of the neck and upper back of gently

restrained pups. After placement, pups were monitored to ensure proper tick attachment before

being returned to their cages. After 72 hours, all engorged ticks naturally detached and were

collected from the bottom of the cages. Detached ticks were stored at −20°C until further

analysis for *B. burgdorferi* colonization via dark-field microscopy.

The infected nymphs used in this study were generated by feeding larval *Ixodes scapularis*

(Oklahoma Tick Laboratory) on a mouse infected with a multi-strain culture of *Borrelia*

burgdorferi sensu stricto. This *B. burgdorferi* culture was recovered from heart and bladder

tissues of Ch3-HeN mice infected with nymphal ticks collected in Massachusetts in 2020. To

maintain infection across generations, the multi-strain *B. burgdorferi* inoculum was first

introduced into C3H mice, after which successive mice were infected through tick-borne

transmission. Larvae were allowed to feed on infected mice, acquire *B. burgdorferi*, and molt

into infected nymphs, which were then used to challenge mice in a cycle that mimics natural

enzootic transmission of *B. burgdorferi*.

ELISA for Anti-OspA and Anti-PepVF IgG. Serum antibody responses to OspA and to *B.*

burgdorferi pepVF were assessed using an enzyme-linked immunosorbent assay (ELISA).

PepVF contains two peptides from *Borrelia burgdorferi* VlsE and flaB proteins in tandem,

separated by a 3aa glycine linker, and is used in our lab to determine *B. burgdorferi* infection

(15). Presence of antibody to PepVF indicates exposure to *B. burgdorferi*. Purified recombinant

OspA (1 μg/mL) or recombinant PepVF (1 μg/mL) was adsorbed onto Nunc MaxiSorp plates

(Thermo Fisher Scientific, Waltham, MA) by overnight incubation at 4°C. Following antigen

coating, the plates were blocked with 5% bovine serum albumin (BSA) in PBS-Tween (0.05%)

for 1 hour at room temperature to prevent nonspecific binding. Serum samples were diluted

1:100 in blocking buffer and incubated on the plates for 2 hours at 37°C. To detect antigen-

specific IgG, a goat anti-mouse IgG antibody conjugated to horseradish peroxidase (HRP)

(1:6000; Jackson ImmunoResearch, West Grove, PA) was added, followed by a 1-hour

incubation at 37°C. After washing, plates were developed using TMB substrate (Sigma-Aldrich)

for 15 minutes before stopping the reaction with 1N H_2SO_4. Optical density (OD) was measured

at 450 nm and blanked against the control using a SpectraMax Plus ELISA reader (Molecular

Devices, San Jose, CA). For Cutoff determination the positivity threshold was defined as mean +

3 SD of negative controls.

Molecular Detection of Borrelia burgdorferi DNA by qPCR. Total DNA was extracted from

heart and bladder tissues and heart culture media using the DNeasy Blood & Tissue Kit (Qiagen,

Germantown, MD) according to the manufacturer's protocol. qPCR was performed to quantify

B. burgdorferi flaB gene copies per mg of tissue or per mL of culture media. Quantitative PCR

(qPCR) was performed to quantify *Borrelia burgdorferi* flaB gene copies in DNA extracted from

heart and bladder tissues, as well as culture supernatants. Reactions were carried out in a 25 μL

volume using 2X TaqMan Master Mix (Thermo Fisher Scientific), with final primer and probe

concentrations of 200 nM. Each reaction contained 5 μL of DNA template, and all samples were

run in duplicate to ensure reproducibility. The primer and probe set used for TaqMan-based

detection of the *flaB* gene was as follows: Forward primer: 5'-AAGCAATCTAGGTCTCAAGC-

3'; Reverse primer: 5'-GCTTCAGCCTGGCCATAAATAG-3'; Probe: 5'-FAM-

AGATGTGGTAGACCCGAAGCCGAG-TAMRA-3'. Amplification was performed using a

QuantStudio 7 Real-Time PCR System (Applied Biosystems) under the following cycling

conditions: an initial denaturation at 95°C for 10 minutes, followed by 45 cycles of 95°C for 15

seconds and 60°C for 1 minute. Fluorescence was measured at each cycle, and *flaB* copy

numbers were determined by comparison to a standard curve generated from known

concentrations of *B. burgdorferi* genomic DNA

Statistical Analysis. Statistical analyses were conducted using GraphPad Prism 9.0 (GraphPad

Software, San Diego, CA). To assess the distribution of the data, the Shapiro-Wilk test was

applied to determine normality. For group comparisons, an unpaired t-test was used when data

followed a normal distribution, while the Mann-Whitney U test was applied to non-normally

distributed data. A significance threshold of $p < 0.05$ was used for all statistical tests.

Results

Maternal Transfer of Anti-OspA Antibodies in Pups. To determine the extent of maternal antibody transfer, serum anti-OspA IgG titers were measured in pups when they reached D24 (pre-challenge). Pups were born to dams undergoing a continuous vaccination schedule. Pups born from dams vaccinated until breeding pairs were formed are grouped under 1st breeding. Pups born from dams vaccinated until giving birth are grouped under 2nd breeding. Pups born from dams vaccinated until the pups were two weeks old are grouped under 3rd breeding. The results are shown in Figure 2. Anti-OspA IgG was significantly increased in D24 pups born from vaccinated dams during the 1st, the 2nd and the 3rd breeding, in contrast to pups born to control unvaccinated dams. Furthermore, levels of anti-OspA IgG were significantly increased in pups from the 2nd and 3rd breeding in comparison to the 1st breeding. The data suggests that longevity of vaccination of the dams past birth of the pups affects the overall level of anti-OspA IgG transferred from dams to pups.

Seroconversion to *B. burgdorferi* Following Tick Challenge. To assess whether pups were protected from tick-mediated *B. burgdorferi* infection, anti-PepVF IgG were measured in serum samples collected at euthanasia (~day 70), at least 4 weeks after the last day of challenge (**Figure 3.** Pups (n=8) from the 1st Breeding of vaccinated dams had equivalent anti-PepVF IgG titers as control pups (n=10), confirming infection with tick-transmitted *B. burgdorferi* in both groups (p = 0.7642) (Fig 3A). Although 2/3 pups from the 2nd Breeding of vaccinated dams did not have anti-PepVF antibodies (Fig 3B, circle), overall differences with the control group (n=5) were not significant due to the low numbers of pups in this experiment (p = 0.9432). In contrast to the

control group, pups from the 3rd Breeding of vaccinated dams (Fig 3C) had anti-PepVF IgG

levels below the cutoff (p = 0.0147), indicating absence of infection. These results suggest that

maternal vaccination that extended to- and post-parturition conferred anti-OspA passive

immunity to pups.

Pathogen Burden in Tissues (qPCR Analysis). To directly assess pathogen dissemination to

tissues, we purified DNA from heart and bladder from pups euthanized > 4 weeks after tick

challenge (~ day 71) and performed qPCR targeting the *B. burgdorferi flaB* gene (**Figure 4**). In

pups from the 1st Breeding, there are no differences in *flaB* loads in heart tissues from vaccinated

and control groups; in pups from the 2nd Breeding, 2/3 mice in the vaccinated group did not have

flaB in heart tissues, whereas 5/5 mice in the control group had *flaB* DNA in heart tissues.

However, differences between the groups are not significant (p=0.3393). However, in pups from

the 3rd Breeding, no detectable *flaB* copies were found in bladder tissues from the vaccinated

group (0/6), in contrast to the pups from the control group that were all positive (6/6) for *B.*

burgdorferi flaB (p=0.0022). Absence of *B. burgdorferi flaB* DNA in pups from the 2nd and 3rd

Breeding of vaccinated dams correlates with absence of PepVF seroconversion, further

confirming the protective effect of maternal antibody transfer when dams were vaccinated until

parturition.

Heart Tissue Culture of *B. burgdorferi* From Pups From the 3rd Breeding. To further assess

B. burgdorferi persistence, heart tissues from pups born from 3rd Breeding dams were cultured in

BSK-H medium at 34°C for 21 days, and live spirochetes were counted using a Petroff-Hausser

chamber under a dark field microscope (**Figure 5A**). Only pups (4/5) from unvaccinated control

dams had detectable motile *B. burgdorferi* in culture, confirming that spirochetes had

successfully colonized cardiac tissue. The control group exhibited up to 4 million spirochetes per

mL, while pups from vaccinated dams had no detectable live bacteria, p=0.0152. qPCR analysis

of culture supernatants (**Figure 5B**) revealed that high *flaB* copy numbers were detected in

control heart cultures (5/5), whereas no *B. burgdorferi* DNA was present in pups from vaccinated

dams (0/5), p=0.0022. This indicates that dams that received vaccine until 2 weeks post-

parturition, produced pups that were protected from cardiac colonization by *B. burgdorferi.*

Absence of OspA IgG in Pups Euthanized at Birth. To determine whether maternal antibodies

were transferred in utero, OspA ELISA was performed on serum from pups euthanized at birth in

comparison to the respective dam (**Figure 6**). In contrast to the respective dam, no anti-OspA

IgG was detected in newborns from vaccinated dams, confirming that placental transfer of

antibodies was absent, and that passive immunity was most likely mediated via breastfeeding.

This observation aligns with previous studies demonstrating that anti-OspA IgG transfer via milk

is a primary mechanism of passive immunity (13).

Anti-OspA IgG transfer from dams to the respective pup litters. To investigate whether

maternal antibody depletion occurred due to antibody transfer to offspring, serum anti-OspA IgG

were measured in dams used for the 3rd breeding at two time points (11Jun and 14Oct) and in the

respective litters, on the latter timepoint, before tick challenge (**Figure 7**). We found that the

reduction in OspA antibody measured between the two time points in the dams corresponds to

the increase in OspA antibody measured in the respective litters in the latter timepoint:

~0.7OD$_{450}$ for Dam 1 (Fig 7A) and ~1.8OD$_{450}$ for Dam 2 (Fig 7B). These findings suggest that

transfer of antibody from dam to pups may be dependent on the efficiency of the pups'

breastfeeding.

Discussion

Our study demonstrates that maternal vaccination against *B. burgdorferi* can lead to the transfer

of anti-OspA IgG to offspring in *Peromyscus leucopus*, though the extent of transfer and

protective efficacy depends on the duration of maternal immunization. Pups born to dams

vaccinated until birth of the pups and until two weeks postpartum exhibited significantly higher

anti-OspA IgG titers than those from dams vaccinated until breeding of the dams. After

challenge with *B. burgdorferi* infected *Ixodes scapularis* ticks, 2/3 pups from dams vaccinated

until birth (2nd Breeding) and all pups (6/6) from dams vaccinated until two weeks postpartum

(3rd Breeding), had no anti-PepVF antibody in serum, no *B. burgdorferi* DNA in bladder or heart

tissues, and in the 3rd Breeding we found no motile *B. burgdorferi* in heart tissue, which was

confirmed by qPCR. The data shows that anti-OspA antibody transferred from *P. leucopus* dams

to pups challenged ~ 4 weeks after birth.

Vaccination of *P. leucopus* dams until the breeding pairs were created (1st Breeding) did not

protect *P. leucopus* pups from tick-mediated infection due to absence of anti-OspA antibodies in

pups born more than 4 weeks after the dams were vaccinated. This was surprising given our

previous studies in *Peromyscus leucopus* (Meirelles Richer, 2011) and C3H-HeN mice (Phillip,

2021) orally vaccinated with our reservoir targeted vaccine (RTV). In the first study, we

vaccinated *P. leucopus* with 20-30 units of live RTV for 1-4 months and found that mice

maintained anti-OspA antibody for up to a year (Meirelles Richer, 2011). In the second study, we

found that C3H-HeN dams vaccinated with 20 units of live RTV over a period of 8 weeks,

produced pups 33-79 days after vaccination that maintained protective titers of OspA antibody

from 2-9 weeks after birth. The major difference between the former studies and the current

study is that in the current study we used RTV inactivated with BPL, whereas in the former

studies we used live RTV. This data indicates that production of anti-OspA antibody by *P.*

leucopus requires persistent vaccination, if the vaccine formulation is inactivated.

Previous studies show that maternal IgG transfer provides passive immunity but does not

necessarily confer sterilizing protection against vector-borne pathogens (11, 12). The degree of

IgG transfer and its impact on early-life immunity can vary depending on maternal antibody

titers, the mechanism of transfer (transplacental vs. lactogenic), and the duration of exposure (16)

and neutralizing capability of the antibody (17-19). In our study, the neutralizing capability of

anti-OspA antibody transferred from *P. leucopus* dams to its pups is evidenced by the complete

absence of *B. burgdorferi* antibody in serum of pups (no exposure), as well as absence of DNA

or motile spirochetes in the tissues (no dissemination). This shows that for tick-transmitted *B.*

burgdorferi if *P. leucopus* is persistently vaccinated and produces sufficient anti-OspA antibody

to be transferred to pups, the offspring is protected to at least 4 weeks post birth. The decline in

maternal anti-OspA titers over time, particularly during lactation, suggests a depletion of

circulating IgG that could have impacted the duration of passive protection provided to pups.

This observation is consistent with findings in other models where maternal antibody levels

decrease postnatally as antibodies are transferred to offspring (11).

From a reservoir-targeted vaccine control perspective, our findings suggest that if maternal anti-

OspA antibodies reduce tick colonization efficiency or delay early-stage infection in *P. leucopus*,

this could have broader ecological implications by disrupting pathogen transmission cycles.

Further studies are needed to assess the effect of passively transferred anti-OspA antibody in

reduction of nymphal infection prevalence.

Acknowledgments

Research reported in this publication was supported by the National Institute of Allergy and

Infectious Diseases of the National Institutes of Health under Awards Number R01AI139267

(MGS), R43AI155211 (MGS) and R44AI167605 (MGS). The content is solely the responsibility

of the authors and does not necessarily represent the official views of the National Institutes of

Health.

References

1.   Steere AC, Sikand VK, Meurice F, Parenti DL, Fikrig E, Schoen RT, et al. Vaccination
against Lyme disease with recombinant Borrelia burgdorferi outer-surface lipoprotein A
with adjuvant. Lyme Disease Vaccine Study Group. N Engl J Med. 1998;339(4):209-15.
2.   Bhattacharya D, Bensaci M, Luker KE, Luker G, Wisdom S, Telford SR, et al.
Development of a baited oral vaccine for use in reservoir-targeted strategies against Lyme
disease. Vaccine. 2011;29(44):7818-25.
3.   Telford SR, 3rd, Cunningham JA, Waltari E, Hu L. Nest box-deployed bait for
delivering oral vaccines to white-footed mice. Ticks Tick Borne Dis. 2011;2(3):151-5.
4.   Tsao J, Barbour AG, Luke CJ, Fikrig E, Fish D. OspA immunization decreases
transmission of Borrelia burgdorferi spirochetes from infected Peromyscus leucopus mice
to larval Ixodes scapularis ticks. Vector Borne Zoonotic Dis. 2001;1(1):65-74.
5.   Tilly K, Bestor A, Rosa PA. Functional Equivalence of OspA and OspB, but Not OspC,
in Tick Colonization by Borrelia burgdorferi. Infect Immun. 2016;84(5):1565-73.
6.   de Silva AM, Telford SR, 3rd, Brunet LR, Barthold SW, Fikrig E. Borrelia burgdorferi
OspA is an arthropod-specific transmission-blocking Lyme disease vaccine. J Exp Med.
1996;183(1):271-5.
7.   Schwan TG, Piesman J. Temporal changes in outer surface proteins A and C of the
lyme disease-associated spirochete, Borrelia burgdorferi, during the chain of infection in
ticks and mice. J Clin Microbiol. 2000;38(1):382-8.
8.   Fikrig E, Barthold SW, Kantor FS, Flavell RA. Protection of mice against the Lyme
disease agent by immunizing with recombinant OspA. Science. 1990;250(4980):553-6.
9.   Gomes-Solecki MJ, Brisson DR, Dattwyler RJ. Oral vaccine that breaks the
transmission cycle of the Lyme disease spirochete can be delivered via bait. Vaccine.
2006;24(20):4440-9.
10.   Richer LM, Brisson D, Melo R, Ostfeld RS, Zeidner N, Gomes-Solecki M. Reservoir
targeted vaccine against Borrelia burgdorferi: a new strategy to prevent Lyme disease
transmission. J Infect Dis. 2014;209(12):1972-80.
11.   Hasselquist D, Nilsson JA. Maternal transfer of antibodies in vertebrates: trans-
generational effects on offspring immunity. Philos Trans R Soc Lond B Biol Sci.
2009;364(1513):51-60.
12.   Boulinier T, Staszewski V. Maternal transfer of antibodies: raising immuno-ecology
issues. Trends Ecol Evol. 2008;23(5):282-8.
13.   Phillip K, Nair N, Samanta K, Azevedo JF, Brown GD, Petersen CA, et al. Maternal
transfer of neutralizing antibodies to B. burgdorferi OspA after oral vaccination of the
rodent reservoir. Vaccine. 2021;39(31):4320-7.
14.   Gomes-Solecki M, Richer L. Recombinant E. coli Dualistic Role as an Antigen-
adjuvant Delivery Vehicle for Oral Immunization. Methods Mol Biol. 2018;1690:347-57.
15.   Gingerich MC, Nair N, Azevedo JF, Samanta K, Kundu S, He B, et al. A parainfluenza
virus 5 (PIV5)-vectored intranasal vaccine for Lyme disease provides long-lasting
protection against tick transmitted Borrelia burgdorferi in mice. Res Sq. 2023.

16.   Fouda GG, Martinez DR, Swamy GK, Permar SR. The Impact of IgG transplacental
transfer on early life immunity. Immunohorizons. 2018;2(1):14-25.
17.   Meirelles Richer L, Aroso M, Contente-Cuomo T, Ivanova L, Gomes-Solecki M.
Reservoir targeted vaccine for lyme borreliosis induces a yearlong, neutralizing antibody
response to OspA in white-footed mice. Clin Vaccine Immunol. 2011;18(11):1809-16.
18.   Gingerich MC, Nair N, Azevedo JF, Samanta K, Kundu S, He B, et al. Intranasal
vaccine for Lyme disease provides protection against tick transmitted Borrelia burgdorferi
beyond one year. NPJ Vaccines. 2024;9(1):33.
19.   Lundberg U, Hochreiter R, Timofoyeva Y, Kanevsky I, Meinke A, Anderson AS, et al.
Preclinical Evidence for the Protective Capacity of Antibodies Induced by Lyme Vaccine
Candidate VLA15 in People. Open Forum Infect Dis. 2024;11(9):ofae467.
363

A. Overall Vaccination Schedule

B. Breeding and Vaccination of Dams

1st Breeding: Dams were vaccinated until breading pairs were created (12 weeks)
2nd Breeding: Dams were vaccinated until birth of first pups (24 weeks)
3rd Breeding: Dams were vaccinated until pups reached 2 weeks of age (36 weeks)

C. Birth of pups, tick challenge and euthanasia

Figure 1. Study timeline. (A) Oral vaccination schedule of adult female mice. Pellets coated with the vaccine were exchanged for regular feed at the end of vaccination weeks. Regular feed was provided during rest periods. Controls mice were fed uncoated control pellets following the same schedule. Mice were bled at the begining and end of each vaccination cycle and anti-OspA antibody titer was measured by ELISA. The vaccination cycle was repeated until vaccinated mice reached a minimum anti-ospA antibody titer of OD (blanked) > 3. B) Timeline for breeding of dams. C) Birth of pups, challenge with B. burgdorferi infected ticks and collection of tissues after euthanasia.

Figure 2. Maternal transfer of anti-OspA IgG antibodies to pups.
ELISA quantification of anti-OspA IgG in sera from D24 pups born to
dams vaccinated for different durations. Optical density (OD) at 450nm
(blanked) is shown for each group. 1st Breeding, pups from dams
vaccinated until breeding; 2nd Breeding, pups from dams vaccinated
until birth; 3rd breeding, pups from dams vaccinated until pups reached
two weeks of age. Each point represents an individual pup, with lines
indicating group means ± SD. Statistical analysis by Unpaited t test or
Mann-Whitney test depending of normal distribution of data, *p<0.05,
** p<0.005, *** p<0.0005.

Figure 3. Anti-PepVF antibody responses in terminal bleeds. (A–C) ELISA quantification of anti-PepVF IgG in terminal sera from pups born to dams vaccinated for different durations. Pups were bled > 4 weeks following tick challenge. Optical density (OD) at 450 nm (blanked) is shown for each group. (A) Pups from dams vaccinated until breeding. (B) Pups from dams vaccinated until birth. (C) Pups from dams vaccinated until pups reached two weeks of age. Each data point represents an individual pup, with lines indicating group means ± SD. Statistics by Unpaired t test or Mann Whitney test depending normal distribution of data: ns, not significant, * p<0.05.

Figure 4. Borrelia DNA detection in heart and bladder tissues of pups from dams vaccinated for different durations. (A–C) qPCR quantification of *Borrelia burgdorferi flaB* gene in tissues collected from pups > 4 weeks post tick challenge. (A) *flaB* qPCR in heart tissue from pups born to dams vaccinated until breeding; (B) *flaB* qPCR in heart tissue from pups born to dams vaccinated until birth; (C) *flaB* qPCR in bladder tissue from pups born to dams vaccinated until pups reached two weeks of age. Each data point represents an individual pup, with lines indicating group means. Statistics by Mann Whitney test: ns, not significant, **p<0.005.

Figure 5. *Borrelia burgdorferi* burden in heart cultures from pups born to dams vaccinated until two weeks postpartum (3rd Breeding). Quantification of *B. burgdorferi* in heart tissue cultures from pups at > 4 weeks post-challenge. (A) Enumeration of motile *B. burgdorferi* in heart tissue cultures using a Petroff-Hausser counting chamber under a dark field microscope. (B) qPCR quantification of *B. burgdorferi* flaB gene in heart culture media. Each data point represents an individual pup, with lines indicating group means ± SD. Statistics by Mann Whitney test, *p<0.05 and ** p<0.005.

Figure 6. Anti-OspA IgG in pups and respective dam, at birth before breastfeeding. ELISA quantification of anti-OspA IgG in sera from pups sacrificed at birth, before the opportunity for breast milk intake. Pups were delivered from dams vaccinated until two weeks postpartum (3rd Breeding). Optical density (OD) at 450 nm (blanked) is shown. Each data point represents an individual pup, with lines indicating group means. The cutoff value (dashed line) represents the mean + 3 SD of negative controls.

Figure 7. Anti-OspA IgG transfer from dams to the respective litters. ELISA quantification of anti-OspA IgG in sera from two dams used in the 3rd breeding and the respective litters. Sera from dams were colected at two time-points ~ 4 months apart. Sera from the respective litters were collected on the latter time point, before tick challenge. Each data point represents an individual pup, with lines indicating group means ± SD.