Accurate identification of medulloblastoma subtypes from diverse data sources with severe
batch effects by RaMBat

Mengtao Sun[1], Jieqiong Wang[2] and Shibiao Wan[1]*

[1]Department of Genetics, Cell Biology and Anatomy, University of Nebraska Medical Center,
Omaha, NE, USA

[2]Department of Neurological Sciences, University of Nebraska Medical Center, Omaha, NE,
USA

*Correspondence: Shibiao Wan, swan@unmc.edu

12 Abstract

As the most common pediatric brain malignancy, medulloblastoma (MB) includes multiple
distinct molecular subtypes characterized by clinical heterogeneity and genetic alterations.
Accurate identification of MB subtypes is essential for downstream risk stratification and tailored
therapeutic design. Existing MB subtyping approaches perform poorly due to limited cohorts and
severe batch effects when integrating various MB data sources. To address these concerns, we
propose a novel approach called RaMBat for accurate MB subtyping from diverse data sources
with severe batch effects. Benchmarking tests based on 13 datasets with severe batch effects
suggested that RaMBat achieved a median accuracy of 99%, significantly outperforming state-of-
the-art MB subtyping approaches and conventional machine learning classifiers. RaMBat could
efficiently deal with the batch effects and clearly separate subtypes of MB samples from diverse
data sources. We believe RaMBat will bring direct positive impacts on downstream MB risk
stratification and tailored treatment design.

Keywords: Medulloblastoma; Batch effect; Intra-sample gene ranking; Cancer subtype
identification; RaMBat.

27 Introduction

As the most common malignant pediatric cancer arising in the cerebellar vermis,
medulloblastoma (MB) accounts for around 20% of all pediatric central nervous system (CNS)
neoplasms and over 70% of patients diagnosed are children under the age of 10 [1,2]. Based on the
genetic heterogeneity and phenotypic diversity [3], four major MB subgroups with distinctive
molecular landscapes, somatic mutations and clinical outcomes are defined by the World Health
Organization (WHO) [4], including Sonic HedgeHog (SHH) activated, Wingless-Type (WNT)
activated, and the numerically designated Group 3 and Group 4 [5]. WNT and SHH are named after
signaling pathway disturbances in the WNT (*CTNNB1* mutation [6,7]) and SHH (*PTCH*, *SUFU*, *SMO*
mutation, or *GLI2* amplification [8]) respectively, which are clearly separable with other two
subtypes. Currently, the long-term survival rate of MB has reached 70% based on the standard
treatment, however, there are wide disparities among patient outcomes due to the significant
difference in histologic characteristics of the tumor, age, residual disease and other factors [9,10].
WNT can be almost completely cured under current therapy schemes (5-year event-free survival
greater than 90% [11]), whereas SHH-activated MB occurs more frequently in infants and largely
dependent on the specific genetic features where *TP53* mutation caused tumor has poorer
prognosis [8]. Group 3 and Group 4 do not exhibit subgroup-defining mutations [12], but still have
distinct clinic-biological features where Group 3 patients have the most unfavorable prognosis due
to a higher incidence of "high-risk" features, such as LCA (large cell/anaplastic) histology and
MYC amplification [5,13–15], and Group 4 has an intermediate prognosis with the feature of
isochromosome 17q (*i17q*) [16]. In other words, the survival rates vary significantly with different
MB subtypes. Conventional radiation and chemotherapy have been demonstrated to have long-
term sequelae. Given the heterogeneity of these molecular subtypes, tailoring individualized
treatment plans not only enhances therapeutic efficacy but also minimizes the long-term side
effects associated with conventional therapies [17]. Therefore, accurately identifying MB subtypes
is crucial for guiding effective treatment strategies.

Conventional methods like morphological analysis, immunophenotyping, or molecular
profiling for identifying MB subtypes are costly, time-consuming, or inaccurate. To overcome
these problems, recent studies have demonstrated the feasibility of using next generation
sequencing (NGS) data including transcriptomics (e.g., RNA-seq [18]) and proteomics [19] for
categorizing MB and developed in-silico predictive methods to facilitate a more efficient and
convenient prediction process. Previous studies indicate that identifying MB subtypes based on

transcriptomic data is an efficient and feasible way [20] as it is an essential tool for expression
profiling and provides a way for comprehensive understanding of biological problems [21]. Most
recently, medulloPackage [22] is constructed for rapid and accurate MB subtype identification based
on the RNA-Seq and microarray datasets. medulloPackage identified discriminating features
through gene ratio analysis and an unweighted mean of normalized scores is employed for sample
classification. However, this approach would introduce bias as the data did not necessarily conform
to normal distribution. In addition, another method named Medullo-Model ToSubtype (MM2S) [23]
is developed to enable classification of individual gene expression profiles from MB samples,
including patient samples, mouse models, and cell lines, against well-established molecular
subtypes. MM2S achieves high accuracy for well-characterized subtypes like WNT and SHH.
However, its performance is limited for Group 3 and Group 4, which remain poorly characterized
and more heterogeneous, leading to challenges in accurate classification. In summary, existing MB
subtyping methods directly use absolute gene expression levels without accounting for batch
effects across different cohorts, which may lead to inaccurate classifications. Although
transcriptomics datasets with varying compositions and sample sizes have been made publicly
available, effectively utilizing diverse data sources remains challenging due to the severe batch
effects. Various methods for batch effect adjustment, such as SVA [24], svaseq [25], DWD [26], XPN [27],
and RUV [28], have been proposed, while normalizing across different batches can still distort true
biological signals, particularly when sample distribution between batches is uneven [29–32].

Instead of relying on absolute gene expression levels, strategies like pair-wise analysis of
gene expression (PAGE) [33] leverages the gene expression ratio (GER) [34] which exhibits consistent
patterns within samples to construct the model [35] for batch effect correction [36,37]. The GER strategy
has been widely utilized in various biomedical domains [35,38,39], as its stability allows GER to be
applied across different technological platforms, enhancing its utility in comparative omics studies.
For instance, Zhao et al. [37] developed DRGpair which based on intra-sample paired GER for
ovarian cancer detection to overcome the limitations of absolute gene expression values including
batch effects and biological heterogeneity. Another transfer learning method, scPAGE2 [36], based
on intra-sample GER using PAGE strategy to enhance generalization for sepsis diagnosis. By
converting discrete gene pair differences into continuous GER values, scPAGE2 integrates both
single-cell and bulk RNA-seq data. Moreover, recent studies [38–40] suggest that ranking gene
expression, unlike absolute values which are sensitive to batch effects, provided a more stable

representation of gene expression patterns. Additionally, ranking is scale-independent and it
proceeds multi-cohort integration regardless of the difference among profile measurements.

Herein, we propose a novel and accurate computational approach called RaMBat for MB
subtyping across diverse transcriptomics datasets with severe batch effects, based on intra-sample
gene expression ranking information as well as a series of steps for gene-rank based information
processing, enabling effective batch effect correction and robust integration of multiple bulk
transcriptomics datasets. Based on 13 heterogeneous transcriptomics datasets, we demonstrate the
accuracy and efficiency of RaMBat by comparing it with state-of-the-art methods like
medulloPackage [22] and MM2S [23] as well as other machine learning (ML) methods equipped with
the conventional batch correction method ComBat [41]. Additionally, we have developed an R
package RaMBat that is available on GitHub at https://github.com/wan-mlab/RaMBat.

Results

Study design for MB identification

To get subtype-specific signatures for MB samples across different cohorts with severe
batch effects, we proposed RaMBat to identify MB subtype accurately from diverse
transcriptomics data sources based on the gene expression ranking. The overall workflow
contained four key steps (**Fig. 1)** which were designed to efficiently capture both global and local
dependencies within high-dimensional intra-sample gene expression data from different cohorts,
including (1) intra-sample gene expression analysis, (2) reversed expression pattern analysis, (3)
informative GERs selection and (4) MB sample subtyping. As input, transcriptomics data was
firstly converted into gene expression ranking followed with intra-sample gene expression analysis
which was a two-step process including gene rank analysis and two-sided t-test (See Methods; **Fig.**
1A). Intra-sample gene expression analysis was performed between each pair of subgroups to
identify genes that were specifically and significantly associated with each subtype. Next, the
reversed expression pattern analysis (See Methods; **Fig. 1B**) was performed as a two-step process
including reversal ratio analysis and Fisher's exact test. Reversal ratio analysis leveraged rank
levels between every possible pair of genes to retrieve subtype-specific GERs, combining each
two genes together to construct the GER matrix. GER contained in each subtype was compared

exhaustively to find the reversed gene expression pattern between each two subtypes cross samples.

To enhance the accuracy for MB subtyping, LASSO [42] (see Methods; **Fig. 1C**) was applied to

identify non-zero coefficients GERs that were strongly associated with subgroups. Similarly, as

described above, comparisons between each pair of subgroups were performed and the identified

GERs formed the foundation of RaMBat.

Fig. 1 The workflow of RaMBat based on GERs for MB subtyping. RaMBat consisted of four

steps for GERs selection and MB sample subtyping. (**A**) Perform gene ranking analysis. (**B**)

Calculate reversal ratio values for GERs filtering. (**C**) Select GERs using LASSO. (**D**) Identify

MB subtypes based on selected GERs.

In this study, a large-scale microarray dataset (GSE85217, 763 samples) was used to

training. To ensure robust performance evaluation, RaMBat was tested on 13 independent test

datasets which were generated using different platforms and exhibited significant scale disparities,

necessitating standardized normalization to enhance comparability. 1035 GERs, corresponding to

738 unique genes (**Fig. 1A-C**), were identified. WNT had the highest number of GERs, totaling

488, followed by 331, 111, 105 GERs for SHH, Group 3 and Group 4, respectively. Subtype

information was determined by calculating the average GER score for each subtype within each

sample, with the subtype corresponding to the highest average GER score being assigned as the

subtype for the sample (see Methods; **Fig. 1D**).

Benchmarking with the state-of-the-art methods for identifying MB subtypes

To demonstrate the superiority of RaMBat, we benchmarked RaMBat with state-of-the-art methods including medulloPackage and MM2S as well as seven ML classifiers for MB subtyping across 13 independent datasets. Among them, medulloPackage was developed based on RNA-seq and microarray datasets for MB subtype prediction and MM2S was constructed based on microarray data for classification of patient samples, mouse models, and cell lines. Additionally, we trained and optimized seven ML classifiers, including SVM, LR, RF, XGBoost, KNN, NB and MLP, as benchmarks for comparison. In general, RaMBat consistently performed the best among all the benchmarking methods for MB subtyping. Specifically, RaMBat outperformed medulloPackage and MM2S in eight (**Fig. 2B-E, G, I, K, N**) of 13 datasets (**Fig. 2B-N**) and 11(**Fig. 2B-F, H, J-N**) out of 13 datasets respectively. For the remaining five (**Fig. 2F, H, J, L, M**) and two (**Fig. 2G, I**) datasets, respectively, RaMBat achieved the same accuracy as these methods. The overall accuracy for RaMBat across all datasets was 99.02%, where three Group 3 samples in GSE21140 and one Group 3 sample in GSE10327 were misclassified as Group 4, and one Group 4 sample in GSE74195 was misclassified as Group 3. As previously noted, the high similarity in transcriptomic and genomic profiles between Group 3 and Group 4 resulted in the molecular ambiguity and histological misclassification. However, the overall accuracy was still 11.8% (absolute) higher than MM2S and 7.9 (absolute) higher than medulloPackage. Additionally, the overall subtyping performance of RaMBat was better than all seven ML classifiers (**Fig. 2A**). The lowest accuracy achieved by RaMBat was 95% in GSE74195 (**Fig. 2J**), and the highest accuracy of 100% was recorded in ten datasets, including GSE73038 (**Fig. 2B**), GSE67850 (**Fig. 2E**), GSE50765 (**Fig. 2F**), GSE49243 (**Fig. 2G**), GSE30074 (**Fig. 2H**), GSE37382 (**Fig. 2I**), GSE62803 (**Fig. 2K**), GSE41842 (**Fig. 2L**), GSE50161 (**Fig. 2M**), GSE12992 (**Fig. 2N**). When evaluating performance by subtype, RaMBat always outperformed medulloPackage and MM2S on all four subtypes across all independent datasets in terms of accuracy, precision, sensitivity, F1-score, MCC, G-measure, AUC, specificity, and Jaccard index (**Fig. 3**).

Fig. 2 RaMBat significantly outperformed the state-of-the-art methods for MB subtyping based on 13 independent test datasets with severe batch effects. (**A**) Overall accuracy of RaMBat, medulloPackage, MM2S and other 7 ML classifiers were compared across all 13 independent datasets. Comparative performance of RaMBat, medulloPackage, MM2S and other 7 ML classifiers based on (**B**) GSE73038, (**C**) GSE10327, (**D**) GSE21140, (**E**) GSE67850, (**F**) GSE50765, (**G**) GSE49243, (**H**) GSE30074, (**I**) GSE37382, (**J**) GSE74195, (**K**) GSE62803, (**L**) GSE41842, (**M**) GSE50161, (**N**) GSE12992 in terms of accuracy, respectively. MLP, multilayer perceptron; KNN, k-nearest neighbors; XGBoost, eXtreme gradient boosting; RF, random forest; LR, logistic regression; SVM, support vector machine; MM2S, medullo-model to subtypes.

Fig. 3 RaMBat achieved superior subtype-specific performance compared to state-of-the-art methods across 13 independent datasets for MB subtyping. Comparing performance of RaMBat with medulloPackage and MM2S across datasets in terms of accuracy, precision, sensitivity, F1-Score, MCC, G-measure, AUC, specificity and Jaccard index for SHH (**A**), WNT (**B**), Group 3 (**C**) and Group 4 (**D**). Metrics were obtained based on the prediction result of 508 samples in all 13 independent test datasets. MCC, Matthews correlation coefficient; AUC, area under the curve; Jaccard, Jaccard index.

To further evaluate the impact of batch effects and the ability of RaMBat to efficiently integrate multi-source data with severe batch effects for MB subtyping, we combined all 13 independent test datasets and visualized the sample distribution using t-SNE (**Fig. 4**). When applied to raw gene expression data, conventional t-SNE visualization exhibited severe batch effects, with samples clustering predominantly by datasets rather than by biological subtypes (**Fig. 4A**). However, these technical variations obscured the biologically meaningful clustering of MB subtypes as the data did not cluster together according to major MB subtypes (**Fig. 4B**). In contrast, RaMBat effectively removed batch effects, enabling a clear separation of MB subtypes across diverse datasets (**Fig. 4C-D**). After integrating predicted subtype information from RaMBat into visualization, samples were well-integrated across different datasets (**Fig. 4C**), indicating that the batch effects were successfully removed. More importantly, RaMBat presented a clear

visualization global landscape of all MB samples with all major subtypes distinctly separable (**Fig.**

4D). These results indicated that RaMBat can accurately integrate diverse bulk transcriptomics

datasets, effectively removing batch effects while maintaining the underlying biological signals

crucial for MB subtype classification.

Fig. 4 RaMBat enabled more effective visualization of MB subtypes than conventional t-SNE

across 13 batch-affected independent datasets. (**A**) Samples were labeled by batch based on all

samples in independent test datasets without normalization. (**B**) Samples were labeled by subtype

based on all samples in independent test datasets without normalization. (**C**) RaMBat visualization

in which samples were labeled by batch based on all samples in normalized independent test

datasets. (**D**) RaMBat visualization in which samples were labeld by subtype based on all samples

in normalized independent test datasets.

Subtype-specific GERs selection

To investigate whether the marker genes identified by RaMBat were biologically meaningful, we used the training dataset as the basis for GER selection. The absolute expression levels were converted into intra-sample rankings, followed by gene rank analysis and two-sided t-test. There were 1055 ranking down genes and 1257 ranking up genes between SHH and Group 3, 1123 ranking down genes and 1233 ranking up genes between SHH and Group 4, 1220 ranking down genes and 1201 ranking up genes between SHH and WNT, 1221 ranking down genes and 1325 ranking up genes between WNT and Group 3, 1476 ranking down genes and 1522 ranking up genes between WNT and Group 4, 632 ranking down genes and 561 ranking up genes between Group 3 and Group 4 (**Supplementary Fig. S1A-F**). After intersecting these ranked up and down genes, 710, 880, 346, and 345 differentially ranked genes were identified for the SHH, WNT, Group 3, and Group 4, respectively.

Based on the differentially ranked genes in each subtype, 251,695, 386,760, 59,685 and 59,340 GERs for SHH, WNT, Group 3 and Group 4 were constructed, respectively. For each subtype, we performed the reversed expression pattern analysis between each pair of subgroups. For each comparison, a contingency table was constructed for the Fisher's exact test. A total of 904, 2,381, 347 and 456 GERs (**Fig. 1C**) were screened for SHH, WNT, Group 3 and Group 4 respectively. Based on the reversed GERs, LASSO feature selection was performed to select the most important features which results in a final list of 1,035 GERs, corresponding to 738 unique genes (**Fig. 1C**). To comprehensively analyze the subtype-specific gene expression landscape in MB, the gene frequency across all 1,035 GERs for each subtype was also analyzed (**Fig. 5A-D**). Specifically, the top 10 most frequent genes in each subtype were labeled. For example, potential marker genes identified for SHH include *NEUROG1* and *ATOH1* [43], while *FZD10* was highlighted as a potential marker for the WNT subtype [44]. To further investigate the gene expression patterns underlying MB subtype differentiation, differential expressed gene (DEG) analysis was performed and DEGs for each subtype were identified (**Fig. 5E-H**). To assess the overlap between RaMBat selected genes and DEGs, the identified DEGs with 738 unique genes were intersected (**Fig. 5I-P**). Notably, all top 10 most frequently ranked genes (**Fig. 5A-D**) were present within the intersected genes, except for *ZDHHC15* in Group 3, demonstrating strong consistency between RaMBat's feature selection and conventional DEG analysis. Notably, RaMBat identified important genes that were not captured by conventional DEG analysis, such as *BOC*, which has been implicated in promoting the progression of early MB to more advanced stage. This finding

suggested that RaMBat effectively captured subtype-specific molecular signatures, boosting its

reliability in MB classification. To further validate the robustness of these findings, the expression

patterns of the top 10 intersected genes of each subtype were tested using heatmaps. Distinct

expression patterns of these 738 differentially ranked genes (**Fig. 6A)** evidenced their ability to

effectively distinguish each MB subtype from others and enable clear separation into specific

subgroups. The hierarchical clustering results further confirmed the biological relevance of the

selected genes in MB subtyping. Specifically, based on data from GSE85217, the expression

profiles of the top 10 most frequently ranked intersected genes were distinct across MB subtypes

(**Fig. 6B-E)**. The results indicated specific expression patterns that effectively distinguished MB

subtypes. Additionally, the expression level of upregulated DEGs, including *PDLIM3*,

TMEM132C, *GABRA5* and *RBM24*, across all 13 independent test datasets highlighted their

specific overexpression in SHH, WNT, Group 3 and Group 4 respectively (**Fig. 6F-I).**

Fig. 5 MB subtype-specific differential expression analysis showed strong concordance between RaMBat-selected features and conventional DEG analysis. The frequency distribution of all unique genes included in GERs for SHH (**A**), WNT (**B**), Group 3 (**C**), and Group 4 (**D**) respectively. The top 10 most frequently ranked genes in SHH, WNT, Group 3, and Group 4 were labeled. Differential expressed genes for SHH (**E**), WNT (**F**), Group 3 (**G**), and Group 4 (**H**) were identified. The x-axis represented the log2 fold change while y-axis showed -log10(p-value). Red and blue dots indicated significantly upregulated and downregulated genes respectively. Triangles highlighted top 5 upregulated and downregulated genes in each comparison. (**I-L**) Genes consistently identified by both DEG analysis and RaMBat. The intersections represented genes identified by both DEG analysis and RaMBat for SHH (**I, M**), WNT (**J, N**), Group 3 (**K, O**), Group 4 (**L, P**), where panels **I-L** showed upregulated genes, and panels **M-P** showed downregulated genes.

Fig. 6 Identified differentially ranked genes showed distinct subtype-specific gene expression patterns. (**A**) The expression pattern of 738 unique genes associated with 4 molecular subtypes across samples in GSE85217. The absolute gene expression levels of the top 10 frequently ranked genes for SHH (**B**), WNT (**C**), Group 3 (**D**), and Group 4 (**E**) respectively. For all heatmaps, rows corresponded to genes, and columns corresponded to samples in GSE85217. Color scale ranged from blue (low expression) to red (high expression). (**F-I**) The expression level of the top upregulated DEG for SHH (**F**), WNT (**G**), Group 3 (**H**), and Group 4 (**I**) across 13 independent datasets with normalization. Each point represented a sample, colored by expression intensity (red: high, grey: low).

Discussion

Medulloblastoma is a highly malignant pediatric brain cancer in cerebellum with peak incidence between ages 3-7 and a slight male predominance [3]. Based on the molecular

characteristics, it is mainly classified into four subtypes: SHH, WNT, Group 3 and Group 4, each

with specific genetic profiles and prognostic outcomes. These four subtypes have varied symptoms

to each other and exhibit significant variations in treatment response and prognosis, therefore,

classifying patients into accurate subtype plays significant roles in the following therapy.

Computational methods are designed to efficiently predict accurate subtype information. However,

the absolute gene expression level deviated due to batch effect and biological heterogeneity even

if using appropriate normalization methods. Previous studies [45,46], like iPAGE [33], has evidenced

the analysis of the intra-sample relative expression level between genes is robust and generalizable

in detecting biological signals. Furthermore, gene pair-based analysis is able to perform cross-

platform comparison by integrating absolute expression from different cohorts and consequently

stable across experimental assays and platforms [33,47]. Building upon these advantages, we develop

RaMBat, which achieves better prediction performance than existing state-of-the-art methods for

MB subtyping. The benchmark comparison among RaMBat, medulloPackage, MM2S and other

optimized ML classifiers demonstrates that analyzing differentially ranked genes provides robust

and reliable insights for MB subtyping, complementing traditional differential expression-based

approaches.

RaMBat presents high accuracy across 13 independent test datasets, with the most

misclassifications occurring between Group 3 and Group 4. Among all test samples, one Group 4

sample is misclassified as Group 3, and four Group 3 samples are misclassified as Group 4.

Previous studies have noted that these two subtypes exhibit highly similarity in transcriptomic and

genomic profiles, contributing to the ambiguity between these two subtypes. Although recent

studies have uncovered the heterogeneity within Group 3 and Group 4 and even suggesting their

division into eight subtypes [16], patients with Group 3 or Group 4 are currently still treated with

same therapeutic regimen [1,3]. However, there are ongoing studies attempting to stratify treatment

by molecular subtype to improve outcomes.

RaMBat effectively identifies subtype-specific marker genes, demonstrating its robustness

in selecting biologically relevant features for MB subtyping. Specifically, selected GERs contain

NEUROG1, GLI1, ATOH1 associated with the SHH signature. *NEUROG1* and *ATOH1* are the top-

selected genes with 40 and 22 GERs respectively, whereas *GLI1* is only involved in one GER. This

aligns with the study published in 2007 which indicates *ATOH1, NEUROG1* and *GLI1* might be

relevant biomarkers for SHH subtype [43]. *GLI1* is a transcription factor and a key downstream effector of SHH signaling, directly linking it to SHH-driven tumorigenesis. *ATOH1* is a well-established marker of cerebellar granule precursor cells, and its expression is strongly associated with medulloblastomas that arise from these cells. In contrast, *NEUROG1* highly expressed tumors are suggested to originate from cerebellar ventricular zone progenitors rather than granule cell precursors. Additionally, canonical over-expressed genes *HHIP, BOC* and *SFRP1* are also included in the GERs [48]. The WNT subtype of medulloblastoma is closely linked to the activation of the WNT signaling pathway, primarily through mutations that stabilize beta-catenin and drive oncogenesis. Five WNT signaling genes are observed within GERs associated with the WNT subtype, including *SP5, WIF1, FZD10, CER1,* and *PRKCB*, which play various roles in the WNT signaling pathway. For example, the activation of the WNT pathway leads to the upregulation of *SP5*, which in turn promotes the expression of genes that contribute to tumorigenesis [49,50]. Its presence indicates active WNT signaling, which is characteristic of the WNT subtype. The *WIF1* plays a significant role in medulloblastoma by acting as a negative regulator of the WNT signaling pathway [51]. As a receptor in the Frizzled family, *FZD10* mediates WNT ligand binding, activating the canonical WNT/β-catenin pathway [44]. Although less is known about the specific drivers, pathways, and biomarkers that discriminate Group 3 and Group 4 subtypes, it is established that these tumors share certain common pathways. For example, *UNC5D* has been identified as a key gene associated with neuronal migration in MB, showing specificity for certain subtypes and being present in GERs [51,52]. Additionally, specific chromosomal aberrations affecting *AFF3* may contribute to tumorigenesis by disrupting regulatory pathways. The *AFF3* gene can potentially discriminate these two subtypes, as it is downregulated in Group 3, but upregulated in Group 4. As expected, it only presents in GERs for Group 3 [51]. Thus, the detailed function of *AFF3* may worth more investigation.

Unlike absolute gene expression levels which are sensitive to batch effects, RaMBat is scale-independent and multi-cohort integration regardless of the difference among profile measurements. The data visualization by RaMBat (**Fig. 4**) further confirmed that it can efficiently eliminate batch effects and clearly separate MB subtypes, unlike conventional approaches such as t-SNE, which are susceptible to batch-induced variations. We admit that one limitation of our study is its high computational cost and time-consuming nature, particularly when handling large-scale datasets, which significantly reduces efficiency. Overall, this study proposes RaMBat, a novel

approach leveraging ranking gene expression to select robust and subtype-specific GERs for

accurate MB subtyping across diverse transcriptomics datasets with severe batch effects.

Benchmarking across 13 datasets demonstrates that RaMBat outperforms state-of-the-art MB

subtyping methods, achieving a high median accuracy of 99.02%, while also providing effective

batch effect correction. We believe that RaMBat will significantly enhance downstream risk

stratification for MB and contribute to the design of more personalized treatment strategies.

Additionally, we also expect that RaMBat will be highly effective in addressing many biomedical

problems with batch effects for various types of pediatric, adolescent, and young adult (AYA)

cancers.

Methods

Datasets

A transcriptomic dataset (GSE85217) was used to train the RaMBat, characterized using

Affymetrix Human Gene 1.1 ST Array and downloaded from the Gene Expression Omnibus [53]

database (GEO, https://www.ncbi.nlm.nih.gov/geo). For model performance evaluation, 13

datasets were downloaded through the *getGEO* function in R with following accession numbers:

GSE10327, GSE12992, GSE21140, GSE30074, GSE37382, GSE41842, GSE49243, GSE50161,

GSE50765, GSE62803, GSE67850, GSE73038, GSE74195. Specifically, to address scale

disparities, datasets were normalized as $(x - \bar{x})/s$, where each gene expression value x was

subtracted by the mean value $\bar{x}$ of the column and dividing it by the standard deviation s of the

column. In both the training and independent test datasets, Group 4 samples represented the largest

proportion, followed by SHH, Group 3, and WNT. Additionally, the number of samples for each

subtype was consistently higher in the training dataset compared to the independent test dataset.

The training dataset, GSE85217, included 763 MB samples (223 SHH, 70 WNT, 144 Group 3, and

326 Group 4), while the independent test datasets contained 508 samples (177 SHH, 42 WNT, 92

Group 3, and 192 Group 4) (**Supplementary Table 1**).

To identify MB subtypes based on multiple bulk transcriptomics data, an accurate MB

subtyping approach based on gene ranking for heterogeneous data with severe batch effects, named

RaMBat, was developed. RaMBat contained four key steps (**Fig. 1A**), including intra-sample gene

expression analysis, reversed expression pattern analysis, informative GERs selection, and finally

MB sample subtyping. It leveraged gene expression rankings instead of absolute gene expression

levels to address batch effects from different data sources. Based on 13 transcriptomics datasets

with severe batch effects, we demonstrated that RaMBat significantly outperformed state-of-the-

art methods like medulloPackage and MM2S as well as seven optimized ML classifiers equipped

with the conventional batch correction method ComBat in terms of prediction accuracy for

identifying MB subtypes.

Intra-sample gene expression analysis

Based on the ascending order, from the smallest to the largest values, we ordered all genes

inside the dataset. Subsequently, we transformed the absolute gene expression levels

$$[G_1, G_2, \ldots, G_n]$$

into rank values,

$$[T_1, T_2, \ldots, T_n]$$

where G was the absolute value of gene expression, n was the total number of genes, T was the

rank value of gene within one sample, where ranking was performed in ascending order. The gene

with the lowest expression value was assigned a rank of 1, while the gene with the highest

expression value was assigned a rank of n. We then divided the rank matrix of GSE85217 into four

submatrices corresponding to the SHH, WNT, Group 3, and Group 4, based on sample labels. For

each gene, we calculated the gene rank difference and a p value between each combination of

submatrices, resulting in six different comparisons: SHH vs. WNT, SHH vs. Group 3, SHH vs.

Group 4, WNT vs. Group 3, WNT vs. Group 4, and Group 3 vs. Group 4. The mean rank of each

gene across all samples within one subtype was calculated and compared to its mean rank in

another subtype, with the gene rank representing the difference between these two mean rank

scores. A two-sided t-test was then performed to assess the statistical significance of the difference

between subtypes. We applied thresholds of $p < 0.05$ and a gene rank > 2314 (representing 11% of

the total number of genes) to identify significant differentially ranked genes.

Reversed expression pattern analysis

To construct gene pairs based on the differentially ranked genes, we followed iPAGE [33] protocols to construct GERs. Specifically, for each sample, the rank values of all possible gene pairs were compared, encoding their relationships as binary indicators, represented by

$$R^{(k)} = \left(R_{1,2}^{(k)}, R_{1,3}^{(k)}, \dots, R_{1,n}^{(k)}, R_{2,3}^{(k)}, R_{2,4}^{(k)}, \dots, R_{2,n}^{(k)}, \dots, R_{(n-1)n}^{(k)} \right) \tag{1}$$

$$= \left(s\left(T_1^{(k)} - T_2^{(k)}\right), s\left(T_1^{(k)} - T_3^{(k)}\right), \dots, s\left(T_1^{(k)} - T_n^{(k)}\right), \dots, s\left(T_2^{(k)} - T_3^{(k)}\right), s\left(T_2^{(k)} \right.$$

$$\left. - T_4^{(k)}\right), \dots, s\left(T_2^{(k)} - T_n^{(k)}\right), \dots, s\left(T_{n-1}^{(k)} - T_n^{(k)}\right) \right)$$

where GERs (R) was constructed by gene rankings (T) in a sample k and function s denoted the process of converting gene rankings into GER values as following:

$$R_{i,j} = s(T_i - T_j) = \begin{cases} 1, if\ T_i > T_j \\ 0, if\ T_i \leq T_j \end{cases} \tag{2}$$

Every two different genes within each sample were subtracted to convert the rank matrix into GER matrix. Then, GERs with a significant difference in their relative expression $R_{i,j}$ between each two subtypes were extracted.

Two situations were taken into consideration, where $R_{i,j} = 1$ and $R_{i,j} = 0$. We counted the number of samples with $R_{i,j} = 1$ and $R_{i,j} = 0$ in the first subtype group as $f_{1,1}$ and $f_{1,2}$ respectively, then in the second subtype group, the number of samples with $R_{i,j} = 1$ and $R_{i,j} = 0$ as $f_{2,1}$ and $f_{2,2}$ respectively. Finally, the contingency table was shown as follows:

	$T_i > T_j$	$T_i \leq T_j$
Selected subtype	$f_{1,1}$	$f_{1,2}$
Rest subtype	$f_{2,1}$	$f_{2,2}$

For example, for the comparison between SHH and WNT, $f_{1,1}$ and $f_{1,2}$ were the number of samples with $R_{i,j} = 1$ and $R_{i,j} = 0$ respectively in SHH, and $f_{2,1}$ and $f_{2,2}$ were the number of samples with $R_{i,j} = 1$ and $R_{i,j} = 0$ respectively in WNT.

$$reversal\ ratio = \frac{f_{1,1} + f_{2,2}}{f_{1,1} + f_{1,2} + f_{2,1} + f_{2,2}} \tag{3}$$

To calculate the reversal ratio, we considered 12 different comparisons, with each subtype having

three pairwise comparisons. For example, when selected specific GERs for SHH, the comparisons

were with WNT, Group 3, and Group 4, respectively. We established different thresholds for the

reversal ratio based on the subtype comparisons. A threshold of 0.95 was set for the six pairwise

comparisons within the SHH and WNT subtypes. For comparisons between Group 3 and SHH,

Group 4 and SHH, we set the threshold at 0.9. A threshold of 0.85 was used for comparisons

between Group 3 and WNT, Group 4 and WNT, Group 4 and Group 3. Lastly, a threshold of 0.8

was applied specifically for the comparison between Group 3 and Group 4. At the same time, we

performed Fisher's exact test to obtain the p value for each of the twelve comparisons. The p value

was calculated by

$$p = \frac{(f_{1,1} + f_{2,2})!\,(f_{2,1} + f_{2,2})!\,(f_{1,1} + f_{2,1})!\,(f_{2,2} + f_{2,2})!}{f_{1,1}!\,f_{2,2}!\,f_{2,1}!\,f_{2,2}!\,t!} \tag{4}$$

432   , where $t = f_{1,1} + f_{1,2} + f_{2,1} + f_{2,2}$ and the threshold for p value was set as 0.05. For each subtype,

433   three sets of selected GERs were obtained which were then intersected to identify GERs

434   consistently presenting across all three pairwise comparisons within that subtype. At the end of

435   this step, subtype-specific GERs were selected for SHH, WNT, Group 3 and Group 4, respectively,

436   based on a reversal ratio exceeding the threshold and a p value > 0.05 (**Fig. 1B**).

Selecting informative GERs

To reduce the computational complexity of the subsequent MB sample subtyping step

based on GERs and boost the stability of this method, we furthermore refined the number of GERs

using LASSO (Least Absolute Shrinkage and Selection Operator) [54]. LASSO naturally performed

feature selection by shrinking the coefficients of less important features to zero, effectively

eliminating them from the model. We conducted LASSO 12 times to select the most important

GERs corresponding to three pairwise comparisons for each of the four subtypes (**Fig. 1C**).

Similarly, we intersected three sets of selected GERs for each subtype. In the end, 1035 GERs,

corresponding to 738 unique genes, with non-zero coefficients were filtered which were then used

as the input for sample subtyping. LASSO was performed using R package "glmnet" [55].

Additionally, based on these 1035 GERs, heatmap was plotted using the R package "pheatmap" [56].

RaMBat Visualization

RaMBat utilized a combination of two key matrices: a subtype score matrix and a sample-

to-subtype matrix derived from prediction results. The former matrix was obtained through the

RaMBat subtype identification workflow (**Fig. 1**), in which GERs were used to calculate subtype-

specific scores for each sample. This matrix captured the continuous prediction confidence of each

sample across the four MB subtypes (SHH, WNT, Group 3, Group 4), forming a $R \times C$ matrix,

where R is the number of samples and C is the number of subtypes. A sample-to-subtype matrix

was constructed using one-hot encoding based on the predicted subtype labels derived from the

subtype score matrix. Specifically, the subtype with the highest score for a given sample was

assigned a value of 1, while all other subtypes were assigned 0. This results in a binary matrix of

the same dimension ($R \times C$), representing discrete subtype assignments. These two matrices were

then concatenated along the feature axis to form a unified visualization matrix, which was used as

input for t-SNE to generate a low-dimensional embedding. This integration strategy enabled the

combination of both quantitative prediction certainty and categorical class identity, thereby

enhancing the biological interpretability and subtype-level separation in the visualization space.

Benchmarking with the state-of-the-art methods

To assess the prediction performance of RaMBat, we initially downloaded 13 independent

test datasets with clear subtype information, containing 508 samples (**Supplementary Table 1**).

The subtyping process for a given sample was composed of three stages, 1) convert the absolute

gene expression into GERs based on the selected unique genes, 2) filter to the selected GERs in

the model and calculate the mean score of GERs for each subtype, 3) determine the subtype with

the highest score and assign it to the sample (**Fig. 1D**). Finally, we compared the performance of

RaMBat with state-of-the-art MB classifiers, including the medulloPackage [22] and MM2S [23,57].

medulloPackage was also developed using GERs but based on the differentially expressed genes

(DEGs) selected by limma package in R. In contrast, MM2S employed a k-nearest neighbor (KNN)

[58] classifier based on the gene-set enrichment analysis ranked matrix. Additionally, to compare the

performance of RaMBat with machine learning methods, 7 classifiers, including support vector

machine (SVM) [59], logistic regression (LR) [60], random forest (RF) [61], XGBoost [62], KNN, naïve

bayes (NB) , multilayer perceptron (MLP) [54] were trained and optimized. The training dataset and

independent test datasets which were normalized and then batch effect corrected by the *combat*

function in R were used for training and testing of these ML classifiers. For performance evaluation,

we considered nine major metrics: accuracy, precision, sensitivity, specificity, area under the curve

(AUC), Jaccard index (Jaccard), Matthews correlation coefficient (MCC), G-measure and F1-

score.

Supplementary Data

Supplementary Fig. S1 Differential rank gene analysis within RaMBat. (A-F) The

differentially ranked gene for specific medulloblastoma subtypes. The x-axis represented the gene

rank difference where y-axis showed -log10(p-value). Red dots indicated significantly up-ranked

genes and blue dots represented significantly down-ranked genes. Triangles represented top 5

up/down-ranked genes. Differentially ranked genes between SHH and WNT (A), SHH and Group

3 (B), SHH and Group 4 (C), WNT and Group 3 (D), WNT and Group 4 (E), Group 3 and Group

4 (F).

Supplementary Table 1 Comparing detailed information for each dataset. Summary of MB

subgroup distribution in training and independent test datasets. Num, number of samples.

Competing interests

498   The authors declare no competing interests.

Funding

Research reported in this publication was supported by the Office Of The Director, National

Institutes Of Health of the National Institutes of Health under Award Number R03OD038391, and

by the National Cancer Institute of the National Institutes of Health under Award Number

P30CA036727. This work was supported by the American Cancer Society under award number

IRG-22-146-07-IRG, and by the Buffett Cancer Center, which is supported by the National Cancer

Institute under award number CA036727. This work was also partially supported by the National

Institute of General Medical Sciences under Award Numbers P20GM103427. This study was in

part financially supported by the Child Health Research Institute at UNMC/Children's Nebraska.

This work was also partially supported by the University of Nebraska Collaboration Initiative

Grant from the Nebraska Research Initiative (NRI). The content is solely the responsibility of the

authors and does not necessarily represent the official views from the funding organizations.

Author contributions

514   SW conceived and designed the study. SW and MS developed the method, performed the

515   experiments and analyzed the data. All authors participated in writing the paper. SW and MS

516   revised the paper. The manuscript was approved by all authors.

Data availability

All the data used in this manuscript are publicly available in the corresponding references.

Code availability

RaMBat is available as an R package in GitHub at https://github.com/wan-mlab/RaMBat.

Reference

1. Juraschka, K. & Taylor, M. D. Medulloblastoma in the age of molecular subgroups: a review. *J Neurosurg Pediatr* **24**, 353–363 (2019).

2. Martin, A. M., Raabe, E., Eberhart, C. & Cohen, K. J. Management of Pediatric and Adult Patients with Medulloblastoma. *Curr. Treat. Options in Oncol.* **15**, 581–594 (2014).

3. Northcott, P. A. *et al.* Medulloblastoma. *Nat Rev Dis Primers* **5**, 11 (2019).

4. Louis, D. N. *et al.* The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol* **131**, 803–820 (2016).

5. Taylor, M. D. *et al.* Molecular subgroups of medulloblastoma: the current consensus. *Acta Neuropathol* **123**, 465–472 (2012).

6. Clifford, S. C. *et al.* Wnt/Wingless pathway activation and chromosome 6 loss characterize a distinct molecular sub-group of medulloblastomas associated with a favorable prognosis. *Cell Cycle* **5**, 2666–2670 (2006).

7. Ellison, D. W. *et al.* beta-Catenin status predicts a favorable outcome in childhood medulloblastoma: the United Kingdom Children's Cancer Study Group Brain Tumour Committee. *J Clin Oncol* **23**, 7951–7957 (2005).

8. Kool, M. *et al.* Genome sequencing of SHH medulloblastoma predicts genotype-related

response to smoothened inhibition. *Cancer Cell* **25**, 393–405 (2014).

9. Thomas, A. & Noël, G. Medulloblastoma: optimizing care with a multidisciplinary approach.

JMDH **Volume 12**, 335–347 (2019).

10. Fellay, C. N. *et al.* Medulloblastomas in adults: prognostic factors and lessons from

paediatrics. *Curr Opin Neurol* **24**, 626–632 (2011).

11. Choi, J. Y. Medulloblastoma: Current Perspectives and Recent Advances. *Brain Tumor Res*

Treat **11**, 28–38 (2023).

12. Northcott, P. A. *et al.* The whole-genome landscape of medulloblastoma subtypes. *Nature*

547, 311–317 (2017).

13. Kool, M. *et al.* Molecular subgroups of medulloblastoma: an international meta-analysis of

transcriptome, genetic aberrations, and clinical data of WNT, SHH, Group 3, and Group 4

medulloblastomas. *Acta Neuropathol* **123**, 473–484 (2012).

14. Northcott, P. A. *et al.* Medulloblastomics: the end of the beginning. *Nat Rev Cancer* **12**, 818–

834 (2012).

15. Ryan, S. L. *et al.* MYC family amplification and clinical risk-factors interact to predict an

extremely poor prognosis in childhood medulloblastoma. *Acta Neuropathol* **123**, 501–513

(2012).

16. Sharma, T. *et al.* Second-generation molecular subgrouping of medulloblastoma: an

international meta-analysis of Group 3 and Group 4 subtypes. *Acta Neuropathol* **138**, 309–

326 (2019).

17. Ramaswamy, V. *et al.* Risk stratification of childhood medulloblastoma in the molecular era:

the current consensus. *Acta Neuropathol* **131**, 821–831 (2016).

18. Gibson, P. *et al.* Subtypes of medulloblastoma have distinct developmental origins. *Nature*

468, 1095–1099 (2010).

19. Godbole, S. *et al.* Multiomic profiling of medulloblastoma reveals subtype-specific

targetable alterations at the proteome and N-glycan level. *Nat Commun* **15**, 6237 (2024).

20. Shih, D. J. H. *et al.* Cytogenetic prognostication within medulloblastoma subgroups. *J Clin*

Oncol **32**, 886–896 (2014).

21. Rani, B. & Sharma, V. K. Transcriptome profiling: methods and applications- A review. *AG*

38, (2017).

22. Rathi, K. S. *et al.* A transcriptome-based classifier to determine molecular subtypes in

medulloblastoma. *PLoS Comput Biol* **16**, e1008263 (2020).

23. Gendoo, D. M. A. & Haibe-Kains, B. MM2S: personalized diagnosis of medulloblastoma

patients and model systems. *Source Code Biol Med* **11**, 6 (2016).

24. Leek, J. T., Johnson, W. E., Parker, H. S., Jaffe, A. E. & Storey, J. D. The sva package for

removing batch effects and other unwanted variation in high-throughput experiments.

Bioinformatics **28**, 882–883 (2012).

25. Leek, J. T. svaseq: removing batch effects and other unwanted noise from sequencing data.

Nucleic Acids Research **42**, e161–e161 (2014).

26. Benito, M. *et al.* Adjustment of systematic microarray data biases. *Bioinformatics* **20**, 105–

114 (2004).

27. Shabalin, A. A., Tjelmeland, H., Fan, C., Perou, C. M. & Nobel, A. B. Merging two gene-

expression studies via cross-platform normalization. *Bioinformatics* **24**, 1154–1160 (2008).

28. Gagnon-Bartsch, J. A. & Speed, T. P. Using control genes to correct for unwanted variation

in microarray data. *Biostatistics* **13**, 539–552 (2012).

29. Weishaupt, H. *et al.* Batch-normalization of cerebellar and medulloblastoma gene expression

datasets utilizing empirically defined negative control genes. *Bioinformatics* **35**, 3357–3364

(2019).

30. Wang, D. *et al.* Extensive up-regulation of gene expression in cancer: the normalised use of

microarray data. *Mol Biosyst* **8**, 818–827 (2012).

31. Nygaard, V., Rødland, E. A. & Hovig, E. Methods that remove batch effects while retaining

group differences may lead to exaggerated confidence in downstream analyses. *Biostatistics*

17, 29–39 (2016).

32. Parker, H. S. *et al.* Preserving biological heterogeneity with a permuted surrogate variable

analysis for genomics batch correction. *Bioinformatics* **30**, 2757–2763 (2014).

33. Zheng, X., Leung, K.-S., Wong, M.-H. & Cheng, L. Long non-coding RNA pairs to assist in

diagnosing sepsis. *BMC Genomics* **22**, 275 (2021).

34. Reddy, A. *et al.* Gene Expression Ratios Lead to Accurate and Translatable Predictors of

DR5 Agonism across Multiple Tumor Lineages. *PLoS ONE* **10**, e0138486 (2015).

35. Li, Q. *et al.* bvnGPS: a generalizable diagnostic model for acute bacterial and viral infection

using integrative host transcriptomics and pretrained neural networks. *Bioinformatics* **39**,

btad109 (2023).

36. Jin, N. *et al.* PAGE-based transfer learning from single-cell to bulk sequencing enhances

model generalization for sepsis diagnosis. *Briefings in Bioinformatics* **26**, bbae661 (2024).

37. Zhao, P. *et al.* Intra-sample reversed pairs based on differentially ranked genes reveal

biosignature for ovarian cancer. *Computers in Biology and Medicine* **172**, 108208 (2024).

38. Wang, R. *et al.* Improving bulk RNA-seq classification by transferring gene signature from

single cells in acute myeloid leukemia. *Briefings in Bioinformatics* **23**, bbac002 (2022).

39. Wu, Q. *et al.* meGPS: a multi-omics signature for hepatocellular carcinoma detection

integrating methylome and transcriptome data. *Bioinformatics* **38**, 3513–3522 (2022).

40. Li, H. *et al.* Whole transcriptome analysis reveals non-coding RNA's competing endogenous

gene pairs as novel form of motifs in serous ovarian cancer. *Comput Biol Med* **148**, 105881

(2022).

41. Johnson, W. E., Li, C. & Rabinovic, A. Adjusting batch effects in microarray expression data

using empirical Bayes methods. *Biostatistics* **8**, 118–127 (2007).

42. Tibshirani, R. Regression Shrinkage and Selection Via the Lasso. *Journal of the Royal*

Statistical Society Series B: Statistical Methodology **58**, 267–288 (1996).

43. Salsano, E. *et al.* Expression of the neurogenic basic helix-loop-helix transcription factor

NEUROG1 identifies a subgroup of medulloblastomas not expressing ATOH1. *Neuro Oncol*

9, 298–307 (2007).

44. Northcott, P. A. *et al.* Medulloblastoma Comprises Four Distinct Molecular Variants. *JCO*

29, 1408–1414 (2011).

45. Heinäniemi, M. *et al.* Gene-pair expression signatures reveal lineage control. *Nat Methods*

10, 577–583 (2013).

46. Wang, H. *et al.* Individual-level analysis of differential expression of genes and pathways for

personalized medicine. *Bioinformatics* **31**, 62–68 (2015).

47. Li, N. *et al.* Pairwise analysis of gene expression for oral squamous cell carcinoma via a

large-scale transcriptome integration. *J Cellular Molecular Medi* **28**, e70153 (2024).

48. Landry, A. P., Samuel, N., Spears, J. & Zador, Z. Integrated computational analyses reveal

novel insights into the stromal microenvironment of SHH-subtype medulloblastoma. *Sci Rep*

11, 20694 (2021).

49. Weidinger, G., Thorpe, C. J., Wuennenberg-Stapleton, K., Ngai, J. & Moon, R. T. The Sp1-

related transcription factors sp5 and sp5-like act downstream of Wnt/beta-catenin signaling

in mesoderm and neuroectoderm patterning. *Curr Biol* **15**, 489–500 (2005).

50. Fujimura, N. *et al.* Wnt-mediated down-regulation of Sp1 target genes by a transcriptional

repressor Sp5. *J Biol Chem* **282**, 1225–1237 (2007).

51. Kool, M. *et al.* Integrated Genomics Identifies Five Medulloblastoma Subtypes with Distinct

Genetic Profiles, Pathway Signatures and Clinicopathological Features. *PLoS ONE* **3**, e3088

(2008).

52. Kaur, K. *et al.* Integrating Molecular Subclassification of Medulloblastomas into Routine

Clinical Practice: A Simplified Approach. *Brain Pathology* **26**, 334–343 (2016).

53. Barrett, T. *et al.* NCBI GEO: archive for functional genomics data sets—update. *Nucleic

Acids Research* **41**, D991–D995 (2012).

54. Kukreja, S. L., Löfberg, J. & Brenner, M. J. A LEAST ABSOLUTE SHRINKAGE AND

SELECTION OPERATOR (LASSO) FOR NONLINEAR SYSTEM IDENTIFICATION.

IFAC Proceedings Volumes **39**, 814–819 (2006).

55. Friedman, J. D., Reece, G. R. & Eldor, L. The Utility of the Posterior Thigh Flap for

Complex Pelvic and Perineal Reconstruction: *Plastic and Reconstructive Surgery* **126**, 146–

155 (2010).

56. Raivo Kolde. pheatmap: Pretty Heatmaps.

57. Gendoo, D. M. A., Smirnov, P., Lupien, M. & Haibe-Kains, B. Personalized diagnosis of

medulloblastoma subtypes across patients and model systems. *Genomics* **106**, 96–106

(2015).

58. Cover, T. & Hart, P. Nearest neighbor pattern classification. *IEEE Trans. Inform. Theory* **13**,

21–27 (1967).

59. Cortes, C. & Vapnik, V. Support-vector networks. *Machine learning* **20**, 273–297 (1995).

60. Tolles, J. & Meurer, W. J. Logistic regression: relating patient characteristics to outcomes.

Jama **316**, 533–534 (2016).

61. Tin Kam Ho. The random subspace method for constructing decision forests. *IEEE Trans.*

Pattern Anal. Machine Intell. **20**, 832–844 (1998).

62. Chen, T. & Guestrin, C. Xgboost: A scalable tree boosting system. in *Proceedings of the*

22nd acm sigkdd international conference on knowledge discovery and data mining 785–794

(2016).

63. Hand, D. J. & Yu, K. Idiot's Bayes: Not So Stupid after All? *International Statistical Review*

/ Revue Internationale de Statistique **69**, 385 (2001).

665