Contact networks of small mammals highlight potential transmission foci of *Lassa mammarenavirus*.

Authors

David Simons [1,2,3], Ravi Goyal [4], Umaru Bangura [5,6], Rory Gibb [2,7] Ben Rushton [8],

Dianah Sondufu [6], Joyce Lamin [6], James Koninga [9], Momoh Jimmy [9], Mike Dawson [6],

Joseph Lahai [6], Rashid Ansumana [6], Elisabeth Fichet-Calvet [5], Richard Kock [1],

Deborah Watson-Jones [3,10], Kate E. Jones [2,7]

[1] Centre for Emerging, Endemic and Exotic Diseases, The Royal Veterinary College, London, United Kingdom

[2] Centre for Biodiversity and Environment Research, Department of Genetics, Evolution and Environment, University College London, London, United Kingdom

[3] Department of Clinical Research, London School of Hygiene and Tropical Medicine, London, United Kingdom

[4] Department of Medicine, University of California, San Diego, USA

[6] Implementation Research, Zoonoses control, Bernard-Nocht Institute for Tropical

Medicine, Hamburg, Germany

[7] Njala University, Bo, Sierra Leone

[8] People & Nature Lab, UCL East, Department of Genetics, Evolution and Environment,

University College London, London, United Kingdom

[9] Panadea Diagnostics GmbH, Hamburg, Germany

[10] Kenema Government Hospital, Kenema, Sierra Leone

[11] Mwanza Intervention Trials Unit, National Institute for Medical Research, Mwanza,

Tanzania

Corresponding author: David Simons, (dzs6259@psu.edu), The Royal Veterinary

College, Royal College Street, London, United Kingdom, NW1 0TU, +46793428490

Word count (abstract): 250

Word count (text): 6831

Figures: 5

Tables: 1

Supplementary Information: 2 pdf files

Supplementary Figures: 20 in 1 document

Key words: Rodent associated zoonoses; Mastomys natalensis; Disease ecology;

Transmission networks; Land use; Sierra Leone

Abstract

Lassa fever, caused by *Lassa mammarenavirus* (LASV), is an endemic zoonosis in several West African countries. Human infections primarily result from transmission from rodent hosts, with the primary reservoir species being *Mastomys natalensis*, a synanthropic rodent. In Sierra Leone, small-mammal communities are structured along land use gradients, and this structure is hypothesized to influence the risk of Lassa fever outbreaks in human populations. However, whether anthropogenic habitats facilitate significant LASV transmission among small mammals is not well understood.

To address this, we conducted rodent sampling over 43,266 trap nights, detecting 684 rodents and shrews in the Lassa fever-endemic Eastern Province of Sierra Leone. We reconstructed small-mammal contact networks to investigate whether contact rates and network structures differed by land use to identify settings more conducive to viral transmission among hosts. We found that small-mammal communities were larger in villages and agricultural settings than in forests, but contact rates were similar across habitats. Network structures differed by land use, with villages exhibiting more disconnected networks compared to agricultural settings. Notably, intra-specific contact among *M. natalensis* was more likely in agricultural environments than in villages, suggesting that land use can influence LASV transmission dynamics.

Overall, LASV seroprevalence was 5.7% across the small-mammal communities, with LASV antibodies detected in nine rodent and shrew species. These findings underscore the importance of expanding rodent surveillance across various habitats to understand habitat-specific pathogen transmission dynamics. A more systematic

58 approach to LASV surveillance in villages and agricultural settings will help identify key

59 host species driving pathogen dynamics.

60 Introduction

Lassa fever caused by *Lassa mammarenavirus* (LASV) is a rodent-associated

zoonoses endemic to West Africa, with an estimated 100,000-900,000 infections

annually[1,2]. Unlike the outbreaks in Nigeria, cases in other endemic countries - Guinea,

Liberia and Sierra Leone - are sporadically documented[3–5]. In Sierra Leone disease

outbreaks frequently go undetected, consistent with findings of up to 80% LASV

seropositivity among human communities in regions previously not considered

endemic[6]. Human infections typically result from transmission from rodent hosts, with

limited subsequent human-to-human transmission[7]. Understanding LASV transmission

in endemic settings requires a detailed characterization of small-mammal community

interactions, through which pathogen transmission occurs and is maintained.

The primary reservoir of LASV, *Mastomys natalensis* is a native synanthropic

rodent species, widespread across sub-Saharan Africa. Pathogen challenge studies on

M. natalensis colonies, suggest that acute infection does not significantly alter rodent

behaviour or cause clinical pathology[8–10]. LASV is transmitted through both direct

contact (e.g., superficial wounds caused by infected conspecifics) and indirect contact

(e.g., exposure to contaminated environments), at low infectious doses[9]. Vertical

transmission (mother-to-pup) may also contribute to transmission dynamics[10,11].

Infected adult rodents exhibit detectable viral RNA as early as 3 days post-infection,

with viral loads peaking within 1-2 weeks and resolving by 40 days[9]. Among individuals

infected within the first 2 weeks of life viral RNA is detectable up to 16-months post

infection[10]. The transient nature of acute infection - outside of neonatal infections - has

led many studies to focus on LASV-specific antibody detection, rather than circulating

virus[12–14].

84   The antibody response dynamics to LASV in rodents are not yet well understood.

Based on data from a similar arenavirus, Morogoro virus, seroconversion is expected to

occur by 7 days post-infection, with antibodies (e.g., IgG) persisting beyond the decline

of detectable RNA[11]. LASV-infected rodents are presumed to develop lifelong immunity

to reinfection upon seroconversion, however, the efficacy of neutralizing antibodies is

unclear and the role of immune or partially immune individuals in transmission networks

is not known[9,10,15]. Although antibody-based studies have limitations, the higher

prevalence of LASV seropositivity compared to acute infections provides valuable

insights into viral dynamics within endemic regions. For example, a recent study in Bo

district, Sierra Leone, reported a 2.8% LASV IgG seroprevalence among rodents and

shrews, compared to a 0.3% prevalence of acute infection detected via PCR,

underscoring the challenges of identifying acute infections in small-mammal

populations[16].

97   While *M. natalensis* is considered the primary host species of LASV, 10 further

rodent species have been identified as acutely or previously infected with LASV in

endemic regions[14,17–19]. The contribution of these additional species to pathogen

transmission into human populations - and their role in viral transmission or

maintenance within their species communities - remains unclear. In species-rich

environments, both direct and indirect contact among rodents may result in incidental

infections of non-reservoir species, which are subsequently detected through

surveillance. Incidental infections of non-primary host species may have little impact on

viral transmission or maintenance[20]. Alternatively, these species could facilitate the

transfer of LASV across landscapes, linking geographically isolated M. natalensis

populations and causing reintroduction of the virus into reservoir species

populations[21,22]. Increasing recognition of multi-species host systems in zoonoses

underscores the importance of expanding surveillance efforts to the wider community in

which the host resides to better understand pathogen prevalence and dynamics[23,24].

Host network structure is a key determinant of pathogen dynamics. Pathogens

are more likely to persist in dense, well-connected networks where frequency

dependent transmission dominates[25]. In contrast, pathogens with limited environmental

transmission are likely to become isolated in fragmented or discontinuous networks[26–28].

Networks that contain nodes (i.e., individual humans or rodents) with high node

betweenness — nodes which stand between other nodes, acting as crucial focal points

between other nodes — can significantly influence pathogen transmission and

maintenance, especially in discontinuous networks where they may be the only linkage

between network subcomponents (i.e., super-spreaders)[29,30]. While the removal of

these high-betweenness nodes can fragment the network and interrupt pathogen

transmission, the presence of infectious high-betweenness nodes can enhance

pathogen spread through the entire network due to their connections with a greater

number of individuals than average[31]. Moreover, the rate of transmission following a

contact event is influenced by host competence and the type of contact (e.g., inter-

specific or intra-specific)[32,33]. An understanding of the composition of rodent contact

networks in LASV-endemic regions has not been systematically reported and could

offer valuable insights into potential transmission networks, even in the absence of

direct observation of transmission events. Previous studies have provided general

descriptions of rodent populations but these cannot be readily translated into detailed

contact networks[16,34,35]. Ethical concerns regarding the release of potentially infectious

individuals into human communities limit the feasibility of capture-mark-release

experiments, which are otherwise effective for directly observing contact networks

among tagged individuals.

Small-mammal communities in LASV-endemic regions are structured along

anthropogenic land use gradients[14,36]. As such, the risk of Lassa fever outbreaks in

human populations is expected to correlate with these gradients[6,37,38]. Within human-

dominated land use types, the prevalence of typically synanthropic rodent hosts of

LASV is anticipated to be higher due to increased food availability, shelter, and reduced

predation pressure[39–41]. These factors influence rodent abundance and population

dynamics which in turn may promote greater pathogen persistence, as observed in

several other rodent-associated zoonosis systems[42–44]. Understanding how rodent

contact networks, like rodent occurrence and abundance, vary along these

anthropogenic gradients could reveal potentially distinct pathogen transmission

networks in different habitats. We hypothesize that rodent contact rates — and the

associated pathogen transmission networks — are greater in human-dominated

habitats where resources are concentrated.

The rich geolocation and temporal data obtained from systematic rodent trapping

enables the estimation of direct or indirect rodent contacts by inferring shared space

utilisation over short time periods[29,45]. Contact networks derived from wildlife data have

been successfully employed to study pathogen transmission. These networks are

particularly valuable for investigating the role of community structure and the impact of

contact rate heterogeneity between species in multi-host pathogen systems[46–48].

In this study, we leverage rodent and shrew trapping data collected over three

years in the Lassa fever-endemic region of Eastern Sierra Leone to reconstruct the

contact networks of small-mammal communities. We characterize potential interactions

— both direct and indirect — within these communities as a network, where the nodes

represent rodents or shrews, and the connections (or edges) between them represent

potential interactions. We hypothesize that the spatial clustering of conspecifics and the

increased abundance of commensal species in anthropogenically dominated

environments will lead to higher intra-specific contact rates compared to inter-specific

contact rates within these communities. We use contact rates within and between

species to explore their variation along an anthropogenic land use gradient with a

particular focus on *M. natalensis*. Finally, we report the prevalence of antibodies against

LASV among individual small mammals in the study region and investigate the

association between contact rates with seropositivity.

Methods

Study area

Rodent trapping surveys were conducted between October 2020-April 2023

within and around four village study sites (Baiama, Lalehun, Lambayama, and Seilama)

in the Lassa fever endemic zone of the Eastern Province of Sierra Leone (Figure 1A).

Surveys were conducted within trapping grids along a land use gradient of

anthropogenic disturbance comprising, forest, agriculture (including fallow and currently

in-use areas), and villages (within and outside of permanent structures) (see

Supplementary Information 1 for study site and trap grid locations). Trapping survey

sessions occurred four times annually with two trapping surveys in each of the rainy and

dry seasons (May to November and December to April, respectively), producing a total

of 10 trapping sessions over the study period.

A)

B)

Species

- Mastomys natalensis
- Crocidura olivieri
- Praomys rostratus
- Mus musculus
- Rattus rattus
- Lophuromys sikapusi
- Mus setulosus
- Crocidura buettikoferi
- Crocidura grandiceps
- Malacomys edwardsi
- Lemniscomys striatus
- Hylomyscus simus
- Hybomys planifrons
- Crocidura theresae
- Mastomys erythroleucus
- Gerbilliscus guineae
- Dasymys rufulus
- Other

Figure 1: A) Location of village study sites (circles) in the Eastern Province of Sierra

Leone. Kenema, the largest city in the province, and the national capital, Freetown, are

also shown (+). The inset map highlights the location of Sierra Leone within West Africa.

B) An example of a rodent contact network derived from trapping data during visit 5 in

village land use. Each coloured node represents an individual small mammal, with lines

(edges) indicating inferred contacts between individuals. The number of edges

connected to a node represents its degree. Betweenness reflects the importance of a

node in connecting different parts of the network. Pale nodes indicate unobserved

individuals, for whom contacts (edges) were not recorded. Species listed in the legend

without colours were never detected in village land use types.

Study sites were selected to represent the range of land use in the Eastern

Province of Sierra Leone, considering both accessibility throughout the year and

acceptability of the study protocol to the village communities (Supplementary

Information 1). At each trapping grid 49 Sherman traps (7.62cm x 8.89cm x 22.86cm)

(H.B. Sherman Traps, Tallahassee, USA), were arranged in a 7 x 7 grid, with traps

placed 7 metres apart in a grid conforming to the local landscape (median trapping grid

area = 3,882 m^2). In permanent structures, trap placement deviated from the grid

structure. At each visit, permanent structures were randomly selected from a grid

projected over the village area, with four traps placed within each structure. The location

of each trap within the grid was geolocated. Traps were baited with a locally produced

mixture of oats, palm oil and dried fish. Each morning, traps were checked, closed for

the day, and re-baited in the evening. Each trapping survey session consisted of four

consecutive trap-nights (TN) at each trapping grid within the village study site. Rodents

and shrews were associated with the coordinates of the trap they were detected.

Geospatial processing was performed using the sf package R (version 4.1.2)[49,50].

All small mammals were handled by trained researchers using appropriate

personal protective equipment. Animals were sedated using halothane and euthanized

according to established protocols[51]. Morphological measurements and samples of

blood and tissue (skin, liver and spleen) were collected. The study was approved by the

Clinical Research Ethical Review Board and Animal Welfare Ethical Review Board of

the Royal Veterinary College, UK (URN: 2019 1949-3), and the ethics committee of

Njala University, Sierra Leone. The study adhered to national and institutional ethical

guidelines for the humane treatment of animals and safe handling of zoonotic

pathogens. Trapping and sampling protocols were designed to minimize stress to the

animals and reduce the risk of pathogen exposure to researchers and communities.

Community engagement sessions were conducted to ensure understanding and

acceptance of the study objectives, and all work complied with the principles outlined in

the ARRIVE guidelines (v2)[52]. All carcasses were incinerated to mitigate pathogen

transmission risks.

Species identification

Species identification was performed in the field based on external morphological

characteristics, including body length, tail length, ear length, and pelage colouration,

following the taxonomic keys of Happold and Kingdon and Monadjem *et al.*

(Supplementary Information 2)[53,54]. Field identification was supplemented by molecular

methods to confirm species identity for individuals identified as *Mastomys sp.*, *Mus sp.*,

Rattus sp. and *Crocidura sp.* alongside a random subset of remaining individuals (50%

of remaining samples).

All samples remained in Sierra Leone and were stored at -20°C until processing.

Genomic DNA was extracted using QIAGEN DNAeasy kits as per the manufacturer's

instructions (Supplementary Information 1)[55]. DNA extracts were amplified using

platinum *Taq* polymerase (Invitrogen) and *cytochrome B* primers[16]. DNA amplification

was assessed through gel electrophoresis, and successful amplification products were

Sanger sequenced (performed by Eurofins Genomics). The sequences were attributed

to rodent species using the BLAST program, comparing the obtained sequences to

cytochrome B records in the NCBI database (accessed 2023-06-30)[56].

Lassa mammarenavirus seroprevalence

Serological status of trapped rodents and shrews was determined using the

BLACKBOX® LASV IgG ELISA Kit (Diagnostics Development Laboratory, Bernhard

Nocht Institute for Tropical Medicine), which has been validated for rodent samples[57,58].

The protocol is reproduced in Supplementary Information 1. Briefly, 1µL of whole blood

was inactivated by mixing with the provided sample dilution buffer (1:50). Where whole

blood was unavailable (21 samples, 3%), blood was extracted from dried blood spots

stored on filter paper by incubating with phosphate-buffered saline containing 0.08%

Sodium Azide and 0.05% Tween-20[59]. Samples, alongside negative and positive

controls, were incubated on the ELISA kit plates for 24 hours at 4-8 °C in a wet

chamber. Following incubation, the plates were washed and incubated for a further hour

with 1:10,000 diluted HRP-labelled streptavidin. A final wash was performed prior to the

addition of 100µL of 3,3',5,5'-Tetramethylbenzidine (TMB) substrate to wells, with

incubation for 10 minutes. The colorimetric reaction was stopped by adding 100µL of a

stop solution.

A deviation from the kit protocol occurred due to local ELISA plate reader

limitations. We measured the optical density (OD) at 450nm and 630nm, as opposed to

450nm and 620nm but this was not expected to have an important effect on absorbance

patterns, as advised by the manufacturer. The index value was calculated by

subtracting OD_{630} from OD_{450} and dividing by the cut-off value (the mean values of the

negative controls + 0.150). Samples were classified as positive if the index value was

greater than or equal to 1.1, negative if the index value was less than or equal to 0.9,

and inconclusive if the index value was between 0.9 and 1.1. Inconclusive results were

retested.

The prevalence of seropositive individuals is reported aggregated by species. A

Bayesian logistic regression model was constructed, using the `brms` package, to

estimate the Odds Ratio (OR) of seropositivity for each species compared to *M.*

natalensis, which served as the reference species[60]. Specifically, a Bernoulli regression

with normal, uninformative priors for population-level effects was used, with a binary

dependent variable for seropositivity and an independent variable of small mammal

species. Only species with more than 10 individuals assayed for antibodies to LASV

were included in this model. Posterior distributions are presented in graphical format,

alongside the posterior mean and 95% Credible Interval (CrI). Unlike frequentist

approaches, Bayesian inference does not rely on p-values; rather, statistical support for

an effect is assessed based on the posterior distribution, with associations interpreted in

terms of the central tendency (e.g., OR) and whether the CrI excludes the null value

(OR = 1).

In addition to the species-level seroprevalence analysis, we conducted post-hoc

exploratory analyses to assess differences in LASV seroprevalence by village and land

use type. These analyses employed Bayesian logistic regression models analogous to

those used in species-level comparisons, with results interpreted cautiously given their

non-pre-specified nature.

Small-mammal community

Species contact networks were reconstructed from the trapping data. Capture-

mark-recapture (CMR) methods have previously been used to identify space-sharing by

individuals[29,61,62]. In our study system, a CMR design was not feasible due to the risk of

releasing an infected animal back into a human community. Therefore, we considered

that rodents experience direct or indirect contact with other rodents through detections

at trapping locations that overlap in both time and space[45]. We assumed these potential

contacts were sufficient to transmit LASV if they were trapped within a buffer zone of 30

meter radius (2,828 m2) from the location of the trap during the same 4 trap night

session. The 30 meter buffer was chosen to encompass the typical home range of an

individual. A key assumption underlying this approach is that an individual was trapped

at the center of their home range[62]. This buffer was applied uniformly across species,

assuming that all species shared the same home range size.

We evaluated the appropriateness of selecting 30 meters as the buffer radius for

the primary analysis using the HomeRange R package (version 1.0.2)[63]. Four rodent

species from our study system had available home range data: *M. natalensis* (our

primary species of interest), *Lemnisomys striatus, Mus musculus* and *Rattus rattus.* The

30 meter buffer is expected to encompass the entire home range of *M. natelensis*

(mean home range = 419m^2) and more than 50% of the area of the home range of the

remaining species (*L. striatus* = 83%, *M. musculus* = 92%, *R. rattus* = 52%)

(Supplementary Figure 1). We further assessed the impact of this assumption on

produced contact networks by performing pre-specified sensitivity analyses with buffer

radii of 15 and 50 meters.

Networks were constructed from observed animals (nodes) and the presence or

absence of contact between them (edges). Data were aggregated by land use type and

sampling visit, producing a potential 32 distinct networks from 201 trapping grid, village

and visit combinations (see Figure 1B for an example network). However, as no rodents

were detected in three networks derived from forest sites, only 29 networks were used

in subsequent analysis.

We first explored the properties of these networks, stratified by land use type,

reporting species richness (number of different species), the number of nodes (number

of individual animals), the number of edges (number of contacts between individual

animals), mean node degree (i.e., the mean number of connections to other nodes for

each animal), and mean betweenness centrality (i.e., the mean number of times a node

lies on the shortest path between other nodes). Descriptions of degree were reported at

the global (i.e., network-level) and node-level. We then describe the contact networks

stratified by small mammal species, reporting the degree distribution of contacts by

species and investigating differences across the land use gradient. Finally, we explore

the species-level network characteristics by reporting the proportion of contacts each

species has with other species (i.e., the proportion of total inter- and intra-specific

contacts) stratified by land use.

Modelling the probablity of inter- and intra-specific contact rates in *Mastomys natalensis* across a land use gradient

To examine the association between land use and species with the probability of

contact between two individuals, we modelled these contacts using Exponential-Family

Random Graph Models (ERGMs)[64]. The analysis was limited to *M. natalensis*, the

primary rodent host of LASV. Estimation of ERGM parameters provide an OR for the

probability of an edge in a network – conditional on the rest of the network - based on

network properties included in the model and nodal attributes. Within our trapping grids,

only a subset of all individuals are detected in traps. Including unobserved individuals —

and thus, unobserved contacts — enhances the interpretability and generalizability of

the network models. This approach allows for a more accurate estimation of the total

population size by accounting for missing data, thereby making the network models

more representative of the entire population from which the analytic sample was

derived.

Incorporating unobserved individuals for modelling inter- and intra-specific *Mastomys natalensis* contacts

Previous analysis of our study system suggests that the probability of detecting a

rodent at each trap is less than 10% for 4 trap nights, provided that the species is

present in the trapping grid[36]. To estimate the abundance of individuals of each species

within a trapping grid, we modelled abundance (i.e., total population size) from repeated

count data using an N-mixture model implemented in the `unmarked` R package (version

1.2.5)[65,66]. The latent abundance distribution was modelled using Poisson, negative

binomial or zero-inflated Poisson random variables. The abundance model included the

number of trap nights and season as replicate-dependent detection covariates, as well

as location (rural vs. peri-urban setting) and land use type (forest, agriculture or village)

as occurrence covariates.

To select the most appropriate model for each species, we compared the Akaike

Information Criterion (AIC) of the Poisson, negative binomial, and zero-inflated Poisson

models. The best-fitting model was then used to derive the estimated abundance. The

median estimated abundance from the distribution produced for each trapping grid was

used to estimate the number of unobserved individuals in each network, aggregated by

land use type (Supplementary Figures 2.1-2.12). The number of observed individuals

was subtracted from the predicted abundance to derive the number of unobserved

individuals for each species. These unobserved individuals were explicitly set to have

missing (i.e., unobserved) edge values.

Finally, the constructed adjacency matrices were converted to networks using the

`network` R package (version 1.13.0.1) for subsequent ERGM modelling[67] (Figure 1B)

and Supplementary Figures 3.1-3.3).

Network models to estimate the probability of inter- and intra-specific contact rates

357   ERGMs were specified for each of our inferred contact networks to compare the

358   probabilities of edges forming based on rodent characteristics (i.e., species). The

359   general model is shown in Equation 1:

$$P(Y = y) \propto \exp\left(\theta_1 g_1(y) + \theta_2 g_2(y) + \ldots + \theta_p g_p(y)\right) \quad (1)$$

Where p is the number of terms in the model, and the values of the coefficients θ

represent the size and direction of the effects of the covariates $g(y)$ on the overall

probability of an edge being present in the network. At the edge level the expression for

the probability of the entire graph can be re-expressed as the conditional log-odds of a

single edge between two nodes (a contact between two rodents) as shown in Equation

2.

$$\text{logit}\left(P\left(Y_{ij} = 1 | y_{ij}^c\right)\right) = \theta' \delta\left(y_{ij}\right) \quad (2)$$

Where Y_{ij} is the random variable for the state of the node pair ij and $y_{i,j}^c$ signifies

all dyads in the network other than $y_{i,j}$. θ' is the coefficient for the change production of

an edge between the two nodes conditional on all other dyads remaining the same

$(\delta\left(y_{ij}\right))$.

ERGMs are implemented using the `ergm` package (version 4.3.2) in R[68]. Three terms

were included in the final ERGM to model the probability of the formation of ties

(Equation 3). The first term (edges), describes the density of the network, representing

the probability of a tie being observed in the network. The second term (species)

represents the conditional probability of a tie forming, conditional on the species of the

nodes. The third term (species homophily) accounts for intraspecific tie formation

among rodent individuals (i.e., the conditional probability of two individuals of the same

species forming a tie). To reduce linear dependency of the nodal terms and due to data

sparsity within our inferred networks all non-*M. natalensis* are grouped as "Other

species" through the `levels` term of the nodal covariates for the analysis of the effect of

land use on the probability of inter- or intra-specific contacts for *M. natalensis*.

$$P(Y = y) \propto \exp\left(\theta_{edges}g_{edges}(y) + \theta_{species}g_{species}(y) + \theta_{homophily}g_{homophily}(y)\right) \ (3)$$

ERGMs were implemented on the individual networks for each land use type at

each visit. The effect sizes from each model were pooled through random-effects meta-

analysis, stratified by land use, to produce a land use-specific summary effect size for

each coefficient[69]. Inclusion in meta-analysis was restricted to ERGMs that produced

stable estimates for each of the model terms (i.e., sufficient detections of *M. natalensis*

within the network). Random-effects models were conducted using the `metafor` package

(version 4.0.0) in R[70]. Heterogeneity across the models was assessed using the Q-test

and the restricted maximum-likelihood estimator (τ^2) with a prediction interval for the

true outcomes produced[69,71]. The Q-test assesses whether there is greater variability

among effect sizes than expected by chance, with a significant result indicating

substantial heterogeneity. The τ^2 statistic estimates the between-study variance,

quantifying the degree of heterogeneity rather than just testing for its presence. Weights

for each network included in meta-analysis were assigned using inverse-variance

weights[72].

The presence of influential networks was assessed using Cook's distance, for

models including influential networks leave-one-out sensitivity analysis were

performed[73]. Forest plots were generated to visualize the summary OR of the probability

of a tie for each model term, stratified by land use type.

Models with unstable estimates for the species homophily term were not included

in the random-effects meta-analysis. No contact networks from forest land use

contributed to meta-analysis as no *M. natalensis* were detected in these settings. Five

models from agricultural settings and eight from village settings were included in meta-

analysis.

Association of *Lassa mammarenavirus* seropositivity and position within a small-mammal community contact network

To investigate pathogen transmission within our networks, using seropositivity as

a proxy for prior exposure to LASV, we first report the small-mammal species found to

contain individuals that were seropositive for LASV. We then compared the nodal

degree of seropositive and seronegative individuals using a Wilcoxon rank-sum test with

continuity correction[74]. This analysis was repeated stratified by species to assess

whether contact rates were associated with an individual being seropositive. Finally, we

compared the node-level betweenness of seropositive and seronegative individuals to

determine whether an individual's position within a structured contact network was

associated with prior exposure to LASV.

Results

Overall 684 small mammals were trapped from 43,266 trap-nights. Seventeen species were identified, 13 of which were rodent species (76%) along with four species of insectivorous shrews (24%). *M. natalensis* was the most commonly detected species (N = 113, 16.5%), followed by *Crocidura olivieri* (N = 105, 15.3%) and *Praomys rostratus* (N = 102, 15%) (Table 1).

Prevalence of *Lassa mammarenavirus* antibodies within small-mammal communities

Antibodies to LASV were identified in 39 rodents and shrews (39/684, 5.7%) from 9 species, including *M. natalensis* (11/39, 28%), 5 *C. olivieri* (8/39, 21%), 8 *Lophuromys sikapusi* (8/39, 21%) and 4 *Rattus rattus* (4/39, 10%) (Table 1).

Compared to the primary rodent host of LASV (*M. natalensis*), the likelihood of LASV seropositivity was higher in *L. sikapusi* (OR = 2.09, 95% Credible Interval (CrI) = 0.86-4.78) but with wide credible intervals suggesting uncertainty in this result (Figure 2). There was weaker evidence for increased seropositivity in the sole positive shrew species *C. olivieri* (OR = 1.15, 95% CrI = 0.47-2.66). In contrast *M. musculus* (OR = -0.31, 95% CrI = 0.22-0.057) and *P. rostratus* (OR = 0.39, 95% CrI = 0.13-1.11) were less likely to be positive for antibodies against LASV than *M. natalensis*. Wide posterior distributions in this analysis reflect the overall low antibody prevalence, relatively small sample sizes, and uninformative priors used in the model.

435   Table 1 provides the number of individuals tested and seroprevalence of LASV

436 antibodies by species, while Figure 2 illustrates the odds ratios of seropositivty for

437 species with sufficient sample sizes compared to *M. natalensis.*

Table 1: The number of individuals detected and antibodies to Lassa mammarenavirus

among those individuals.

Species	Order	Individuals (N)	LASV Antibody detected (%)	Percentage of all positive individuals	OR (95% CrI)
Mastomys natalensis	Rodentia	113	11 (9.7%)	28.2%	ref
Crocidura olivieri	Eulipotyphla	105	8 (7.6%)	20.5%	1.15 (0.47-2.66)
Praomys rostratus	Rodentia	102	2 (2%)	5.1%	0.39 (0.13-1.11)
Mus musculus	Rodentia	90	0 (0%)	0%	0.22 (0.06-0.73)
Rattus rattus	Rodentia	88	4 (4.5%)	10.3%	0.73 (0.27-1.85)
Lophuromys sikapusi	Rodentia	57	8 (14%)	20.5%	2.09 (0.86-4.78)
Mus setulosus	Rodentia	43	3 (7%)	7.7%	1.01 (0.31-2.96)
Crocidura buettikoferi	Eulipotyphla	23	0 (0%)	0%	0.44 (0.09-1.78)
Crocidura grandiceps	Eulipotyphla	15	0 (0%)	0%	0.51 (0.1-2.33)
Malacomys edwardsi	Rodentia	11	1 (9.1%)	2.6%	1.08 (0.22-4.55)
Lemniscomys striatus	Rodentia	11	0 (0%)	0%	0.59 (0.11-2.68)
Hylomyscus simus	Rodentia	9	0 (0%)	0%	
Hybomys planifrons	Rodentia	7	1 (14.3%)	2.6%	
Mastomys erythroleucus	Rodentia	4	1 (25%)	2.6%	
Crocidura theresae	Eulipotyphla	3	0 (0%)	0%	
Gerbilliscus guineae	Rodentia	2	0 (0%)	0%	
Dasymys rufulus	Rodentia	1	0 (0%)	0%	

Figure 2: Odds Ratios of seropositivity to LASV among small-mammal species,

compared to Mastomys natalensis. Only species with more than 10 individuals assayed

for antibodies to LASV were included in this analysis.

Small mammals with antibodies to LASV were detected in three of the study

villages, Lalehun (N = 18, 46%), Seilama (N = 12, 31%) and Baiama (N = 9, 23%).

Lalehun had the highest percentage of antibody positive rodents (18/157, 12%),

followed by Baiama (9/121, 7%) and Seilama (12/263, 5%). No positive individuals were

detected in Lambayama, the most urbanised village study site, suggesting a potential

association between lower seroprevalence and urban settings. Antibody positive

individuals were detected during all study visits except visit 9 (2023-February).

Antibody positive small mammals were detected in all land use types. The

proportion of antibody positive individuals among all small mammals trapped were

similar across forest (2/44, 4.5%), agricultural (24/379, 6.3%) and village (13/261, 5%)

land use types.

In exploratory analyses conducted post hoc, we observed potential differences in

LASV seroprevalence by village and land use type. While these analyses were not pre-

specified and should be interpreted with caution, seroprevalence appeared to be higher

in the most rural village, Lalehun (OR = 2.56, 95% CrI = 1.27-5.27), than the more

urbanised village Lambayama (OR = 0.21, 95% CrI = 0.06-0.68). Baiama showed an

intermediate level of seroprevalence (OR = 1.57, 95% CrI = 0.67-3.52). Similarly,

agricultural land use (OR = 1.27, 95% CrI = 0.66-2.5) showed the highest proportion of

LASV antibody-positive small mammals, with lower seroprevalence observed in forest

settings (OR = 0.85, 95% CrI = 0.24-2.71). Differences for land use crossed an OR of 1

and so were not considered importantly different.

Small-mammal community contact networks

Contact networks produced from observations in agricultural land use contained

the highest small mammal species richness (12), followed by villages (9) and forests (6

species). The number of individuals (nodes) was also greatest in agricultural settings (n

= 379) compared to villages (n = 261) and forests (n = 44). Networks in village settings

exhibited the highest overall connectivity, as indicated by the highest mean degree

(mean = 6.2, standard deviation (SD) = 4.6), followed by forests (mean = 5.1, SD = 3.3)

and agricultural settings (mean = 4.9, SD = 5.4).

Additionally, the highest degree centrality values - indicating individuals with the

greatest number of direct connections - were observed in agricultural (degree = 24) and

village (degree = 20) settings. This suggests that while villages had the highest overall

connectivity, certain individuals in agricultural settings acted as highly connected hubs

within their networks.

Mean betweenness centrality followed an anthropogenic gradient, highest in

villages (mean = 3.06, SD = 10.2), decreasing in agricultural settings (mean = 0.46, SD

= 2.6) and lowest in forests (mean = 0.07, SD = 0.16). This suggests that animals in

human-modified environments played a greater role in connecting otherwise separate

parts of the network.

Degree centrality varied across species, with the highest values observed in

species more common in agricultural settings. *L. sikapusi*, *M. setulosus*, *P. rostratus*,

and *C. olivieri* — three native rodents and one shrew — had individuals with degree

centrality values up to 24, though most had lower values (Figure 3). In villages, *Mus*

musculus, a synanthropic invasive species, had a maximum degree of 20 and a high

median degree. In contrast, *M. natalensis*, despite being abundant in both agricultural

and village settings, had a lower maximum degree (12 in villages, 9 in agriculture) and a

similar median degree across both (5 and 4, respectively).

Figure 3: The degree (number of contacts) of individual small mammals stratified by

species and land use type. Boxes contain the median and inter-quartile range of the

degree distribution. Whiskers include the upper and lower quartile with outliers shown

as points.

497   No consistent trend in degree centrality was observed across all species when

498   stratified by land use type (Figure 3). For commensal species such as *M. natalensis*,

499   *Rattus rattus* and *M. musculus* the median degree centrality was elevated in villages.

500   However, for *M. natalensis* and *R. rattus*, no statistically significant differences were

501   detected in degree distributions when stratified by land use.

Describing inter- and intra-specific contact within small mammal communities

Species with more detected individuals generally exhibited a greater number of

inter-specific contacts ($r = 0.62$, $p = 0.007$, $n = 15$, Pearson correlation). For instance,

the frequently detected species, *M. natalensis*, *P. rostratus* and *R. rattus* each had

contacts with more than 12 other species. An exception to this trend was *M. musculus*,

which, despite being the fourth most frequently detected species, only had observed

contacts with four other species (Figure 4 and Supplementary Figures 4.1 and 4.2).

Intra-specific contacts were common for most species, but notable differences

emerged across land use types. For example, *M. natalensis* interacted with 12 other

species in agricultural land use settings, with 45% of all observed contacts involving

other individuals of the same species (Figure 3). However, in village settings, where *M.*

natalensis interacted with fewer species (9), the proportion of intra-specific contacts

decreased to 31% (Supplementary Figure 4.2). Not all species showed a dominance of

intra-specific contacts. For example, in agricultural areas, *L. sikapusi* interacted with 12

other species. However, a similar proportion of contacts for this species came from *P.*

rostratus (27%) as from other individuals of *L. sikapusi* (26%).

Figure 4: The proportion of contacts between individual small mammals in agricultural

land use. Darker colours indicate increasing proportions of observed contacts to a

species (Contact to) from named species (Contact from). Numbers in the cells

correspond to the proportion of contacts to a species from a named species. For

example, 45% of all contacts to Mastomys natalensis are from other M. natalensis while

9% of contacts are from Lophuromys sikapusi. Percentages sum to 100% in the Contact

to axis, while they may exceed 100% in Contact from. Species are ordered by the total

number detected in this study with M. natalensis (N = 113) in the bottom left.

The probability of inter- and intraspecific contact rates of

Mastomys natalensis across a land use gradient

Focusing on the reservoir species of LASV, *M. natalensis*, we analysed the

probability of contact using 13 ERGM models (8/10 village networks, 5/10 agricultural

networks), which were deemed suitable for random-effects meta-analysis. The odds of

a contact being observed for *M. natalensis* were low compared to all potential contacts

(observed and unobserved edges) and similar across both agricultural (Odds Ratio =

0.14, 95% Confidence Interval = 0.09-0.23, $p < 0.001$) and village land use (OR = 0.24,

95% C.I. = 0.17-0.36, $p < 0.001$) (Figure 5A). There was substantial heterogeneity in

contact odds between networks across different visits for both land use types ($\hat{\tau}^2_{\text{agriculture}}$

= 0.26, $Q = 112$, $p < 0.001$ and $\hat{\tau}^2_{\text{village}} = 0.23$, $Q = 54$, $p < 0.001$). Compared to other

rodent species, *M. natalensis* was observed to form fewer contacts overall.

When examining inter-specific contacts, *M. natalensis* showed a marginally

reduced, non-statistically significant odds of interacting with other species in agricultural

(OR = 0.49, 95% C.I. = 0.24-1.01, $p = 0.054$) and village settings (OR = 0.74, 95% C.I. =

0.55-1.01, $p = 0.055$) compared to inter-specific contacts among other rodent species

(Figure 5B). Notably, there was substantial heterogeneity in inter-specific contact odds

between networks ($\hat{\tau}^2_{\text{agriculture}} = 0.59$, $Q = 31$, $p < 0.001$ and $\hat{\tau}^2_{\text{village}} = 0.09$, $Q = 15$, $p = $

0.03). Land use type did not appear to significantly influence the probability of inter-

specific contact for *M. natalensis* relative to other species.

Conversely, *M. natalensis* showed a statistically significantly increase in the odds

of forming intra-specific contacts in agricultural settings (OR = 7.5, 95% C.I. = 3.42-16.5,

p < 0.001), but not in village settings (OR = 1.69, 95% C.I. = 0.85-3.36, *p* = 0.13) (Figure

5C). Heterogeneity in intra-specific contact odds was low in both land use types

($\hat{\tau}^2_{\text{agriculture}}$ = 0.22, Q = 5.6, *p* = 0.23 and $\hat{\tau}^2_{\text{village}}$ = 0.39, Q = 12, *p* = 0.1). These findings

suggest that *M. natalensis* was statistically significantly more likely to engage in intra-

specific interactions in agricultural settings compared to other small mammals but not in

villages.

Sensitivity analyses revealed no changes in the direction of effect sizes when

altering the contact radius (Supplementary Figures 5.1 and 5.2, for 15 and 50 meters

respectively). Additionally, leave-one-out sensitivity analyses for influential networks did

not indicate meaningful changes in the effect size magnitude or direction. These results

support the robustness of the findings concerning contact buffer assumptions and

community composition changes across visits.

Figure 5: Random effects meta-analysis of ERGM network models reporting the odds of

a contact being observed for M natalensis. A) The odds ratio of a contact being

observed for M. natalensis in Agricultural or Village land use types. B) The odds ratio of

a contact being observed between M. natalensis and an individual of a different rodent

species. C) The odds ratio of a contact being observed between M. natalensis and

another M. natalensis.

Association of *Lassa mammarenavirus* seropositivity and position within a small-mammal community contact network

Rodents and shrews that were LASV seropositive had a significantly lower mean degree centrality (mean = 3.7, SD = 2.9) compared to seronegative individuals (mean = 5.5, SD = 5.1) (W = 10018, p = 0.039). Species-specific analyses, performed only for species with more than five seropositive individuals (*M. natalensis*, *L. sikapusi* and *C. olivieri*) showed significantly lower mean degree centrality in seropositive compared to seronegative individuals for *M. natalensis* (W = 354, *p* = 0.048) and *L. sikapusi* (W = 99.5, *p* = 0.03), but not for *C. olivieri* (W = 429.5, *p* = 0.54). In contrast, betweenness centrality did not differ significantly between seropositive and seronegative individuals, either overall or within any of the three species examined individually. This suggests that while seropositive individuals had fewer direct contacts (lower degree centrality), their role in connecting otherwise unconnected individuals within the network (betweenness centrality) was not substantially different from that of seronegative individuals.

The lower mean degree centrality observed in seropositive individuals suggests that individuals with fewer connections were more likely to be seropositive. Given the low overall seroprevalence, this pattern may reflect limited recent pathogen transmission

587   within the population, potentially due to stochastic transmission events rather than

588   sustained intraspecific transmission.

Discussion

In the Eastern province of Sierra Leone, we found that small-mammal contact

networks were generally larger in village and agricultural settings, yet overall contact

rates were similar across land use types. Most individual animals had fewer than five

contacts, forming relatively sparse networks. While degree centrality did not differ

systematically by species, *M. natalensis* exhibited a high probability of intra-specific

contacts in agricultural settings, suggesting that these areas may serve as focal points

for LASV transmission, consistent with findings linking anthropogenic environments to

zoonotic disease risk[75,76]. LASV seropositivity was detected at low prevalence across

six rodent and shrew species, with *M. natalensis* comprising most seropositive

individuals. However, seropositive individuals had lower degree centrality overall,

though this pattern was not consistent within species.

We hypothesised that contact rates would be higher in anthropogenic habitats,

but global degree centrality was similar across land use types. However, the highest-

degree individuals were primarily found in village and agricultural settings, highlighting

spatial heterogeneity within networks. This finding underscores the limitations of relying

solely on aggregated network metrics in capturing individual-level variation, which is

critical for understanding transmission pathways[77,78].

Agricultural settings hosted the highest species richness, with a greater

proportion of interspecific contacts, reflecting edge effects that promote interactions

between synanthropic and sylvatic species[79,80]. Notably, native rodent species such as

P. rostratus in agricultural settings exhibited high degree centrality, suggesting that

agricultural habitats may amplify interspecific interactions, potentially increasing

zoonotic transmission risk[39]. These findings highlight the heterogeneity in contact rates

within species, often obscured in aggregated analyses[77].

Network structure also varies across land use types. Village networks exhibited

higher betweenness centrality, indicating greater fragmentation, with key individuals

bridging disconnected sub-components. These bottlenecks in network structure could

constrain LASV transmission by limiting sustained contact chains[81]. Conversely, the

lower betweenness centrality in agricultural networks suggests more cohesive

connectivity, potentially facilitating more efficient transmission.

Despite overall lower contacts, *M. natalensis* in agricultural settings exhibited

strong intra-specific clustering, whereas in villages, interspecific contacts were more

frequent. These findings align with prior studies showing that *M. natalensis* exhibits

weak territorial behaviour, similar to *R. rattus* but contrasting with *M. musculus*[10,82–84].

This intra-specific homophily in agricultural settings may promote LASV transmission

within *M. natalensis* populations, where competent hosts can sustain transmission

chains[85]. In contrast, higher interspecific contacts in villages could facilitate spillover but

may not sustain transmission within *M. natalensis*. Rodent movement between habitats

may further influence these dynamics[86].

The structure of agricultural networks could enable rapid LASV spread,

potentially leading to local pathogen extinction if transmission exceeds population

replenishment[81,87]. By contrast, fragmented village networks may allow for slower

transmission and long-term persistence[88]. These differences suggest interventions

should be tailored to land use contexts, particularly given agriculture's potential role in

amplifying transmission.

The seroprevalence of LASV (5.7%), was consistent with prior estimates from

Eastern Sierra Leone[16]. Our study included forest sites further from human habitation,

yet the proportion of M. natalensis individuals testing positive was similar (~9%).

However, *M. natalensis* represented a smaller fraction of total seropositive individuals

(28% vs. 75%), and we detected LASV antibodies in additional species (*C. olivieri*, *M.

setulosus*, *Hybomys planifrons*, *Mastomys erythroleucus*), suggesting broader host

involvement than previously documented[16]. These findings align with evidence from

other LASV-endemic regions, where multiple rodent and shrew species have been

implicated in LASV maintenance, necessitating multispecies approaches to surveillance

and control[12,16,18].

Several key assumptions must be considered. First, direct and indirect contacts

among rodents were inferred from co-location in space and time rather than directly

observed[45]. This approach assumes individuals who are detected at the centroid of their

home range move uniformly within it, which is unlikely to be accurate[62]. As a result,

contact rates may be under- or overestimated, particularly for species with variable

movement behaviours[29].Future studies using real-time tracking (e.g., radio tagging)

could refine these estimates, though ethical challenges remain[89,90]. Second, only a

subset of rodents and shrews active at a study site were captured[91,92]. We account

somewhat for the impact this will have on our network models by inferring the total

abundance of species within these sites[78,93]. If captured individuals behave differently

from those not detected, network inferences may be biased. For example, if trap

shyness influences space sharing, contact rates could be overestimated. Replicating

this study across additional sites would help assess these potential biases.

In conclusion, this study highlights the variability in contact rates and network

structures among rodent and shrew species across land-use types in LASV-endemic

settings. The complex contact networks involving multiple species may facilitate LASV

transmission in unexpected ways, particularly in agricultural settings where intra- and

inter-specific interactions are frequent. These findings underscore the importance of

tailoring control strategies to specific ecological contexts and host species behaviours to

mitigate Lassa fever risks effectively.

References

1.   McCormick JB, Webb PA, Krebs JW, Johnson KM, Smith ES., 1987. A

prospective study of the epidemiology and ecology of lassa fever. *J Infect Dis 155*: 437–

44

2.   Basinski AJ, Fichet-Calvet E, Sjodin AR, Varrelman TJ, Remien CH, Layman NC,

Bird BH, Wolking DJ, Monagin C, Ghersi BM, Barry PA, Jarvis MA, Gessler PE,

Nuismer SL., 2021. Bridging the gap: Using reservoir ecology and human serosurveys

to estimate lassa virus spillover in west africa. *PLoS Comput Biol 17*: e1008811

3.   Jetoh RW, Malik S, Shobayo B, Taweh F, Yeabah TO, George J, Gbelee B,

Teahton J, Jaryan F, Tegli M, Umeokonkwo CD, MaCauley J., 2022. Epidemiological

characteristics of lassa fever cases in liberia: A retrospective analysis of surveillance

data, 2019-2020. *International Journal of Infectious Diseases*

4.   Shaffer JG, Schieffelin JS, Momoh M, Goba A, Kanneh L, Alhasan F, Gbakie M,

Engel EJ, Bond NG, Hartnett JN, Nelson DKS, Bush DJ, Boisen ML, Heinrich ML,

Rowland MM, et al., 2021. Space-time trends in lassa fever in sierra leone by ELISA

serostatus, 2012–2019. *Microorganisms 9*: 586

5.   Bausch DG, Demby AH, Coulibaly M, Kanu J, Goba A, Bah A, Condé N, Wurtzel

HL, Cavallaro KF, Lloyd E, Baldet FB, Cissé SD, Fofona D, Savané IK, Tolno RT, et al.,

2001. Lassa fever in guinea: I. Epidemiology of human disease and clinical

observations. *Vector Borne Zoonotic Dis 1*: 269–281

6.   Grant DS, Engel EJ, Yerkes NR, Kanneh L, Koninga J, Gbakie MA, Alhasan F,

Kanneh FB, Kanneh IM, Kamara FK, Momoh M, Yillah MS, Foday M, Okoli A, Zeoli A,

et al., 2023. Seroprevalence of anti-lassa virus IgG antibodies in three districts of sierra

leone: A cross-sectional, population-based study. *PLOS Neglected Tropical Diseases

17*: e0010938

7.   Lo Iacono G, Cunningham AA, Fichet-Calvet E, Garry RF, Grant DS, Khan SH,

Leach M, Moses LM, Schieffelin JS, Shaffer JG, Webb CT, Wood JLN., 2015. Using

modelling to disentangle the relative contributions of zoonotic and anthroponotic

transmission: The case of lassa fever. *PLOS NEGLECTED TROPICAL DISEASES 9*

8.   Walker DH, Wulff H, Lange JV, Murphy FA., 1975. Comparative pathology of

lassa virus infection in monkeys, guinea-pigs, and mastomys natalensis. *Bull World*

Health Organ 52: 523–534

9.   Safronetz D, Rosenke K, Meade-White K, Sloan A, Maiga O, Bane S, Martellaro

C, Scott DP, Sogoba N, Feldmann H., 2022. Temporal analysis of lassa virus infection

and transmission in experimentally infected mastomys natalensis. *PNAS Nexus 1*:

pgac114

10.   Hoffmann C, Krasemann S, Wurr S, Hartmann K, Adam E, Bockholt S, Müller J,

Günther S, Oestereich L., 2024. Lassa virus persistence with high viral titers following

experimental infection in its natural reservoir host, Mastomys natalensis. *Nature*

Communications 15: 9319

11.   Borremans B, Vossen R, Becker-Ziaja B, Gryseels S, Hughes N, Van Gestel M,

Van Houtte N, Günther S, Leirs H., 2015. Shedding dynamics of morogoro virus, an

african arenavirus closely related to lassa virus, in its natural reservoir host mastomys

natalensis. *Sci Rep 5*: 10445

12.   Demby AH, Inapogui A, Kargbo K, Koninga J, Kourouma K, Kanu J, Coulibaly M,

Wagoner KD, Ksiazek TG, Peters CJ, Rollin PE, Bausch DG., 2001. Lassa fever in

guinea: II. Distribution and prevalence of lassa virus infection in small mammals. *Vector-*

Borne and Zoonotic Diseases 1: 283–297

13.   Kerneis S, Koivogui L, Magassouba N, Koulemou K, Lewis R, Aplogan A, Grais

RF, Guerin PJ, Fichet-Calvet E., 2009. Prevalence and risk factors of lassa

seropositivity in inhabitants of the forest region of guinea: A cross-sectional study. *PLoS*

Neglected Tropical Diseases [electronic resource] 3: e548

14.   Fichet-Calvet E, Becker-Ziaja B, Koivogui L, Gunther S., 2014. Lassa serology in

natural populations of rodents and horizontal transmission. *Vector Borne & Zoonotic*

Diseases 14: 665–674

15.   Mariën J, Borremans B, Gryseels S, Soropogui B, De Bruyn L, Bongo GN,

Becker-Ziaja B, Bellocq JG de, Günther S, Magassouba N, Leirs H, Fichet-Calvet E.,

2017. No measurable adverse effects of lassa, morogoro and gairo arenaviruses on

their rodent reservoir host in natural conditions. *Parasites & Vectors 10*: 210

16.   Bangura U, Buanie J, Lamin J, Davis C, Bongo GN, Dawson M, Ansumana R,

Sondufu D, Thomson EC, Sahr F, Fichet-Calvet E., 2021. Lassa virus circulation in

small mammal populations in bo district, sierra leone. *BIOLOGY-BASEL 10*

17.   Monath TP, Newhouse VF, Kemp GE, Setzer HW, Cacciapuoti A., 1974. Lassa

virus isolation from mastomys natalensis rodents during an epidemic in sierra leone.

Science 185: 263–265

18.   Olayemi A, Cadar D, Magassouba N, Obadare A, Kourouma F, Oyeyiola A,

Fasogbon S, Igbokwe J, Rieger T, Bockholt S, Jérôme H, Schmidt-Chanasit J,

Garigliany M, Lorenzen S, Igbahenah F, et al., 2016. New Hosts of The Lassa Virus.

Scientific Reports 6: 25280

19.   Simons D, Attfield LA, Jones KE, Watson-Jones D, Kock R., 2023. Rodent

trapping studies as an overlooked information source for understanding endemic and

novel zoonotic spillover. *PLOS Neglected Tropical Diseases 17*: e0010772

20.   Gilbert AT, Fooks AR, Hayman DTS, Horton DL, Müller T, Plowright R, Peel AJ,

Bowen R, Wood JLN, Mills J, Cunningham AA, Rupprecht CE., 2013. Deciphering

serology to understand the ecology of infectious diseases in wildlife. *EcoHealth 10*:

298–313

21.   Caron A, Cappelle J, Cumming GS, Garine-Wichatitsky M de, Gaidet N., 2015.

Bridge hosts, a missing link for disease ecology in multi-host systems. *Veterinary*

Research 46: 83

22.   Cardenas NC, Sykes AL, Lopes FPN, Machado G., 2022. Multiple species

animal movements: Network properties, disease dynamics and the impact of targeted

control actions. *Veterinary Research 53*: 14

23.   Keesing F, Ostfeld RS., 2021. Impacts of biodiversity and biodiversity loss on

zoonotic diseases. *Proceedings of the National Academy of Sciences 118*:

e2023540118

24.   Albery GF, Becker DJ, Brierley L, Brook CE, Christofferson RC, Cohen LE,

Dallas TA, Eskew EA, Fagre A, Farrell MJ, Glennon E, Guth S, Joseph MB, Mollentze

N, Neely BA, et al., 2021. The science of the host–virus network. *Nat Microbiol 6*: 1483–

1492

25.   Begon M, Hazel SM, Baxby D, Bown K, Cavanagh R, Chantrey J, Jones T,

Bennett M., 1999. Transmission dynamics of a zoonotic pathogen within and between

wildlife host species. *Proceedings of the Royal Society of London Series B: Biological*

Sciences 266: 1939–1945

26.   Swinton J, Harwood J, Grenfell BT, Gilligan CA., 1998. Persistence thresholds

for phocine distemper virus infection in harbour seal phoca vitulina metapopulations.

Journal of Animal Ecology 67: 54–68

27.   Almberg ES, Cross PC, Dobson AP, Smith DW, Hudson PJ., 2012. Parasite

invasion following host reintroduction: A case study of yellowstone's wolves.

Philosophical Transactions of the Royal Society B: Biological Sciences 367: 2840–2851

28.   Kozakiewicz CP, Burridge CP, Lee JS, Kraberger SJ, Fountain-Jones NM, Fisher

RN, Lyren LM, Jennings MK, Riley SPD, Serieys LEK, Craft ME, Funk WC, Crooks KR,

VandeWoude S, Carver S., 2023. Habitat connectivity and host relatedness influence

virus spread across an urbanising landscape in a fragmentation-sensitive carnivore.

Virus Evolution 9: veac122

29.   Clay CA, Lehmer EM, Previtali A, St. Jeor S, Dearing MD., 2009. Contact

heterogeneity in deer mice: Implications for sin nombre virus transmission. *Proc Biol Sci*

276: 1305–1312

30.   VanderWaal KL, Ezenwa VO., 2016. Heterogeneity in pathogen transmission:

Mechanisms and methodology. *Functional Ecology 30*: 1606–1622

31.   Chen S, White BJ, Sanderson MW, Amrine DE, Ilany A, Lanzas C., 2014. Highly

dynamic animal contact network and implications on disease transmission. *Sci Rep 4*:

4472

32.   Faust CL, Dobson AP, Gottdenker N, Bloomfield LSP, McCallum HI, Gillespie

TR, Diuk-Wasser M, Plowright RK., 2017. Null expectations for disease dynamics in

shrinking habitat: Dilution or amplification? *Philosophical Transactions of the Royal*

Society B: Biological Sciences 372: 20160173

33.   Young HS, Parker IM, Gilbert GS, Sofia Guerra A, Nunn CL., 2017. Introduced

species, disease ecology, and biodiversity–disease relationships. *Trends in Ecology &*

Evolution 32: 41–54

34.   Fichet☐Calvet E, Audenaert L, Barrière P, Verheyen E., 2010. Diversity,

dynamics and reproduction in a community of small mammals in upper guinea, with

emphasis on pygmy mice ecology. *African Journal of Ecology 48*: 600–614

35.   Happi AN, Olumade TJ, Ogunsanya OA, Sijuwola AE, Ogunleye SC, Oguzie JU,

Nwofoke C, Ugwu CA, Okoro SJ, Otuh PI, Ngele LN, Ojo OO, Adelabu A, Adeleye RF,

Oyejide NE, et al., 2022. Increased prevalence of lassa fever virus-positive rodents and

diversity of infected species found during human lassa fever epidemics in nigeria.

Microbiol Spectr 10: e0036622

36.   Simons D, Gibb R, Bangura U, Sondufu D, Lamin J, Koninga J, Jimmy M,

Dawson M, Lahai J, Ansumana R, Fichet-Calvet E, Watson-Jones D, Kock R, Jones K.,

2025. Land use gradients drive spatial variation in Lassa fever host communities in the

Eastern Province of Sierra Leone.

37.   Klitting R, Kafetzopoulou LE, Thiery W, Dudas G, Gryseels S, Kotamarthi A,

Vrancken B, Gangavarapu K, Momoh M, Sandi JD, Goba A, Alhasan F, Grant DS,

Okogbenin S, Ogbaini-Emovo E, et al., 2022. Predicting the evolution of the lassa virus

endemic area and population at risk over the next decades. *Nature communications 13*:

5596

38.   Longet S, Leggio C, Bore JA, Key S, Tipton T, Hall Y, Koundouno FR, Bower H,

Bhattacharyya T, Magassouba N, Günther S, Henao-Restrapo A-M, Rossman JS,

Konde MK, Fornace K, et al., 2023. Influence of landscape patterns on exposure to

lassa fever virus, guinea. *Emerg Infect Dis 29*: 304–313

39.   Gibb R, Redding DW, Chin KQ, Donnelly CA, Blackburn TM, Newbold T, Jones

KE., 2020. Zoonotic host diversity increases in human-dominated ecosystems. *Nature

584*: 398–402

40.   Albery GF, Carlson CJ, Cohen LE, Eskew EA, Gibb R, Ryan SJ, Sweeny AR,

Becker DJ., 2022. Urban-adapted mammal species have more known pathogens.

Nature Ecology & Evolution 6: 794–801

41.   Ecke F, Han BA, Hörnfeldt B, Khalil H, Magnusson M, Singh NJ, Ostfeld RS.,

2022. Population fluctuations and synanthropy explain transmission risk in rodent-borne

zoonoses. *Nat Commun 13*: 7532

42.   Sauvage F, Langlais M, Yoccoz NG, Pontier D., 2003. Modelling hantavirus in

fluctuating populations of bank voles: The role of indirect transmission on virus

persistence. *Journal of Animal Ecology 72*: 1–13

43.   Laverty SM, Adler FR., 2009. The role of age structure in the persistence of a

chronic pathogen in a fluctuating population. *Journal of Biological Dynamics 3*: 224–234

44.   Salkeld DJ, Salathé M, Stapp P, Jones JH., 2010. Plague outbreaks in prairie

dog populations explained by percolation thresholds of alternate host abundance.

Proceedings of the National Academy of Sciences 107: 14247–14250

45.   Perkins SE, Cagnacci F, Stradiotto A, Arnoldi D, Hudson PJ., 2009. Comparison

of social networks derived from ecological data: Implications for inferring infectious

disease dynamics. *Journal of Animal Ecology 78*: 1015–1022

46.   Böhm M, Hutchings MR, White PCL., 2009. Contact networks in a wildlife-

livestock host community: Identifying high-risk individuals in the transmission of bovine

TB among badgers and cattle. *PLOS ONE 4*: e5016

47.   Drewe JA, Eames KTD, Madden JR, Pearce GP., 2011. Integrating contact

network structure into tuberculosis epidemiology in meerkats in south africa:

Implications for control. *Prev Vet Med 101*: 113–120

48.   White LA, Forester JD, Craft ME., 2017. Using contact networks to explore

mechanisms of parasite transmission in wildlife. *Biological Reviews 92*: 389–409

49.   Pebesma E., 2018. Simple features for r: Standardized support for spatial vector

data. *The R Journal 10*: 439–446

50.   R Core Team., 2021. R: A language and environment for statistical computing.

Vienna, Austria: R Foundation for Statistical Computing

51.   Fichet-Calvet E., 2014. Chapter 5 - lassa fever: A rodent-human interaction.

Johnson N, ed. *The role of animals in emerging viral diseases.* Boston: Academic

Press, 89–123

52.   Percie Du Sert N, Ahluwalia A, Alam S, Avey MT, Baker M, Browne WJ, Clark A,

Cuthill IC, Dirnagl U, Emerson M, Garner P, Holgate ST, Howells DW, Hurst V, Karp

NA, et al., 2020. Reporting animal research: Explanation and elaboration for the

ARRIVE guidelines 2.0. *PLOS Biology 18*: e3000411

53.   Happold DCD, Kingdon J eds., 2013. *Mammals of Africa. Vol. 3: Rodents, hares*

and rabbits. London: Bloomsbury

54.   Monadjem A, Taylor PJ, Denys C, Cotterill FPD., 2015. *Rodents of sub-saharan*

africa: A biogeographic and taxonomic synthesis. Berlin, München, Boston: DE

GRUYTER

55.   QIAGEN., 2023. DNeasy blood & tissue kits. Available at:

https://www.qiagen.com/us/products/discovery-and-translational-research/dna-rna-

purification/dna-purification/genomic-dna/dneasy-blood-and-tissue-kit. Accessed.

January 20, 2023

56.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ., 1990. Basic local

alignment search tool. *J Mol Biol 215*: 403–410

57.   Gabriel M, Adomeh DI, Ehimuan J, Oyakhilome J, Omomoh EO, Ighodalo Y,

Olokor T, Bonney K, Pahlmann M, Emmerich P, Lelke M, Brunotte L, Ölschläger S,

Thomé-Bolduan C, Becker-Ziaja B, et al., 2018. Development and evaluation of

antibody-capture immunoassays for detection of lassa virus nucleoprotein-specific

immunoglobulin m and g. *PLOS Neglected Tropical Diseases 12*: e0006361

58.   Soubrier H, Bangura U, Hoffmann C, Olayemi A, Adesina AS, Günther S,

Oestereich L, Fichet-Calvet E., 2022. Detection of lassa virus-reactive IgG antibodies in

wild rodents: Validation of a capture enzyme-linked immunological assay. *Viruses 14*:

59.   Grüner N, Stambouli O, Ross RS., 2015. Dried blood spots - preparing and

processing for use in immunoassays and in molecular techniques. *J Vis Exp*: 52619

60.   Bürkner P-C., 2017. brms: An R package for Bayesian multilevel models using

Stan. *Journal of Statistical Software 80*: 1–28

61.   Carslake D, Bennett M, Bown K, Hazel S, □fer S, Begon M., 2005. Space–time

clustering of cowpox virus infection in wild rodent populations. *Journal of Animal

Ecology 74*: 647–655

62.   Wanelik KM, Farine DR., 2022. A new method for characterising shared space

use networks using animal trapping data. *Behav Ecol Sociobiol 76*: 127

63.   Broekman MJE, Hoeks S, Freriks R, Langendoen MM, Runge KM, Savenco E,

Harmsel R ter, Huijbregts MAJ, Tucker MA., 2023. HomeRange: A global database of

mammalian home ranges. *Global Ecology and Biogeography 32*: 198–205

64.   Hunter DR, Handcock MS, Butts CT, Goodreau SM, Morris M., 2008. Ergm: A

package to fit, simulate and diagnose exponential-family models for networks. *Journal

of Statistical Software 24*: 1–29

65.   Royle JA., 2004. N-mixture models for estimating population size from spatially

replicated counts. *Biometrics 60*: 108–115

66.   Fiske I, Chandler R., 2011. Unmarked: An r package for fitting hierarchical

models of wildlife occurrence and abundance. *Journal of Statistical Software 43*: 1–23

67.   Butts CT., 2008. **Network** : A package for managing relational data in *r*. *J Stat

Soft 24*

68.   Handcock MS, Hunter DR, Butts CT, Goodreau SM, Krivitsky PN, Morris M.,

2022. Ergm: Fit, simulate and diagnose exponential-family models for networks. The

Statnet Project (https://statnet.org)

69.   Riley RD, Higgins JPT, Deeks JJ., 2011. Interpretation of random effects meta-

analyses. *BMJ 342*: d549

70.   Viechtbauer W., 2010. Conducting meta-analyses in r with the metafor package.

Journal of Statistical Software 36: 1–48

71.   Cochran WG., 1954. The combination of estimates from different experiments.

Biometrics 10: 101–129

72.   Borenstein M, Hedges LV, Higgins JPT, Rothstein HR., 2010. A basic

introduction to fixed-effect and random-effects models for meta-analysis. *Research

Synthesis Methods 1*: 97–111

73.   Cheung MW-L., 2019. A guide to conducting a meta-analysis with non-

independent effect sizes. *Neuropsychol Rev 29*: 387–396

74.   Bauer DF., 1972. Constructing confidence sets using rank statistics. *Journal of*

the American Statistical Association 67: 687–690

75.   Pei S, Yu P, Raghwani J, Wang Y, Liu Z, Li Y, Cheng Y, Lin Q, Song C,

Dharmarajan G, Faust CL, Tian Y, Xu Y, Liang Y, Qu J, et al., 2024. Anthropogenic land

consolidation intensifies zoonotic host diversity loss and disease transmission in human

habitats. *Nature Ecology & Evolution 9*: 99–110

76.   Carlson CJ, Brookson CB, Becker DJ, Cummings CA, Gibb R, Halliday FW,

Heckley AM, Huang ZYX, Lavelle T, Robertson H, Vicente-Santos A, Weets CM, Poisot

T., 2025. Pathogens and planetary change. *Nature Reviews Biodiversity 1*: 32–49

77.   Farine DR, Whitehead H., 2015. Constructing, conducting and interpreting animal

social network analysis. *Journal of Animal Ecology 84*: 1144–1163

78.   Silk MJ, Fisher DN., 2017. Understanding animal social structure: Exponential

random graph models in animal behaviour research. *Animal Behaviour 132*: 137–146

79.   Despommier D, Ellis BR, Wilcox BA., 2007. The Role of Ecotones in Emerging

Infectious Diseases. *EcoHealth 3*: 281–289

80.   Pruvot M, Chea S, Hul V, In S, Buor V, Ramassamy J-L, Fillieux C, Sek S, Sor R,

Ros S, Nuon S, San S, Ty Y, Chao M, Sours S, et al., 2024. Small mammals at the

edge of deforestation in Cambodia: Transient community dynamics and potential

pathways to pathogen emergence. *One Earth 7*: 123–135

81.   Keeling MJ, Eames KTD., 2005. Networks and epidemic models. *J R Soc*

Interface 2: 295–307

82.   Anderson PK., 1961. Density, social structure, and nonsocial environment in

house-mouse populations and the implications for regulation of numbers. *Trans N Y*

Acad Sci 23: 447–451

83.   Whisson DA, Quinn JH, Collins KC., 2007. Home range and movements of roof

rats (rattus rattus) in an old-growth riparian forest, california. *Journal of Mammalogy 88*:

589–594

84.   Borremans B, Hughes NK, Reijniers J, Sluydts V, Katakweba AAS, Mulungu LS,

Sabuni CA, Makundi RH, Leirs H., 2014. Happily together forever: Temporal variation in

spatial patterns and complete lack of territoriality in a promiscuous rodent. *Population*

Ecology 56: 109–118

85.   Luis AD, Kuenzi AJ, Mills JN., 2018. Species diversity concurrently dilutes and

amplifies transmission in a zoonotic host–pathogen system through competing

mechanisms. *Proceedings of the National Academy of Sciences 115*: 7979–7984

86.   Marien J, Kourouma F, Magassouba N, Leirs H, Fichet-Calvet E., 2018.

Movement Patterns of Small Rodents in Lassa Fever-Endemic Villages in Guinea.

Ecohealth 15: 348–359

87.   Messinger SM, Ostling A., 2009. The consequences of spatial structure for the

evolution of pathogen transmission rate and virulence. *The American Naturalist 174*:

441–454

88.   Peel AJ, Pulliam JRC, Luis AD, Plowright RK, O'Shea TJ, Hayman DTS, Wood

JLN, Webb CT, Restif O., 2014. The effect of seasonal birth pulses on pathogen

persistence in wild mammal populations. *Proceedings of the Royal Society B: Biological Sciences 281*: 20132962

89.   Mohr K, Leirs H, Katakweba A, Machang'u R., 2007. Monitoring rodents movements with a biomarker around introduction and feeding foci in an urban environment in tanzania. *afzo 42*: 294–298

90.   Borremans B, Reijniers J, Hughes NK, Godfrey SS, Gryseels S, Makundi RH, Leirs H., 2017. Nonlinear scaling of foraging contacts with rodent population density. *Oikos 126*: 792–800

91.   Parmenter RR, Yates TL, Anderson DR, Burnham KP, Dunnum JL, Franklin AB, Friggens MT, Lubow BC, Miller M, Olson GS, Parmenter CA, Pollard J, Rexstad E, Shenk TM, Stanley TR, et al., 2003. Small-mammal density estimation: A field comparison of grid-based vs. Web-based density estimators. *Ecological Monographs 73*: 1–26

92.   Moore JE, Swihart RK., 2005. Modeling patch occupancy by forest rodents: Incorporating detectability and spatial autocorrelation with hierarchically structured data. *The Journal of Wildlife Management 69*: 933–949

93.   Vega Yon GG, Slaughter A, Haye K de la., 2021. Exponential random graph models for little networks. *Social Networks 64*: 225–238