Endogenously generated Dutch-type Aβ nonfibrillar aggregates dysregulate presynaptic neurotransmission in the absence of detectable inflammation

Emilie L. Castranio[1], Merina Varghese[2], Elentina K. Argyrousi[3], Kuldeep Tripathi[4], Linda Söderberg[5], Erin Bresnahan[6,7,8], David Lerner[6,7], Francesca Garretti[6,7], Hong Zhang[3], Jonathan van de Loo[3], Cheryl D. Stimpson[9,10], Ronan Talty[6,7,11], Charles Glabe[12], Efrat Levy[13], Minghui Wang[14], Marjan Ilkov[14], Bin Zhang[14], Lars Lannfelt[3], Brigitte Guérin[15], William D. Lubell[16], Shai Rahimipour[4], Dara L. Dickstein[6,7,9,10], Sam Gandy[1,7,17,18]†, Ottavio Arancio[3]†, and Michelle E. Ehrlich[1,13,19,*]†

[1]Department of Neurology, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA.

[2]George & Anne Ryan Institute for Neuroscience, University of Rhode Island, RI, 02881, USA.

[3]Department of Pathology and Cell Biology, Department of Medicine, and The Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Vagelos College of Physicians and Surgeons, Columbia University, New York, NY, 10032, USA.

[4]Bar-Ilan University, Department of Chemistry, Ramat Gan, 5290002 Israel.

[5]BioArctic, Stockholm, SE-112 51 Sweden

[6]Nash Family Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA.

[7]Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA.

[8]Department of Hematology and Oncology, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA.

[9]Department of Pathology, Uniformed Services University of Health Sciences, Bethesda, MD, 20814, USA.

[10]Henry M. Jackson Foundation for the Advancement of Military Medicine Inc, Bethesda, MD, 20817, USA.

[11]Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA

[12]Department of Molecular Biology and Biochemistry, University of California, Irvine, CA, 92697, USA.

[13]Center for Dementia Research, The Nathan S. Kline Institute for Psychiatric Research, 140 Old Orangeburg Road, Orangeburg, NY, 10962, USA, and New York University Grossman School of Medicine, NYU Langone Medical Center, New York, NY, 10016, USA.

[14]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA.

[15]Department of Radiochemistry, University of Sherbrooke, Sherbrooke, Quebec, J1K 2R1, Canada

[16]Department of Chemistry, University of Montreal, Montreal, Quebec, H3C 3J7, Canada

[17]Department of Psychiatry and Alzheimer's Disease Research Center, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA.

[18]James J Peters VA Medical Center, Bronx, NY, 10468, USA.

[19]Department of Pediatrics, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA.

† Contributed equally to supervision

* Correspondence: michelle.ehrlich@mssm.edu

Abstract (166 words)

APP^{E693Q} transgenic mice develop aging-related learning deficits and accumulate endogenously generated nonfibrillar aggregates of Aβ (NFA-Aβ) and APP α-carboxy terminal fragments. The APP^{E693Q} mutation disrupts amyloid fibril formation, and no plaques develop in these mice. In the current study, the aging-related accumulation of NFA-Aβ in APP^{E693Q} mice was revealed by A11 immunohistochemistry and NFA-Ab-detecting cyclic D,L-α-peptide-FITC microscopy. The presynaptic termini of APP^{E693Q} mice developed aging-related physiological abnormalities in post-tetanic potentiation, synaptic fatigue, and synaptic vesicle replenishment. Single-cell RNA sequencing showed that excitatory neurons exhibited the most altered transcriptomic profile, especially involving "protein translation" and "oxidative phosphorylation". Direct measurements of electron transport chain catalysis revealed reduction in mitochondrial complex I activity in Dutch mice. Microglial transcript analysis revealed no evidence of inflammation. The incomplete clinical response to fibrillar Aβ immunodepletion in human patients may be attributable to residual NFA-Aβ that are undetectable by currently available biofluid or neuroimaging biomarkers. The depletion or neutralization of both fibrillar and NFA-Aβ may be needed for complete elimination of Aβ toxicity.

Teaser

APP[E693Q] "NFA-Aβ only" mice reveal clinically relevant mechanisms despite the absence of detectable inflammation

Introduction

Synaptic dysfunction is believed to be a key pathological hallmark of Alzheimer's disease (AD) that begins gradually and parallels clinical cognitive decline (*1-4*). Many studies have demonstrated that the burden of fibrillar amyloid-β (Aβ) proteinopathy does not correlate with synaptic loss (*5-7*) or with cognitive status(*8*). Instead, some investigators have proposed that brain levels of synaptotoxic nonfibrillar aggregates of Aβ (NFA-Aβ) including Ab oligomers and protofibrils correlate more closely with synaptic loss (*9, 10*). Synaptic NFA-Aβ has been detected in AD brains, and NFA-Aβ levels in human brain synaptosomes can differentiate subjects with early AD from cognitively intact controls with AD pathology (*11-13*). *In vitro*, synthetic NFA-Aβ peptides can cause inhibition of long-term potentiation (LTP) in hippocampal slices (*14, 15*), while *in vivo* injections of NFA-Aβ into the cerebral ventricles of healthy rats leads to cognitive deficits (*16*). Studies using a transgenic mouse model of AD have demonstrated that neuronal loss in the hippocampus correlates with NFA-Aβ levels but not with amyloid plaque burden (*17*). In 2010, we reported a transgenic mouse model overexpressing Dutch mutant human amyloid precursor protein (APP^{E693Q}) on the pan-neuronal Thy1 promoter (*18*). This mutation within the Aβ domain substitutes a glutamine for glutamic acid, thereby disrupting fibril assembly and favoring the formation of NFA-Aβ. The APP[E693Q] mutation causes hereditary cerebral hemorrhage with amyloidosis - Dutch type (HCHWAD) (*19*). The neuropathology of HCHWAD shows prominent cerebrovascular amyloidosis and hemorrhage as well

as diffuse parenchymal Aβ deposition (*20*). Although the APP^{E693Q} (or "Dutch") transgenic mouse failed to recapitulate the full neuropathology of either AD or HCHWAD, we were struck by the aging-related development of learning behavior impairment that was observed in the absence of detectable fibrillar Aβ. Analysis of the Dutch mouse brain revealed that this learning behavior impairment was directly related to the brain levels of NFA-Aβ (*18, 21*), an observation supported by imaging data in the current paper using epitomic histochemistry with anti-prefibrillar oligomer-specific antibody A11 (*97*) and NFA-Aβ-detecting cyclic peptide-FITC fluorescence microscopy (*22*). The Dutch mice also exhibit detectable cerebrovascular Aβ deposition and traces of perivascular iron (presumably due to the extravasation of red blood cells) without overt hemorrhage. The reported accumulation of APP α- and β-carboxyl-terminal fragments (APP-CTFs) in the Dutch mouse brain has raised the possibility that the learning impairment might be at least partially attributable to APP-CTF proteinopathy alone or in combination with the NFA-Aβ proteinopathy (*23*).

A detailed study of hippocampal CA1 neurons reported that significant decreases in dendritic length and complexities of apical, but not basal, dendrites were observed in Dutch compared to wildtype mice (*24*). The synaptic ultrastructure in the same brain region revealed no differences in overall synaptic density or morphology. A significant reduction in the postsynaptic density (PSD) length of synapses was identified on mushroom spines in Dutch mice compared to those of wildtype littermates. In the current study, electrophysiological and immuno-electron microscopy approaches were employed to investigate the excitatory synapses in the hippocampus of the Dutch mouse.

Bulk sequencing of RNA from the entorhinal cortex and dentate gyrus of Dutch mice revealed entorhinal cortex profiles that correlated with several human late-onset AD signatures (*25*). Microglial signatures were enriched in the differentially expressed genes (DEGs). Differential exon usage was also prominent in the Dutch mouse entorhinal cortex and the PSAPP ($APP^{KM670/671NL}$/$PSEN1^{\Delta exon9}$) mouse dentate gyrus. The transcriptomes from the entorhinal cortex and the dentate gyrus of Dutch mice were also enriched in markers of glutamatergic neurons, synaptic transmission, and protein translation. To study transcriptomics in the Dutch mouse hippocampus, single-cell RNA sequencing (scRNA-seq) was employed and revealed dysregulation of pathways for protein translation, oxidative phosphorylation, and the electron transport chain. Direct measurements of electron transport chain catalysis revealed changes in activity of mitochondrial complex I but not in that of complexes II, III, or IV.

In summary, a multi-disciplinary analysis of the Dutch mouse has now revealed alterations in synaptic physiology, gene expression, and protein translation in excitatory neurons, as well as changes in the transcriptomic profile and mitochondrial morphology and activity. Moreover, the high sensitivity of scRNA-seq has enabled neuroinflammation to be ruled out as a major contributor to pathogenesis in the hippocampi of Dutch mice. The present studies are among few focusing on endogenous generation of NFA-Aβ without the concurrent presence

of potential confounds from either Aβ fibrils or synthetic peptides. These findings support the notion that fibrillar Aβ is not necessary for the aging-related learning deficit observed in Dutch mice. These data may have important implications for the genesis of cognitive impairment in humans. The clinical response to anti-amyloid antibody therapy may be dampened due to the persistence of toxic NFA-Aβ conformers that go undetected in contemporary analysis of fluid biomarkers and in amyloid fibril positron emission tomography (PET) imaging. Depletion or neutralization of NFA-Aβ may be required for complete elimination of clinically relevant Aβ toxicity.

Results

Dutch mice show increased levels of nonfibrillar aggregate with aging.

Aging-dependent investigation of the Dutch mice revealed that 12 months is the first age at which deficits attributable to NFA-Aβ can be detected (*18*). To determine the involvement of NFA-Aβ in brain changes, multiple analyses were conducted at different ages, before and after the appearance of NFA-Aβ. Epitomic characterization was performed on cortical tissue from 2- and 24-month-old Dutch mice. Interestingly, this characterization revealed a significant increase with aging in the A11/6E10 ratio (ratio of prefibrillar oligomer content to total Aβ content; $F_{(3,16)} = 17.05$; $p < 0.0001$), but no such aging-related change in the OC/6E10 ratio (ratio of fibrillar oligomer content to total Aβ content; $F_{(3,16)} = 2.557$; $p = 0.0916$) in Dutch mice (**Figure 1, Supplementary Figure 1A**). Furthermore, brains were analyzed for the presence of protofibrils (PF) using the homogenization protocol described in Tucker *et al.* (*26*) and assayed for PF levels in the TBS soluble fraction using a mAb158-mAb158 sandwich MSD assay employing an in-house synthetic protofibril as calibrator. Both assays showed very low levels of Aβ PF. In the mAb158 sandwich assay, Aβ PF levels were around 100 pM, levels indistinguishable from those in homogenates of wildtype mice (**Supplementary Figure 1B**). No PF were detected in the cerebellum. Levels of PF in Dutch mice were approximately 10- to 100-fold lower than those measured in Tg-APP[ArcSwe] and Tg2576 (depending on age analyzed). In addition, $Aβ_{38/40/42}$ fragments were analyzed in the TBS insoluble pellet fraction after dissolution in formic acid and neutralization. Levels of Aβ PF were again very low, consistent with the absence of parenchymal plaques in Dutch mice.

The elevation of NFA-Aβ levels in the cortex, thalamus, and hippocampus of Dutch mice was visually and morphologically confirmed using a combination of immunohistochemistry using A11 antibody and fluorescence microscopy employing cyclic D,L-α-peptide-2 (CP-2)-FITC as a fluorescein-tagged NFA-Aβ-detecting ligand (*22*) (**Figure 1, Supplementary Figures 1C and 2-3**). The hypothesis that CP-2 and A11 bind to similar or identical domains on NFA-Aβ was confirmed by cross-competition blocking experiments in which pre-incubation of brain slices with CP-2 peptide prevented the subsequent binding of the prefibrillar oligomer-specific antibody A11 (**Supplementary Figure 4**). Furthermore, the interaction of CP-2 with NFA-Aβ has been characterized (*99-*

103): CP-2 was demonstrated (1) to stabilize Aβ dimers and trimers using a photoinduced cross-linking of unmodified proteins (PICUP) assay (*99-102*) and (2) to bind 10 nm particles typical of NFA-Aβ (*103*). In the current study, CP-2 exhibited similar K_d values for both fibrillar and nonfibrillar forms of Aβ in surface plasmon resonance (SPR) assays (**Supplementary Methods**): CP-2 interacted with Aβ fibrils and NFA-Aβ with K_d values of 6 and 4 μM, respectively.

Dutch mice showed no significant impairment in basal synaptic transmission and long-term synaptic plasticity in the hippocampus.

The relationship between synaptic input strength and the resulting electrical response, the so-called "input-output relation" and long-term potentiation (LTP) were tested in 7- to 7.5-month-old and 10.5- to 11-month-old Dutch mice. The two genotypes displayed similar input-output relationships and LTP in both age groups, suggesting that basal synaptic transmission at the CA1 region of the hippocampus and long-term synaptic plasticity at the Schaffer collateral were unimpaired in Dutch mice (**Figure 2**).

Dutch mice showed no change of hippocampal mGluR-2/3 expression or of mGluR-2/3+ synapse density.

Comparable levels of mGluR-2/3 protein were observed in Western blots of the insoluble membrane fraction from the hippocampus of 12-month-old wildtype and Dutch mice (**Supplementary Figure 5A**). Since Western blots may not detect regional differences, array tomography and immunofluorescence were used to analyze mGluR-2/3 puncta after staining serial sections of CA1 neurons with Neurotrace (**Supplementary Figure 5B**). Comparable densities in the CA1 of wildtype and Dutch mice were observed in 3D reconstructions of mGluR-2/3 synaptic puncta by array tomography (**Supplementary Figure 5C**, $F_{(4,3)} = 1.476$, $p = 0.7802$). Quantification of the total mGluR-2/3 puncta output yielded an average tissue density (puncta/μm^3) of 0.275 ± 0.087 for wildtype mice and 0.322 ± 0.112 for Dutch mice. Both genotypes exhibited comparable puncta size ranges, and no significant difference in average mGluR-2/3 puncta size (**Supplementary Figure 5D**; $F_{(1,3)} = 0.751$, $p = 0.538$) per percentile per sample. Similarly, neither the density nor size of GluN1 puncta were altered in Dutch mice (**Supplementary Figure 5E-F**; $F_{(4,3)} = 1.431$, $p = 0.8003$ for density and $F_{(1,3)} = 0.116$, $p = 0.950$ for puncta size).

Dutch mice had larger PSD area on small mGluR-2/3+ synapses in the CA1 region.

Changes in mGluR-2/3+ excitatory neurons were examined at the ultrastructural level in 12-month-old mice. Comparable percentages of mGluR-2/3+ synapses among total, non-perforated, and perforated synapses in the CA1 were observed in Dutch and wildtype mice using immunogold EM against mGluR-2/3 (**Figure 3A-C**). Moreover, Dutch and wildtype mice exhibited no difference in the total and perforated synapses (**Figure 3D, F**).

Higher postsynaptic density (PSD) area of non-perforated mGluR-2/3$^+$ synapses was detected in the Dutch compared to wildtype mice (**Figure 3E**), both of which had comparable distributions of mGluR-2/3$^+$ particles per PSD area (**Figure 3G-I**).

Dutch mice exhibited defects in PTP, SF, and replenishment of the vesicle pool after depletion.

We assessed several mechanisms related to function of the presynaptic terminal. 11- to 13-month-old Dutch mice and their wildtype littermates showed the same response in paired-pulse facilitation (PPF) **(Figure 4A)**, a type of short-term plasticity that is most likely dependent upon the build-up of Ca^{2+} in the presynaptic neuron due to the close pairing of two action potentials. Accumulation of Ca^{2+} in the presynaptic neuron results in increased neurotransmitter release and subsequently increased response after the arrival of the second action potential[27]. Accordingly, facilitation is no longer present when a long-time interval (1 second) is introduced between the two stimuli. Both wildtype and Dutch mice showed increased facilitation at short time intervals, and no difference was observed between the two genotypes. In contrast to PPF, Dutch mice showed decreased post-tetanic potentiation (PTP) when compared to wildtype mice **(Figure 4B)**. PTP is performed in the presence of D-2-amino-5-phosphonovalerate (D-APV) that blocks NMDA receptors, thereby preventing the onset of plasticity through NMDA-dependent postsynaptic mechanisms that might interfere with its measurement. Due to the DAPV, the fEPSP slope declines back to baseline a few seconds after stimulation. PTP is another form of short-term plasticity and reflects the enhanced release of neurotransmitter due to the increased levels of Ca^{2+} at the presynaptic terminal during the tetanic stimulation[28].

We also studied the effect of the Dutch mutation on the efficacy of presynaptic mechanisms by examining synaptic fatigue (SF) and replenishment of the vesicle pool after its depletion. Both mechanisms appeared to differ between wildtype and Dutch mice. Specifically, the rate of decay examined with SF was faster for the Dutch mice as compared to wildtype **(Figure 4C),** whereas the growth rate during the replenishment phase was higher in the Dutch in comparison to wildtype **(Figure 4D).** The decline in the slope that is observed during SF is thought to represent the depletion of the readily releasable pool (RRP) of neurotransmitter that resides in vesicles at the presynaptic cells, while replenishment depicts the refilling process of the RRP after its depletion with high frequency stimulation. Mechanistically, it has been suggested that both SF and replenishment are dependent on accumulation of residual Ca^{2+} at the presynaptic terminal[27, 28].

Dutch mice exhibited no change in vesicle size or number of docked vesicles but show a trend towards fewer vesicles in the reserve pool at presynaptic terminals in the CA1.

The size, number, and distribution of vesicles were quantified within the presynaptic terminal of 12-month-old mice. No differences were observed in synaptic vesicle sizes at the presynaptic terminal nor the number of

docked vesicles at the synapses of the active zone in Dutch and wildtype mice (**Supplemental Figure 6A-C, E, G**). A decline in the number of synaptic vesicles was found in the reserve pool beyond 300 nm from the active zone of Dutch compared to wildtype mice, but did not reach statistical significance and appeared to be driven by vesicles in large synapses (**Supplemental Figures 6D and 6H**). The hippocampi of Dutch and wildtype mice exhibited no differences in the levels of either the postsynaptic density protein of 95kDa (PSD95) or the presynaptic marker synaptophysin (**Supplemental Figure 7**).

Dutch mice had no change in synaptic distribution of mGluR-2/3+ in the CA1.

Comparable distributions of mGluR-2/3 were observed by immunogold labeling and electron microscopy (EM) at the synapse (**Figures 5A-C**), pre- and postsynaptic compartments (**Figure 5D**), presynaptic active zone, synaptic cleft, peri-synaptic area, postsynaptic membrane, and postsynaptic cytoplasm (**Figure 5E**) in 12-month-old wildtype and Dutch mice.

Dutch mice display shared gene expression profiles across multiple cell types.

Employing brains from Dutch mice alongside sex- and age-matched wildtype littermates, scRNA-seq was performed. Hippocampi were collected from mice at 12 months, an age characterized by established cognitive deficits and high levels of NFA-Aβ and α-CTF in Dutch mice(*18*). After quality control filtering (**Supplementary Figure 8**), 61,642 (32,661 Dutch and 28,981 wildtype) cells were isolated from thirteen distinct clusters which were identified based on the expression of well-known cell-type-specific markers **(Figure 6A-B**, **Supplementary Tables 1-2).** Pseudo-bulk differential expression analysis was used to identify gene expression changes within each cell type. For further analysis (C0-6), clusters of interest were identified based on differentially expressed genes (DEGs; adj. p-value cutoff <0.05) (**Figure 6C-D**, **Supplementary Figures 9-12, Supplementary Table 3**).

Faced with considerable overlap in DEGs, the clusterProfiler(*29*) approach was employed to compare biological themes and identify the Gene Ontology (GO) term enrichment shared among the gene clusters. Among the shared terms were "structural constituent of ribosome" (example genes include: *Rpl23, Rpl30, Rpl37a, Rpl6, Rpl9, Rps24*) **(Figure 7A-B)**, and "amyloid-beta binding" (*Apoe, Atp1a3, Clu, Cst3)* **(Figure 7A, C)**. The shared DEGs and GO terms across clusters were similar to a previously identified "aging signature" in mouse brain(*30*), in which ribosomal protein genes, long noncoding RNA genes (e.g., *Malat1*) as well as ribosome-related pathways and processes, all were significantly affected in 21- to 22-month-old mouse brain across multiple cell types compared to that of their 2-to 3-month-old counterparts. Across clusters, 453 DEGs were found to overlap with the aging-related gene list **(Figure 7D)**, from which 435 genes displayed the same directionality across all clusters (389 upregulated and 46 downregulated, **Figure 7E**) as reported.(*30*).

Dutch mice do not show obvious differences in microglial gene expression.

In many Aβ-depositing mouse models of AD, microglia show prominent transcriptional responses and increased microgliosis with conspicuous peri-plaque localization(*31, 32*). Additionally, the microglial immune response to NFA-Aβ has been implicated as an early, key step in AD pathogenesis. To investigate the microglial responses in more detail, we performed subclustering analysis, which revealed three subpopulations of microglia in the hippocampus (**Figure 8A**). None of the subpopulations showed higher abundance in either genotype. The subpopulations also did not segregate based on expression of established microglial profile markers, such as homeostatic, disease-associated microglia (DAM)(*33*), type I interferon-induced (IFN), or major histocompatibility complex-expressing (MHC) signatures(*34*) (**Figure 8B**). Expression of homeostatic markers was high in all microglia subpopulations and did not change in microglia from the hippocampus of Dutch mice (**Figure 8C**). Additionally, no differences were seen in IBA1 IHC between 12-month-old Dutch and wildtype mice **(Supplementary Figure 13A-B)**. Notably, we also did not find an increase in colocalization of C1q and PSD95 puncta in 12-month-old Dutch mice **(Supplementary Figure 13C)**, as was reported in J20 mice(*35*).

Dutch mice show changes in gene expression associated with translation in excitatory neurons.

Cluster 6 was identified as excitatory neurons and showed the most extensive and unique transcriptional response in Dutch mice (**Figure 6C-D, Figure 9A**). In the GO term analysis, upregulated Cluster 6 DEGs included "cytoplasmic translation" (*Eif2s2, Eef1a1, Eef1b2, Rpl22, Rpl26, Rpl27*) and "negative regulation of amyloid fibril formation" (*Cryab, Pfdn6, Slc25a4, Grin2b, Stmn1, Smad4, Pik3r1*) **(Figure 9B, Supplementary Table 4)**. Notably, "regulation of synapse organization" was also among the top GO terms (*Cask, Cdh8, Gpc4, Grin2b, Mef2c, Nrxn1, Pafah1b1, Ptprd, Tuba1a, Ube2v2, Malat1, Opa1, Mycbp2, Nrcam, Il1rapl1, Arhgef9*), a selection of which are also highlighted in **Figure 9A**. In the GO term analysis, downregulated Cluster 6 DEGs included those related to processes associated with mitochondrial bioenergetic function, i.e., "electron transport chain" (*Mt-Co1, Mt-Co3, Mt-Cytb, Ppp1cb, Calm2*) and "oxidative phosphorylation" (*mt-Atp6, mt-Nd1, mt-Nd3)* (**Figure 9C, Supplementary Table 5**). A selection of unique DEGs from Cluster 6 is shown in **Figure 9D.**

Gene expression was also examined for biological processes associated with presynaptic mechanisms and synaptic transmission, specifically "regulation of synapse organization", a subset of "regulation of cell morphogenesis" (**Supplementary Figure 14A**). Myocyte Enhancer Factor 2C (*Mef2c*) is among this gene list and has been identified as a candidate risk gene for developing late onset Alzheimer's disease (LOAD)(*36, 37*). Considering that oligodendrocytes can modulate presynaptic properties and synaptic transmission(*38, 39*), relevant biological processes such as "modulation of chemical synaptic transmission" were assessed and found among the top biological processes in Cluster 4 oligodendrocytes (**Supplementary Figure 14B, Supplementary**

Table 6). Approximately 25% of the genes present in this biological process are among DEGs in Cluster 6. Genes associated with synaptic transmission that were among those specific to Cluster 4 included transferrin (*Trf*), a recently identified biomarker of AD(*40*), and carbonic anhydrase 2 (*Car2*). Carbonic anhydrase inhibitors have been considered for prevention of neurovascular dysfunction in CAA and AD due in part to their ability to prevent mitochondrial dysfunction and oxidative stress (*41, 42*).

A cell-cell communication analysis was performed using the software tool Cellphone DB to compute interactions between clusters based on the known Ligand-Receptor (LR) pairs (**Supplementary Table 7**). For each LR in each sample, the software estimated the interaction strength from Cluster 4 oligodendrocytes to Cluster 6 excitatory neurons, from which differences in interaction strengths between Dutch and WT mice were evaluated by t-test. Twelve LR pairs showed significant interaction difference at p-value ≤ 0.05, and all twelve presented higher interactions in Dutch mice. Among these LRs, *Gabbr2* was up-regulated in the Cluster 6 excitatory neurons in Dutch mice compared to those of wildtype (log2FC = 0.19, adj. p-value = 3.6E-4). In Dutch compared to wildtype mice, stronger interactions were exhibited by *Gabbr2* with multiple GABA inhibitory neurotransmitter complex partners including GAD2, SLC6A11, SLC6A1, SLC6A8, SLC6A6, and SLC32A1.

Dutch mice have fewer and smaller mitochondria in CA1 presynaptic boutons.

The number and morphology of mitochondria were assessed in the presynaptic terminals of excitatory neurons in the *stratum radiatum* of wildtype and Dutch mice (**Figure 10A, B**). Previously, NFA-Aβ has been reported to cause imbalance in mitochondrial fission-fusion, leading to fragmented morphology and altered function(*43-47*). Fewer presynaptic mitochondria were found in Dutch compared to wildtype mice (**Figure 10C**, mean ± SD for total of mitochondria per presynaptic bouton = 106.3 ± 20.06 in wildtype and 75.60 ± 14.62 in Dutch, p= 0.0324 by two-tailed unpaired t test). This deficiency in the Dutch mice was likely due to fewer straight mitochondria at the presynaptic terminals, because curved mitochondria numbers were unchanged between genotypes (**Figure 10D-F**, mean ± SD for number of straight mitochondria = 101.3 ± 17.95 in wildtype and 71.80 ± 15.06 in Dutch, p= 0.314 by two-tailed unpaired t test. Mean ± SD for number of curved mitochondria = 5.00 ± 2.16 in wildtype and 3.80 ± 2.59 in Dutch). In assessments of differences in morphology, mitochondrial diameters were comparable (**Figure 10G**, mean ± SD = 328.6 ± 19.29 in wildtype and 301.8 ± 28.73 in Dutch), but shorter lengths were found in 12-month-old Dutch compared to wildtype mice, as estimated from the number of layers traversed in the consecutive sections by each mitochondrion (**Figure 10H**). Each layer traversed was ~ 80 nm using 3-dimensional reconstruction of measurements from serial sections.

Dutch mice have reduced mitochondrial electron transport chain complex I activity.

The activities of electron transport chain complexes I-IV and citrate synthase were assessed in hippocampi from 7- and 12-month-old wildtype and Dutch mice as described(*104, 105*). Each complex activity was standardized to either µg protein (p=0.0355) or to mU/ml of citrate synthase (CS) and analyzed using an unpaired t test (p=0.0197) (**Supplemental Figure S15, Supplemental Tables 8, 9**). Compared to that of wildtype, mitochondrial complex I activity was reduced in 12- but not 7-month-old Dutch mice.

Discussion

The purpose of this study was to elucidate the molecular pathogenesis underlying the learning behavioral deficits exhibited by the Dutch mouse model which accumulates NFA-Aβ and APP α-CTFs but never forms amyloid fibrils. Our study adds to the few that have detected endogenously generated NFA-Aβ without concurrent presence of fibrils or the presence of synthetic peptides. The location of Dutch NFA-Aβ in the mouse brain was ascertained using immunohistochemistry with A11 antibody and fluorescence microscopy with a cyclic peptide-FITC ligand(*22*). In contrast to the apparently normal density and morphology of synapses in excitatory neurons, we observed abnormalities in the apical dendrite structure in the CA1 of Dutch mice(*24*). Therefore, we sought to determine whether Dutch mice exhibit plasticity impairment during growth from 7 to 13 months, before and after NFA-Aβ aggregation can be detected in the brain. A variety of electrophysiological techniques were used to examine short- and long-term synaptic plasticity. Although no changes in basal synaptic transmission nor long-term synaptic plasticity were observed at the CA1-CA3 area of the hippocampus in slices obtained from young (7- to 7.5-month-old) and old (10.5- to 11-month-old) Dutch mice, impaired short-term synaptic plasticity developed at the CA1-CA3 synapses first appearing at the age of 11 to 13 months.

Assessment of the I-O curve and the LTP response after high frequency stimulation in the CA1-CA3 synapses, indicated no difference between Dutch and wildtype genotypes, suggesting that NFA-Aβ and/or APP α-CTFs affect neither spontaneous activity nor the long-term plasticity in the examined age window. In contrast, intracerebroventricular injection of synthetic Dutch and Arctic (APP^{E693G}) Aβ peptides have been previously reported to block LTP in rats(*48*). Long-term plasticity changes may possibly be manifested only in older Dutch mice. In addition, changes may occur in other types of long-term synaptic plasticity such as long-term depression, as well as in other areas of the hippocampus, such as mossy fiber-CA3 synapses. Despite normal *long-term* synaptic plasticity in young or old Dutch mice, defects in several types of *short-term* plasticity emerged in the aging 11- to 13-month-old Dutch mice. Specifically, mice carrying the Dutch mutation exhibited impaired PTP, synaptic fatigue (SF), and impaired replenishment of the synaptic pool, but unaffected PPF. Both PPF and PTP are related to accumulation of Ca^{2+} at the presynaptic terminal, due (respectively) to two closely paired pulses or to high frequency stimulation, both of which augment neurotransmitter release. Although impaired PTP without

changes in PPF might seem contradictory, previous studies have shown that PPF and PTP are governed by different synaptic vesicle proteins. Specifically, the absence of the synaptic vesicle protein synaptogyrin I in mice causes impaired PTP, but its absence has no effect on PPF(*49*). More in-depth investigation of synaptic vesicle proteins is required to elucidate the molecular mechanisms underlying impaired plasticity in Dutch mice.

Our earlier studies identified abnormalities in the dendritic structure of excitatory neurons in the CA1 of Dutch mice(*24*). Although mGluR2 is necessary for synaptic binding of oAβ, and mGluR3 has been reported to mediate the effects of NFA-Aβ on synaptic function (*50, 51*), the CA1 of Dutch and wildtype mice exhibited similar synaptic density of mGluR-2/3$^+$. Moreover, similar levels of mGluR-2/3 were observed in the hippocampus as well as the pre- and postsynaptic compartments of axospinous synapses in the CA1 of Dutch and wildtype mice. In contrast, mGluR-2/3 expression was previously reported to be reduced in the CA1 of AD patients(*52, 53*), depending on which hippocampal subfield was examined(*53, 54*). Furthermore, in primary rat neurons treated with NFA-Aβ, mGluR-2/3 levels initially increased as the immediate response but decreased following activation, suggesting that synaptic activity contributes to the effect of NFA-Aβ on receptor subunit expression (*55*). In CA1 synapses in the Dutch mice, differences were identified in the greater PSD area of small mGluR-2/3$^+$ synapses and in the reduced number and size of presynaptic mitochondria (discussed below in the context of the transcriptomic data). Recruitment of mGluR1 or NMDA receptors to the PSD in the presence of NFA-Aβ could be a possible explanation for our finding of increased PSD area on non-perforated mGluR-2/3$^+$ synapses in the Dutch mice.

Dutch mice showed altered presynaptic plasticity mechanisms related to the RRP of neurotransmitters. Specifically, Dutch mice exhibited respectively faster rates of decay and recovery during high frequency stimulation that causes fatigue at the synapses and after synaptic depression. The depletion and replenishment of the RRP of neurotransmitters appear more profound in Dutch mice in comparison to wildtype mice. The RRP of neurotransmitters refers to a small subset of neurotransmitter-containing vesicles residing close to the presynaptic terminal (either docked or non-docked) and disposed for immediate released during stimulation. Besides accumulation of Ca^{2+}, several parameters could affect the decay of the RRP including the number of active zones and the number of docking sites at the presynaptic terminal. In turn, replenishment of the RRP could be mediated by the Ca^{2+}-mediated recruitment of vesicles from the reserve pool to the RRP. The detailed mechanism of the observed changes in the RRP of neurotransmitter requires further investigation.

The required interplay between pre- and postsynaptic neurons during synaptic plasticity suggests that presynaptic changes could affect activation of postsynaptic neurons eventually leading to long-term synaptic impairments. Despite the observed changes in the release probability and the RRP of neurotransmitters, high-frequency stimulation resulted in normal LTP, suggesting that Dutch mice do not display derangement of postsynaptic mechanisms at the ages studied, an observation that could be attributed to compensatory

mechanisms. In one possibility, the decrease in release probability with PTP may be balanced by the increased depletion and replenishment of the RRP.

In contrast to previous reports, hippocampal microglia (Cluster 2) did not show a high number of DEGs, nor were differences in expression of microglia DAM and homeostatic signatures observed between Dutch and wildtype mice. The highest total and unique number of DEGs were found in excitatory neurons (Cluster 6) and were associated with processes such as "protein translation" and "electron transport chain". Knowing that presynaptic function can be impacted by oligodendrocytes(*38, 39*), and that Cluster 4 had the second highest total number of DEGs, LR pairs were examined between oligodendrocytes and excitatory neurons, which indicated that *Gabbr2* showed stronger interaction with multiple GABA inhibitory neurotransmitter complex partners in Dutch mice.

Finally, prior studies using bulk RNAseq on the entorhinal cortex and dentate gyrus of wildtype and Dutch mice were re-examined using scRNA-seq, revealing no obvious differences in microglial transcriptional profiles. Moreover, the Dutch mouse hippocampal transcriptome did not present neuroinflammatory and microglial transcriptomic responses that are commonly observed in models with senile plaques and amyloid fibrils (e.g., PSAPP, 5XFAD). Failure to capture such a response may be due to the cell populations selected for profiling. Indeed, previous characterization indicated that microgliosis in other Dutch mice (*56*) was not generalized but was limited to the immediate proximity of Aβ-laden vessels. In this study, no differences were observed in microgliosis nor were increases detected in C1q colocalization with PSD95, a process reported to be induced by NFA-Aβ, that drives activation of complement and synapse elimination(*35*). In a transcriptomic analysis on postmortem brain tissue from humans with HCHWAD, inflammatory genes and pathways were also not prominently represented(*57*). As detected in the Dutch mouse, the main pathways over-represented in the human HCHWAD brains were related to mitochondrial dysfunction (*57*). The principal up-regulated pathways entail interactions of the extracellular matrix and transforming growth factor beta signaling. In a clustering analysis to identify molecular subtypes of AD, gene expression profiles have been gleaned from the Mount Sinai Brain Bank (MSBB-AD) and Religious Orders Study-Memory and Aging Project (ROSMAP)(*58*). The Dutch mouse gene signature significantly correlated with the subtype of "typical" or "Aβ predominant" AD presentation, and was characterized by increased immune activation and inflammation, as well as decreased synaptic signaling and dysregulation in neuronal processes and synaptic signaling (*58*). Considering our current results, the overlap between the Dutch model and the AD subtypes may be more closely associated with alterations in synaptic signaling than immune activation. Taken together, the results minimize the contribution of neuroinflammation in the pathogenesis leading to learning disorder in Dutch mice.

Among the most dysregulated pathways in the Dutch mouse brain was "structural constituent of ribosome", which is associated with the most significant GO term in Cluster 6 "protein translation". Increasing evidence

suggests that dysregulation of mRNA translation contributes to the pathogenesis of several neurological diseases, including AD(*59*). Accumulation of Aβ is linked to increased protein synthesis through loss of interactions from key translation regulators, particularly translation repressors(*60*). When the cell-type specific transcriptional responses to NFA-Aβ were examined, altered transcriptional profiles were identified in multiple cell types. In these profiles, overlap with GO term enrichment analysis indicated "amyloid-beta binding" among the shared processes. Previously, both DEGs and processes altered in the Dutch mouse were identified in an aging-related signature(*30*), and a recent study revealed that Aβ accumulation triggers fragmentation of the Golgi apparatus, a key organelle needed for protein processing(*61*). A vicious cycle can be created because Golgi fragmentation exacerbates Aβ production(*61*). On the contrary, prevention of the phosphorylation of Golgi structural proteins, such as GRASP65, has rescued apparatus structure and reduced Aβ secretion(*61*).

The Cluster 6 neuron DEGs downregulated in Dutch mice compared to wildtype were associated with mitochondrial functions, specifically GO terms "electron transport chain" and "oxidative phosphorylation". In concordance, cellular respiration pathways were the most significantly altered pathways identified in the transcriptomic analysis from human patients with HCHWAD(*57*). Moreover, fewer and smaller mitochondria were found at the presynaptic terminals of excitatory neurons in the CA1 of Dutch mice compared to wildtype. Strong evidence linking NFA-Aβ to mitochondrial complex I endpoints has come from the combined use of a genetically encoded fluorescent redox sensor and multiphoton microscopy to image and monitor *in vivo* the mitochondrial redox imbalance in neurons from the APP/PS1 transgenic mouse model of AD(*62*). The deposition of Aβ plaque and particularly soluble NFA-Aβ species were shown to lead to increased mitochondrial oxidative stress levels within neuronal mitochondria(*62*). Mitochondrial redox dysregulation in neurons was prevented by inhibition of mitochondrial calcium influx, which is pathologically exacerbated in Aβ plaque-depositing mice(*62*). Furthermore, mitochondria-targeting antioxidants decreased mitochondrial oxidative stress back to basal levels and reverted Aβ plaque-associated dystrophic neurites in the APP/PS1 mice(*62*). Defective mitochondrial complex I activity has been detected in living human AD brain(*63*). Altered mitochondrial fission and fusion has been observed in the hippocampus of human AD and in Tg2576 mice and may explain the observations of decreased mitochondrial density and smaller mitochondria in Dutch mice(*46, 65*). Increased mitochondrial fragmentation has been reported in neurons exposed to NFA-Aβ(*66*). Elevated levels of NFA-Aβ impaired transport of mitochondria in dendrites and axons and increased mitochondrial permeability in mice harboring the Osaka mutation of APP (APP^{E693D})(*72*). Mitochondria-targeting therapy holds promise for treating neurodegenerative disease (*108, 109*) and could be used in combination with other anti-Aβ and anti-tau therapies to arrest AD progression. Our finding of reduced density of presynaptic mitochondria in the CA1 of Dutch mice suggests that NFA-Aβ might also affect transport of mitochondria to axon terminals. The dysfunction of

mitochondria and endoplasmic reticulum (ER) protein translation observed in Dutch mice is consistent with observations in other mouse models of AD(*62, 69*).

In conclusion, the current study provides several notable observations. Two independent methods visualized for the first time *endogenously generated* NFA-Aβ *in vivo* in the absence of artefacts related to the intracerebral injection of synthetic peptides or to the concurrent presence of Aβ fibrils. Changes related to the *APP^{E693Q}*-genotype were revealed in NFA-Aβ histopathology, in expression of mitochondrial and neuronal translation-related genes, in presynaptic physiology, and in the ultrastructure of presynaptic mitochondria. Neuroinflammation is not a major contributor to the pathogenesis in Dutch mice. Neither fibrillar Aβ nor neuroinflammation is required for the aging-related learning behavior phenotype.

This study of APP^{E693Q} mice is particularly timely considering the promising outcome of clinical trials for lecanemab(*70*) and donanemab (*71*). Unlike other anti-amyloid antibodies, lecanemab selectively binds to NFA-Aβ, leading to its neutralization and elimination(*70, 71*). After 18 months of treatment with either antibody, participants experiencing mild cognitive impairment due to AD showed a modest but statistically significant 27% slower rate of cognitive decline with lecanemab and 35% slower rate with donanemab. The modest clinical response to therapeutic infusion of anti-amyloid antibodies may be attributable, at least in part, to the persistence of residual toxic NFA-Aβ that goes undetected by current fluid biomarker analysis and amyloid fibril imaging. The potential clinical importance of NFA-Aβ is reinforced by the recent discovery of a new FAD mutation (*APP^{D678N}*) that also favors NFA-Aβ accumulation (*72*). Therapeutic depletion of NFA-Aβ may be required for optimum and more robust cognitive benefits from anti-amyloid antibody infusion. Newer methods for imaging NFA-Aβ (e.g., [^{64}Cu]-NOTA cyclic D,L-α-peptides) in the living human brain (*22*) and assays for detection of NFA-Aβ in biofluids (*106, 107, 110*) may provide earlier diagnosis and improved monitoring of patients undergoing Aβ immunodepletion therapy by assuring that both fibrillar and nonfibrillar Aβ conformers are cleared as completely as possible.

Materials and Methods

Experimental animals

Male and female 2- to 24-month-old mice were used in this study. All animals were group housed in microisolator cages under a 24-hour light/dark cycle and given *ad libitum* access to food and water. The Dutch mice used in the study express the human *APP* gene harboring the Dutch mutation (*APP*^{E693Q}) under control of the *Thy1* promoter(*18*). All animal procedures were conducted in accordance with the National Institute of Health Guidelines for the Care and Use of Experimental Animals and were approved by the Institutional Animal Care and Use Committees at the Icahn School of Medicine at Mount Sinai.

Dot blot assay

The cerebral cortices of mouse brain samples were thawed and lysed by adding 2x volume of lysis buffer [5.0 mM NaCl, 5.0 mM EDTA, 1.0 mM dithiothreitol (DTT), and protease inhibitor mixture in 25 mM Tris buffer, pH 7.5] containing 0.5 mm Zirconium Oxide Beads (ADV-ZrOBO5, ORNAT, Israel), homogenized with a tissue grinder motor pestle (Alex-Red, Israel) on ice, and the mixtures were centrifuged at 4°C at 14,000 rpm for 15 minutes. The total protein in each sample was quantified by Nano-Drop absorbance quantification at 280 nm.

For dot-blot analysis, equal amounts of samples of diluted proteins (100 ng/μL) were directly applied in duplicate onto nitrocellulose membrane and dried at room temperature for 30 minutes. The membrane was then blocked with 5% bovine serum albumin (BSA) in tris-buffered saline (TBS) (pH 7.4) containing 0.1% (v/v) Tween 20 (TBST). Blots were incubated overnight at 4°C with: 6E10 mouse monoclonal antibody (1:1000); rabbit polyclonal antibody recognizing prefibrillar oligomers, A11 (1:5000); rabbit monoclonal antibody OC recognizing fibrillar oligomers (1:1000); anti-murine-α-tubulin antibody (1:5000) in TBST containing 5% BSA. The membranes were washed with TBST three times and incubated for 1 hour at room temperature with horseradish peroxidase (HRP) conjugated either to an anti-mouse IgG (1:10000) or anti-rabbit IgG (1:5000) in 5% BSA in TBST. The blots were washed with TBST and developed using the enhanced chemiluminescence (ECL) reagent kit (Pierce) in Amersham Imager 680 machine. Images were then used to analyze the mean of integrated densities of each blot using ImageJ software (National Institutes of Health, USA)(*73*).

Immunohistochemistry (IHC) and CP-2-FITC incubation

All staining was performed in paraformaldehyde (PFA)-fixed 30 μm thick sections of Dutch and wildtype mouse brains. For A11 immunohistochemistry and cyclic D,L-α-peptide-2 (CP-2)-FITC fluorescence, free-floating sections were washed with 0.1 M phosphate buffer saline (PBS) in 0.1% Triton (PBST) three times for 15 minutes and blocked with 2% bovine serum albumin (BSA) for 1 hour. Sections were incubated with a rabbit polyclonal antibody A11, recognizing Aβ prefibrillar oligomers (1:2500) in antibody buffer (PBST containing 2% BSA) overnight at 4°C. Sections were washed and incubated with Alexa Fluor 680 secondary goat anti-rabbit antibody (1:1000 Abcam) in antibody buffer for 1 hour at room temperature. Sections were then incubated with fluorescein-5-isothiocyanate (FITC)-conjugated CP-2 [0.15 mg/mL in 2% dimethyl sulfoxide (DMSO)] for 3 hours at room temperature. Sections were washed for 10 min in 1% PBST and kept for 15 min to dry. Sections were then mounted on super frost glass slides, embedded with Fluro-Gel II mounting medium containing DAPI (4',6-diamidino-2-phenylindole dihydrochloride) (Electron Microscopy Sciences, Hatfield PA), cover-slipped, and stored in the dark for microscopic imaging. Immunofluorescence images containing the hippocampus, cortex, and thalamus were obtained using the 20x objective of a Leica DM6000 microscope (Leica Microsystems, Wetzlar, Germany), coupled to a controller module and a high sensitivity 3CCD video camera system (MBF

Biosciences, VT) and an Intel Xeon workstation (Intel). All images were analyzed by the Image J program(*73*), and one section from each mouse (from a total of 5 mice in each group) were taken for analysis. The average intensity was calculated from the area of the hippocampus, cortex, and thalamus.

Brain tissue homogenization and Aβ ELISAs

Fresh frozen brain tissue (cerebrum) was homogenized at a 1:5 tissue:buffer ratio, using a Precellys Evolution homogenizer (Bertin Technologies, Montigny-Le-Bretonneux, France) to sequentially extract TBS, TBS with 0.5% Triton X-100 (TBS-T) and formic acid (FA) soluble fractions of Aβ. After homogenization in TBS, samples were centrifuged at $16,000 \times g$ for 1 h at 4 °C (TBS_{16K}). For a subset of experiments, a fraction of this supernatant was further centrifuged at $100,000 \times g$ followed by collection of the supernatant (TBS_{100K}). Homogenization of the remaining tissue pellet was repeated according to the same procedure with TBS-T, then followed by FA. Brain extracts were analyzed with Aβ1-40 and Aβ1-42 ELISAs, as previously described(*74*). In brief, 96-well half-area plates were coated over night with 50 ng/well of anti-Aβ40 (custom production, Agrisera) or anti-Aβ42 (700254, Thermo Fisher Scientific, USA), then blocked with 1% BSA in PBS for 3 h at RT. TBS_{16K}, TBS-T and FA brain extracts were diluted in ELISA incubation buffer (PBS, 0.1% BSA, 0.05% Tween) and incubated over night at 4 °C, followed by detection with 0.5 µg/ml biotinylated 3D6 and HRP-conjugated streptavidin (Mabtech AB, Nacka, Sweden). Signals were developed with K Blue Aqueous TMB substrate (Neogen Corp., Lexington, KA, US). Plates were developed and read with a spectrophotometer at 450 nm. The TBS and TBS-T extracts were also analyzed for soluble Aβ aggregates using two different sandwich ELISAs. The first, based on 3D6 as both capture and detection antibody, allows for detection of Aβ aggregates from the size of a dimer, but not for monomeric Aβ. The second preferentially detects larger soluble Aβ aggregates as it utilizes the Aβ protofibril selective antibody mAb158 for capture and 3D6 for detection(*75*). Ninety-six-well half-area plates were coated over night with 3D6 (50 ng/well) and blocked with 1% BSA in PBS for 3 h at RT. Brain extracts were diluted in ELISA incubation buffer (PBS, 0.1% BSA, 0.05% Tween) and incubated over night at 4 °C, followed by 3D6biotin detection and development as above.

Slice preparation, electrophysiological recordings, and analysis

For electrophysiological experiments, hippocampi from 7- to 13-month-old Dutch and wildtype mice were excised on ice and coronal hippocampal slices (400 µm) were cut with a tissue chopper. Slices were transferred to a recording chamber where they were allowed to recover for 2 hours at 29°C and perfused with artificial cerebrospinal fluid (ACSF) containing (124.0 mM), KCl (4.4 mM), Na_2HPO_4 (1.0 mM), $NaHCO_3$ (25.0 mM), $CaCl_2$ (2.0 mM), $MgCl_2$ (2.0 mM), and glucose (10.0 mM). ACSF was bubbled with carbogen (95% O_2 and 5% CO_2) and perfused at a flow rate of 2 ml/minute.

Field excitatory post-synaptic potentials (fEPSP) were measured after stimulating the Schaffer collateral fibers with a bipolar tungsten electrode placed at the CA3 and recording at the *stratum radiatum* of CA1 with a glass pipette filled with ACSF. Basal synaptic transmission was evaluated by plotting the relationship between increased voltages (5 V to 35 V) and evoked fEPSP responses, generating an input-output (I-O) curve. Recordings were measured at approximately 35% of the maximum evoked response as determined by the I-O curve. For LTP recordings, a 30-minute baseline was recorded prior to theta-burst stimulation (4 pulses at 100 Hz, with the bursts repeated at 5 Hz and 3 tetani of 10-burst trains administered at 15-second intervals), responses were recorded for 2 hours after tetanization and measured as fEPSP slope expressed as percentage of baseline. Paired-pulse facilitation (PPF) was induced by pairing two pulses at intervals of 15, 20, 30, 40, 50, 100, 200, 300, 500 and 1000 ms. PPF was expressed as the ratio between the second and the first fEPSP slopes. Post-tetanic potentiation (PTP) was induced by the above-described theta-burst stimulation protocol in presence of the NMDA receptor blocker D-APV (50 µM). Synaptic fatigue (SF) and replenishment were measured by 100 Hz stimulation followed by 7 stimuli spaced at 125 msec from each other in the presence of D-APV (50 µM).

Results were analyzed in pClamp 11 (Molecular Devices) and fEPSP slopes were compared by 2-way ANOVA for repeated measures for the I-O curve, LTP, PPF and PTP recordings. For statistical comparison of synaptic fatigue and replenishment, the rates of exponential growth/decay (k) extra sum-of-squares F test was used to test whether k differed between data sets vs. the data sets sharing one k-value. Statistics were performed using GraphPad Prism 9 software. Data were expressed as mean ± SEM. The level of significance was set at $p < 0.05$.

Perfusions and tissue processing for electron microscopy and array tomography

For electron microscopy (EM) and array tomography synapse studies, 12-month-old Dutch and wildtype mice were anesthetized intraperitoneally, and transcardially perfused with 1% PFA in phosphate-buffered saline (PBS; pH 7.4) followed by 4% PFA with 0.125% glutaraldehyde in PBS as described previously[24, 76, 77]. The brains were carefully removed, hemisected, postfixed overnight at 4°C in 4% PFA in PBS with 0.125% glutaraldehyde for synapse and immunogold analyses, or by 2% PFA with 2% glutaraldehyde in PBS for presynaptic vesicle analysis, and sectioned on a Vibratome (Leica VT1000S, Bannockburn, IL) into 250-µm thick sections. All sections were stored at 4°C in PBS with 0.1% sodium azide until use. Coronal sections encompassing the CA1 region of the hippocampus were prepared for EM as reported[24, 77]. For synapse and immunogold analyses, slices were cryoprotected in graded phosphate buffer/glycerol washes at 4°C, and manually microdissected to obtain blocks containing the CA1 region. The blocks were rapidly freeze-plunged into liquid propane cooled by liquid nitrogen (-190°C) in a Universal cryofixation System KF80 (Reichert-Jung) and subsequently immersed in 1.5% uranyl acetate dissolved in anhydrous methanol at -90°C for 24 hours in a cryosubstitution unit (Leica). Block temperatures were raised from -90°C to -45°C in steps of 4°C/hour, washed with anhydrous methanol, and infiltrated with Lowicryl resin (Electron Microscopy Sciences) at -45°C. The resin was polymerized by exposure

to ultraviolet light (360 nm) for 48 hours at -45°C followed by 24 hours at 0°C. For presynaptic vesicle studies, tissue blocks were washed overnight in 0.1 M sodium cacodylate buffer, placed in 1% tannic acid/0.1 M cacodylate buffer solution at room temperature for one hour and washed 3 times for 10 minutes each in 0.1 M sodium cacodylate. The samples were then incubated for 1 hour at room temperature in 2% osmium tetroxide/double distilled water (ddH$_2$O), washed again in 0.1 M sodium cacodylate 3 times for 10 minutes then subjected to a graduated ethanol dehydration series. After dehydration, the tissue samples were infiltrated with Spurr Low Viscosity Embedding Kit (Sigma Aldrich) and polymerized for 12 hours at 70°C. For both EM preparations, block faces were trimmed, ultrathin sections (90 nm) were cut with a diamond knife (Diatome) on an ultramicrotome (Reichert-Jung), and sections were collected on formvar/carbon-coated nickel slot grids (Electron Microscopy Sciences).

Post-embedding immunogold labeling for EM

Immuno-gold labeling was performed using a previously optimized method(*78*). Briefly, the EM sections were treated with 0.1% sodium borohydride in 50 mM glycine prepared in Tris buffer for 6 minutes in the dark, to remove aldehydes. Sections were then subjected to six quick rinses and two 10-minute washes in TBS. The sections were incubated for 30 min in blocking buffer consisting of 2% human serum albumin (HSA) in TBS with 0.1% Triton X-100. After three short washes in TBS, the sections were incubated in anti-glutamate receptor 2 & 3 rabbit polyclonal antibody (Millipore AB1506, diluted 1:100 in 2% HSA in TBS) at room temperature overnight with gentle shaking. Following six quick rinses and two 5-minute washes in TBS, the grids were incubated in blocking buffer for 20 minutes, washed three times, and incubated for 90 minutes at room temperature in goat anti-rabbit IgG conjugated to 10-nm gold particles (Aurion 25362), diluted 1:40 in 2% HSA and 5 mg/ml polyethylene glycol in TBS. The grids were rinsed six times and washed for 5 minutes in TBS and then in deionized water. Imaging was done using an H-7000 electron microscope (Hitachi High Technologies America, Inc.) with an AMT Advantage CCD camera (Advanced Microscopy Techniques) at 15,000x. The electron micrographs were adjusted for brightness and sharpness with Photoshop CS5 (Adobe Systems).

Quantifying synapse density, ultrastructure, and distribution of immunogold particles at mGluR-2/3-labeled synapses

All EM analyses were conducted by an experimenter blinded to the genotypes of the mice. For synapse quantification and ultrastructural analysis, nine sets of serial images across the same set of 5 consecutive ultrathin sections were taken from the *stratum radiatum* of CA1 and imported into Adobe Photoshop (version CS5). To obtain a stereologically unbiased population of synapses for quantitative morphologic analysis, we used a disector approach on ultrathin sections as described(*24, 76, 77, 79*). Briefly, all axospinous synapses were identified within the first and last 2 images of each 5-section serial set and counted if they were contained in the reference image

(layers 1 and 4) but not in the corresponding look-up image (layers 2 and 5). To increase sampling efficiency, the reference image and look-up image were then reversed; thus, each animal included in the current study contributed synapse density data from a total of 18 disector pairs. The total volume examined was 11.317 μm^3, and the height of the disector was 180 nm. Axospinous synapse density (per μm^3) was calculated as the total number of counted synapses from both images divided by the total volume of the disector. The criteria for inclusion as an axospinous synapse included the presence of synaptic vesicles in the presynaptic terminal and a distinct asymmetric PSD separated by a clear synaptic cleft. For a synapse to be scored as perforated it had to display two or more separate PSDs.

To measure structural parameters such as spine head diameter and PSD length, we identified all axospinous synapses in the middle portion of three serial sections. The synapse and spine head were followed through layers 2-5. In each layer, the PSD length (if present) and the spine head diameter at its widest point parallel to the synapse were measured. PSD area was determined by summing the PSD lengths through all sections and multiplying by section thickness (90 μm). This same process was repeated for unique synapses in layer 4 (reference layer 5), following the synapses from layer 4 to layer 1. Each of these synapses were also classified as non-perforated or perforated. A total of 734 total synapses (341 from Dutch mice and 393 from wildtype) were analyzed in this manner. An average of 81 synapses per animal and a minimum of 71 synapses per animal were sampled. For each unique synapse in layer 2 in each series (as identified for PSD length and spine head analysis above), the synapse was followed through layers 2 though 5 and the number of immunogold particles in each of the following areas were counted: active zone (presynaptic, within 45 nm of the membrane across from the PSD; region A), synaptic cleft (region B), PSD (region C), peri-synaptic (peri—PSD, within 200 nm of the PSD on the post-synaptic spine head; region D), postsynaptic membrane (>200 nm away from the PSD; region E), and postsynaptic cytoplasm (region F). A cutoff of 45 nm away from the membrane was used to differentiate between receptors at the membrane and receptors in the cytosol.

For presynaptic vesicle analysis, 100 synapses per animal were analyzed from at least 30 different images taken at a magnification of 8,000x. PSD length, maximum spine head diameter, and cytosolic area of the presynaptic terminal were measured using Adobe Photoshop 2023. Vesicles in the presynaptic terminal were labeled with a dot in the center of the vesicle, and a line was drawn over the PSD. An algorithm was written in Python that measured the shortest distance between each vesicle (represented by the dot in the center) and its corresponding PSD (represented by the manually drawn line over the PSD) and reported a list of measurements for all vesicles within an adjustable threshold distance (**Supplemental Figure 5**). For each synapse, the total number of vesicles was determined with output from this algorithm, along with their individual distances from the PSD. The vesicles were divided into bins of 10 and 50 nm for comparison of spatial distribution within the

presynaptic terminal. Additionally, vesicles within 10 nm of the PSD were classified as docked, based on a 17 nm

average minimum diameter of vesicles in our samples(*80-82*).

ScRNA-seq and data analysis

For the scRNA-seq experiments, fresh hippocampi were dissected from 12- to 13-month-old Dutch and

wildtype mice, and the tissue was dissociated into a single-cell suspension following the 10X Genomics sample

preparation protocol. We pooled the hippocampi from 3 mice of the same gender, genotype, and age to reach the

target concentration of 2000 cells/µl. The single cell cDNA libraries were generated using a V3 Chromium kit

(10X Genomics) and sequenced using a NovaSeq S4 (Illumina).

The raw sequencing data were processed using the standard Cell Ranger pipeline to generate unique molecular

identifier (UMI) count matrices. Starting from UMI matrices, we first performed quality control (QC) by removing

low-quality cells with either too few (< 200) or too many (> 4000) genes detected, retaining 61,642 cells after

filtering. We then removed insufficiently detected genes by keeping 20,548 genes expressed in more than one cell.

After QC, we obtained 5,266 ~ 10,949 cells per individual, and an average of 618 unique genes per cell per

individual. We performed a clustering analysis using a well-established scRNA-seq data integration workflow

based on R package Seurat (v3)(*83*). Briefly, the UMIs data were first normalized by sequencing depth and log-

transformed using the LogNormalize method implemented in Seurat. The 2,000 most variable gene features were

identified, scaled, and centered. The scRNA-seq data across all sequencing libraries were combined based on the

canonical correlation analysis (CCA) integration method available from Seurat. Next, dimensional reduction was

performed using principal component analysis (PCA) based on the 2,000 most variable genes. The top 25 principal

components (PCs) collectively explaining more than 90% of the variance were selected for calculating 2D

dimensional reductions by Uniform Manifold Approximation and Projection for Dimension Reduction

(UMAP)(*84*). The same top 25 PCs were also used to compute the nearest neighbor graph and the subsequent cell

clusters with the Louvain algorithm implemented in Seurat (*83*). This clustering analysis resulted in 14 clusters

(named as Clusters 0 – 13, sorted by cluster size in a decreasing order) at a clustering resolution of 0.1. These

clusters contained 267 to 20,702 cells each.

Cluster cell type annotation

For each cluster, we first interrogated the expression patterns of known gene markers to annotate clusters into

major cell-types: inhibitory neurons (*Gad1*, *Gad2*), excitatory neurons (*Slc17a7*), astrocytes (*Aqp4*),

oligodendrocytes (*Mog*), microglia (*Csf1r*, *Tyrobp,* and *Trem2*), oligodendrocyte progenitor cells (*Vcan* and

Pdgfra), endothelial cells (*Flt1*), and pericytes (*Vtn* and *Pdgfrb*). Next, we calculated *de novo* cluster signatures

by comparing the cells in a cluster against the cells of the other clusters using Wilcoxon rank sum test in Seurat.

We defined cluster marker genes as those upregulated by at least 1.2-fold and with Bonferroni adjusted p-value

less than 0.05. To assist the annotation of cell type of each cluster, we overlapped the *de novo* cluster marker gene

signatures with a large-scale collection of cell type markers curated from over 1,054 single-cell experiments(*85*), with p-value significance of the overlaps computed by a hypergeometric test.

Cluster-specific differential gene expression and functional enrichment analysis

DEGs between Dutch and wild type sample groups in each cluster were identified using MAST(*86, 87*) implemented in Seurat. Similarly, DEGs between males and females, and DEGs between Dutch and wild type in each sex group were identified using MAST. DEGs were determined at the cutoff of Bonferroni-corrected p-value ≤ 0.05. Functional enrichment of DEGs with MSigDB gene annotation collections was examined by a hypergeometric test. Results with Benjamini-Hochberg (BH) correction p-value ≤ 0.05 were considered statistically significant.

Microglia subclustering analysis

We studied a subset of microglial cells from the full data matrix. PCA was performed and then the top PCs collectively explaining more than 90% of the variance were selected for calculating dimensional reduction by UMAP (*84*) as well as for clustering analysis using Seurat (*83*). To test if the microglia subclusters represent known subtypes, we computed single cell signature scores (*88*) corresponding to homeostatic, disease-associated microglia (DAM), type I IFN-induced (IFN) and MHC-expressing (MHC) microglia based on signature genes curated by Chen and Colonna (*34*).

Quantifying presynaptic mitochondrial numbers and morphology

For analysis of mitochondria, 12 series of 9 consecutive 90-nm thick sections were selected from each mouse using a systematic-random approach for EM imaging. All mitochondrial analyses were performed by the same experimenter, blinded to the group designation of subjects as described (*89*). For each series of 9 consecutive sections, the middle fifth section was used as a reference, and all pre-synaptic axonal boutons containing a mitochondrion were identified. Each of these boutons was followed throughout the series, and the number, length and morphology of mitochondria were recorded. Mitochondria were classified as either straight or curved. Mitochondria that were bent at least 90° were identified as curved, while all other mitochondria were classified as straight.

Statistics

Data sets from the dot blot assays were analyzed using a one-way ANOVA followed by Tukey's multiple comparisons test. Data sets from the A11 immunohistochemistry and CP-2 cyclic peptide-FITC (CP-2-FITC) (*22*) fluorescence microscopy were analyzed using a two-way ANOVA followed by Sidak's multiple comparisons test. Data sets were analyzed using a two-tailed Student's t-test. Threshold for significance was set at $\alpha = 0.05$. Distribution of mGluR-2/3 immunogold particles was analyzed using repeated measures ANOVA. Data are

represented as mean ± SD. Statistical analyses were carried out using Prism software (version 9.2.0, GraphPad

Software, San Diego, CA).

Data Availability

The scRNA-seq data (cell ranger processed count matrices) are in a Synapse repository (accession

syn52752404, with direct link https://www.synapse.org/#!Synapse:syn52752404/files/).

References

1. S. W. Scheff, D. A. Price, F. A. Schmitt, S. T. DeKosky, E. J. Mufson. Synaptic alterations in CA1 in mild Alzheimer
disease and mild cognitive impairment. *Neurology* **68**, 1501-1508 (2007).
2. T. Arendt, Synaptic degeneration in Alzheimer's disease. *Acta Neuropathol* **118**, 167-179 (2009).
3. G. M. Shankar, D. M. Walsh. Alzheimer's disease: synaptic dysfunction and Abeta. *Mol Neurodegener* **4**, 48 (2009).
4. A. John, P. H. Reddy. Synaptic basis of Alzheimer's disease: Focus on synaptic amyloid beta, P-tau and
mitochondria. *Ageing Res Rev* **65**, 101208 (2021).
5. M. Pang, L. Zhu, A. Gabelle, A. R. Gafson, R. W. Platt, J. E. Galvin, P. Krolak-Salmon, I. Rubino, C. de Moor, S.
Belachew, C. Shen. Effect of reduction in brain amyloid levels on change in cognitive and functional decline in
randomized clinical trials: An instrumental variable meta-analysis. *Alzheimers Dement*, (2022).
6. T. Bussiere, P. D. Friend, N. Sadeghi, B. Wicinski, G. I. Lin, C. Bouras, P. Giannakopoulos, N. K. Robakis, J. H.
Morrison, D. P. Perl, P. R. Hof. Stereologic assessment of the total cortical volume occupied by amyloid deposits
and its relationship with cognitive status in aging and Alzheimer's disease. *Neuroscience* **112**, 75-91 (2002).
7. A. Akram, D. Christoffel, A. B. Rocher, C. Bouras, E. Kovari, D. P. Perl, J. H. Morrison, F. R. Herrmann, V.
Haroutunian, P. Giannakopoulos, P. R. Hof. Stereologic estimates of total spinophilin-immunoreactive spine number
in area 9 and the CA1 field: relationship with the progression of Alzheimer's disease. *Neurobiol Aging* **29**, 1296-
1307 (2008).
8. P. T. Nelson, I. Alafuzoff, E. H. Bigio, C. Bouras, H. Braak, N. J. Cairns, R. J. Castellani, B. J. Crain, P. Davies,
K. Del Tredici, C. Duyckaerts, M. P. Frosch, V. Haroutunian, P. R. Hof, C. M. Hulette, B. T. Hyman, T. Iwatsubo,
K. A. Jellinger, G. A. Jicha, E. Kövari, W. A. Kukull, J. B. Leverenz, S. Love, I. R. Mackenzie, D. M. Mann, E.
Masliah, A. C. McKee, T. J. Montine, J. C. Morris, J. A. Schneider, J. A. Sonnen, D. R. Thal, J. Q. Trojanowski, J.
C. Troncoso, T. Wisniewski, R. L. Woltjer, T. G. Beach. Correlation of Alzheimer disease neuropathologic changes
with cognitive status: a review of the literature. *J Neuropathol Exp Neurol* **71**, 362-381 (2012).
9. K. Ishibashi, T. Tomiyama, K. Nishitsuji, M. Hara, H. Mori. Absence of synaptophysin near cortical neurons
containing oligomeric Abeta in Alzheimer's disease brain. *J Neurosci Res* **84**, 632-636 (2006).
10. M. P. Lambert, A. K. Barlow, B. A. Chromy, C. Edwards, R. Freed, M. Liosatos, T. E. Morgan, I. Rozovsky, B.
Trommer, K. L. Viola, P. Wals, C. Zhang, C. E. Finch, G. A. Krafft, W. L. Klein. Diffusible, nonfibrillar ligands
derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci U S A* **95**, 6448-6453
(1998).
11. J. A. Fein, S. Sokolow, C. A. Miller, H. V. Vinters, F. Yang, G. M. Cole, K. H. Gylys. Co-localization of amyloid
beta and tau pathology in Alzheimer's disease synaptosomes. *Am J Pathol* **172**, 1683-1692 (2008).
12. S. Sokolow, K. M. Henkins, T. Bilousova, C. A. Miller, H. V. Vinters, W. Poon, G. M. Cole, K. H. Gylys. AD
synapses contain abundant Abeta monomer and multiple soluble oligomers, including a 56-kDa assembly.
Neurobiol Aging **33**, 1545-1555 (2012).
13. T. Bilousova, C. A. Miller, W. W. Poon, H. V. Vinters, M. Corrada, C. Kawas, E. Y. Hayden, D. B. Teplow, C. Glabe,
R. Albay, 3rd, G. M. Cole, E. Teng, K. H. Gylys. Synaptic Amyloid-beta Oligomers Precede p-Tau and Differentiate
High Pathology Control Cases. *Am J Pathol* **186**, 185-198 (2016).
14. D. M. Walsh, I. Klyubin, J. V. Fadeeva, W. K. Cullen, R. Anwyl, M. S. Wolfe, M. J. Rowan, D. J. Selkoe. Naturally
secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo. *Nature* **416**,
535-539 (2002).
15. G. M. Shankar, B. L. Bloodgood, M. Townsend, D. M. Walsh, D. J. Selkoe, B. L. Sabatini. Natural oligomers of the
Alzheimer amyloid-beta protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-
dependent signaling pathway. *J Neurosci* **27**, 2866-2875 (2007).

16. J. P. Cleary, D. M. Walsh, J. J. Hofmeister, G. M. Shankar, M. A. Kuskowski, D. J. Selkoe, K. H. Ashe. Natural
oligomers of the amyloid-beta protein specifically disrupt cognitive function. *Nat Neurosci* **8**, 79-84 (2005).
17. B. DaRocha-Souto, T. C. Scotton, M. Coma, A. Serrano-Pozo, T. Hashimoto, L. Sereno, M. Rodriguez, B. Sanchez,
B. T. Hyman, T. Gomez-Isla. Brain oligomeric beta-amyloid but not total amyloid plaque burden correlates with
neuronal loss and astrocyte inflammatory response in amyloid precursor protein/tau transgenic mice. *J Neuropathol*
Exp Neurol **70**, 360-376 (2011).
18. S. Gandy, A. J. Simon, J. W. Steele, A. L. Lublin, J. J. Lah, L. C. Walker, A. I. Levey, G. A. Krafft, E. Levy, F.
Checler, C. Glabe, W. B. Bilker, T. Abel, J. Schmeidler, M. E. Ehrlich. Days to criterion as an indicator of toxicity
associated with human Alzheimer amyloid-beta oligomers. *Ann Neurol* **68**, 220-230 (2010).
19. E. Levy, M. D. Carman, I. J. Fernandez-Madrid, M. D. Power, I. Lieberburg, S. G. van Duinen, G. T. Bots, W.
Luyendijk, B. Frangione. Mutation of the Alzheimer's disease amyloid gene in hereditary cerebral hemorrhage,
Dutch type. *Science* **248**, 1124-1126 (1990).
20. M. L. Maat-Schieman, S. G. van Duinen, J. Haan, R. A. Roos. Morphology of cerebral plaque-like lesions in
hereditary cerebral hemorrhage with amyloidosis (Dutch). *Acta Neuropathol* **84**, 674-679 (1992).
21. E. M. Knight, S. H. Kim, J. C. Kottwitz, A. Hatami, R. Albay, A. Suzuki, A. Lublin, C. M. Alberini, W. L. Klein, P.
Szabo, N. R. Relkin, M. Ehrlich, C. G. Glabe, S. Gandy, J. W. Steele. Effective anti-Alzheimer Abeta therapy
involves depletion of specific Abeta oligomer subtypes. *Neurol Neuroimmunol Neuroinflamm* **3**, e237 (2016).
22. M. Habashi, S. Vutla, K. Tripathi, S. Senapati, P. S. Chauhan, A. Haviv-Chesner, M. Richman, S.-A. Mohand, V.
Dumulon-Perreault, R. Mulamreddy, E. Okun, J. H. Chill, B. Guérin, W. D. Lubell, S. Rahimipour. Early diagnosis
and treatment of Alzheimer's disease by targeting toxic soluble Aβ oligomers. *Proceedings of the National Academy*
of Sciences **119**, e2210766119 (2022).
23. G. Kaur, M. Pawlik, S. E. Gandy, M. E. Ehrlich, J. F. Smiley, E. Levy. Lysosomal dysfunction in the brain of a
mouse model with intraneuronal accumulation of carboxyl terminal fragments of the amyloid precursor protein.
Mol Psychiatry **22**, 981-989 (2017).
24. K. A. Price, M. Varghese, A. Sowa, F. Yuk, H. Brautigam, M. E. Ehrlich, D. L. Dickstein,.Altered synaptic structure
in the hippocampus in a mouse model of Alzheimer's disease with soluble amyloid-beta oligomers and no plaque
pathology. *Mol Neurodegener* **9**, 41 (2014).
25. B. Readhead, J. V. Haure-Mirande, B. Zhang, V. Haroutunian, S. Gandy, E. E. Schadt, J. T. Dudley, M. E. Ehrlich.
Molecular systems evaluation of oligomerogenic APP(E693Q) and fibrillogenic APP(KM670/
671NL)/PSEN1(Deltaexon9) mouse models identifies shared features with human Alzheimer's brain molecular
pathology. *Mol Psychiatry* **21**, 1153-1154 (2016).
26. S. Tucker, C. Möller, K. Tegerstedt, A. Lord, H. Laudon, J. Sjödahl, L. Söderberg, E. Spens, C. Sahlin, E. R. Waara,
A. Satlin, P. Gellerfors, G. Osswald, L. Lannfelt. The murine version of BAN2401 (mAb158) selectively reduces
amyloid-β protofibrils in brain and cerebrospinal fluid of tg-ArcSwe mice. *J Alzheimers Dis* **43**, 575-588 (2015).
27. R. S. Zucker, W. G. J. A. r. o. p. Regehr. Short-term synaptic plasticity. **64**, 355-405 (2002).
28. H. Zhang, B. Gong, S. Liu, M. Fa, I. Ninan, A. Staniszewski, O. J. C. A. R. Arancio. Synaptic fatigue is more
pronounced in the APP/PS1 transgenic mouse model of Alzheimer's disease. **2**, 137-140 (2005).
29. G. Yu, L.-G. Wang, Y. Han, Q.-Y. He, clusterProfiler: an R Package for Comparing Biological Themes Among Gene
Clusters. *OMICS: A Journal of Integrative Biology* **16**, 284-287 (2012).
30. M. Ximerakis, S. L. Lipnick, B. T. Innes, S. K. Simmons, X. Adiconis, D. Dionne, B. A. Mayweather, L. Nguyen,
Z. Niziolek, C. Ozek, V. L. Butty, R. Isserlin, S. M. Buchanan, S. S. Levine, A. Regev, G. D. Bader, J. Z. Levin, L.
Rubin. Single-cell transcriptomic profiling of the aging mouse brain. *Nature Neuroscience* **22**, 1696-1708 (2019).
31. S. Krasemann, C. Madore, R. Cialic, C. Baufeld, N. Calcagno, R. El Fatimy, L. Beckers, E. O'Loughlin, Y. Xu,
Z. Fanek, D. J. Greco, S. T. Smith, G. Tweet, Z. Humulock, T. Zrzavy, P. Conde-Sanroman, M. Gacias, Z. Weng, H.
Chen, E. Tjon, F. Mazaheri, K. Hartmann, A. Madi, J. D. Ulrich, M. Glatzel, A. Worthmann, J. Heeren, B. Budnik,
C. Lemere, T. Ikezu, F. L. Heppner, V. Litvak, D. M. Holtzman, H. Lassmann, H. L. Weiner, J. Ochando, C. Haass,
O. Butovsky. The TREM2-APOE Pathway Drives the Transcriptional Phenotype of Dysfunctional Microglia in
Neurodegenerative Diseases. *Immunity* **47**, 566-581.e569 (2017).
32. G. D. Manocha, A. M. Floden, K. Rausch, J. A. Kulas, B. A. McGregor, L. Rojanathammanee, K. R. Puig, K. L.
Puig, S. Karki, M. R. Nichols, D. C. Darland, J. E. Porter, C. K. Combs. APP Regulates Microglial Phenotype in a
Mouse Model of Alzheimer's Disease. *J Neurosci* **36**, 8471-8486 (2016).

33.   H. Keren-Shaul, A. Spinrad, A. Weiner, O. Matcovitch-Natan, R. Dvir-Szternfeld, T. K. Ulland, E. David, K.
Baruch, D. Lara-Astaiso, B. Toth, S. Itzkovitz, M. Colonna, M. Schwartz, I. Amit. A Unique Microglia Type
Associated with Restricting Development of Alzheimer's Disease. *Cell* **169**, 1276-1290.e1217 (2017).
34.   Y. Chen, M. Colonna. Microglia in Alzheimer's disease at single-cell level. Are there common patterns in humans
and mice? *J Exp Med* **218**, (2021).
35.   S. Hong, V. F. Beja-Glasser, B. M. Nfonoyim, A. Frouin, S. Li, S. Ramakrishnan, K. M. Merry, Q. Shi, A. Rosenthal,
B. A. Barres, C. A. Lemere, D. J. Selkoe, B. Stevens. Complement and microglia mediate early synapse loss in
Alzheimer mouse models. *Science* **352**, 712-716 (2016).
36.   P. Sunderaraman, S. Cosentino, N. Schupf, J. Manly, Y. Gu, S. Barral. MEF2C Common Genetic Variation Is
Associated With Different Aspects of Cognition in Non-Hispanic White and Caribbean Hispanic Non-demented
Older Adults. *Front Genet* **12**, 642327 (2021).
37.   J. C. Lambert, C. A. Ibrahim-Verbaas, D. Harold, A. C. Naj, R. Sims, C. Bellenguez, A. L. DeStafano, J. C. Bis, G.
W. Beecham, B. Grenier-Boley, G. Russo, T. A. Thorton-Wells, N. Jones, A. V. Smith, V. Chouraki, C. Thomas, M.
A. Ikram, D. Zelenika, B. N. Vardarajan, Y. Kamatani, C. F. Lin, A. Gerrish, H. Schmidt, B. Kunkle, M. L.
Dunstan, A. Ruiz, M. T. Bihoreau, S. H. Choi, C. Reitz, F. Pasquier, C. Cruchaga, D. Craig, N. Amin, C. Berr, O.
L. Lopez, P. L. De Jager, V. Deramecourt, J. A. Johnston, D. Evans, S. Lovestone, L. Letenneur, F. J. Morón, D. C.
Rubinsztein, G. Eiriksdottir, K. Sleegers, A. M. Goate, N. Fiévet, M. W. Huentelman, M. Gill, K. Brown, M. I.
Kamboh, L. Keller, P. Barberger-Gateau, B. McGuiness, E. B. Larson, R. Green, A. J. Myers, C. Dufouil, S. Todd,
D. Wallon, S. Love, E. Rogaeva, J. Gallacher, P. St George-Hyslop, J. Clarimon, A. Lleo, A. Bayer, D. W. Tsuang,
L. Yu, M. Tsolaki, P. Bossù, G. Spalletta, P. Proitsi, J. Collinge, S. Sorbi, F. Sanchez-Garcia, N. C. Fox, J. Hardy,
M. C. Deniz Naranjo, P. Bosco, R. Clarke, C. Brayne, D. Galimberti, M. Mancuso, F. Matthews, S. Moebus, P.
Mecocci, M. Del Zompo, W. Maier, H. Hampel, A. Pilotto, M. Bullido, F. Panza, P. Caffarra, B. Nacmias, J. R.
Gilbert, M. Mayhaus, L. Lannefelt, H. Hakonarson, S. Pichler, M. M. Carrasquillo, M. Ingelsson, D. Beekly, V.
Alvarez, F. Zou, O. Valladares, S. G. Younkin, E. Coto, K. L. Hamilton-Nelson, W. Gu, C. Razquin, P. Pastor, I.
Mateo, M. J. Owen, K. M. Faber, P. V. Jonsson, O. Combarros, M. C. O'Donovan, L. B. Cantwell, H. Soininen, D.
Blacker, S. Mead, T. H. Mosley, Jr., D. A. Bennett, T. B. Harris, L. Fratiglioni, C. Holmes, R. F. de Bruijn, P.
Passmore, T. J. Montine, K. Bettens, J. I. Rotter, A. Brice, K. Morgan, T. M. Foroud, W. A. Kukull, D. Hannequin,
J. F. Powell, M. A. Nalls, K. Ritchie, K. L. Lunetta, J. S. Kauwe, E. Boerwinkle, M. Riemenschneider, M. Boada,
M. Hiltuenen, E. R. Martin, R. Schmidt, D. Rujescu, L. S. Wang, J. F. Dartigues, R. Mayeux, C. Tzourio, A.
Hofman, M. M. Nöthen, C. Graff, B. M. Psaty, L. Jones, J. L. Haines, P. A. Holmans, M. Lathrop, M. A.
PericakVance, L. J. Launer, L. A. Farrer, C. M. van Duijn, C. Van Broeckhoven, V. Moskvina, S. Seshadri, J.
Williams, G. D. Schellenberg, P. Amouyel. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci
for Alzheimer's disease. *Nat Genet* **45**, 1452-1458 (2013).
38.   M. Jang, E. Gould, J. Xu, E. J. Kim, J. H. Kim. Oligodendrocytes regulate presynaptic properties and
neurotransmission through BDNF signaling in the mouse brainstem. *Elife* **8**, (2019).
39.   C. Habermacher, M. C. Angulo, N. Benamer. Glutamate versus GABA in neuron-oligodendroglia communication.
Glia **67**, 2092-2106 (2019).
40.   K. Hoshi, H. Ito, E. Abe, T. J. Fuwa, M. Kanno, Y. Murakami, M. Abe, T. Murakami, A. Yoshihara, Y. Ugawa, T.
Saito, T. C. Saido, K. Matsumoto, Y. Yamaguchi, K. Furukawa, H. Arai, M. Kanai, M. Miyajima, H. Arai, N. Ogawa,
H. Akatsu, Y. Hashizume, H. Tateno, T. Honda, Y. Hashimoto. Transferrin Biosynthesized in the Brain Is a Novel
Biomarker for Alzheimer's Disease. *Metabolites* **11**, (2021).
41.   N. Lemon, E. Canepa, M. A. Ilies, S. Fossati. Carbonic Anhydrases as Potential Targets Against Neurovascular Unit
Dysfunction in Alzheimer's Disease and Stroke. *Front Aging Neurosci* **13**, 772278 (2021).
42.   X. Wang, B. E. Figueroa, I. G. Stavrovskaya, Y. Zhang, A. C. Sirianni, S. Zhu, A. L. Day, B. S. Kristal, R. M.
Friedlander. Methazolamide and melatonin inhibit mitochondrial cytochrome C release and are neuroprotective in
experimental models of ischemic injury. *Stroke* **40**, 1877-1885 (2009).
43.   A. Lee, C. Kondapalli, D. M. Virga, T. L. Lewis, Jr., S. Y. Koo, A. Ashok, G. Mairet-Coello, S. Herzig, M. Foretz,
B. Viollet, R. Shaw, A. Sproul, F. Polleux. Aβ42 oligomers trigger synaptic loss through CAMKK2-
AMPKdependent effectors coordinating mitochondrial fission and mitophagy. *Nat Commun* **13**, 4444 (2022).
44.   B. A. Györffy, V. Tóth, G. Török, P. Gulyássy, R. Kovács, H. Vadászi, A. Micsonai, M. E. Tóth, M. Sántha, L.
Homolya, L. Drahos, G. Juhász, K. A. Kékesi, J. Kardos. Synaptic mitochondrial dysfunction and septin

accumulation are linked to complement-mediated synapse loss in an Alzheimer's disease animal model. *Cell Mol*
Life Sci **77**, 5243-5258 (2020).

45.   A. Norambuena, H. Wallrabe, R. Cao, D. B. Wang, A. Silva, Z. Svindrych, A. Periasamy, S. Hu, R. E. Tanzi, D. Y.
Kim, G. S. Bloom, A novel lysosome-to-mitochondria signaling pathway disrupted by amyloid-β oligomers. *The*
EMBO Journal **37**, e100241 (2018).

46.   M. J. Calkins, M. Manczak, P. Mao, U. Shirendeb, P. H. Reddy. Impaired mitochondrial biogenesis, defective axonal
transport of mitochondria, abnormal mitochondrial dynamics and synaptic degeneration in a mouse model of
Alzheimer's disease. *Hum Mol Genet* **20**, 4515-4529 (2011).

47.   J. S. da Silva, Y. Nonose, F. Rohden, P. C. Lukasewicz Ferreira, F. U. Fontella, A. Rocha, A. W. Brochier, R. V.
Apel, T. M. de Lima, B. Seminotti, A. U. Amaral, A. Galina, D. O. Souza. Guanosine Neuroprotection of Presynaptic
Mitochondrial Calcium Homeostasis in a Mouse Study with Amyloid-beta oligomers. *Mol Neurobiol* **57**, 4790-
4809 (2020).

48.   I. Klyubin, D. M. Walsh, W. K. Cullen, J. V. Fadeeva, R. Anwyl, D. J. Selkoe, M. J. Rowan. Soluble Arctic amyloid
beta protein inhibits hippocampal long-term potentiation in vivo. *Eur J Neurosci* **19**, 2839-2846 (2004).

49.   R. Janz, T. C. Südhof, R. E. Hammer, V. Unni, S. A. Siegelbaum, V. Y. J. N. Bolshakov. Essential roles in synaptic
plasticity for synaptogyrin I and synaptophysin I. **24**, 687-700 (1999).

50.   W. Q. Zhao, F. Santini, R. Breese, D. Ross, X. D. Zhang, D. J. Stone, M. Ferrer, M. Townsend, A. L. Wolfe, M. A.
Seager, G. G. Kinney, P. J. Shughrue, W. J. Ray. Inhibition of calcineurin-mediated endocytosis and alpha-amino3-
hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors prevents amyloid beta oligomer-induced synaptic
disruption. *J Biol Chem* **285**, 7619-7632 (2010).

51.   N. R. Reinders, Y. Pao, M. C. Renner, C. M. da Silva-Matos, T. R. Lodder, R. Malinow, H. W. Kessels. Amyloidbeta
effects on synapses and memory require AMPA receptor subunit GluA3. *Proc Natl Acad Sci U S A* **113**, E6526E6534
(2016).

52.   M. D. Ikonomovic, R. Sheffield, D. M. Armstrong. AMPA-selective glutamate receptor subtype immunoreactivity
in the hippocampal formation of patients with Alzheimer's disease. *Hippocampus* **5**, 469-486 (1995).

53.   T. L. Carter, R. A. Rissman, A. J. Mishizen-Eberz, B. B. Wolfe, R. L. Hamilton, S. Gandy, D. M. Armstrong.
Differential preservation of AMPA receptor subunits in the hippocampi of Alzheimer's disease patients according
to Braak stage. *Exp Neurol* **187**, 299-309 (2004).

54.   J. H. Y. Yeung, J. L. Walby, T. H. Palpagama, C. Turner, H. J. Waldvogel, R. L. M. Faull, A. Kwakowsky.
Glutamatergic receptor expression changes in the Alzheimer's disease hippocampus and entorhinal cortex. *Brain*
Pathol **31**, e13005 (2021).

55.   B. G. Jang, S. In, B. Choi, M. J. Kim. Beta-amyloid oligomers induce early loss of presynaptic proteins in primary
neurons by caspase-dependent and proteasome-dependent mechanisms. *Neuroreport* **25**, 1281-1288 (2014).

56.   M. C. Herzig, D. T. Winkler, P. Burgermeister, M. Pfeifer, E. Kohler, S. D. Schmidt, S. Danner, D. Abramowski, C.
Sturchler-Pierrat, K. Burki, S. G. van Duinen, M. L. Maat-Schieman, M. Staufenbiel, P. M. Mathews, M. Jucker.
Abeta is targeted to the vasculature in a mouse model of hereditary cerebral hemorrhage with amyloidosis. *Nat*
Neurosci **7**, 954-960 (2004).

57.   L. Grand Moursel, W. M. C. van Roon-Mom, S. M. Kiełbasa, H. Mei, H. P. J. Buermans, L. M. van der Graaf, K.
M. Hettne, E. J. de Meijer, S. G. van Duinen, J. F. J. Laros, M. A. van Buchem, P. A. C. t Hoen, S. M. van der
Maarel, L. van der Weerd. Brain Transcriptomic Analysis of Hereditary Cerebral Hemorrhage With Amyloidosis
Dutch Type. *Front Aging Neurosci* **10**, 102 (2018).

58.   R. A. Neff, M. Wang, S. Vatansever, L. Guo, C. Ming, Q. Wang, E. Wang, E. Horgusluoglu-Moloch, W.-m. Song,
A. Li, E. L. Castranio, J. Tcw, L. Ho, A. Goate, V. Fossati, S. Noggle, S. Gandy, M. E. Ehrlich, P. Katsel, E. Schadt,
D. Cai, K. J. Brennand, V. Haroutunian, B. Zhang. Molecular subtyping of Alzheimer's disease using RNA
sequencing data reveals novel mechanisms and targets. *Science Advances* **7**, eabb5398 (2021).

59.   A. Jishi, X. Qi, H. C. Miranda. Implications of mRNA translation dysregulation for neurological disorders. *Semin*
Cell Dev Biol **114**, 11-19 (2021).

60.   A. Ghosh, K. Mizuno, S. S. Tiwari, P. Proitsi, B. Gomez Perez-Nievas, E. Glennon, R. T. Martinez-Nunez, K. P.
Giese. Alzheimer's disease-related dysregulation of mRNA translation causes key pathological features with ageing.
Translational Psychiatry **10**, 192 (2020).

61. G. Joshi, Y. Chi, Z. Huang, Y. Wang. Aβ-induced Golgi fragmentation in Alzheimer's disease enhances Aβ
production. *Proc Natl Acad Sci U S A* **111**, E1230-1239 (2014).

62. M. Calvo-Rodriguez, S. S. Hou, A. C. Snyder, E. K. Kharitonova, A. N. Russ, S. Das, Z. Fan, A. Muzikansky, M.
Garcia-Alloza, A. Serrano-Pozo, E. Hudry, B. J. Bacskai. Increased mitochondrial calcium levels associated with
neuronal death in a mouse model of Alzheimer's disease. *Nat Commun* **11**, 2146 (2020).

63. A.V. Venkataraman, A. Mansur, G. Rizzo, C. Bishop, Y. Lewis, E. Kocagoncu, A. Lingford-Hughes, M. Huiban, J.
Passchier, J.B. Rowe, H. Tsukada, D. J. Brooks, L. Martarello, R. A. Comley, L. Chen, A.J. Schwarz, R. Hargreaves,
R.N. Gunn, E. A. Rabiner, P. M. Matthew. Widespread cell stress and mitochondrial dysfunction occur in patients
with early Alzheimer's disease. *Sci Transl Med.* **14**(658), eabk1051. doi: 10.1126/scitranslmed.abk1051. (2022).

64. M. Balietti, B. Giorgetti, T. Casoli, M. Solazzi, F. Tamagnini, C. Burattini, G. Aicardi, P. Fattoretti. Early selective
vulnerability of synapses and synaptic mitochondria in the hippocampal CA1 region of the Tg2576 mouse model
of Alzheimer's disease. *J Alzheimers Dis* **34**, 887-896 (2013).

65. X. Wang, B. Su, H. G. Lee, X. Li, G. Perry, M. A. Smith, X. Zhu. Impaired balance of mitochondrial fission and
fusion in Alzheimer's disease. *J Neurosci* **29**, 9090-9103 (2009).

66. C. D. Sanmartin, T. Adasme, C. Hidalgo, A. C. Paula-Lima. The antioxidant N-acetylcysteine prevents the
mitochondrial fragmentation induced by soluble amyloid-beta peptide oligomers. *Neurodegener Dis* **10**, 34-37
(2012).

67. T. Umeda, T. Tomiyama, N. Sakama, S. Tanaka, M. P. Lambert, W. L. Klein, H. Mori. Intraneuronal amyloid beta
oligomers cause cell death via endoplasmic reticulum stress, endosomal/lysosomal leakage, and mitochondrial
dysfunction in vivo. *J Neurosci Res* **89**, 1031-1042 (2011).

68. T. Umeda, E. M. Ramser, M. Yamashita, K. Nakajima, H. Mori, M. A. Silverman, T. Tomiyama. Intracellular
amyloid β oligomers impair organelle transport and induce dendritic spine loss in primary neurons. *Acta
Neuropathol Commun* **3**, 51 (2015).

69. L. Hedskog, C. M. Pinho, R. Filadi, A. Rönnbäck, L. Hertwig, B. Wiehager, P. Larssen, S. Gellhaar, A. Sandebring,
M. Westerlund, C. Graff, B. Winblad, D. Galter, H. Behbahani, P. Pizzo, E. Glaser, M. Ankarcrona. Modulation of
the endoplasmic reticulum–mitochondria interface in Alzheimer's disease and related models. *Proceedings of the
National Academy of Sciences* **110**, 7916-7921 (2013).

70. C. H. van Dyck, C. J. Swanson, P. Aisen, R. J. Bateman, C. Chen, M. Gee, M. Kanekiyo, D. Li, L. Reyderman, S.
Cohen, L. Froelich, S. Katayama, M. Sabbagh, B. Vellas, D. Watson, S. Dhadda, M. Irizarry, L. D. Kramer, T.
Iwatsubo. Lecanemab in Early Alzheimer's Disease. *New England Journal of Medicine*, (2022).

71. J.R. Sims, J.A. Zimmer, C.D. Evans, M. Lu, P. Ardayfio, J. Sparks, A.M. Wessels, S. Shcherbinin, H.
Wang, E.S. Monkul Nery, E.C. Collins, P. Solomon, S. Salloway, L.G. Apostolova, O. Hansson, C. Ritchie,
D.A. Brooks, M. Mintun, D.M. Skoyronsky; TRAILBLAZER-AL2 Investigators. Donanemab in Early
Symptomatic Alzheimer Disease: The TRAILBLAZER-ALZ 2 Randomized Clinical Trial. *JAMA.* **330**,
512-527. Doi: 10.001/jama.2023.13239 (2023).

72. H. Sun, Y. Wang, Y. Wang, F. Zeng. Case of Autosomal Dominant Alzheimer Disease With Negative Findings From
PiB-PET Examination. *Neurol Genet* **10**, e200119 (2024).

73. C. A. Schneider, W. S. Rasband, K. W. Eliceiri. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* **9**,
671-675 (2012).

74. M. Pagnon de la Vega, V. Giedraitis, W. Michno, L. Kilander, G. Güner, M. Zielinski, M. Löwenmark, R. Brundin,
T. Danfors, L. Söderberg, I. Alafuzoff, L. N. G. Nilsson, A. Erlandsson, D. Willbold, S. A. Müller, G. F. Schröder,
J. Hanrieder, S. F. Lichtenthaler, L. Lannfelt, D. Sehlin, M. Ingelsson. The Uppsala APP deletion causes early onset
autosomal dominant Alzheimer's disease by altering APP processing and increasing amyloid β fibril formation. *Sci
Transl Med* **13**, (2021).

75. S. R. Meier, D. Sehlin, G. Hultqvist, S. Syvänen. Pinpointing Brain TREM2 Levels in Two Mouse Models of
Alzheimer's Disease. *Mol Imaging Biol* **23**, 665-675 (2021).

76. J. W. Steele, H. Brautigam, J. A. Short, A. Sowa, M. Shi, A. Yadav, C. M. Weaver, D. Westaway, P. E. Fraser, P. H.
St George-Hyslop, S. Gandy, P. R. Hof, D. L. Dickstein. Early fear memory defects are associated with altered
synaptic plasticity and molecular architecture in the TgCRND8 Alzheimer's disease mouse model. *J Comp Neurol
522, 2319-2335 (2014).

77. M. J. Lazarczyk, J. E. Kemmler, B. A. Eyford, J. A. Short, M. Varghese, A. Sowa, D. R. Dickstein, F. J. Yuk, R. Puri, K. E. Biron, M. Leist, W. A. Jefferies, D. L. Dickstein. Major Histocompatibility Complex class I proteins are critical for maintaining neuronal structural complexity in the aging brain. *Sci Rep* **6**, 26199 (2016).

78. Y. Hara, M. Punsoni, F. Yuk, C. S. Park, W. G. Janssen, P. R. Rapp, J. H. Morrison. Synaptic distributions of GluA2 and PKMzeta in the monkey dentate gyrus and their relationships with aging and memory. *J Neurosci* **32**, 73367344 (2012).

79. Y. Hara, C. S. Park, W. G. Janssen, M. T. Roberts, J. H. Morrison, P. R. Rapp. Synaptic correlates of memory and menopause in the hippocampal dentate gyrus in rhesus monkeys. *Neurobiol Aging* **33**, 421 e417-428 (2012).

80. Y. Han, P. S. Kaeser, T. C. Südhof, R. Schneggenburger. RIM determines Ca²+ channel density and vesicle docking at the presynaptic active zone. *Neuron* **69**, 304-316 (2011).

81. C. Tan, S. S. H. Wang, G. de Nola, P. S. Kaeser. Rebuilding essential active zone functions within a synapse. *Neuron* **110**, 1498-1515.e1498 (2022).

82. P. G. Camoletto, H. Vara, L. Morando, E. Connell, F. P. Marletto, M. Giustetto, M. Sassoè-Pognetto, P. P. Van Veldhoven, M. D. Ledesma. Synaptic vesicle docking: sphingosine regulates syntaxin1 interaction with Munc18. *PLoS One* **4**, e5310 (2009).

83. A. Butler, P. Hoffman, P. Smibert, E. Papalexi, R. Satija. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nature Biotechnology* **36**, 411 (2018).

84. L. McInnes, J. Healy, J. Melville. UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction. *arXiv* **1802.03426**, (2018).

85. O. Franzén, L.-M. Gan, J. L. M. Björkegren. PanglaoDB: a web server for exploration of mouse and human singlecell RNA sequencing data. *Database* **2019**, (2019).

86. A. Butler, P. Hoffman, P. Smibert, E. Papalexi, R. Satija. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nature biotechnology* **36**, 411-420 (2018).

87. R. Satija, J. A. Farrell, D. Gennert, A. F. Schier, A. Regev. Spatial reconstruction of single-cell gene expression data. *Nat Biotechnol* **33**, 495-502 (2015).

88. C. Neftel, J. Laffy, M. G. Filbin, T. Hara, M. E. Shore, G. J. Rahme, A. R. Richman, D. Silverbush, M. L. Shaw, C. M. Hebert, J. Dewitt, S. Gritsch, E. M. Perez, L. N. Gonzalez Castro, X. Lan, N. Druck, C. Rodman, D. Dionne, A. Kaplan, M. S. Bertalan, J. Small, K. Pelton, S. Becker, D. Bonal, Q. D. Nguyen, R. L. Servis, J. M. Fung, R. Mylvaganam, L. Mayr, J. Gojo, C. Haberler, R. Geyeregger, T. Czech, I. Slavc, B. V. Nahed, W. T. Curry, B. S. Carter, H. Wakimoto, P. K. Brastianos, T. T. Batchelor, A. Stemmer-Rachamimov, M. Martinez-Lage, M. P. Frosch, I. Stamenkovic, N. Riggi, E. Rheinbay, M. Monje, O. Rozenblatt-Rosen, D. P. Cahill, A. P. Patel, T. Hunter, I. M. Verma, K. L. Ligon, D. N. Louis, A. Regev, B. E. Bernstein, I. Tirosh, M. L. Suva. An Integrative Model of Cellular States, Plasticity, and Genetics for Glioblastoma. *Cell* **178**, 835-849 e821 (2019).

89. Y. Hara, F. Yuk, R. Puri, W. G. Janssen, P. R. Rapp, J. H. Morrison. Presynaptic mitochondrial morphology in monkey prefrontal cortex correlates with working memory and is improved with estrogen treatment. *Proc Natl Acad Sci U S A* **111**, 486-491 (2014).

90. Y. He, P. R. Hof, W. G. M. Janssen, J. D. Rothstein, J. H. Morrison. Differential synaptic localization of GluR2 and EAAC1 in the macaque monkey entorhinal cortex: a postembedding immunogold study. *Neuroscience Letters* **311**, 161-164 (2001).

91. Y. Zhou, B. Zhou, L. Pache, M. Chang, A. H. Khodabakhshi, O. Tanaseichuk, C. Benner, S. K. Chanda. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun* **10**, 1523 (2019).

92. K. D. Micheva, N. O'Rourke, B. Busse, S. J. Smith. Array tomography: high-resolution three-dimensional immunofluorescence. *Cold Spring Harb Protoc* **2010**, pdb top89 (2010).

93. K. D. Micheva, N. O'Rourke, B. Busse, S. J. Smith. Array tomography: imaging-stained arrays. *Cold Spring Harb Protoc* **2010**, pdb prot5526 (2010).

94. J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D. J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, A. Cardona. Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**, 676-682 (2012).

95. K. R. Kay, C. Smith, A. K. Wright, A. Serrano-Pozo, A. M. Pooler, R. Koffie, M. E. Bastin, T. H. Bak, S. Abrahams, K. J. Kopeikina, D. McGuone, M. P. Frosch, T. H. Gillingwater, B. T. Hyman, T. L. Spires-Jones. Studying synapses in human brain with array tomography and electron microscopy. *Nat Protoc* **8**, 1366-1380 (2013).

984   96.   A. Cardona, S. Saalfeld, J. Schindelin, I. Arganda-Carreras, S. Preibisch, M. Longair, P. Tomancak, V. Hartenstein,
985   R. J. Douglas. TrakEM2 software for neural circuit reconstruction. *PLoS One* **7**, e38011 (2012).
986   97.   R. Kayed, E. Head, J. L. Thompson, T. M. McIntire, S. C. Milton, C. W. Cotman, C. G. Glabe, Common
987   structure of soluble amyloid oligomers implies common mechanism of pathogenesis. *Science* **300**, 486-489
988   (2003).
989   98.   R. Kayed, E. Head, F. Sarsoza, T. Saing, C.W. Cotman, M. Necula, L. Margol, J. Wu, L. Breydo, J.L. Thompson,
990   S. Rasool, T. Gurlo, P. Butler, C.G. Glabe. Fibril specific, conformation dependent antibodies recognize a generic
991   epitope common to amyloid fibrils and fibrillar oligomers that is absent in prefibrillar oligomers. *Mol*
992   *Neurodegener.* **2**, 18. doi: 10.1186/1750-1326-2-18 (2007).
993   99.   M. Richman, S. Wilk, M. Chemerovski, S.K. Wärmländer, A. Wahlström, A. Gräslund, S. Rahimipour. *In vitro*
994   and mechanistic studies of an antiamyloidogenic self- assembled cyclic D, L-α-peptide architecture. *J Am Chem*
995   *Soc.* **135**, 3474-84 (2016).
996   100.   M. Chemerovski-Glikman, M. Richman, S. Rahimipour. Structure-based study of antiamyloidogenic cyclic D,L-α-
997   peptides. *Tetrahedron* **70**, 7639–7644 (2014).
998   101.   M. Chemerovski-Glikman, E. Rozentur-Shkop, M. Richman, A. Grupi, A. Getler, H. Y. Cohen, H. Shaked, C.
999   Wallin, S. K. Wärmländer, E. Haas, A. Gräslund, J. H. Chill, S. Rahimipour. Self-assembled cyclic D,L-alpha-
000   peptides as generic conformational inhibitors of the alpha-synuclein aggregation and toxicity: In vitro and
001   mechanistic studies. *Chemistry*, **22**, 14236-14246 (2016).
002   102.   A. Belostozky, M. Richman, E. Lisniansky, A. Tovchygrechko , J. Chill, S. Rahimipour. Inhibition of tau-derived
003   hexapeptide aggregation and toxicity by a self-assembled cyclic d,l-α-peptide conformational inhibitor. *Chem*
004   *Commun (Camb).* **54**, 5980-5983. doi: 10.1039/c8cc01233d (2018).
005   103.   K. Ono, M.M. Condron, D. B. Teplow. Structure-neurotoxicity relationships of amyloid beta-protein oligomers.
006   *Proc. Natl. Acad. Sci. U. S. A.* **106,** 14745-14750 (2009).
007   104.   R.J. Rodenburg. Biochemical diagnosis of mitochondrial disorders. *J Inherit Metab Dis* **34**,283-92 (2011).
008   105.   T.L. Emmerzaal, R.J. Rodenburg, H. Tanila, V. Verweij, A.J. Kiliaan, T. Kozicz T. Age-Dependent Decrease of
009   Mitochondrial Complex II Activity in a Familial Mouse Model for Alzheimer's Disease. *J Alzheimers Dis.* **66**, 75-
010   82. doi: 10.3233/JAD-180337 (2018).
011   106.   M. Noguchi-Shinohara, K. Shuta, H. Murakami, Y. Mori, J. Komatsu, C. Kobayashi, S. Hersch, K. Horie, K. Ono.
012   Lecanemab-Associated Amyloid-β Protofibril in Cerebrospinal Fluid Correlates with Biomarkers of
013   Neurodegeneration in Alzheimer's Disease. *Ann Neurol.* doi: 10.1002/ana.27175. (2025).
014   107.   D. Shea, E. Colasurdo, A. Smith, C. Paschall, S. Jayadev, C.D. Keene, D. Galasko, A. Ko, G. Li, E. Peskind, V.
015   SOBA: Development and testing of a soluble oligomer binding assay for detection of amyloidogenic toxic
016   oligomers. *Proc Natl Acad Sci U S A.* **119**: e2213157119. doi: 10.1073/pnas.2213157119 (2022).
017   108.   Choi, E.H., Kim, M.H., Park, S.J. Targeting mitochondrial dysfunction and reactive oxygen species for
018   neurodegenerative disease treatment. *International journal of molecular sciences*, 25(14), p.7952. 109 (2024).
019   109.   Xu, J., Du, W., Zhao, Y., Lim, K., Lu, L., Zhang, C., Li, L. Mitochondria targeting drugs for neurodegenerative
020   diseases—Design, mechanism and application. *Acta Pharmaceutica Sinica B*, 12(6), pp.2778-2789, (2022).
021   110.   E. Andersson, N. Lindblom, S. Janelidze, G. Salvadó, E. Gkanatsiou, L. Söderberg, C. Möller, L. Lannfelt, J. Ge,
022   J. Hanrieder, K. Blennow, T. Deierborg, N. Mattsson-Carlgren, H. Zetterberg, G. Gouras, O. Hansson. Soluble
023   cerebral Aβ protofibrils link Aβ plaque pathology to changes in CSF $A\beta_{42}/A\beta_{40}$ ratios, neurofilament light and tau
024   in Alzheimer's disease model mice. *Nat Aging*. doi: 10.1038/s43587-025-00810-8. (2025).

Acknowledgments

We would like to thank all the members of the Dickstein, Arancio, Levy, Rahimipour, Lubell, Guerin, and Ehrlich/Gandy laboratories, and William G. M. Janssen, Allison Sowa, Daniel R. Dickstein, Frank Yuk, and Michael Ragusa for technical assistance and discussion. The authors would like to thank Yanzhuang Wang (Shenzhen Bay Laboratory), and Gary Gibson, Giovanni Manfredi, and Csaba Konrad (all from Weill Cornell Medical College) for valuable conversations and suggestions. We also thank Tamas Kozicz and Graeme Preston (Icahn School of Medicine at Mount Sinai) for performing the electron transport chain activity assays.

Funding:

National Institutes of Health grant F32AG077900 (ELC)

National Institutes of Health grant P30 AG066514 to Mary Sano with Developmental Pilot Award (ELC, SG)

National Institutes of Health grant U01AG046170 (SG, MEE, MW, BZ)

National Institutes of Health grant RF1AG058469 (SG, MEE)

National Institutes of Health grant RF1AG059319 (SG, MEE)

National Institutes of Health grant R01AG061894 (SG, MEE)

Cure Alzheimer's Fund (SG, MEE, SR, BG, WL)

National Institutes of Health grant RF1AG057440 (MW, BZ)

National Institutes of Health grant R56AG056507 (CG)

National Institutes of Health grant AG056732 (EL)

National Institutes of Health grant R01NS110024 (OA)

Alzheimer's Association NIRG-12-242386 (DLD)

Medical Student Training in Aging Research Program Scientific Fellowship from the American Federation for Aging Research (EB)

The Linda Wachner Foundation (SG, MEE)

National Institutes of Health grant RF1AG077828 (MW)

National Institutes of Health grant R21AG077168 (MW)

Alzheimer's Association AARG-22-928419 (MW)

Author contributions:

Conceptualization: MEE, SG, EL, DLD, OA

Methodology: EB, MV, DLD, KT, SH, JVDL, OA, EA, ELC, CDS, MW, BZ, CG, SR

Investigation: RT, MV, DL, EB, DLD, FG, ELC, KT, SH, JVDL, OA, EA, HZ, BZ, MW, CDS, LS, LL, MI, SR

Visualization: MV, DLD, KT, SH, ELC, MW

Supervision: MEE, SG, EL, DLD, OA

Writing—original draft: MV, DLD, EB, ELC, SG, EA, OA, MW

Writing—review & editing: MV, DLD, ELC, SG, MEE, EL, CG, SR, WL, BG

Competing interests: Dr. Gandy is a co-founder of Recuerdo Pharmaceuticals. He has served as a consultant in the past for J&J, Diagenic, and Pfizer, and he currently consults for Cognito Therapeutics, GLG Group, SVB Securities, Guidepoint, Third Bridge, MEDACORP, Altpep, Vigil Neurosciences, Memory Garden, and Eisai. He has received research support in the past from Warner-Lambert, Pfizer, Baxter, and Avid. The authors declare no other competing interests.

Data and materials availability: The scRNA-seq data (cell ranger processed count matrices) are in a Synapse repository (accession syn52752404, with direct link https://www.synapse.org/#!Synapse:syn52752404/files/). All data are available in the main text or the supplementary materials." The APP^{E693Q} transgenic mouse can be provided by Thomas Jefferson University pending scientific review and a completed material transfer agreement. Requests for the APP^{E693Q} transgenic mouse should be submitted to michelle.ehrlich@mssm.edu.

073 **Disclaimer**

074 The opinions expressed herein are those of the authors and are not necessarily representative of those of the
075 government of the United States, the Uniformed Services University of the Health Sciences, the Department of
076 Defense (DoD), or the United States Army, Navy or Air Force or the Henry M. Jackson Foundation for the
077 Advancement of Military Medicine, Inc.

1073 **Supplementary Materials Table of Contents**

Figure 1. Aging-related accumulation of NFA-Aβ in the cerebral cortex of APP^{E693Q} transgenic mice revealed by double labeling with A11 epitomic immunohistochemistry and CP-2 cyclic peptide-FITC fluorescence microscopy. Representative images from 24-month-old Dutch mice showing fluorescence microscopy using DAPI, A11 staining with AlexaFluor-680 secondary antibody (red), and CP-2 cyclic peptideFITC (CP-2-FITC, green)(*22*). Lower right panel is a merge of A11 and CP-2-FITC images. Scale bar = 100 μm. Quantification of A11-immunoreactive and CP-2-FITC-immunoreactive NFA-Aβ aggregates in 2- and 24mo Dutch shown in Figure S1C. ** p < 0.01.

Figure 2. Dutch mice do not exhibit impairment in basal synaptic transmission or LTP. (A, B) Basal neurotransmission **(A)** and LTP **(B)** are unimpaired in hippocampal slices from Dutch mice as compared to slices from wildtype littermates at the age of 7 to 7.5 months ($F_{(1,28)}$ =0.0026, p=0.959 and $F_{(1,28)}$ =0.219, p=0.644, respectively). **(C-D)** Slices from Dutch mice do not exhibit impairment of basal neurotransmission **(C)** and LTP **(D)** at the age of 10.5 to 11 months compared to wildtype mice ($F_{(1,18)}$ =0.116, p=0.737 and $F_{(1,18)}$ =0.017, =0.899, respectively).

Figure 3. Distribution and ultrastructure of mGluR-2/3-labeled synapses were comparable in the CA1 of 12-monthold Dutch and wildtype mice. mGluR-2/3-labeled synapses were defined as having ≥1 immunogold particle located in the PSD or synaptic cleft. **(A-C)** No significant differences were observed between Dutch and wildtype mice in percentage of mGluR-2/3-labeled total, perforated, and non-perforated synapses, respectively. **(D-F)** Average PSD area of mGluR-2/3labeled synapses for all synapses, non-perforated synapses, and perforated synapses, respectively. Dutch mice had larger PSD area of mGluR-2/3-labeled non-perforated synapses than wildtype (p= 0.04, by unpaired t test), whereas perforated synapses or total synapses were comparable between the two groups. **(G-I)** No significant difference was observed in density of immuno-gold particles in mGluR-2/3-labeled synapses, calculated as number of immuno-gold particles/ PSD area, for all synapses, non-perforated synapses, and perforated synapses, respectively. Data are mean ± SD from n = 5 wildtype, and n = 4 Dutch mice.

Figure 4. Dutch mice showed impairment of certain measurements of short-term synaptic plasticity. **(A)** Dutch mice showed no impairment in paired-pulse facilitation at the age of 11 to 13 months old ($F_{(1, 220)}$ =0.06364, p=0.8011; 2-way ANOVA). **(B)** Post-tetanic potentiation was significantly reduced in slices from 11- to 13-month-old Dutch mice compared to wildtype ($F_{(1,36)}$ =4.771, p=0.036; 2-way ANOVA for the first 6 points after potentiation). **(C)** Significant differences were observed in synaptic fatigue when comparing slices from Dutch mice and wildtype littermates (k=2.247±0.1622 for wildtype and k=2.782±0.1634 for Dutch, p=0.015, k:

rate of exponential decay). **(D)** The rate of synaptic vesicle replenishment following a 100 Hz stimulation was found to be significantly increased in hippocampal slices from Dutch mice as compared to slices from wildtype littermates (k=7.509±0.7277 for wildtype, k=13.64±1.682 for Dutch, $p<0.0001$, k: rate of exponential growth).

Figure 5. Synaptic distribution of mGluR-2/3 in the CA1 is comparable in 12-month-old wildtype and Dutch mice. (A-C) Examples of mGluR-2/3 immuno-gold staining. Note the distribution of immuno-gold particles within the presynaptic terminal (purple masks) and the postsynaptic dendritic spine (cyan masks). Scale bars = 500 nm. **(D)** Schematic diagram demonstrating the 6 synaptic bins used to categorize the localization of each gold particle (adapted from (*90*)). Bin A, active zone (presynaptic, within 45 nm of the membrane across from the PSD); B, synaptic cleft; C, postsynaptic density (PSD); D, peri-synaptic (Peri-PSD, on the postsynaptic spine head, within 200 nm of the PSD); E, postsynaptic membrane (>200 nm away from the PSD) and F, postsynaptic cytoplasm (cytosol). **(E)** Distribution of immuno-gold labeled mGluR-2/3 among the various regions of the synapse showed no significant differences between wildtype and Dutch mice. Data are mean ± SD from wildtype (n = 5) and Dutch (n = 4) (repeated measures ANOVA).

Figure 6. Transcriptional responses in Dutch mouse brains. (A) scRNA-seq of hippocampal tissues from 12-month-old male and female Dutch and wildtype mice (n=2 samples per sex per genotype with 3 mice pooled per sample). UMAP plot showing 14 distinguishable clusters (0-13). **(B)** Violin plots illustrating expression of marker genes for each cell type. **(C)** UpSet plot highlighting the overlapping and unique genes between the clusters. **(D)** Bar plot showing number of DEGs per cluster by directionality.

Figure 7. Overlapping transcriptomic profiles highlight processes impacted in multiple cell types. (A) Dot plot highlighting shared GO term enrichment across clusters. **(B-C)** Dot plot of genes associated with shared GO terms **(B)** "structural constituent of ribosome" and **(C)** "amyloid-beta binding", split into Dutch and wildtype mice across all clusters. Wildtype mice are in blue, Dutch mice are in purple. **(D)** UpSet plot comparing overlap of cluster DEGs (adj. p-value <0.05) with previously defined aging gene list. **(E)** Heatmap of log2FC of genes shared with aging gene list.

Figure 8. Microglia transcriptomic profiles are similar between Dutch and wildtype mice. (A) UMAP plot showing sub-clustering analysis results for Cluster 2 microglia, identifying 3 subpopulations of microglia with UMAP plot split by genotype to the right. **(B)** UMAP plots showing expression of previously identified microglial signatures; DAM(*33*), homeostatic, IFN and MHC(*34*). **(C)** Violin plots showing expression across subclusters by genotypes of selected homeostatic microglial genes.

Figure 9. Transcriptional profile of Cluster 6 excitatory neurons is altered in Dutch mice. (A) Volcano plot

of genes enriched in Cluster 6 as calculated by pseudobulk differential expression analysis. Red genes are Log2FC

> 0.25 and adj. p-value < 0.05, blue Log2FC < 0.25 and adj. p-value < 0.05, green genes are Log2FC > 0.25 and

adj. p-value > 0.05, and grey genes are Log2FC < 0.25 and adj. p-value > 0.05. **(B-C)** GO term analysis of **(B)**

upregulated and **(C)** downregulated genes enriched in Cluster 6 (adj. p-value < 0.05) using metascape(*91*). **(D)**

Dot plot of DEGs unique to Cluster 6.

Figure 10. Fewer mitochondria were present in presynaptic boutons at excitatory synapses in the CA1 of

12-month-old Dutch mice compared to wildtype. **(A)** Electron micrographs showing representative examples

of presynaptic mitochondria (blue masks). **(B)** Curved presynaptic mitochondria demonstrate a bend of at least

90°. Straight presynaptic mitochondria did not exhibit any curvature. Scale bar = 500 nm. **(C)** Total number, **(D)**

number of straight mitochondria, **(E)** number of curved mitochondria, **(F)** percentage of curved mitochondria,

(G) average diameter, and **(H)** number of layers traversed by each mitochondrion in wildtype and Dutch mice.

Data are mean ± SD from n = 4 wildtype and n = 5 Dutch mice.

 Supplementary Tables Titles and Captions

Supplementary Table 1. Cluster Cell Numbers. Number of cells per cluster per sample.

Supplementary Table 2. Cluster Markers. De novo cluster signatures calculated by comparing cell clusters
using Wilcox rank sum test in Seurat.

Supplementary Table 3. Cluster DEGs. Differentially expressed genes across all 14 clusters using MAST
implemented in Seurat.

Supplementary Table 4. Cluster 6 GO-terms. GO-term analysis of genes enriched in Cluster 6 (l2f>0, padj
<0.05) using metascape.

Supplementary Table 5. Cluster 6 GO-terms. GO-term analysis of genes enriched in Cluster 6 (l2f<0, padj
<0.05) using metascape.

Supplementary Table 6. Cluster 4 GO-terms. GO-term analysis of genes enriched in Cluster 6 (l2f>0, padj
<0.05) using metascape.

Supplementary Table 7. Ligand-Receptor Pairs for Clusters 6 and 4. Cell-cell communication analysis
calculated by Cellphone DB to compute interaction between Clusters 6 and Clusters 9 based on known Ligand-
Receptor pairs. t-test was used to compare interaction strength differences between Dutch and WT mice

Supplementary Table 8. Mitochondrial electron transport chain (mETC) enzymology. mETC complexes
standardized to μg protein.

Supplementary Table 9. Mitochondrial electron transport chain (mETC) enzymology. mETC complexes
standardized to mU/ml citrate synthase.

Supplementary Materials for

Endogenously generated Dutch-type Aβ nonfibrillar aggregates dysregulate presynaptic neurotransmission in the absence of detectable inflammation.

Emilie L. Castranio *et al.*

*Corresponding author. Email: michelle.ehrlich@mssm.edu

This PDF file includes:

Supplemental Text, Methods, and Data

Cyclic peptide interactions with Aβ oligomers (NFA-Aβ)

      The concept that NFA-Aβ of various polypeptide sequences might share structural features was first introduced by Kayed *et al.* (2007) (*97,98*) who showed that oligomers of Aβ, IAPP, lysozyme, α-synuclein, prion protein, insulin, and polyglutamine are equally recognized by antibodies that were generated against Aβ oligomers and fibrils. A11 and OC are two antibodies that can each detect unique oligomer subtypes by binding to generic higher order structural epitopes (*97,98*). Richman *et al.* (2013) (*99*) screened a library of six-residue cyclic D,L-α-peptides and identified short peptides that could accomplish a similar goal. Cyclic D,L-α-peptides comprised of even numbers of alternating D- and L-α-amino acids self-assemble into nanotube structures that replicate the intermolecular hydrogen-bonding patterns of cross β-sheets with features of amyloid proteins such as interstrand distances of 4.8 Å (*99*). At and above stoichiometric concentrations, cyclic peptide CP-2 (*c*-[Lys^1-D-Leu^2-Nle^3-D-Trp^4-His^5-D-Ser^6]) was shown to engage soluble and low molecular weight Aβ species, modulate aggregation, disassemble fibrils, and inhibit cell toxicity by an "off-pathway" mechanism (*99, 100*). CP-2 also bound other aggregation-prone polypeptides including PD-associated α-synuclein (α-syn) and tau-derived hexapeptide AcPHF6 (*101, 102*). Various biochemical methods have been employed to demonstrate either the binding of CPs to oligomers or the catalytic effects of cyclic peptides on oligomer structure. These include photoinduced cross-linking of unmodified proteins (PICUP) combined with SDS/PAGE) (*99*) and dynamic light scattering (DLS) (*22*). In some experiments, an aza derivative of CP-2 (CP-7) was prepared, and DLS was used to determine its effect on the size of Aβ40 NFA-Aβ. According to different microscopy techniques including atomic force microscopy and electron microscopy, the size of the NFA-Aβ is around 10 nm (*103*).

Array tomography

      Array tomography (AT) is a high-resolution imaging technique that permits immunofluorescence imaging of multiple targets with unprecedented resolution and quantitative reliability (*92*). To prepare samples for AT, sections from CA1 were blocked in ~2 mm x ~1 mm sections, embedded in the acrylic resin LR White, dehydrated in graded series of ethanol (30%, 40%, 50%, 70% for 10 minutes each at 4 °C) and then transferred to a mixture of 1:3 70% ethanol and LR White (Electron Microscopy Sciences, EMS #14384-UC) for 15 minutes at 4 °C. Subsequently, tissues were placed in 100% LR White for two cycles of 15 minutes at 4 °C and then transferred to fresh 100% LR White overnight. On the second day, the tissue was incubated in pure 100% LR White for two 2-hour intervals at 4°C. Each block of tissue was placed at the bottom of a flow-through capsule and inserted into a BEEM capsule (EMS #7000B) with a corresponding label. A master solution of 10 ml of LR White and one drop of LR White Accelerator (London Resin Company) was used to fill the BEEM capsules and left to

polymerize overnight in a water bath at 4°C. Embedded tissue samples were stored at room temperature until they were cut. A histology knife (DiATOME) was used to remove excess resin and to expose the tissue surface. Sample sections of 70 nm were stained with toluidine blue to confirm the region of interest and to plan trimming area. Next, an ultratrim 45° knife (DiATOME) was used to trim the block face to an area of 2 mm x 1 mm. A Coat-Quick G pen (EMS #70624) was used to apply a small layer of glue to the bottom of the block face to hold the serial sections together as they form a strip. Strips of 15-50 sequential, ultra-thin, 70 nm serial sections were cut using a Jumbo histology knife and mounted on Superfrost Plus charged slides.

Tissue strips were circled with Aqua Hold Pap Pen (EMS #71311) and rehydrated with glycine (50 nM in Tris buffered saline, TBS) for 5 minutes in a humid chamber. Following a rinse with blocking solution (94% TBS, 1% bovine serum albumin BSA, and 5% Tween 20) samples were incubated in additional blocking solution for 5 minutes. Anti-mGluR-2/3 antibody (1:100, Millipore, AB1506) was applied to serial strips and incubated at 4 °C overnight in a humidifying chamber.

The following day, the strips were rinsed with TBS for 4 x 5 minutes at room temperature. Secondary antibody Alexa Fluor 488 donkey anti-rabbit IgG (1:400, Life Technologies, A-21206) and NeuroTrace 640/660 deep-red fluorescent Nissl stain (1:400, N-21483 Life Technologies) were applied to serial strips for 30 minutes at room temperature. The serial sections were rinsed with TBS for 4 x 5 minutes followed by one rinse in distilled water, mounted with Vecta Shield (Vector Labs, H-1000), coverslipped (CSHP-No 1.5-24 × 60; Bioscience Tools), and imaged immediately.

Images were acquired as described (*93*). Briefly, serial sections were imaged on an AxioImager.M2 microscope (Zeiss), equipped with an ORCA-R2 C10600-10B digital camera (Hamamatsu), and the Zen 2012 Blue Edition version 1.1.1.0 (Zeiss) imaging software. Areas of interest were initially identified using a 5x, N.A. 0.16 Plan-Apochromat objective (Zeiss). Once the region comprising the pyramidal neurons of the CA1 region was located, the tissue was imaged at higher magnification with a 63x, 1.4NA Plan-Apochromat objective (Zeiss). The exposure was adjusted for both channels (488 and 640 nm) and a blood vessel or a large cell body that would be visible in the following section was chosen as a landmark. The image of the first section was taken, and the stage was repositioned and aligned to the same landmark in the next section. If needed, the focus and/or exposure were adjusted, and the second image was collected. This process was repeated for the entire length of the serial strip, collecting 7-40 images per slide. The original images collected were converted from .czi to .tif format for subsequent analysis.

The serial images acquired for each sample were processed for analysis with FIJI (ImageJ, U. S. National Institutes of Health, Bethesda, Maryland, USA) (*94*). First, stacks were generated for each channel per sample using the Bioformats Importer Plug-In. Next, the background was subtracted (rolling ball diameter 50 pixels), and the image stacks were normalized (pixel saturation 0.1%) to adjust for variation in fluorescence between sections (*95*). The processed image stacks were aligned using the MultiStackReg plug-in (courtesy of the Smith

laboratory, Stanford, CA). The NeuroTrace channel was aligned and used as a reference for subsequent alignments of mGluR-2/3. Stacks that could not be automatically aligned were aligned manually using TrakEM2 plug-in (*96*). To analyze the puncta, a threshold was set for each channel using FIJI. To exclude imaging artifacts and cell bodies, thresholds were adjusted so that only punctate signal for the synaptic protein channel remained. Then, images were processed through the Watershed program (Smith laboratory, Stanford, CA), which generates an output with the total number of puncta per image stack and the size, in voxels (3D pixels), of each punctum. These puncta data were used for statistical analysis.

Western blotting

Mouse brains were dissected, and the hippocampus was immediately snap-frozen and stored at -80 ˚C until processed. Tissue samples were suspended at 10% w/v in Tris-buffered saline (TBS), pH 7.6 with 1% v/v Triton X-100 with protease and phosphatase inhibitors (Halt protease inhibitor cocktail 1862209 and 1861274, Thermo Scientific, Rockford, IL and phosphatase inhibitor P5276, Sigma) and homogenized for 20 s at 5,000 rpm using a Precellys 24 homogenizer (Bertin Instruments, Montigny-le-Bretonneux, France). The homogenates were centrifuged at 1,200 x *g* for 10 minutes. Equal volumes of the supernatant were transferred to ultracentrifuge tubes, placed in a TLA-55 fixed-angle rotor, and centrifuged for 1 hour at 100,000xg at 4 °C using an Optima Max-XP tabletop ultracentrifuge (Beckman Coulter, Brea, CA). The supernatant was saved as the soluble fraction. The insoluble pellet was resuspended in 5 μl of TBS + 1% sodium dodecyl sulfate (SDS) with protease and phosphatase inhibitors for every mg of original tissue weight. This insoluble fraction was incubated for 20 minutes and centrifuged for 10 minutes at 16,000 x *g* using a 5254 R table-top centrifuge (Eppendorf, Hamburg, Germany). Protein concentrations were estimated with BCA protein assay (Pierce^{TM} BCA Protein Assay Kit, Thermo Scientific). Samples were stored in -20 °C until ready for use. Samples were separated by electrophoresis on 12% Tris-Glycine SDS-polyacrylamide gels followed by transfer to PVDF membranes (Thermo Scientific, Rockford, IL). The membranes were then incubated in anti-glutamate receptor 2 & 3 (anti-mGluR-2/3, 0.2 μg/ml, rabbit polyclonal, AB1506, Millipore). The secondary antibody used was horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG (1:20,000, Santa Cruz Biotechnology sc-3837). Actin-HRP (1:1,000) was used to detect the housekeeping protein. Membranes were exposed to ECL (Western Lighting^{TM} Plus-ECL, PerkinElmer) and imaged using ChemiDoc MP imaging system and Image Lab software (BioRad). When necessary, blots were stripped of primary antibody using low pH stripping buffer (25 mM glycine, pH 2.0 with 1% w/v SDS). Optical densities of bands were quantified using ImageJ software(*73*). Density was normalized to actin, and results are presented as percentages of wildtype average.

For the synaptic marker western blots, protein was extracted from hippocampi by homogenization in RIPA buffer (Thermo Fisher Scientific) containing phosphatase/protease inhibitor cocktail (Thermo Fisher Scientific) and centrifuged at 15,000 × *g* for 20 minutes at 4°C before collecting the supernatant. A 30 μg aliquot of total

protein was separated by electrophoresis in precast 4% to 12% Bis-Tris Gels (Bio-Rad) and transferred to nitrocellulose membranes. Membranes were blocked in 5% milk before probing with the following antibodies: mouse anti-PSD95 (clone 6G6-1C9, 1:4000, EMD Millipore, MAB1596), rabbit anti-synaptophysin (1:1000, Abcam, ab16659), and mouse anti-GAPDH (1:2000, SantaCruz, sc-32233). The secondary antibody used was peroxidase-labelled anti-rabbit or anti-mouse IgG (1:5000, Vector Laboratories). SuperSignal West Femto ECL (Pierce, #34096) was used to reveal the immunoreactive proteins, and images were acquired using a ChemiDoc MP (Bio-Rad). Luminescent immunoreactive protein bands were quantified using Fiji software (ImageJ)*(94)*. Density was normalized to GAPDH, and results are presented as the fold-change compared to wildtype average.

Immunohistochemistry

For cross-competition blocking experiments, PFA-fixed 40μm thick sections from a 5XFAD transgenic mouse (B6SJL-Tg(APPSwFlLon,PSEN1*M146L*L286V)6799Vas/Mmjax) were used. Free-floating sections were washed with 0.1 M phosphate buffer saline (PBS) in 0.1% triton (PBST) three times for 15 min and blocked with 2% bovine serum albumin (BSA) for 1h. For blocking experiments, prior to antibodies reactivity, sections were incubated with unlabeled CP-2 (25 μM) or PBS overnight. Sections were then washed with DDW three times for 10 min and incubated with NFA-Aβ-detecting A11 antibody (1:2500) in antibody buffer (PBST containing 2% BSA) overnight at 4°C. Sections were then washed with 1% PBST and incubated with Alexa Fluor 568 secondary goat anti-rabbit antibody (1:1000 Abcam) in antibody buffer for 1 h at room temperature. Sections were then stained with DAPI (4',6-diamidino-2-phenylindole dihydrochloride; 1:500, MP Biomedicals) and washed for 10 min in PBS with 0.1% TBST. Stained sections were mounted on super frost glass slides, embedded with ImmunoMount (Sigma-Aldrich, Switzerland), coverslipped, and stored in dark for microscopic images.

Free-floating sections were stained for immunofluorescence with anti-IBA1 immunohistochemistry (IHC) and were mounted onto slides for double-labeling. For anti-IBA1 IHC, sections were washed in PBS three times for 10 minutes and then blocked in PBST with 5% normal goat serum (NGS) for 1 hour at room temperature. Sections were incubated overnight at 4 °C in primary antibody anti-IBA1 (1:500, Wako, 019-19741) in PBST with 2% NGS. The tissues were subsequently washed in PBST, then incubated in PBS with 1% NGS and biotinylated secondary antibody. Sections were washed in PBS, then mounted and dried, before coverslipping and imaging with an Olympus BX61 microscope.

For double-labeling with anti-C1q and anti-PSD95, slides were rinsed three times for 10 minutes in PBS and then incubated in a citrate buffer for antigen retrieval. Sections for C1q complex (C1q) were incubated in the water bath at 85 °C in citrate buffer with a pH of 9.5 for 30 minutes, followed by 20 minutes at room temperature. They were then rinsed again and placed in a dilution buffer containing 10% NGS and 0.4% Triton-X for 1 hour. One section from each animal was placed into primary antibody for anti-rat C3 (1:100, MP Biosciences, 55730) and another for anti-rabbit C1q (1:100, Abcam, ab182451). Both antibody solutions also contained anti-mouse

postsynaptic density protein (PSD95; 1:100, Millipore, MAB1596). Slides were incubated overnight at 4°C. Following incubation, slides were rinsed and then incubated for one hour in secondary antibody solutions of anti-rabbit Alexafluor 488 (Invitrogen, A11034) and anti-mouse Alexafluor 594 (Invitrogen, A11032) or anti-rat Alexafluor 488 (Invitrogen, A11006) and anti-mouse Alexfluor 594. Slides were rinsed and coverslipped using Vectashield Vibrance with DAPI (Vector Laboratories, H-1800). Z-stack multi-channel images were taken at 63x using a Zeiss LSM 800 Confocal and Zen Blue software (v. 2.6, Carl Zeiss, Inc, Munich, Germany) and analyzed with Fiji Software (ImageJ).

Dynamic light scattering

To determine the size of NFA-Aβ before and after incubation with CP-7, a solution of Aβ (30 µM) in PBS was incubated in the absence or presence of cyclic peptide **7** (150 µM) for 72 h and the size of the particles was monitored by a Malvern Zetasizer Nano ZS instrument (Malvern, UK) operated with a 3 mW He−Ne laser source operating at a wavelength of 633 nm. Scattered light was detected at an angle of 173° (*99*).

Mitochondrial electron transport chain (mtETC) activities

The activities of mitochondrial complexes I (CI), II (CII), III (CIII), and IV (CIV) were assayed using a spectrophotometric enzyme activity assay (*104, 105*) performed on a FLUOstar Omega spectrophotometric plate reader (BMG). mtETC complex activities were measured and normalized to both citrate synthase (CS) activity and protein concentration. CS is a common readout for mitochondrial mass, and normalization of complex activities to CS activity can reveal actual alterations in activities of the complexes themselves, or alterations in their abundances within the mitochondria, while normalization of complex activities to total protein concentration of the homogenates can potentially reveal alterations in the mitochondrial mass within the tissue. About 10-20 mg of frozen brain tissue was homogenized in SEF Buffer (10 mM K_2HPO_4/KH_2PO_4, 250 mM sucrose, 2 mM EDTA), using a bead mill homogenizer and 1.5 mL microtubes pre-filled with 1.4 mm ceramic beads (Omni International).

Complex I (CI): Cell homogenates were incubated in a potassium phosphate (K_2HPO_4)-buffered solution of bovine serum albumin (BSA) in the presence of nicotinamide adenine dinucleotide (NADH), ubiquinone (CoQ), and the artificial electron receptor 2,6- dichlorophenolindophenol (DCPIP). Ubiquinone reduced to ubiquinol (QH2) by the oxidation of NADH to NAD+ by CI rapidly reduces DCPIP (blue) to DCIPIH2 (colorless). The NADH oxidation was assayed through spectrophotometric measurement of the extinction of DCPIP absorption at 600 nm over 17 minutes. Non-specific NADH oxidation was determined by simultaneously assaying NADH oxidation in each tissue homogenate in the presence of the potent CI inhibitor rotenone, and CI activity was calculated by subtracting the nonspecific (rotenone-inhibited) NADH oxidation from the total (rotenone-uninhibited) NADH oxidation.

Complex II (CII): Cell homogenates were incubated in a K_2HPO_4-buffered solution of BSA, ethylenediaminetetraacetic (EDTA), and sodium azide (NaAz) in the presence of succinate, decylubiquinone (DUB), adenosine triphosphate (ATP) and DCPIP. DUB reduced to DUH2 by the oxidation of succinate by CII rapidly oxidizes DCPIP (blue) to DCIPIH2 (colorless). Succinate oxidation was assayed through spectrophotometric measurement of the extinction of DCPIP absorbance at 600 nm for 15 minutes. Non-specific reduction of DCPIP was corrected for by simultaneously assaying each tissue homogenate in the presence of the potent CII inhibitor malonate, and CII activity was calculated by subtracting the nonspecific (malonate-inhibited) succinate oxidation from the total (malonate-uninhibited) succinate oxidation.

Complex III (CIII): Cell homogenates were incubated in a K_2HPO_4-buffered solution of EDTA, NaAz, and polysorbate 20 in the presence of reduced DUH2 and cytochrome C (CytC). Reduction of CytC by CIII via oxidation of DUH2 was assayed via spectrophotometric measurement of reduced CytC absorbance at 550 nm over 15 minutes. Non-specific reduction of CytC was corrected for by first assaying for CytC reduction in the presence of DUH2 but in the absence of the homogenate.

Complex IV (CIV): Cell homogenates were incubated in a K_2HPO_4-buffered solution in the presence of reduced CytC. The oxidation of reduced CytC by CIV was assayed via spectrophotometric measurement of the extinction of absorption by reduced CytC at 550 nm over 15 minutes. The reaction endpoint was assessed by artificially oxidizing all reduced CytC in the reaction mixture via the addition of the potent oxidizer potassium ferricyanide ($K_3Fe(CN)_6$), and the total reduced CytC in the reaction was quantified using a triplicate of blank wells lacking tissue homogenate.

Citrate Synthase (CS): Cell homogenates were incubated in a TrisHCl-buffered solution in the presence of oxaloacetic acid, acetyl-CoA, and 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB). CoA-SH generated by the condensation of acetyl-CoA and oxaloacetic acid by CS to citric acid rapidly cleaves DTNB (colorless) to TNB-, which rapidly ionizes to TNB2- (yellow). Condensation of acetyl-CoA and oxaloacetic acid by CS was assayed via spectrophotometric measurement of TNB2- at 411nm over 15 minutes. Protein concentration of cell homogenates was assayed using the Pierce BCA Protein Assay (Millipore).

Surface plasmon resonance (SPR) assays

All experiments were performed using a Biacore T200 (Cytiva) and carried out at 25°C. The instrument was desorbed before use with the Biacore automated 'Desorb' routine. A CM5 sensor (Cytiva) was docked, running buffer changed to PBS-T and the sensor cleaned with a 30 second injection of 50 mM NaOH (0.22 μm filtered; 10 μl/min flow rate). Aβ42 fibrils were prepared at 20 μg/ml and Aβ42 NFAs at 4 or 20 μg/ml in 10 mM sodium acetate buffer at pH 5.5, 5.0, 4.5 or 4.0. The proteins were injected onto the sensor surface at 10 μl/min for 120 seconds to confirm the optimum immobilization pH. The sensor surface was then cleaned with a 30 second

injection of 50 mM NaOH. Aβ42 oligomers could not be pre-concentrated onto the sensor surface and could not be immobilized by amine coupling.

Aβ42 NFAs, Aβ42 fibrils, A11 antibody and BSA were immobilized on the sensor surface by direct amine coupling. Reference or active channel surfaces were activated for 7 minutes with a 1:1 mixture comprising 200 mM 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) and 50 mM N-Hydroxysuccinimide (NHS). Aβ42 fibrils (Fc2), A11 antibody (Fc3) and Aβ42 NFAs were coupled to the active channel surfaces. BSA was coupled to the reference channel surface on Fc1. All unreacted EDC-NHS groups were then inactivated with a 7-minute injection of 1 M ethanolamine. All solutions were injected onto the relevant flow channel at a rate of 10 µl/min.

CP-2 was prepared as a stock solution in DMSO. For CP-2 binding assays, running buffer was changed to PBS-T with 1% DMSO. For all other binding assays running buffer was PBS-T. Analytes were injected onto reference (Fc1) and protein immobilized channels (Fc2, Fc3, Fc4) for 120 seconds and dissociated in running buffer for 180 seconds. The sensor surface was regenerated with a 30 second injection of 10 mM glycine pH 1.5. Samples were injected in ascending concentration series and in duplicate at each concentration. Vehicle control injections were included before and after the analyte binding series for buffer blank subtraction.

Binding response data were analyzed using Biacore T200 Evaluation software. Binding responses were reference channel subtracted (i.e., Fc2-Fc1, Fc3-Fc1, Fc4-Fc1 as appropriate) and buffer blank subtracted (double referenced). Response amplitude was then averaged over a 5 second window, 5 seconds before the end of analyte sample injection of the double reference subtracted binding responses.

Figure S1. Quantification of dot blot assay and quantification of A11 and OC epitomic IHC and CP-2-FITC fluorescence microscopy from Figure 1. (A) Dot blot assay with quantification for A11/6E10 ratio, OC/6E10 ratio and α-tubulin levels across all groups. (B) Quantification of mAb158 sandwich assay showing no differences in Aβ PF concentration between TBS soluble brain fractions extracted from 12-month-old Dutch (n=3/sex) and wildtype (n=1/sex) mice. (C) Quantification of A11 IHC, and CP-2-FITC(22) comparing 2- and 24-month old Dutch mice.

Figure S2. Dutch mice show aging-related accumulation of NFA-Aβ in the thalamus using DAPI, A11 immunohistochemistry, and CP-2-FITC fluorescence microscopy(*22*).

Figure S3. Dutch mice show aging-related elevation of levels of NFA-Aβ in the hippocampus using DAPI, A11 immunohistochemistry, and CP-2-FITC fluorescence microscopy(*22*)**.**

Figure S4. CP-2 cyclic peptide binding alters binding of NFA-Aβ-detecting A11 antibody. Representative immunofluorescence image of adjacent sections of the (A-B) hippocampus and (C-D) cortex of the same 7-month-old 5xFAD mouse brain stained with NFA-Aβ-detecting A11 antibody in presence of either PBS (A and C) or excess cyclic peptide CP-2 (B and D).

Figure S5. No change in expression of mGluR-2/3 in the hippocampus of wildtype and Dutch mice. (A) Densitometric quantification of mGluR-2/3 normalized to actin from 12-month-old wildtype and Dutch mice and expressed as a percentage of the protein level in wildtype mice with example western blot. Data are mean ± SD from n = 5 wildtype and n = 5 Dutch mice. **(B)** Higher magnification of a single 70 nm section, showing synaptic puncta of mGluR-2/3 (green) and GluN1 (red), along with a nucleus stained with DAPI. Cell bodies stained with Neurotrace were used as landmarks for alignment. Scale bar = 10 μm. Large Inset is an example of serial sections stained with Neurotrace and imaged at 10x for array tomography. The images were stitched together to render a full-length depiction of a typical sample. Small Inset is magnifying the area of serial sectioning in the box; arrow indicates the CA1 cell layer. Scale bars = 1 mm and 0.5 mm (inset). **(C)** Density of synaptic puncta immunoreactive for mGluR-2/3 were similar in 12-month-old wildtype and Dutch mice. Data are mean ± SD. **(D)** Distribution of puncta sizes was comparable between the two genotypes for mGluR-2/3 synapses. Data are mean ± SEM from n = 4 wildtype and n = 5 Dutch mice. **(E)** Density of synaptic puncta immunoreactive for GluN1 were similar in 12-month-old wildtype and Dutch mice. Data are mean ± SD. **(F)** Distribution of puncta sizes was comparable between the two genotypes for GluN1 synapses. Data are mean ± SEM. The puncta were assessed from n = 4 wildtype n = 5 Dutch mice.

Figure S6. Synaptic vesicle size, number, and distribution are comparable in the CA1 of 12-month-old wildtype and Dutch mice. (A) Representative EM micrograph of presynaptic vesicles. Green dots indicate vesicles in the presynaptic terminal and the green line indicates the PSD, as described in the Methods section. Scale bars = 400 nm. **(B)** Frequency distribution of vesicle size. **(C-D)** Average counts of synaptic vesicles within 50 nm from the active zone or total vesicles at the presynaptic terminal of all synapses. **(E-F)** Average counts of synaptic vesicles within 50 nm from the active zone or total vesicles at the presynaptic terminal of synapses with a head diameter <0.4mm (small, thin synapses). **(G-H)** Average counts of synaptic vesicles within 50 nm from the active zone or total vesicles at the presynaptic terminal of synapses with a head diameter >0.4mm (large, mushroom synapses). Data are mean ± SD from wildtype (n = 4) and Dutch (n = 4) (Multiple Mann-Whitney).

Figure S7. No differences in levels of synaptic markers between Dutch and wildtype mice. (A) Western blot images for postsynaptic density protein 95 (PSD95) of hippocampal lysates from 12-month-old wildtype and Dutch mice of both sexes. **(B)** Densitometric quantification of PSD95 normalized to GAPDH and normalized to the protein level in wildtype mice. **(C)** Western blot images for synaptophysin (SYP) of hippocampal lysates from 12-month-old wildtype and Dutch mice of both sexes. **(D)** Densitometric quantification of SYP normalized to GAPDH and normalized to the protein level in wildtype mice. **(E)** Western blot images for GAPDH of hippocampal lysates from 12-month-old wildtype and Dutch mice of both sexes. Data are mean ± SEM from n = 10 wildtype (5M and 5F) and n = 9 Dutch mice (5M and 4F). *DU was excluded from all quantification as an outlier.

Figure S8. Quality control filtering of scRNA-Seq data. (A) Raw data values and **(B)** filtered values for nFeatures (left), nCounts (center) and % mitochondria reads (right) for the samples. **(C)** UMAP from Figure 6 with split UMAPs for Dutch and wildtype samples.

Figure S9. Transcriptional profile of Cluster 0- excitatory neurons is altered in Dutch mice. (A) Volcano plot of genes enriched in Cluster 0 as calculated by pseudobulk differential expression analysis. Red genes are Log2FC > 0.25 and adj. p-value < 0.05, blue Log2FC < 0.25 and adj. p-value < 0.05, green genes are Log2FC > 0.25 and adj. p-value > 0.05, and grey genes are Log2FC < 0.25 and adj. p-value > 0.05. **(B-C)** GO term analysis of **(B)** up-regulated and **(C)** down-regulated genes enriched in Cluster 0 (adj. p-value < 0.05) using metascape(*91*).

Figure S10. Transcriptional profile of Cluster 1- astrocytes is altered in Dutch mice. (A) Volcano plot of genes enriched in Cluster 1 as calculated by pseudobulk differential expression analysis. Red genes are Log2FC > 0.25 and adj. p-value < 0.05, blue Log2FC < 0.25 and adj. p-value < 0.05, green genes are Log2FC > 0.25 and adj. p-value > 0.05, and grey genes are Log2FC < 0.25 and adj. p-value > 0.05. **(B-C)** GO term analysis of **(B)** upregulated and **(C)** downregulated genes enriched in Cluster 1 (adj. p-value < 0.05) using metascape(*91*).

Figure S11. Transcriptional profile of Cluster 3 endothelial cells is altered in Dutch mice. (A) Volcano plot of genes enriched in Cluster 3 as calculated by pseudobulk differential expression analysis. Red genes are Log2FC > 0.25 and adj. p-value < 0.05, blue Log2FC < 0.25 and adj. p-value < 0.05, green genes are Log2FC > 0.25 and adj. p-value > 0.05, and grey genes are Log2FC < 0.25 and adj. p-value > 0.05. **(B-C)** GO term analysis of **(B)** upregulated and **(C)** downregulated genes enriched in Cluster 3 (adj. p-value < 0.05) using metascape(*91*).

Figure S12. Transcriptional profile of Cluster 4 oligodendrocytes is altered in Dutch mice. (A) Volcano plot of genes enriched in Cluster 4 as calculated by pseudobulk differential expression analysis. Red genes are Log2FC > 0.25 and adj. p-value < 0.05, blue Log2FC < 0.25 and adj. p-value < 0.05, green genes are Log2FC > 0.25 and adj. p-value > 0.05, and grey genes are Log2FC < 0.25 and adj. p-value > 0.05. **(B-C)** GO term analysis of **(B)** upregulated and **(C)** downregulated genes enriched in Cluster 4 (adj. p-value < 0.05) using metascape(*91*).

Figure S13. No differences in microgliosis or C1q co-localization with synaptic puncta in Dutch mice. (A-B) Representative images of anti-IBA1 IHC in hippocampus (HIP, left) and prefrontal cortex (PFC, right) from 12-month-old wildtype (A) and Dutch mice (B) showed no difference in microgliosis. **(C)** Confocal images of PSD95 (red) and C1q (green) in 12-month-old wildtype and Dutch mice CA1 with quantification of colocalized PSD95- and C1q-immunoreactive puncta (right) revealing differences were not significant. Data are mean ± SEM from n = 4 males for each genotype.

Figure S14. Biological Processes associated with synaptic transmission are altered in excitatory neurons and oligodendrocytes. (A-B) Dot plot of genes associated with upregulated GO terms **(A)** "regulation of synapse organization" from Cluster 6 excitatory neurons and **(B)** "modulation of chemical synaptic transmission", from Cluster 4 oligodendrocytes, split into Dutch and wildtype mice across all clusters. Wildtype mice are in blue, Dutch mice are in purple. In B, genes marked with an asterisk have adj. p<0.05 in Cluster 6.

Figure S15. **Mitochondrial complex I (CI), complex II (CII), complex III (CIII), complex IV (CIV), and citrate synthase activities in the hippocampi of wildtype (open circle) and Dutch (solid circle) mice at age 12 months.** See also Supplemental Tables 8 and 9. For comparison of CI in wildtype vs Dutch, p=0.0355 when standardized to µg protein or p=0.0197 when standardized to mU/ml of citrate synthase (CS) and analyzed using an unpaired t test.