Title: Urbanisation and habitat fragmentation favour thermophilic and monogynous ant species

Authors: Basile Finand[1,2], Thibaud Monnin[1], Céline Bocquet[1], Angélique Bultelle[1], Pierre Fédérici[1],

Léa Darmedru[1], Solène Gouffault[1], Joséphine Ledamoisel[1], Nicolas Loeuille[1]

[1]Sorbonne Université, Université Paris Cité, Université Paris Est Créteil, CNRS, INRAE, IRD, Institut

d'Ecologie et des Sciences de l'Environnement de Paris (UMR7618), 75005 Paris, France

[2]Ecosystems and Environment Research Program, Faculty of Biological and Environmental Sciences,

University of Helsinki, Helsinki, Finland

Corresponding author: Basile Finand, University of Helsinki Yliopistonkatu 3 00100 Helsinki Finland,

finand.basile@gmail.com

Acknowledgements: We would like to thank the City of Paris and the Natural History Museum of

Paris for allowing us to sample the parks of the city. We also thank Xavier Espadaler and Bernard

Kaufmann for their help in the identification of some species.

Conflict of interest: The authors declare no conflict of interest.

Authorship statement: Basile Finand, Thibaud Monnin, and Nicolas Loeuille conceived the study.

Basile Finand, Thibaud Monnin, Nicolas Loeuille, Léa Darmedru, Joséphine Ledamoisel, Céline

Bocquet, Angélique Bultelle, Pierre Fédérici, and Solène Gouffault collected the data. Basile Finand

analysed the data. Basile Finand, Thibaud Monnin, and Nicolas Loeuille wrote the paper. All the

other co-authors agreed on the last version of the paper.

Data statement: Data and codes are available on Zenodo: https://zenodo.org/records/14792981

Abstract:

Environmental changes such as urbanisation and habitat fragmentation profoundly impact ecological

communities by altering habitats, resources, and microclimate. Ants, with diverse life histories and

strong ecological effects, are ideal models to study these pressures. We investigated the response of

ant communities, including taxonomic and functional diversity, to urbanisation and habitat

fragmentation in the Paris region, comparing 24 urban parks vs 25 rural forests outside the city. We

found a clear difference in species composition between urban and rural environments, with a

higher prevalence of monogynous and thermophilic species in the city. Forest communities are

homogeneous across the three fragmentation levels we studied, while park communities differ

noticeably depending on park size, with larger parks harbouring more species. Our findings suggest

that urbanisation selects specific ant traits and favours more thermophilic species, thereby

increasing the mean thermal preference of urban communities. These selective effects influence

which species can colonise and survive in different patches, shaping metacommunity structure and

potentially affecting the resilience of ant communities under climate change.

Keywords: Ants, heat island effect, metacommunity, trait selection, urbanisation

Introduction

Habitat fragmentation stands as a primary driver of biodiversity decline, exerting manifold impacts on ecosystems worldwide (Alberti, 2015; Aronson et al., 2014; Diamond & Martin, 2021; Haddad et al., 2015; Johnson & Munshi-South, 2017; Piano et al., 2020). This phenomenon is characterised by a reduction in total habitat size, with an increase in patch number, but smaller and more isolated (Fahrig, 2003). Habitat fragmentation impedes dispersal between populations, leading to direct impacts on individual and gene flows, resulting in population declines and reductions in genetic variability (Hastings, 1983; Travis et al., 2013; Young et al., 1996). These effects reverberate through ecological communities, altering their composition and dynamics. Habitat fragmentation gives rise to metacommunities, i.e. interconnected communities linked through dispersal. Metacommunity dynamics can happen in various ways depending on species and environmental characteristics (Leibold et al., 2004). In scenarios where variations in environmental conditions among patches do not play a key role, biodiversity hinges exclusively on the competitive abilities and colonisation capacities of species, a dynamic termed patch dynamic. Colonising species can exploit and establish populations in patches where competitors cannot persist (Tilman, 1994). Consequently, habitat fragmentation, through habitat destruction and isolation, is expected to favour colonising species while disadvantaging more competitive species, thereby reducing total diversity (Finand, Monnin, et al., 2024; Tilman et al., 1994). Conversely, in environments where variability among patches drives ecological dynamics, two dispersal-dependent mechanisms emerge. In scenarios with restricted dispersal, species diversity is governed by species sorting, where species whose ecological niches best match local conditions competitively exclude others within each patch. Conversely, when dispersal between patches is extensive, mass effects come into play (Mouquet & Loreau, 2003), allowing less-adapted species to persist in small numbers, being continually resupplied from surrounding patches, akin to source-sink dynamics. Although species sorting typically yields lower species richness per patch compared to mass effect, when dispersal is exceedingly high and mass effects dominate, all species compete for average metacommunity

conditions, leading to competitive exclusion at this scale. Diversity is then low, at the local and

metacommunity scale (Mouquet & Loreau, 2003). The heightened fragmentation of habitat, by

augmenting patch isolation, is expected to foster species sorting, consequently diminishing local

species richness and increasing species turnover among patches. Finally, the neutral theory, akin to

island biogeography (MacArthur & Wilson, 1967), posits that diversity is solely dependent on species

loss through emigration and extinction balanced by gains through immigration and speciation, with

no differentiation in species abilities (Hubbell, 2001). Importantly, these four paradigms can be

disentangled based on variations in species composition and patterns of local (alpha), regional

(gamma) diversity and variations in species composition among patches (beta diversity). For

instance, if species sorting dominates, local composition should differ in contrasted environments so

that beta diversity is mostly associated with a "turnover" component (sensu (Baselga, 2010)).

Conversely, local dispersal in mass effects should lead to a spatial autocorrelation in species

composition so that communities in harsher environments or smaller patches should be a subset of

communities developing in better or larger patches. This enhances the "nestedness" component of

beta diversity (sensu (Baselga, 2010)).

Urbanisation is another major factor affecting biodiversity. This transformative process

reshapes environments in many ways, fostering novel ecological dynamics and posing significant

challenges to native flora and fauna. First, urbanisation is most often associated with high habitat

fragmentation (Haddad et al., 2015). Second, it also generates varied types of pollution, including

chemicals, noise disturbance, and altered light regimes, often favouring species with specific

adaptations or tolerances (Hölker et al., 2010; Newport et al., 2014). These pollutants permeate air,

water, and soil, exerting deleterious effects on biotic components of ecosystems. Third, changes in

resource availability represent another hallmark of urbanisation, including variations in quantity,

quality, and seasonality (Hantak et al., 2021; Wilson & Jamieson, 2019). Finally, the creation of

distinct microclimates within cities, characterised by elevated temperatures (urban heat island

effect), exacerbates thermal stress on organisms and modifies ecosystem dynamics (Kaiser et al.,

2016; McGlynn et al., 2019). This thermal gradient can influence species distributions, phenology,

and physiological processes, driving community composition and structure shifts.

Beyond diversity indices, the specific environmental factors and the low connectivity

encountered in cities should select specific traits. Habitat fragmentation may favour low dispersal

(Cheptou et al., 2008; Finand et al., 2023), while higher temperatures promote species and

phenotypes whose thermal preference matches these conditions (Campbell-Staton et al., 2021;

Diamond et al., 2018). For instance, longer-legged species are selected in hotter parks in Taichung

City because the distance between the body and soil limits desiccation (Liu et al., 2019). In general,

urban and rural species may differ in many other traits (Buchholz & Egerer, 2020; Croci et al., 2008)

reflecting the multidimensional changes in such environments.

Insects and in particular ants have an important impact on ecosystems by dispersing seeds,

pollinating, participating in nutrient cycles, structuring the soil, and participating in food chains

(Hölldobler & Wilson, 1990). They are a very good model group to study the effects of urbanisation

and habitat fragmentation on communities. First, their life history traits are variable between

species. Some of these traits are strongly related to their dispersal abilities (size of individuals,

presence or absence of wings in queens) and competitiveness (size of colonies, diet), and therefore

play a key role as functional traits in metacommunity dynamics. Second, ants are abundantly present

and diverse in and outside cities, which allows direct comparative studies. Most ant species depend

on green spaces, which makes the city a very fragmented habitat compared to the countryside, even

if some species can live in built areas. Last, they are easy to sample. Several empirical studies have

shown that urbanisation and habitat fragmentation affect the species richness of ants, with a

decrease in species richness as urbanisation increases and a change in community composition

(Carpintero & Reyes-López, 2014; Ješovnik & Bujan, 2021; Leal et al., 2012; Pacheco & Vasconcelos,

2007; Theunis et al., 2005). For instance, climate and resource specialist ants are more sensitive to

habitat fragmentation (Leal et al., 2012). Moreover, this change in community composition due to

habitat fragmentation impacts ecosystem services provided by ants influencing decomposition and

soil structure (Sanford et al., 2009).

Paris and its region offer a good opportunity to study urbanisation and habitat

fragmentation, with a marked gradient from the very densely populated city centre to low-density

areas thirty kilometres away. The city has numerous urban parks of various sizes, surrounded by a

matrix of streets and buildings unsuitable for most ant species. This makes Parisian parks a perfect

setup to understand the impact of fragmentation on ant communities. Moreover, the urban heat

island effect within Paris is well-documented (Lemonsu et al., 2015). The surrounding region is more

rural with many natural forests of various sizes allowing an interesting comparison with the city.

We here investigate the impact of urbanisation and habitat fragmentation on ant

communities at different scales by comparing forest (low urbanisation and low fragmentation) and

urban communities (high urbanisation and high fragmentation). We measured community

differences between the two habitats, and the effects of the two main components of

fragmentation, isolation, and patch size, within each habitat. We hypothesise that fragmentation

decreases species richness and changes community structure, with more turnover than nestedness

due to higher species sorting and lower mass effects. Furthermore, we studied variations in traits

linked to thermal preference, colony social structure (queen number) and dispersal abilities. We

hypothesise that the city harbours species that are more thermal-resistant due to the urban heat

island effect, and that have short dispersal distances because of the high habitat fragmentation.

Material and methods

Patch selection

To study the impact of urbanisation and particularly of habitat fragmentation on litter ants,

we sampled two different habitats: 25 rural forests in the Paris region and 24 wooded parts of urban

parks within the city (Figure 1). Forests are considered minimally fragmented environments, as they

offer a favourable habitat for litter ants and are surrounded by a matrix of habitats that, while less

favourable, can still support ant species, e.g. hedges around agricultural fields or private gardens. On

the contrary, parks are islands of favourable habitats (woods and shrubs) surrounded by a mostly

hostile matrix (streets and buildings), though local refugia may exist. To compare the same litter ant

communities in forests and parks, we targeted park areas with natural litter for sampling, i.e., areas

under trees or shrubs. This allowed us to focus on the specific community of ants living in the litter

and to make habitats as comparable as possible. Evidence that we sampled the forest litter ant

community in parks is that the two species most abundant in our park samples are *Temnothorax*

nylanderi and *Myrmecina graminicola*, which are particularly abundant in forests (Blatrix et al., 2013;

Seifert, 2018) and also the most abundant ants in our forest samples (see results). Forests adjacent

to the city are used here as reference communities to assess the effect of urbanisation, which

intrinsically involves habitat change, as it is the closest environment to litter. Differences between

park and forest habitats reflect spatial variations in niche conditions, which is precisely what this

study aims to examine. We study the effect of urbanisation at the regional scale (city parks vs

forests) and the effect of habitat fragmentation at the local scale (parks of various sizes and

isolation, forests of various sizes).

We selected forest patches (n = 24) of three fragmentation classes (Figure 1a). Minimally

fragmented forests (n = 10) had between 75 and 95% of forest cover within 1 Km, medium

fragmented forests (n = 8) had between 30 and 70% of forest cover, and highly fragmented forests

(n = 6) had less than 5% of forest cover in their surroundings. We used the database BD Forêt®

Version 2.0 from the "National Institute of Geographic and Forest Information" of France (IGN,

https://inventaire-forestier.ign.fr/carto/afficherCarto/V2, last visit: 19/07/2022) to assess forest

cover. Forest classes are evenly distributed around Paris, and we chose well-separated locations that

likely represent independent plots (mean distance = 52.1 +/- 23.7 Km, min = 6.4 Km, max = 122.1

Km).

Parks fragmentation was measured using two metrics, size and isolation. We selected parks

(n = 25) of three size classes (Figure 1b), namely large (n = 8, 80 000-250 000 m²), medium (n = 9, 5

000-12 000 m²), and small parks (n = 8, 500-2 000 m²), and measured their isolation as the quantity

of green area (other parks) within 1 Km around the focal park boundaries. As for forests, we selected

parks that were evenly distributed within Paris and well separated, assumed to represent

independent points (mean distance = 4.9 +/- 2.1 Km, min = 552 m, max = 9.5 Km).

Sampling

We sampled forests between May and July 2020, between 11 am and 4 pm, on rainless days.

We distributed 12 quadrats of one square meter each, equally spaced every 9 m, along a transect of

100 m. We independently collected the litter and superficial soil on each of the 12 quadrats. We

collected organisms using the Winkler extraction method (Agosti et al., 2000). We detail the

procedure below.

We sampled parks between May and August 2021, between 11 am and 4 pm, on rainless

days. To compare the same litter ant communities in forests and parks, we targeted park areas with

natural litter for sampling, i.e. areas under trees or shrubs. It allows us to focus on the specific

community of ants living in the litter and make the habitats as comparable as possible with forests.

We purposefully placed the 12 quadrats into these areas and dispersed the quadrats within parks as

much as possible (considering the configuration of the parks, transect was not possible). One park

(OLB) was so small that only five quadrats could be placed and sampled, and owing to a disturbance

only ten quadrats were sampled in another park (MAS). The sampling and extraction protocols were

the same for parks and forests.

For each quadrat, we sifted the litter and the first centimetres of soil using a 1 cm² sieve

(Figure S1a). This removed leaves, twigs and stones and retained the soil and organisms within,

including ants. We kept what passed the sieve in large plastic bags until we returned to the

laboratory. There, we moved the litter to sieve bags that were hung in a Winkler extractor for 48

hours (Agosti et al., 2000) (Figure S1b). As the litter gradually dried out, organisms moved away and

fell into an alcohol-filled collecting vial. This method samples all the macrofauna living in the

collected soil and litter, including ant foragers present in the litter and parts of the colonies that nest

in the litter or superficially in the soil.

We sampled litter ants in forests and parks in two separate years (2020 and 2021) due to

logistical constraints and movement restrictions related to the COVID-19 pandemic. Note that ant

queens and colonies live several years, and that the sampling was done during the same season.

While fluctuations of abundances from one year to another are certainly possible, our analyses focus

on species presence/absence in each quadrat rather than abundance. We further discuss the

implications of our sampling scheme in the discussion section.

Identification of ant species

We identified ants at the species level using a binocular microscope and three identification

keys (Blatrix et al., 2013; Seifert, 2018); Galkowski & Lebas "Identification des *Myrmica*",

unpublished). In addition, *Temnothorax* and *Hypoponera* species were identified by Xavier

Espadaler, while *Tetramorium* species were identified by Bernard Kaufmann.

Ant species traits

We selected different species-level traits to test for the effect of urbanisation. Queen size,

worker size and colony size (number of individuals) were obtained from the bibliography for all

collected species (Blatrix et al., 2013; Seifert, 2018). In addition, colony mass was estimated as

worker size[3] X colony size. Because colony mass is directly linked to metabolic requirements (Brown

et al., 2004), we expect that it could vary between urban and forest sites, either due to changes in

resources or temperature regimes.

We also analysed the social structure of a colony, with two levels: monogyny (species with

typically one queen per colony) and polygyny (species with typically several queens per colony). This

trait is to some extent linked to dispersal as monogynous species often perform independent colony

foundation (long dispersal distance strategy but low competitivity) whereas polygynous species do

colony fission (short dispersal distance strategy but high competitivity) (Cronin et al., 2013; Keller,

1991).

Finally, for each species, we estimated thermophily based on a meridional index from

Daufresne et al. (2004) (Daufresne et al., 2004). Recent advances in ant thermal ecology have

highlighted more precise traits such as critical thermal limits, activity temperatures, or behavioural

tolerances (Nascimento et al., 2022; Parr & Bishop, 2022; Roeder et al., 2021). They are well-suited

to characterise precisely the thermal performances of a few species in a given environment.

However, they require live workers in significant numbers (20 to 30 workers per colony, more for

polymorphic species, and several colonies per species), making them incompatible with our sampling

protocol with Winklers. The meridional index is better suited for analysing diverse assemblages

across broad spatial scales, like in our study. In addition, it integrates factors other than thermal

resistance per se that may relate to thermal preference, e.g. circadian rhythm of activity, and it is

generalisable across taxa and compatible with studies on biogeographical shifts under climate

change. It provides a useful, scalable metric for detecting broad patterns of community shifts along

environmental gradients. Species were considered meridional if most of their distribution lies south

of Paris, with meridionality being calculated by (Paris latitude – minimum observed latitude) /

(maximum observed latitude – minimum observed latitude). The meridional index for a given species

is, therefore, close to one when Paris is at the northern limit of its distribution, and close to zero

when Paris is at the southern limit of its distribution. Minimum and maximum latitudes were

obtained for each species distribution from the antmap.org website (last visit: 28/02/2024; (Guénard

et al., 2017; Janicki et al., 2016)). Only the records considered to pertain to the native distribution

were considered. Following previous works, we use this index as a proxy of the thermal preference

of a species. A Parisian species at the northern limit of its distribution is more thermophilic than one

at the southern limit. Paris is farther north than the northern range limit for two species. We have

attributed 1, the highest value, to them.

To understand which traits are linked to urbanisation, we assign a value of urbanisation

preference for each species, calculated as the number of urban patches where the species has been

found divided by the total number of patches (urban and forest patches) where it has been found.

Strictly urban and forest species have urbanisation indices of one and zero, respectively.

Analyses

Statistical analyses were done with R v3.6.2 (R Core Team 2016). We checked that our

sampling was sufficient to sample the litter ant biodiversity with rarefaction curves and chao analysis

using the vegan package.

We studied alpha diversity as a function of urbanisation at the regional scale (forests vs.

parks) using ANOVAs. At the local scale (among parks and among forests), the observed species

richness and the Shannon index were analysed as a function of park size and isolation, and forest

fragmentation using ANOVAs. We checked that the data complied with application conditions using

the DHARMa package (Hartig, 2022), and log-transformed them when normality or homoscedasticity

were not satisfied.

We analysed beta diversity with Jaccard's dissimilarity index because we here focus on

species presence/absence rather than on abundance. Direct use of individual densities could be

problematic, as our sampling procedure can recover isolated individuals as well as whole colonies for

some *Temnothorax* species and *Myrmecina graminicola*. The second instance would inflate recorded

abundances of superficial species compared to species nesting deeper in the soil for which we only

collect foragers. We performed db-RDA (distance-based redundancy analysis), an RDA on

dissimilarity distances, to assess if our environmental factors affect dissimilarity between our

communities (Legendre & Anderson, 1999). We used ANOVAs to test the significance using the

vegan package. We used the package *betapart* to assess the percentage of turnover and nestedness

composing the dissimilarities between our communities (Baselga & Orme, 2012).

To link the urbanisation preference to species traits, we first checked the correlation among

traits using a Pearson test. The different size measurements were highly correlated (Figure S2). We

kept the total mass of a colony, as it contains the size of a colony and the weight of individuals. In

the end, variations in three traits were explored: the total mass of a colony, the social structure, and

the meridional index. To assess if the urbanisation preference of species is correlated with each trait,

we performed a linear model for the colony mass and the meridional index and a generalised linear

model with a binomial link function for the social structure. We tested the application conditions

(normality, homoscedasticity) using the DHARMa package (Hartig, 2022). We log-transformed the

total mass.

Results

We collected and identified 36 361 ants from 12 genera and 29 species. Abundances are

similar in forests and parks (18 094 and 18 267 ants, respectively). Similarly, the number of species

recovered per habitat type is comparable (Figure 2, 23 species in forests and 20 species in parks). 14

species occurred in both parks and forests, albeit in most cases with different relative abundances,

while 9 species occurred in forests and 6 in parks only (Figure 2). One invasive species only was

recorded (*Lasius neglectus*). Removing it from the analyses does not qualitatively change the results.

		Sampling effort

The rarefaction curves and the chao analyses show that the sampling was sufficient to

estimate litter ant diversity (Figure S3). Indeed, we sampled 95.19 +/- 25.41 % of the theoretical

number of species in forests and 91.13 +/- 13.12 % in parks. Only two forests (CSM and GIF) and

three parks (HEA, HEG and PDG) have a percentage of observed species representing less than 80%

of the theoretical number of species. However, this theoretical number of species is higher in these

patches (around or above 15) than in the other patches (usually around 10, seldom higher than 12)

so it may be overestimated. Therefore, we use the observed number of species per patch in the

analyses.

Urbanisation effect at the regional scale (forests vs parks)

Unexpectedly, we found more species locally with increased urbanisation (F value = 6.67, p =

0.013, R² = 10.6%, Figure 3a). Indeed, parks support 8.96 +/- 1.74 species while forest sites have 7.58

+/- 2.30 species. In agreement with this, the Shannon index is higher in the city environment

showing a higher diversity with less dominant species (T value = 2.808, p = 0.007, R² = 13%, Figure

3d).

Urbanisation also influences species composition (F value = 28.47, p = 0.001, R² = 37.7%,

Figure 4a). Communities are clearly different between parks and forests, with 89% of the beta

diversity due to turnover (Jaccard index = 0.517, Figure 2), suggesting a dominant role of species

sorting.

Fragmentation effect at the local scale

The fragmentation level at the local scale has no effect. In forests, it does not affect species

richness (F value = 0.17, p = 0.84, Figure 3c), Shannon index (F value = 0.488, p = 0.62, Figure 3f) or

the composition of ant communities (F value = 1.37, p = 0.128, Figure 4c). Similarly, for parks,

isolation does not affect species richness (t value = -1.677, p = 0.107, Figure S4a), Shannon index (t

value = -1.866, p = 0.075, Figure S4b) or ant communities composition (F = 0.77, p = 0.69).

Effect of park size

As expected, larger parks harbour more species than smaller parks (F = 3.74, p = 0.039, R^2 =

0.19, Figure 3b). The effect is significant between small and large parks (pairwise comparison: large-

medium p = 0.21, medium-small p = 0.57, large-small p = 0.03). Large parks have on average 8 +/- 2.7

species, medium parks have 7.38 +/- 2.07 species and small ones have 7.17 +/- 2.14 species. Patch

size has a similar but marginal influence on the Shannon index (F value = 2.83, p = 0.081, Figure 3e).

Patch size also affects the spatial turnover of ant communities (F value = 2.14, p = 0.008, R^2 =

16.3%, Figure 4b). The beta diversity is composed of 100% of turnover between large and medium

parks, 84% between medium and small, and 93% between large and small (Jaccard index large-

medium = 0.222; medium-small = 0.278; large-small = 0.450) highlighting a dominant role of species

sorting.

Effect of urbanisation on species traits

There is no effect of urbanisation on the total mass of a colony (t value = -0.76, p = 0.46,

Figure 5a). However, it is correlated with the social structure (z value = -2.26, p = 0.02, R2 = 0.15,

Figure 5b), with more monogynous species in urban habitats. It is also correlated with a higher

meridional index, meaning that urban environments tend to select thermophilic ant species (t value

= 2.89, p = 0.008, R2 = 0.21, Figure 5c).

Discussion

Our study shows that urbanisation and the associated habitat fragmentation influence the

litter ant communities at the metacommunity scale. Surprisingly, the ant biodiversity assessed by

the total number of species per environment (gamma diversity) is stable, and the species richness

per patch is slightly higher in parks than in forests. Communities differ markedly between the two

environments, with a strong turnover in species compositions. At a local scale (alpha diversity),

species richness and community composition are affected by park size and do not vary among

forests. Finally, we highlight that the turnover in species composition between urban and forest

patches is associated with systematic differences in species phenotypes. Urbanisation selects species

that tend to be monogynous and thermophilic.

A possible limitation of our study is that forests and urban parks were sampled in different

years (2020 vs. 2021). While we acknowledge this issue, we however think that potential biases are

here limited. A first reason stems from the fact that our method relies on community composition

(presence/absence of species), not on species abundances. While abundances may fluctuate from a

year to another, we do not expect such variations when looking at the presence/absence of long-

lived species. Ant queens and colonies are indeed long-lived (Hölldobler & Wilson, 1990). Ant queens

live on average 10 years, up to 28 years, and colonies outlive queens in species where colonies have

several queens or where queens can be replaced, e.g. by adoption of daughter queens (Keller, 1998).

The composition of ant species community is therefore relatively stable from one year to the other

(Donoso, 2017; Herbers, 2011). Focusing on species presence/absence, therefore reduces sensitivity

to yearly fluctuations in colony size or activity. The clear differences observed between forests and

parks align with ecological gradients (urbanisation, fragmentation, microclimate) rather than

temporal effects. Notably, 2020 (forest sampling) was warmer in the Paris region than 2021 (park

sampling, 2020: 14.3 °C, 2021: 12.9 °C. Source: opendata.paris.fr), yet we still detected

thermophilisation in parks, which strengthens our inference. Finally, both habitats were sampled in

the same season during no rain days, with sufficient effort to capture nearly all expected species

(Fig. S3). Taken together, these points support that the observed differences reflect genuine

ecological responses to urbanisation and fragmentation rather than temporal sampling artefacts.

At a regional scale, comparing rural versus urban environments reveals no differences in the

total number of species but a clear difference in community composition, with beta diversity

dominated by its turnover component. This suggests that dispersal is low between the two

environments, creating different communities adapted to each, and highlights species sorting as a

strong regional-scale mechanism in this metacommunity. Indeed, if dispersal between forest and

urban sites was high, we would expect either community composition to be similar (low beta

diversity) or, if colonisation success differed between the environments, that the communities of

one environment would be a subset of the communities of the other environment (high nestedness,

(Baselga, 2010)). Differences in community composition between forests and parks can also reflect

differences in fragmentation and shifts in ecological niches. In parks, wooded areas represent only a

portion of the park and are often non-contiguous and surrounded by open habitats, allowing

workers from these patches to forage in the sampled litter. In contrast, even the most fragmented

forest patches were large enough to allow sampling away from the forest edge. Climate represents

another niche axis: parks are warmer due to the urban heat island effect, attracting warm-adapted

species, as confirmed by our results. Urbanisation also introduces various pollutants, which may

select for particular species and traits. These niche changes, inherent to urban environments, likely

contribute to the observed differences in community composition.

Surprisingly, more species are present per urban park than per forest. Several explanations

can be proposed. First, as mentioned above, species sorting dominates, suggesting restricted

dispersal between forests and city parks, and parks may thus host more ant species than forests if

they harbour a higher diversity of habitats and microenvironments. Although our sampling within

parks targeted shaded litter patches under bushes or trees to mimic forest conditions, they were

often surrounded by more open and trampled areas, potentially creating a range of microhabitats.

This aspect, not fully quantifiable in our study due to a lack of direct microhabitat characterisation,

could contribute to higher species richness in parks. However, if parks were richer simply because of

a higher microhabitat heterogeneity, we would expect to find a high nestedness. Instead, our data

reveal a near-complete turnover in species composition between forests and parks, suggesting that

parks do not merely add species but rather replace forest species with species better adapted to

urban conditions. Since warmer climates support more species, parks may receive a larger pool of

colonisers, potentially increasing their species richness. This is confirmed with the trait analysis,

where we have more warm-adapted species in the parks. Another possibility is that species richness

in parks might be "inflated" by continuous colonisation processes such as soil inputs during park

maintenance or transport of material within urban areas. These hypotheses would implicitly

question our view of forests being good patches compared to parks. An alternative explanation

would be that forests are (or have been until relatively recently) well-connected and form a

metacommunity of their own. If so, high dispersal among forests could homogenise ant forest

communities through mass effects, leading to a depressed lower alpha diversity reflecting partial

homogenization (Leibold et al., 2004; Mouquet & Loreau, 2003). In contrast, when dispersal is low,

as may be the case between parks, competition occurs independently within each park resulting in

different communities hence higher alpha and gamma diversities. This increases turnover between

patches as observed in our data. This is likely given that parks were made for human leisure and

differ more from one another than natural forests do.

Our observations of local community compositions support the idea of strong mass effects

among forests. Indeed, there is no difference in species richness and species composition among

forests, even for the most fragmented ones. This suggests a high degree of dispersal between forest

sites, which homogenises the communities. This is plausible because the matrix separating forests is

less hostile than the one separating parks. Indeed, field hedges, roadsides and private gardens may

connect forests by a network of trees and shrubs producing leaf litter and facilitating the dispersal of

ants. Moreover, forests may be more similar to one another than parks. The former evolved from

natural forests and have been managed largely for similar usages (e.g. wood production, leisure,

hunting), whereas the latter are artificial and have to some extent been conceived to differ from one

another to provide varied scenic environments. This could be confirmed by a more systematic study

of biotic and abiotic environmental variations. If so, such homogeneity could limit the number of

local niches, hence species sorting.

In urban areas, on a park scale, species richness is generally lower in smaller parks, and

community composition differs due to high species turnover. This pattern aligns with the classical

species-area relationship, which links larger areas to higher diversity (Preston, 1962), potentially

reflecting intensified species sorting in smaller parks (Leibold et al., 2004; Mouquet & Loreau, 2003).

In smaller parks, competitive exclusion may be more pronounced, as only the most competitive

species persist, while low dispersal limits the arrival of new species. This limited dispersal is

exacerbated by the hostile urban matrix surrounding these parks, which restricts movement and

colonisation, particularly in smaller parks less accessible through natural or human-mediated

dispersal (e.g., fewer plants or materials introduced). Consequently, these combined niche and

dispersal effects contribute to lower alpha diversity and higher turnover between smaller parks.

These patterns are consistent with other studies, including other ant communities. Smaller parks in

highly urbanized zones tend to harbour fewer ant species than those in peri-urban, less isolated

areas (Pacheco & Vasconcelos, 2007), while within cities, larger parks generally support greater

alpha diversity (Carpintero & Reyes-López, 2014). This trend extends to other ecosystems and taxa.

For example, forests show similar patterns (Leal et al., 2012). Other studies on isolated zooplankton

communities in controlled environments also report reduced diversity due to limited dispersal

(Steiner & Asgari, 2022), mirroring the low mass effects in our urban ant metacommunity. Such

findings are consistent with island biogeography theory, which suggests that isolated habitats, like

urban parks or fragmented forests, host fewer species than more contiguous areas (Ramalho et al.,

2022).

Beyond the turnover in species composition, we highlight that phenotypic traits are

deterministically associated with community variations. Urban species are more often monogynous

and more thermophilic. While the second trait was expected, given the heat island effect in cities,

the first was not. Monogyny is somewhat linked to dispersal capacity (Keller, 1991; Sundström et al.,

2005). Monogynous colonies are often founded by winged queens that disperse alone by flying over

long distances. In contrast, polygynous colonies are either founded by related non-flying queens that

disperse on foot over short distances with workers or arise secondarily by non-flying queens that

remain in their natal colony. Theoretically, habitat fragmentation can select for high dispersal (i.e.

winged queens), for instance, when dispersal is traded off with competition capacity (Finand,

Monnin, et al., 2024; Tilman, 1994), when fragmentation leads to high kin competition in remaining

patches, or to avoid inbreeding (Cote et al., 2017; Gandon, 1999; Hamilton & May, 1977). Some

empirical studies support these theoretical results, including in ants. For instance, at the intraspecific

level, colonisation ability can be in a trade-off with competition in an ant species (Finand, Loeuille, et

al., 2024). This selection for high dispersal in a fragmented context could explain, as a byproduct,

that city parks harbour more monogynous species because polygynous species disperse less.

However, some studies have shown opposite results, with habitat fragmentation in cities selecting

for lower dispersal (Cheptou et al., 2008; Finand et al., 2023). Indeed, while habitat fragmentation

usually favours high dispersal, it may have the opposite effect under some circumstances (e.g. highly

hostile matrix increasing dispersal costs, Bonte et al. 2012; Hastings 1983). A second explanation is

that monogynous species can be considered as pursuing spatial bet-hedging because flying queens

can access and distribute the risks on more patches than walking queens. Indeed, a colony producing

flying queens can attempt to establish new colonies on many patches. In contrast, a colony

producing apterous queens produces a few only, as each requires the allocation of workers, and

they have restricted dispersal, hence it has few establishment attempts. A bet-hedging strategy is

selected under temporally variable environments (Slatkin, 1974). City patches may be more variable,

due to many disturbances associated with human activities, and parks surrounded by unfavourable

environments. Having winged queens will increase the probability of having offspring in a more

favourable environment. A third hypothesis is that urban parks are relatively new, particularly

regarding ants that have a longer generation time than most other insects. The first species

colonising a new patch are the most dispersive ones, which in ants have winged queens and thus are

monogynous. Dispersal limitation also likely lasts in this highly fragmented context. Forests being

much older, the less dispersive but more competitive species had the time to colonise them and

progressively exclude better colonisers. All these hypotheses are not exclusive. Other taxa have

shown this shift in community composition with more dispersive species in the city, like carabid

beetles (Piano et al., 2017).

The high abundance of thermophilic ants in urban areas aligns with numerous studies

indicating that cities, such as Paris, are subject to the Urban Heat Island effect (Lemonsu et al.,

2015). Heat-susceptible species may disappear, and thermophilic species may dominate

communities in cities. Parr and Bishop (2022) suggest that temperate ant species may be less

vulnerable to climate warming than tropical species, and in some cases might even benefit from

rising temperatures. However, our findings—showing a near-complete replacement of forest ant

communities by species with more southern affinities in urban parks—do not fully support this idea.

On the contrary, although temperature is not the only factor differing between urban and rural

forests, our results suggest that our temperate ant communities living in forests may be negatively

impacted by rising temperatures. The prevalence of thermophilic species in urban areas has been

documented not only in ants but also in other groups, such as carabid beetles (Piano et al., 2017)

and moths (Franzén et al., 2020). More broadly, the process of thermophilisation, where

communities shift toward species that prefer warmer conditions due to rising temperatures, has also

been observed in non-urban ecosystems, including fish (Daufresne et al., 2004) and plants (Feeley et

al., 2020). This result is particularly relevant in the context of climate change, as temperatures are

expected to rise in the coming decades, leading to significant ecological consequences (McCarty,

2001). Cities offer a valuable natural laboratory for studying the impacts of climate change, with

urban communities serving as ideal experimental units to investigate these effects. Furthermore, our

findings suggest that climate change could act as a critical filter for community composition,

potentially leading to the decline of less thermophilic species. While urban areas might serve as

reservoirs for future biodiversity under changing climates, they will likely become increasingly

inhospitable for many species as climate change progresses.

References

References

Agosti, D., Majer, J. D., Alonso, L. E., & Schultz, T. R. (2000). *Ants: standard methods for measuring*
and monitoring biodiversity (Smithsonia).

Alberti, M. (2015). Eco-evolutionary dynamics in an urbanizing planet. *Trends in Ecology & Evolution*,
30(2), 114–126. https://doi.org/10.1016/j.tree.2014.11.007

Aronson, M. F. J., La Sorte, F. A., Nilon, C. H., Katti, M., Goddard, M. A., Lepczyk, C. A., Warren, P. S.,
Williams, N. S. G., Cilliers, S., Clarkson, B., Dobbs, C., Dolan, R., Hedblom, M., Klotz, S.,
Kooijmans, J. L., Kühn, I., MacGregor-Fors, I., McDonnell, M., Mörtberg, U., … Winter, M.
(2014). A global analysis of the impacts of urbanization on bird and plant diversity reveals key
anthropogenic drivers. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1780),
20133330. https://doi.org/10.1098/rspb.2013.3330

Baselga, A. (2010). Partitioning the turnover and nestedness components of beta diversity. *Global*
Ecology and Biogeography, *19*(1), 134–143. https://doi.org/10.1111/j.1466-8238.2009.00490.x

Baselga, A., & Orme, C. D. L. (2012). betapart: an R package for the study of beta diversity. *Methods*
in Ecology and Evolution, *3*(5), 808–812. https://doi.org/10.1111/j.2041-210X.2012.00224.x

Blatrix, R., Galkowski, C., Lebas, C., & Wegnez, P. (2013). *Fourmis de France* (Delachaux).

Bonte, D., Dyck, H. Van, Bullock, J. M., Delgado, M., Gibbs, M., Lehouck, V., Matthysen, E., Mustin, K.,
Schtickzelle, N., Stevens, V. M., Vandewoestijne, S., Baguette, M., Barton, K., Benton, T. G.,
Chaput-bardy, A., Dytham, C., Hovestadt, T., Meier, C. M., Steve, C. F., … Travis, J. M. J. (2012).
Costs of dispersal. *Biological Reviews*, *87*, 290–312. https://doi.org/10.1111/j.1469-
185X.2011.00201.x

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a metabolic
theory of ecology. *Ecology*, *85*(7), 1771–1789. https://doi.org/10.1890/03-9000

Buchholz, S., & Egerer, M. H. (2020). Functional ecology of wild bees in cities: towards a better
understanding of trait-urbanization relationships. *Biodiversity and Conservation*, *29*(9–10),
2779–2801. https://doi.org/10.1007/s10531-020-02003-8

Campbell-Staton, S. C., Velotta, J. P., & Winchell, K. M. (2021). Selection on adaptive and
maladaptive gene expression plasticity during thermal adaptation to urban heat islands. *Nature*
Communications, *12*(1), 6195. https://doi.org/10.1038/s41467-021-26334-4

Carpintero, S., & Reyes-López, J. (2014). Effect of park age, size, shape and isolation on ant
assemblages in two cities of Southern Spain. *Entomological Science*, *17*, 41–51.
https://doi.org/10.1111/ens.12027

Cheptou, P.-O., Carrue, O., Rouifed, S., & Cantarel, A. (2008). Rapid evolution of seed dispersal in an
urban environment in the weed *Crepis sancta*. *Proceedings of the National Academy of*
Sciences, *105*(10), 3796–3799. https://doi.org/10.1073/pnas.0708446105

Cote, J., Bestion, E., Jacob, S., Travis, J., Legrand, D., & Baguette, M. (2017). Evolution of dispersal
strategies and dispersal syndromes in fragmented landscapes. *Ecography*, *40*, 56–73.
https://doi.org/10.1111/ecog.02538

Croci, S., Butet, A., & Clergeau, P. (2008). Does urbanization filter birds on the basis of their
biological traits? *The Condor*, *110*(2), 223–240. https://doi.org/10.1525/cond.2008.8409

Cronin, A. L., Molet, M., Doums, C., Monnin, T., & Peeters, C. (2013). Recurrent Evolution of
Dependent Colony Foundation Across Eusocial Insects. *Annual Review of Entomology*, *58*(1),
37–55. https://doi.org/10.1146/annurev-ento-120811-153643

Daufresne, M., Roger, M. C., Capra, H., & Lamouroux, N. (2004). Long-term changes within the
invertebrate and fish communities of the Upper Rhône River: effects of climatic factors. *Global*
Change Biology, *10*(1), 124–140. https://doi.org/10.1046/j.1529-8817.2003.00720.x

Diamond, S. E., Chick, L. D., Perez, A., Strickler, S. A., & Martin, R. A. (2018). Evolution of thermal
tolerance and its fitness consequences: parallel and non-parallel responses to urban heat
islands across three cities. *Proceedings of the Royal Society B: Biological Sciences*, *285*(1882),
20180036. https://doi.org/10.1098/rspb.2018.0036

Diamond, S. E., & Martin, R. A. (2021). Evolution in Cities. *Annual Review of Ecology, Evolution, and*
Systematics, *52*(1), 519–540. https://doi.org/10.1146/annurev-ecolsys-012021-021402

Donoso, D. A. (2017). Tropical ant communities are in long-term equilibrium. *Ecological Indicators*,
83, 515–523. https://doi.org/10.1016/j.ecolind.2017.03.022

Fahrig, L. (2003). Effects of habitat fragmentation on biodiversity. *Annual Review of Ecology,*
Evolution, and Systematics, *34*, 487–515.
https://doi.org/10.1146/annurev.ecolsys.34.011802.132419

Feeley, K. J., Bravo-Avila, C., Fadrique, B., Perez, T. M., & Zuleta, D. (2020). Climate-driven changes in
the composition of New World plant communities. *Nature Climate Change*, *10*(10), 965–970.
https://doi.org/10.1038/s41558-020-0873-2

Finand, B., Loeuille, N., Bocquet, C., Fédérici, P., Ledamoisel, J., & Monnin, T. (2023). Habitat
fragmentation through urbanization selects for low dispersal in an ant species. *Oikos*.
https://doi.org/10.1111/oik.10325

Finand, B., Loeuille, N., Bocquet, C., Fédérici, P., & Monnin, T. (2024). Solitary foundation or colony
fission in ants: an intraspecific study shows that worker presence and number increase colony
foundation success. *Oecologia*, *204*(3), 517–527. https://doi.org/10.1007/s00442-023-05509-4

Finand, B., Monnin, T., & Loeuille, N. (2024). Evolution of dispersal and the maintenance of
fragmented metapopulations. *Oikos*, *2024*(2). https://doi.org/10.1111/oik.10490

Franzén, M., Betzholtz, P.-E., Pettersson, L. B., & Forsman, A. (2020). Urban moth communities
suggest that life in the city favours thermophilic multi-dimensional generalists. *Proceedings of*
the Royal Society B: Biological Sciences, *287*(1928), 20193014.
https://doi.org/10.1098/rspb.2019.3014

Gandon, S. (1999). Kin Competition, the Cost of Inbreeding and the Evolution of Dispersal. *Journal of*
Theoretical Biology, *200*(4), 345–364. https://doi.org/10.1006/jtbi.1999.0994

Guénard, B., Weiser, M. D., Gómez, K., Narula, N., & Economo, E. P. (2017). The Global Ant
Biodiversity Informatics (GABI) database: synthesizing data on the geographic distribution of
ant species (Hymenoptera: Formicidae). *Myrmecological News*, *24*, 83–89.
https://doi.org/https://doi.org/10.25849/myrmecol.news_024:083

Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., Lovejoy, T. E., Sexton,
J. O., Austin, M. P., Collins, C. D., Cook, W. M., Damschen, E. I., Ewers, R. M., Foster, B. L.,
Jenkins, C. N., King, A. J., Laurance, W. F., Levey, D. J., Margules, C. R., … Townshend, J. R.
(2015). Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances*,
1(2). https://doi.org/10.1126/sciadv.1500052

Hamilton, W. D., & May, R. M. (1977). Dispersal in stable habitats. *Nature*, *269*(5629), 578–581.
https://doi.org/10.1038/269578a0

Hantak, M. M., McLean, B. S., Li, D., & Guralnick, R. P. (2021). Mammalian body size is determined by
interactions between climate, urbanization, and ecological traits. *Communications Biology*,
4(1), 972. https://doi.org/10.1038/s42003-021-02505-3

Hartig, F. (2022). *DHARMa: Residual diagnostics for hierarchical (multi-Level / mixed) regression
models*. R package version 0.4.6. https://CRAN.R-project.org/package=DHARMa

Hastings, A. (1983). Can spatial variation alone lead to selection for dispersal? *Theoretical Population
Biology*, *24*, 244–251. https://doi.org/10.1016/0040-5809(83)90027-8

Herbers, J. M. (2011). Nineteen years of field data on ant communities (Hymenoptera: Formicidae):
What can we learn? *Myrmecological News*, *15*, 43–52.
https://doi.org/https://doi.org/10.25849/myrmecol.news_015:043

Hölker, F., Wolter, C., Perkin, E. K., & Tockner, K. (2010). Light pollution as a biodiversity threat.
Trends in Ecology & Evolution, *25*(12), 681–682. https://doi.org/10.1016/j.tree.2010.09.007

Hölldobler, B., & Wilson, E. O. (1990). *The ants* (Harvard University Press, Ed.).

Hubbell, S. P. (2001). *The Unified Neutral Theory of Biodiversity and Biogeography*. Princeton
University Press.

Janicki, J., Narula, N., Ziegler, M., Guénard, B., & Economo, E. P. (2016). Visualizing and interacting
with large-volume biodiversity data using client–server web-mapping applications: The design
and implementation of antmaps.org. *Ecological Informatics*, *32*, 185–193.
https://doi.org/10.1016/j.ecoinf.2016.02.006

Ješovnik, A., & Bujan, J. (2021). Wooded areas promote species richness in urban parks. *Urban
Ecosystems*, *24*(6), 1305–1315. https://doi.org/10.1007/s11252-021-01112-8

Johnson, M. T. J., & Munshi-South, J. (2017). Evolution of life in urban environments. *Science*,
358(6363), eaam8327. https://doi.org/10.1126/science.aam8327

Kaiser, A., Merckx, T., & Van Dyck, H. (2016). The Urban Heat Island and its spatial scale dependent
impact on survival and development in butterflies of different thermal sensitivity. *Ecology and
Evolution*, *6*(12), 4129–4140. https://doi.org/10.1002/ece3.2166

Keller, L. (1991). Queen number, mode of colony founding, and queen reproductive success in ants
(Hymenoptera Formicidae). *Ethology Ecology & Evolution*, *3*(4), 307–316.
https://doi.org/10.1080/08927014.1991.9525359

Keller, L. (1998). Queen lifespan and colony characteristics in ants and termites. *Insectes Sociaux*,
45(3), 235–246. https://doi.org/10.1007/s000400050084

Leal, I. R., Filgueiras, B. K. C., Gomes, J. P., Iannuzzi, L., & Andersen, A. N. (2012). Effects of habitat
fragmentation on ant richness and functional composition in Brazilian Atlantic forest.
Biodiversity and Conservation, *21*(7), 1687–1701. https://doi.org/10.1007/s10531-012-0271-9

Legendre, P., & Anderson, M. J. (1999). Distance-based redundancy analysis: testing multispecies
responses in multifactorial ecological experiments. *Ecological Monographs*, *69*(1), 1–24.
https://doi.org/https://doi.org/10.1890/0012-9615(1999)069[0001:DBRATM]2.0.CO;2

Leibold, M. A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J. M., Hoopes, M. F., Holt, R. D.,
Shurin, J. B., Law, R., Tilman, D., Loreau, M., & Gonzalez, A. (2004). The metacommunity
concept: a framework for multi-scale community ecology. *Ecology Letters*, *7*(7), 601–613.
https://doi.org/10.1111/j.1461-0248.2004.00608.x

Lemonsu, A., Viguié, V., Daniel, M., & Masson, V. (2015). Vulnerability to heat waves: Impact of
urban expansion scenarios on urban heat island and heat stress in Paris (France). *Urban
Climate*, *14*, 586–605. https://doi.org/10.1016/j.uclim.2015.10.007

Liu, K.-L., Peng, M.-H., Hung, Y.-C., & Neoh, K.-B. (2019). Effects of park size, peri-urban forest
spillover, and environmental filtering on diversity, structure, and morphology of ant
assemblages in urban park. *Urban Ecosystems*, *22*(4), 643–656.
https://doi.org/10.1007/s11252-019-00851-z

MacArthur, R. H., & Wilson, E. O. (1967). *The Theory of Island Biogeography*. Princeton University
Press. https://doi.org/10.1515/9781400881376

McCarty, J. P. (2001). Ecological Consequences of Recent Climate Change. *Conservation Biology*,
15(2), 320–331. https://doi.org/10.1046/j.1523-1739.2001.015002320.x

McGlynn, T. P., Meineke, E. K., Bahlai, C. A., Li, E., Hartop, E. A., Adams, B. J., & Brown, B. V. (2019).
Temperature accounts for the biodiversity of a hyperdiverse group of insects in urban Los
Angeles. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1912), 20191818.
https://doi.org/10.1098/rspb.2019.1818

Mouquet, N., & Loreau, M. (2003). Community patterns in source-sink metacommunities. *The
American Naturalist*, *162*(5), 544–557. https://doi.org/10.1086/378857

Nascimento, G., Câmara, T., & Arnan, X. (2022). Critical thermal limits in ants and their implications
under climate change. *Biological Reviews*, *97*(4), 1287–1305.
https://doi.org/10.1111/brv.12843

Newport, J., Shorthouse, D. J., & Manning, A. D. (2014). The effects of light and noise from urban
development on biodiversity: Implications for protected areas in Australia. *Ecological
Management & Restoration*, *15*(3), 204–214. https://doi.org/10.1111/emr.12120

Pacheco, R., & Vasconcelos, H. L. (2007). Invertebrate conservation in urban areas: Ants in the
Brazilian Cerrado. *Landscape and Urban Planning*, *81*(3), 193–199.
https://doi.org/10.1016/j.landurbplan.2006.11.004

Parr, C. L., & Bishop, T. R. (2022). The response of ants to climate change. *Global Change Biology*,
28(10), 3188–3205. https://doi.org/10.1111/gcb.16140

Piano, E., De Wolf, K., Bona, F., Bonte, D., Bowler, D. E., Isaia, M., Lens, L., Merckx, T., Mertens, D.,
van Kerckvoorde, M., De Meester, L., & Hendrickx, F. (2017). Urbanization drives community

shifts towards thermophilic and dispersive species at local and landscape scales. *Global Change*
Biology, *23*(7), 2554–2564. https://doi.org/10.1111/gcb.13606

Piano, E., Souffreau, C., Merckx, T., Baardsen, L. F., Backeljau, T., Bonte, D., Brans, K. I., Cours, M.,
Dahirel, M., Debortoli, N., Decaestecker, E., De Wolf, K., Engelen, J. M. T., Fontaneto, D.,
Gianuca, A. T., Govaert, L., Hanashiro, F. T. T., Higuti, J., Lens, L., … Hendrickx, F. (2020).
Urbanization drives cross-taxon declines in abundance and diversity at multiple spatial scales.
Global Change Biology, *26*(3), 1196–1211. https://doi.org/10.1111/gcb.14934

Preston, F. W. (1962). The Canonical Distribution of Commonness and Rarity: Part I. *Ecology*, *43*(2),
185. https://doi.org/10.2307/1931976

R Core Team. (2022). R: A language and environment for statistical computing. *R Foundation for*
Statistical Computing, Vienne, Au. https://www.R-project.org/

Ramalho, W. P., With, K. A., de Sousa Mesquita, G., de Arruda, F. V., Guerra, V., Ferraz, D., Andrade,
M. S., & do Prado, V. H. M. (2022). Habitat fragmentation rather than habitat amount or
habitat split reduces the diversity and abundance of ground-dwelling anurans within forest
remnants of the Brazilian Cerrado. *Journal for Nature Conservation*, *69*, 126259.
https://doi.org/10.1016/j.jnc.2022.126259

Roeder, K. A., Roeder, D. V, & Bujan, J. (2021). Ant Thermal Tolerance: A Review of Methods,
Hypotheses, and Sources of Variation. *Annals of the Entomological Society of America*, *114*(4),
459–469. https://doi.org/10.1093/aesa/saab018

Sanford, M. P., Manley, P. N., & Murphy, D. D. (2009). Effects of Urban Development on Ant
Communities: Implications for Ecosystem Services and Management. *Conservation Biology*,
23(1), 131–141. https://doi.org/10.1111/j.1523-1739.2008.01040.x

Seifert, B. (2018). *The Ants of Central and Northern Europe*. Lutra Verlags- und Vertriebsgesellschaft.

Slatkin, M. (1974). Hedging one's evolutionary bets. *Nature*, *250*(5469), 704–705.
https://doi.org/10.1038/250704b0

Steiner, C. F., & Asgari, M. (2022). Habitat isolation reduces intra- and interspecific biodiversity and
stability. *Royal Society Open Science*, *9*(2). https://doi.org/10.1098/rsos.211309

Sundström, L., Seppä, P., & Pamilo, P. (2005). Genetic population structure and dispersal patterns in
Formica ants — a review. *Ann. Zool. Fennici*, *42*, 163–177.

Theunis, L., Gilbert, M., Roisin, Y., & Leponce, M. (2005). Spatial structure of litter-dwelling ant
distribution in a subtropical dry forest. *Insectes Sociaux*, *52*, 366–377.
https://doi.org/10.1007/s00040-005-0822-0

Tilman, D. (1994). Competition and Biodiversity in Spatially Structured Habitats. *Ecology*, *75*(1), 2–16.
https://doi.org/10.2307/1939377

Tilman, D., May, R. M., Lehman, C. L., & Nowak, M. A. (1994). Habitat destruction and the extinction
debt. *Nature*, *371*(6492), 65–66. https://doi.org/10.1038/371065a0

Travis, J. M. J., Delgado, M., Bocedi, G., Baguette, M., Bartoń, K., Bonte, D., Boulangeat, I., Hodgson,
J. A., Kubisch, A., Penteriani, V., Saastamoinen, M., Stevens, V. M., & Bullock, J. M. (2013).
Dispersal and species' responses to climate change. *Oikos*, *122*, 1532–1540.
https://doi.org/10.1111/j.1600-0706.2013.00399.x

Wilson, C. J., & Jamieson, M. A. (2019). The effects of urbanization on bee communities depends on
floral resource availability and bee functional traits. *PLOS ONE*, *14*(12), e0225852.
https://doi.org/10.1371/journal.pone.0225852

Young, A., Boyle, T., & Brown, T. (1996). The population genetic consequences of habitat
fragmentation for plants. *Trends in Ecology & Evolution*, *11*(10), 413–418.
https://doi.org/10.1016/0169-5347(96)10045-8

683
684

Figure 1: Map of the sampling sites in forests around Paris (Ile-de-France region) (a) and parks in Paris (b). The colour of the circles indicates the fragmentation level of forests or the size of parks. minimally fragmented forests (n = 10) have between 75 and 95% of forest cover within 1 Km, medium fragmented forests (n = 8) have between 30 and 70% of forest cover, and highly fragmented forests (n = 6) have less than 5% of forest cover in their surroundings. Parks are divided

based on surface: large (80 000 to 250 000 m² (n = 8)), medium (5 000 to 12 000 m² (n = 9)), and

small (500-2 000 m² (n = 8)). Site information is detailed in Table S1.

Figure 2: Variations in the presence and abundance of ant species depending on patches. Darker

Levels show higher abundances, i.e., of the percentage of quadrats where the species was found. On

the Y-axis, green patches correspond to forests and grey patches correspond to parks. Red, orange,

and yellow show fragmentation levels for forests and patch sizes for parks. Species are ranked based

on their urbanisation preference along the X-axis.

687 **Figure 3:** Species richness (a-c) and Shannon index (d-f) between forests and parks (a, d), between

688 parks of different sizes (b, e), and between forests of different fragmentation levels (c, f). Significant

689 differences are observed in species richness and Shannon index between forests and parks (Species

690 richness: F value = 6.67, p = 0.013, R^2 = 10.6%; Shannon index: T value = 2.808, p = 0.007, R^2 = 13%),

691 and in species richness between parks of different sizes (F = 3.74, p = 0.039, R^2 = 0.19).

Figure 4: Beta diversity (db-RDA analysis) between forests and parks (a), between parks (b), and between forests (c). Blue acronyms are the species. Capital acronyms are the sites. For (c), yellow

represents high fragmentation, orange medium fragmentation, and red low fragmentation. Ant communities are markedly different between parks and forests, with communities from the two habitats being separated with no overlap (F value = 28.47, p = 0.001, R^2 = 37.7%). Communities also clearly differ between parks of different sizes, with a gradual change in composition from small to medium to large parks (F value = 2.14, p = 0.008, R^2 = 16.3%). No difference is observed between forests of different fragmentation level (F = 0.77, p = 0.69).

Figure 5: Relation between urbanisation preference of the species and their traits: colony mass (a), Social structure (b), Meridional index as a proxy of thermophily (c). Urbanisation preference varies between zero for strictly forest species to one for strictly urban species. It is correlated to the social structure (z value = -2.26, p = 0.02, R^2 = 0.15) and thermophily (t value = 2.89, p = 0.008, R^2 = 0.21), but not to the mass of the colony (t value = -0.76, p = 0.46).

692