Title: Immunodominant structural proteins Gc and N drive T cell-mediated protection

against La Crosse virus

Authors: Reem Alatrash[1,2], Bobby Brooke Herrera[1,2]

Affiliations:

[1]Rutgers Global Health Institute, Rutgers University, New Brunswick, NJ, USA

[2]Department of Medicine, Division of Allergy, Immunology, and Infectious Diseases and

Child Health Institute of New Jersey, Rutgers Robert Wood Johnson Medical School,

New Brunswick, NJ, USA

Key words: La Crosse, virus-specific T cells, immunogenicity, vaccines.

Abstract

La Crosse virus (LACV), a negative-sense RNA bunyavirus, is a major cause of pediatric encephalitis in the United States, disproportionately affecting children under the age of 16 years. Although LACV can lead to severe morbidity and mortality in this vulnerable pediatric population, no vaccines or specific antiviral therapies are currently available. Murine models recapitulate the age-dependent susceptibility observed in humans, whereby weanling mice (3 weeks old) succumb to LACV-induced neurological disease and adult mice (≥8 weeks old) exhibit resistance. In this study, we systematically characterized the T cell-mediated responses that underlie this differential susceptibility. We show that adult mice mount robust, polyfunctional LACV-specific CD4$^+$ and CD8$^+$ T cell responses against structural and non-structural proteins as early as 6 days post-infection (dpi), with sustained production of interferon-gamma (IFN-γ), granzyme B, interleukin-2 (IL-2), and tumor necrosis factor alpha (TNF-α). We further show that these T cells expand significantly in adult mice and display in vivo cytotoxicity against targeted cells pulsed with immunogenic antigens derived from the envelope Gc and nucleocapsid (N) proteins. In contrast, weanling mice exhibit significantly weaker T cell responses, as evidenced by reduced expansion of both CD4$^+$ and CD8$^+$ T cells and diminished cytokine production, correlating with 100% mortality by 7 dpi. Importantly, immunization studies with LFn-LACV-Gc and -N significantly improve in vivo T cell cytotoxicity and survival in weanlings, highlighting their potential as vaccine candidates. Collectively, our findings define key cellular correlates of protection against LACV and establish a framework for the development of pediatric vaccines designed to mitigate the public health burden of LACV-induced encephalitis.

Introduction

La Crosse virus (LACV) is a single-stranded, negative-sense RNA virus belonging to the *Bunyavirales* order (1). Its genome is segmented into three strands referred to as small (S), medium (M), and large (L), distinguished by their relative sizes. The S segment encodes the nucleocapsid protein (N) and nonstructural protein s (NSs), the latter of which has been shown to modulate the host cell antiviral response through innate immune pathways (2). The M segment encodes a glycosylated polyprotein precursor (GPC) that is cleaved by host cell proteases to generate the envelope glycoproteins Gn and Gc (3). Cleavage of GPC also produces the nonstructural protein m (NSm), which has been implicated in viral assembly (2). The L segment encodes an RNA-dependent RNA polymerase (RdRp), responsible for both transcription and replication of the viral genome.

Human infection with LACV occurs predominantly through the bite of infected Eastern tree hole mosquitoes (*Aedes triseriatus*), which are endemic throughout North America. Other mosquito species can also harbor LACV, potentially facilitating viral spread into new regions and hosts (4). Following entry into the human host, LACV replicates in skin muscle cells, enabling subsequent viral dissemination to multiple tissues and organs (5, 6). Although most LACV-infected individuals experience only mild, febrile symptoms, the virus can invade the central nervous system (CNS) causing a subset to experience severe neurological manifestations, including altered behavior, seizures, coma, or even death (7). Pediatric populations, particularly children under the age of 16, exhibit the highest susceptibility to LACV encephalitis, implicating age-associated factors in disease severity (8).

Currently, there are no approved antiviral therapies or vaccines against LACV,

necessitating reliance on supportive care alone to mitigate disease symptoms. The

absence of licensed vaccines partly reflects a limited understanding of how the host

immune response mediates protection, particularly against neurological sequelae. This

gap in knowledge has been difficult to address in humans, where clinical samples,

especially those collected during the acute phase of infection, are challenging to obtain.

However, murine models accurately recapitulate the age-dependent susceptibility seen

in humans, with weanling mice (3 weeks old) succumbing to LACV-induced

neuropathology, whereas adults (≥8 weeks old) resist peripheral LACV infection (5, 9,

10).

Previous studies in adult mice indicate a critical role of the peripheral immune

system in preventing LACV neuroinvasion, given that inoculation of C57BL/6 wild type

mice via intraperitoneal (IP) or intradermal (ID) routes does not induce overt

neurological disease, whereas intracranial inoculation does (6, 11-13). Additional

research reveals that type I interferon responses are indispensable for protecting mice

from lethal peripheral LACV challenge at all ages, highlighting the importance of innate

immunity in early antiviral control (12, 14). However, innate immune responses play a

dual role in LACV infection, offering both protective and pathogenic effects depending

on the tissue context (15) . In peripheral tissues, activation of antiviral recognition

pathways induces type I interferon responses, such as IFN-β, which suppress viral

replication, control LACV infection, and limit its spread to the CNS (14, 16, 17).

Conversely, similar signaling in the CNS can also lead to neuronal damage and cell

death (18). Myeloid dendritic cells, which bridge innate and adaptive immunity, are

critical for early protective responses in adult mice infected with LACV (12). Additionally,

treatment of young mice with type I interferon, particularly IFN-β, enhances protection

mediated by myeloid dendritic cells (12). Notably, both innate and adaptive immune

responses appear to cooperate in controlling LACV infection in adult mice. The

depletion of $CD4^+$ and $CD8^+$ T cells in adult mice reverse their resistance phenotype,

emphasizing that T cell-mediated responses are critical for protection and prevention of

neurological complications (11). This study also highlighted that while lymphocytes,

including NK cells and $CD4^+$ and $CD8^+$ T cells infiltrate the CNS of susceptible weanling

mice, their depletion does not impact neurological disease, suggesting they are not

contributors to pathogenesis in young mice (19). Conversely, in adult mice, the

depletion of both T cells and B cells results in increased viral loads in the brain and

heightened neurological disease (19). Despite this work, the precise immunological

correlates of protective immunity remain insufficiently characterized, particularly with

respect to identifying virus-specific T cell responses and their antigenic targets.

We recently demonstrated that LACV-infected weanling mice produce a Th2-

skewed cytokine profile and exhibit few IFN-γ producing T cells as compared to adults

(20). However, the passive transfer of splenic leukocytes from immune adult mice to

naïve weanlings prior to LACV challenge significantly improved survival (20). We also

recently showed that adult mice have delayed spread of LACV into the CNS as

compared to weanlings, further supporting adaptive immunity in peripheral control of the

virus (21). We hypothesize that T cell-mediated immunity is essential for protection

against LACV neuropathogenesis. Thus, a better understanding of age-dependent T

cell responses is pivotal for designing effective vaccines and immunotherapies that

target pediatric populations.

In this study, we provide a detailed analysis of T cell responses in adult and

weanling mice across the LACV proteome. We demonstrate robust, polyfunctional CD4$^+$

and CD8$^+$ T cell responses directed against both structural and non-structural proteins

in LACV-infected adult mice, whereas these responses are significantly blunted in

weanlings. We additionally show that upon immunization structural proteins Gc and N

elicit high in vivo cytotoxic activity in weanling mice that significantly improve survival.

These findings underscore the need for vaccine development targeting these

immunodominant antigens and highlight key immune determinants that could be

leveraged to protect children from LACV-induced encephalitis.

Results

Adult mice mount robust LACV-specific cellular response while weanling mice

lack these responses.

Previous studies have documented high immunogenicity of structural antigens in

bunyaviruses, but the immunogenicity of non-structural proteins has been less

thoroughly investigated (1). To identify LACV T cell targets, we performed in silico

peptide analysis of the entire LACV proteome, evaluating potential epitopes restricted

by both human and mouse MHC Class I and Class II molecules. Our analysis

highlighted multiple immunogenic peptides in structural and non-structural proteins. Out

of 176 total peptides identified as binders to MHC Class I molecules, an average of 12,

18, and 7 peptides were associated with Gn, Gc, and N, respectively. For the non-

structural proteins, we identified 5, 6, and 59 peptides binding to NSs, NSm, and RdRp,

respectively, suggesting that these proteins may also serve as dominant targets for T

cell-mediated immunity (Table 1, Supplementary 1). Among the peptides identified, Gc

and RdRp contained the highest number of highly immunogenic sequences

(Supplementary 1). Notably, 12 peptides were predicted to bind to both human and

mouse MHC Class I molecules, underscoring their potential as cross-species

immunogenic epitopes (Table 1). Based on these predictions, we proceeded with ex

vivo analyses using a panel of immunogens, consisting of a modified *Bacillus anthracis*

lethal factor (LFn) fused to each of the LACV structural or non-structural proteins. LFn

has been shown to transport full-length antigens into the cytosol to induce virus-specific

T cell responses via the MHC Class I and Class II pathways (22-25).

To quantify the LACV-specific cellular responses, we measured IFN-γ production

by splenic leukocytes from LACV-infected and uninfected C57BL/6 wild type mice in

enzyme-linked immunospot (ELISPOT) assays at 6 and 14 days post-infection. Adult

mice mounted a robust cellular response to multiple LACV proteins by 6 dpi, which

persisted by 14 dpi (Fig.1A, Supplementary Fig. 2 A-B). Structural proteins Gc and N

induced the strongest responses, followed by NSm as compared to the negligible

responses observed in uninfected adult controls (Fig. 1B, Supplementary Fig. 2 A-B). In

contrast, weanling mice at 6 dpi showed minimal IFN-γ responses to any LACV proteins,

which were similar to the uninfected controls (Fig. 1A, Supplementary. 3 A). Because all

weanling mice succumbed to infection by 7 dpi, data at 14 dpi could not be obtained.

These data indicate that the age-dependent immunological disparity in LACV infection

extends to both the magnitude and breadth of T cell responses.

Adult mice exhibit significantly higher percentage of both CD4$^+$ and CD8$^+$ splenic

T cells following LACV infection compared to weanlings

We next evaluated whether the limited LACV-specific cellular responses in

weanlings reflect a smaller pool of available T cells. Splenocytes from adult and

weanling mice were analyzed by flow cytometry at 6 and 14 dpi for the frequency of

CD4$^+$ and CD8$^+$ T cells. In adult mice, LACV infection triggered a significant increase in

CD4$^+$ T cell numbers at 6 dpi compared to age-matched uninfected controls at the same

timepoint (Fig. 2A). This population expansion persisted through 14 dpi. Similarly, CD8$^+$

T cells in adults exhibited a more pronounced expansion, reaching statistical

significance at both time points (Fig. 2B). By contrast, the percentage of CD4$^+$ and CD8$^+$

T cells in weanlings at 6 dpi remained unchanged from uninfected age-matched controls

and significantly lower percentages observed in infected adult mice at 6 dpi,

corresponding with their poor ex vivo IFN-γ response. These findings suggest that both

CD4$^+$ and CD8$^+$ T cell numbers expand in adults but not weanlings post LACV-

challenge.

Adult mice mount functional LACV-specific CD4$^+$ and CD8$^+$ T cell responses,

whereas weanling mice do not

To determine the functional capacity of virus-specific T cells in adult and

weanling mice following LACV challenge, we performed intracellular cytokine staining of

splenic leukocytes pulsed with LFn-LACV immunogens. We simultaneously measured

the production of IFN-γ, granzyme B, IL-2, and TNF-α in CD4$^+$ and CD8$^+$ T cells (Fig. 3,

Supplementary Fig. 4-8). Adult mice displayed robust IFN-γ responses across structural

and most non-structural proteins by 6 dpi. In adult mice, CD4$^+$ and CD8$^+$ T cells

displayed a robust IFN-γ response to all structural and most non-structural LACV

proteins as early as 6 dpi, with levels significantly higher than in age-matched

uninfected controls (Fig. 3 A-B). CD4$^+$ T cell responses began to decline by 14 dpi,

while CD8$^+$ T cell responses increased at 14 dpi against most LACV proteins, indicating

sustained effector activity of CD8$^+$ T cells (Fig. 3 A-B). Importantly, weanling mice

exhibited negligible cytokine responses by CD4$^+$ and CD8$^+$ T cell cells, mirroring their

poor splenic T cell expansion and heightened mortality (Fig. 3 C-D, Supplementary Fig.

7).

A similar trend was observed for other cytokines. Granzyme B expression was

significantly elevated in adult mice but was minimal in weanlings (Supplementary Fig. 4,

7). IL-2 and TNF-α responses were generally weak across both age groups, though

adult mice consistently showed higher levels than weanlings (Supplementary Fig. 5-7).

TNF-α expression was detectable in adults and weanlings but remained low overall

compared to other analyzed cytokines (Supplementary Fig. 6). These results indicate

that virus-specific T cells in adult mice mount functional effector responses during a

LACV infection, whereas such responses are notably weak or absent in weanling mice.

We further examined polyfunctionality by assessing the capacity of T cells to

concurrently produce multiple cytokines. In adult mice, ~20–34% of CD8$^+$ T cells co-

produced IFN-γ and granzyme B at 6 dpi, rising to 54-69% by 14 dpi. (Fig. 4 A-B). A

smaller proportion of CD8$^+$ T cells demonstrated dual functionality involving granzyme B

with IL-2 or IFN-γ with IL-2 (Fig. 4 A-B). These polyfunctional responses were most

pronounced following pulsing cells with the LFn-LACV Gc, Gn, or RdRp immunogens,

underscoring the immunogenicity of these proteins (Fig. 4 A-B, Supplementary Fig. 9).

CD4$^+$ T cells also exhibited multi-cytokine functionality, although IFN-γ was the

dominant effector molecule (Fig. 4 C-D). These CD4$^+$ T cell effector functions were

sustained at 14 dpi (Supplementary Fig. 9 C-D), underscoring their role in orchestrating

immune responses during LACV infection. Thus, in contrast to weanlings, adult mice

generated a robust, polyfunctional T cell response that likely contributes to effective viral

clearance and survival.

T cell responses against LACV Gc and N exhibit potent in vivo cytotoxicity and

confer protection in weanlings upon vaccination

We next assessed T cell cytotoxic activity using an in vivo cytotoxicity assay

based on carboxyfluorescein succinimidyl ester (CFSE). Briefly, CFSE-labeled

splenocytes pulsed with individual LFn-LACV proteins (LFn-LACV-Gc or -N) were

transferred into LACV-infected adult mice at 14 dpi, and LACV-infected weanlings at 6

dpi, and the percentage of target cell killing was calculated (Fig. 5A). In adult mice,

target cells pulsed with LFn-LACV-Gc or -N were eliminated at rates of ~53% and ~40%,

respectively (Fig. 5A, 5D). In contrast, weanling mice exhibited minimal killing of target

cells pulsed with LFn-LACV-Gc (~5%) or LFn-LACV-N (~7%) at 6 dpi, the latest

timepoint that could be assessed (Fig. 5A, D). These results show that LACV-specific T

cells mediate cytotoxicity in LACV-infected adult mice but not in weanlings.

To further translate these findings into a vaccination strategy for susceptible

weanling mice, we immunized neonatal mice (1 week old) with either LFn-LACV-Gc or -

N, with a boosting dose at 2 weeks old, then subsequently challenged them with LACV

at weanling age (Fig. 5B-C). The two-dose immunization regimen significantly improved

survival compared to control groups, increasing survival rates to ~70–80% (Fig. 5C). To

assess whether immunization impacted T cell cytotoxicity in weanlings, we repeated the

CFSE-based in vivo cytotoxicity assay post-immunization. CFSE-labeled splenocytes

pulsed with individual LFn-LACV proteins were transferred into their respective LFn-

LACV-Gc or -N immunized recipient mice, and cytotoxicity was measured. Strikingly,

LFn-LACV-Gc immunization enhanced target cell killing to ~21% in weanling mice, a

significant increase compared to the negligible cytotoxic response observed post-

infection. Similarly, LFn-LACV-N immunization improved target cell elimination to ~19%,

demonstrating a clear enhancement in cytotoxic capacity relative to unvaccinated

weanling mice (Fig. 5D). These results confirm that vaccine-elicited T cell responses

contribute to protection against LACV-induced disease in weanling mice by enhancing T

cell cytotoxic activity.

Discussion

In this study, we characterized virus-specific T cells in adult and weanling mice

providing a critical step toward developing a pediatric vaccine for this clinically

significant arbovirus. Our findings demonstrate significant differences in T cell

responses between adult and weanling mice. Adult mice mounted robust LACV-specific

T cell responses, characterized by strong IFN-γ production against both structural and

nonstructural antigens, with the highest responses targeting the structural proteins Gc

and N. In contrast, weanling mice displayed significantly weaker T cell responses at 6

dpi, underscoring the inability of their immature immune systems to mount effective

cellular immunity. This disparity likely contributes to the heightened susceptibility of

weanlings to severe LACV infection.

Further analysis revealed significant expansion of CD4$^+$ and CD8$^+$ T cell

populations in adult mice following LACV infection, accompanied by robust cytokine

production, including IFN-γ, granzyme B, IL-2, and TNF-α. Polyfunctional CD4$^+$ and

CD8$^+$ T cells, which co-produced cytokines especially IFN-γ and granzyme B, were

prominent in adults, supporting effective viral control and clearance. By contrast,

weanlings exhibited limited T cell expansion and functional responses, suggesting an

immature adaptive immune system unable to mount sufficient antiviral defenses.

Consistent with prior studies, our findings reaffirm the essential role of CD4$^+$ and

CD8$^+$ T cells in mediating protection in adult mice (19). Depletion or knockout of these

subsets led to disease progression and neurological complications at later stages of

LACV infection in adult mice, emphasizing the necessity of adaptive immunity in

sustaining the protective effects initially provided by innate responses (19). On the other

hand, the poor immune outcomes in weanlings may stem from deficiencies in both

innate and adaptive immunity in younger mice. Evidence suggests that dendritic cells in

young mice exhibit impaired functionality, potentially due to reduced maturity or lower

expression of co-stimulatory molecules (26). Additionally, a limited T cell receptor (TCR)

repertoire, immature germinal center reactions, and suboptimal T follicular helper (Tfh)

cell activity further compromise the adaptive immune response in weanlings (26).

Understanding these mechanisms is critical for informing vaccine strategies aimed at

enhancing immunity in vulnerable populations.

In our efforts to inform vaccine design, we explored the immunogenicity against

the entire LACV proteome. Our ex vivo analysis, which included T cell analysis against

both structural and nonstructural proteins, represents a significant advancement over

previous studies on members of *Bunyavirales* which focused only on structural antigens

(1). Immunogenic epitopes were identified across the proteome, providing a broad array

of potential T cell targets. The strong immunogenicity of the structural proteins Gc and N

was validated through their ability to elicit robust T cell responses with significant in vivo

cytotoxicity. Moreover, immunization with these proteins conferred protection in

weanling mice upon LACV challenge, demonstrating their potential as vaccine

candidates. These findings align with studies on genetically related viruses within the

Peribunyaviridae family, such as Jamestown Canyon Virus (JCV), Oropouche Virus

(OROV), and Bunyamwera Virus (BUNV), where glycoproteins (Gn/Gc) and

nucleocapsid (N) proteins were identified as key targets for T cell responses based

solely on immunoinformatic analysis (27-29). Similarly, comprehensive analyses of Rift

Valley Fever Virus (RVFV) and Severe Fever with Thrombocytopenia Syndrome Virus

(SFTSV) have highlighted the immunogenicity of structural and nonstructural proteins

(30, 31). Notably, cross-reactive N-specific CD8[+] T cells have been observed across

hantaviruses, and T cell responses targeting Gn and Gc have been identified in patients

infected with Hantaan virus (HTNV) and Andes Virus (ANDV) (32-34).

While the current study did not fully determine the specific mechanisms of

vaccine-induced protection, we have established a foundation for future research to

identify correlates of immunity for LFn-LACV-Gc or -N vaccines. These findings served

as a validation of the high immunogenicity of identified viral proteins in this study and

their protective potential in weanling mice. Notably, this work represents one of the few

attempts to develop a vaccine against LACV, building on early studies that

demonstrated protection using DNA vaccines encoding Gc/Gn and N proteins in adult

IFNAR1$^{-/-}$ mice (35). Mice vaccinated with DNA encoding the glycoproteins Gn and Gc

produced neutralizing antibodies and exhibited a high degree of protection against

challenge with high doses of LACV (35). Depletion of CD4$^+$ T cells in mice vaccinated

with DNA encoding Gn/Gc reduced their capacity to control the infection (35).

While most prior studies have utilized mouse models to investigate LACV

pathogenesis, our understanding of T cell responses in LACV-infected humans remains

limited. Surveillance data from LACV-endemic regions reveal that a majority of

individuals with detectable neutralizing antibodies do not report a history of clinically

significant infection (7). Among those who do develop encephalitis, clinical

manifestations range from mild aseptic meningitis to transient or permanent

neurological complications (36). This variability in susceptibility to infection and disease

severity suggests a potential role for MHC-restricted T cell responses. Supporting this

hypothesis, an earlier study examining MHC antigen frequencies in patients with LACV

encephalitis identified HLA-specific risk factors associated with disease, such as the

prevalence of HLA-B49 and/or the absence of HLA-DR5 (37). These findings

underscore the importance of T cell-mediated immunity in determining disease

susceptibility and severity in human LACV infections. Although there is a convenient

mouse model available for studying LACV, there is a critical need to understand human

HLA restriction and disease susceptibility and to characterize HLA-specific T cell

responses in both young and adult populations.

By emphasizing the age-dependent nature of T cell responses and identifying

immunogenic viral targets, this work provides a foundation for the development of LACV

vaccines and therapeutics. Future research should focus on addressing the mechanistic

basis of weak T cell responses in juvenile models, including deficits in antigen

presentation and T cell priming. Bridging these gaps will further inform vaccine

strategies to mitigate the burden of LACV disease, particularly in children, the

population most at risk.

Methods

Infection of mice with LACV and neurological disease progression criteria.

All animal studies were approved by the Rutgers IACUC (protocol number

202300007). Wild-type (C57Bl/6) mice were purchased from Jackson Laboratories and

maintained in a breeding colony at the Child Health Institute of New Jersey. LACV (La

Crosse Virus, NR-540), a human isolate was purchased and obtained from BEI

Resources. Mice at 3 (weanling) or 8 (adult) weeks of age were inoculated with 10^3 PFU

of LACV in phosphate-buffered saline (PBS) intraperitoneally in a volume of 200

µL/mouse. Mice were observed daily for signs of neurological disease that included

hunched posture, seizures, reluctance, or inability to move normally, or paralysis.

Animals with clear clinical signs of neurological disease were scored as clinical and

euthanized immediately.

Bioinformatic analyses.

All 9mer peptides encoded in the LACV proteome were predicted for their

binding affinity to MHC class I alleles A*0201, A*0101, A*1101, and B*0702. C57BL/6

MHC I was also included for analyses (H2Kb). Binding predictions are performed using

NetMHCpan 4.1 EL MHC class I prediction tool available on the Immune Epitope

Database (IEDB) (http://www.iedb.org) Web site (38). Peptides are selected if they were

in the top 1% of binders for each MHC molecule analyzed for further analysis for MHC I

immunogenicity available on the IEDB website. Immunogenic peptides were predicted

as previously described (39). Immunogenic peptides were assigned a positive

immunogenic score. A total of 176 9-mer peptides were identified by MHC class I

predictions, as described above.

LFn-LACV immunogens

LFn-LACV immunogens were purchased from Mir Biosciences, Inc., (New Jersey,

USA). In brief, gene fragments encoding the LACV Gc, Gn, N, NSm, NSs and RdRp

were cloned into the LFn expression vector. The constructs were transformed

into *Escherichia coli* BLR (DE3) (Millipore, Medford, MA). Selected transformants were

sequenced to verify in-frame cloning. The LFn-LACV immunogens were expressed

upon induction with isopropyl-β-d-thiogalactopyranoside (IPTG; Sigma-Aldrich, St. Louis,

MO, USA) in 5 liters of Luria broth containing Kanamycin for 4 h. Cells were pelleted by

centrifugation and resuspended in imidazole (1 mM) binding buffer (Novagen, Madison,

WI) in the presence of a protease inhibitor cocktail (Thermo Fisher Scientific, Waltham,

MA). Cell pellets were sonicated and centrifuged at 4°C, and the supernatants were

loaded in an equilibrated nickel-charged column for affinity purification. The bound

proteins were eluted in 100 to 200 mM imidazole, desalted with a Sephadex G-25M

column (Sigma-Aldrich, St. Louis, MO, USA), and eluted in phosphate-buffered saline

(PBS) (Sigma-Aldrich, St. Louis, MO, USA). The PBS-eluted proteins were passed

through Detoxi-Gel (Thermo Fisher Scientific, Rockford, IL). Protein concentrations

were determined, and samples were stored at −80°C.

Splenic leukocytes harvest

Spleens from LACV-infected and uninfected mice were harvested at 6 and 14 dpi,

as described previously (10). Spleens were processed into single-cell suspensions by

passing them through a 70 µM mesh filter, followed by red blood cell lysis using RBC

Lysis Buffer (Thermo Fisher Scientific). Isolated cells were then seeded for subsequent

ELISPOT and flow cytometry/ICS assays.

ELISPOT test.

Ex vivo ELISPOT tests ELISPOT assays were performed as previously

described (23). Briefly, 96-well plates (Millipore) are coated with anti-mouse IFN-γ

antibodies (BD Biosciences) and incubated overnight at 4°C. Plates are blocked with

RPMI 10% FBS and washed 2 times. Splenic leukocytes harvested from LACV infected

mice, as well as uninfected controls are seeded in triplicate at 2 x 10^5 cells/well. LFn-

LACV immunogens are added to each well (5 µg/ml) and plates are incubated at 37°C

with 5% CO2 for 18-24 hours. Cells stimulated with PMA (Phorbol 12-myristate 13-

acetate) will serve as a positive control and cells stimulated with LFn alone as a

negative control. After incubation and removal of cells, the detection antibody (BD

Biosciences) is added and incubated at room temperature. Plates are washed then

incubated with streptavidin HRP (BD Biosciences). After washing, AEC substrate (BD

Biosciences) is added before rinsing with water and air-drying. Digitized images are

analyzed using CTL ImmunoSpot reader (Cellular Technology Limited). LFn-LACV

immunogen specific spots is calculated by subtracting the mean of the negative control

values from the mean values of the specific stimulation. Positive responses are greater

than four times the mean background, three standard deviations above the background,

and >55 spot-forming cells (SFC)/ 2×10^5 total splenic leukocytes.

Flow cytometry and ICS assay.

Briefly, isolated cells are seeded at a density of $2 \times 10\square$ cells per well in a 96-well

U-bottom plate, with LFn-LACV immunogens added to each well at a concentration of

10 µg/ml, followed by incubation at 37°C with 5% CO_2 for 18 hours. Cells stimulated

with PMA will serve as a positive control and cells stimulated with LFn alone as a

negative control. Brefeldin A inhibitor is added during the last 4 hours of incubation.

Post-incubation, cells are stained for viability using a viability dye (LIVE/DEAD Fixable

Aqua Dead Cell Stain Kit, Thermo Fisher Scientific) and Fc receptors are blocked using

anti-mouse CD16/CD32 Fcγ III/II (Thermo fisher sciences) and normal mouse serum.

Cells are then stained for extracellular markers using a panel of antibodies (CD8a-

Alexa Fluor 488, CD19- Brilliant Violet 650, CD4- APC-eFluor 780, CD3- Alexa Fluor

700, CD45.2-PE-Cy7 (only for CNS sample); all from Thermo Fisher Scientific), washed,

and fixed and permeabilized using the FIX & PERM Cell Fixation and Permeabilization

Kit (Thermo Fisher Scientific). Following permeabilization, cells are stained for

intracellular cytokines (Granzyme B- PE-Cyanine 5.5, IL-2-PE, TNF-alpha- Brilliant

Violet 421, IFN-γ-eFluor 660; all from Thermo Fisher Scientific), washed, and

resuspended in FACS buffer. Flow cytometry data are acquired using the Aurora

system (Cytek® Biosciences) and analyzed using FlowJo software (version 10.2;

TreeStar, Inc), retaining live cells and excluding doublets using SSC-A and FSC-A

gating.

In vivo cytotoxicity assay.

Wild-type (C57Bl/6) mice (recipients) were either infected with 10^3 LACV, or

immunized with LACV-Gc or N. Splenocytes (targets) were harvested from donor naïve

wild-type mice. RBC were lysed, and the target cells were pulsed with 10 µg/ml of the

irrelevant LFn-immunogen (ZIKV), LFn-Gc, LFn-N for 18 h at 37°C. The cells were then

washed and labeled with Carboxyfluorescein succinimidyl ester (CFSE) (Invitrogen) in

PBS for 10 min at 37°C. LFn-LACV-pulsed cells were labeled with 5 µM CFSE

(CFSE^{high}) and the irrelevant protein-pulsed cells with 0.5 µM CFSE (CFSE^{low}). After

washing, the two cell populations were mixed at a 1:1 ratio and 5×10^6 cells were

injected i.v. into naive ,infected recipient mice at 14 dpi or immunized weanling mice.

After 4 h, the mice were sacrificed and splenocytes were analyzed by flow cytometry,

gating on lymphocyte cells using SSC-A and FSC-A gating. The percentage killing was

calculated as follows: 100 – ((percentage LACV peptide-pulsed in infected

mice/percentage irrelevant peptide-pulsed in infected mice)/(percentage LACV peptide-

pulsed in naive mice/percentage irrelevant peptide-pulsed in naive mice)) × 100.

LFn-LACV immunizations.

1 week old mice were immunized s.c. with 50 µg of each LFn-Gc or LFn-N

protein. Mock-immunized mice received LFn-ZIKV-NS3 protein. Mice were boosted with

the same dose after 7 days following the first immunization. Mice were then challenged

7 days after second dose immunization with 1000 PFU of LACV. Mice were monitored

for neurological signs development.

Statistical analysis

All statistical analyses were performed using Prism software Version 7.01 (GraphPad)

and are described in the figure legends.

Conflict of Interest

BBH is a co-founder of the company Mir Biosciences, Inc., which had no role in

the study. The remaining authors declare that the research was conducted in the

absence of any commercial or financial relationships that could be construed as a

potential conflict of interest.

Funding

The study was funded in part by New Jersey Health Foundation grant award

(IFHA17-24) to BBH. The funder had no role in the writing of the manuscript or decision

to publish.

References

1.   Alatrash R, Herrera BB. 2024. The Adaptive Immune Response against

Bunyavirales. Viruses 16:483.

2.   Leventhal SS, Wilson D, Feldmann H, Hawman DW. 2021. A Look into

Bunyavirales Genomes: Functions of Non-Structural (NS) Proteins. Viruses

13:314.

3.   Hulswit RJG, Paesen GC, Bowden TA, Shi X. 2021. Recent Advances in

Bunyavirus Glycoprotein Research: Precursor Processing, Receptor Binding and

Structure. Viruses 13:353.

4.   Westby KM, Fritzen C, Paulsen D, Poindexter S, Moncayo AC. 2015. La Crosse

Encephalitis Virus Infection in Field-Collected Aedes albopictus, Aedes japonicus,

and Aedes triseriatus in Tennessee. J Am Mosq Control Assoc 31:233-41.

5.   Bennett RS, Cress CM, Ward JM, Firestone CY, Murphy BR, Whitehead SS.

2008. La Crosse virus infectivity, pathogenesis, and immunogenicity in mice and

monkeys. Virol J 5:25.

6.   Schneider CA, Leung JM, Valenzuela-Leon PC, Golviznina NA, Toso EA,

Bosnakovski D, Kyba M, Calvo E, Peterson KE. 2024. Skin muscle is the initial

site of viral replication for arboviral bunyavirus infection. Nat Commun 15:1121.

7.   Haddow AD, Odoi A. 2009. The incidence risk, clustering, and clinical

presentation of La Crosse virus infections in the eastern United States, 2003-

2007. PLoS One 4:e6145.

8.   Balfour HH, Jr., Siem RA, Bauer H, Quie PG. 1973. California arbovirus (La

Crosse) infections. I. Clinical and laboratory findings in 66 children with

meningoencephalitis. Pediatrics 52:680-91.

9.   Hartman AL, Myler PJ. 2023. Bunyavirales: Scientific Gaps and Prototype

Pathogens for a Large and Diverse Group of Zoonotic Viruses. The Journal of

Infectious Diseases 228:S376-S389.

10.   López K, Wilson SN, Coutermash-Ott S, Tanelus M, Stone WB, Porier DL,

Auguste DI, Muller JA, Allicock OM, Paulson SL, Erasmus JH, Auguste AJ. 2021.

Novel murine models for studying Cache Valley virus pathogenesis and in utero

transmission. Emerg Microbes Infect 10:1649-1659.

11.   Winkler CW, Myers LM, Woods TA, Carmody AB, Taylor KG, Peterson KE. 2017.

Lymphocytes have a role in protection, but not in pathogenesis, during La Crosse

Virus infection in mice. J Neuroinflammation 14:62.

12.   Taylor KG, Woods TA, Winkler CW, Carmody AB, Peterson KE. 2014. Age-

dependent myeloid dendritic cell responses mediate resistance to la crosse virus-

induced neurological disease. J Virol 88:11070-9.

13.   Basu R, Nair V, Winkler CW, Woods TA, Fraser IDC, Peterson KE. 2021. Age

influences susceptibility of brain capillary endothelial cells to La Crosse virus

infection and cell death. Journal of Neuroinflammation 18:125.

14.   Hefti HP, Frese M, Landis H, Di Paolo C, Aguzzi A, Haller O, Pavlovic J. 1999.

Human MxA protein protects mice lacking a functional alpha/beta interferon

system against La crosse virus and other lethal viral infections. J Virol 73:6984-

91.

15.   Taylor KG, Peterson KE. 2014. Innate immune response to La Crosse virus

493   infection. Journal of NeuroVirology 20:150-156.

494   16.   Blakqori G, Delhaye S, Habjan M, Blair CD, Sánchez-Vargas I, Olson KE,

495   Attarzadeh-Yazdi G, Fragkoudis R, Kohl A, Kalinke U, Weiss S, Michiels T,

496   Staeheli P, Weber F. 2007. La Crosse bunyavirus nonstructural protein NSs

497   serves to suppress the type I interferon system of mammalian hosts. J Virol

498   81:4991-9.

499   17.   Pavlovic J, Schultz J, Hefti HP, Schuh T, Mölling K. 2000. DNA vaccination

500   against La Crosse virus. Intervirology 43:312-21.

501   18.   Mukherjee P, Woods TA, Moore RA, Peterson KE. 2013. Activation of the innate

502   signaling molecule MAVS by bunyavirus infection upregulates the adaptor protein

503   SARM1, leading to neuronal death. Immunity 38:705-16.

504   19.   Winkler CW, Myers LM, Woods TA, Carmody AB, Taylor KG, Peterson KE. 2017.

505   Lymphocytes have a role in protection, but not in pathogenesis, during La Crosse

506   Virus infection in mice. Journal of Neuroinflammation 14:62.

507   20.   Alatrash R, Vaidya V, Herrera BB. 2024. Age-specific dynamics of neutralizing

508   antibodies, cytokines, and chemokines in response to La Crosse virus infection in

509   mice. J Virol 98:e0176224.

510   21.   Lumkong L, Alatrash R, Sridhar S, Herrera BB. 2025. Development of an RT-

511   RPA assay for La Crosse virus detection provides insights into age-dependent

512   neuroinvasion in mice. bioRxiv

513   doi:10.1101/2025.01.21.634131:2025.01.21.634131.

514   22.   Herrera BB, Tsai WY, Chang CA, Hamel DJ, Wang WK, Lu Y, Mboup S, Kanki

515   PJ. 2018. Sustained Specific and Cross-Reactive T Cell Responses to Zika and

516   Dengue Virus NS3 in West Africa. J Virol 92.

517   23.   Herrera BB, Tsai WY, Brites C, Luz E, Pedroso C, Drexler JF, Wang WK, Kanki

518   PJ. 2018. T Cell Responses to Nonstructural Protein 3 Distinguish Infections by

519   Dengue and Zika Viruses. mBio 9.

520   24.   Herrera BB, Hamel DJ, Oshun P, Akinsola R, Akanmu AS, Chang CA, Eromon P,

521   Folarin O, Adeyemi KT, Happi CT, Lu Y, Ogunsola F, Kanki PJ. 2018. A modified

522   anthrax toxin-based enzyme-linked immunospot assay reveals robust T cell

523   responses in symptomatic and asymptomatic Ebola virus exposed individuals.

524   PLoS Negl Trop Dis 12:e0006530.

525   25.   Akanmu S, Herrera BB, Chaplin B, Ogunsola S, Osibogun A, Onawoga F, John-

526   Olabode S, Akase IE, Nwosu A, Hamel DJ, Chang CA, Kanki PJ. 2023. High

527   SARS-CoV-2 seroprevalence in Lagos, Nigeria with robust antibody and cellular

528   immune responses. J Clin Virol Plus 3:100156.

529   26.   Dakic A, Shao Q-x, D'Amico A, O'Keeffe M, Chen W-f, Shortman K, Wu L. 2004.

530   Development of the Dendritic Cell System during Mouse Ontogeny1. The Journal

531   of Immunology 172:1018-1027.

532   27.   Nelluri KDD, Ammulu MA, Durga ML, Sravani M, Kumar VP, Poda S. 2022. In

533   silico multi-epitope Bunyumwera virus vaccine to target virus nucleocapsid N

534   protein. Journal of Genetic Engineering and Biotechnology 20:89.

535   28.   Shahab M, Aiman S, Alshammari A, Alasmari AF, Alharbi M, Khan A, Wei D-Q,

536   Zheng G. 2023. Immunoinformatics-based potential multi-peptide vaccine

designing against Jamestown Canyon Virus (JCV) capable of eliciting cellular

and humoral immune responses. International Journal of Biological

Macromolecules 253:126678.

29. Adhikari UK, Tayebi M, Rahman MM. 2018. Immunoinformatics Approach for

Epitope-Based Peptide Vaccine Design and Active Site Prediction against

Polyprotein of Emerging Oropouche Virus. Journal of Immunology Research

2018:6718083.

30. Adhikari UK, Rahman MM. 2017. Overlapping CD8+ and CD4+ T-cell epitopes

identification for the progression of epitope-based peptide vaccine from

nucleocapsid and glycoprotein of emerging Rift Valley fever virus using

immunoinformatics approach. Infection, Genetics and Evolution 56:75-91.

31. Suleman M, Asad U, Arshad S, Rahman Au, Akbar F, Khan H, Hussain Z, Ali SS,

Mohammad A, Khan A, Ali L, Wei D-Q. 2022. Screening of immune epitope in

the proteome of the Dabie bandavirus, SFTS, to design a protein-specific and

proteome-wide vaccine for immune response instigation using an

immunoinformatics approaches. Computers in Biology and Medicine 148:105893.

32. Maeda K, West K, Toyosaki-Maeda T, Rothman AL, Ennis FA, Terajima M. 2004.

Identification and analysis for cross-reactivity among hantaviruses of H-2b-

restricted cytotoxic T-lymphocyte epitopes in Sin Nombre virus nucleocapsid

protein. Journal of General Virology 85:1909-1919.

33. Wang M, Zhu Y, Wang J, Lv T, Jin B. 2011. Identification of Three Novel CTL

Epitopes within Nucleocapsid Protein of Hantaan Virus. Viral Immunology

24:449-454.

34.   Wang M, Wang J, Kang Z, Zhao Q, Wang X, Hui L. 2015. Kinetics and

561   Immunodominance of Virus-Specific T Cell Responses During Hantaan Virus

562   Infection. Viral Immunology 28:265-271.

563   35.   Schuh T, Schultz J, Moelling K, Pavlovic J. 1999. DNA-Based Vaccine against La

564   Crosse Virus: Protective Immune Response Mediated by Neutralizing Antibodies

565   and CD4+ T Cells. Human Gene Therapy 10:1649-1658.

566   36.   Boutzoukas AE, Freedman DA, Koterba C, Hunt GW, Mack K, Cass J, Yildiz VO,

567   de Los Reyes E, Twanow J, Chung MG, Ouellette CP. 2023. La Crosse Virus

568   Neuroinvasive Disease in Children: A Contemporary Analysis of

569   Clinical/Neurobehavioral Outcomes and Predictors of Disease Severity. Clin

570   Infect Dis 76:e1114-e1122.

571   37.   Case KL, West RM, Smith MJ. 1993. Histocompatibility antigens and La Crosse

572   encephalitis. J Infect Dis 168:358-60.

573   38.   Lundegaard C, Lamberth K, Harndahl M, Buus S, Lund O, Nielsen M. 2008.

574   NetMHC-3.0: accurate web accessible predictions of human, mouse and monkey

575   MHC class I affinities for peptides of length 8-11. Nucleic Acids Res 36:W509-12.

576   39.   Calis JJ, Maybeno M, Greenbaum JA, Weiskopf D, De Silva AD, Sette A, Keşmir

577   C, Peters B. 2013. Properties of MHC class I presented peptides that enhance

578   immunogenicity. PLoS Comput Biol 9:e1003266.

581   **Tables**

583   **Table 1. Predicted immunogenic LACV derived epitopes with human and/or MHC**

584   **restriction.**

	Restriction

Epitope	Sequence	HLA A*01:01	HLA A*02:01	HLA A*11:01	HLA B*07:02	H2Kb
Gc$_{1-9}$	YTNIATQLK	+		+		
Gc$_{69-78}$	YSQTTWRVY	+	+			+
Gc$_{156-164}$	KSNDDLIAY	+		+		
Gc-$_{180-189}$	AIIDYIAYM	+	+	+		+
Gc$_{249-258}$	LSPRSGAIY	+			+	
Gc$_{633-642}$	YSLPRIVAV		+		+	+
N$_{335-344}$	FLQKFGIRL		+			+
NSs$_{49-58}$	VSQRGRLTL		+		+	+
NSs$_{71-80}$	FLETGTTQL	+	+			
NSm$_{13-24}$	SIITAVLVL		+			+
NSm$_{73-82}$	VIIGLLIGL		+			+
RdRp$_{711-720}$	LSDYDITQK	+		+		
RdRp$_{910-919}$	KVFDRLYEL	+	+	+	+	+
RdRp$_{1065-1074}$	SAQDVFYKY	+		+		
RdRp$_{1084-1093}$	YPQEKERILY	+			+	
RdRp$_{1166-1175}$	ILKEAITLL	+	+			+
RdRp$_{1456-1465}$	GMVELFTYL	+	+			+
RdRp$_{1860-1869}$	SINLTITGY	+		+		
RdRp$_{1960-1969}$	RIYNEITPV	+	+			+
RdRp$_{2092-3001}$	STPIFNIYY	+		+	+	+

Figure Legends

Figure 1. LACV-specific ex vivo cellular response in wildtype mice. (A) Top:
Quantified spot forming cells (SFC) detected by ELISPOT assay using splenic
leukocytes from adult and weanling mice infected with LACV at 6 days post-infection
(dpi). Bottom: Splenic leukocytes were treated with LFn-LACV proteins and IFN-γ
responses were detected by ELISPOT assay. (-) negative control: cells stimulated with
LFn alone, (+) positive control: cells stimulated with PMA (phorbol 12-myristate 13-

acetate). Statistical significance was determined using ANOVA test. **(B)** Percent of total
responses generated against each LFn-LACV protein in adult mice at 14 dpi.

Figure 2. LACV-induced CD4$^+$ and CD8$^+$ T cell Expansion in LACV infected mice.
Total percentage of CD4$^+$ T cells in **(A)** and CD8$^+$ T cell populations (**B**) among total
splenocytes of adult and weanling mice at 6 and 14 days post-infection (dpi) with LACV
analyzed by Flow cytometry. Statistical significance was determined using ANOVA test,
*p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

Figure 3. LACV-specific ex vivo splenic CD4$^+$ and CD8$^+$ T cell Functional response
in LACV infected mice. Functional LACV-specific CD4$^+$ and CD8$^+$ T cell responses
shown as the percent of CD4$^+$ or CD8$^+$ cells positive for IFN☐ in CD3$^+$ gate in FC/ICS in
adult **(A, B)** and weanling mice **(C, D)** at 6 , 14 dpi compared to age-matched uninfected
controls. Statistical significance was determined using ANOVA test, *p < 0.05, **p <
0.01, ***p < 0.001, ****p < 0.0001.

Figure 4. Polyfunctional profiles of LACV-specific CD8$^+$ and CD4$^+$ T cells at 6 dpi.
Pie charts depicting the distribution of LACV-specific CD8$^+$ T cell **(A)** and CD4$^+$ T cell
(C) populations with single or dual functional profiles, defined by cytokine production**. (B,**
D) Bar graph showing the total percentage of LACV-specific CD8$^+$ T or CD4$^+$ T cells
producing one or two cytokines in response to infection.

Figure 5. In vivo cytotoxicity and protection mediated by LACV-specific T cells.
(A) Schematic representation of the in vivo cytotoxicity assay. Equal numbers of naïve
splenocytes pulsed with LFn-LACV-Gc or LFn-LACV-N (CFSE^{hi} target cells) and
splenocytes pulsed with an irrelevant protein, LFn-ZIKV-NS3 (CFSE^{lo} marker cells),
were transferred into either naïve or LACV-immune recipient mice. Percent killing was
quantified by flow cytometry in recipient spleens 4 hours post-transfer. **(B)** Schematic of
the LFn-LACV immunization strategy. One-week-old mice were immunized with 50 µg
of either LFn-LACV-Gc or LFn-LACV-N. A booster dose was administered 7 days later,
followed by LACV challenge at weanling age. **(C)** Survival curves of weanling mice
immunized with LFn-LACV-Gc or LFn-LACV-N compared with LFn-ZIKV-NS3 and
subsequently challenged with LACV (n = 5 per group). **(D)** Bar chart quantifying in vivo
cytotoxicity in previously LACV-infected adult and weanling mice, as well as in LFn-
LACV-Gc- or LFn-LACV-N-immunized weanling mice. Percent killing was determined by
flow cytometry.

Figures

655   **Figure. 1. LACV-specific ex vivo cellular response in wildtype mice.**

671 **Figure. 2. LACV-induced CD4+ and CD8+ T cell Expansion in LACV infected mice.**

688   **Figure. 3. LACV-specific ex vivo splenic CD4[+] and CD8[+] T cell Functional**

689   **response in LACV infected mice.**

Figure. 4. Polyfunctional profiles of LACV-specific CD8+ and CD4+ T cells at 6 dpi.

716   **Figure. 5. In vivo cytotoxicity and protection mediated by LACV-specific T cells.**

Supplementary Figure Legends

Supplementary Figure. 1. In silico prediction of immunogenic peptides within LACV proteome with HLA-restriction. MHC I immunogenicity score of top 1% binders to **(A)** HLA A*0101, **(B)** HLA A*0201, **(C)** HLA A*1101 and **(D)** HLA B*0702. Number of top 1% of binders and percent immunogenic peptides within each LACV protein is indicated as total counts and counts with positive value of MHC I immunogenicity score respectively. **(F)** MHC II immunogenicity score of peptides predicted to bind using 7-allele method.

Supplementary Figure. 2. LACV-specific ex vivo cellular response in adult wildtype mice. Top: Quantified spot forming cells (SFC) detected by ELISPOT assay using splenic leukocytes from adult mice infected with LACV at 6 dpi **(A)** and 14 dpi **(B)** compared to age-matched non-infected controls. Bottom: Splenic leukocytes were treated with LFn-LACV proteins and IFN-γ responses were detected by ELISPOT assay. (-) negative control: cells stimulated with LFn alone, (+) positive control: cells stimulated with PMA (phorbol 12-myristate 13-acetate). Statistical significance was determined using ANOVA or t-test. **(C)** Percent of total responses generated against each LFn-LACV protein in adult mice at 6 dpi.

Supplementary Figure. 3. LACV-specific ex vivo cellular response in weanling wildtype mice. Top: Quantified spot forming cells (SFC) detected by ELISPOT assay using splenic leukocytes from weanling mice infected with LACV at 6 dpi compared to age-matched non-infected controls. Bottom: Splenic leukocytes were treated with LFn-LACV proteins and IFN-γ responses were detected by ELISPOT assay. (-) negative control: cells stimulated with LFn alone, (+) positive control: cells stimulated with PMA (phorbol 12-myristate 13-acetate). Statistical significance was determined using ANOVA or t-test. **(B)** Percent of total responses generated against each LFn-LACV protein in weanling mice at 6 dpi.

Supplementary Figure. 4. LACV-specific ex vivo splenic CD4$^+$ and CD8$^+$ T cell Functional response in LACV infected mice. Functional LACV-specific CD4+ and CD8+ T cell responses shown as the percent of CD4 or CD8 cells positive for Granzyme B in CD3+ gate in FC/ICS in adult (A, B) and weanling mice (C, D) at 6, 14 dpi compared to age-matched uninfected controls. Statistical significance was determined using ANOVA test, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

Supplementary Figure. 5. LACV-specific ex vivo splenic CD4⁺ and CD8⁺ T cell
Functional response in LACV infected mice. Functional LACV-specific CD4+ and
CD8+ T cell responses shown as the percent of CD4 or CD8 cells positive for IL-2 in
CD3+ gate in FC/ICS in adult (A, B) and weanling mice (C, D) at 6, 14 dpi compared to
age-matched uninfected controls. Statistical significance was determined using ANOVA
test, $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$, $^{****}p < 0.0001$.

Supplementary Figure. 6. LACV-specific ex vivo splenic CD4⁺ and CD8⁺ T cell
Functional response in LACV infected mice. Functional LACV-specific CD4+ and
CD8+ T cell responses shown as the percent of CD4 or CD8 cells positive for TNF-□ in
CD3+ gate in FC/ICS in adult (A, B) and weanling mice (C, D) at 6, 14 dpi compared to
age-matched uninfected controls. Statistical significance was determined using ANOVA
test, $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$, $^{****}p < 0.0001$.

Supplementary Figure. 7. Functional differences in virus-specific CD4⁺ and CD8⁺
T cells between adult and weanling mice at 6 dpi. Heatmap illustrating p-values
derived from the Mann-Whitney U test comparing the percent of virus-specific CD4⁺ **(A)**
and CD8⁺ T **(B)** cells producing individual cytokines analyzed in FC/ICS assay in adult
versus weanling mice. Columns correspond to the functional markers analyzed, and
rows indicate T cell specificity. Darker colors indicate lower p value.

Supplementary Figure. 8. Representation of gating strategy for intracellular
staining quantification of LACV-specific CD8⁺ Functional responses in adult
wildtype mice using ICS/Flow cytometry. Gating for cytokine-positive populations is
shown as an example for infected and uninfected adult mice at 14 dpi.

Supplementary Figure. 9. Polyfunctional profiles of LACV-specific CD8⁺ and CD4⁺ T
cells at 14 dpi. Pie charts depicting the distribution of LACV-specific CD8⁺ T cell **(A)** and
CD4+ T cell **(C)** populations with single or dual functional profiles, defined by cytokine
production. **(B, D)** Bar graph showing the total percentage of LACV-specific CD8⁺ T or
CD4+ T cells producing one or two cytokines in response to infection.

Supplementary Figures

Supplementary Figure. 1. In silico prediction of immunogenic peptides within LACV proteome with HLA-restriction.

Supplementary Figure. 2. LACV-specific ex vivo cellular response in adult

wildtype mice.

838 **Supplementary Figure. 3. LACV-specific ex vivo cellular response in weanling**
839 **wildtype mice.**

Supplementary Figure. 4. LACV-specific ex vivo splenic CD4$^+$ and CD8$^+$ T cell Functional response in LACV infected mice.

888 **Supplementary Figure. 6. LACV-specific ex vivo splenic CD4$^+$ and CD8+ T cell**
889 **Functional response in LACV infected mice.**

904 **Supplementary Figure. 7. Functional differences in virus-specific CD4$^+$ and CD8$^+$**

905 **T cells between adult and weanling mice at 6 dpi.**

925 **Supplementary Figure 8. Representation of gating strategy for intracellular**
926 **staining quantification of LACV-specific CD8+ Functional responses in adult**
927 **wildtype mice using ICS/Flow cytometry.**

**Supplementary Figure 9. Polyfunctional profiles of LACV-specific CD8+ and CD4+
T cells at 14 dpi.**