Historic mine waste contains diverse microbial communities that reflect waste type

and geochemistry

Mackenzie B. Best[1], Zohreh Kazemi Motlagh[2], Virginia T. McLemore[3], and Daniel S. Jones[1,4]

[1]Department of Earth and Environmental Science, New Mexico Institute of Mining and

Technology, Socorro, New Mexico, USA

[2]Department of Mineral Engineering, New Mexico Institute of Mining and Technology,

Socorro, New Mexico, USA

[3]New Mexico Bureau of Geology and Mineral Resources, New Mexico Institute of Mining

and Technology, Socorro, New Mexico, USA

[4]National Cave and Karst Research Institute, Carlsbad, New Mexico, USA

Correspondence: mackenzie.best@student.nmt.edu, daniel.s.jones@nmt.edu

ABSTRACT

Waste rock and tailings left behind by historic mining operations can contain substantial critical mineral resources. However, over the decades and centuries since these deposits were emplaced, microbial communities developed that can catalyze rock weathering and elemental cycling, which could have impacted the economic resources but also might be harnessed for future biomining or other metal recovery efforts. Here we combined cell counting, rRNA gene and transcript sequencing, and whole rock geochemistry to compare the composition and abundance of microbial communities from five inactive mine sites in South-Central New Mexico that contain critical minerals. While acidic seeps and adits at the sites contained organisms commonly found in acid rock drainage and bioleaching operations, these organisms were only present at very low abundance in the waste rock and tailings, which were instead dominated by bacteria and archaea that mostly represent inorganic nitrogen- and organic carbon-oxidizing populations. Generally, rRNA transcript libraries contained many of the same organisms as rRNA gene libraries, indicating that most of these microbial populations are active. Differences among total and active communities correspond to waste rock geochemistry, including concentrations of sulfur, iron, and other variables such as copper, lead, and rare earth elements. Nevertheless, many of the rRNA gene and transcript sequences in these deposits were from groups without cultured representatives, and these unknown microorganisms are likely important for biogeochemical cycling over the long lifetime of these waste deposits. We also discuss recommendations for microbiological assessment of similar large historic mine waste deposits.

IMPORTANCE

New Mexico has a long history of mining, with hundreds of mining districts across the state, many of which contain inactive operations with historic tailings and waste rock. Because metallurgical processing was in its infancy when most of these mines were active, they

contain substantial metal resources in tailings and waste rock that could be used to

support domestic demand for critical minerals. We found that microbial communities

associated with these deposits do not represent typical bioleaching communities, and

instead are dominated by taxa not typically associated with mine wastes. However, the

deposits do contain rare iron and sulfur-cycling taxa that could catalyze metal mobilization,

as well as active populations of novel microorganisms that are likely important for

biogeochemical cycling. These mine waste-associated microbial communities could

represent important resources for bioremediation and other biotechnological applications

to recover valuable elements from these and other historic mine waste deposits.

INTRODUCTION

Many of the low-carbon technologies at the heart of energy-efficient technologies are more metal and mineral intensive than traditional fossil fuel-based infrastructure, resulting in a huge increase in global demand and price for these metals and mineral commodities (1). These non-fuel commodities have been termed 'critical minerals' by the U.S. Geological Survey (USGS) and Department of Energy (DOE) due to their importance in the development of new technologies and large-scale infrastructure projects (2–4). However, many of the high-grade shallow ore deposits we have historically relied on to supply these minerals are dwindling, with some element reserves projected to be completely exhausted within 50-100 years (5). The result is that this increase in demand must be met with deeper, lower grade deposits (6) and has prompted a search for non-traditional repositories of critical minerals such as recycling pre-existing metal resources (7, 8) and reevaluating mineral deposits and mine wastes (9–16). The reliance on imported critical minerals means the supply chain is highly susceptible to disruption due to import costs as well as geopolitical factors such as trade negotiations and agreements (17). Given this vulnerability, there is a push to locate and quantify critical minerals within the US at both active and inactive mine sites to better support domestic demand. New Mexico has a long history with mining and contains over 9000 inactive mines, many of which contain waste material in the form of bulk waste rock or tailings that were mined from precious- and base- metal deposits that have elevated concentrations of critical minerals (9, 18). These inactive mines vary widely in age and production history, and were processed exclusively for select base and precious metals at a time when metallurgical processing was in its infancy, meaning that they could contain substantial critical mineral resources within these wastes.

Microorganisms matter for each step in the mining lifecycle due to their capacity to enhance mineral weathering and mobilize and precipitate metals (19–23). When sulfide-rich minerals are exposed to the surface, microbially-catalyzed redox reactions can lead to the formation of acid rock drainage (ARD) or acid mine drainage (AMD) (24–35). Biological

or abiotic strategies to prevent or limit sulfide mineral-oxidizing populations can help

prevent AMD and secure mine wastes (36, 37), and acidophilic populations can be used to

immobilize and even recover metals using active and passive remediation strategies such

as low pH iron oxidation and anaerobic sulfate reduction (38–43).

Microorganisms can also be used to extract metals during active mining in

'bioleaching' or 'biooxidation' processes that are collectively known as "biomining" (44–

50). The ability of microbes to oxidize metal sulfide and other minerals by producing

oxidants, acids, and lixiviants, either during direct mineral contact or in a leaching solution

(51–53) (54, 55), lowers the processing cost, which has made it easier for companies to

process non-traditional metal resources such as refractory and low-grade ores (49, 52, 56,

57), electronic waste (58–60), and mine waste (61–63). Currently, approximately 15% of

global copper, as well as smaller amounts of gold and other metals, is extracted using

microorganisms (64, 65). Microbes have also been evaluated for in situ leaching concepts,

in which the rock is first fractured and treated with a biologically-regenerated leach

solution (e.g., (65–67)). Although mostly untested, mine wastes are potential targets for in

situ bioleaching because it is often relatively permeable (68, 69) and contains leftover

metals. Small-scale studies have shown that biomining process does work with tailings

(61, 62).

Mine wastes can be defined as any product from a mine that is left over once mining

has ceased. This can include bulk waste rock and tailings. Bulk waste rock ranges in size

from pebble to boulder and includes materials that were above or within the ores

(overburden and gangue, respectively) and was excavated and stored in waste piles on the

surface. Bulk waste rock has not been processed for metal extraction, and may contain

variable metal content (70–72). Tailings represent material that has been processed and

metals extracted, leaving a fine-grained sand-sized residue. Although this material has

been processed, many of these tailings still contain some metals (11–13, 73–77). Despite

the importance of microbes for metal mobilization, relatively few studies have been carried

out on these mine wastes to determine what microbial communities are present. From the

limited number of studies addressing this knowledge gap, we know that these sources do

contain robust communities of microorganisms, many of which can catalyze metal

oxidation and reduction (23, 53, 73, 75, 78–87). These communities are highly variable and

demonstrate a heterogeneity that is often overlooked when remediation strategies and

modeling metal removal and cycling dynamics (23, 74), as these communities vary not just

with metal availability, but with depth (83, 86, 88, 89). However, many of these studies have

relied on traditional culture-based microbiological techniques, introducing bias by

selecting for organisms that are easier to grow in lab settings and do not provide an

accurate representation for environmental communities (73, 78, 81, 85). Understanding

microbial community structure and variability across these types of mine waste is crucial

to understanding metal cycling dynamics for these mining-impacted environments and

may play a role in assessing the bioleaching potential of these low-grade non-traditional

metal resources.

In order to evaluate the microbiological communities associated with historic mine

wastes, we sampled waste rock, tailings, and sediment from five inactive mines in south-

central New Mexico: Copper Flat, Center, Carlisle, Alhambra, and Black Hawk. This work

was paired with geochemical studies assessing critical minerals in these wastes (90, 91).

All five sites experienced production at some point in the past, and all contain critical

minerals in their waste (90, 91). The goals of this study were to (i) evaluate the drivers

behind the composition and structure of microbial communities in different waste types;

(ii) evaluate methods for culture-independent analysis of historic mine wastes, including

approaches for obtaining representative microbiological samples of large waste piles; and

(iii) explore microbial resources for future in situ leaching opportunities and potential

bioremediation strategies for these and other historical mining areas.

Geologic context and mine waste characteristics. Samples for this study come from five

inactive mines in three different mining districts: the Carlisle and Center mines in the

Steeple Rock district, the Copper Flat mine in the Hillsboro district, and the Black Hawk

and Alhambra mines in the Black Hawk district (Fig. 1).

The deposits at the Center and Carlisle mines are characterized by a low-sulfidation

volcanic epithermal system with Au-Ag veins (92, 93). Exploration began in 1860, and

production was first reported in 1880 and continued sporadically until 1994. An estimated

$10 million worth of metals was produced including gold (Au), silver (Ag), copper (Cu), lead

(Pb), and zinc (Zn) in its 100 years of activity (92). The workings at the Center mine includes

a central flooded adit, and a mix of low-grade ore material and bulk waste rock (gangue

and low-grade mineralization) that was piled up, flattened, and covered with 15 cm of soil

cover. Recent work by (90) has revealed the presence of critical minerals in the mine waste

at these sites, including arsenic (As), bismuth (Bi), tellurium (Te), fluorite (F), and Zn, all of

which are associated with these types of mineral deposits.

The Copper Flat mine in the Hillsboro District was developed around the

mineralized Copper Flat quartz-monzonite porphyry (CFQM) deposit that was first

identified in 1975 (94, 95). Despite being in production for only 4 months in 1982, the

Copper Flat mine produced 1.2 million short tons of ore containing Cu, Au, and Ag (96).

This site features an open pit lake, a seasonal acidic seep, and extensive bulk waste rock

piles and tailings deposits that contain substantial critical minerals endowments of Bi,

cobalt (Co), gallium (Ga), and Te (90).

The Black Hawk mine is an arsenide 5-element vein mineral system. This deposit

type is associated with critical minerals such as Co, nickel (Ni), As, Bi, tungsten (W), Te,

and Zn. The site experienced sporadic production from 1881-1960, producing Cu, Au, and

Pb (97). The nearby Alhambra mine also experienced production starting in 1881 (98). It is

estimated that the district produced more than $1,000,000 silver from 1883 to 1893 (98).

Recent geochemical work by (91) has revealed the presence of critical minerals in the mine

wastes at both sites including cadmium (Cd), Zn, and Te. The mine waste at both inactive

mines is in the form of bulk waste rock piles mixed with ore material.

RESULTS

Field sampling and observations. We collected samples from all five inactive mine sites: the Carlisle and Center mines in the Steeple Rock District (CY and CN, respectively), the Copper Flat mine in the Hillsboro district (CF), and the Black Hawk and Alhambra mines in the Black Hawk District (BH and AH, respectively) (Fig. 1). The waste types sampled at each location are summarized in Table 1, with additional details in Table S1.

Bulk waste rock was collected from all five sites. In order to take a representative sample of a large waste rock pile, a grid of approximately equal-sized squares was superimposed onto the waste rock pile and the center of each grid square demarcated by flags. Individual samples are collected at each flag (following protocols established by the USGS, outlined in (9). The overburden was removed, and the sample was sieved through a 4mm sieve for subsequent geochemical analyses. To get a representative composite sample for the entirety of the waste rock pile, a small but equal volume of material from each individual sample hole was sieved through a 4mm sieve and combined into one composite sample. For microbiological sampling, six individual samples were chosen, and samples collected from each individual hole. Similarly, a composite sample was collected by combining small amounts of material from each of the selected individual geochemistry holes. Geochemical data from these sites were reported in (91) and (90).

Tailings were sampled at CF and CY. The tailings at CF are horizontally and vertically extensive, and so we employed the same sampling strategy as with the bulk waste rock piles: six individual samples and a composite sample were collected. In addition, tailings were collected above, at, and below a visible oxidation front from one deep sampling pit (Fig. S1). The distance from the top of the pile to the oxidation front was 30 cm, and the distance from the oxidation front to the bottom of the hole where the third sample was collected was 70 cm. The tailings at CY are much less extensive both horizontally and vertically, so only three samples were collected: above, at, and below a visible oxidation interface like at CF (Fig. 1a).

Sediment and biofilms were sampled from flooded adits at CN and CY. At the time

of sampling, the pH of the CN adit was pH 6.5, while the two flooded adits at CY were pH

3.1 and 3.38 (Table S1). Sediments were also sampled from the seasonal acidic seep at CF

where the pH measured 3.55 in 2022 and between 1.54 and 1.95 in 2020.

16S rRNA gene libraries. We prepared 93 successful rRNA gene libraries from 64 of 73

total samples attempted. Libraries ranged from 10,094 to 104,001 reads per sample with

an average library size of 43,269 reads (standard deviation 24,048). In total, 15,648

operating taxonomic units (OTUs) were identified at 97% similarity.

The three types of waste were associated with different microbial communities. The

acidic seep samples from CF and the flooded adits of CN and CY were dominated by

genera commonly found in ARD such as *Acidiphilium, Leptospirillum, Acidibacter,*

Metallibacterium, and *Ferrithrix* (Fig. S2). In contrast, the waste rock samples from all sites

show much more diverse communities compared to the seeps and tailings (Fig. 2), and

include abundant bacteria in the genera *Bradyrhizobium* and *Solirubrobacter*, along with

abundant archaea such as *Candidatus Nitrocosmicus* and *Nitrososphaeraceae* spp. (Fig.

S3). The tailings also contain some of these same genera, as well as *Ramlibacter*,

Paucibacter, *Enhydrobacter*, *Kocuria*, unclassified *Ktedonobacteraceae*, and

Spingomonas. Some of the genera associated with iron and sulfur oxidation that occur in

the seep and adit sediments are present at low levels in other waste types, such as

Acidiphilium (Fig. 3), which also contain some other rare taxa associated with metal and

sulfur oxidation or reduction such as *Sulfuriferula*, *Acidibacter,* and *Sulfurifustis* (Fig. 3).

Overall, the microbial communities from both waste rock and tailings are dominated at all

sites by uncultured and sometimes unclassified microorganisms (Fig. 3, S4).

Non-metric multidimensional scaling (NMDS) ordination shows that rRNA gene

libraries from sites AH, BH, and CN are similar to each other and cluster tightly at one

extreme of the ordination, while libraries from sites CF and CY were much more

heterogeneous (Fig. 4). Sites AH, BH, and CN also had the highest diversity (Fig. 2), perhaps

because these are the three sites that had a soil cover. ANalysis Of SIMilarities (ANOSIM)

analysis showed that libraries from different sites were statistically significantly different (R

= 0.28, p < 0.001), as were libraries from bulk waste rock versus tailings tailings, for all sites

(R = 0.28, p < 0.001) as well as for only libraries from sites CY and CF for which both tailings

and waste rock were sampled (R = 0.40, p < 0.001). These differences were also statistically

significant by PERMANOVA (by site, PERMANOVA F = 6.8, p < 0.001; waste rock versus

tailings for all sites, PERMANOVA F = 6.0, p < 0.001; waste rock versus tailings for site CY

and CF, PERMANOVA F = 3.6, p < 0.001), and all pairwise comparisons of libraries by site

are also all statistically significant (PERMANOVA F = 2.3 - 8.7, p < 0.002).

In order to evaluate potential geochemical drivers of microbial community

composition, we used an environmental overlay to evaluate the relationships among

geochemical variables and rRNA gene and transcript libraries in NMDS ordinations (Fig. 5).

When libraries from all sites are included, %S, %C, SiO_2, and copper content are

statistically significant and oriented with the first ordination axis. This aligns with the

separation between libraries from CN, AH, and BH, which cluster tightly at the extreme of

the first ordination axis, and libraries CY and CF, which are more variable. When only

libraries from only sites AH, BH, and CN were ordinated, libraries cluster by site, with %S,

As, Pb, and U oriented with the first ordination axis, and Cu, Co, Ag, and TREE (total rare

earth element) concentration oriented positively with the first ordination axis and

negatively with the second. When sites were compared based on geochemical variables

alone, sites AH and BH overlap, and separate from CF and CY along the first ordination

axis, indicating distinct geochemical characteristics that could be driving differences in

microbial communities among the sites (Fig. S5). In order to further explore the relationship

between specific taxa and geochemical variables, we correlated the most abundant OTUs

with individual geochemical variables in order to determine if specific populations might be

responding to specific geochemical drivers. Select correlations are shown in Fig. S6.

Evaluation of sampling methods for historic waste rock and tailings piles. To attempt to

get a representative sample of large waste rock and tailings piles, six samples were taken

from separate random locations around the perimeter and surface of the pile. A small

volume of material from each of these individual samples were combined into a composite

sample and homogenized, and libraries prepared from both the individual and composite

samples. Non-metric multidimensional scaling (NMDS) ordination shows that the

composite samples plot with the individual samples from the same waste pile for all five

sites (Fig. 6). In addition, because waste rock is heterogeneous, we extracted each

samples twice and, in some cases, sequenced individual sample DNA extracts along with

the combined DNA extracts. Overall, 16S rRNA gene libraries from the individual extracts

were similar to those from the combined extract (Fig. S7). We also found that the Qiagen

PowerSoil Pro kit resulted in higher DNA yields than other kits tested, so all DNA library

analyses reported in this study were extracted using this kit using a bead-beating protocol

described in (99).

16S rRNA transcript libraries. In order to assess active microbial populations in the mine

waste, rRNA transcript libraries were prepared from composite samples of all waste rock

piles and the CF tailings. rRNA transcript libraries from all except for one sample (CY23-13)

were successful, and successful libraries contained between 34,975 and 57,123 with an

average library size of 38,759 reads (standard deviation 8,140).

NMDS ordinations show that 16S rRNA transcript libraries from the composite

waste rock, as well as the tailings composite from CF, are overall similar to 16S rRNA gene

libraries from those locations (Fig. 4, 7, 8); NMDS analyses of rRNA gene libraries alone

have similar structure to those with rRNA transcripts (Fig. S7). Of the seven composite

samples with successful 16S rRNA transcript libraries, six had successful 16S rRNA gene

libraries and could be compared directly. For the other composite tailings sample, rRNA

transcript libraries were compared to individual samples from the same tailings pile. These

comparisons show that, overall, most taxa in the rRNA gene libraries are present in the

rRNA transcript libraries but there are individual OTUs and phyla that be either over- or

under-represented (Fig. 7, 8). In general, archaea such as *Candidatus* Nitrososphaera and

Candidatus Nitrocosmicus are represented at much higher abundances in the rRNA gene

libraries compared to the rRNA transcript libraries. Other genera, such as *Bradyrhizobium*

and *Streptomyces*, appear in higher abundance in the rRNA transcript libraries compared

to the rRNA gene libraries. The majority of organisms appear similarly abundant in rRNA

transcripts as in the gene libraries, such as *Solirubrobacter* spp., *Sulfurifustis* spp.,

Nitrospira spp., and *Kocuria* spp.

Cell counts. We performed cell counts on composite waste rock samples from sites AH,

BH, CN, and CY, and for site CF, we counted four individual samples. Bulk waste rock had

between 6.1×10^7 and 1.1×10^{10} cells g^{-1}, with the lowest cell abundance at site CF and the

highest at site CN (Table 2). Overall, tailings had lower cell abundance, between 2.0×10^6

and 6.9×10^8 cells g^{-1} (Table 2). At sites CF and CN, cell counts were lowest in tailings

sampled below the visible oxidation fronts, but were at a similar order of magnitude to

samples from at and above the oxidation front (Table 2).

DISCUSSION

Different waste types have distinct microbial communities. Overall, microbial

communities varied strongly among the different waste types (seeps and adits, bulk waste

rock, and tailings). The acidic seeps and adits had the least diverse communities, with

abundant taxa of known acidophilic iron- and sulfur-oxidizing microorganisms that are

common in acid rock drainage, such as *Leptospirillum* spp., *Sulfuritalea* spp., and

Acidiphilium spp., *Cuniculiplasma* spp., *Acidibacter* spp., and *Ferrithrix* spp. (24, 26, 27,

31, 100). There are also photosynthetic primary producers present, based on chloroplasts

from the unicellular algal taxa *Chlamydomonas* and *Euglena*. Overall, the most abundant

organisms in the acidic seeps and adits only occur at low levels in the waste rock and

tailings (Fig. S2).

In contrast, microbial communities in the bulk waste rock and tailings were more

diverse, and were dominated by different microorganisms, including abundant taxa from

genera and families without cultured representatives (Fig. 3, S3). Many of the OTUs

identified in the bulk waste and tailings that were classified to the genus level are more commonly associated with soil or other non-mining environments. For example, based on rRNA gene and transcript libraries, *Solirubrobacter* spp. and *Kocuria* spp. are abundant and active in the bulk waste piles sampled here, and are commonly found in desert soils that have been found to be highly UV-C resistant and dessication resistant (101–103). Bulk waste also contains unclassified members of the *Ktedonobacteraceae* (Fig. S3), which has been found to have a degree of heavy metal resistance and are often associated with metal-impacted soils (104, 105). Waste rock communities also include members of the Crenarchaeota and Thaumarchaeota such as *Candidatus* Nitrososphaera, *Candidatus* Nitrocosmicus, and *Candidatus* Nitrosotalea in rRNA gene and transcript libraries. These are three groups of ammonia-oxidizing archaea (AOA) that have been found in a variety of environments including ARD (106), wastewater treatment plants (107), and, more commonly, soils (108–113). There are other genera of common soil nitrogen cyclers present in our bulk waste sample microbial communities including *Noviherbaspirillum* (114, 115), *Bradyrhizobium* (116, 117), and *Nitrospira* spp. (111, 118). Some of the taxa that are abundant in tailings, but not seeps or the bulk waste, also include organisms common to soils. For example, *Ramlibacter* spp. have been observed in soils and associated with plant rhizomes (119–122). *Sphingomonas* spp. have been found in a wide variety of environments including freshwater (123) and soils (124, 125), including reclaimed amended tailings piles (87), and *Enhydrobacter* sp. has been identified in metal-impacted soils specifically soils containing high concentrations of cadmium (Cd) (126, 127).

The waste rock communities at sites AH, BH, and CN had both the highest diversity and species richness compared to CF and CY (Fig. 2). This is likely because the waste rock at AH, BH, and CN had 10-15 cm of soil cover over the top of the waste rock piles, while waste rock and tailings piles at CF and CY had no such cover. A cover provides additional opportunities for microbial colonization from the overlying soil, and increases nutrient availability and substrate attachment opportunities. There has been limited work assessing the microbial communities of bulk waste rock, both with and without a soil cover. Our results are consistent with work by (128) which found that bulk waste rock that

was covered with a soil overburden had significant microbial contributions from the soil

cover, and that samples close to the soil-waste interface were more diverse than samples

from deeper in the waste rock pile. In our samples, the soil cover appears to have created a

more homogenous microbial environment, as libraries from individual CF and CY samples

are much more variable compared to individual samples from AH, BH, and CN (Fig. 2, 4).

In this study, we focused on the shallow surface of the waste rock and tailings.

Chen et al. (83) found that *Proteobacteria* were primarily dominant in the upper layers of

the tailings pile, and that the community shifted at depth to contain more *Firmicutes* and

Desulfobacterota. This community change was attributed to geochemical variables such

as pore-water pH increases with depth and dissolved metal ion concentrations, most

notably iron. Other studies have suggested that community changes with depth also hinge

on the disappearance of oxides and oxygen, leading to an increase in sulfate- and iron-

reducing microorganisms (53, 88, 129). The only areas where we sampled with depth were

in tailings piles at sites CF and CN, where we were able to sample above, below, and at a

visible redox interface (Fig. 1a, S1). *Acidibacter* spp., which is a genus of iron-reducing

bacteria (130), are present at and below the redox interface at CY, but otherwise we don't

see any clear trends in the specific phyla identified in (83) or any abundant known sulfate

or metal reducers (Fig. S8). However, many libraries contain OTUs from genera, families,

and higher taxonomic groupings, so it is possible that these samples contain unrecognized

sulfate or iron reducers.

Overall, ordination analyses showed that differences among microbial communities

correspond to elemental content and other geochemical variables among the sites (Fig. 5).

In addition, some OTUs are statistically significantly correlated with geochemical variables

such as percent sulfur, iron, and other metals that could indicate a response to rock-

associated chemical energy resources or toxic compounds (Fig. S9). For example, OTUs

corresponding to "*Candidatus* Nitrososphaera" and the group KD4-96 in the *Chloroflexi*ota

are abundant in samples with high iron and low sulfur concentrations that have low copper

and lead (Fig. S9), and uncultured members of order Vicinamibacterales (phylum

Acidobacteriota) and *Gaiellales* (*Actinobacteriota*) are similarly negatively correlated with

sulfur, copper, and lead content (Fig. S9), seemingly independent of site. Perhaps these

taxa are sensitive to metals. In contrast, there are not any abundant OTUs that show clear

correlations with increasing sulfur content. Some taxa from samples with the highest sulfur

include *Dechloromonas*, *Rothia*, *Corynebacterium*, *Enhydrobacter*, *Sphingomonas*, which

are not genera typically associated with inorganic sulfur compound oxidation, but these

taxa are only abundant in a few high sulfur samples. There are some OTUs that are

positively correlated with rare earth element content, such as OTUs classified Archangium

(*Myxococcota*) and TRA3−20 in the *Gammaproteobacteria* (Fig. S9), although these taxa

could instead be responding to other factors, such as high concentrations of other

elements like As or Co that are also abundant in these samples (Table S1). OTUs classified

as IMCC26256 (*Actinobacteriota*) and B12-WMSP1 (*Ktedonobacteria*, *Chloroflexi*ota) are

abundant in rRNA gene and transcript libraries from samples with high Cu and low pH from

uncovered waste rock at CY and CF (Fig. S9), which could indicate an association with

copper sulfide minerals at these sites. However, future meta-omic or culture-based

analyses will be necessary to evaluate the genomic and metal-cycling capabilities of these

and other uncultured and unclassified organisms associated with these historic wastes.

Implications for microbiological characterization of historic waste rock and tailings.

Waste rock and tailings can be challenging environments for microbiological analysis due

to their heterogeneity, metal content, and significant spatial extent on the order of 10s of

meters up to kilometers. To attempt to capture this heterogeneity, we compared individual

versus composite samples, and showed that the composite samples were effective at

representing the heterogeneity of the waste pile microbial communities for the waste piles

sampled here (Fig. 6). On one hand, collecting and analyzing multiple samples from each

waste rock pile provided an opportunity to draw more careful correlations between

geochemical variables and help parse out site specific effects (Fig. 5, S9). However, on the

other hand, a large number of samples for microbiological analysis can be cost prohibitive,

so collecting a composite sample is a reasonable approach for characterizing the waste-

associated microbial communities from large waste rock and tailings piles, at least for

piles that are on the order of several 10s of meters like those sampled here.

Implications for mine waste biogeochemistry and metal mobilization. Microbial

communities associated with acid mine drainage and acidic, sulfide mineral-rich ores

often have abundant iron- and sulfur-oxidizing populations. These organisms are important

for metal release in biomining operations. Iron-oxidizers serve as "oxidant generators" that

produce Fe(III), which is the primary oxidant for metal sulfide minerals; sulfur oxidizers are

"acid generators" that produce sulfuric acid and contribute to the breakdown of some

sulfide minerals; and heterotrophs act as "janitors" that remove low molecular weight

organic carbon compounds that are often toxic to the autotrophic Fe and S oxidizing

populations (131, 132). While we found this type of community in the acidic seeps and

adits, the waste rock and tailings had a very different structure that was dominated by taxa

more common to soil and that are associated with the oxidation of organic carbon and

inorganic nitrogen compounds. Some taxa in the waste rock and tailings are known to

oxidize or reduce iron and sulfur compounds, such as *Sulfurifustis* (133) and *Acidibacter*

(130) (Fig. 3), but these were not very abundant. This differs from previous work on

circumneutral mine waste, especially work utilizing culture-dependent methods, where a

higher number of acidophilic and neutrophilic iron- and sulfur-oxidizing microorganisms

have been identified (73, 78, 80, 85, 86, 88, 89, 134). It is likely that the communities within

the waste rock piles sample here are not mobilizing metals to the same extent we would

see in ARD and bioleaching heaps.

The presence of a soil-like microbial community may indicate successful

reclamation. The revegetation of tailings piles has been shown to change microbial

community assemblage in favor of nitrogen-cycling and rhizome-associated

microorganisms (81), thereby halting metal mobilization and preventing the formation of

ARD (41). For example, prior studies have found that reclaimed vegetated tailings piles

contain high abundances of *Nitrospira* spp. and *Firmicutes*. (79, 81, 84). In semi-arid areas

in particular, microorganisms common to soil microbiomes were found in high abundance

on abandoned tailings piles, and 47% of the taxa present in these communities were

associated with drought and UV stress conditions (87), like the *Solirubrobacter* spp. and

Kocuria spp. observed here.

With critical minerals in high demand and the push to identify and quantify non-

traditional metal resources, waste rock and tailings represent a substantial resource that

could theoretically be accessed using in situ leaching or other biomining techniques.

However, we don't know if the more abundant sulfur- and metal-cycling organisms such as

Sulfurifustis and *Acidibacter*, or some of the rare populations like *Acidiphilium*, could be

stimulated to mobilize metals and critical minerals in these deposits. Changing

geochemical conditions to favor these organisms, such as by increasing the temperature

or moisture conditions or adding elemental sulfur or ferrous iron to favor

chemolithoautotrophic microbial growth, could stimulate leaching (55, 65, 134–136). In

addition, like other historic mine waste resources (83), these deposits have many unknown

populations that may be mobilizing or immobilizing metals via metabolic pathways that are

not yet well-understood or recognized. A more complete understanding of the microbial

assemblages that develop *in situ* may provide new insight into the most appropriate

organisms and processes for metal recovery and recycling from these and other historic

mine waste deposits.

MATERIALS AND METHODS

Field sites and sampling procedures. Samples were collected from five inactive mine

sites: Copper Flat mine (Sierra County), Carlisle mine, Center mine, Alhambra mine, and

Black Hawk mine (Grant County), abbreviated CF, CY, CN, AH, and BH, respectively.

Samples of all available waste types were collected at each site, and included acidic

seeps, flooded adits, bulk waste rock, and tailings (Table 1).

For waste rock sampling, material for geochemical analyses was sampled from the

top 15 cm as described in (90), (91), and (9). Samples for microbiological analysis were

collected by using sterile implements to expose a fresh surface and scrape approximately

5 mL of material from these same holes into a sterile 15 or 50 mL tube. Samples were

mixed by shaking, and approximately 500µL was removed into a 2 mL microcentrifuge tube

for cell counting. The larger sample was then immediately frozen on dry ice for nucleic acid

extraction and stored at -80°C upon return to the NMT Geobiology Lab. The smaller subset

was fixed with paraformaldehyde (PFA) as in (137) and stored at 4°C. For large waste rock

piles, samples were collected from six surface sites selected randomly around the

perimeter in an attempt to evaluate and capture the heterogeneity of the large pile.

Samples were preserved individually, along with a composite sample that was prepared by

combining approximately 5 mL from each of the six locations and preserving it as for the

individual samples. Four sampling locations had a soil cover: the tailings at CF, and bulk

waste rock at CN, BH, and AH. Where a soil cover was present, the hole was dug so that

waste rock material was sampled and not the overlying soil cover.

Because of the larger size of the tailings pile at CF, individual and composite

samples were collected as for bulk rock above. At sites CF and CN, tailings displayed

oxidation fronts that were clearly visible by color changes, and were sampled above, at,

and below these fronts. Sediments of seep and adit sediments were collected from the top

0.5 cm and preserved as described above.

Nucleic acid extraction and rRNA gene and transcript library preparation. DNA was

extracted using a DNeasy PowerSoil Pro DNA isolation kit (Qiagen, Hilden, Germany)

following the standard kit protocol with one modification: samples were subject to bead

beating for 20, 40, and 60 seconds at 2500 RPM using a Mini-Beadbeater (Biospec

Products, Bartlesville, OK, USA), and a 250 µL aliquot was removed at each time point as in

(99). To avoid the "nugget effect", each sample was extracted at least twice. The V4 region

of the 16S rRNA gene (DNA) was amplified using 10mM V4 Nextera primers (Integrated

DNA Technologies, Coralville, IA, USA; forward primer tail

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG; reverse primer tail

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG). PCR was performed with the AllTaq

PCR Core Kit (Qiagen) with initial denaturation for 5 minutes at 94ºC followed by 30 or 35

cycles of 94ºC for 30s, 52ºC for 45s, and 72ºC for 30s, with a final elongation at 72ºC for 10

minutes. Samples were then held within the thermocycler at 4ºC until they were moved to

a -20ºC freezer. Samples were run for 30 cycles initially, and if no amplification was

observed, the sample was re-amplified for 35 cycles. A DNA extraction blank was also

subjected to PCR to check for contamination during the DNA extraction process, and a

PCR reagent blank was amplified to check for PCR reagent contamination. Extractions with

visible PCR produce in blanks were not processed further. We also tested DNeasy

PowerBiofilm Kit (Qiagen) and DNeasy PowerSoil Kit (Qiagen), but had overall better

performance with the DNeasy PowerSoil Pro kit (Qiagen).

RNA extraction was performed using the RNeasy PowerBiofilm RNA isolation kit

(Qiagen), using the same bead-beating protocol as for DNA extractions, above, with a 150

µL aliquot removed after 20, 40, and 60 s of bead beating. DNA was removed using two

DNase treatments, an on-column DNase treatment using DNase I (Qiagen) that was

provided in the RNeasy PowerBiofilm RNA isolation kit, and any remaining DNA in the

sample was digested using the Invitrogen TURBO DNA-free kit kit (Thermo Fisher Scientific,

Waltham, MA, USA) and subsequently cleaned with the RNA Clean and Concentrator-5 kit

(Zymo Research, Irvine, CA, USA). Reverse-transcription PCR was conducted using a

OneStep *Ahead* RT-PCR Kit (Qiagen) with the same V4 primers as described above under

the following program: 50ºC for 10 minutes, followed by an initial denaturation for 5

minutes at 94ºC, and then 30 or 35 cycles of 94ºC for 10s, 50ºC for 10 s, and 72ºC for 15 s

before finishing with 5 minutes at 72ºC and holding at 12ºC until the PCR products were

moved to the -20ºC freezer for storage. Samples were run for 30 PCR cycles initially, and

were re-run at 35 cycles if no RNA amplification was observed. To ensure that samples did

not contain residual DNA, we amplified the RNA extract according to the DNA PCR

specifications described above. If DNA amplification was observed after 35 cycles, the

Invitrogen TURBO DNA-free (Qiagen) treatment and RNA Clean and Concentrator (Zymo)

process was repeated until no amplification was observed (usually no more than one

additional treatment was necessary). A PCR reagent blank was also included in each PCR

run to check for PCR reagent contamination.

PCR products were barcoded and sequenced at the University of Minnesota Core

Genomics Center (UMGC) on an Illumina MiSeq with version 3 chemistry and 300 paired

end cycles. DNA extraction blanks and PCR reagent blanks were sequenced alongside

samples, even though no PCR products were visible.

Cell counting. In order to detach cells from mineral and soil particle surfaces, a small

amount of the PFA-fixed sample and sonicated using a FisherBrand Model CL-18

dismembrator (Fisher Scientific, Hampton, NH, USA) for 45 seconds on 100% amplitude.

Cells were then stained with DAPI (4',6-diamidino-2-phenylindole) (Biotium, Fremont, CA,

USA) and filtered onto a 0.2 µM membrane filter (Cytiva Whatman, Marlborough, MA, USA).

The filter was mounted on a glass slide, covered using Vectashield antifade mounting

medium (Vector Laboratories, Newark, CA, USA), and observed under an Olympus BX63F

compound epifluorescence microscope (Olympus Life Science, Waltham, MA, USA). Cells

were counted from 10 randomly-selected grid sections on the slide, averaged, and

converted to cells g^{-1} sediment based on the mass of sediment sonicated and accounting

for the dilution at each step.

Bioinformatic and statistical analyses. Quality filtering, trimming, and OTU calling

followed the same procedures as in (137) and described below. Raw sequences were

trimmed and filtered using Sickle (https://github.com/najoshi/sickle) to an average quality

score of >28 (5' trimming only) and a minimum length of 100 bp. Cutadapt v. 4.2 (138) was

used to trim residual adaptors, and PEAR (138) was used to assemble the forward (R1) and

reverse (R2) reads. Primer sequences were removed with prinseq v.0.20.4 (140). OTU

calling (97% similarity) and chimera removal also followed procedures in (137) using

USEARCH v. 11 (141) and VSEARCH v. 2.21 (142). Representative OTUs were determined

with mothur v. 1.47 (143) and SILVA v. 138.1 (144) with a confidence cut-off score of 50.

Following quality filtering and trimming, libraries with <10,000 reads were excluded from

subsequent community analyses.

R Statistical Software RStudio v. 4.2.1 (145) was used to conduct statistical

analyses. Additional R packages used included vegan (v. 2.6-4) (145), cluster (2.1.4) (147),

ggplot2 (v. 3.5.1) (148). Rare OTUs (<0.01%) were excluded, and the remaining data was

normalized using the angular transformation to deemphasize abundant OTUs. Non-metric

multidimensional scaling (NMDS) ordinations were calculated using the *metaMDS()*

function with four dimensions (k=4). ANOSIM and PERMANOVA analyses were calculated

with the vegan package using default parameters (Bray-Curtis distance, 999 permutations,

with *anosim()* and *adonis2()* functions). The statistical significance of comparisons

between 3 or more groups was assessed with a pairwise analysis using the

pairwise.adonis() function (149). Diversity was calculated using richness and the Shannon-

Wiener Index, which takes into account the number of OTUs and their evenness, using

functions *rarefy()* and *diversity()* after subsampling the OTU matrix to 10,000 reads.

Stacked bar charts of OTU relative abundance were created using code from

https://jkzorz.github.io/2019/06/05/stacked-bar-plots.html and color palettes for the

stacked bar charts from Paletteer (https://github.com/EmilHvitfeldt/paletteer).

Data availability. Raw sequences are available on the NCBI SRA under BioProject

PRJNA1226971.

ACKNOWLEDGMENTS

We would like to thank researchers and students from the economic geology research

group at the New Mexico Bureau of Geology and Mineral Resources for their help with field

work and site access including Bonnie Frey, Evan Owen, Mark Leo-Russell, Stellah

Cherotich, Abena Acheampong-Mensah, Kyle Stafford, and Bob and Jakob Newcomer. We

also thank Robert Seal and Kate Campbell from the USGS for assistance with field work

and insightful discussion. We thank Jimmy Swift for his help with cell counts and

microscopy. Site access was provided by Themac Resources and Santa Fe Gold Corp.

This work was supported by a USGS Earth MRI grant award #G22AC00510, and a

Grant-in-Aid from the New Mexico Geological Society.

REFERENCES

1. IEA. 2023. World Energy Outlook 2023. Paris.

2. White House. 2022. Securing a made in America supply chain of critical minerals. Washington D.C.

3. Burton J. 2022. U.S. Geological Survey releases 2022 list of critical minerals. USGS National News Releases. https://www.usgs.gov/news/national-news-release/us-geological-survey-releases-2022-list-critical-minerals.

4. McLemore VT, Gysi A. 2023. Critical minerals in New Mexico. *New Mexico Earth Matters*, v. Winter 2023. New Mexico Bureau of Geology and Mineral Resources. Available at https://geoinfo.nmt.edu/publications/periodicals/earthmatters/23/n1/em_v23_n1.pdf

5. Watling H. 2015. Review of biohydrometallurgical metals extraction from polymetallic mineral resources. Minerals 5:1–60. https://doi.org/10.3390/min5010001.

6. Calvo G, Mudd G, Valero A, Valero A. 2016. Decreasing ore grades in global metallic mining: A theoretical issue or a global reality? Resources 5:1–14. https://doi.org/10.3390/resources5040036.

7. Tabelin CB, Park I, Phengsaart T, Jeon S, Villacorte-Tabelin M, Alonzo D, Yoo K, Ito M, Hiroyoshi N. 2021. Copper and critical metals production from porphyry ores and E-wastes: A review of resource availability, processing/recycling challenges, socio-environmental aspects, and sustainability issues. Resour Conserv Recycl 170:105610. https://doi.org/10.1016/j.resconrec.2021.105610.

8. Srichandan H, Mohapatra RK, Parhi PK, Mishra S. 2019. Bioleaching approach for extraction of metal values from secondary solid wastes: A critical review. Hydrometallurgy 189:105122. https://doi.org/10.1016/j.hydromet.2019.105122.

9. McLemore VT, Owen EJ. 2024. Geochemistry of critical minerals in mine wastes in New Mexico. SME Annual Conference Preprint 24-043:347-355. Available at https://store.smenet.org/20slp36/347

10. Hobart KK, Piatak NM, Seal RR. 2023. Evaluating sulfide minerals in mine tailings from volcanogenic massive sulfide and sedimentary exhalative zinc-lead deposits as a potential source of critical mineral resources. Goldschmidt Annual Conference Abstracts, Lyon, France, July 9-14, 2023. https://doi.org/10.7185/gold2023.17061

628   11.   Ceniceros-Gómez AE, Macías-Macías KY, de la Cruz-Moreno JE, Gutiérrez-Ruiz ME,
629   Martínez-Jardines LG. 2018. Characterization of mining tailings in México for the
630   possible recovery of strategic elements. J South Am Earth Sci 88:72–79.
631   https://doi.org/10.1016/j.jsames.2018.08.013.
632

633   12.   Sarker SK, Haque N, Bhuiyan M, Bruckard W, Pramanik BK. 2022. Recovery of
634   strategically important critical minerals from mine tailings. J Environ Chem Eng 10:
635   107622. https://doi.org/10.1016/j.jece.2022.107622.
636

637   13.   Vitti C, Arnold BJ. 2022. The Reprocessing and revalorization of critical minerals in
638   mine tailings. Min Metall Explor 39:49-54. https://doi.org/10.1007/s42461-021-
639   00524-6.
640

641   14.   USGS Communications and Publishing. 2021. US Mines produced an estimated
642           $82.3 billion in minerals during 2020. USGS National News Releases.
643   https://www.usgs.gov/news/national-news-release/us-mines-produced-estimated-
644   823-billion-minerals-during-2020.
645

646   15.   Ladislaw S, Carey L, Bright H. 2019. Critical minerals and the role of U.S. mining in a
647   low-carbon future. *Center for Strategic & International Studies*. Available at
648   https://csis-website-prod.s3.amazonaws.com/s3fs-
649   public/publication/191218_CriticalMinerals_layout_v3.pdf
650

651   16.   Dudley B. 2019. BP Statistical Review of World Energy 2019. Pureprint Group Limited,
652   UK.
653

654   17.   Whitworth AJ, Forbes E, Verster I, Jokovic V, Awatey B, Parbhakar-Fox A. 2022. Review
655   on advances in mineral processing technologies suitable for critical metal recovery
656   from mining and processing wastes. Clean Eng Technol 7: 100451.
657   https://doi.org/10.1016/j.clet.2022.100451.
658

659   18.   Boulet MP, Larocque ACL. 1998. A comparative mineralogical and geochemical study
660   of sulfide mine tailings at two sites in New Mexico, USA. Environmental Geology
661   33:130–142.
662

663   19.   Hobart KK, Greensky ZZ, Hernandez K, Feinberg JM, Bailey J V., Jones DS. 2023.
664   Microbial communities from weathered outcrops of a sulfide-rich ultramafic
665   intrusion, and implications for mine waste management. Environ Microbiol 25:3512–
666   3526. https://doi.org/10.1111/1462-2920.16489.
667

668   20.   Jones DS, Walker GM, Johnson NW, Mitchell CPJ, Coleman Wasik JK, Bailey J V. 2019.
669   Molecular evidence for novel mercury methylating microorganisms in sulfate-
670   impacted lakes. ISME J 13:1659–1675. https://doi.org/10.1038/s41396-019-0376-1.
671

672   21.   Oremland RS, Kulp TR, Blum JS, Hoeft SE, Baesman S, Miller LG, Stolz JF. 2005.
673   Microbiology: A microbial arsenic cycle in a salt-saturated, extreme environment.
674   Science 308:1305–1308. https://doi.org/10.1126/science.1110832.
675

676   22.   Turpeinen R, Kairesalo T, Häggblom MM. 2004. Microbial community structure and
677   activity in arsenic-, chromium- and copper-contaminated soils. FEMS Microbiol Ecol
678   47:39–50. https://doi.org/10.1016/S0168-6496(03)00232-0.
679

680   23.   Frascoli F, Hudson-Edwards KA. 2018. Geochemistry, mineralogy and microbiology
681   of molybdenum in mining-affected environments. Minerals 8:42.
682   https://doi.org/10.3390/min8020042.
683

684   24.   Grettenberger CL, Hamilton TL. 2021. Metagenome-assembled genomes of novel
685   taxa from an acid mine drainage environment. Appl Environ Microbiol 87:1–17.
686   https://doi.org/10.1128/AEM.00772-21.
687

688   25.   Grettenberger CL, Havig JR, Hamilton TL. 2020. Metabolic diversity and co-
689   occurence of multiple *Ferrovum* species at an acid mine drainage site. BMC
690   Microbiol 20:119. https://doi.org/10.1186/s12866-020-01768-w.
691

692   26.   Wakao N, Nagasawa N, Matsuura T, Matsukura H, Matsumoto T, Hiraishi A, Sakurai Y,
693   Shiota H. 1994. *Acidiphilum multivorum* sp. nov., an acidophilic
694   chemoorganotrophic bacterium from pyritic acid mine drainage. J Gen Appl
695   Microbiol 40:143–159. https://doi.org/10.2323/jgam.40.143.
696

697   27.   Dopson M, Baker-Austin C, Hind A, Bowman JP, Bond PL. 2004. Characterization of
698   *Ferroplasma* isolates and *Ferroplasma acidarmanus* sp. nov., extreme acidophiles
699   from acid mine drainage and industrial bioleaching environments. Appl Environ
700   Microbiol 70:2079–2088. https://doi.org/10.1128/AEM.70.4.2079-2088.2004.
701

702   28.   Watkin ELJ, Keeling SE, Perrot FA, Shiers DW, Palmer M-L, Watling HR. 2009. Metals
703   tolerance in moderately thermophilic isolates from a spent copper sulfide heap ,
704   closely related to *Acidithiobacillus caldus*, *Acidimicrobium ferrooxidans* and
705   *Sulfobacillus thermosulfidooxidans*. J Microbiol Biotechnol 36:461–465.
706   https://doi.org/10.1007/s10295-008-0508-5.
707

708   29.   Bevilaqua D, Leite ALLC, Garcia O, Tuovinen OH. 2002. Oxidation of chalcopyrite by
709   *Acidithiobacillus ferrooxidans* and *Acidithiobacillus thiooxidans* in shake flasks.
710   Process Biochem 38:587–592. https://doi.org/10.1016/S0032-9592(02)00169-3.
711

712   30.   Galleguillos Perez PA. 2011. Biodiversity and stress response of extremophilic
713   prokaryotes isolated from the Escondida copper mine, Chile. Doctoral dissertation,
714   Bangor University (United Kingdom).
715

716 31. Mesa V, Gallego JLR, González-Gil R, Lauga B, Sánchez J, Méndez-García C, Peláez
717   AI. 2017. Bacterial, archaeal, and eukaryotic diversity across distinct microhabitats
718   in an acid mine drainage. Front Microbiol 8:1756.
719   https://doi.org/10.3389/fmicb.2017.01756.
720

721 32. Johnson DB. 2003. Chemical and microbiological characteristics of mineral spoils
722   and drainage waters as abandoned coal and metal mines. Water Air Soil Pollut 3:47–
723   66. https://doi.org/10.1023/A:1022107520836.
724

725 33. Nordstrom DK, Blowes DW, Ptacek CJ. 2015. Hydrogeochemistry and microbiology
726   of mine drainage: An update. Applied Geochemistry 57:3–16.
727   https://doi.org/10.1016/j.apgeochem.2015.02.008.
728

729 34. Tyson GW, Chapman J, Hugenholtz P, Allen EE, Ram RJ, Richardson PM, Solovyev V
730   V., Rubin EM, Rokhsar DS, Banfield JF. 2004. Community structure and metabolism
731   through reconstruction of microbial genomes from the environment. Nature 428:37–
732   43. https://doi.org/10.1038/nature02340.
733

734 35. Abramov SM, Straub D, Tejada J, Grimm L, Schädler F, Bulaev A, Thorwarth H, Amils
735   R, Kappler A, Kleindienst S. 2022. Biogeochemical niches of Fe-cycling communities
736   influencing heavy metal transport along the Rio Tinto, Spain. Appl Environ Microbiol
737   88:e02290-21. https://doi.org/10.1128/aem.02290-21.
738

739 36. Mahabub MS, Alahi F, Al Imran M. 2023. Unlocking the potential of microbes:
740   biocementation technology for mine tailings restoration—a comprehensive review.
741   Environ Sci Pollut Res 30:91676–91709.
742

743 37. Johnson DB, Yajie L, Okibe N. 2008. "Bioshrouding"—A novel approach for securing
744   reactive mineral tailings. Biotechnol Lett 30:445–449.
745   https://doi.org/10.1007/s10529-007-9574-4.
746

747 38. Larson LN, Fitzgerald M, Singha K, Gooseff MN, Macalady JL, Burgos W. 2013.
748   Hydrogeochemical niches associated with hyporheic exchange beneath an acid
749   mine drainage-contaminated stream. J Hydrol 501:163–174.
750   https://doi.org/10.1016/j.jhydrol.2013.08.007.
751

752 39. Jones DS, Kohl C, Grettenberger C, Larson LN, Burgos WD, Macalady JL. 2015.
753   Geochemical niches of Iron-oxidizing acidophiles in acidic coal mine drainage. Appl
754   Environ Microbiol 81:1242–1250. https://doi.org/10.1128/AEM.02919-14.
755

756 40. Johnson DB, Hallberg KB. 2005. Acid mine drainage remediation options: A review.
757   Sci Tot Environ 338:3–14. https://doi.org/10.1016/j.scitotenv.2004.09.002.
758

759 41. Akcil A, Koldas S. 2006. Acid Mine Drainage (AMD): causes, treatment and case
760 studies. J Clean Prod 14:1139–1145. https://doi.org/10.1016/j.jclepro.2004.09.006.
761

762 42. Hedrich S, Schippers A. 2020. Distribution of acidophilic microorganisms in natural
763 and man-made acidic environments. Curr Issues Mol Biol 40:25–48.
764 https://doi.org/10.21775/CIMB.040.025.
765

766 43. Johnson DB, Kanao T, Hedrich S. 2012. Redox transformations of iron at extremely
767 low pH: Fundamental and applied aspects. Front Microbiol 3:96.
768 https://doi.org/10.3389/fmicb.2012.00096.
769

770 44. Bosecker K. 1997. Bioleaching: Metal solubilization by microorganisms. FEMS
771 Microbiol Rev 20:591–604. https://doi.org/10.1016/S0168-6445(97)00036-3.
772

773 45. Schippers A, Breuker A, Blazejak A, Bosecker K, Kock D, Wright TL. 2010. The
774 biogeochemistry and microbiology of sulfidic mine waste and bioleaching dumps
775 and heaps, and novel Fe(II)-oxidizing bacteria, p. 342–350. *In* Hydrometallurgy.
776 https://doi.org/10.1016/j.hydromet.2010.01.012.
777

778 46. Olson GJ, Brierley JA, Brierley CL. 2003. Bioleaching review part B: Progress in
779 bioleaching: Applications of microbial processes by the minerals industries. Appl
780 Microbiol Biotechnol 63:249–257. https://doi.org/10.1007/s00253-003-1404-6.
781

782 47. Cárdenas JP, Quatrini R, Holmes DS. 2016. Genomic and metagenomic challenges
783 and opportunities for bioleaching: a mini-review. Res Microbiol 167:529–538.
784 https://doi.org/10.1016/j.resmic.2016.06.007.
785

786 48. Giebner F, Kaden L, Wiche O, Tischler J, Schopf S, Schlömann M. 2019. Bioleaching
787 of cobalt from an arsenidic ore. Miner Eng 131:73–78.
788 https://doi.org/10.1016/j.mineng.2018.10.020.
789

790 49. Panda S, Akcil A, Pradhan N, Deveci H. 2015. Current scenario of chalcopyrite
791 bioleaching: A review on the recent advances to its heap-leach technology.
792 Bioresour Technol 196:694–706. https://doi.org/10.1016/j.biortech.2015.08.064.
793

794 50. Sand W, Schippers A, Hedrich S, Vera M. 2022. Progress in bioleaching:
795 fundamentals and mechanisms of microbial metal sulfide oxidation – part A. Appl
796 Microbiol Biotechnol 106: 6933–6952. https://doi.org/10.1007/s00253-022-12168-7.
797

798 51. Zhu J, Li Q, Jiao W, Jiang H, Sand W, Xia J, Liu X, Qin W, Qiu G, Hu Y, Chai L. 2012.
799 Adhesion forces between cells of *Acidithiobacillus ferrooxidans, Acidithiobacillus*
800 *thiooxidans* or *Leptospirillum ferrooxidans* and chalcopyrite. Colloids Surf B
801 Biointerfaces 94:95–100. https://doi.org/10.1016/j.colsurfb.2012.01.022.
802

803 52. Jia Y, Tan Q, Sun H, Zhang Y, Gao H, Ruan R. 2019. Sulfide mineral dissolution
804   microbes: Community structure and function in industrial bioleaching heaps. Green
805   Energy Environ 4:29–37. https://doi.org/10.1016/j.gee.2018.04.001.
806

807 53. Lu X, Wang H. 2012. Microbial oxidation of sulfide tailings and the environmental
808   consequences. Elements 8:119–124. https://doi.org/10.2113/gselements.8.2.119.
809

810 54. Jorjani E, Askari Sabzkoohi H. 2022. Gold leaching from ores using biogenic lixiviants
811   – A review. Curr Res Biotechnol 4:10-20.
812   https://doi.org/10.1016/j.crbiot.2021.12.003.
813

814 55. Pakostova E, Johnson DB. 2019. Microbial generation of sulfuric acid from granular
815   elemental sulfur in laboratory-scale bioreactors. Hydrometallurgy 190:105152.
816   https://doi.org/10.1016/j.hydromet.2019.105152.
817

818 56. Azizitorghabeh A, Mahandra H, Ramsay J, Ghahreman A. 2022. A sustainable
819   approach for gold recovery from refractory source using novel BIOX-TC system. J Ind
820   Eng Chem 115: 209–218. https://doi.org/10.1016/j.jiec.2022.08.002.
821

822 57. Remonsellez F, Galleguillos F, Moreno-Paz M, Parro V, Acosta M, Demergasso C.
823   2009. Dynamic of active microorganisms inhabiting a bioleaching industrial heap of
824   low-grade copper sulfide ore monitored by real-time PCR and oligonucleotide
825   prokaryotic acidophile microarray. Microb Biotechnol 2:613–624.
826   https://doi.org/10.1111/j.1751-7915.2009.00112.x.
827

828 58. Hong Y, Valix M. 2014. Bioleaching of electronic waste using acidophilic sulfur
829   oxidising bacteria. J Clean Prod 65:465–472.
830   https://doi.org/10.1016/j.jclepro.2013.08.043.
831

832 59. Arshadi M, Mousavi SM. 2015. Multi-objective optimization of heavy metals
833   bioleaching from discarded mobile phone PCBs: Simultaneous Cu and Ni recovery
834   using Acidithiobacillus ferrooxidans. Sep Purif Technol 147:210–219.
835   https://doi.org/10.1016/j.seppur.2015.04.020.
836

837 60. Moazzam P, Boroumand Y, Rabiei P, Baghbaderani SS, Mokarian P, Mohagheghian F,
838   Mohammed LJ, Razmjou A. 2020. Lithium bioleaching: an emerging approach for the
839   recovery of Li from spent lithium ion batteries. Chemosphere 277:135907.
840   https://doi.org/10.1016/j.chemosphere.2021.130196.
841

842 61. Lee E, Han Y, Park J, Hong J, Silva RA, Kim S, Kim H. 2015. Bioleaching of arsenic from
843   highly contaminated mine tailings using *Acidithiobacillus thiooxidans*. J Environ
844   Manage 147:124–131. https://doi.org/10.1016/j.jenvman.2014.08.019.
845

846 62. Pineda YS, Devries SL, Steiner NC, Block-Cora KA. 2023. Bioleaching of gold in mine
847   tailings by *Alcaligenes faecalis*. Minerals 13:410.
848   https://doi.org/10.3390/min13030410.
849

850 63. Park J, Han Y, Lee E, Choi U, Yoo K, Song Y, Kim H. 2014. Bioleaching of highly
851   concentrated arsenic mine tailings by *Acidithiobacillus ferrooxidans*. Sep Purif
852   Technol 133:291–296. https://doi.org/10.1016/j.seppur.2014.06.054.
853

854 64. Brierley CL, Brierley JA. 2013. Progress in bioleaching : part B : applications of
855   microbial processes by the minerals industries. Appl Microbiol Biotechnol 97:7543–
856   7552. https://doi.org/10.1007/s00253-013-5095-3.
857

858 65. Johnson DB. 2014. Biomining-biotechnologies for extracting and recovering metals
859   from ores and waste materials. Curr Opin Biotechnol 30:24–31.
860   https://doi.org/10.1016/j.copbio.2014.04.008.
861

862 66. Filippov L, Izart C, Hejny H, Hirsch KA, Johnson B, Kahnt R, Märten H, Slabbert W.
863   2017. The BIOMOre Project - A new mining concept for extracting metals from deep
864   ore deposits using biotechnology. Mining Report 153:436–445.
865

866 67. Götze K, Haseneder R, Braeuer AS. 2022. Investigations on strategic element
867   recovery by an underground membrane pilot plant from in-situ extracted bioleaching
868   solutions. Minerals 12:46. https://doi.org/10.3390/min12010046.
869

870 68. Lima VHS, Moura JP, Pissarra TCT, do Valle Junior RF, Silva MMAP de M, Valera CA, De
871   Melo MC, Fernandes LFS, Da Costa AM, Pacheco FAL. 2024. Groundwater flow and
872   transport of metals under deposits of mine tailings: A case study in Brumadinho,
873   Minas Gerais, Brazil. Case Stud Chem Environ Eng 9:100690.
874   https://doi.org/10.1016/j.cscee.2024.100690.
875

876 69. West AC, Van Geel PJ, Raven KG, Nguyen TS, Belfadhel M Ben, Flavelle P. 2003.
877   Groundwater flow and solute transport in a laboratory-scale analogue of a
878   decommissioned in-pit tailings management facility. Can Geotech J 40:326–341.
879   https://doi.org/10.1139/t02-108.
880

881 70. Schaider LA, Senn DB, Brabander DJ, Mccarthy KD, Shine JP. 2007. Characterization
882   of zinc, lead, and cadmium in mine waste: Implications for transport, exposure, and
883   bioavailability. Environ Sci Technol 41:4164–4171.
884   https://doi.org/10.1021/es0626943.
885

886 71. Lapakko K. 2002. Mining, minerals and sustainable development metal mine rock
887   and waste characterization tools: An overview. Mining, Minerals and Sustainable
888   Development (MMSD) working paper 67, 31 p.
889

890 72. Jamieson HE, Walker SR, Parsons MB. 2015. Mineralogical characterization of mine
891   waste. Appl Geochem 57:85-105. https://doi.org/10.1016/j.apgeochem.2014.12.014.
892

893 73. Benidire L, Pereira SIA, Loqman S, Castro PML, Boularbah A. 2022. Physical,
894   chemical, and microbiological characterization of Kettara Mine Tailings, Morocco.
895   Soil Syst 6:23. https://doi.org/10.3390/soilsystems6010023.
896

897 74. Wielinga B, Lucy JK, Moore JN, Seastone OF, Gannon JE. 1999. Microbiological and
898   Geochemical Characterization of Fluvially Deposited Sulfidic Mine Tailings. Appl
899   Environ Microbiol 65:1548-55. https://doi.org/10.1128/AEM.65.4.1548-1555.1999
900

901 75. Wielinga B, Lucy JK, Moore JN, Seastone OF, Gannon JE. 1999. Microbiological and
902   geochemical characterization of fluvially deposited sulfidic mine tailings. Appl
903   Environ Microbiol 65:1548–1555. https://doi.org/10.1128/AEM.65.4.1548-1555.1999.
904

905 76. Kinnunen PHM, Kaksonen AH. 2019. Towards circular economy in mining:
906   Opportunities and bottlenecks for tailings valorization. J Clean Prod 228:153–160.
907   https://doi.org/10.1016/j.jclepro.2019.04.171.
908

909 77. Nieva NE, Borgnino L, García MG. 2018. Long term metal release and acid generation
910   in abandoned mine wastes containing metal-sulphides. Environ Pollut 242:264–276.
911   https://doi.org/10.1016/j.envpol.2018.06.067.
912

913 78. Fortin D, Davis B, Beveridge TJ. 2006. Role of *Thiobacillus* and sulfate-reducing
914   bacteria in iron biocycling in oxic and acidic mine tailings. FEMS Microbiol Ecol
915   21:11–24. https://doi.org/10.1111/j.1574-6941.1996.tb00329.x.
916

917 79. Pepper IL, Zerzghi HG, Bengson SA, Iker BC, Banerjee MJ, Brooks JP. 2012. Bacterial
918   populations within copper mine tailings: Long-term effects of amendment with Class
919   A biosolids. J Appl Microbiol 113:569–577. https://doi.org/10.1111/j.1365-
920   2672.2012.05374.x.
921

922 80. Pakostova E, McAlary M, Marshall S, McGarry S, Ptacek CJ, Blowes DW. 2022.
923   Microbiology of a multi-layer biosolid/desulfurized tailings cover on a mill tailings
924   impoundment. J Environ Manage 302.
925   https://doi.org/10.1016/j.jenvman.2021.114030.
926

927 81. Li Y, Jia Z, Sun Q, Zhan J, Yang Y, Wang D. 2016. Ecological restoration alters microbial
928   communities in mine tailings profiles. Sci Rep 6: 25193.
929   https://doi.org/10.1038/srep25193.
930

82. Pérez-De-Mora A, Burgos P, Madejón E, Cabrera F, Jaeckel P, Schloter M. 2006.
Microbial community structure and function in a soil contaminated by heavy metals:

Effects of plant growth and different amendments. Soil Biol Biochem 38:327–341.
https://doi.org/10.1016/j.soilbio.2005.05.010.

83. Chen M, Grégoire DS, Bain JG, Blowes DW, Hug LA. 2024. Legacy copper/nickel mine
tailings potentially harbor novel iron/sulfur cycling microorganisms within highly
variable communities. Appl Environ Microbiol 90:e00143-24.
https://doi.org/10.1128/aem.00143-24.

84. Mendez MO, Neilson JW, Maier RM. 2008. Characterization of a bacterial community
in an abandoned semiarid lead-zinc mine tailing site. Appl Environ Microbiol
74:3899–3907. https://doi.org/10.1128/AEM.02883-07.

85. Korehi H, Blöthe M, Sitnikova MA, Dold B, Schippers A. 2013. Metal mobilization by
iron- and sulfur-oxidizing bacteria in a multiple extreme mine tailings in the Atacama
Desert, Chile. Environ Sci Technol 47:2189–2196.
https://doi.org/10.1021/es304056n.

86. Chen LX, Li JT, Chen YT, Huang LN, Hua ZS, Hu M, Shu WS. 2013. Shifts in microbial
community composition and function in the acidification of a lead/zinc mine tailings.
Environ Microbiol 15:2431–2444. https://doi.org/10.1111/1462-2920.12114.

87. Mghazli N, Sbabou L, Hakkou R, Ouhammou A, El Adnani M, Bruneel O. 2021.
Description of microbial communities of phosphate mine wastes in Morocco, a
semi-arid climate, using high-throughput sequencing and functional prediction.
Front Microbiol 12:666936. https://doi.org/10.3389/fmicb.2021.666936.

88. Blowes DW, Jambor JL, Hanton-Fong CJ, Lortie L, Douglas Gould W. 1998.
Geochemical, mineralogical and microbiological characterization of a sulphide-
bearing carbonate-rich gold-mine tailings impoundment, Joutel, Québec. Applied
Geochemistry 13:687–705.

89. Diaby N, Dold B, Pfeifer HR, Holliger C, Johnson DB, Hallberg KB. 2007. Microbial
communities in a porphyry copper tailings impoundment and their impact on the
geochemical dynamics of the mine waste. Environ Microbiol 9:298–307.
https://doi.org/10.1111/j.1462-2920.2006.01138.x.

90. Acheampong-Mensah AS, McLemore VT. 2024. Geochemistry of critical minerals in
mine wastes at Hillsboro and Steeple Rock districts, New Mexico. SME Annual
Conference Preprint 24-042:340-346. Available at
https://store.smenet.org/20slp36/340

91. Motlagh ZK, Mclemore VT. 2024. Critical minerals and acid mine drainage in Black
Hawk mine waste, Grant County, New Mexico. SME Annual Conference Preprint 24-
022:176-183. Available at https://store.smenet.org/20slp36/176

92.   McLemore VT. 2008. Geochemistry and statistical analyses of epithermal veins at the Carlisle and Center mines, Steeple Rock District, New Mexico, USA. Arizona Geological Society Digest 22:485–496.

93.   McLemore VT. 1993. Geology and geochemistry of the mineralization and alteration in the Steeple Rock district, Grant County, New Mexico and Greenlee County, Arizona. Doctoral dissertation, University of Texas at El Paso.

94.   Dunn PG. 1984. Geologic studies during the development of the Copper Flat porphyry deposit. Min Eng 36:151–159.

95.   Dunn PG, Titley SR. 1982. Geology of the Copper Flat porphyry copper deposit, Hillsboro, Sierra County, New Mexico, p. 313–326. *In* Advances in Geology of the Porphyry-Copper Deposits. University of Arizona Press, Tucson.

96.   McLemore VT, Munroe EA, Heizler MT, McKee C. 1999. Geochemistry of the Copper Flat porphyry and associated deposits in the Hillsboro mining district, Sierra County, New Mexico, USA. J Geochem Explor 67:167–189. https://doi.org/10.1016/S0375-6742(99)00072-2.

97.   McLemore VT, Sutphin DM, Hack DR, Pease TC. 1996. Mining history and mineral resources of the Mimbres Resource Area. Doña Ana, Luna, Hidalgo, and Grant Counties, New Mexico: New Mexico Bureau of Miners and Mineral Resources Open File Report 424:252. https://geoinfo.nmt.edu/publications/openfile/downloads/400-499/424/ofr_424.pdf

98.   Gillerman E, Whitebread DH. 1953. The uranium-bearing nickel-cobalt-native silver deposits in the Black Hawk District, Grant County, New Mexico. 1009-K US Geological Survey Bulletin 313:1–41.

99.   Havlena ZE, Hose LD, DuChene HR, Baker GM, Powell JD, Labrado AL, Brunner B, Jones DS. 2024. Origin and modern microbial ecology of secondary mineral deposits in Lehman Caves, Great Basin National Park, NV, USA. Geobiology 22:e12594. https://doi.org/10.1111/gbi.12594.

100. Li L, Liu Z, Zhang M, Meng D, Liu X, Wang P, Li X, Jiang Z, Zhong S, Jiang C, Yin H. 2020. Insights into the metabolism and evolution of the genus *Acidiphilium*, a typical acidophile in acid mine drainage. mSystems 5. https://doi.org/10.1128/msystems.00867-20.

101. Jiang ZM, Mou T, Sun Y, Su J, Yu LY, Zhang YQ. 2023. Environmental distribution and genomic characteristics of *Solirubrobacter*, with proposal of two novel species. Front Microbiol 14:1267771. https://doi.org/10.3389/fmicb.2023.1267771.

1021

102. Li WJ, Zhang YQ, Schumann P, Chen HH, Hozzein WN, Tian XP, Xu LH, Jiang CL. 2006. *Kocuria aegyptia* sp. nov., a novel actinobacterium isolated from a saline, alkaline desert soil in Egypt. Int J Syst Evol Microbiol 56:733–737. https://doi.org/10.1099/ijs.0.63876-0.

103. Gholami M, Etemadifar Z, Bouzari M. 2015. Isolation a new strain of *Kocuria rosea* capable of tolerating extreme conditions. J Environ Radioact 144:113–119. https://doi.org/10.1016/j.jenvrad.2015.03.010.

104. Chang Y juan, Land M, Hauser L, Chertkov O, del Rio TG, Nolan M, Copeland A, Tice H, Cheng JF, Lucas S, Han C, Goodwin L, Pitluck S, Ivanova N, Ovchinikova G, Pati A, Chen A, Palaniappan K, Mavromatis K, Liolios K, Brettin T, Fiebig A, Rohde M, Abt B, Göker M, Detter JC, Woyke T, Bristow J, Eisen JA, Markowitz V, Hugenholtz P, Kyrpides NC, Klenk HP, Lapidus A. 2011. Non-contiguous finished genome sequence and contextual data of the filamentous soil bacterium *Ktedonobacter racemifer* type strain (SOSP1-21 T). Stand Genomic Sci 5:97–111. https://doi.org/10.4056/sigs.2114901.

105. Yan B, Guo X, Liu M, Huang Y. 2020. *Ktedonosporobacter rubrisoli* gen. nov., sp. nov., a novel representative of the class *Ktedonobacteria*, isolated from red soil, and proposal of *Ktedonosporobacteraceae* fam. nov. Int J Syst Evol Microbiol 70:1015–1025. https://doi.org/10.1099/ijsem.0.003864.

106. Gupta A, Saha A, Sar P. 2021. *Thermoplasmata* and *Nitrososphaeria* as dominant archaeal members in acid mine drainage sediment of Malanjkhand Copper Project, India. Arch Microbiol 203:1833–1841. https://doi.org/10.1007/s00203-020-02130-4.

107. Sauder LA, Albertsen M, Engel K, Schwarz J, Nielsen PH, Wagner M, Neufeld JD. 2017. Cultivation and characterization of *Candidatus* Nitrosocosmicus exaquare, an ammonia-oxidizing archaeon from a municipal wastewater treatment system. ISME J 11:1142–1157. https://doi.org/10.1038/ismej.2016.192.

108. Stieglmeier M, Klingl A, Alves RJE, Rittmann SKMR, Melcher M, Leisch N, Schleper C. 2014. *Nitrososphaera viennensis* gen. nov., sp. nov., an aerobic and mesophilic, ammonia-oxidizing archaeon from soil and a member of the archaeal phylum Thaumarchaeota. Int J Syst Evol Microbiol 64:2738–2752. https://doi.org/10.1099/ijs.0.063172-0.

109. Wan S, Liao X, Zhou TT, Wu Y, Hu A, Yan D, Zhang J, Long XE. 2021. Shift in archaeal community along a soil profile in coastal wheat-maize rotation fields of different reclamation ages. Land Degrad Dev 32:4162–4173. https://doi.org/10.1002/ldr.4022.

1064   110.   Clark IM, Hughes DJ, Fu Q, Abadie M, Hirsch PR. 2021. Metagenomic approaches
1065   reveal differences in genetic diversity and relative abundance of nitrifying bacteria
1066   and archaea in contrasting soils. Sci Rep 11:15905. https://doi.org/10.1038/s41598-
1067   021-95100-9.
1068

1069   111.   Marusenko Y, Garcia-Pichel F, Hall SJ. 2015. Ammonia-oxidizing archaea respond
1070   positively to inorganic nitrogen addition in desert soils. FEMS Microbiol Ecol 91.
1071   https://doi.org/10.1093/femsec/fiu023.
1072

1073   112.   Leininger S, Urich T, Schloter M, Schwark L, Qi J, Nicol GW, Prosser JI, Schuster SC,
1074   Schleper C. 2006. Archaea predominate among ammonia-oxidizing prokaryotes in
1075   soils. Nature 442:806–809. https://doi.org/10.1038/nature04983.
1076

1077   113.   He JZ, Hu HW, Zhang LM. 2012. Current insights into the autotrophic thaumarchaeal
1078   ammonia oxidation in acidic soils. Soil Biol Biochem 55:146–154.
1079   https://doi.org/10.1016/j.soilbio.2012.06.006.
1080

1081   114.   Ishii S, Ashida N, Ohno H, Segawa T, Yabe S, Otsuka S, Yokota A, Senoo K. 2017.
1082   *Noviherbaspirillum denitrificans* sp. nov., a denitrifying bacterium isolated from rice
1083   paddy soil and *Noviherbaspirillum autotrophicum* sp. nov., a denitrifying,
1084   facultatively autotrophic bacterium isolated from rice paddy soil and proposal to
1085   reclassify *Herbaspirillum massiliense* as *Noviherbaspirillum massiliense* comb. nov.
1086   Int J Syst Evol Microbiol 67:1841–1848. https://doi.org/10.1099/ijsem.0.001875.
1087

1088   115.   Chaudhary DK, Kim J. 2017. *Noviherbaspirillum agri* sp. nov., isolated from reclaimed
1089   grassland soil, and reclassification of *Herbaspirillum massiliense* (Lagier et al., 2014)
1090   as *Noviherbaspirillum massiliense* comb. nov. Int J Syst Evol Microbiol 67:1508–1515.
1091   https://doi.org/10.1099/ijsem.0.001747.
1092

1093   116.   Delamuta JRM, Ribeiro RA, Ormeño-Orrillo E, Melo IS, Martínez-Romero E, Hungria
1094   M. 2013. Polyphasic evidence supporting the reclassification of *Bradyrhizobium
1095   japonicum* group Ia strains as *Bradyrhizobium diazoefficiens* sp. nov. Int J Syst Evol
1096   Microbiol 63:3342–3351. https://doi.org/10.1099/ijs.0.049130-0.
1097

1098   117.   Griebsch A, Matschiavelli N, Lewandowska S, Schmidtke K. 2020. Presence of
1099   *Bradyrhizobium* sp. under continental conditions in Central Europe. Agriculture
1100   (Switzerland) 10:1–17. https://doi.org/10.3390/agriculture10100446.
1101

1102   118.   Mehrani MJ, Sobotka D, Kowal P, Ciesielski S, Makinia J. 2020. The occurrence and
1103   role of *Nitrospira* in nitrogen removal systems. Bioresour Technol 303.
1104   https://doi.org/10.1016/j.biortech.2020.122936.
1105

1106   119.   Lee HJ, Lee SH, Lee SS, Lee JS, Kim Y, Kim SC, Jeon CO. 2014. *Ramlibacter solisilvae*
1107   sp. nov., isolated from forest soil, and emended description of the genus

1108   *Ramlibacter*. Int J Syst Evol Microbiol 64:1317–1322.
1109   https://doi.org/10.1099/ijs.0.058396-0.

1110

1111   120.   Khan SA, Kim HM, Baek JH, Jung HS, Jeon CO. 2021. *Ramlibacter terrae* sp. nov. And
1112   *Ramlibacter montanisoli* sp. nov., Isolated from soil. J Microbiol Biotechnol 31:1210–
1113   1217. https://doi.org/10.4014/jmb.2105.05023.

1114

1115   121.   Heulin T, Barakat M, Christen R, Lesourd M, Sutra L, De Luca G, Achouak W. 2003.
1116   *Ramlibacter tataouinensis* gen. nov., sp. nov., and *Ramlibacter henchirensis* sp. nov.,
1117   cyst-producing bacteria isolated from subdesert soil in Tunisia. Int J Syst Evol
1118   Microbiol 53:589–594. https://doi.org/10.1099/ijs.0.02482-0.

1119

1120   122.   Yan ZF, Trinh H, Moya G, Lin P, Li CT, Kook M, Yi TH. 2017. *Ramlibacter rhizophilus* sp.
1121   nov., isolated from rhizosphere soil of national flower Mugunghwa from South Korea.
1122   Int J Syst Evol Microbiol 67:3773–3777. https://doi.org/10.1099/ijsem.0.002191.

1123

1124   123.   Lee JH, Kim DI, Choe HN, Lee SD, Seong CN. 2017. *Sphingomonas*
1125   *limnosediminicola* sp. nov. and *Sphingomonas palustris* sp. nov., isolated from
1126   freshwater environments. Int J Syst Evol Microbiol 67:2834–2841.
1127   https://doi.org/10.1099/ijsem.0.002029.

1128

1129   124.   Huang HD, Wang W, Ma T, Li GQ, Liang FL, Liu RL. 2009. *Sphingomonas*
1130   *sanxanigenens* sp. nov., isolated from soil. Int J Syst Evol Microbiol 59:719–723.
1131   https://doi.org/10.1099/ijs.0.000257-0.

1132

1133   125.   Hussain S, Devers-Lamrani M, El Azhari N, Martin-Laurent F. 2011. Isolation and
1134   characterization of an isoproturon mineralizing *Sphingomonas* sp. strain SH from a
1135   French agricultural soil. Biodegradation 22:637–650. https://doi.org/10.1007/s10532-
1136   010-9437-x.

1137

1138   126.   Qiong W, Fengshan P, Xiaomeng X, Rafiq MT, Xiao'e Y, Bao C, Ying F. 2021. Cadmium
1139   level and soil type played a selective role in the endophytic bacterial community of
1140   hyperaccumulator *Sedum alfredii* Hance. Chemosphere 263:127986.
1141   https://doi.org/10.1016/j.chemosphere.2020.127986.

1142

1143   127.   An F, Li H, Diao Z, Lv J. 2019. The soil bacterial community in cropland is vulnerable
1144   to Cd contamination in winter rather than in summer. Environ Sci Pollut Res 26:114–
1145   125. https://doi.org/10.1007/s11356-018-3531-8.

1146

1147   128.   McNeill BJ, Pakostova E, Bain JG, Gould WD, Amos RT, Wilson GW, Ptacek CJ, Blowes
1148   DW. 2020. Microbial community structure within a weathered waste-rock pile
1149   overlain by a monolayer soil cover. Appl Geochem 114:104531.
1150   https://doi.org/10.1016/j.apgeochem.2020.104531.

1151

1152 129. Liu J, Hua ZS, Chen LX, Kuang JL, Li SJ, Shu WS, Huang LN. 2014. Correlating
1153 microbial diversity patterns with geochemistry in an extreme and heterogeneous
1154 environment of mine tailings. Appl Environ Microbiol 80:3677–3686.
1155 https://doi.org/10.1128/AEM.00294-14.
1156

1157 130. Falagán C, Johnson DB. 2014. *Acidibacter ferrireducens* gen. nov., sp. nov.: an
1158 acidophilic ferric iron-reducing gammaproteobacterium. Extremophiles 18:1067–
1159 1073. https://doi.org/10.1007/s00792-014-0684-3.
1160

1161 131. Johnson DB. 2014. Recent developments in microbiological approaches for securing
1162 mine wastes and for recovering metals from mine waters. Minerals 4:279–292.
1163 https://doi.org/10.3390/min4020279.
1164

1165 132. Johnson DB. 2018. The evolution, current status, and future prospects of using
1166 biotechnologies in the mineral extraction and metal recovery sectors. Minerals
1167 8:343. https://doi.org/10.3390/min8080343.
1168

1169 133. Kojima H, Shinohara A, Fukui M. 2015. *Sulfurifustis variabilis* gen. nov., sp. nov., a
1170 sulfur oxidizer isolated from a lake, and proposal of *Acidiferrobacteraceae* fam. nov.
1171 and *Acidiferrobacterales* ord. nov. Int J Syst Evol Microbiol 65:3709–3713.
1172 https://doi.org/10.1099/ijsem.0.000479.
1173

1174 134. Blowes DW, Al T, Lortie L, Gould WD, Jambor JL. 1995. Microbiological, chemical,
1175 and mineralogical characterization of the Kidd Creek mine tailings impoundment,
1176 Timmins area, Ontario. Geomicrobiol J 13:13–31.
1177 https://doi.org/10.1080/01490459509378001.
1178

1179 135. Dopson M, Johnson DB. 2012. Biodiversity, metabolism and applications of
1180 acidophilic sulfur-metabolizing microorganisms. Environ Microbiol 14:2620–2631.
1181 https://doi.org/10.1111/j.1462-2920.2012.02749.x.
1182

1183 136. Rawlings DE, Johnson DB. 2007. The microbiology of biomining: Development and
1184 optimization of mineral-oxidizing microbial consortia. Microbiology 153:315–324.
1185 https://doi.org/10.1099/mic.0.2006/001206-0.
1186

1187 137. Jones DS, Lapakko KA, Wenz ZJ, Olson MC, Roepke EW, Sadowsky MJ, Novak PJ,
1188 Bailey J V. 2017. Novel microbial assemblages dominate weathered sulfide bearing
1189 rock from copper-nickel deposits in the Duluth Complex, Minnesota, USA. Appl
1190 Environ Microbiol 83:e00909-17. https://doi.org/10.1128/AEM.00909-17.
1191

1192 138. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput
1193 sequencing reads. EMBnet J 17:10. https://doi.org/10.14806/ej.17.1.200.
1194

1195 139. Zhang J, Kobert K, Flouri T, Stamatakis A. 2014. PEAR: A fast and accurate Illumina
1196   Paired-End reAd mergeR. Bioinformatics 30:614–620.
1197   https://doi.org/10.1093/bioinformatics/btt593.
1198

1199 140. Schmieder R, Edwards R. 2011. Quality control and preprocessing of metagenomic
1200   datasets. Bioinformatics 27:863–864. https://doi.org/10.1093/bioinformatics/btr026.
1201

1202 141. Edgar RC. 2013. UPARSE: Highly accurate OTU sequences from microbial amplicon
1203   reads. Nat Methods 10:996–998. https://doi.org/10.1038/nmeth.2604.
1204

1205 142. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016. VSEARCH: A versatile open
1206   source tool for metagenomics. PeerJ 2016. https://doi.org/10.7717/peerj.2584.
1207

143. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA,
Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ,
Weber CF. 2009. Introducing mothur: Open-source, platform-independent,
community-supported software for describing and comparing microbial
communities. Appl Environ Microbiol 75:7537–7541.
https://doi.org/10.1128/AEM.01541-09.
1214

1215 144. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glöckner FO. 2007.
1216   SILVA: A comprehensive online resource for quality checked and aligned ribosomal
1217   RNA sequence data compatible with ARB. Nucleic Acids Res 35:7188–7196.
1218   https://doi.org/10.1093/nar/gkm864.
1219

1220 145. R Development Core Team. 2023. R Statistical Software RStudio. 4.2.1.
1221

146. Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB,
Solymos P, Stevens MHH, Szoecs E, Barbour M, Bedward M, Bolker B, Borcard D,
Carvalho G, Chirico M, De Caceres M, Durand S, Antoniazi Evangelista HB, FitzJohn
R, Friendly M, Furneaux B, Hannigan G, Hill MO, Lahti L, McGlinn D, Ouellette M-H,
Ribeiro Cunha E, Smith T, Stier A, Ter Braak CJF, Weedon J. 2024. Community ecology
package. 2. R package version.
1228

1229 147. Maechler M, Rousseeuw P, Struyf A, Hubert M, Hornik K, Studer M, Roudier P,
1230   Gonzalez J, Kozlowski K, Schubert E, Murphy K. 2022. Finding groups in data: Cluster
1231   analysis extended Rousseeuw et al. Documentation for package 'cluster' version
1232   2.1.8
1233

1234 148. Wickham H, Chang W, Henry L, Pedersen TL, Takahashi K, Wilke C, Woo K, Yutani H,
1235   Dunnington D, van den Brand T. 2024. Create elegant data visualisations using the
1236   grammar of graphics. 2.1. R package version.
1237

1238   149.   Martinez Arbizu P. 2020. pairwiseAdonis: Pairwise multilevel comparison using
1239   adonis. 0.4. R package version 04. R package version.

1240
1241
1242

1243 **Figures and Tables**

TABLE 1 Waste rock types sampled at inactive mine sites

Mine	Bulk waste rock	Acidic seep	Flooded adit	Tailings
Copper Flat (CF)	Y	Y	-	Y
Center (CN)	Y	-	Y	-
Carlisle (CY)	Y	-	Y	Y
Black Hawk (BH)	Y	-	-	-
Alhambra (AH)	Y	-	-	-

1252 "-" indicates this sample type was not present (and therefore not collected)
1253
1254
1255
1256

TABLE 2 Cell counts from select waste rock and tailings

Sample	Location	Sample type	Cells/g wet sediment	Sample notes
AH24-13	Alhambra	bulk waste	9.9×10^8	soil cover, waste composite
BH24-13	Black Hawk	bulk waste	1.6×10^9	soil cover, waste composite
BH24-23	Black Hawk	bulk waste	1.5×10^9	soil cover, waste composite
CN23-13	Center	bulk waste	1.1×10^{10}	soil cover, waste composite
CY23-13	Carlisle	bulk waste	3.9×10^9	waste composite
CF22-601*	Copper Flat	bulk waste	2.4×10^8	
CF22-603*	Copper Flat	bulk waste	1.2×10^8	
CF22-604*	Copper Flat	bulk waste	6.1×10^7	
CF22-606*	Copper Flat	bulk waste	4.7×10^8	
CF23-621	Copper Flat	tailings	1.0×10^8	above oxidation front
CF23-622	Copper Flat	tailings	9.0×10^7	at oxidation front
CF23-623	Copper Flat	tailings	2.0×10^7	below oxidation front
CF23-631	Copper Flat	tailings	4.8×10^7	above oxidation front, tailings composite
CY23-20	Carlisle	tailings	9.7×10^7	above oxidation front
CY23-21	Carlisle	tailings	6.9×10^8	at oxidation front
CY23-22	Carlisle	tailings	1.6×10^8	below oxidation front

*Individual (composite sample was not available for cell counting from the CF composite)

1257

1258

1259

1260

1261

1262

1263
1264
1265   **FIG 1**   (a) Map of New Mexico showing the three districts sampled. The Hillsboro district
1266   contains the Copper Flat mine (CF), the Steeple Rock district contains the Center (CN) and
1267   Carlisle mines (CY), and the Black Hawk District contains the Black Hawk (BH) and
1268   Alhambra mines (AH). Representative photos show (b) bulk waste rock (white arrows) at CY,
1269   (c) a seasonal acidic seep at CF, and (d) tailings at CY.
1270

1274 **FIG 2** Shannon diversity (top) and expected species richness (bottom) boxplots of bulk
1275 waste rock and tailings samples from the Alhambra mine (AH), Black Hawk mine (BH),
1276 Copper Flat mine (CF), Center mine (CN), and Carlisle mine (CY). Sites CF and CY did not
1277 have a soil cover.

FIG 3 Bar chart showing the relative abundance of specific taxa in waste samples by sample type and location. 'Unclassified' or 'uncultured' refers to OTUs that were either unclassified at the genus level or higher, or from groups with no known cultured representative.

FIG 4   A non-metric multidimensional scaling (NMDS) ordination of samples collected from waste rock and tailings all five inactive mine sites (seep and adit sediments are not included). In the left panel, samples are coded by sample location. Unfilled shapes are rRNA gene libraries, and filled shapes are rRNA transcript libraries from composite samples. In the right panel, samples are coded by sample type (either waste rock or tailings).

FIG 5 Non-metric multidimensional scaling (NMDS) ordinations of samples from (a) the Alhambra, Black Hawk, Copper Flat, Carlisle, and Center mines (AH, BH, CF, CY, and CN, respectively) and (b) only AH, BH, and CN, for which bulk waste rock was topped with a soil cover. Vectors of fitted geochemical variables indicate by asterisks are statistically significant (*** = p <0.001, ** = p <0.01, and * = p <0.05). RNA transcript samples are represented by filled shapes, and DNA samples are represented by open shapes.

1305
1306
FIG 6 Non-metric multidimensional scaling (NMDS) ordinations of samples from each
mine site comparing composite samples (filled symbols) and individual samples (open
symbols) from the same locations. Squares indicate rRNA gene libraries, and triangles
indicate rRNA transcript libraries. The bottom right panel is tailings from site CF, and the
others are all waste rock. In the bottom middle panel, the upper and lower bench at site BH
refers to samples collected from two distinct benches from previous mining activity. The
upper bench refers to the bench that is higher in elevation (by 5m) that is the 'top' of the
waste rock piles. In the legends, "comp" = composite sample.
1315

1316
1317
1318 **FIG 7** Phyla-level community composition of samples by sample type and rRNA gene
1319 versus rRNA transcript libraries from composite samples.
1320
1321
1322

1323
1324
FIG 8 A two-way cluster analysis comparing 16S rRNA gene and transcript libraries from
bulk waste rock. The points are scaled according to relative abundance (%). The taxonomic
affiliation shows phylum and genus for each OTU, except for those that unclassified at the
genus level, for which the highest available taxonomic classification is provided. BH waste
1 and 2 refer to samples from the upper and lower bench, respectively, and the two CF
waste RNA libraries are from separate composite samples from that site (Fig 6; Table S1).
1331
1332
1333